1

Million years old recombination suppression and balancing selection in a region

partially linked to the mating-type locus in the invasive chestnut blight fungal pathogen

Cryphonectria parasitica

Fanny E. Hartmann[1]*, Ricardo C. Rodriguez de la Vega[1], Arthur Demené[2], Thomas Badet[3], Jean-Philippe Vernadet[1], Quentin Rougemont[1], Amandine Labat[1], Alodie Snirc[1], Lea Stauber[5], Daniel Croll[3], Simone Prospero[4], Cyril Dutech[2], Tatiana Giraud[1]

[1] Ecologie Systematique et Evolution, CNRS, Université Paris-Saclay, AgroParisTech, 91198 Gif-sur-Yvette, France

[2] BIOGECO, INRAE, Univ. Bordeaux, 69 route d'Arcachon, Cestas F-33610, France.

[3] Laboratory of Evolutionary Genetics, Institute of Biology, University of Neuchâtel, Neuchâtel, Switzerland

[4] Swiss Federal Research Institute WSL, Birmensdorf, Switzerland

[5] Institute for Infectious Diseases, University of Bern, Switzerland

*Corresponding author: Fanny E. Hartmann

E-mail: fanny.hartmann@universite-paris-saclay.fr

Key words: recombination suppression, fungi, mating-type chromosome, inversion, deleterious allele sheltering, supergenes, evolutionary strata, sex chromosomes

Abstract

Recombination suppression often evolves in sex chromosomes and around mating-type loci.

In the invasive chestnut blight fungus *Cryphonectria parasitica* (Ascomycota), a genomic

region was previously suggested to lack recombination and to be partially linked to the

mating-type (MAT) locus based on the analysis of a few progenies. Using hundreds of

available *C. parasitica* genomes and generating several new high-quality genome assemblies

from the native and introduced range of the pathogen, we show that a ca. 1.2 Mb genomic

region proximal to the mating-type locus lacks recombination worldwide. In invasive

populations, this MAT-proximal region displayed two highly differentiated haplotypes, that

were strongly associated to mating types, but not completely. High-quality assemblies

revealed an inversion in one of the haplotypes and footprints of degeneration worldwide, the

MAT-proximal region being enriched in gene disruptions, non-synonymous substitutions and

transposable elements in both haplotypes. The divergence between the two haplotypes was

estimated to have occurred at least 1.5 million years ago and two haplotypes segregate in all

continents, including the native range. High differentiation between haplotypes, their

occurrence on different continents, their balanced frequencies within populations, their

genomic rearrangements and degeneration worldwide, altogether suggest an ancient

recombination suppression maintained by selection. The MAT-Proximal region carries

multiple genes upregulated under virus infection or vegetative incompatibility reaction. This

study sheds light on a case of a large non-recombining region partially linked to a mating

compatibility locus, and on balancing selection maintaining differentiated haplotypes,

possibly involving deleterious mutations and/or host or virus adaptation in a devastating tree

pathogen.

Introduction

Recombination increases the efficiency of selection, the purging of deleterious alleles and the generation of potentially beneficial allelic combinations (Otto and Lenormand 2002). However, recombination can also break up beneficial allelic combinations and, therefore, can be selected against, generating supergenes, i.e., large regions without recombination encompassing multiple genes (Schwander et al. 2014). This is often the case in genomic regions controlling mating compatibility, such as regions determining sex, mating type or self-incompatibility in plants, animals, fungi, algae and oomycetes (Charlesworth et al. 2005; Bergero and Charlesworth 2009; Umen 2011; Charlesworth 2016; Dussert et al. 2020; Hartmann et al. 2021). Such recombination suppression keeps alleles linked across different genes that prevent self-compatibility or intermediate sexual phenotypes.

Recombination suppression can sometimes extend away from sex-determining genes or mating-type loci (Bergero and Charlesworth 2009; Furman et al. 2020; Hartmann et al. 2021). Such extension can be due to antagonistic selection, i.e. selection for different alleles in different sexes or mating types (Charlesworth 2023). Extension of recombination suppression can also occur near mating-type loci or sex-determining genes because of selection for inversions carrying fewer deleterious mutations than average in the population, their few deleterious mutations being sheltered when linked to a permanently heterozygous allele, such as on a Y chromosome or a mating-type chromosome (Jay et al. 2022; Jay, D.L. Jeffries, Hartmann, et al. 2024).

On the long term, however, because of less efficient selection, genomic regions without recombination show signs of degeneration, such as transposable element accumulation, rearrangements, non-synonymous substitutions, decreased gene expression, reduced frequency of optimal codons and gene losses (Bachtrog 2013; Carpentier et al. 2022; Duhamel et al. 2023). The non-recombining regions can nevertheless be maintained by balancing selection (Jay et al. 2021; Berdan et al. 2022). Chromosomal rearrangements such as inversions are often considered to be the mechanistic cause of recombination suppression, although they may also be a mere consequence of recombination suppression. The evolutionary mechanisms leading to recombination suppression and allowing their persistence remain however poorly known (Jay, D. Jeffries, Hartmann, et al. 2024). Reporting more

diverse constellations of recombination suppression in a variety of organisms with contrasting
life-history traits is important for understanding the general patterns of recombination
suppression, their evolution and maintenance (Ironside 2010; Charlesworth 2016; Furman et
al. 2020; Hartmann et al. 2021).

In the invasive chestnut blight fungal pathogen *Cryphonectria parasitica*, mating
compatibility is determined by a mating-type locus that displays two alleles, called *MAT-1*
and *MAT-2* that are both permanently heterozygous in the diploid and dikaryotic stages
(McGuire et al. 2001). A large region without recombination was suggested to occur near the
mating-type locus, although not completely linked to it (*i.e.,* at 3.9 cM of the mating-type
locus), based on segregation of RAPD and RFLP markers in three crosses involving five
different parents, from the USA, Japan or Italy (Kubisiak and Milgroom 2006a). Recent
genome-wide association studies in a worldwide collection and in local *C. parasitica*
populations in southern Switzerland found significant association of SNPs with the mating-
type locus across a large region (>1 Mb), further suggesting the existence of reduced
recombination in this region (Stauber et al. 2021; Stauber et al. 2022). This genomic region
was enriched in SNPs, transposable elements and copy-number variants, such as deletions,
which are consistent with sequence degeneration and a lack of recombination (Stauber et al.
2021), and which prevented its assembly so far.

The evolutionary history of the species is well documented as it is an invasive and highly
damaging pathogen, having almost caused the extinction of American chestnut (*Castanea*
dentata) in North America (Anagnostakis 1987). From its center of origin in Asia and its
original hosts, the Chinese chestnut (*Ca. mollissima*) and the Japanese chestnut (*Ca. crenata*),
C. parasitica has first invaded North America, killing millions of American chestnuts (*Ca.*
dentata). In Europe, at least two distinct introduction events occurred, on the European
chestnut (*Ca. sativa*): one introduction from North America to Italy, and one directly from
Asia, probably to the Pyrenees Mountains. European strains will be therefore hereafter
referred to as invasive European strains introduced from North America or invasive European
strains introduced directly from Asia. The chestnut blight symptoms have been less severe in
Europe than in North America, due to lower susceptibility of *C. sativa* and a virus infecting *C.*
parasitica and causing hypovirulence (Dutech et al. 2010; Dutech et al. 2012). The genetic
determinants of the adaptation of the pathogen to its new environments and hosts remain
largely unknown (Lovat and Donnelly 2019).

Lineages with an apparently clonal structure have been identified in invasive populations while sexually reproducing populations occur both in the native and introduced ranges (Dutech et al. 2012; Stauber et al. 2021). The lineages with a predominant clonal structure likely still undergo rare sex events, as shown by the introgression of the missing mating-type allele in most of them (Demené et al. 2019). Furthermore, although this ascomycete fungus is mostly found as haploid mycelia, some isolates have been reported to be heterokaryotic at the mating-type locus in several invasive European and North American populations, i.e., with cells carrying different nuclei, of the opposite mating types *MAT-1* and *MAT-2* (McGuire et al. 2004; McGuire et al. 2005; Dutech et al. 2010; Stauber et al. 2021; Stauber et al. 2022). The theoretical model of recombination suppression based on recessive deleterious mutations showed that non-recombining fragments could be selected for even with a substantial fraction of the life cycle being haploid (Jay et al. 2022). Hence, due to previous evidence of recombination suppression, the natural occurrence of a heterokaryotic stage and its archetypical evolutionary history of invasive pathogen, *C. parasitica* seems an excellent biological model to study the evolution of recombination suppression and test recent theoretical predictions about the existence of non-recombining regions only partially linked to sex- or mating-type determining loci.

Here, we therefore studied the occurrence of recombination suppression in the genomic region proximal to the mating-type locus in *C. parasitica,* using the previously published genomes and further generated six new high-quality long-read based genome assemblies of strains from the native and introduced range of the pathogen. More specifically, we tested whether recombination was completely suppressed in this region partially linked to the mating-type locus in *C. parasitica*. We first analyze sexually reproducing invasive European populations, for which extensive genomic datasets are available in local populations. We assessed, in these populations, the degree of linkage disequilibrium within the MAT-proximal region, the level of differentiation between haplotypes, as well as the frequency of the two non-recombining haplotypes and their association with mating types. We investigated whether other regions of the genome showed reduced recombination rates and whether genomic footprints of degeneration were present in the non-recombining region, such as genomic rearrangements, transposable element accumulation, non-synonymous substitutions and gene disruptions. We also tested whether this non-recombining region has been gradually expanding, estimated the age of recombination suppression and investigated whether the

predicted gene functions in the MAT-proximal region could help understanding the
evolutionary cause for recombination suppression and partial linkage to the mating-type
locus. We then extended the analyses to other populations with less extensive genomic
resources, i.e. other invasive populations and native Asian populations, investigating the
presence of recombination suppression footprints and the occurrence of the two differentiated
haplotypes worldwide, as well as the possibility of introgression, using available genomes of
closely related species, pathogenic and non-pathogenic (Stauber et al. 2021). Finally, we used
available expression data to investigate whether the MAT-Proximal region carried genes
upregulated under infection by the virus responsible for hypovirulence or during vegetative
incompatibility reactions, a phenomenon avoiding hyphal fusions between individuals,
considered to protect against virus transmission.

Results

Footprints of recombination suppression in a large region (> 1 Mb) proximal to the
mating-type locus in sexually reproducing invasive populations

We first performed population genomic analyses on available genome sequences within local
populations sampled in southern Switzerland over two temporal frames (early 1990 and
2019), and within the more broadly geographically distributed CL1 cluster in central and
southeastern Europe, both having a recombining genetic structure and being invasive
European populations introduced from North America (Table S1 (Stauber et al. 2021)). We
studied only monokaryotic strains, i.e. having either *MAT-1* or *MAT-2* in their genomes, for
phasing haplotypes and performed stringent SNPs filtering by masking repeats on the EP155
reference genome (Crouch et al. 2020) and removing missing data and rare variants to ensure
robustness of our analyses. We identified 8,900 SNPs segregating among 71 strains of the
1990 Swiss population (35 MAT-1; 36 MAT-2; Table S1), 9,646 SNPs segregating among 62
strains of the Swiss 2019 population (20 MAT-1; 42 MAT-2; Table S1) and 15,104 SNPs
segregating among 88 strains of the CL1 cluster (41 MAT-1; 47 MAT-2; Table S1), which
represent an average density of about 20 SNPs/100 kb, in agreement with previous studies in
this invasive fungus (Stauber et al. 2021; Stauber et al. 2022). We investigated the genome-
wide linkage disequilibrium (LD) landscape and looked for large blocks (>1 Mb) of high LD
(r2>0.9) among SNPs within contigs, to look for signatures of reduced recombination along
the genome. LD indeed decays by half across a few 100 kb on average in C. parasitica in
recombining regions of the genome (Demené et al. 2019). Therefore, high LD beyond >1 Mb

represents strong evidence of recombination cessation and the SNP density is sufficient to
study such LD variation.
190
We found a large block of high LD on the chromosome called scaffold_2 of the EP155
genome in the three populations (red arrow in Fig 1 and Fig S1), corresponding to the mating-
type chromosome. The high-LD block was located near the mating-type locus but did not
encompass the mating-type locus itself and will be called hereafter the "MAT-proximal
region". In the 1990 and 2019 Swiss populations, SNPs between 0.493 Mb and 1.720 Mb in
the mating-type chromosome were all in high LD (with $r^2>0.9$ between SNPs distant by up to
1 Mb; in this region, mean r^2 of all SNP pairs in the Swiss 1990 population = 0.89; mean r^2 of
all SNP pairs in the Swiss 2019 population = 0.83; Fig S1). In the CL1 cluster, the large high-
LD block was also present, but was split into two blocks, a large one and a smaller, peripheral
one (orange arrows in Fig 1). We indeed detected values of $r^2>0.9$ between SNPs distant by
up to 1 Mb, between 0.532 Mb and 1.535 Mb (mean r^2 of all SNP pairs = 0.94). Between
1.535 and 1.717 Mb, SNPs were also in high LD (mean r^2 = 0.84). The LD between the two
high-LD blocks was lower (r^2= 0.59) than within blocks but was still higher than elsewhere
along the genome (orange rectangle at the right border of the red triangle; Fig 1). A similar
pattern also existed in the Swiss populations (orange rectangle at the right border of the red
triangle; Fig S1), although less marked. This is likely due to rare recombination events at one
specific locus near the edge of the large fully non-recombining region, lowering the LD level
in populations between the two parts of the otherwise fully non-recombining region. The
mating-type locus was located at 1.737 Mb in the EP155 genome, distant by 17 kb and 20 kb
from the LD block in the Swiss populations and CL1 cluster, respectively. SNP density was
on average 48 SNPs/100 kb in the high-LD block region, i.e. higher than in the rest of the
chromosome (Fig S2C). When sampling SNPs distant of at least 50 kb in this region to have
the same SNP density all over the chromosome, we also detected the high-LD block, which
indicates that the SNP density did not generated biases in the LD pattern (Fig S2D).
215
We found no other regions of the genomes with $r^2>0.9$ between SNPs across such a large
genomic distance in any population. On scaffold_6 in the CL1 cluster, two smaller SNP
blocks had a $r^2>0.9$ between each other despite being distant (Fig S2C; between SNPs blocks
at 3-20 kb and 2.151-2.278 kb), which is likely due to the major intra-scaffold translocation
described in this region between the EP155 and ESM15 strains (Demené et al. 2022). The
maximum size of the blocks with $r^2>0.9$ on other scaffolds ranged from 135 to 540 kb in the

CL1 cluster, as previously described (Demené et al. 2019). The average distance between
SNPs in high LD (r^2>0.9) across the genome was higher on the scaffold_2 than on other
scaffolds (in the CL1 cluster, pairwise Wilcoxon test p-value < 2e-16 with Bonferroni
correction). Furthermore, the proportion of SNPs in high LD (r^2>0.9) across the genome was
also much higher on the scaffold_2 than on other scaffolds due to the high LD in the MAT-
proximal region (Fig S2D, Table S2). Therefore, the large (1Mb) and localized block with
maximal LD values proximal to the mating-type locus both in the Swiss populations and the
CL1 genetic cluster stands out as exceptional in the genome and indicates full recombination
cessation, which supports previous inferences from progeny segregation analyses (Kubisiak
and Milgroom 2006b). Indeed, even low rates of recombination homogenize alleles and
prevents LD building (Dufresnes et al. 2015). The MAT-proximal non-recombining region
was actually larger (1.23 Mb), but with likely rare recombination events at one precise locus
near its edge, lowering LD between its two fully non-recombining parts.

Consistent with a lack of recombination, the MAT-proximal region formed two genetically
highly differentiated haplotypes in these invasive Swiss and CL1 populations (Fig 2A; Fig S3
A(1)-C(1)), as shown by the two clusters on the principal component analysis (PCA) using
only SNPs located in the MAT-proximal region, while no structure was detected in the rest of
the genome (Fig 2B-C; Fig S3A(2)-C(2)). The neighbor-net networks further supported the
existence of two differentiated haplotypes in the MAT-Proximal region, contrasting with an
otherwise recombining structure genome-wide (Fig 2D-E; Fig S3B-D). In the Swiss 2019
population, a few reticulations between haplotypes were found and three strains (LU3,
Nov10, Nov4) appeared to have an intermediate sequence in the MAT-Proximal region
between the two haplotypes (Fig S3D). These intermediates were rare and may correspond to
incomplete lineage sorting, gene conversion, infrequent recombination between haplotypes
and/or may represent a third haplotype.

248
In agreement with the view of a partial linkage with the mating-type locus, the two genetic
clusters in the MAT-proximal region appeared strongly associated with mating types,
although the association did not reach 100% (Figs 2A and D; Fig S3, Table 1). Indeed, the
distribution of mating types between the two PCA clusters strongly deviated from
expectations under random association (Table 1; chi-squared test: χ^2 = 33.3; p-value = 5.876e-
08, with r^2 = 0.38 in CL1). We hereafter call the two differentiated haplotypes MAT-Prox1

and MAT-Prox2, referring to the mating-type association. The EP155 genome used as reference for SNP calling above carried the MAT-Prox1 haplotype.

Inversion in the non-recombining MAT-proximal region

To investigate whether rearrangements between haplotypes were present in the MAT-proximal region, we sequenced the genome, using PacBio HiFi, of two strains with alternative MAT-proximal haplotypes: M1400 (MAT-2; MAT-Prox2) and M6697 (MAT-1; MAT-Prox1), originating from the Swiss population in Gnosca and belonging to the CL1 cluster, i.e. European invasive populations introduced from North America (Stauber et al. 2021; Stauber et al. 2022). We built high-quality genome assemblies: statistics of the assemblies were in the same range as those of the genome assemblies of the ESM15 and EP155 strains previously sequenced (Table S3; Demené et al 2022; Crouch et al 2020). The mating-type chromosome could be assembled as a single contig for the first time, in the M1400 genome, likely corresponding to a full chromosome, and was assembled into two contigs in the M6697 genome. We therefore used the M1400 genome as a reference for subsequent analyses. To identify the location of the MAT-proximal region in the M1400 genome, we computed LD by read mapping in the 1990 Swiss populations. We found the large block (> 1 Mb) of high LD (r^2>0.9) between 7,285,137 and 8,828,934 bp (red arrow; Fig. S4). The MAT-proximal region was also located 20 kb away from the mating-type locus (located at 7.265 Mb) and was 1.540 Mb long using the M1400 genome as reference. Consistent with the results using the EP155 genome as reference, we found that the high-LD region was divided into two higher-LD blocks near the edge at 7.392 Mb (orange arrows in Fig. S4).

The two new high-quality genome assemblies from the invasive Swiss population (introduced from North America) revealed an inversion between the two haplotypes in the MAT-proximal region (Fig 3). The two newly sequenced PacBio genomes were indeed collinear except for the mating-type chromosome (Fig S5A-B-C), where we found a large inversion (>1 Mb) between the M1400 and M6697 genomes (Fig 3; Fig S5A-B-C). Breakpoints of the inversion were located at ca. 7.455 and 8.563 Mb of the tig00000001 contig in the M1400 genome and at ca. 3.380 and 4.185 Mb of the tig00000060 contig in the M16697 genome. The region affected by the inversion was ca. 300 kb smaller in the M6697 genome than in the M1400 genome, while the whole MAT-proximal region was ca. 100 kb smaller. The split into the two high-LD blocks in natural populations reported above was outside of the inversion, i.e. 70 kb from the inversion breakpoint (orange arrows in Fig S4). LD in natural populations was

higher within the inversion (median value of $r^2 = 1$) than in other regions of the MAT-
proximal region (median value of $r^2 = 0.64$; Wilcoxon rank sum test with continuity
correction; $W = 4.8714e+10$; p-value $< 2e-16$).

We looked for centromeres, as the non-recombining regions near the mating-type locus in
other ascomycetes, when they occur, either capture the centromere (Menkis et al. 2008; Sun et
al. 2017), or are associated to the occurrence of a single crossing-over between the centromere
and the mating-type locus (Grognet et al. 2014; Hartmann et al. 2021; Vittorelli et al. 2022).
The centromere in *C. parasitica* may be at 4.380-4.536 Mb on the mating-type chromosome
in the M1400 genome, as we detected here a peak in TE density and a drop in GC content
(Fig 4A-B). A dotplot of repeats in this region also presented a pattern typical of centromeres
(Fig S5D). The MAT-proximal region and the inversion thus did not include the putative
centromere and was instead located about 2.9 Mb away of the putative centromere.

Higher genetic differentiation and lower genetic diversity within the MAT-proximal region
than in recombining regions
When analyzing polymorphism within the invasive 1990 Swiss population using the M1400
genome as reference, we found much higher genetic differentiation between strains of the two
MAT-proximal haplotypes in this region than elsewhere along the genome (Fig 4C; Fig S6A;
Table S4), as expected for a non-recombining region in LD with the mating-type locus. For
example, the F_{ST} values between the two non-recombining haplotypes within the MAT-
proximal region in the 1990 Swiss population were nearly maximal (median $F_{ST}=0.93$ per 50
kb window), while they were near zero in the rest of the genome (median $F_{ST}=0.02$; Wilcoxon
test p-value $< 2e-16$ with Bonferroni correction; Fig 4C; Fig S6A). Such F_{ST} values close to 1
indicate a lack of shared polymorphism and therefore support the inference of recombination
suppression in the MAT-proximal region.

Within each haplotype, the genetic diversity at the MAT-proximal region was lower than in
the rest of the genome (Wilcoxon test p-value $< 2e-16$ for the MAT-Prox1 haplotype; p-value
$= 2.3e-07$ for the MAT-Prox2 haplotype; Fig 4E-F; Fig S6C; Table S4), as expected for a
region without recombination associated with the mating-type locus, as its effective
population size is half as in recombining regions. The diversity was especially low in the pool
of isolates with the MAT-Prox1 haplotype (see all MAT-Prox1 sequences clustering on a
single point on the PCA in Fig 2A and in the neighbor-net network in Fig 2D, in contrast to

the more scattered MAT-Prox2 sequences). Such a very low diversity may be due to a recent

selective sweep or to particularly strong bottleneck in this haplotype during the invasion.

We detected signals for long-term balancing selection in the MAT-proximal haplotypes.

Computation of the Tajima's D statistics within all strains suggested signatures of balancing

selection in the MAT-proximal region as expected in a region without recombination

associated with the mating-type locus maintained at a frequency close to 1/2. Tajima's D

values in the MAT-proximal region pooling all sequences (median D=3.0 per 50 kb window)

were higher than in the rest of the genome (median D=-0.14; Wilcoxon test p-value < 2e-16

with Bonferroni correction; Fig 4G; Fig S6D). In contrast, Tajima's D was significantly lower

than the rest of the genome within each pool of haplotype, including in the MAT-Prox2

haplotype, suggestive of positive selection (Wilcoxon test p-value = 0.0037 for the MAT-

Prox1 haplotype; p-value = 0.0022 for the MAT-Prox2 haplotype with Bonferroni correction;

Fig 4H-I; Fig S6D; Table S4). We found no difference in population diversity statistics

between the recombining part of the mating-type contig and the other contigs (Table S4A).

The study of synonymous divergence (d_S) between M1400 and M6697 haplotypes suggests

that the recombination suppression is at least 1.5 million years old and that there is no pattern

of evolutionary strata within the MAT-proximal region, i.e. segments with different levels of

differentiation between haplotypes that would indicate stepwise expansion of recombination

suppression away from the mating-type locus. Per-gene synonymous divergence (d_S) is

typically used for detecting gradual expansion of recombination cessation and for estimating

its age, as it is considered a good proxy for the time since recombination suppression. Indeed,

when recombination is suppressed, mutations accumulate independently in the two non-

recombining haplotypes. We plotted the per-gene synonymous divergence (d_S) between

M1400 and M6697 along the M1400 genome, as its mating-type contig is likely being

assembled as a full chromosome (Fig S7A&E; Table S4). Consistent with recombination

suppression, we found significantly higher d_S values in the MAT-proximal region (computed

for 71 genes) than in the other regions of the mating-type contig (pairwise Wilcoxon test p-

value <2e-16 with Bonferroni correction) and other contigs (pairwise Wilcoxon test p-value

<2e-16 with Bonferroni correction). We found no significant differences between the other

regions of the mating-type contig and other contigs (pairwise Wilcoxon test p-value = 0.23

with Bonferroni correction). The d_S pattern displayed no indication of gradual expansion of

recombination cessation within the MAT-proximal region, as there was no stair-like pattern.

Note that the per-gene d_S could not be computed for the genes with a copy not predicted in
one of the haplotypes and therefore mostly in the oldest, most degenerated regions (see
below). Using synonymous substitution rate estimates in fungi (Kasuga et al. 2002; Taylor
and Berbee 2006) and the mean d_S values across genes of the MAT-proximal region, we
estimated the age of the divergence between the two haplotypes to be at least 1.5 Million
years when considering the mean d_S values of all genes in the MAT-proximal region (d_S
mean= 0.0495) (Table S4).

Enrichment in transposable elements in the non-recombining region and inversion
breakpoints
When studying the M1400 and M6697 high-qualities assemblies, we detected an enrichment
in transposable elements (TEs) in the MAT-proximal region in the two haplotypes compared
to the rest of the genome. The percentage of bp occupied by TEs (TE load) was higher than
50% in the MAT-proximal region, while it was only 9% on average in other regions. TE load
in the MAT proximal region was higher in the M6697 haplotype (76%, MAT-1; MAT-Prox1)
than the M1400 haplotype (68% MAT-2; MAT-Prox2). Class I retrotransposons with Gypsy
(LTR-Ty3) and LARD elements were the most abundant TEs in all genomic regions
(autosomes, MAT-Proximal region and the rest of the mating-type contig), representing >70%
of TE annotations (Fig 5A-B), as shown previously (Demené et al. 2022). The TE load
however varied significantly among TE families and genomic regions (ANOVA; Table S5).
In M6697 for example, TIR elements appeared less abundant in the non-recombining than
recombining regions while LTR-Ty3 elements were more frequent in the MAT-proximal
region (Fig 5B). In M1400, LARD elements were more frequent in the MAT-proximal region
than in recombining regions (Fig 5B).

The presence of transposable elements at the two inversion breakpoints in the two genomes
(Fig 5A) suggests that the inversion may have occurred via non-homologous recombination
mediated by these elements. By dating retrotransposon insertions using between-copy
divergence and within-copy LTR divergence in both M1400 and M6697 genomes (Fig 5C
and D), we found that TEs within the inversion and around the inversion breakpoints were
older than TEs farther from the breakpoints. This is expected in non-recombining regions as
selection is there less efficient to purge TE insertions so that they remain there longer
(Duhamel et al. 2023). The date estimates of LTR element insertions (Fig 5D) further
indicated that the TE accumulation in the MAT-proximal region was as old as a few million

years, confirming the age estimated based on sequence divergence between MAT-proximal haplotypes.

Gene functions, pseudogenization and degeneration in the non-recombining region

Recombination suppression could allow linking genes with mating-type antagonistic effects to the mating-type locus, or link together co-adapted alleles, for example for parasitizing different host genotypes. Out of 127 genes functionally annotated in the MAT-proximal region of the EP155 genome, 50 genes were supported by RNAseq data acquired *in vitro* for this strain (Chun et al. 2020), suggesting that the region contain functionally active genes. Among these 50 genes, nine were upregulated during barrage allorecognition of the EP155 strain with a compatible strain (Belov et al. 2021) and five were upregulated during infection of the EP155 strain with the hypovirus CHV1 (Chun et al. 2020), suggesting that the MAT-Proximal region may have a role in host-parasite interactions and vegetative incompatibility. We found no gene function known to be involved in mating or mating compatibility, nor in pathogenicity, in the MAT-proximal region (Table S6). We nevertheless identified genes potentially coding for proteins with kinase domain or proteins with homeodomains (Table S6), that are often involved in the regulation of developmental pathways.

By comparing gene predictions in the M6697 and M1400 newly assembled genomes, we identified a higher prevalence of gene disruptions (i.e. genes predicted in a single of the two haplotypes) in the MAT-proximal region than elsewhere in the genomes. Out of 175 groups of single-copy orthologous genes in the MAT-proximal region, 80 (46%) were only present in M1400 and 24 (14%) only present in M6697. In other genomic regions, only 3% of genes were only present in one genome. This suggests that gene disruption occurred in the non-recombining region, in higher proportions than elsewhere in the genome (Fisher's exact test p-value < 2.2e-16), and especially in the M6697 genome (MAT-1; MAT-Prox1). The enrichment in TEs and in gene disruptions in the MAT-Proximal region explains the lower gene density in this region (Fig 4) and supports the inference of an old full recombination suppression. The occurrence of high TE load and gene degeneration in both MAT-Proximal haplotypes further confirms the lack of homozygotes in this region, in which recombination could purge deleterious mutations, or indicates that recombination suppression occurs even in homozygotes, although it remains to be investigated.

We found overall higher non-synonymous substitution rate compared to the baseline substitution rates (d_N/d_S values) between the M1400 and M6697 genomes in the MAT-proximal region (median per gene $d_N/d_S = 0.80$) than in other contigs ($d_N/d_S = 0.43$; pairwise Wilcoxon test p-value = 1.7e-07 with Bonferroni correction; Fig S7A; Table S4). This is consistent with relaxed purifying selection due to recombination suppression in the MAT-proximal region. In addition, 13 genes displayed $d_N/d_S > 1$, suggesting that they may evolve under positive selection.

Recombination suppression in the MAT-Proximal region in other populations assessed from high-quality genome assemblies

We analyzed five additional high-quality genome assemblies for strains from the Asian native range (CL2 and CL4 clusters) and the invasion range (including two North American strains belonging to the CL1 cluster and one European strain introduced directly from Asia and belonging to the CL2 cluster), three of which were generated for the present study (Table S3). Most of the high-quality genomes displayed the same MAT-Proximal chromosomal arrangement as M1400 (MAT-Prox2, Swiss population and belonging to the CL1 cluster), suggesting that it is the ancestral state (Fig S8A-D), although the incomplete assemblies may prevent detecting other rearrangements. The inversion detected in M6697 (MAT-Prox1, Swiss population and belonging to CL1 cluster) was also present in the EP155 strain (MAT-Prox1, North America and belonging to CL1 cluster, Crouch et al, 2020; Fig S8D), indicating that this inversion was already segregating in early invasive populations in North America. In other genomic regions outside of the MAT-Proximal region, all analyzed high-quality genomes were colinear to the ESM15 reference genome, except the highly rearranged EP155 genome, as previously reported (Demené et al. 2022).

The MAT-Proximal region in all the high-quality genomes, from Asian native strains and invasive strains, displayed signs of degeneration, with a high TE load and gene disruptions, as seen from lower gene densities and missing orthologs (Fig6; Fig S8), confirming the ancient age of recombination suppression and its occurrence in the native range. Breaks of synteny between genome assemblies were associated with presence or absence of transposable elements and genes (Fig S8). Study of d_S and d_N/d_S between the MRC10 strain, belonging to the European clonal lineages of direct Asian origin (CL2 cluster), the M6697, M1400 and the two Asian native strains further confirmed recombination suppression in the native range and in invasive strains from different origins (North America or directly from Asia). The d_S and

d_N/d_S values were indeed significantly higher in the MAT-proximal region than other
genomic regions (Fig S7 B-E). Mean d_S values in the MAT-proximal region were in the same
range as d_S values between the M6697 and M1400 genomes, suggesting a similar date of
recombination suppression and, therefore, that recombination suppression was ancestral. The
MAT-proximal region appeared smaller in the two Asian strains XIM9508 and ESM15 (<1
Mb) than in invasive strains (Table S3), but this may be due to the incomplete genome
assemblies.

Presence of the two MAT-Proximal haplotypes in other populations
By analyzing available additional Illumina genomes in the native range and in invasive
populations, with less dense sampling within populations, we recovered the two haplotypes in
the MAT-proximal region, i.e., high differentiation between two clusters in the MAT-
Proximal region, contrasting with a lack of genetic subdivision in other genomic regions or a
completely different genetic subdivision, corresponding to previously described population
genetic structures (Figs 7 and FigS9). In particular, the neighbor-net network with multiple
strains sequenced previously from the CL2 and CL3 clusters (from the introduced range and
the native range, South Korean and Japan) and from the CL4 cluster (from the native range,
China;) reinforces the view of balancing selection on the MAT-Proximal haplotypes.
Sequences from the CL1 and CL2 clusters were indeed intermingled in the MAT-proximal
region (Fig 7A), while the structure in the rest of the genome instead corresponded to the
different population clusters, i.e., CL1, CL2, CL3 and CL4 (Fig 7B).

However, the two haplotypes were less differentiated in the native range and in invasive
populations originating directly from Asia than in invasive populations from the first invasion
wave via North America. For example, branch lengths on the neighbor-net indicate a lower
differentiation between haplotypes within CL2 (Fig 7A) than within the European CL1 or
Swiss invasive populations (Fig 2D; Fig S3). Some reticulations were present between MAT-
proximal haplotypes in CL2, which may again correspond to incomplete lineage sorting, gene
conversion or rare recombination events.

We found no association of the MAT-Proximal haplotypes with mating types at the world
scale (Figs 7 and S9), which may be due to the lack of extensive sampling within local
populations. As the MAT-Proximal region is only partially linked to the mating-type locus,
founder effects may yield different associations between MAT-Proximal haplotypes and

mating types in different populations. Only in the S12 European invasive clonal lineage (only

MAT-1), we found a single MAT-Proximal haplotype (MAT-Prox1; Fig S9B). In two

invasive European lineages of American origin with a predominantly clonal structure (re092

and re103), the strains with a mating type different from their clone-mates due to localized

introgression from other clonal lineages (Demené et al. 2019) also had their MAT-Proximal

region introgressed (Fig S9C); this finding reinforces the view of the strong linkage between

the mating-type locus and the MAT-Proximal region, and of an advantage of maintaining an

association between MAT-proximal haplotypes and mating types within populations. The

three 2019 Swiss strains that exhibited an intermediate sequence in the MAT-Proximal region

clustered with strains from Europe with a direct Asian origin (Fig S9C). This may represent a

third MAT-Proximal haplotype, associated with the second invasion in Europe, directly from

Asia.

No evidence of introgression in the MAT-proximal region from other species

The stronger differentiation in the MAT-Proximal region in the Swiss and CL1 invasive

populations than in other populations could also be due to an introgression event from a

closely related species; however, the small genetic distance and the reticulations observed in

the neighbor-net network between the M6697 Swiss reference genome and the Asian

genomes M8444 and XA19 rather indicate that the MAT-Prox1 haplotype in Switzerland

originated from Asian *C. parasitica* populations, likely a population different from those at

the origin of the CL2 invasive cluster (Fig 7).

Gene genealogies also provided no evidence of introgression in the MAT-proximal region

from closely related *Cryphonectria* species. We performed a gene orthology analysis

including three genomes of the closely related species *C. japonica*, C. *carpinicola*, and *C.*

naterciae that were previously sequenced with short read technologies (Stauber et al. 2020).

We retrieved only nine genes from the MAT proximal region that were shared between the

M6697 and M1400 genomes that had orthologs in one of the related species. This may be due

to low quality assemblies in the outgroups (Stauber et al. 2020) and/or a region without

recombination also in the outgroups with gene losses and many repeats. We found no gene

with a topology consistent with introgression from another species into the MAT-Proximal

region, i.e. with a MAT-Prox1 or MAT-Prox2 haplotype that would branch with an outgroup

allele rather than with the alternative haplotype of *C. parasitica*. The high load of TEs and

gene disruptions in the two MAT-Proximal haplotypes does not fit with the introgression

hypothesis either, unless the introgression occurred from a species with recombination
suppression in the MAT-Proximal region.
528
Discussion
A large and million-year-old non-recombining region partially linked to the mating-type
***locus in* Cryphonectria parasitica**
We found strong evidence for the complete cessation of recombination in a region of more
than 1 Mb (1.2 Mb to 1.5 Mb depending on the haplotype), strongly but not completely linked
to the mating-type locus in *C. parasitica*. Indeed, we detected maximal levels of linkage
disequilibrium in otherwise recombining populations and the existence of two highly
differentiated haplotypes, without shared polymorphism, while even low rates of
recombination can homogenize alleles and prevent LD build-up (Dufresnes et al. 2015). The
full cessation of recombination was further supported by the existence of a previously
unknown, large rearrangement (an inversion) in the center of the non-recombining region in
invasive strains. The higher TE load in this region constitutes additional evidence for full
recombination cessation, as the lack of recombination renders selection against TE
accumulation less efficient, in particular due to Muller's ratchet: two chromosomes with
different TE insertions cannot recombine to produce a chromosome free of TE insertions. TE
takes hundreds of thousands of years to accumulate in non-recombining regions (Duhamel et
al. 2023), so that the high TE load in the MAT-Proximal region in *C. parasitica* indicates old
recombination cessation. The particularly high frequency of missing genes and non-
synonymous substitutions in the MAT-Proximal region worldwide further indicates
degeneration and ancient recombination suppression. The estimates for the age of
recombination suppression based on the divergence between haplotypes and on TE insertions
were in fact at least a few million years. However, in contrast to other known non-
recombining regions on sex or mating-type chromosomes (Bergero and Charlesworth 2009;
Hartmann et al. 2021), the region without recombination was not completely linked to the
mating-type locus. Indeed, the occurrence of rare recombination events between the non-
recombining MAT-Proximal region and the mating-type locus are supported by previous
segregation analyses (Kubisiak and Milgroom 2006b) and the lack of association found here
between the MAT-Proximal haplotypes and the mating types at the worldwide scale, as well
as their incomplete association within populations.
558
Large inversion in the MAT-proximal non-recombining region

We found a large inversion between the two highly differentiated MAT-proximal haplotypes
in invasive populations originating from North America (CL1 genetic cluster, Swiss 1990 and
2019 populations and US populations). The chromosomal arrangement of the M1400 MAT-
Prox2 haplotype is likely the ancestral state as it is the most frequent worldwide, although a
more complete sampling in the native range is required to obtain definitive conclusion. The
inversion would therefore be the M6697 MAT-Prox1 haplotype and may contribute to
recombination suppression. The non-recombining region was defined based on LD pattern
and was larger than the inversion. This may indicate that additional proximal mechanisms
suppress recombination. The inversion may even be a consequence rather than a cause of
recombination cessation, as previously found in other fungal mating-type chromosomes
(Grognet et al. 2014; Sun et al. 2017; Branco et al. 2018; Vittorelli et al. 2022). Other
proximal causes of recombination cessation may for example be genetic recombination
modifiers, transposable elements, epigenetic marks, histone modifications or lack of
recruitment of proteins responsible for double-strand breaks (Maloisel and Rossignol 1998;
Boideau et al. 2022; Legrand et al. 2024). The lack of rearrangements at the edge of the
MAT-proximal region may allow rare recombination or gene conversion events, which may
explain the presence of two distinct LD blocks with lower LD between them and the lower
differentiation at the edges of the MAT-proximal region. Alternatively, the LD around the
inversion may be formed by less frequent recombination as recombination is often modified
around inversion breakpoints (Pegueroles et al. 2010; Stevison et al. 2011) and the actual size
of the non-recombining region may be smaller than suggested from the LD pattern.

The MAT-proximal non-recombining region is million-year-old and degenerated
Following recombination suppression, selection efficacy decreases. We detected footprints of
degeneration, with particularly high occurrence of gene disruptions and of TE insertions in the
MAT-proximal region, both in Europe, North America and in the native Asian range of the
fungus. A previous study (Demené et al. 2022) had already reported an enrichment of LTR-
Ty3 (*Gypsy*) element in the region near the mating-type locus in *C. parasitica*. We found
LTR-Ty3 elements at the inversion breakpoints and an enrichment in the MAT-Prox1
haplotype. Interestingly, the same TE family (LTR-Ty3) seems to have expanded in the non-
recombining regions in other fungi with recombination suppression on their mating-type
chromosomes (Duhamel et al. 2023), suggesting they may be more prone to accumulate in
non-recombining regions. Accumulations of transposable elements have also been detected in
other mating-type chromosomes, sex chromosomes and supergenes (Bachtrog 2013; Fontana

et al. 2020; Carpentier et al. 2022; Duhamel et al. 2023); some other large supergenes do not
seem to exhibit TE load, likely due to recent introgression, gene conversion or double
crossovers (Harringmeyer and Hoekstra 2022; Huang et al. 2022; Matschiner et al. 2022; Hill
et al. 2023).

The transposable element accumulation in the MAT-proximal region and the enrichment in
gene disruptions support the estimated age of recombination suppression at a few million
years ago based on mean pairwise gene sequence divergence between haplotypes. The finding
of the two haplotypes and of the inversion in the two invaded continents, Europe and North
America, as well as the occurrence of degeneration footprints in the MAT-Proximal region in
the native Asian range, further support the inference of an ancient recombination suppression
maintained by balancing selection. Indeed, even rare recombination events allow purging
deleterious mutations and TE insertions. This is in agreement with previous genetic maps
reporting lack of recombination near the mating-type locus in crosses involving Japanese,
North American and Italian isolates (Kubisiak and Milgroom 2006a). The high TE load and
degeneration level in terms of gene disruption in strains from Asian and North American
origins indicate a worldwide recombination cessation, so likely ancestral to all these native
and invasive populations. The strong degeneration, and in both haplotypes, further indicates
that even crosses between strains carrying the same MAT-Proximal haplotype does not allow
recombination in this region, which may be due to recombination suppression in the MAT-
Proximal region regardless of the haplotype, i.e., not being due to the inversion.

The differentiation between the MAT-Proximal haplotypes appeared greater in the CL1
cluster and Swiss populations (from the first invasion event, from North America) than in the
other clusters, which may be due to a founder effect or the lack of sampling in the source
population in Asia. The MAT-Prox1 haplotype in the populations from the first invasion
event from North America shows an inversion, a very low diversity, and the strongest
footprints of degeneration. Neighbornet networks show some reticulations between and
within haplotypes in some populations (e.g. CL2 that include the invasive strains from Asia,
but also the Swiss 2019 populations), and a few, rare intermediates between the two MAT-
Proximal haplotypes, which is likely due to incomplete lineage sorting, or, alternatively, to
rare recombination or gene conversion events, without leading to haplotype homogenization,
likely due to lower fitness of intermediates. These findings point to a strong balancing
selection maintaining the two haplotypes differentiated and non-recombining, and perhaps

different selection regimes or successive selection events during the various invasion waves
of *C. parasitica.*

The higher differentiation between the MAT-Proximal haplotypes in the first invasion wave
in Europe could also have resulted from introgression from a closely related species. The
evolution of non-recombining regions by introgression has been reported for example in ants
and butterflies (Jay et al. 2018; Helleu et al. 2022; Stolle et al. 2022). However, we found
little evidence for an introgression: i) the two haplotypes are present in the CL2 cluster of
invasive European populations directly originating from Asia, ii) the MAT-Prox1 haplotype
in the CL1 cluster more differentiated from the MAT-Prox2 haplotype than in the CL2 cluster
is genetically close to an Asian haplotype, iii) footprints of recombination suppression are
present in the native range in terms of degeneration and particular genomic structure in the
MAT-Proximal region, and iv) we found no signatures of introgression from closely related
species in gene genealogies.

The selective forces potentially maintaining the two haplotypes in the MAT-proximal
region
The genes found missing in the non-recombining region in one or the other haplotype may
help maintaining the two haplotypes by selection due to the sheltering of combinations of
deleterious mutations in repulsion, i.e. thanks to a heterozygote advantage called pseudo-
overdominance (Abu-Awad and Waller 2023). Such pseudo-overdominance advantage may
only be a consequence of recombination suppression, and contribute to its maintenance, or
may even be its initial cause (Branco et al. 2017; Jay et al. 2021; Jay et al. 2022; Charlesworth
2023; Jay, D.L. Jeffries, Hartmann, et al. 2024). Indeed, this situation of a non-recombining
region partially linked to the mating-type locus and with two haplotypes maintained by
selection fits predictions from a recent theoretical model showing that recombination
suppression can be selected for near permanently heterozygous alleles for protecting
haplotypes with fewer deleterious mutations than average in the population, while sheltering
their few recessive deleterious alleles. Importantly, the model predicts that selection can also
occur for regions partially linked to heterozygous alleles (Jay et al. 2022). Similar pseudo-
overdominance effect due to the sheltering of deleterious mutations can lead to balancing
selection in the context of local adaptation (Jay, Aubier, et al. 2024). These models require the
existence of a substantial diploid or dikaryotic phase for sheltering recessive deleterious
alleles in a heterozygous state, although the mechanism also works with a substantial haploid

phase (Jay et al. 2022). In fact, although *C. parasitica* has a predominant haploid phase,
strains can be found as dikaryons heterozygous for the mating-type locus in some natural
populations, and not only as monokaryons (McGuire et al. 2004; McGuire et al. 2005;
Stauber et al. 2021; Stauber et al. 2022). The importance and frequency of heterokaryons in
C. parasitica in nature would deserve further investigations. It may be sufficient if recessive
deleterious alleles are sheltered in ca 20% of individuals for selecting recombination
suppression (Jay et al. 2022), and/or if the genes involved are preferentially expressed during
the diploid or dikaryotic phases. However, it is challenging to assess whether the degree of
linkage between the mating-type locus and the MAT-Proximal region and the frequency of
dikaryotic stages are sufficient to fit the model. This hypothesis of balancing selection, and in
particular the hypothesis of overdominance (i.e. heterozygote advantage), does not require the
same association between the MAT-Proximal haplotypes and mating types across all
populations worldwide, it is sufficient that the MAT-Proximal haplotypes are strongly
associated to mating types within populations. Further sampling will be required to test
whether the MAT-Proximal haplotypes are associated to mating types within local
populations in the CL3 and CL4 clusters and in the native Asian range.

It could indeed also be that genes evolve under overdominance in this region for other reasons
than deleterious mutations, so that a partial linkage to a permanently heterozygous locus is
beneficial, perhaps in relationship with the pathogenic lifestyle of the fungus. One could
imagine, for example, that heterozygosity could be advantageous at genes involved in
virulence against the tree host, and especially the novel hosts colonized in invasive ranges, or
in resistance against a parasitic virus known to negatively affect fitness in *C. parasitica* (Choi
and Nuss 1992; Brusini et al. 2017). As a matter of fact, multiple genes present in the MAT-
Proximal region are up-regulated under infection by the hypovirus CHV1 or during the
vegetative incompatibility reaction considered to play a role in the prevention of virus
transmission (Choi et al. 2012; Rigling and Prospero 2018). As an alternative to a
heterozygote advantage, the selection maintaining the recombination suppression and the two
haplotypes in the MAT-proximal region could be some kind of negative-frequency dependent
selection of beneficial allelic combinations, possibly linked to a trench-warfare-like arms race
with the host tree, the virus or the microbial community, where partial linkage to the mating-
type locus would help maintenance in balanced proportions and therefore would prevent allele
loss (Tellier et al. 2014; Jay, Aubier, et al. 2024). Such a role in host-pathogen interactions
would also be consistent with the selective sweep footprints detected in the MAT-proximal

region in a previous study (Stauber et al. 2021), if there is recurrent positive selection for
improving the efficiency of the pathogen weapons within each of one of the two MAT-
proximal haplotypes. The detection of the selective sweep may have, however, been impacted
by particular demographic effect during the invasion (Moinet et al. 2022). Balancing selection
and location adjacent to a non-recombining region was recently found for loci involved in
host resistance in the *Daphnia–Pasteuria* system (Fredericksen et al. 2023). The MAT-
Proximal region may in this case even include genes involved different traits under balancing
selection, and would then constitute a supergene (Schwander et al. 2014). The transposable
elements insertion polymorphism observed from the high quality assemblies may also be
adaptive (Casacuberta and González 2013; Orteu et al. 2024).

Another hypothesis for explaining such extension of recombination suppression beyond the
mating-type locus is antagonistic selection, i.e. linkage of alleles that would improve fitness
of a MAT-1 gamete while being deleterious in a MAT-2 gamete, or vice-versa. However, full
recombination suppression with the mating-type locus would be expected under such
antagonistic selection, as well as the same association between the MAT-Proximal alleles and
the mating-type in all populations, in contrast to our findings. In addition, we found no
particular predicted function in the MAT-proximal region that could be related to mating
compatibility and there are very little functions, if any, with possible antagonistic functions
between mating types in fungi (Bazzicalupo et al. 2019; Hartmann et al. 2021).

Conclusion
In conclusion, we provide strong evidence for the existence of a non-recombining region
partially linked to the mating-type locus and maintained by selection in the chestnut blight
fungus *C. parasitica*. The non-recombining region in *C. parasitica* displayed contrasting
differentiation levels between haplotypes and different chromosomal arrangements in native
and invasive populations. We found footprints of balancing selection in the MAT-proximal
region in the first introduction of the pathogen in Europe from North America and a
chromosomal inversion. The non-recombining region displayed lower differentiation between
haplotypes in Asia and in the second invasion wave directly from Asia. This non-recombining
region may thus underlie important adaptive traits and thereby provide important applications
for the control of a devastating tree pathogen. In addition, the high-quality genome assemblies
provided here, from the native and introduced ranges, will more generally be useful for
studies aiming at understanding the evolution of this invasive and damaging pathogen.

Material and Methods

Strains and genomic data

For population genomic analyses, we studied a collection of 386 monokaryotic *C. parasitica* strains sampled worldwide, from the native and introduced ranges of the pathogen, and sequenced previously with the short-read Illumina technology (strain information are presented in Table S1; (Demené et al. 2019; Stauber et al. 2021; Stauber et al. 2022)). Data were downloaded from NCBI Bioproject numbers PRJNA604575, PRJNA644891 and PRJNA706885. We focused our analyses first on European invasive strains originating from North America. We studied 88 strains belonging to the CL1 genetic cluster, in central and southeastern Europe (Stauber et al. 2021). We excluded the putative clonal genotypes previously identified (Stauber et al. 2021). We also studied 71 strains sampled in the 1990 and 62 strains sampled in 2019 in southern Switzerland (Stauber et al. 2022). Mating-type ratios close to ½ and population structure analyses of the CL1 genetic cluster and the Swiss 1990 population suggest frequent recombination in these populations (Stauber et al. 2021; Stauber et al. 2022). The mating-type ratio was 1/3 *MAT-1* - 2/3 *MAT-2* in the 2019 Swiss population and population structure analyses suggested regular sexual reproduction and recent population bottleneck. To study more broadly the presence of the two haplotypes in the MAT-proximal region, we analyzed monokaryotic strains belonging to the CL2, CL3, CL4 genetic clusters, as well as the S12 European invasive lineage (Stauber et al. 2021) and additional monokaryotic strains from the US and Europe (with an Asian or North American origin (Demené et al. 2019)). We excluded heterokaryotic strains, i.e. strains having both MAT-1 and MAT-2 alleles, as the phase was challenging to infer.

For mapping and SNP calling, we first used as reference the 43.9 Mb genome sequence EP155 v2.0 of the strain EP155 (*MAT-2,* North America, CL1 cluster) available at the Joint Genome Institute (http://jgi.doe.gov/)(Crouch et al. 2020). For comparative genomics, we used the published genome of the strain ESM15 (MAT-2, Japan, CL2 cluster; (Demené et al. 2022)) available at DDBJ/ENA/GenBank on the bioproject PRJNA700518 under the accession JAGDFO000000000. We additionally sequenced *de novo,* with long-read technologies, five strains from the native and introduced ranges of the pathogen. We sequenced with PacBio Hifi the genomes of the strains M1400 (MAT-2) and M6697 (MAT-1) sampled in Gnosca, southern Switzerland (Stauber et al. 2021). Mycelia were stored as

glycerol stocks at -80 C after strain isolation. To produce mycelium for DNA extraction, isolates were inoculated onto cellophane-covered potato dextrose agar plates (PDA, 39 g/L; BD Becton, Dickinson and Company, Franklin Lakes, USA) (Hoegger et al. 2000) and incubated for a minimum of 1 week at 24°C, at a 14 hr light and 10 hr darkness cycle. Mycelium and spores were harvested by scratching the mycelial mass off the cellophane, transferring it into 2 mL tubes and freeze-drying it for 24 hr (Stauber et al. 2021). DNA extraction was performed with the NucleoBond High Molecular Weight DNA kit from Macherey-Nagel, with the mechanical disruption of about 30 mg of lyophilized mycelium with two beads of 3 mm for 5 min at 30 Hz. Sequencing was outsourced to Novogene, the Netherlands. We additionally sequenced with Oxford Nanopore MinION technology the strains XIM9508 (China, CL4 cluster, MAT-1), MRC10 (South Western France introduced directly from Asia, CL2 cluster, MAT-2) and DUM-005 (USA, CL1 cluster, MAT-2). These isolates had been collected for previous studies (Milgroom et al. 1996; Liu et al. 2003; Dutech et al. 2012). The protocols for mycelium production, DNA extraction and sequencing for these three strains were the same as in (Demené et al. 2022).

Short-read data processing and SNP calling

We used SNP calling datasets of the genomes from monokaryotic strains of the CL1, CL2, CL3, CL4 genetic clusters, the S12 invasive lineage and the Swiss populations against the EP155 reference v2.0 genome obtained in (Stauber et al. 2021; Stauber et al. 2022). We performed mapping and raw SNP calling of short-read data using the M1400 new genome assembly as a reference as described in (Stauber et al. 2021; Stauber et al. 2022). Briefly, we trimmed reads with Trimmomatic v0.36 (Bolger et al. 2014) and aligned them with Bowtie 2 v2.3.5.1 (Langmead et al. 2009) and SAMtools v1.9 (Li et al. 2009) to the EP155 v2.0 genome. Raw SNP calling and filtration for quality were conducted with the genome analysis toolkit GATK v3.8 and v4.0.2.0 (McKenna et al. 2010). We used the filtration parameters described in (Stauber et al. 2021): QUAL>=100, MQRankSum (lower) >= -2.0, QD:_20.0, MQRankSum (upper) <=2.0, MQ:_20.0, BaseQRankSum (lower) >=-2.0, ReadPosRankSum (lower) >=-2.0, ReadPosRankSum (upper) <=2.0, BaseQRankSum (upper) <=2.0. We further removed SNPs overlapping with transposable elements predicted *de novo* in the EP155 v2.0 genome by (Stauber et al. 2021).

Population genomics analyses

For all population genomics analyses, we excluded SNPs with missing data in more than 10%
of the strains and kept only polymorphic strains with vcftools v0.1.16 (Danecek et al. 2011).
To study linkage disequilibrium, we further excluded rare variants (minor allele frequency
<0.1) with the vcftools (Danecek et al. 2011) option --maf 0.1. We computed LD with the --
hap-r2 option of vcftools (Danecek et al. 2011) for each scaffold and each population
separately. We used the --thin 50000 option of vcftools (Danecek et al. 2011) to sample SNPs
distant of at least 50 kb. We used the R package LDheatmap v1.0-6 (Shin et al. 2006) to plot
LD r^2 values among SNP pairs. To perform principal component analyses (PCAs), we first
used vcftools (Danecek et al. 2011) to convert VCF format files in Plink format. We then used
the Plink v1.90b5.3 (Purcell et al. 2007) –pca command to run PCA analysis. We used the R
package PopGenome v2.7.5 (Pfeifer et al. 2014) to compute nucleotide diversity, Tajima's D
values and the F_{ST} index in 50 kb window overlapping over 10 kbp. Windows containing
fewer than 5 SNPs were removed from the analysis. We used the R package ggplot2 (2_3.5.0)
to plot results. Pairwise Wilcoxon tests were performed in R with Bonferroni or false
discovery rate correction.
812
813 We built neighbornet networks using SplitsTree4 v 4.19.2 (Huson 1998). VCF file were
814 converted to nexus format using PGDSpider v1.0 tool (Lischer and Excoffier 2012). For the
815 study of other populations, we included only one strain of each haplotype of the invasive
816 European population but not all CL1 and Swiss strains for network readability; we
817 nevertheless checked that the results were the same with all CL1 and Swiss strains.
818

Long-read based assemblies
PacBio Hifi reads of strains M1400 and M6697 were both assembled using canu v1.8 (Koren
et al. 2017) program with a set genome size of 44 Mb. Multiple assembly pipelines were used
for the other strains. Oxford Nanopore MinION reads and Illumina reads of the genomes of
both XIM9508 and MRC10 strains were assembled using HybridSPAdes (Antipov et al.
2016). Assemblies were manually curated and scaffolds were cut when an evidence of a
chimeric connection was detected (i.e. mis-paired short reads) as previously described
(Demené et al. 2022). For DUM005, the assembly was generated by Ra with basic parameters
that uses Oxford Nanopore MinION reads and corrects the assembly with Illumina reads
(https://github.com/lbcb-sci/ra) as it outperformed the HybridSPAdes assembly. As the
HybridSpades assembly of the MRC10 strain suggested absence of collinearity with the
MAT-proximal region M1400 and M6697, we further checked the assembly of the MAT-

proximal region of MRC010 by generating a meta-assembly of this strain. We used the

assembler tool canu v2.2 with an estimated genome size of 42 Mb. We also used Flye v2.9.3-

b1797 (Kolmogorov et al. 2019) with an estimated genome size of 42 Mb, --nano-raw for

reads with an error rate below 20 and a coverage for initial disjointig assembly of 50. Then we

used ragtag (Alonge et al. 2019) to patch the canu assembly with the Flye assembly as query

in a first loop. In a second loop, the canu assembly was patched with the first loop ragtag

assembly. Finally, we polished this second output assembly of ragtag with short reads and the

consensus part of medaka v1.11.3 with no change in the parameters

(https://github.com/nanoporetech/medaka). Assemblies statistics were obtained with quast

v5.1 (Gurevich et al. 2013). We assessed the completeness of each assembly using the

Benchmarking of Universal Single-Copy Orthologue (BUSCO) tools with the Sordariomyceta

ortholog set (sordariomycetes_odb10, 2020-08-05, n = 3817 orthologous groups searched)

(Manni et al. 2021). Gene models were predicted with the Helixer v0.3.1 pipeline (Holst et al.

2023). We also run Helixer pipeline for ESM15 and EP155 strains for the gene orthology

analysis. Statistics of the obtained gene annotation was obtained with the Agat v1.0.0 tool

(Dainat et al. 2020). Transposable elements of all genomes were annotated using

Repeatmasker v4-0-7 (Smit et al. 2013) and the customized library built for *C. parasitica* in

(Demené et al. 2022) contained in the "Curated_TEs_database_ESM015_EP155.fa" file

(available on the "Portail Data INRAE: Chromosomal rearrangements but no change of genes

and transposable elements repertoires in an invasive forest-pathogenic fungus" at

https://doi.org/10.15454/UTIB8U). The class Gipsy invader was renamed LTR-Ty3. We

filtered out TE copies shorter than 100bp. We used the HybridSPAdes preliminary assembly

of MRC10 for gene model annotations. To study gene functions of the EP155 genome, we

used the gene annotation available at http://jgi.doe.gov/ (Crouch et al. 2020). To study support

from RNAseq data, we used RNAseq data from the EP155 strain cultivated *in vitro* available

at Genebank under Project ID number PRJNA588887 and accessions numbers

SRR10428542, SRR10428543, SRR10428544 (Chun et al. 2020). Raw reads were mapped

using STAR v2.7.10a (Dobin et al. 2013) with the settings. --alignIntronMax 1000 --

limitBAMsortRAM 1629219965--quantMode GeneCounts. We used the program

featureCounts (Liao et al. 2013) v2.0.6 with the options -p --countReadPairs -M - -B -O --

largestOverlap. We considered a gene to be supported for read count >10.

Estimation of retrotranposon insertion time and nucleotide divergence time
864

To get an estimation of the insertion date of the transposable elements present in the MAT-
proximal non-recombining region, we applied two complementary methods. We first used the
divergence between the LTR sequences in retrotransposons, as these LTR sequences at their
edges are identical at the time of TE insertion and then diverge with time. For this, we used a
de-novo prediction of LTR retrotransposons using LTRharvest GenomeTools 1.6.2
(Ellinghaus et al. 2008; Gremme et al. 2013) and LTR_Finder v1.07 (Xu and Wang 2007). To
prepare data for LTRharvest, we first created enhanced suffix for the M1400 and M6697
genome assemblies using the GenomeTools Suffixerator (-dna -suf -lcp -bwt -bck -mirrored
parameters). We ran LTRharvest using two pipelines, designed to identify LTR
retrotransposons with a minimum and maximum LTR length of 100 bp and 7000 bp
respectively and at least 85% identity between two LTR regions, with and without canonical
TGCA motifs, respectively: i) -minlenltr 100 -maxlenltr 7000 -mintsd 4 -maxtsd 6 -similar
85 -vic 10 -seed 20 -motif TGCA -motifmis 1; ii) -minlentltr 100, -maxlenltr 7000, -mintsd 4,
-maxtsd 6, -similar 85, -vic 10, -seed 20. Similarly, we ran LTR_Finder on the M1400 and
M6697 genome assemblies to retrotransposons with both TGCA and non-TGCA motifs and a
minimum and maximum LTR length of 100 bp and 7000 bp respectively and at least 85%
identity between two LTR regions (-D 15000, -d 1000, -l 100, -L 7000, -p 20, -C, -M 0.85).
Finally, we used LTR_retriever v2.8.5 (Ou and Jiang 2018) with default parameters to filter
out false positive LTR candidates identified by LTRharvest and LTR_Finder and get an
estimation of each element insertion date.
885
As a second method to estimate the insertion date of the transposable elements present in the
MAT-proximal non-recombining region, we used the set of curated consensus sequences from
(Demené et al. 2022) to annotate the inversion sequence or its surroundings. We first used
samtools faidx v1.9 (Li et al. 2009) to extract the sequence corresponding to inversion and
their 50 kb surroundings in M1400 and M6697 genome assemblies. We annotated the
sequences corresponding to the inversions and the concatenated 50 kb surroundings in both
isolates using RepeatMasker version 4.1.5 and rmblast as search engine (v2.10.0) with -no_is
-pa 20 -cutoff 250 -a parameters. Finally, we parsed the RepeatMasker .out file using the
helped script parseRM_merge_interrupted.pl and omitting Simple_repeat and
Low_complexity regions (https://github.com/4ureliek/Parsing-RepeatMasker-Outputs). We
then built a summary of the alignments using the RepeatMasker helper script
buildSummary.pl and calculated sequence divergence using the calcDivergenceFromAlign.pl
script to finally render results with the createRepeatLandscape.pl from the same helper suite.

Comparative genomics analyses

Genome synteny between long reads assemblies were studied using the nucmer v3.1 program (Marçais et al. 2018). Outputs were plotted with the R programs ggplot2 (Wickham 2009), genoPlotR (Guy et al. 2010) and RIdeogram (Hao et al. 2020). The dotplot in the putative centromere region of the M1400 mating-type contig was performed using the online megablast alignment tool available at https://blast.ncbi.nlm.nih.gov/(last accessed 13[th] May 2024).

To study gene disruption, synonymous and non-synonymous divergence (d_S and d_N) and testing introgression in the MAT-proximal region, we build orthology relationships for genes of the genome assemblies of *C. parasitica* strains M1400, M6697, XIM9508, MRC10, DUM-005, ESM15 and EP155 and included three genomes of the closely related species *Cryphonectria japonica* (IF-6), *Cryphonectria carpinicola* (CS3), and *Cryphonectria naterciae* (M3664). Genome, gene annotation and species tree of these closely related species were previously published by (Stauber et al. 2020). Genome data were retrieved from NCBI bioproject number PRJNA644891 and accession IDs JACWRX000000000 for IF-6, JACWRQ000000000 for CS3 and JACWST000000000 for M3664. We run OrthoFinder v2.3.7 (Emms and Kelly 2019) analysis on protein sequences. We used the translatorX v1.1 program (Abascal et al. 2010) with default parameters that use a codon-based approach to align orthologous gene coding sequences. To compute d_S and d_N vaues of one-to-one orthologs between pairs of genome assemblies, we use the yn00 program of PAML (Yang 2007). Estimation of divergence time between haplotypes was performed using computed gene d_S values and the formula Tgenerations $= dS/2\mu$. We used previous estimates of substitution rates in fungi (Kasuga et al. 2002; Taylor and Berbee 2006) and considered that *C. parasitica* undergoes one generation a year (Guerin et al. 2001). We build gene coding sequences trees with the outgroup genomes in the MAT-proximal region using iqtree2 v2.2.2.6 (Minh et al. 2020) with 1000 bootstraps and used the Newick Utilities (https://github.com/tjunier/newick_utils) for displaying phylogenetics tree.

References

Abascal F, Zardoya R, Telford MJ. 2010. TranslatorX: multiple alignment of nucleotide sequences guided by amino acid translations. *Nucleic Acids Res*. 38:W7-13.

Abu-Awad D, Waller D. 2023. Conditions for maintaining and eroding pseudo-
overdominance and its contribution to inbreeding depression. *Peer Community J.*
[Internet] 3. Available from:
https://peercommunityjournal.org/articles/10.24072/pcjournal.224/

Alonge M, Soyk S, Ramakrishnan S, Wang X, Goodwin S, Sedlazeck FJ, Lippman ZB,
Schatz MC. 2019. RaGOO: fast and accurate reference-guided scaffolding of draft
genomes. *Genome Biol.* 20:224.

Anagnostakis SL. 1987. Chestnut blight: The classical problem of an introduced pathogen.
Mycologia [Internet]. Available from:
https://www.tandfonline.com/doi/abs/10.1080/00275514.1987.12025367

Antipov D, Korobeynikov A, McLean JS, Pevzner PA. 2016. hybridSPAdes: an algorithm for
hybrid assembly of short and long reads. *Bioinformatics* 32:1009–1015.

Bachtrog D. 2013. Y-chromosome evolution: emerging insights into processes of Y-
chromosome degeneration. *Nat. Rev. Genet.* 14:113–124.

Bazzicalupo AL, Carpentier F, Otto SP, Giraud T. 2019. Little evidence of antagonistic
selection in the evolutionary strata of fungal mating-type chromosomes
(*Microbotryum lychnidis-dioicae*). *G3 Genes Genomes Genet.* 9:1987–1998.

Belov AA, Witte TE, Overy DP, Smith ML. 2021. Transcriptome analysis implicates
secondary metabolite production, redox reactions, and programmed cell death during
allorecognition in Cryphonectria parasitica. *G3 GenesGenomesGenetics* 11:jkaa021.

Berdan EL, Blanckaert A, Butlin RK, Flatt T, Slotte T, Wielstra B. 2022. Mutation
accumulation opposes polymorphism: supergenes and the curious case of balanced
lethals. *Philos. Trans. R. Soc. B Biol. Sci.* 377:20210199.

Bergero R, Charlesworth D. 2009. The evolution of restricted recombination in sex
chromosomes. *Trends Ecol. Evol.* 24:94–102.

Boideau F, Richard G, Coriton O, Huteau V, Belser C, Deniot G, Eber F, Falentin C, Ferreira
de Carvalho J, Gilet M, et al. 2022. Epigenomic and structural events preclude
recombination in Brassica napus. *New Phytol.* 234:545–559.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* 30:2114–2120.

Branco S, Badouin H, Rodríguez de la Vega RC, Gouzy J, Carpentier F, Aguileta G, Siguenza
S, Brandenburg J-T, Coelho MA, Hood ME, et al. 2017. Evolutionary strata on young
mating-type chromosomes despite the lack of sexual antagonism. *Proc. Natl. Acad.*
Sci. 114:7067–7072.

Branco S, Carpentier F, Rodríguez de la Vega RC, Badouin H, Snirc A, Prieur SL, Coelho
MA, Vienne DM de, Hartmann FE, Begerow D, et al. 2018. Multiple convergent
supergene evolution events in mating-type chromosomes. *Nat. Commun.* 9:2000.

Brusini J, Wayne ML, Franc A, Robin C. 2017. The impact of parasitism on resource
allocation in a fungal host: the case of *Cryphonectria parasitica* and its mycovirus,
Cryphonectria Hypovirus 1. *Ecol. Evol.* 7:5967–5976.

Carpentier F, Rodríguez de la Vega RC, Jay P, Duhamel M, Shykoff JA, Perlin MH, Wallen
RM, Hood ME, Giraud T. 2022. Tempo of degeneration across independently evolved
nonrecombining regions. *Mol. Biol. Evol.* 39:msac060.

Casacuberta E, González J. 2013. The impact of transposable elements in environmental
adaptation. *Mol. Ecol.* 22:1503–1517.

Charlesworth B. 2023. The fitness consequences of genetic divergence between polymorphic
gene arrangements. *Genetics*:iyad218.

Charlesworth D. 2016. The status of supergenes in the 21st century: recombination
suppression in Batesian mimicry and sex chromosomes and other complex
adaptations. *Evol. Appl.* 9:74–90.

Charlesworth D, Vekemans X, Castric V, Glémin S. 2005. Plant self-incompatibility systems:
a molecular evolutionary perspective. *New Phytol.* 168:61–69.

Choi GH, Dawe AL, Churbanov A, Smith ML, Milgroom MG, Nuss DL. 2012. Molecular
characterization of vegetative incompatibility genes that restrict hypovirus
transmission in the chestnut blight fungus *Cryphonectria parasitica*. *Genetics*
190:113–127.

Choi GH, Nuss DL. 1992. Hypovirulence of chestnut blight fungus conferred by an infectious
viral cDNA. *Science* 257:800–803.

Chun J, Ko Y-H, Kim D-H. 2020. Transcriptome Analysis of Cryphonectria parasitica
Infected With Cryphonectria hypovirus 1 (CHV1) Reveals Distinct Genes Related to
Fungal Metabolites, Virulence, Antiviral RNA-Silencing, and Their Regulation.
Front. Microbiol. 11:1711.

Crouch JA, Dawe A, Aerts A, Barry K, Churchill ACL, Grimwood J, Hillman BI, Milgroom
MG, Pangilinan J, Smith M, et al. 2020. Genome sequence of the chestnut blight
fungus Cryphonectria parasitica EP155: A fundamental resource for an archetypical
invasive plant pathogen. *Phytopathology*® 110:1180–1188.

Dainat J, Hereñú D, Pucholt P. 2020. AGAT:Another Gff Analysis Toolkit to handle
annotations in any GTF. *GFF Format*.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter
G, Marth GT, Sherry ST, et al. 2011. The variant call format and VCFtools.
Bioinformatics 27:2156–2158.

Demené A, Laurent B, Cros-Arteil S, Boury C, Dutech C. 2022. Chromosomal
rearrangements with stable repertoires of genes and transposable elements in an
invasive forest-pathogenic fungus. *Peer Community J.* [Internet] 2. Available from:
https://peercommunityjournal.org/articles/10.24072/pcjournal.127/

Demené A, Legrand L, Gouzy J, Debuchy R, Saint-Jean G, Fabreguettes O, Dutech C. 2019. Whole-genome sequencing reveals recent and frequent genetic recombination between clonal lineages of Cryphonectria parasitica in western Europe. *Fungal Genet. Biol.* 130:122–133.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29:15–21.

Dufresnes C, Borzée A, Horn A, Stöck M, Ostini M, Sermier R, Wassef J, Litvinchuck SN, Kosch TA, Waldman B, et al. 2015. Sex-chromosome homomorphy in palearctic tree frogs results from both turnovers and X-Y recombination. *Mol. Biol. Evol.* 32:2328–2337.

Duhamel M, Hood ME, Vega RCR de la, Giraud T. 2023. Dynamics of transposable element accumulation in the non-recombining regions of mating-type chromosomes in anther-smut fungi. :2022.08.03.502670. Available from: https://www.biorxiv.org/content/10.1101/2022.08.03.502670v2

Dussert Y, Legrand L, Mazet ID, Couture C, Piron M-C, Serre R-F, Bouchez O, Mestre P, Toffolatti SL, Giraud T, et al. 2020. Identification of the first oomycete mating-type locus sequence in the grapevine downy mildew pathogen, *Plasmopara viticola*. *Curr. Biol.* 30:3897-3907.e4.

Dutech C, Barrès B, Bridier J, Robin C, Milgroom MG, Ravigné V. 2012. The chestnut blight fungus world tour: successive introduction events from diverse origins in an invasive plant fungal pathogen. *Mol. Ecol.* 21:3931–3946.

Dutech C, Fabreguettes O, Capdevielle X, Robin C. 2010. Multiple introductions of divergent genetic lineages in an invasive fungal pathogen, Cryphonectria parasitica, in France. *Heredity* 105:220–228.

Ellinghaus D, Kurtz S, Willhoeft U. 2008. LTRharvest, an efficient and flexible software for de novo detection of LTR retrotransposons. *BMC Bioinformatics* 9:18.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome Biol.* 20:238.

Fontana S, Chang N-C, Chang T, Lee C-C, Dang V-D, Wang J. 2020. The fire ant social supergene is characterized by extensive gene and transposable element copy number variation. *Mol. Ecol.* 29:105–120.

Fredericksen M, Fields PD, Pasquier LD, Ricci V, Ebert D. 2023. QTL study reveals candidate genes underlying host resistance in a Red Queen model system. *PLOS Genet.* 19:e1010570.

Furman BLS, Metzger DCH, Darolti I, Wright AE, Sandkam BA, Almeida P, Shu JJ, Mank JE. 2020. Sex chromosome evolution: So many exceptions to the rules. *Genome Biol. Evol.*

Gremme G, Steinbiss S, Kurtz S. 2013. GenomeTools: A Comprehensive Software Library for Efficient Processing of Structured Genome Annotations. *IEEE/ACM Trans. Comput. Biol. Bioinform.* 10:645–656.

Grognet P, Bidard F, Kuchly C, Tong LCH, Coppin E, Benkhali JA, Couloux A, Wincker P, Debuchy R, Silar P. 2014. Maintaining two mating types: structure of the mating type locus and its role in heterokaryosis in *Podospora anserina. Genetics* 197:421–432.

Guerin L, Froidefond G, Xu X-M. 2001. Seasonal patterns of dispersal of ascospores of Cryphonectria parasitica (chestnut blight). *Plant Pathol.* 50:717–724.

Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075.

Guy L, Roat Kultima J, Andersson SGE. 2010. genoPlotR: comparative gene and genome visualization in R. *Bioinformatics* 26:2334–2335.

Hao Z, Lv D, Ge Y, Shi J, Weijers D, Yu G, Chen J. 2020. RIdeogram: drawing SVG graphics to visualize and map genome-wide data on the idiograms. *PeerJ Comput. Sci.* 6:e251.

Harringmeyer OS, Hoekstra HE. 2022. Chromosomal inversion polymorphisms shape the genomic landscape of deer mice. *Nat. Ecol. Evol.* 6:1965–1979.

Hartmann FE, Duhamel M, Carpentier F, Hood ME, Foulongne-Oriol M, Silar P, Malagnac F, Grognet P, Giraud T. 2021. Recombination suppression and evolutionary strata around mating-type loci in fungi: documenting patterns and understanding evolutionary and mechanistic causes. *New Phytol.* 229:2470–2491.

Helleu Q, Roux C, Ross KG, Keller L. 2022. Radiation and hybridization underpin the spread of the fire ant social supergene. *Proc. Natl. Acad. Sci.* 119:e2201040119.

Hill J, Enbody ED, Bi H, Lamichhaney S, Lei W, Chen J, Wei C, Liu Y, Schwochow D, Younis S, et al. 2023. Low mutation load in a supergene underpinning alternative male mating strategies in ruff (*Calidris pugnax*). *Mol. Biol. Evol.* 40:msad224.

Hoegger PJ, Rigling D, Holdenrieder O, Heiniger U. 2000. Genetic structure of newly established populations of *Cryphonectria parasitica. Mycol. Res.* 104:1108–1116.

Holst F, Bolger A, Günther C, Maß J, Triesch S, Kindel F, Kiel N, Saadat N, Ebenhöh O, Usadel B, et al. 2023. Helixer–de novo prediction of primary Eukaryotic gene models combining deep learning and a hidden markov model. :2023.02.06.527280. Available from: https://www.biorxiv.org/content/10.1101/2023.02.06.527280v2

Huang K, Ostevik KL, Elphinstone C, Todesco M, Bercovich N, Owens GL, Rieseberg LH. 2022. Mutation load in sunflower inversions is negatively correlated with inversion heterozygosity.Takahashi A, editor. *Mol. Biol. Evol.* 39:msac101.

Huson DH. 1998. SplitsTree: analyzing and visualizing evolutionary data. *Bioinforma. Oxf. Engl.* 14:68–73.

Ironside JE. 2010. No amicable divorce? Challenging the notion that sexual antagonism
drives sex chromosome evolution. *BioEssays* 32:718–726.

Jay P, Aubier TG, Joron M. 2024. The interplay of local adaptation and gene flow may lead to
the formation of supergenes. *Mol. Ecol.*:e17297.

Jay P, Chouteau M, Whibley A, Bastide H, Parrinello H, Llaurens V, Joron M. 2021.
Mutation load at a mimicry supergene sheds new light on the evolution of inversion
polymorphisms. *Nat. Genet.* 53:288–293.

Jay P, Jeffries D, Hartmann FE, Véber A, Giraud T. 2024. Why do sex chromosomes
progressively lose recombination? *Trends Genet.* [Internet]. Available from:
https://www.sciencedirect.com/science/article/pii/S0168952524000672

Jay P, Jeffries DL, Hartmann FE, Véber A, Giraud T. 2024. Why do sex chromosomes
progressively lose recombination? *Trends in Genetics*:in press.

Jay P, Tezenas E, Véber A, Giraud T. 2022. Sheltering of deleterious mutations explains the
stepwise extension of recombination suppression on sex chromosomes and other
supergenes. *PLOS Biol.* 20:e3001698.

Jay P, Whibley A, Frézal L, Rodríguez de Cara MÁ, Nowell RW, Mallet J, Dasmahapatra
KK, Joron M. 2018. Supergene evolution triggered by the introgression of a
chromosomal inversion. *Curr. Biol.* 28:1839-1845.e3.

Kasuga T, White TJ, Taylor JW. 2002. Estimation of nucleotide substitution rates in
eurotiomycete fungi. *Mol. Biol. Evol.* 19:2318–2324.

Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, error-prone reads using
repeat graphs. *Nat. Biotechnol.* 37:540–546.

Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu: scalable
and accurate long-read assembly via adaptive k-mer weighting and repeat separation.
Genome Res. 27:722–736.

Kubisiak TL, Milgroom MG. 2006a. Markers linked to vegetative incompatibility (vic) genes
and a region of high heterogeneity and reduced recombination near the mating type
locus (MAT) in *Cryphonectria parasitica. Fungal Genet. Biol.* 43:453–463.

Kubisiak TL, Milgroom MG. 2006b. Markers linked to vegetative incompatibility (*vic*) genes
and a region of high heterogeneity and reduced recombination near the mating type
locus (*MAT*) in *Cryphonectria parasitica. Fungal Genet. Biol. FG B* 43:453–463.

Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. *Genome Biol.* 10:R25.

Legrand S, Saifudeen A, Bordelet H, Vernerey J, Guille A, Bignaud A, Thierry A, Acquaviva
L, Gaudin M, Sanchez A, et al. 2024. Absence of chromosome axis protein
recruitment prevents meiotic recombination chromosome-wide in the budding yeast
Lachancea kluyveri. *Proc. Natl. Acad. Sci.* 121:e2312820121.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin
R. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics*
25:2078–2079.

Liao Y, Smyth GK, Shi W. 2013. The Subread aligner: fast, accurate and scalable read
mapping by seed-and-vote. *Nucleic Acids Res.* 41:e108–e108.

Lischer HEL, Excoffier L. 2012. PGDSpider: an automated data conversion tool for
connecting population genetics and genomics programs. *Bioinformatics* 28:298–299.

Liu YC, Linder-Basso D, Hillman BI, Kaneko S, Milgroom MG. 2003. Evidence for
interspecies transmission of viruses in natural populations of filamentous fungi in the
genus Cryphonectria. *Mol. Ecol.* 12:1619–1628.

Lovat C-A, Donnelly DJ. 2019. Mechanisms and metabolomics of the host–pathogen
interactions between Chestnut (Castanea species) and Chestnut blight (Cryphonectria
parasitica). *For. Pathol.* 49:e12562.

Maloisel L, Rossignol JL. 1998. Suppression of crossing-over by DNA methylation in
Ascobolus. *Genes Dev.* 12:1381–1389.

Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021. BUSCO update: Novel
and streamlined workflows along with broader and deeper phylogenetic coverage for
scoring of eukaryotic, prokaryotic, and viral genomes. *Mol. Biol. Evol.* 38:4647–4654.

Marçais G, Delcher AL, Phillippy AM, Coston R, Salzberg SL, Zimin A. 2018. MUMmer4:
A fast and versatile genome alignment system. *PLOS Comput. Biol.* 14:e1005944.

Matschiner M, Barth JMI, Tørresen OK, Star B, Baalsrud HT, Brieuc MSO, Pampoulie C,
Bradbury I, Jakobsen KS, Jentoft S. 2022. Supergene origin and maintenance in
Atlantic cod. *Nat. Ecol. Evol.* 6:469–481.

McGuire IC, Davis JE, Double ML, MacDonald WL, Rauscher JT, McCawley S, Milgroom
MG. 2005. Heterokaryon formation and parasexual recombination between
vegetatively incompatible lineages in a population of the chestnut blight fungus,
Cryphonectria parasitica. *Mol. Ecol.* 14:3657–3669.

McGuire IC, Marra RE, Milgroom MG. 2004. Mating-type heterokaryosis and selfing in
Cryphonectria parasitica. *Fungal Genet. Biol.* 41:521–533.

McGuire IC, Marra RE, Turgeon BG, Milgroom MG. 2001. Analysis of mating-Type genes
in the chestnut blight fungus, *Cryphonectria parasitica*. *Fungal Genet. Biol.* 34:131–
144.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K,
Altshuler D, Gabriel S, Daly M, et al. 2010. The Genome Analysis Toolkit: a
MapReduce framework for analyzing next-generation DNA sequencing data. *Genome
Res.* 20:1297–1303.

Menkis A, Jacobson DJ, Gustafsson T, Johannesson H. 2008. The mating-type chromosome
in the filamentous ascomycete *Neurospora tetrasperma* represents a model for early
evolution of sex chromosomes. *PLOS Genet.* 4:e1000030.

Milgroom MG, Wang K, Zhou Y, Lipari SE, Kaneko S. 1996. Intercontinental population structure of the chestnut blight fungus, *Cryphonectria parasitica*. *Mycologia* 88:179–190.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear R. 2020. IQ-TREE 2: New models and efficient methods for phylogenetic inference in the genomic era. *Mol. Biol. Evol.* 37:1530–1534.

Moinet A, Schlichta F, Peischl S, Excoffier L. 2022. Strong neutral sweeps occurring during a population contraction. *Genetics* 220:iyac021.

Orteu A, Kucka M, Gordon IJ, Ng'iru I, van der Heijden ESM, Talavera G, Warren IA, Collins S, ffrench-Constant RH, Martins DJ, et al. 2024. Transposable element insertions are associated with Batesian mimicry in the pantropical butterfly *Hypolimnas misippus*. *Mol. Biol. Evol.* 41:msae041.

Otto SP, Lenormand T. 2002. Resolving the paradox of sex and recombination. *Nat. Rev. Genet.* 3:252–261.

Ou S, Jiang N. 2018. LTR_retriever: A highly accurate and sensitive program for identification of long terminal repeat retrotransposons. *Plant Physiol.* 176:1410–1422.

Pegueroles C, Ordóñez V, Mestres F, Pascual M. 2010. Recombination and selection in the maintenance of the adaptive value of inversions. *J. Evol. Biol.* 23:2709–2717.

Pfeifer B, Wittelsbürger U, Ramos-Onsins SE, Lercher MJ. 2014. PopGenome: an efficient Swiss army knife for population genomic analyses in R. *Mol. Biol. Evol.* 31:1929–1936.

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P, de Bakker PIW, Daly MJ, et al. 2007. PLINK: A tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.* 81:559–575.

Rigling D, Prospero S. 2018. Cryphonectria parasitica, the causal agent of chestnut blight: invasion history, population biology and disease control. *Mol. Plant Pathol.* 19:7–20.

Schwander T, Libbrecht R, Keller L. 2014. Supergenes and complex phenotypes. *Curr. Biol.* 24:R288–R294.

Shin J-H, Blay S, McNeney B, Graham J, others. 2006. LDheatmap: an R function for graphical display of pairwise linkage disequilibria between single nucleotide polymorphisms. *J. Stat. Softw.* 16:1–10.

Smit AFA, Hubley R, Green P. 2013. RepeatMasker Open-4.0. Available at: http://www.repeatmasker.org Accessed: November 30, 2017.

Stauber L, Badet T, Feurtey A, Prospero S, Croll D. 2021. Emergence and diversification of a highly invasive chestnut pathogen lineage across southeastern Europe.Castric V, Weigel D, Laine A-L, Castric V, editors. *eLife* 10:e56279.

Stauber L, Croll D, Prospero S. 2022. Temporal changes in pathogen diversity in a perennial plant–pathogen–hyperparasite system. *Mol. Ecol.* 31:2073–2088.

Stauber L, Prospero S, Croll D. 2020. Comparative genomics analyses of lifestyle transitions
at the origin of an invasive fungal pathogen in the genus *Cryphonectria*. *mSphere*
5:10.1128/msphere.00737-20.

Stevison LS, Hoehn KB, Noor MAF. 2011. Effects of Inversions on Within- and Between-
Species Recombination and Divergence. *Genome Biol. Evol.* 3:830–841.

Stolle E, Pracana R, López-Osorio F, Priebe MK, Hernández GL, Castillo-Carrillo C, Arias
MC, Paris CI, Bollazzi M, Priyam A, et al. 2022. Recurring adaptive introgression of a
supergene variant that determines social organization. *Nat. Commun.* 13:1180.

Sun Y, Svedberg J, Hiltunen M, Corcoran P, Johannesson H. 2017. Large-scale suppression
of recombination predates genomic rearrangements in *Neurospora tetrasperma*. *Nat.*
Commun. 8:1–8.

Taylor JW, Berbee ML. 2006. Dating divergences in the Fungal Tree of Life: review and new
analyses. *Mycologia* 98:838–849.

Tellier A, Moreno-Gámez S, Stephan W. 2014. Speed of adaptation and genomic footprints of
host–parasite coevolution under arms race and trench warfare dynamics. *Evolution*
68:2211–2224.

Umen JG. 2011. Evolution of sex and mating loci: An expanded view from Volvocine algae.
Curr. Opin. Microbiol. 14:634–641.

Vittorelli N, Hartmann, Giraud T. 2022. Stepwise recombination suppression around the
mating-type locus in the fungus *Schizothecium tetrasporum* (Ascomycota,
Sordariales).

Wickham H. 2009. ggplot2: elegant graphics for data analysis. Springer Science & Business
Media

Xu Z, Wang H. 2007. LTR_FINDER: an efficient tool for the prediction of full-length LTR
retrotransposons. *Nucleic Acids Res.* 35:W265–W268.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.*
24:1586–1591.

1224

1225

Acknowledgments

1227 This work was supported by the European Research Council (ERC) EvolSexChrom (832352) grant

1228 and a Fondation Louis D grant from the Institut de France to TG, and by the ANR PIA grant # ANR-

1229 20-IDEES-0002 to F.E.H. We acknowledge the GenOuest bioinformatics core facility

1230 (https://www.genouest.org) for providing the computing infrastructure, (GPU) for genome

1231 annotation". The authors declare that they have no competing interests.

1232

1233 TG and FEH conceptualized the study and acquired funding. TG supervised the study. AL, AS, CD,
1234 AD, SP, DC, LS and TB contributed to data and their availability. AL and AS performed *in vitro*
1235 culture and DNA extraction. FEH, RRDLV, QR, JPV, LS and TB analyzed genomes. FEH and TG
1236 wrote the original draft. All authors edited the manuscript.

1237

1238 The raw data and the new assemblies produced in this study will be published pending scientific
1239 review.

1240

1241 **Tables**

1242 **Table 1: Distribution of mating types (*MAT-1* and *MAT-2*) among non-recombining**
1243 **haplotypes of the MAT-proximal region in the European invasive *Cryphonectria***
1244 ***parasitica* CL1 genetic cluster and the Swiss populations.** MAT-Prox1 and MAT-Prox2
1245 haplotypes were defined based on the clusters of the principal component analysis.

1246

1247 **Figures**

1248

Figure 1: Linkage disequilibrium (LD) analysis along the contig carrying the mating-
type locus in a *Cryphonectria parasitica* European invasive population. LD heatmaps
using single nucleotide polymorphisms (SNPs; n=3815) located on the contig carrying the
mating-type locus (scaffold_2 of the EP155 genome) in the CL1 genetic cluster (European
invasive population introduced from North America); pairs of SNPs with high LD, i.e. r^2
>0.9, correspond to the red triangle. The mating-type locus location is shown with a green
triangle and the MAT-proximal region lacking recombination is shown with a red arrow. The
two high-LD blocks within the MAT-proximal region are shown with orange arrows. SNPs at
the limit of the MAT-proximal region and the two high LD blocks were manually highlighted
with red dotted lines and an orange line.

1259

Figure 2: Genetic structure using single nucleotide polymorphisms (SNPs) in the MAT-
proximal region lacking recombination and other genomic regions in a *Cryphonectria*
***parasitica* European invasive population. A-C.** Principal component analysis (PCA). Two
principal components are presented. Percentage of variance explained by each PC is indicated
into brackets. Strains are colored according to their mating type (*MAT-1* or *MAT-2*). **D-E.**
Neighbor-net network from a SplitsTree analysis. These analyses were performed in the CL1
genetic cluster (European invasive populations introduced from North America) based on: **A**

and D. SNPs (n=2,220) located within the MAT-proximal region along the contig carrying

the mating-type locus (scaffold_2 of the EP155 genome).; **B.** SNPs (n=1,595) located in other

regions along the contig carrying the mating-type locus; **C and E** SNPs (n=11,289) located on

other contigs of the EP155 genome. On panels A and D, the two clusters corresponding to the

MAT-Prox1 and MAT-Prox2 haplotypes are shown with red circles. The identified haplotype

of each strain is indicated in Table S1. The number of strains within the MAT-Prox1

haplotype is indicated on panel A by the letter n. The newly sequenced M1400 and M6697

strains are highlighted with a black rectangle

1275

1276 **Figure 3: Synteny and rearrangements between the two newly sequenced genomes of**

1277 **European invasive strains introduced from North America (M1400 and M6697) in the**

1278 **chromosome carrying the MAT-proximal region lacking recombination in**

1279 *Cryphonectria parasitica.* The mating-type locus is located with a green diamond. The MAT-

1280 proximal region defined from LD analyses is indicated with red arrows.

1281

Figure 4: Genetic diversity and divergence between non-recombining haplotypes of the

MAT-proximal region along the contig carrying the mating-type locus in a

Cryphonectria parasitica **European invasive population.** A-B. GC content (A) and

transposable element densities (B) along the contig carrying the mating-type locus

(tig_0000001) of the M1400 *C. parasitica* genome. C. Relative divergence (F_{ST}) between

strains of the MAT-Prox1 and MAT-Prox2 haplotypes: D-E. Nucleotide diversity within

pools of strains for each MAT-Proximal haplotype; F-G-H Tajima's D for all strains pooled

and within pools of strains for each MAT-Proximal haplotype; The MAT-proximal region

defined from LD analyses and the inversion between M1400 and M6697 genomes are

indicated with red and blue arrows respectively. The mating-type locus location is shown with

a green triangle. The location of the putative centromere is indicated by a yellow circle. All

population statistics were computed for the 1990 Swiss population along the mating-type

contig (tig_0000001) of the M1400 genome per 50-kb window overlapping over 10 kbp.

Windows containing fewer than 5 SNPs were removed from the analysis.

1296

1297 **Figure 5: Annotation of transposable elements (TEs) and estimates of their insertion**

1298 **time in the MAT-proximal region lacking recombination and other genomic regions in**

1299 *Cryphonectria parasitica.* A. Annotation of transposable elements (TEs) and gene density

1300 along the MAT-proximal region in M1400 and M6697 genomes. The figure only shows the

genomic region from 7 Mb on tig_0000001 of the M1400 genome and from 2.8 Mb on tig_0000060 of the M6697 genome. The mating-type locus is located with a green triangle. The two high-LD blocks within the MAT-proximal region are shown with orange arrows. Synteny for 10 kb segments with identity > 90 % is shown in red and inversion in blue. Transposons larger than 5 kb are shown in purple for the transposons annotated as LTR-Ty3 by and in green for other transposons. Orthologous genes shared between genomes are shown in black and unique to each each genome in grey. B. Relative percentage of each TE annotation (in %) in the non-recombining MAT-proximal region and other recombining regions of M6697 and M1400 genomes. C. Pairwise genetic distance between TE copies within the inversion and around (50 kb) in both M1400 and M6697 genomes. Distribution of the Kimura substitution levels computed using the consensus sequence for the TEs. D. Estimates of the age of intact copies of LTR within and around the inversion.

Figure 6: Transposable elements content and gene content variation in seven high quality genome assemblies from other _Cryphonectria parasitica_ populations from the native and the introduced range. A. TE load (percentage of base pairs occupied by transposable elements) in the seven high quality genome assemblies in the MAT proximal region and other recombining regions. **B.** Number of orthologous genes present in the seven high quality genome assemblies. Only single copy orthologous genes shared between M6697 and M1400 or present only in one this genome are shown.

Figure 7: Genetic structure using single nucleotide polymorphisms (SNPs) in the MAT-proximal region lacking recombination (A) and other genomic regions (B) in other resequenced _Cryphonectria parasitica_ populations from the native and the introduced range. Neighbor-net network from a SplitsTree analysis based on: **A** SNPs (n=4,120) located within the MAT-proximal region along the contig carrying the mating-type locus (scaffold_2 of the EP155 genome); **B** SNPs (n= 103,058) located on other contigs of the EP155 genome. Color of empty circles around strain ID indicate the genetic clusters the strains belong to (CL1, CL2, CL3, CL4; n= 33 strains**)**. Dotted circles indicate strains of the native range and plain circles indicate strains of the introduced range. Red and blue dots near strain ID indicate the mating type (_MAT-1_ or _MAT-2_, respectively).

1334   Table 1

1335

	Population	CL1 genetic cluster		Swiss 1990 population		Swiss 2019 population	
	Haplotype	MAT-Prox1	MAT-Prox2	MAT-Prox1	MAT-Prox2	MAT-Prox1	MAT-Prox2
Number of strains in each PCA cluster	total	36	52	20	51	8	54
	MAT-1	30	11	18	17	5	15
	MAT-2	6	41	2	34	3	39

1336

1337

1338

1339 Fig 1

1340

1341
1342

1343   Fig 2

1344

1345
1346

1347
1348

1349   Fig 3

1350
1351
1352

1353   Fig 4

1354

1355

1356

1357   Fig5

1358

1359

1360

1361   Fig 6

A

B

Orthologs present
and absent in M66

Orthologs present
and absent in M14

Orthologs present
M1400 and M6697

Number of present orthologs

* MAT-prox
** MAT-pro

1362
1363

1364   Fig 7

1365

1366

1367

1368

1369

1370