1

Million years old recombination suppression and balancing selection in a region linked to the mating-type locus in the invasive chestnut blight fungal pathogen *Cryphonectria parasitica*

Fanny E. Hartmann[1]*, Ricardo C. Rodriguez de la Vega[1], Quentin Rougemont[1], Arthur Demené[2], Thomas Badet[3], Daniel Croll[3], Amandine Labat[1], Simone Prospero[4], Alodie Snirc[1], Lea Stauber[5], Cyril Dutech[2], Tatiana Giraud[1]

[1] Ecologie Systematique et Evolution, CNRS, Université Paris-Saclay, AgroParisTech, 91198 Gif-sur-Yvette, France

[2] BIOGECO, INRAE, Univ. Bordeaux, 69 route d'Arcachon, Cestas F-33610, France.

[3] Laboratory of Evolutionary Genetics, Institute of Biology, University of Neuchâtel, Neuchâtel, Switzerland

[4] Swiss Federal Research Institute WSL, Birmensdorf, Switzerland

[5] Institute for Infectious Diseases, University of Bern, Switzerland

*Corresponding author: Fanny E. Hartmann

E-mail: fanny.hartmann@universite-paris-saclay.fr

Key words: recombination suppression, fungi, mating-type chromosome, inversion, deleterious allele sheltering, supergenes, evolutionary strata, sex chromosomes

Abstract (short version)

Recombination suppression often evolves around mating compatibility loci, generating supergenes. Using hundreds of available genomes and generating high-quality genome assemblies from the invasive chestnut blight fungus *Cryphonectria parasitica*, we show that a 1.2 Mb genomic region proximal to the mating-type locus lacks recombination worldwide, thus constituting a supergene, but only partially linked to the mating-type locus. In invasive populations, this MAT-proximal region displayed two highly differentiated haplotypes, strongly associated to mating types, but not completely. High-quality genome assemblies revealed an inversion in the MAT-proximal region, and footprints of degeneration worldwide, with gene disruptions, non-synonymous substitutions and transposable elements. The two haplotypes segregate in all continents and have diverged at least 1.5 million years ago. This study reveals a large supergene partially linked to a mating compatibility locus, with balancing selection maintaining differentiated haplotypes, possibly involving deleterious mutations and/or host adaptation in a devastating tree pathogen.

Teaser:

A new supergene discovered near the mating-type locus in the devastating tree pathogen responsible for the chestnut blight disease

Abstract (longer version)

Recombination suppression often evolves in sex chromosomes and around mating-type loci. In the invasive chestnut blight fungus *Cryphonectria parasitica* (Ascomycota), a genomic region was previously suggested to lack recombination and to be partially linked to the mating-type (MAT) locus based on the analysis of a few progenies. Using hundreds of available *C. parasitica* genomes and generating several new high-quality genome assemblies from the native and introduced range of the pathogen, we show that a ca. 1.2 Mb genomic region proximal to the mating-type locus lacks recombination worldwide, thus constituting a supergene. In invasive populations with extensive local genomic data and a recombining genome-wide structure, this MAT-proximal region displayed two highly differentiated haplotypes, that were strongly associated to mating types, but not completely, and with footprints of selective sweeps. High-quality assemblies of the two haplotypes in invasive and

native populations revealed an inversion in one of the two MAT-proximal haplotypes and
footprints of degeneration worldwide, the MAT-proximal region being enriched in gene
disruptions, non-synonymous substitutions and transposable elements in both haplotypes. The
divergence between the two haplotypes in the supergene was estimated to have occurred at
least 1.5 million years ago and two haplotypes segregate in all continents, including the native
range. Footprints of selective sweep, high differentiation between haplotypes, their
occurrence on different continents, their balanced frequencies within populations, their
genomic rearrangements and degeneration worldwide, altogether suggest an ancient
recombination suppression maintained by selection. This study sheds light on an interesting
case of a large supergene partially linked to a mating compatibility locus, and on balancing
selection maintaining differentiated haplotypes, possibly involving deleterious mutations
and/or host adaptation in a devastating tree pathogen.

Introduction:

Recombination increases the efficiency of selection, the purging of deleterious alleles and the generation of potentially beneficial allelic combinations (*1*). However, recombination can also break up beneficial allelic combinations and, therefore, be selected against, generating supergenes, i.e., large regions without recombination encompassing multiple genes (*2*). This is often the case in genomic regions controlling mating compatibility, such as regions determining sex, mating type or self-incompatibility in plants, animals, fungi, algae and oomycetes (*3–8*). Such recombination suppression keeps alleles linked across different genes that prevent self-compatibility or intermediate sexual phenotypes. Even beyond sex and mating-type chromosomes, there is an increasing number of supergene reports (*2*). Striking examples include supergenes controlling social structure in ants, wing color patterns in butterflies, reproductive morphs in birds or host-parasite interactions (*9–12*).

On the long term, however, because of less efficient selection, genomic regions without recombination show signs of degeneration, such as transposable element accumulation, rearrangements, non-synonymous substitutions, decreased gene expression, reduced frequency of optimal codons and gene losses (*13–15*). Recombination suppression in sex chromosomes and other supergenes can nevertheless be maintained by balancing selection or heterozygote advantage (*11, 16*). The mechanisms leading to recombination suppression and allowing their persistence remain poorly. Reporting more diverse supergene constellations in a variety of organisms with contrasting life-history traits is important for understanding the general patterns of their evolution and maintenance (*6, 8, 17, 18*).

Recombination suppression can extend progressively away from sex-determining or mating-type loci, and around other types of supergenes (*4, 8, 18*). This is usually detected by plotting the synonymous differentiation (d_S) between alleles along the chromosome and detecting "evolutionary strata" of differentiation (i.e. with different d_S levels), as differentiation between alleles increases with time since recombination suppression (*4*). A recent theoretical model showed that, even when only partially linked to a mating-type or sex-determining allele, a recombination suppression event trapping a region with fewer deleterious alleles than average could be selected for and be maintained as heterozygous in linkage disequilibrium with the mating-type or sex-determining locus (*19–21*).

A situation that may correspond to this theoretically predicted case has been reported in the

invasive chestnut blight fungal pathogen *Cryphonectria parasitica*, where a large region

without recombination was suggested to occur near the mating-type locus, although not

completely linked to it (*i.e.,* at 3.9 cM of the mating-type locus), based on segregation of

RAPD and RFLP markers in three crosses involving five different parents, from the USA,

Japan or Italy (*22*). Recent genome-wide association studies in a worldwide collection and in

local *C. parasitica* populations in southern Switzerland found significant association of SNPs

with the mating-type locus across a large region (>1 Mb), further suggesting the existence of

reduced recombination in this region (*23, 24*). This genomic region was enriched in SNPs,

transposable elements and copy-number variants, such as deletions, which are consistent with

sequence degeneration and a lack of recombination (*23*), and which prevented its assembly so

far. The strongest footprint of selective sweeps detected genome-wide in *C. parasitica* was

found matching this region (*23*), indicating recent positive selection, and thus possibly

selection for recombination suppression. In *C. parasitica*, the mating-type locus displays two

alleles (*MAT-1* and *MAT-2*) that are both permanently heterozygous in the diploid and

dikaryotic stages (*25*). Although this ascomycete fungus is mostly found as haploid mycelia,

some isolates have been reported to be heterokaryotic at the mating-type locus in several

invasive European and North American populations, i.e., carrying different nuclei, of the

opposite mating types *MAT-1* and *MAT-2* (*23, 24, 26–28*). The theoretical model based on

recessive deleterious mutations showed that non-recombining fragments could be selected for

even with a substantial fraction of the life cycle being haploid (*19*). Hence, *C. parasitica*

seems an excellent biological model to test recent theoretical predictions about the existence

of non-recombining regions partially linked to sex- or mating-type determining loci.

The evolutionary history of the species is well documented as it is an invasive and highly

damaging pathogen, having almost caused the extinction of American chestnut (*Castanea*

dentata) in North America (*29*). From its center of origin in Asia and its original hosts, the

Chinese chestnut (*Ca. mollissima*) and the Japanese chestnut (*Ca. crenata*), *C. parasitica* has

first invaded North America, killing millions of American chestnuts (*Ca. dentata*). In Europe,

at least two distinct introduction events occurred, on the European chestnut (*Ca. sativa*): one

introduction from North America to Italy, and one directly from Asia, probably to the

Pyrenees Mountains. European strains will be therefore hereafter referred to as invasive

European strains introduced from North America or invasive European strains introduced

directly from Asia. The chestnut blight symptoms have been less severe in Europe than in

North America, due to lower susceptibility of *C. sativa* and a virus infecting *C. parasitica* and

causing hypovirulence (*28*, *30*). Clonal lineages have been identified in invasive populations

while sexually reproducing populations occur both in the native and introduced ranges (*23*,

30). The clonal lineages likely still undergo rare sex events, as shown by the introgression of

the missing mating-type allele in most of them (*31*).

Here, we studied the occurrence of recombination suppression in the genomic region

proximal to the mating-type locus in *C. parasitica,* using the previously published genomes

and further generated six new high-quality long-read based genome assemblies of strains from

the native and introduced range of the pathogen. More specifically, we tested whether

recombination was completely suppressed in this region partially linked to the mating-type

locus in *C. parasitica*. We first analyze sexually reproducing invasive European populations,

for which extensive genomic datasets are available in local populations. We assessed, in these

populations, the degree of linkage disequilibrium within the MAT-proximal region, the level

of differentiation between haplotypes, as well as the frequency of the two non-recombining

haplotypes and their association with mating types. We investigated whether other regions of

the genome showed reduced recombination rates and whether genomic footprints of

degeneration were present in the non-recombining region, such as genomic rearrangements,

transposable element accumulation, non-synonymous substitutions and gene disruptions. We

also tested whether this non-recombining region has been gradually expanding, estimated the

age of recombination suppression and investigated whether the predicted gene functions in the

MAT-proximal region could help understanding the evolutionary cause for recombination

suppression and partial linkage to the mating-type locus. We then extended the analyses to

other populations with less extensive genomic resources, i.e. other invasive populations and

native Asian populations, investigating the presence of recombination suppression footprints

and the occurrence of the two differentiated haplotypes worldwide, as well as the possibility

of introgression, using available genomes of closely related species, pathogenic and non-

pathogenic (*23*).

Results

Footprints of recombination suppression in a large region (> 1 Mb) proximal to the

mating-type locus in sexually reproducing invasive populations

We first performed population genomic analyses on available genome sequences within local

populations sampled in southern Switzerland over two temporal frames (early 1990 and

2019), and within the more broadly geographically distributed CL1 cluster in central and southeastern Europe, both having a recombining genetic structure and being invasive European populations introduced from North America (Table S1 (*23*)). We studied only monokaryotic strains, i.e. having either *MAT-1* or *MAT-2* in their genomes, for phasing haplotypes. After masking repeats on the EP155 reference genome (*32*) and removing missing data and rare variants, we identified 8,900 SNPs segregating among 71 strains of the 1990 Swiss population (35 *MAT-1*; 36 *MAT-2*; Table S1), 9,646 SNPs segregating among 62 strains of the Swiss 2019 population (20 *MAT-1*; 42 *MAT-2*; Table S1) and 15,104 SNPs segregating among 88 strains of the CL1 cluster (41 *MAT-1*; 47 *MAT-2*; Table S1) (*23*, *24*). Such a low number of SNPs is consistent with expectations and previous studies in this invasive clonal fungus. We investigated the genome-wide linkage disequilibrium (LD) landscape and looked for large blocks (>1 Mb) of high LD (r^2>0.9) among SNPs within contigs, to look for signatures of reduced recombination along the genome. LD indeed decays by half across a few 100 kb on average in *C. parasitica* in recombining regions of the genome (*31*), so that high LD beyond this distance is indicative of recombination cessation.

We found a large block of high LD on the chromosome called scaffold_2 of the EP155 genome in the three populations (red arrow in Fig 1 and Fig S1), corresponding to the mating-type chromosome. The high-LD block was located near the mating-type locus but did not encompass the mating-type locus itself and will be called hereafter the "MAT-proximal region". In the 1990 and 2019 Swiss populations, SNPs between 0.493 Mb and 1.720 Mb in the mating-type chromosome were all in high LD (with r^2>0.9 between SNPs distant by up to 1 Mb; in this region, mean r^2 of all SNP pairs in the Swiss 1990 population = 0.89; mean r^2 of all SNP pairs in the Swiss 2019 population = 0.83; Fig S1). In the CL1 cluster, the large high-LD block was also present, but was split into two blocks, a large one and a smaller, peripheral one (orange arrows in Fig 1). We indeed detected values of r^2>0.9 between SNPs distant by up to 1 Mb, between 0.532 Mb and 1.535 Mb (mean r^2 of all SNP pairs = 0.94). Between 1.535 and 1.717 Mb, SNPs were also in high LD (mean r^2 = 0.84). The LD between the two high-LD blocks was lower (r^2= 0.59) than within blocks but was still higher than elsewhere along the genome (orange rectangle at the right border of the red triangle; Fig 1). A similar pattern also existed in the Swiss populations (orange rectangle at the right border of the red triangle; Fig S1), although less marked. This is likely due to rare recombination events at one specific locus near the edge of the large fully non-recombining region, lowering the LD level in populations between the two parts of the otherwise fully non-recombining region. The

mating-type locus was located at 1.737 Mb in the EP155 genome, distant 17 kb and 20 kb

from the LD block in the Swiss populations and CL1 cluster, respectively.

We found no other regions of the genomes with $r^2>0.9$ between SNPs across such a large

genomic distance in any population. On scaffold_6 in the CL1 cluster, two smaller SNP

blocks had a $r^2>0.9$ between each other despite being distant (Fig S2A; between SNPs blocks

at 3-20 kb and 2.151-2.278 kb), which is likely due to the major intra-scaffold translocation

described in this region between the EP155 and ESM15 strains (*33*). The maximum size of

the blocks with $r^2>0.9$ on other scaffolds ranged from 135 to 540 kb in the CL1 cluster, as

previously described (*31*). The average distance between SNPs in high LD ($r^2>0.9$) across the

genome was higher on the scaffold_2 than on other scaffolds (in the CL1 cluster, pairwise

Wilcoxon test p-value < 2e-16 with Bonferroni correction). Furthermore, the proportion of

SNPs in high LD ($r^2>0.9$) across the genome was also much higher on the scaffold_2 than on

other scaffolds due to the high LD in of the MAT-proximal region (Fig S2B, Table S2).

Therefore, the large (1Mb) and localized block with maximal LD values proximal to the

mating-type locus both in the Swiss populations and the CL1 genetic cluster stands out as

exceptional in the genome and indicates full recombination cessation, which supports

previous inferences from progeny segregation analyses (*34*). Indeed, even low rates of

recombination homogenize alleles and prevents LD building (*35*). The MAT-proximal non-

recombining region was actually larger (1.23 Mb), but with likely rare recombination events

at one precise locus near its edge, lowering LD between its two fully non-recombining parts.

Noteworthy, we also found blocks of high LD ($r^2>0.9$) around three *vic* loci (i.e., *vic2*, *vic6*

and *vic7* on scaffold_3, scaffold_6, and scaffold_7; Fig S2C), known to be involved in

vegetative incompatibility (vic) and under balancing selection in *C. parasitica* (*24*, *36*). These

high-LD blocks were smaller than the MAT-proximal region but nevertheless larger than

elsewhere in the genome (across more than 100 kb, Fig S2C).

Consistent with a lack of recombination, the MAT-proximal region formed two genetically

highly differentiated haplotypes in these invasive Swiss populations (Fig 2A; Fig S3), as

shown by the two clusters on the principal component analysis (PCA) using only SNPs

located in the MAT-proximal region (Fig 2A), while no structure was detected in the rest of

the genome (Figs 2B and C; Fig S3). The neighbor-net trees further supported the existence of

two differentiated haplotypes in the MAT-Proximal region, contrasting with an otherwise

recombining structure genome-wide (Fig 2D and E). Three strains from the Swiss 2019 population appeared to have an intermediate sequence in the MAT-Proximal region between the two haplotypes (Fig S3D).-These intermediates were rare and may correspond to ancestral polymorphism and/or a third haplotype.

In agreement with the view of a partial linkage with the mating-type locus, the two genetic clusters in the MAT-proximal region appeared strongly associated with mating types, although the association did not reach 100% (Figs 2A and D; Fig S3, Table 1). Indeed, the distribution of mating types between the two PCA clusters strongly deviated from expectations under random association (Table 1; chi-squared test: χ^2 = 33.3; p-value = 5.876e-08 in CL1). We hereafter call the two differentiated haplotypes MAT-Prox1 and MAT-Prox2, referring to the mating-type association. The EP155 genome used as reference for SNP calling above carried the MAT-Prox1 haplotype.

Inversion in the non-recombining MAT-proximal region

To investigate whether rearrangements between haplotypes were present in the MAT-proximal region, we sequenced the genome, using PacBio HiFi, of two strains with alternative MAT-proximal haplotypes: M1400 (MAT-2; MAT-Prox2) and M6697 (MAT-1; MAT-Prox1), originating from the Swiss population in Gnosca and belonging to the CL1 cluster, i.e. European invasive populations introduced from North America (*23*, *24*). We build high-quality genome assemblies: statistics of the assemblies were in the same range as those of the genome assemblies of the ESM15 and EP155 strains previously sequenced (Table S3; Demené et al 2022; Crouch et al 2020). The mating-type chromosome could be assembled as a single contig for the first time, in the M1400 genome, likely corresponding to a full chromosome, and was assembled into two contigs in the M6697 genome. We therefore used the M1400 genome as a reference for subsequent analyses. To identify the location of the MAT-proximal region in the M1400 genome, we computed LD by read mapping in the 1990 Swiss populations. We found the large block (> 1 Mb) of high LD (r^2>0.9) between 7,285,137 and 8,828,934 bp (red arrow; Fig. S4). The MAT-proximal region was also located 20 kb away from the mating-type locus (located at 7.265 Mb) and was 1.540 Mb long using the M1400 genome as reference. Consistent with the results using the EP155 genome as reference, we found that the high-LD region was divided into two higher-LD blocks near the edge at 7.392 Mb (orange arrows in Fig. S4).

The two new high-quality genome assemblies from the invasive Swiss population (introduced

from North America) revealed an inversion between the two haplotypes in the MAT-proximal

region (Fig 3). The two newly sequenced PacBio genomes were indeed collinear except for

the mating-type chromosome (Fig S5), where we found a large inversion (>1 Mb) between

the M1400 and M6697 genomes (Fig 3; Fig S5). Breakpoints of the inversion were located at

ca. 7.455 and 8.563 Mb of the tig00000001 contig in the M1400 genome and at ca. 3.380 and

4.185 Mb of the tig00000060 contig in the M16697 genome. The region affected by the

inversion was ca. 300 kb smaller in the M6697 genome than in the M1400 genome, while the

whole MAT-proximal region was ca. 100 kb smaller. The split into the two high-LD blocks in

natural populations reported above was outside of the inversion, i.e. 70 kb from the inversion

breakpoint (orange arrows in Fig S4). LD in natural populations was higher within the

inversion (median value of $r^2 = 1$) than in other regions of the MAT-proximal region (median

value of $r^2 = 0.64$; Wilcoxon rank sum test with continuity correction; $W = 4.8714e+10$; p-

value < 2e-16).

We looked for centromeres, as the non-recombining regions near mating-type loci in other

ascomycetes, when they occur, either capture the centromere (*37, 38*), or are associated to the

occurrence of a single crossing-over between the centromere and the mating-type locus (*8, 39,*

40). The centromere in *C. parasitica* was likely at 4.380-4.536 Mb on the mating-type

chromosome in the M1400 genome, as we detected here a peak in TE density and a drop in

GC content (Fig 4A-B). The MAT-proximal region and the inversion thus did not include the

putative centromere and was instead located about 2.9 Mb away of the putative centromere.

Higher genetic differentiation and lower genetic diversity within the MAT-proximal

region than in recombining regions

When analyzing polymorphism within the invasive 1990 Swiss population using the M1400

genome as reference, we found higher genetic differentiation between strains of the two

MAT-proximal haplotypes in this region than elsewhere along the genome (Fig 4C-D; Fig

S6A-B; Table S4), as expected for a non-recombining region in LD with the mating-type

locus. For example, the F_{ST} values between the two non-recombining haplotypes within the

MAT-proximal region in the 1990 Swiss population were nearly maximal (median F_{ST}=0.93

per 50 kb window), while they were near zero in the rest of the genome (median F_{ST}=0.02;

Wilcoxon test p-value < 2e-16 with Bonferroni correction; Fig 4C; Fig S6A). Such F_{ST} values

close to 1 indicate a lack of shared polymorphism and therefore support the inference of

recombination suppression in the MAT-proximal region. We also found higher absolute genetic divergence (d_{XY}) in the MAT-proximal region compared to other regions (Wilcoxon test p-value < 2e-16 with Bonferroni correction), with a peak of number of fixed differences in the middle of the MAT-Proximal region and of the inversion (Fig 4D; Fig S6A; Table S4), suggesting a pattern of gradual expansion of recombination cessation within the MAT-proximal region, from its center to the edges. The 95th percentile of highest d_{XY} values was for the windows encompassing the genomic region between 8.190 à 8.280 Mb.

Within each haplotype, the genetic diversity at the MAT-proximal region was lower than in the rest of the genome (Wilcoxon test p-value < 2e-16 for the MAT-Prox1 haplotype; p-value = 2.3e-07 for the MAT-Prox2 haplotype; Fig 4E-F; Fig S6C; Table S4), as expected for a region without recombination associated with the mating-type locus, as its effective population size is half as in recombining regions. The diversity was especially low in the pool of isolates with the MAT-Prox1 haplotype (see all MAT-Prox1 sequences clustering on a single point on the PCA in Fig 2A and in the neighbor-net tree in Fig 2D, in contrast to the more scattered MAT-Prox2 sequences); such a lack of diversity contrasting with other genomic regions and the other haplotype suggests that a selective sweep occurred in this MAT-Prox1 haplotype. This supports previous inferences of a selective sweep in this region based on site-frequency spectrum analyses (*23*).

In addition to footprints of selective sweeps at short time scales, we detected signals for long-term balancing selection in the MAT-proximal haplotypes. Computation of the Tajima's D statistics within all strains suggested signatures of balancing selection in the MAT-proximal region as expected in a region without recombination associated with the mating-type locus maintained at a frequency close to 1/2. Tajima's D values in the MAT-proximal region pooling all sequences (median D=3.0 per 50 kb window) were higher than in the rest of the genome (median D=-0.14; Wilcoxon test p-value < 2e-16 with Bonferroni correction; Fig 4G; Fig S6D). In contrast, Tajima's D was significantly lower than the rest of the genome within each pool of haplotype, including in the MAT-Prox2 haplotype, further supporting the inference of positive selection (Wilcoxon test p-value = 0.0037 for the MAT-Prox1 haplotype; p-value = 0.0022 for the MAT-Prox2 haplotype with Bonferroni correction; Fig 4H-I; Fig S6D; Table S4). We found no difference in population diversity statistics between the recombining part of the mating-type contig and the other contigs (Table S4A).

The study of synonymous divergence (d_S) between M1400 and M6697 haplotypes suggests

that the recombination suppression is at least 1.5 million years old and that there may be a

pattern of evolutionary strata within the MAT-proximal region, i.e. segments with different

levels of differentiation between haplotypes that could indicate stepwise expansion of

recombination suppression away from the mating-type locus. Per-gene synonymous

divergence (d_S) is typically used for detecting gradual expansion of recombination cessation

and for estimating its age, as it is considered a good proxy for the time since recombination

suppression. Indeed, when recombination is suppressed, mutations accumulate independently

in the two non-recombining haplotypes. We plotted the per-gene synonymous divergence (d_S)

between M1400 and M6697 along the M1400 genome, as its mating-type contig is likely

being assembled as a full chromosome (Fig S7A&E; Table S4). Consistent with

recombination suppression, we found significantly higher d_S values in the MAT-proximal

region (computed for 71 genes) than in the other regions of the mating-type contig (pairwise

Wilcoxon test p-value <2e-16 with Bonferroni correction) and other contigs (pairwise

Wilcoxon test p-value <2e-16 with Bonferroni correction). We found no significant

differences between the other regions of the mating-type contig and other contigs (pairwise

Wilcoxon test p-value = 0.23 with Bonferroni correction). The region with higher d_{XY} values

within the MAT-proximal region (between 8.190 à 8.280 Mb) showed significantly higher d_S

values than the rest of the MAT-Proximal region (Wilcoxon test p-value = 0.014) supporting

the view of a gradual expansion of recombination cessation within the MAT-proximal region,

from its center to the edges. Note that the per-gene d_S could not be computed for the genes

with a copy not predicted in one of the haplotypes and therefore mostly in the oldest, most

degenerated regions (see below), which may prevent detecting the oldest stratum using d_S

values, while the d_{XY} values were computed in sliding windows all along the MAT-proximal

region. Using synonymous substitution rate estimates in fungi (*41, 42*) and the mean d_S values

across genes of the MAT-proximal region, we estimated the age of the divergence between

the two haplotypes to be at least 1.5 Million years when considering the mean d_S values of all

genes in the MAT-proximal region (dS mean= 0.0495) and at least 7.2 Million years when

considering only the 15% genes with highest d_S values dS mean= 0.24), under the hypothesis

of gradual expansion of the non-recombining region (Table S4).

Enrichment of transposable elements in the non-recombining region and inversion

breakpoints

When studying the M1400 and M6697 high-qualities assemblies, we detected an enrichment of transposable elements (TEs) in the MAT-proximal region in the two haplotypes compared to the rest of the genome. The percentage of bp occupied by TEs (TE load) was higher than 50% in the MAT-proximal region, while it was only 9% on average in other regions. TE load in the MAT proximal region was higher in the M6697 haplotype (76%, MAT-1; MAT-Prox1) than the M1400 haplotype (68% MAT-2; MAT-Prox2). Class I retrotransposons with Gypsy (LTR-Ty3) and LARD elements were the most abundant TEs in all genomic regions (autosomes, MAT-Proximal region and the rest of the mating-type contig), representing >70% of TE annotations (Fig 5A-B), as shown previously (*33*). The TE load however varied significantly among TE families and genomic regions (ANOVA; Table S5). In M6697 for example, TIR elements appeared less abundant in the non-recombining than recombining regions while LTR-Ty3 elements were more frequent in the MAT-proximal region (Fig 5B). In M1400, LARD elements were more frequent in the MAT-proximal region than in recombining regions (Fig 5B).

The presence of transposable elements at the two inversion breakpoints in the two genomes (Fig 5A) suggests that the inversion may have occurred via non-homologous recombination mediated by these elements. By dating retrotransposon insertions using between-copy divergence and within-copy LTR divergence in both M1400 and M6697 genomes (Fig 5C and D), we found that TEs within the inversion and around the inversion breakpoints were older than TEs farther from the breakpoints. This is expected in non-recombining regions as selection is there less efficient to purge TE insertions so that they remain there longer (Duhamel et al. 2023). The date estimates of LTR element insertions (Fig 5D) further indicated that the TE accumulation in the MAT-proximal region was as old as a few million years, confirming the age estimated based on sequence divergence between MAT-proximal haplotypes.

Gene functions, pseudogenization and degeneration in the non-recombining region

Recombination suppression could allow linking genes with mating-type antagonistic effects to the mating-type locus, or link together co-adapted alleles, for example for parasitizing different host genotypes. We however found no gene function known to be involved in mating or mating compatibility, nor in pathogenicity, in the MAT-proximal region (Table S6). We nevertheless identified genes potentially coding for proteins with kinase domain or proteins with homeodomains (Table S6), that are often involved in the regulation of

developmental pathways. Gene functions in the d_{XY} peak (95th percentile) included general functions such as protein kinase-like domain, SET domain (i.e. with methyltransferase activity) and RNA-binding domain families (Table S6).

By comparing gene predictions in the M6697 and M1400 newly assembled genomes, we identified a higher prevalence of gene disruptions (i.e. genes predicted in a single of the two haplotypes) in the MAT-proximal region than elsewhere in the genomes. Out of 175 groups of single-copy orthologous genes in the MAT-proximal region, 80 (46%) were only present in M1400 and 24 (14%) only present in M6697. In other genomic regions, 3% of genes were only present in one genome. This suggests that gene disruption occurred in the non-recombining region, in higher proportions than elsewhere in the genome (Fisher's exact test p-value < 2.2e-16), and especially in the M6697 genome (MAT-1; MAT-Prox1). These gene disruptions are mainly in the middle of the MAT-proximal region and of the inversion (Fig 5A), i.e. the region corresponding to the d_{XY} peak in the Swiss population and to the high d_S values between the M1400 and M6697 genomes (Figs 4D and S7). In the M1400 genome, gene disruptions are more scattered all along in the MAT-proximal region (Fig 5A). The enrichment in TEs and in gene disruptions in the MAT-Proximal region explains the lower gene density in this region (Fig 4) and supports the inference of an old full recombination suppression. The occurrence of high TE load and gene degeneration in both MAT-Proximal haplotypes further confirms the lack of homozygotes in this region, in which recombination could purge deleterious mutations, or indicates that recombination suppression occurs even in homozygotes, although it remains to be investigated.

We found higher non-synonymous substitution rate between the M1400 and M6697 genomes compared to the baseline substitution rates (d_N/d_S values) in the MAT-proximal region (median per gene $d_N/d_S = 0.80$) than in other contigs ($d_N/d_S = 0.43$; pairwise Wilcoxon test p-value = 1.7e-07 with Bonferroni correction), consistent with relaxed purifying selection due to recombination suppression in the MAT-proximal region (Fig S7A; Table S4).

Recombination suppression in the MAT-Proximal region in other populations assessed from high-quality genome assemblies

We analyzed five additional high-quality genome assemblies for strains from the Asian native range (CL2 and CL4 clusters) and the invasion range (including two North American strains belonging to CL1 cluster and one European strain introduced directly from Asia and

belonging to CL2 cluster), three of which were generated for the present study (Table S3).
Most of the high-quality genomes displayed the same MAT-Proximal chromosomal
arrangement as M1400 (MAT-Prox2, Swiss population and belonging to CL1 cluster),
suggesting that it is the ancestral state (Fig S8A-C), although the incomplete assemblies may
prevent detecting other rearrangements. The inversion detected in M6697 (MAT-Prox1, Swiss
population and belonging to CL1 cluster) was also present in the EP155 strain (MAT-Prox1,
North America and belonging to CL1 cluster, Crouch et al, 2020; Fig S8D), indicating that
this inversion was already segregating in early invasive populations in North America. In
other genomic regions outside of the MAT-Proximal region, all analyzed high-quality
genomes were colinear to the ESM15 reference genome, except the highly rearranged EP155
genome, as previously reported (*33*).

The MAT-Proximal region in all the high-quality genomes, from Asian native strains and
invasive strains, displayed signs of degeneration, with a high TE load (Fig 6B) and gene
disruptions, as seen from lower gene densities (Fig S8A) and missing orthologs (Fig 6C),
confirming the ancient age of recombination suppression and its occurrence in the native
range. Study of d_S and d_N/d_S between the MRC10 strain, belonging to the European clonal
lineages of direct Asian origin (CL2 cluster), the M6697, M1400 and the two Asian native
strains further confirmed recombination suppression in the native range and in invasive strains
from different origins (North America or directly from Asia). The d_S and d_N/d_S values were
indeed significantly higher in the MAT-proximal region than other genomic regions (Fig S7
B-E). Mean d_S values in the MAT-proximal region were in the same range as d_S values
between the M6697 and M1400 genomes, suggesting a similar date of recombination
suppression and, therefore, that recombination suppression was ancestral. The MAT-proximal
region appeared smaller in the two Asian strains XIM9508 and ESM15 (<1 Mb) than in
invasive strains (Table S3), but this may be due to the incomplete genome assemblies.

Presence of the two MAT-Proximal haplotypes in other populations
By analyzing available additional Illumina genomes in the native range and in invasive
populations, with less dense sampling within populations, we recovered the two haplotypes in
the MAT-proximal region, i.e., high differentiation between two clusters in the MAT-
Proximal region, contrasting with a lack of genetic subdivision in other genomic regions or a
completely different genetic subdivision, corresponding to previously described population
genetic structures (Figs 7 and FigS9). In particular, the neighbor-net tree with multiple strains

sequenced previously from the clusters CL2 and CL3 (from the introduced range and the
native range, South Korean and Japan) and from the cluster CL4 (from the native range,
China;) reinforces the view of balancing selection on the MAT-Proximal haplotypes.
Sequences from the CL1 and CL2 clusters were indeed intermingled in the MAT-proximal
region (Fig 7A), while the structure in the rest of the genome instead corresponded to the
different population clusters, i.e., CL1, CL2, CL3 and CL4 (Fig 7B).

However, the two haplotypes were less differentiated in the native range and in invasive
populations originating directly from Asia than in invasive populations from the first invasion
wave via North America. For example, branches length on the neighbor-net indicates a lower
differentiation between haplotypes within CL2 (Fig 7A) than within the European CL1 or
Swiss invasive populations (Fig 2D; Fig S3). Within CL2, F_{ST} and d_{xy} values between the two
haplotypes in the MAT-proximal region were lower than in the invasive population coming
from North America (Table S7).

We found no association of the MAT-Proximal haplotypes with mating types at the world
scale (Figs 7 and S9), which may be due to the lack of extensive sampling within local
populations. As the MAT-Proximal region is only partially linked to the mating-type locus,
founder effects may yield different associations between MAT-Proximal haplotypes and
mating types in different populations. Only in the S12 European invasive clonal lineage (only
MAT-1), we found a single MAT-Proximal haplotype (MAT-Prox1; Fig S9B). In two
invasive European clonal lineages of American origin (re092 and re103), the strains with a
mating type different from their clone-mates due to localized introgression from other clonal
lineages (*31*) also had their MAT-Proximal region introgressed (Fig S9C), reinforcing the
view of the strong linkage between the mating-type locus and the MAT-Proximal region, and
of an advantage of maintaining an association between MAT-proximal haplotypes and mating
types within populations. The few 2019 Swiss strains that exhibited an intermediate sequence
in the MAT-Proximal region clustered with strains from Europe with a direct Asian origin
(Fig S10C). This may represent a third MAT-Proximal haplotype, associated with the second
invasion in Europe, directly from Asia.

No evidence of introgression in the MAT-proximal region from other species
The stronger differentiation in the MAT-Proximal region in the Swiss invasive population
than in other populations could also be due to an introgression event from a closely related

species; however, the small genetic distance and the reticulations observed in the neighbor-net tree between the M6697 Swiss reference genome and the Asian genomes M8444 and XA19 rather indicate that the MAT-Prox1 haplotype in Switzerland originated from Asian *C. parasitica* populations, likely a population different from those at the origin of the CL2 invasive cluster (Fig 7).

Gene genealogies also provided no evidence of introgression in the MAT-proximal region from closely related *Cryphonectria* species. We performed a gene orthology analysis including three genomes of the closely related species *C. japonica*, C. *carpinicola*, and *C. naterciae* that were previously sequenced with short read technologies (*43*). We retrieved only nine genes from the MAT proximal region that were shared between the M6697 and M1400 genomes that had orthologs in one of the related species. This may be due to low quality assemblies in the outgroups (*43*) and/or a region without recombination also in the outgroups with gene losses and many repeats. We found no gene with a topology consistent with introgression from another species into the MAT-Proximal region, i.e. with a MAT-Prox1 or MAT-Prox2 haplotype that would branch with an outgroup allele rather than with the alternative haplotype of *C. parasitica*. The high load of TEs and gene disruptions in the two MAT-Proximal haplotypes does not fit with the introgression hypothesis either, unless the introgression occurred from a species with recombination suppression in the MAT-Proximal region.

Discussion

***A large and million-year-old non-recombining region partially linked to the mating-type locus in* Cryphonectria parasitica**

We found strong evidence for the complete cessation of recombination in a region of more than 1 Mb (1.2 Mb to 1.5 Mb depending on the haplotype), strongly but not completely linked to the mating-type locus in *C. parasitica*. Indeed, we detected maximal levels of linkage disequilibrium in otherwise recombining populations and the existence of two highly differentiated haplotypes, without shared polymorphism, while even low rates of recombination can homogenize alleles and prevent LD build-up (*35*). The full cessation of recombination was further supported by the existence of a previously unknown, large rearrangement (an inversion) in the center of the non-recombining region in invasive strains. The higher TE load in this region constitutes additional evidence for full recombination cessation, as the lack of recombination renders selection against TE accumulation less

efficient, in particular due to Muller's ratchet: two chromosomes with different TE insertions
cannot recombine to produce a chromosome free of TE insertions. TE takes hundreds of
thousands of years to accumulate in non-recombining regions (*15*), so that the high TE load in
the MAT-Proximal region in *C. parasitica* indicates old recombination cessation. The
particularly high frequency of missing genes and non-synonymous substitutions in the MAT-
Proximal region worldwide further indicates degeneration and ancient recombination
suppression. The estimates for the age of recombination suppression based on the divergence
between haplotypes and on TE insertions were in fact at least a few million years. However,
in contrast to other known evolutionary strata on sex or mating-type chromosomes (*4*, *8*), the
region without recombination was not completely linked to the mating-type locus. Indeed, the
occurrence of rare recombination events between the non-recombining MAT-Proximal region
and the mating-type locus are supported by previous segregation analyses (*34*) and the lack of
association found here between the MAT-Proximal haplotypes and the mating types at the
worldwide scale, as well as their incomplete association within populations.

Large inversion in the MAT-proximal non-recombining region

We found a large inversion between the two highly differentiated MAT-proximal haplotypes
in invasive populations originating from North America (CL1 genetic cluster, Swiss 1990 and
2019 populations and US populations). The chromosomal arrangement of the M1400 MAT-
Prox2 haplotype is likely the ancestral state as it is the most frequent worldwide, although a
more complete sampling in the native range is required to obtain definitive conclusion. The
inversion would therefore be the M6697 MAT-Prox1 haplotype and may contribute to
recombination suppression. However, the non-recombining region was larger than the
inversion, which indicates that additional proximal mechanisms suppress recombination. The
inversion may even be a consequence rather than a cause of recombination cessation, as
previously found in other fungal mating-type chromosomes (*37*, *39*, *40*, *44*). Other proximal
causes of recombination cessation may for example be genetic recombination modifiers,
epigenetic marks or histone modifications (*45*, *46*). Alternatively, the inversion may have
constituted a cause for a first step of recombination suppression (i.e., for a first evolutionary
stratum), as actually suggested by the higher d_S and d_{XY} values in the middle of the inversion
compared to the rest of the non-recombining region. Evolutionary strata have been reported in
many sex chromosomes in plants and animals (*13*, *47–49*) and also around mating-type loci in
fungi (*8*, *50*). The lack of rearrangements at the edge of the MAT-proximal region may allow
rare recombination or gene conversion events, which may explain the presence of two distinct

LD blocks with lower LD between them and the lower differentiation at the edges of the
MAT-proximal region.

The MAT-proximal non-recombining region is million-year-old and degenerated

Following recombination suppression, selection efficacy decreases. We detected footprints of
degeneration, with particularly high occurrence of gene disruptions and of TE insertions in the
MAT-proximal region, both in Europe and in the native Asian range of the fungus. (*33*) had
already reported an enrichment of LTR-Ty3 (*Gypsy*) element in the region near the mating-
type locus in *C. parasitica*. We found LTR-Ty3 elements at the inversion breakpoints and an
enrichment in the MAT-Prox1 haplotype. Interestingly, the same TE family (LTR-Ty3) seem
to have expanded in the non-recombining regions in other fungi with recombination
suppression on their mating-type chromosomes (*15*), suggesting they may be more prone to
accumulate in non-recombining regions. Accumulations of transposable elements have also
been detected in other mating-type or sex chromosomes and supergenes (*13–15*, *51*), although
some other large supergenes do not seem to exhibit TE load, likely due to recent
introgression, gene conversion or double crossovers (*52–55*).

The transposable element accumulation in the MAT-proximal region and the enrichment in
gene disruptions support the estimated age of recombination suppression at a few million
years ago based on mean pairwise gene sequence divergence between haplotypes. The finding
of the two haplotypes and of the inversion in the two invaded continents, Europe and North
America, as well as the occurrence of degeneration footprints in the MAT-Proximal region in
the native Asian range, further support the inference of an ancient recombination suppression
maintained by balancing selection. Indeed, even rare recombination events allow purging
deleterious mutations and TE insertions. This means that the association of the MAT-
Proximal haplotypes with mating types is very strong within all recombining populations
(even if not at the worldwide scale, as founding effects may generate different associations
between mating types and MAT-Proximal haplotypes); it may also be that recombination is
suppressed within the MAT-Proximal region even for homozygotes in this region. This is in
agreement with previous genetic maps reporting lack of recombination near the mating-type
locus in crosses involving Japanese, North American and Italian isolates (*22*). The high TE
load and degeneration level in terms of gene disruption in strains from Asian and North

American origins indicate a worldwide recombination cessation, so likely ancestral to all these native and invasive populations.

The differentiation between the MAT-Proximal haplotypes appeared greater in the CL1 cluster and Swiss populations (from the first invasion event, from North America) than in the other clusters, which may be due to a founder effect, the lack of sampling in the source population in Asia and/or a selective sweep. The MAT-Prox1 haplotype in the invasive European population from the first introduction event shows an inversion, a strong selective sweep, and the most degenerated locus. These findings point to a strong selection maintaining the two haplotypes differentiated and non-recombining, and perhaps different selection regimes or successive selection events during the various invasion waves of *C. parasitica.*

The higher differentiation between the MAT-Proximal haplotypes in the first invasion wave in Europe could also have resulted from introgression from a closely related species. The evolution of supergenes by introgression has been reported for example in ants and butterflies (*56–58*). However, we found little evidence for an introgression: i) the two haplotypes are present in the CL2 cluster of invasive European populations directly originating from Asia, ii) the MAT-Prox1 haplotype in the CL1 cluster more differentiated from the MAT-Prox2 haplotype than in the CL2 cluster is genetically close to an Asian haplotype, iii) footprints of recombination suppression are present in the native range in terms of degeneration and particular genomic structure in the MAT-Proximal region, and iv) we found no signatures of introgression from closely related species in gene genealogies.

The high differentiation into two distinct haplotypes, their occurrence in at least two genetic clusters, their balanced frequencies within populations, their degeneration in terms of TE load and disrupted genes on different continents, and the strikingly different genomic subdivision in the MAT-Proximal regions compared to the rest of the genome, altogether suggest an ancient recombination suppression maintained by balancing selection for at least a few million years. The degeneration in both haplotypes indeed indicates a lack of homozygotes at the diploid stage, in which recombination could occur and purge deleterious mutations, or suppressed recombination even in homozygotes.

The selective forces potentially maintaining the two haplotypes in the MAT-proximal region

The genes found missing in the non-recombining region in one or the other haplotype may
help maintaining the two haplotypes by selection due to deleterious mutation sheltering, i.e.
thanks to a heterozygote advantage. Such overdominance advantage may only be a
consequence of recombination suppression, and contribute to its maintenance, or may even be
its initial cause (*11*, *19–21*). Indeed, this situation of a non-recombining region partially
linked to the mating-type locus and with two haplotypes maintained by selection fits
predictions from a recent theoretical model showing that recombination suppression can be
selected for near permanently heterozygous alleles for protecting haplotypes with fewer
deleterious mutations than average in the population, while sheltering their few recessive
deleterious alleles. Importantly, the model predicts that selection can also occur for regions
partially linked to heterozygous alleles (*19*). Similar overdominance effect due to the
sheltering of deleterious mutations can lead to balancing selection in the context of local
adaptation (*59*). These models require the existence of a substantial diploid or dikaryotic
phase for sheltering recessive deleterious alleles in a heterozygous state, although the
mechanism also works with a substantial haploid phase (*19*). In fact, although *C. parasitica*
has a predominant haploid phase, strains can be found as dikaryons heterozygous for the
mating-type locus in some natural populations, and not only as monokaryons (*23*, *24*, *26*, *27*).
The importance and frequency of heterokaryons in *C. parasitica* in nature would deserve
further investigations. It may be sufficient if recessive deleterious alleles are sheltered in ca
20% of individuals for selecting recombination suppression (*19*), and/or if the genes involved
are preferentially expressed during the diploid or dikaryotic phases. This hypothesis would
explain the long-term balancing selection, and also the selective sweep by a recent extension
of recombination suppression, possibly corresponding to the inversion found in invasive
population from the first introduction. This hypothesis of balancing selection, or the
hypothesis of overdominance (i.e. heterozygote advantage), does not require the same
association between the MAT-Proximal haplotypes and mating types across all populations
worldwide, it is sufficient that the MAT-Proximal haplotypes are strongly associated to
mating types within populations. Further sampling will be required to test whether the MAT-
Proximal haplotypes are associated to mating types within local populations in the CL3 and
CL4 clusters and in the native Asian range.
680
It could indeed also be that genes evolve under overdominance in this region for other reasons
than deleterious mutations, so that a partial linkage to a permanently heterozygous locus is
beneficial, perhaps in relationship with the pathogenic lifestyle of the fungus. One could

imagine, for example, that heterozygosity could be advantageous at genes involved in
virulence against the tree host, and especially the novel hosts colonized in invasive ranges, or
in resistance against a parasitic virus known to negatively affect fitness in *C. parasitica* (*60*,
61). As an alternative to a heterozygote advantage, the selection maintaining the
recombination suppression and the two haplotypes in the MAT-proximal region could be
some kind of negative-frequency dependent selection of beneficial allelic combinations,
possibly linked to a trench-warfare-like arms race with the host tree or the microbial
community, where partial linkage to the mating-type locus would help maintenance in
balanced proportions and therefore would prevent allele loss (*59*, *62*). Such a role in host-
pathogen interactions would also be consistent with the selective sweep footprints detected in
the MAT-proximal region if there is recurrent positive selection for improving the efficiency
of the pathogen weapons within each of one of the two MAT-proximal haplotypes. Balancing
selection and location adjacent to a supergene was recently found for loci involved in host
resistance in the *Daphnia–Pasteuria* system (*12*).

Another hypothesis for explaining such extension of recombination suppression beyond the
mating-type locus is antagonistic selection, i.e. linkage of alleles that would improve fitness
of a MAT-1 gamete while being deleterious in a MAT-2 gamete, or vice-versa. However, full
recombination suppression with the mating-type locus would be expected under such
antagonistic selection, as well as the same association between the MAT-Proximal alleles and
the mating-type in all populations, in contrast to our findings. In addition, we found no
particular predicted function in the MAT-proximal region that could be related to mating
compatibility and there are very little functions, if any, with possible antagonistic functions
between mating types in fungi (*8*, *63*).

Conclusion
In conclusion, we provide strong evidence for the existence of a non-recombining region
partially linked to the mating-type locus and maintained by selection in the chestnut blight
fungus *C. parasitica*. Such supergenes are increasingly reported in a variety of organisms
thanks to new sequencing technologies (*2*, *10*, *56*, *57*, *64*). This study therefore contributes to
our understanding of how selection for recombination suppression promotes the maintenance
of two divergent haplotypes despite degeneration. The resulting supergene in *C. parasitica*
displayed contrasting differentiation levels between haplotypes and different chromosomal
arrangements in native and invasive populations. We found strong selection footprints in the

MAT-proximal region in the first introduction of the pathogen in Europe from North America
and a chromosomal inversion. The supergene displayed lower differentiation between
haplotypes in a second invasion wave directly from Asia. This supergene may thus underlie
important adaptive traits and thereby provide important applications for the control of a
devastating tree pathogen. In addition, the high-quality genome assemblies provided here,
from the native and introduced ranges, will more generally be useful for studies aiming at
understanding the evolution of this pathogen.

Material and Methods

Strains and genomic data

For population genomic analyses, we studied a collection of 386 monokaryotic *C. parasitica*
strains sampled worldwide, from the native and introduced ranges of the pathogen, and
sequenced previously with the short-read Illumina technology (strain information are
presented in Table S1; (*23*, *24*, *31*)). Data were downloaded from NCBI Bioproject numbers
PRJNA604575, PRJNA644891 and PRJNA706885. We focused our analyses first on
European invasive strains originating from North America. We studied 88 strains belonging
to the CL1 genetic cluster, in central and southeastern Europe (*23*). We excluded the putative
clonal genotypes previously identified (*23*). We also studied 71 strains sampled in the 1990
and 62 strains sampled in 2019 in southern Switzerland (*24*). Mating-type ratios close to ½
and population structure analyses of the CL1 genetic cluster and the Swiss 1990 population
suggest frequent recombination in these populations (*23*, *24*). The mating-type ratio was 1/3
MAT-1 - 2/3 *MAT-2* in the 2019 Swiss population and population structure analyses suggested
regular sexual reproduction and recent population bottleneck . To study more broadly the
presence of the two haplotypes in the MAT-proximal region, we analyzed monokaryotic
strains belonging to the CL2, CL3, CL4 genetic clusters, as well as the S12 European invasive
lineage (*23*) and additional monokaryotic strains from the US and Europe (with an Asian or
North American origin (*31*)). We excluded heterokaryotic strains, i.e. strains having both
MAT-1 and MAT-2 alleles, as the phase was challenging to infer.

For mapping and SNP calling, we first used as reference the 43.9 Mb genome sequence
EP155 v2.0 of the strain EP155 (*MAT-2,* North America, CL1 cluster) available at the Joint
Genome Institute (http://jgi.doe.gov/)(*32*). For comparative genomics, we used the published
genome of the strain ESM15 (MAT-2, Japan, CL2 cluster; (*33*)) available at

DDBJ/ENA/GenBank   on   the   bioproject   PRJNA700518   under   the   accession
JAGDFO000000000. We additionally sequenced *de novo,* with long-read technologies, five
strains from the native and introduced ranges of the pathogen. We sequenced with PacBio
Hifi the genomes of the strains M1400 (MAT-2) and M6697 (MAT-1) sampled in Gnosca,
southern Switzerland (*23*). Mycelia were stored as glycerol stocks at -80 C after strain
isolation. To produce mycelium for DNA extraction, isolates were inoculated onto
cellophane-covered potato dextrose agar plates (PDA, 39 g/L; BD Becton, Dickinson and
Company, Franklin Lakes, USA) (*65*) and incubated for a minimum of 1 week at 24°C, at a
14 hr light and 10 hr darkness cycle. Mycelium and spores were harvested by scratching the
mycelial mass off the cellophane, transferring it into 2 mL tubes and freeze-drying it for 24 hr
(*23*). DNA extraction was performed with the NucleoBond High Molecular Weight DNA kit
from Macherey-Nagel, with the mechanical disruption of about 30 mg of lyophilized
mycelium with two beads of 3 mm for 5 min at 30 Hz. Sequencing was outsourced to
Novogene, the Netherlands. We additionally sequenced with Oxford Nanopore MinION
technology the strains XIM9508 (China, CL4 cluster, MAT-1), MRC10 (South Western
France introduced directly from Asia, CL2 cluster, MAT-2) and DUM-005 (USA, CL1
cluster, MAT-2). These isolates had been collected for previous studies (*30*, *66*, *67*). The
protocols for mycelium production, DNA extraction and sequencing for these three strains
were the same as in (*33*).

Short-read data processing and SNP calling
We used SNP calling datasets of the genomes from monokaryotic strains of the CL1, CL2,
CL3, CL4 genetic clusters, the S12 invasive lineage and the Swiss populations against the
EP155 reference v2.0 genome obtained in (*23*, *24*). We performed mapping and raw SNP
calling of short-read data using the M1400 new genome assembly as a reference as described
in (*23*, *24*). Briefly, we trimmed reads with Trimmomatic v0.36 (*68*) and aligned them with
Bowtie 2 v2.3.5.1 (*69*) and SAMtools v1.9 (*70*) to the EP155 v2.0 genome. Raw SNP calling
and filtration for quality were conducted with the genome analysis toolkit GATK v3.8 and
v4.0.2.0 (*71*). We used the filtration parameters described in (*23*): QUAL>=100,
MQRankSum (lower) >= -2.0, QD:_20.0, MQRankSum (upper) <=2.0, MQ:_20.0,
BaseQRankSum (lower) >=-2.0, ReadPosRankSum (lower) >=-2.0, ReadPosRankSum
(upper) <=2.0, BaseQRankSum (upper) <=2.0. We further removed SNPs overlapping with
transposable elements predicted *de novo* in the EP155 v2.0 genome by (*23*).

Population genomics analyses

For all population genomics analyses, we excluded SNPs with missing data in more than 10% of the strains and kept only polymorphic strains with vcftools v0.1.16 (*72*). To study linkage disequilibrium, we further excluded rare variants (minor allele frequency <0.1) with the vcftools (*72*) option --maf 0.1. We computed LD with the --hap-r2 option of vcftools (*72*) for each scaffold and each population separately. We used the R package LDheatmap v1.0-6 (*73*) to plot LD r^2 values among SNP pairs. To perform principal component analyses (PCAs), we first used vcftools (*72*) to convert VCF format files in Plink format. We then used the Plink v1.90b5.3 (*74*) –pca command to run PCA analysis. We used the R package PopGenome v2.7.5 (*75*) to compute nucleotide diversity, Tajima's D values and the F_{ST} and d_{xy} index. We used the R package ggplot2 (2_3.5.0) to plot results. Pairwise Wilcoxon tests were performed in R with Bonferroni or false discovery rate correction.

We built neighbornet trees using SplitsTree4 v 4.19.2 (*76*). VCF file were converted to nexus format using PGDSpider v1.0 tool (*77*). For the study of other populations, we included only one strain of each haplotype of the invasive European population but not all CL1 and Swiss strains for tree readability; we nevertheless checked that the results were the same with all CL1 and Swiss strains.

Long-read based assemblies

PacBio Hifi reads of strains M1400 and M6697 were assembled using canu v1.8 (*78*) program with a set genome size of 44 Mb. Oxford Nanopore MinION reads and Illumina reads of the genomes of XIM9508 and MRC10 strains were assembled using HybridSPAdes (*79*). Assemblies were manually curated as previously described (*33*) and scaffolds were cut when an evidence of a chimeric connection was detected (i.e. mis-paired short reads). For DUM005, the assembly was generated by Ra (https://github.com/lbcb-sci/ra) with basic parameters as the Ra assembly quality outperformed the HybridSPades assembly quality (*33*). Ra uses Oxford Nanopore MinION reads and corrects the assembly with Illumina reads Assemblies statistics were obtained with quast v5.1 (*80*). We assessed the completeness of each assembly using the Benchmarking of Universal Single-Copy Orthologue (BUSCO) tools with the Sordariomyceta ortholog set (sordariomycetes_odb10, 2020-08-05, n = 3817 orthologous groups searched) (*81*). Gene models were predicted with the Helixer v0.3.1 pipeline (*82*). We also run Helixer pipeline for ESM15 and EP155 strains for the gene

orthology analysis. Statistics of the obtained gene annotation was obtained with the Agat

v1.0.0 tool (*83*). Transposable elements of all genomes were annotated using Repeatmasker

v4-0-7 (*84*) and the customized library built for *C. parasitica* in (*33*) contained in the

"Curated_TEs_database_ESM015_EP155.fa" file (available on the "Portail Data INRAE:

Chromosomal rearrangements but no change of genes and transposable elements repertoires

in an invasive forest-pathogenic fungus" at https://doi.org/10.15454/UTIB8U). The class

Gipsy invader was renamed LTR-Ty3. We filtered out TE copies shorter than 100bp.

Estimation of retrotranposon insertion time and nucleotide divergence time

828
829

To get an estimation of the insertion date of the transposable elements present in the MAT-

proximal non-recombining region, we applied two complementary methods. We first used the

divergence between the LTR sequences in retrotransposons, as these LTR sequences at their

edges are identical at the time of TE insertion and then diverge with time. For this, we used a

de-novo prediction of LTR retrotransposons using LTRharvest GenomeTools 1.6.2 (*85, 86*)

and LTR_Finder v1.07 (*87*). To prepare data for LTRharvest, we first created enhanced suffix

for the M1400 and M6697 genome assemblies using the GenomeTools Suffixerator (-dna -suf

-lcp -bwt -bck -mirrored parameters). We ran LTRharvest using two pipelines, designed to

identify LTR retrotransposons with a minimum and maximum LTR length of 100 bp and

7000 bp respectively and at least 85% identity between two LTR regions, with and without

canonical TGCA motifs, respectively: i) -minlenltr 100 -maxlenltr 7000 -mintsd 4 -maxtsd 6

-similar 85 -vic 10 -seed 20 -motif TGCA -motifmis 1; ii) -minlentltr 100, -maxlenltr 7000, -

mintsd 4, -maxtsd 6, -similar 85, -vic 10, -seed 20. Similarly, we ran LTR_Finder on the

M1400 and M6697 genome assemblies to retrotransposons with both TGCA and non-TGCA

motifs and a minimum and maximum LTR length of 100 bp and 7000 bp respectively and at

least 85% identity between two LTR regions (-D 15000, -d 1000, -l 100, -L 7000, -p 20, -C, -

M 0.85). Finally, we used LTR_retriever v2.8.5 (*88*) with default parameters to filter out false

positive LTR candidates identified by LTRharvest and LTR_Finder and get an estimation of

each element insertion date.

As a second method to estimate the insertion date of the transposable elements present in the

MAT-proximal non-recombining region, we used the set of curated consensus sequences from

(*33*) to annotate the inversion sequence or its surroundings. We first used samtools faidx v1.9

(*70*) to extract the sequence corresponding to inversion and their 50 kb surroundings in

M1400 and M6697 genome assemblies. We annotated the sequences corresponding to the

inversions and the concatenated 50 kb surroundings in both isolates using RepeatMasker version 4.1.5 and rmblast as search engine (v2.10.0) with -no_is -pa 20 -cutoff 250 -a parameters. Finally, we parsed the RepeatMasker .out file using the helped script parseRM_merge_interrupted.pl and omitting Simple_repeat and Low_complexity regions (https://github.com/4ureliek/Parsing-RepeatMasker-Outputs). We then built a summary of the alignments using the RepeatMasker helper script buildSummary.pl and calculated sequence divergence using the calcDivergenceFromAlign.pl script to finally render results with the createRepeatLandscape.pl from the same helper suite.

Comparative genomics analyses

Genome synteny between long reads assemblies were studied using the nucmer v3.1 program (*89*). Outputs were plotted with the R programs ggplot2 (*90*), genoPlotR (*91*) and RIdeogram (*92*).

For MRC010, we investigate the rearrangements by mapping the longest reads (>10 kb) from the ONT fastq files using minimap2 (-ax map-ont option) (*93*). Reads were sorted and indexed with samtools (*70*) and Jbrowse2 (*94*) was used for displaying the pileups. We parsed the reads mapping to the different contigs of the MAT-proximal region (i.e. split reads) and obtained the overlaps again with minimap (-x ava-ont option) and plot with pafr to visually study the pileup of split reads. A total of 56 reads join contigs MRC010_10 and MRC010_15 on the same orientation as presented in Fig 6A. We found no split reads with contig MRC010_01 but we found telomeric repeats in the same orientation on contig MRC010_15 as the telomeric repeats in contig MRC010_01, supporting that mating-type chromosome of MRC010 starts with contig MRC010_01 and finishes with contig MRC010_15. For contig MRC010_14 reads mapping at both ends of this contig also map on contig MRC010_10, indicating it should be in between MRC010_01 and MRC010_10, but its orientation was not resolved.

To study gene disruption, synonymous and non-synonymous divergence (dS and dN) and testing introgression in the MAT-proximal region, we build orthology relationships for genes of the genome assemblies of *C. parasitica* strains M1400, M6697, XIM9508, MRC10, DUM-005, ESM15 and EP155 and included three genomes of the closely related species *Cryphonectria japonica* (IF-6), *Cryphonectria carpinicola* (CS3), and *Cryphonectria*

naterciae (M3664). Genome, gene annotation and species tree of these closely related species

were previously published by (*43*). Genome data were retrieved from NCBI bioproject

number PRJNA644891 and accession IDs JACWRX000000000 for IF-6,

JACWRQ000000000 for CS3 and JACWST000000000 for M3664. We run OrthoFinder

v2.3.7 (*95*) analysis on protein sequences. We used the translatorX v1.1 program (*96*) with

default parameters that use a codon-based approach to align orthologous gene coding

sequences. To compute dS and dN vaues of one-to-one orthologs between pairs of genome

assemblies, we use the yn00 program of PAML (*97*). Estimation of divergence time between

haplotypes was performed using computed gene dS values and the formula Tgenerations =

dS/2μ. We used previous estimates of substitution rates in fungi (*41*, *42*) and considered that

C. parasitica undergoes one generation a year (*98*). We build gene coding sequences trees

with the outgroup genomes in the MAT-proximal region using iqtree2 v2.2.2.6 (*99*) with 1000

bootraps and used the Newick Utilities (https://github.com/tjunier/newick_utils) for

displaying phylogenetics tree.

References

1. S. P. Otto, T. Lenormand, Resolving the paradox of sex and recombination. *Nat. Rev.*
Genet. **3**, 252–261 (2002).

2. T. Schwander, R. Libbrecht, L. Keller, Supergenes and complex phenotypes. *Curr. Biol.*
24, R288–R294 (2014).

3. D. Charlesworth, X. Vekemans, V. Castric, S. Glémin, Plant self-incompatibility
systems: a molecular evolutionary perspective. *New Phytol.* **168**, 61–69 (2005).

4. R. Bergero, D. Charlesworth, The evolution of restricted recombination in sex
chromosomes. *Trends Ecol. Evol.* **24**, 94–102 (2009).

5. J. G. Umen, Evolution of sex and mating loci: An expanded view from Volvocine algae.
Curr. Opin. Microbiol. **14**, 634–641 (2011).

6. D. Charlesworth, The status of supergenes in the 21st century: recombination
suppression in Batesian mimicry and sex chromosomes and other complex adaptations.
Evol. Appl. **9**, 74–90 (2016).

7. Y. Dussert, L. Legrand, I. D. Mazet, C. Couture, M.-C. Piron, R.-F. Serre, O. Bouchez,
P. Mestre, S. L. Toffolatti, T. Giraud, F. Delmotte, Identification of the first oomycete
mating-type locus sequence in the grapevine downy mildew pathogen, *Plasmopara*
viticola. *Curr. Biol.* **30**, 3897-3907.e4 (2020).

8. F. E. Hartmann, M. Duhamel, F. Carpentier, M. E. Hood, M. Foulongne-Oriol, P. Silar,
F. Malagnac, P. Grognet, T. Giraud, Recombination suppression and evolutionary strata

around mating-type loci in fungi: documenting patterns and understanding evolutionary
and mechanistic causes. *New Phytol.* **229**, 2470–2491 (2021).

9.   C. Küpper, M. Stocks, J. E. Risse, N. dos Remedios, L. L. Farrell, S. B. McRae, T. C.
Morgan, N. Karlionova, P. Pinchuk, Y. I. Verkuil, A. S. Kitaysky, J. C. Wingfield, T.
Piersma, K. Zeng, J. Slate, M. Blaxter, D. B. Lank, T. Burke, A supergene determines
highly divergent male reproductive morphs in the ruff. *Nat. Genet.* **48**, 79–83 (2016).

10. Z. Yan, S. H. Martin, D. Gotzek, S. V. Arsenault, P. Duchen, Q. Helleu, O. Riba-
Grognuz, B. G. Hunt, N. Salamin, D. Shoemaker, K. G. Ross, L. Keller, Evolution of a
supergene that regulates a trans-species social polymorphism. *Nat. Ecol. Evol.* **4**, 240–
249 (2020).

11. P. Jay, M. Chouteau, A. Whibley, H. Bastide, H. Parrinello, V. Llaurens, M. Joron,
Mutation load at a mimicry supergene sheds new light on the evolution of inversion
polymorphisms. *Nat. Genet.* **53**, 288–293 (2021).

12. M. Fredericksen, P. D. Fields, L. D. Pasquier, V. Ricci, D. Ebert, QTL study reveals
candidate genes underlying host resistance in a Red Queen model system. *PLOS Genet.*
19, e1010570 (2023).

13. D. Bachtrog, Y-chromosome evolution: emerging insights into processes of Y-
chromosome degeneration. *Nat. Rev. Genet.* **14**, 113–124 (2013).

14. F. Carpentier, R. C. Rodríguez de la Vega, P. Jay, M. Duhamel, J. A. Shykoff, M. H.
Perlin, R. M. Wallen, M. E. Hood, T. Giraud, Tempo of degeneration across
independently evolved nonrecombining regions. *Mol. Biol. Evol.* **39**, msac060 (2022).

15. M. Duhamel, M. E. Hood, R. C. R. de la Vega, T. Giraud, Dynamics of transposable
element accumulation in the non-recombining regions of mating-type chromosomes in
anther-smut fungi. bioRxiv [Preprint] (2023).
https://doi.org/10.1101/2022.08.03.502670.

16. E. L. Berdan, A. Blanckaert, R. K. Butlin, T. Flatt, T. Slotte, B. Wielstra, Mutation
accumulation opposes polymorphism: supergenes and the curious case of balanced
lethals. *Philos. Trans. R. Soc. B Biol. Sci.* **377**, 20210199 (2022).

17. J. E. Ironside, No amicable divorce? Challenging the notion that sexual antagonism
drives sex chromosome evolution. *BioEssays* **32**, 718–726 (2010).

18. B. L. S. Furman, D. C. H. Metzger, I. Darolti, A. E. Wright, B. A. Sandkam, P. Almeida,
J. J. Shu, J. E. Mank, Sex chromosome evolution: So many exceptions to the rules.
Genome Biol. Evol., doi: 10.1093/gbe/evaa081 (2020).

19. P. Jay, E. Tezenas, A. Véber, T. Giraud, Sheltering of deleterious mutations explains the
stepwise extension of recombination suppression on sex chromosomes and other
supergenes. *PLOS Biol.* **20**, e3001698 (2022).

20. B. Charlesworth, The fitness consequences of genetic divergence between polymorphic
gene arrangements. *Genetics*, iyad218 (2023).

21. P. Jay, D. L. Jeffries, F. E. Hartmann, A. Véber, T. Giraud, Why do sex chromosomes
progressively lose recombination? *Trends in Genetics*, in press (2024).

22. T. L. Kubisiak, M. G. Milgroom, Markers linked to vegetative incompatibility (vic)
genes and a region of high heterogeneity and reduced recombination near the mating
type locus (MAT) in *Cryphonectria parasitica*. *Fungal Genet. Biol.* **43**, 453–463 (2006).

23. L. Stauber, T. Badet, A. Feurtey, S. Prospero, D. Croll, Emergence and diversification of
a highly invasive chestnut pathogen lineage across southeastern Europe. *eLife* **10**,
e56279 (2021).

24. L. Stauber, D. Croll, S. Prospero, Temporal changes in pathogen diversity in a perennial
plant–pathogen–hyperparasite system. *Mol. Ecol.* **31**, 2073–2088 (2022).

25. I. C. McGuire, R. E. Marra, B. G. Turgeon, M. G. Milgroom, Analysis of mating-Type
genes in the chestnut blight fungus, *Cryphonectria parasitica*. *Fungal Genet. Biol.* **34**,
131–144 (2001).

26. I. C. McGuire, R. E. Marra, M. G. Milgroom, Mating-type heterokaryosis and selfing in
Cryphonectria parasitica. *Fungal Genet. Biol.* **41**, 521–533 (2004).

27. I. C. McGuire, J. E. Davis, M. L. Double, W. L. MacDonald, J. T. Rauscher, S.
McCawley, M. G. Milgroom, Heterokaryon formation and parasexual recombination
between vegetatively incompatible lineages in a population of the chestnut blight fungus,
Cryphonectria parasitica. *Mol. Ecol.* **14**, 3657–3669 (2005).

28. C. Dutech, O. Fabreguettes, X. Capdevielle, C. Robin, Multiple introductions of
divergent genetic lineages in an invasive fungal pathogen, Cryphonectria parasitica, in
France. *Heredity* **105**, 220–228 (2010).

29. S. L. Anagnostakis, Chestnut blight: The classical problem of an introduced pathogen.
Mycologia (1987).

30. C. Dutech, B. Barrès, J. Bridier, C. Robin, M. G. Milgroom, V. Ravigné, The chestnut
blight fungus world tour: successive introduction events from diverse origins in an
invasive plant fungal pathogen. *Mol. Ecol.* **21**, 3931–3946 (2012).

31. A. Demené, L. Legrand, J. Gouzy, R. Debuchy, G. Saint-Jean, O. Fabreguettes, C.
Dutech, Whole-genome sequencing reveals recent and frequent genetic recombination
between clonal lineages of Cryphonectria parasitica in western Europe. *Fungal Genet.*
Biol. **130**, 122–133 (2019).

32. J. A. Crouch, A. Dawe, A. Aerts, K. Barry, A. C. L. Churchill, J. Grimwood, B. I.
Hillman, M. G. Milgroom, J. Pangilinan, M. Smith, A. Salamov, J. Schmutz, J. S.
Yadav, I. V. Grigoriev, D. L. Nuss, Genome sequence of the chestnut blight fungus
Cryphonectria parasitica EP155: A fundamental resource for an archetypical invasive
plant pathogen. *Phytopathology®* **110**, 1180–1188 (2020).

33. A. Demené, B. Laurent, S. Cros-Arteil, C. Boury, C. Dutech, Chromosomal
rearrangements with stable repertoires of genes and transposable elements in an invasive
forest-pathogenic fungus. *Peer Community J.* **2** (2022).

34. T. L. Kubisiak, M. G. Milgroom, Markers linked to vegetative incompatibility (*vic*) genes and a region of high heterogeneity and reduced recombination near the mating type locus (*MAT*) in *Cryphonectria parasitica. Fungal Genet. Biol. FG B* **43**, 453–463 (2006).

35. C. Dufresnes, A. Borzée, A. Horn, M. Stöck, M. Ostini, R. Sermier, J. Wassef, S. N. Litvinchuck, T. A. Kosch, B. Waldman, Y. Jang, A. Brelsford, N. Perrin, Sex-chromosome homomorphy in palearctic tree frogs results from both turnovers and X-Y recombination. *Mol. Biol. Evol.* **32**, 2328–2337 (2015).

36. M. G. Milgroom, M. L. Smith, M. T. Drott, D. L. Nuss, Balancing selection at nonself recognition loci in the chestnut blight fungus, *Cryphonectria parasitica*, demonstrated by trans-species polymorphisms, positive selection, and even allele frequencies. *Heredity* **121**, 511–523 (2018).

37. Y. Sun, J. Svedberg, M. Hiltunen, P. Corcoran, H. Johannesson, Large-scale suppression of recombination predates genomic rearrangements in *Neurospora tetrasperma. Nat. Commun.* **8**, 1–8 (2017).

38. A. Menkis, D. J. Jacobson, T. Gustafsson, H. Johannesson, The mating-type chromosome in the filamentous ascomycete *Neurospora tetrasperma* represents a model for early evolution of sex chromosomes. *PLOS Genet.* **4**, e1000030 (2008).

39. P. Grognet, F. Bidard, C. Kuchly, L. C. H. Tong, E. Coppin, J. A. Benkhali, A. Couloux, P. Wincker, R. Debuchy, P. Silar, Maintaining two mating types: structure of the mating type locus and its role in heterokaryosis in *Podospora anserina. Genetics* **197**, 421–432 (2014).

40. N. Vittorelli, Hartmann, T. Giraud, Stepwise recombination suppression around the mating-type locus in the fungus *Schizothecium tetrasporum* (Ascomycota, Sordariales). (2022).

41. T. Kasuga, T. J. White, J. W. Taylor, Estimation of nucleotide substitution rates in eurotiomycete fungi. *Mol. Biol. Evol.* **19**, 2318–2324 (2002).

42. J. W. Taylor, M. L. Berbee, Dating divergences in the Fungal Tree of Life: review and new analyses. *Mycologia* **98**, 838–849 (2006).

43. L. Stauber, S. Prospero, D. Croll, Comparative genomics analyses of lifestyle transitions at the origin of an invasive fungal pathogen in the genus *Cryphonectria. mSphere* **5**, 10.1128/msphere.00737-20 (2020).

44. S. Branco, F. Carpentier, R. C. Rodríguez de la Vega, H. Badouin, A. Snirc, S. L. Prieur, M. A. Coelho, D. M. de Vienne, F. E. Hartmann, D. Begerow, M. E. Hood, T. Giraud, Multiple convergent supergene evolution events in mating-type chromosomes. *Nat. Commun.* **9**, 2000 (2018).

45. L. Maloisel, J. L. Rossignol, Suppression of crossing-over by DNA methylation in *Ascobolus. Genes Dev.* **12**, 1381–1389 (1998).

46. F. Boideau, G. Richard, O. Coriton, V. Huteau, C. Belser, G. Deniot, F. Eber, C. Falentin, J. Ferreira de Carvalho, M. Gilet, M. Lodé-Taburel, L. Maillet, J. Morice, G.

Trotoux, J.-M. Aury, A.-M. Chèvre, M. Rousseau-Gueutin, Epigenomic and structural
events preclude recombination in Brassica napus. *New Phytol.* **234**, 545–559 (2022).

47. C. L. Peichel, S. R. McCann, J. A. Ross, A. F. S. Naftaly, J. R. Urton, J. N. Cech, J.
Grimwood, J. Schmutz, R. M. Myers, D. M. Kingsley, M. A. White, Assembly of the
threespine stickleback Y chromosome reveals convergent signatures of sex chromosome
evolution. *Genome Biol.* **21**, 177 (2020).

48. N. Hu, Evolution of a ZW sex chromosome system in willows. *Nat. Commun.* (2023).

49. C. Moraga, C. Branco, Q. Rougemont, P. Veltsos, P. Jedlička, A. Muyle, M. Hanique, E.
Tannier, X. Liu, E. Mendoza-Galindo, C. Lemaitre, P. D. Fields, C. Cruaud, K. Labadie,
C. Belser, J. Briolay, S. Santoni, R. Cegan, R. Linheiro, R. C. R. de la Vega, G. Adam,
A. E. Filali, V. Mossion, A. Boualem, R. Tavares, A. Chebbi, R. Cordaux, C. Fruchard,
D. Prentout, A. Velt, B. Spataro, S. Delmotte, L. Weingartner, H. Toegelová, Z.
Tulpová, P. Cápal, H. Šimková, H. Štorchová, M. Krüger, O. A. J. Abeyawardana, D. R.
Taylor, M. S. Olson, D. B. Sloan, S. Karrenberg, L. F. Delph, D. Charlesworth, T.
Giraud, A. Bendahmane, A. D. Genova, A. Madoui, R. Hobza, G. A. B. Marais, The
Silene latifolia genome and its giant Y chromosome. bioRxiv [Preprint] (2023).
https://doi.org/10.1101/2023.09.21.558754.

50. S. Branco, H. Badouin, R. C. Rodríguez de la Vega, J. Gouzy, F. Carpentier, G.
Aguileta, S. Siguenza, J.-T. Brandenburg, M. A. Coelho, M. E. Hood, T. Giraud,
Evolutionary strata on young mating-type chromosomes despite the lack of sexual
antagonism. *Proc. Natl. Acad. Sci.* **114**, 7067–7072 (2017).

51. S. Fontana, N.-C. Chang, T. Chang, C.-C. Lee, V.-D. Dang, J. Wang, The fire ant social
supergene is characterized by extensive gene and transposable element copy number
variation. *Mol. Ecol.* **29**, 105–120 (2020).

52. O. S. Harringmeyer, H. E. Hoekstra, Chromosomal inversion polymorphisms shape the
genomic landscape of deer mice. *Nat. Ecol. Evol.* **6**, 1965–1979 (2022).

53. K. Huang, K. L. Ostevik, C. Elphinstone, M. Todesco, N. Bercovich, G. L. Owens, L. H.
Rieseberg, Mutation load in sunflower inversions is negatively correlated with inversion
heterozygosity. *Mol. Biol. Evol.* **39**, msac101 (2022).

54. M. Matschiner, J. M. I. Barth, O. K. Tørresen, B. Star, H. T. Baalsrud, M. S. O. Brieuc,
C. Pampoulie, I. Bradbury, K. S. Jakobsen, S. Jentoft, Supergene origin and maintenance
in Atlantic cod. *Nat. Ecol. Evol.* **6**, 469–481 (2022).

55. J. Hill, E. D. Enbody, H. Bi, S. Lamichhaney, W. Lei, J. Chen, C. Wei, Y. Liu, D.
Schwochow, S. Younis, F. Widemo, L. Andersson, Low mutation load in a supergene
underpinning alternative male mating strategies in ruff (*Calidris pugnax*). *Mol. Biol.*
Evol. **40**, msad224 (2023).

56. P. Jay, A. Whibley, L. Frézal, M. Á. Rodríguez de Cara, R. W. Nowell, J. Mallet, K. K.
Dasmahapatra, M. Joron, Supergene evolution triggered by the introgression of a
chromosomal inversion. *Curr. Biol.* **28**, 1839-1845.e3 (2018).

57. Q. Helleu, C. Roux, K. G. Ross, L. Keller, Radiation and hybridization underpin the
spread of the fire ant social supergene. *Proc. Natl. Acad. Sci.* **119**, e2201040119 (2022).

58. E. Stolle, R. Pracana, F. López-Osorio, M. K. Priebe, G. L. Hernández, C. Castillo-
Carrillo, M. C. Arias, C. I. Paris, M. Bollazzi, A. Priyam, Y. Wurm, Recurring adaptive
introgression of a supergene variant that determines social organization. *Nat. Commun.*
13, 1180 (2022).

59. P. Jay, T. G. Aubier, M. Joron, The interplay of local adaptation and gene flow may lead
to the formation of supergenes. *Mol. Ecol.*, e17297 (2024).

60. G. H. Choi, D. L. Nuss, Hypovirulence of chestnut blight fungus conferred by an
infectious viral cDNA. *Science* **257**, 800–803 (1992).

61. J. Brusini, M. L. Wayne, A. Franc, C. Robin, The impact of parasitism on resource
allocation in a fungal host: the case of *Cryphonectria parasitica* and its mycovirus,
Cryphonectria Hypovirus 1. *Ecol. Evol.* **7**, 5967–5976 (2017).

62. A. Tellier, S. Moreno-Gámez, W. Stephan, Speed of adaptation and genomic footprints
of host–parasite coevolution under arms race and trench warfare dynamics. *Evolution*
68, 2211–2224 (2014).

63. A. L. Bazzicalupo, F. Carpentier, S. P. Otto, T. Giraud, Little evidence of antagonistic
selection in the evolutionary strata of fungal mating-type chromosomes (*Microbotryum*
lychnidis-dioicae). *G3 Genes Genomes Genet.* **9**, 1987–1998 (2019).

64. J. Gutiérrez-Valencia, P. W. Hughes, E. L. Berdan, T. Slotte, The genomic architecture
and evolutionary fates of supergenes. *Genome Biol. Evol.* **13** (2021).

65. P. J. Hoegger, D. Rigling, O. Holdenrieder, U. Heiniger, Genetic structure of newly
established populations of *Cryphonectria parasitica*. *Mycol. Res.* **104**, 1108–1116
(2000).

66. M. G. Milgroom, K. Wang, Y. Zhou, S. E. Lipari, S. Kaneko, Intercontinental
population structure of the chestnut blight fungus, *Cryphonectria parasitica*. *Mycologia*
88, 179–190 (1996).

67. Y. C. Liu, D. Linder-Basso, B. I. Hillman, S. Kaneko, M. G. Milgroom, Evidence for
interspecies transmission of viruses in natural populations of filamentous fungi in the
genus Cryphonectria. *Mol. Ecol.* **12**, 1619–1628 (2003).

68. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

69. B. Langmead, C. Trapnell, M. Pop, S. L. Salzberg, Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. *Genome Biol.* **10**, R25 (2009).

70. H. Li, B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G.
Abecasis, R. Durbin, The Sequence Alignment/Map format and SAMtools.
Bioinformatics **25**, 2078–2079 (2009).

71. A. McKenna, M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis, A. Kernytsky, K.
Garimella, D. Altshuler, S. Gabriel, M. Daly, M. A. DePristo, The Genome Analysis
Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data.
Genome Res. **20**, 1297–1303 (2010).

72.   P. Danecek, A. Auton, G. Abecasis, C. A. Albers, E. Banks, M. A. DePristo, R. E.
Handsaker, G. Lunter, G. T. Marth, S. T. Sherry, G. McVean, R. Durbin, 1000 Genomes
Project Analysis Group, The variant call format and VCFtools. *Bioinformatics* **27**, 2156–
2158 (2011).

73.   J.-H. Shin, S. Blay, B. McNeney, J. Graham, others, LDheatmap: an R function for
graphical display of pairwise linkage disequilibria between single nucleotide
polymorphisms. *J. Stat. Softw.* **16**, 1–10 (2006).

74.   S. Purcell, B. Neale, K. Todd-Brown, L. Thomas, M. A. R. Ferreira, D. Bender, J.
Maller, P. Sklar, P. I. W. de Bakker, M. J. Daly, P. C. Sham, PLINK: A tool set for
whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.*
81, 559–575 (2007).

75.   B. Pfeifer, U. Wittelsbürger, S. E. Ramos-Onsins, M. J. Lercher, PopGenome: an
efficient Swiss army knife for population genomic analyses in R. *Mol. Biol. Evol.* **31**,
1929–1936 (2014).

76.   D. H. Huson, SplitsTree: analyzing and visualizing evolutionary data. *Bioinforma. Oxf.*
Engl. **14**, 68–73 (1998).

77.   H. E. L. Lischer, L. Excoffier, PGDSpider: an automated data conversion tool for
connecting population genetics and genomics programs. *Bioinformatics* **28**, 298–299
(2012).

78.   S. Koren, B. P. Walenz, K. Berlin, J. R. Miller, N. H. Bergman, A. M. Phillippy, Canu:
scalable and accurate long-read assembly via adaptive k-mer weighting and repeat
separation. *Genome Res.* **27**, 722–736 (2017).

79.   D. Antipov, A. Korobeynikov, J. S. McLean, P. A. Pevzner, hybridSPAdes: an algorithm
for hybrid assembly of short and long reads. *Bioinformatics* **32**, 1009–1015 (2016).

80.   A. Gurevich, V. Saveliev, N. Vyahhi, G. Tesler, QUAST: quality assessment tool for
genome assemblies. *Bioinformatics* **29**, 1072–1075 (2013).

81.   M. Manni, M. R. Berkeley, M. Seppey, F. A. Simão, E. M. Zdobnov, BUSCO update:
Novel and streamlined workflows along with broader and deeper phylogenetic coverage
for scoring of eukaryotic, prokaryotic, and viral genomes. *Mol. Biol. Evol.* **38**, 4647–
4654 (2021).

82.   F. Holst, A. Bolger, C. Günther, J. Maß, S. Triesch, F. Kindel, N. Kiel, N. Saadat, O.
Ebenhöh, B. Usadel, R. Schwacke, M. Bolger, A. P. M. Weber, A. K. Denton, Helixer–
de novo prediction of primary Eukaryotic gene models combining deep learning and a
hidden markov model. bioRxiv [Preprint] (2023).
https://doi.org/10.1101/2023.02.06.527280.

83.   J. Dainat, D. Hereñú, P. Pucholt, AGAT:Another Gff Analysis Toolkit to handle
annotations in any GTF. *GFF Format* (2020).

84.   A. F. A. Smit, R. Hubley, P. Green, RepeatMasker Open-4.0. Available at:
http://www.repeatmasker.org Accessed: November 30, 2017. (2013).

85. D. Ellinghaus, S. Kurtz, U. Willhoeft, LTRharvest, an efficient and flexible software for
de novo detection of LTR retrotransposons. *BMC Bioinformatics* **9**, 18 (2008).

86. G. Gremme, S. Steinbiss, S. Kurtz, GenomeTools: A Comprehensive Software Library
for Efficient Processing of Structured Genome Annotations. *IEEE/ACM Trans. Comput.*
Biol. Bioinform. **10**, 645–656 (2013).

87. Z. Xu, H. Wang, LTR_FINDER: an efficient tool for the prediction of full-length LTR
retrotransposons. *Nucleic Acids Res.* **35**, W265–W268 (2007).

88. S. Ou, N. Jiang, LTR_retriever: A highly accurate and sensitive program for
identification of long terminal repeat retrotransposons. *Plant Physiol.* **176**, 1410–1422
(2018).

89. G. Marçais, A. L. Delcher, A. M. Phillippy, R. Coston, S. L. Salzberg, A. Zimin,
MUMmer4: A fast and versatile genome alignment system. *PLOS Comput. Biol.* **14**,
e1005944 (2018).

90. H. Wickham, *Ggplot2: Elegant Graphics for Data Analysis* (Springer Science &
Business Media, 2009).

91. L. Guy, J. Roat Kultima, S. G. E. Andersson, genoPlotR: comparative gene and genome
visualization in R. *Bioinformatics* **26**, 2334–2335 (2010).

92. Z. Hao, D. Lv, Y. Ge, J. Shi, D. Weijers, G. Yu, J. Chen, RIdeogram: drawing SVG
graphics to visualize and map genome-wide data on the idiograms. *PeerJ Comput. Sci.*
6, e251 (2020).

93. H. Li, Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**,
3094–3100 (2018).

94. C. Diesh, G. J. Stevens, P. Xie, T. De Jesus Martinez, E. A. Hershberg, A. Leung, E.
Guo, S. Dider, J. Zhang, C. Bridge, G. Hogue, A. Duncan, M. Morgan, T. Flores, B. N.
Bimber, R. Haw, S. Cain, R. M. Buels, L. D. Stein, I. H. Holmes, JBrowse 2: a modular
genome browser with views of synteny and structural variation. *Genome Biol.* **24**, 74
(2023).

95. D. M. Emms, S. Kelly, OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biol.* **20**, 238 (2019).

96. F. Abascal, R. Zardoya, M. J. Telford, TranslatorX: multiple alignment of nucleotide
sequences guided by amino acid translations. *Nucleic Acids Res.* **38**, W7-13 (2010).

97. Z. Yang, PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* **24**,
1586–1591 (2007).

98. L. Guerin, G. Froidefond, X.-M. Xu, Seasonal patterns of dispersal of ascospores of
Cryphonectria parasitica (chestnut blight). *Plant Pathol.* **50**, 717–724 (2001).

99. B. Q. Minh, H. A. Schmidt, O. Chernomor, D. Schrempf, M. D. Woodhams, A. von
Haeseler, R. Lanfear, IQ-TREE 2: New models and efficient methods for phylogenetic
inference in the genomic era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

1198

1199

Acknowledgments

This work was supported by the European Research Council (ERC) EvolSexChrom (832352) grant and a Fondation Louis D grant from the Institut de France to TG, and by the ANR PIA grant # ANR-20-IDEES-0002 to F.E.H. The authors declare that they have no competing interests.

TG and FEH conceptualized the study and acquired funding. TG supervised the study. AL, AS, CD, AD, SP, DC, LS and TB contributed to data and their availability. AL and AS performed *in vitro* culture and DNA extraction. FEH, RRDLV, QR, LS and TB analyzed genomes. FEH and TG wrote the original draft. All authors edited the manuscript.

The raw data and the new assemblies produced in this study will be published pending scientific review.

Tables

Table 1: Distribution of mating types among non-recombining haplotypes of the MAT-proximal region in the European invasive *Cryphonectria parasitica* CL1 genetic cluster and the Swiss populations.

Figures

Figure 1: Linkage disequilibrium (LD) analysis along the contig carrying the mating-type locus in a *Cryphonectria parasitica* European invasive population. LD heatmaps using single nucleotide polymorphisms (SNPs; n=3815) located on the contig carrying the mating-type locus (scaffold_2 of the EP155 genome) in the CL1 genetic cluster (European invasive population introduced from North America); pairs of SNPs with high LD, i.e. r^2 >0.9, correspond to the red triangle. The mating-type locus location is shown with a green triangle and the MAT-proximal region lacking recombination is shown with a red arrow. The two high-LD blocks within the MAT-proximal region are shown with orange arrows. SNPs at the limit of the MAT-proximal region and the two high LD blocks were manually highlighted with red dotted lines and an orange line.

Figure 2: Genetic structure using single nucleotide polymorphisms (SNPs) in the MAT-proximal region lacking recombination and other genomic regions in a *Cryphonectria*

parasitica **European invasive population. A-C.** Principal component analysis (PCA). Two principal components are presented. Percentage of variance explained by each PC is indicated into brackets. Strains are colored according to their mating type (*MAT1-1* or *MAT1-2*). **D-E.** Neighbor-net tree from a SplitsTree analysis. These analyses were performed in the CL1 genetic cluster (European invasive populations introduced from North America) based on: **A and D.** SNPs (n=2,220) located within the MAT-proximal region along the contig carrying the mating-type locus (scaffold_2 of the EP155 genome).; **B.** SNPs (n=1,595) located in other regions along the contig carrying the mating-type locus; **C and E** SNPs (n=11,289) located on other contigs of the EP155 genome. On panels A and D, the two clusters corresponding to the MAT-Prox1 and MAT-Prox2 haplotypes are shown with red circles. The identified haplotype of each strain is indicated in Table S1. The number of strains within the MAT-Prox1 haplotype is indicated on panel A by the letter n. The newly sequenced M1400 and M6697 strains are highlighted with a black rectangle

Figure 3: Synteny and rearrangements between the two newly sequenced genomes of European invasive strains introduced from North America (M1400 and M6697) in the chromosome carrying the MAT-proximal region lacking recombination in *Cryphonectria parasitica*. The mating-type locus is located with a green diamond. The MAT-proximal region defined from LD analyses is indicated with red arrows.

Figure 4: Genetic diversity and divergence between non-recombining haplotypes of the MAT-proximal region along the contig carrying the mating-type locus in a *Cryphonectria parasitica* European invasive population. A-B. GC content (A) and transposable element densities (B) along the contig carrying the mating-type locus (tig_0000001) of the M1400 *C. parasitica* genome. C. Relative divergence (F_{ST}) between strains of the MAT-Prox1 and MAT-Prox2 haplotypes: D. Absolute divergence d_{XY} between strains of the two MAT-Proximal haplotypes. E-F. Nucleotide diversity within pools of strains for each MAT-Proximal haplotype; G-H-I. Tajima's D for all strains pooled and within pools of strains for each MAT-Proximal haplotype; The MAT-proximal region defined from LD analyses and the inversion between M1400 and M6697 genomes are indicated with red and blue arrows respectively. The mating-type locus location is shown with a green triangle. The location of the putative centromere is indicated by a yellow circle. All population statistics were computed for the 1990 Swiss population along the mating-type contig (tig_0000001) of

1266 the M1400 genome per 50-kb window overlapping over 10 kbp. Windows containing fewer

1267 than 5 SNPs were removed from the analysis.

1268

Figure 5: Annotation of transposable elements (TEs) and estimates of their insertion

time in the MAT-proximal region lacking recombination and other genomic regions in

Cryphonectria parasitica. A. Annotation of transposable elements (TEs) and gene density

along the MAT-proximal region in M1400 and M6697 genomes. The figure only shows the

genomic region from 7 Mb on tig_0000001 of the M1400 genome and from 2.8 Mb on

tig_0000060 of the M6697 genome. The mating-type locus is located with a green triangle.

The two high-LD blocks within the MAT-proximal region are shown with orange arrows.

Synteny for 10 kb segments with identity > 90 % is shown in red and inversion in blue.

Transposons larger than 5 kb are shown in purple for the transposons annotated as LTR-Ty3

by and in green for other transposons. Orthologous genes shared between genomes are shown

in black and unique to each each genome in grey. B. Relative percentage of each TE

annotation (in %) in the non-recombining MAT-proximal region and other recombining

regions of M6697 and M1400 genomes. C. Pairwise genetic distance between TE copies

within the inversion and around (50 kb) in both M1400 and M6697 genomes. Distribution of

the Kimura substitution levels computed using the consensus sequence for the TEs. D.

Estimates of the age of intact copies of LTR within and around the inversion.

1285

Figure 6: Transposable elements content and gene content variation in seven high

quality genome assemblies from other *Cryphonectria parasitica* populations from the

native and the introduced range. A. TE load (percentage of base pairs occupied by

transposable elements) in the seven high quality genome assemblies in the MAT proximal

region and other recombining regions. **B.** Number of orthologous genes present in the seven

high quality genome assemblies. Only single copy orthologous genes shared between M6697

and M1400 or present only in one this genome are shown.

1293

Figure 7: Genetic structure using single nucleotide polymorphisms (SNPs) in the MAT-

proximal region lacking recombination (A) and other genomic regions (B) in other

resequenced *Cryphonectria parasitica* populations from the native and the introduced

range. Neighbor-net tree from a SplitsTree analysis based on: **A** SNPs (n=4,120) located

within the MAT-proximal region along the contig carrying the mating-type locus (scaffold_2

of the EP155 genome); **B** SNPs (n= 103,058) located on other contigs of the EP155 genome.

Color of empty circles around strain ID indicate the genetic clusters the strains belong to
(CL1, CL2, CL3, CL4; n= 33 strains). Dotted circles indicate strains of the native range and
plain circles indicate strains of the introduced range. Red and blue dots near strain ID indicate
the mating type (*MAT-1* or *MAT-2*, respectively).
1304
1305

1306　Table 1

1307

	Population	CL1 genetic cluster		Swiss 1990 population		Swiss 2019 population	
	PCA cluster	PCA cluster 1 (PC>0)	PCA cluster 2 (PC<0)	PCA cluster 1 (PC>0)	PCA cluster 2 (PC<0)	PCA cluster 1 (PC>0)	PCA cluster 2 (PC<0)
	Haplotype (based on PCA cluster)	MAT-Prox1	MAT-Prox2	MAT-Prox1	MAT-Prox2	MAT-Prox1	MAT-Prox2
Number of strains in each PCA cluster	total	36	52	20	51	8	54
	mat-1	30	11	18	17	5	15
	mat-2	6	41	2	34	3	39
% of strains in each PCA cluster	mat-1	83	21	90	33	63	28
	mat-2	17	79	10	67	38	72

1308

1309

1310

1311   Fig 1

1312

1313
1314

1315   Fig 2

1316

CL1 genetic cluster (European invasive population)

1317
1318

1319
1320

1321   Fig 3

1322

1323

1324

1325   Fig 4

1326

1327

1328   Fig5

1329

1330

1331

1332   Fig 6

1333

1334

1335   Fig 7

1336

1337

1338

1339

1340

1341