1

An old inversion polymorphism involving giant mobile elements in an invasive fungal

pathogen

Fanny E. Hartmann[1]*, Ricardo C. Rodriguez de la Vega[1], Arthur Demené[2], Thomas Badet[3],
Jean-Philippe Vernadet[1], Quentin Rougemont[1], Amandine Labat[1], Alodie Snirc[1], Lea
Stauber[5], Daniel Croll[3], Simone Prospero[4], Cyril Dutech[2], Tatiana Giraud[1]

[1] Ecologie Systematique et Evolution, CNRS, Université Paris-Saclay, AgroParisTech, 91198 Gif-sur-
Yvette, France

[2] BIOGECO, INRAE, Univ. Bordeaux, 69 route d'Arcachon, Cestas F-33610, France.

[3] Laboratory of Evolutionary Genetics, Institute of Biology, University of Neuchâtel, Neuchâtel,
Switzerland

[4] Swiss Federal Research Institute WSL, Birmensdorf, Switzerland

[5] Institute for Infectious Diseases, University of Bern, Switzerland

*Corresponding author: Fanny E. Hartmann

E-mail: fanny.hartmann@universite-paris-saclay.fr

Key words: recombination suppression, fungi, mating-type chromosome, inversion,

deleterious allele sheltering, supergenes, evolutionary strata, sex chromosomes, starships

Abstract

Recombination suppression can evolve in sex or mating-type chromosomes, or in autosomal supergenes, with different haplotypes being maintained by balancing selection. In the invasive chestnut blight fungus *Cryphonectria parasitica*, a genomic region was suggested to lack recombination and to be partially linked to the mating-type (MAT) locus based on segregation analyses. Using hundreds of available *C. parasitica* genomes and generating new high-quality genome assemblies, we show that a ca. 1.2 Mb genomic region proximal to the mating-type locus lacks recombination, with the segregation of two highly differentiated haplotypes in balanced proportions in invasive populations. High-quality genome assemblies further revealed an inversion in one of the haplotypes in the invaded range. The two haplotypes were estimated to have diverged 1.5 million years ago, and each harboured specific genes, some of which likely belonging to *Starship* elements, that are large mobile elements, mobilized by tyrosine recombinases, able to move accessory genes, and involved in adaptation in multiple fungi. The MAT-proximal region carried genes upregulated under virus infection or vegetative incompatibility reaction. In the native range, the MAT-proximal region also appeared to have a different evolutionary history than the rest of the genome. In all continents, the MAT-Proximal region was enriched in non-synonymous substitutions, in gene presence/absence polymorphism, in tyrosine recombinases and in transposable elements. This study thus sheds light on a case of a large non-recombining region partially linked to a mating compatibility locus, with likely balancing selection maintaining differentiated haplotypes, possibly involved in adaptation in a devastating tree pathogen.

Introduction

Recombination increases the efficiency of selection, the purging of deleterious alleles and the generation of potentially beneficial allelic combinations (Otto and Lenormand 2002). However, recombination can also break up beneficial allelic combinations and, therefore, can be selected against, generating supergenes, i.e., large regions without recombination encompassing multiple genes (Schwander et al. 2014). This is often the case in genomic regions controlling mating compatibility, such as regions determining sex, mating type or self-incompatibility in plants, animals, fungi, algae and oomycetes (Charlesworth et al. 2005; Bergero and Charlesworth 2009; Umen 2011; Charlesworth 2016; Dussert et al. 2020; Hartmann et al. 2021). Such recombination suppression keeps alleles linked across different genes that prevent self-compatibility or intermediate sexual phenotypes, and can sometimes extend away from sex-determining genes or mating-type loci (Bergero and Charlesworth 2009; Furman et al. 2020; Hartmann et al. 2021). Even beyond sex and mating-type chromosomes, there is an increasing number of supergene reports in autosomes (Schwander et al. 2014) with highly differentiated haplotypes maintained by balancing selection. Striking examples include supergenes controlling social structure in ants, wing color patterns in butterflies, reproductive morphs in birds or host-parasite interactions (Küpper et al. 2016; Yan et al. 2020; Jay et al. 2021; Fredericksen et al. 2023).

In the long term, however, because of less efficient selection, genomic regions without recombination show signs of degeneration, such as transposable element accumulation, rearrangements, non-synonymous substitutions, decreased gene expression, reduced frequency of optimal codons and gene losses (Bachtrog 2013; Carpentier et al. 2022; Duhamel et al. 2023). The non-recombining regions can nevertheless be maintained by balancing selection, with different allelic combinations having contrasting advantages in different situations or experiencing negative frequency dependent selection (Jay et al. 2021; Berdan et al. 2022). The evolutionary mechanisms leading to recombination suppression and allowing their persistence remain however poorly known (Jay, D. Jeffries, Hartmann, et al. 2024). Reporting more diverse constellations of recombination suppression in a variety of organisms with contrasting life-history traits is important for understanding the general patterns of recombination suppression, their evolution and maintenance (Ironside 2010; Charlesworth 2016; Furman et al. 2020; Hartmann et al. 2021).

In the invasive chestnut blight fungal pathogen *Cryphonectria parasitica*, mating compatibility is determined by a mating-type locus that displays two alleles, called *MAT-1* and *MAT-2,* that are both permanently heterozygous in the diploid and dikaryotic stages (McGuire et al. 2001). A large region without recombination was suggested to occur near the mating-type locus, although not completely linked to it (*i.e.,* at 3.9 cM of the mating-type locus), based on segregation of RAPD and RFLP markers in three crosses involving five different parents, from the USA, Japan or Italy (Kubisiak and Milgroom 2006a). Recent genome-wide association studies in a worldwide collection and in local *C. parasitica* populations in southern Switzerland found significant association of SNPs with the mating-type locus across a large region (>1 Mb), further suggesting the existence of reduced recombination in this region (Stauber et al. 2021; Stauber et al. 2022). This genomic region was enriched in SNPs, transposable elements and copy-number variants, such as deletions, which are consistent with sequence degeneration and a lack of recombination (Stauber et al. 2021), and which prevented its assembly so far.

The evolutionary history of the species is well documented as it is an invasive and highly damaging pathogen, having almost caused the extinction of American chestnut (*Castanea dentata*) in North America (Anagnostakis 1987). From its center of origin in Asia and its original hosts, the Chinese chestnut (*Ca. mollissima*) and the Japanese chestnut (*Ca. crenata*), *C. parasitica* has first invaded North America, killing millions of American chestnuts (*Ca. dentata*). In Europe, at least two distinct introduction events occurred, on the European chestnut (*Ca. sativa*): one introduction from North America to Italy, and one directly from Asia, probably to the Pyrenees Mountains. European strains will be therefore hereafter referred to as invasive European strains introduced from North America or invasive European strains introduced directly from Asia. The chestnut blight symptoms have been less severe in Europe than in North America, due to lower susceptibility of *Ca. sativa* and a virus infecting *C. parasitica* and causing hypovirulence (Dutech et al. 2010; Dutech et al. 2012). The genetic determinants of the adaptation of the pathogen to its new environments and hosts remain largely unknown (Lovat and Donnelly 2019).

Lineages with an apparently clonal structure have been identified in invasive populations while sexually reproducing populations occur both in the native and invaded ranges (Dutech et al. 2012; Stauber et al. 2021). The lineages with a predominant clonal structure likely still

undergo rare sex events, as shown by the presence of the two mating-type alleles in most of
them (Demené et al. 2019). Furthermore, although this ascomycete fungus is mostly found as
haploid mycelia, some isolates have been reported to be heterokaryotic at the mating-type
locus in several invasive European and North American populations, i.e., with cells carrying
different nuclei, of the opposite mating types *MAT-1* and *MAT-2* (McGuire et al. 2004;
McGuire et al. 2005; Dutech et al. 2010; Stauber et al. 2021; Stauber et al. 2022).

Here, we therefore studied the occurrence of recombination suppression in the genomic region
proximal to the mating-type locus in *C. parasitica,* using the previously published genomes
and further generated six new high-quality long-read based genome assemblies of strains from
the native and invaded range of the pathogen. More specifically, we tested whether
recombination was completely suppressed in the genomic region proximal to the mating-type
locus in *C. parasitica*. We first analyze sexually reproducing invasive European populations,
for which extensive genomic datasets are available in local populations. We assessed, in these
populations, the level of differentiation between haplotypes, as well as the frequency of the
two non-recombining haplotypes and their association with mating types. We investigated
whether other regions of the genome showed reduced recombination rates and whether
genomic footprints of degeneration were present in the non-recombining region, such as
genomic rearrangements, non-synonymous substitutions and transposable element
accumulation. We also tested whether this non-recombining region has been gradually
expanding, estimated the age of recombination suppression and investigated whether the
predicted gene functions in the MAT-proximal region could help understanding the
evolutionary cause for recombination suppression, polymorphism maintenance and partial
linkage to the mating-type locus. We looked in particular for *Starships*, that are giant mobile
elements recently discovered in ascomycete fungi and able to move accessory genes as cargo
within and between genomes (Gluck-Thaler et al. 2022; Urquhart et al. 2024). These cargo
genes can be involved in adaptation (Gluck-Thaler et al. 2022; Urquhart et al. 2024), as
shown in fungal pathogens of coffee (Peck et al. 2023) and of wheat (Bucknell et al. 2024;
Tralamazza et al. 2024), as well as in molds used for making cheeses (Cheeseman et al. 2014;
Ropars et al. 2015) and those maturing dry-cured meat (Lo et al. 2023). *Starships* are
characterized by their captain, a tyrosine recombinase with a DUF3435 domain, being the first
gene at the 5' edge of the elements and allowing their excision and insertion. *Starships* can
contain "cargo" genes, highly variable in nature and number among *Starship* elements
(Urquhart et al. 2024). *Starship* diversity is partitioned into 11 major families, targeting

specific genomic niches for their insertions, e.g., AT-rich regions (Gluck-Thaler et Vogan, 2024). We then extended the analyses to other populations with less extensive genomic resources, i.e. other invasive populations and native Asian populations, investigating the presence of recombination suppression footprints and the occurrence of the two differentiated haplotypes worldwide, the presence of *Starships*, as well as the possibility of introgression, using available genomes of closely related species, pathogenic and non-pathogenic (Stauber et al. 2021). Finally, we used available expression data to investigate whether the MAT-Proximal region carried genes upregulated under infection by the virus responsible for hypovirulence or during vegetative incompatibility reactions, a phenomenon avoiding hyphal fusions between individuals, considered to protect against virus transmission.

Results

Footprints of recombination suppression in a large region (> 1 Mb) proximal to the mating-type locus in sexually reproducing invasive populations

We first performed population genomic analyses on available genome sequences within local populations sampled in southern Switzerland over two temporal frames (early 1990 and 2019), and within the more broadly geographically distributed CL1 cluster in central and southeastern Europe, both having a recombining genetic structure and being invasive European populations introduced from North America (Table S1 (Stauber et al. 2021)). We studied only monokaryotic strains, i.e. having either *MAT-1* or *MAT-2* in their genomes, for phasing haplotypes and performed stringent SNPs filtering by masking repeats on the EP155 reference genome (Crouch et al. 2020) and removing missing data and rare variants to ensure robustness of our analyses. We identified 8,900 SNPs segregating among 71 strains of the 1990 Swiss population (35 MAT-1; 36 MAT-2; Table S1), 9,646 SNPs segregating among 62 strains of the Swiss 2019 population (20 MAT-1; 42 MAT-2; Table S1) and 15,104 SNPs segregating among 88 strains of the CL1 cluster (41 MAT-1; 47 MAT-2; Table S1), which represent an average density of about 20 SNPs/100 kb, in agreement with previous studies in this invasive fungus (Stauber et al. 2021; Stauber et al. 2022). We investigated the genome-wide linkage disequilibrium (LD) landscape and looked for large blocks (>1 Mb) of high LD (r2>0.9) among SNPs within contigs, to look for signatures of reduced recombination along the genome. LD indeed decays by half across a few 100 kb on average in *C. parasitica* in recombining regions of the genome (Demené et al. 2019). Therefore, high LD beyond >1 Mb represents strong evidence of recombination cessation and the SNP density is sufficient to study such LD variation.

We found a large block of high LD on the chromosome called scaffold_2 of the EP155 genome in the three populations, corresponding to the mating-type chromosome (see Fig 1 for the CL1 population, the pattern in the other populations being similar). The high-LD block was located near the mating-type locus but did not encompass the mating-type locus itself and will be called hereafter the "MAT-proximal region". In the 1990 and 2019 Swiss populations, SNPs between 0.493 Mb and 1.720 Mb in the mating-type chromosome were all in high LD (with $r^2>0.9$ between SNPs distant by up to 1 Mb; in this region, mean r^2 of all SNP pairs in the Swiss 1990 population = 0.89; mean r^2 of all SNP pairs in the Swiss 2019 population = 0.83; Fig S1). In the CL1 cluster, the large high-LD block was also present, but was split into two blocks, a large one and a smaller, peripheral one (orange arrows in Fig 1). We indeed detected values of $r^2>0.9$ between SNPs distant by up to 1 Mb, between 0.532 Mb and 1.535 Mb (mean r^2 of all SNP pairs = 0.94). Between 1.535 and 1.717 Mb, SNPs were also in high LD (mean r^2 = 0.84). The LD between the two high-LD blocks was lower ($r^2= 0.59$) than within blocks but was still higher than elsewhere along the genome (orange rectangle at the right border of the red triangle; Fig 1). A similar pattern also existed in the Swiss populations (orange rectangle at the right border of the red triangle; Fig S1), although less marked. This is likely due to rare recombination events at one specific locus near the edge of the large fully non-recombining region, lowering the LD level in populations between the two parts of the otherwise fully non-recombining region. The mating-type locus was located at 1.737 Mb in the EP155 genome, distant by 17 kb and 20 kb from the LD block in the Swiss populations and CL1 cluster, respectively. SNP density was on average 48 SNPs/100 kb in the high-LD block region, i.e. higher than in the rest of the chromosome (Fig S2C). When sampling SNPs distant of at least 50 kb in this region to have the same SNP density all over the chromosome, we also detected the high-LD block, which indicates that the SNP density did not generated biases in the LD pattern (Fig S2D).

We found no other regions of the genomes with $r^2>0.9$ between SNPs across such a large genomic distance in any population. On scaffold_6 in the CL1 cluster, two smaller SNP blocks had a $r^2>0.9$ between each other despite being distant (Fig S2C; between SNPs blocks at 3-20 kb and 2.151-2.278 kb), which is likely due to the major intra-scaffold translocation described in this region between the EP155 and ESM15 strains (Demené et al. 2022). The maximum size of the blocks with $r^2>0.9$ on other scaffolds ranged from 135 to 540 kb in the CL1 cluster, as previously described (Demené et al. 2019). The average distance between

SNPs in high LD ($r^2 > 0.9$) across the genome was higher on the scaffold_2 than on other
scaffolds (in the CL1 cluster, pairwise Wilcoxon test p-value < 2e-16 with Bonferroni
correction). Furthermore, the proportion of SNPs in high LD ($r^2 > 0.9$) across the genome was
also much higher on the scaffold_2 than on other scaffolds due to the high LD in the MAT-
proximal region (Fig S2D, Table S2). Therefore, the large (1Mb) and localized block with
maximal LD values proximal to the mating-type locus both in the Swiss populations and the
CL1 genetic cluster stands out as exceptional in the genome and indicates full recombination
cessation, which supports previous inferences from progeny segregation analyses (Kubisiak
and Milgroom 2006b). Indeed, even low rates of recombination homogenize allelic
frequencies and prevents LD building (Dufresnes et al. 2015). The MAT-proximal non-
recombining region was actually larger (1.23 Mb), but with likely rare recombination events
at one precise locus near its edge, lowering LD between its two fully non-recombining parts.

Consistent with a lack of recombination, the MAT-proximal region formed two genetically
highly differentiated haplotypes in these invasive Swiss and CL1 populations (Fig 2A; Fig S3
A(1)-C(1)), as shown by the two clusters on the principal component analysis (PCA) using
only SNPs located in the MAT-proximal region, while no structure was detected in the rest of
the genome (Fig 2B-C; Fig S3A(2)-C(2)). The neighbor-net networks further supported the
existence of two differentiated haplotypes in the MAT-Proximal region, contrasting with an
otherwise recombining structure genome-wide (Fig 2D-E; Fig S3B-D). In the Swiss 2019
population, a few reticulations between haplotypes were found and three strains (LU3, Nov10,
Nov4) appeared to have an intermediate sequence in the MAT-Proximal region between the
two haplotypes (Fig S3D). This intermediate haplotype was also found in a few other invasive
strains introduced directly from Asia (see below).

The two genetic clusters in the MAT-proximal region appeared associated with mating types
in the CL1 and Swiss 1990 recombining populations (Figs 2A and D; Fig S3, Table 1).
Indeed, the distribution of mating types between the two PCA clusters strongly deviated from
expectations under random association (Table 1; chi-squared test: $\chi^2 = 33.3$; p-value = 5.876e-
08 in CL1; $\chi^2 = 18.76$; p-value = 8.441e-05 in Swiss 1990). We thus hereafter call the two
differentiated haplotypes MAT-Prox1 and MAT-Prox2, referring to the mating-type
association. However, the association between mating types and MAT-Proximal haplotypes
was not complete, with for example only 83% of MAT1-1 strains carrying the MAT-Prox1
haplotype in CL1. The two MAT-Proximal haplotypes were not significantly associated with

mating types in the Swiss 2019 recombining populations (Table 1; chi-squared test: χ^2 =

3.8427; p-value = 0.1464). The EP155 genome used as reference for SNP calling above

carried the MAT-Prox1 haplotype but the MAT1-2 allele.

Inversion in the non-recombining MAT-proximal region

To investigate whether rearrangements between haplotypes were present in the MAT-

proximal region, we sequenced the genome, using PacBio HiFi, of two strains with alternative

MAT-proximal haplotypes: M1400 (MAT-2; MAT-Prox2) and M6697 (MAT-1; MAT-

Prox1), originating from the Swiss population in Gnosca and belonging to the CL1 cluster, i.e.

European invasive populations introduced from North America (Stauber et al. 2021; Stauber

et al. 2022). We built high-quality genome assemblies: statistics of the assemblies were in the

same range as those of the genome assemblies of the ESM15 and EP155 strains previously

sequenced (Table S3; Demené et al 2022; Crouch et al 2020). The mating-type chromosome

could be assembled as a single contig for the first time, in the M1400 genome, likely

corresponding to a full chromosome, and was assembled into two contigs in the M6697

genome. We therefore used the M1400 genome as a reference for subsequent analyses. To

identify the location of the MAT-proximal region in the M1400 genome, we computed LD by

mapping the reads of the 1990 Swiss population to the new M1400 genome. We found the

large block (> 1 Mb) of high LD (r^2>0.9) between 7,285,137 and 8,828,934 bp (red arrow;

Fig. S4), showing that calling SNPs on either a MAT-Prox1 or MAT-Prox2 haplotype yielded

similar LD patterns. The MAT-proximal region was also located 20 kb away from the mating-

type locus (located at 7.265 Mb) and was 1.540 Mb long using the M1400 genome as

reference. Consistent with the results using the EP155 genome as reference, we found that the

high-LD region was divided into two higher-LD blocks near the edge at 7.392 Mb (orange

arrows in Fig. S4).

The two new high-quality genome assemblies from the invasive Swiss population (introduced

from North America) revealed an inversion between the two haplotypes in the MAT-proximal

region (Fig 3). The two newly sequenced PacBio genomes were indeed collinear except for

the mating-type chromosome (Fig S5A-B-C), where we found a large region (>1 Mb) that

seem inverted between the M1400 and M6697 genomes (Fig 3; Fig S5A-B-C). Breakpoints of

the inversion were located at ca. 7.455 and 8.563 Mb of the tig00000001 contig in the M1400

genome and at ca. 3.380 and 4.185 Mb of the tig00000060 contig in the M16697 genome. The

region affected by the inversion was ca. 300 kb smaller in the M6697 genome than in the

M1400 genome, while the whole MAT-proximal region was ca. 100 kb smaller. The split into

the two high-LD blocks in natural populations reported above was outside of the inversion,

i.e. 70 kb from the inversion breakpoint (orange arrows in Fig S4). LD in natural populations

was higher within the inversion (median value of $r^2 = 1$) than in other regions of the MAT-

proximal region (median value of $r^2 = 0.64$; Wilcoxon rank sum test with continuity

correction; W = 4.8714e+10; p-value < 2e-16).

We looked for centromeres, as the non-recombining regions near the mating-type locus in

other ascomycetes, when they occur, either capture the centromere (Menkis et al. 2008; Sun et

al. 2017), or are associated to the occurrence of a single crossing-over between the centromere

and the mating-type locus (Grognet et al. 2014; Hartmann et al. 2021; Vittorelli et al. 2022).

The centromere in *C. parasitica* may be at 4.380-4.536 Mb on the mating-type chromosome

in the M1400 genome, as we detected here a peak in TE density and a drop in GC content

(Fig 4A-B). A dotplot of repeats in this region also presented a pattern typical of centromeres

(Fig S5D). The MAT-proximal region and the inversion thus did not include the putative

centromere and was instead located about 2.9 Mb away of the putative centromere.

Higher genetic differentiation and lower genetic diversity within the MAT-proximal region

than in recombining regions

When analyzing polymorphism within the invasive 1990 Swiss population using the M1400

genome as reference, we found much higher genetic differentiation between strains of the two

MAT-proximal haplotypes in this region than elsewhere along the genome (Fig 4C; Fig S6A;

Table S4), as expected for a non-recombining region in LD with the mating-type locus. For

example, the F_{ST} values between the two non-recombining haplotypes within the MAT-

proximal region in the 1990 Swiss population were nearly maximal (median F_{ST}=0.93 per 50

kb window), while they were near zero in the rest of the genome (median F_{ST}=0.02; Wilcoxon

test p-value < 2e-16 with Bonferroni correction; Fig 4C; Fig S6A). Such F_{ST} values close to 1

indicate a lack of shared polymorphism and therefore support the inference of recombination

suppression in the MAT-proximal region.

Within each haplotype, the genetic diversity at the MAT-proximal region was lower than in

the rest of the genome (Wilcoxon test p-value < 2e-16 for the MAT-Prox1 haplotype; p-value

= 2.3e-07 for the MAT-Prox2 haplotype; Fig 4E-F; Fig S6C; Table S4), as expected for a

region without recombination associated with the mating-type locus, as its effective

population size is half as in recombining regions. The diversity was especially low in the pool
of isolates with the MAT-Prox1 haplotype (see all MAT-Prox1 sequences clustering on a
single point on the PCA in Fig 2A and in the neighbor-net network in Fig 2D, in contrast to
the more scattered MAT-Prox2 sequences). Such a very low diversity may be due to a recent
selective sweep or to particularly strong bottleneck in this haplotype during the invasion.
326
We detected signals for long-term balancing selection in the MAT-proximal haplotypes.
Computation of the Tajima's D statistics within all strains suggested signatures of balancing
selection in the MAT-proximal region as expected in a region without recombination
associated with the mating-type locus maintained at a frequency close to 0.5. Tajima's D
values in the MAT-proximal region pooling all sequences (median D=3.0 per 50 kb window)
were higher than in the rest of the genome (median D=-0.14; Wilcoxon test p-value < 2e-16
with Bonferroni correction; Fig 4G; Fig S6D). In contrast, Tajima's D was significantly lower
than the rest of the genome within each pool of haplotype, including in the MAT-Prox2
haplotype, suggestive of positive selection (Wilcoxon test p-value = 0.0037 for the MAT-
Prox1 haplotype; p-value = 0.0022 for the MAT-Prox2 haplotype with Bonferroni correction;
Fig 4H-I; Fig S6D; Table S4). We found no difference in population diversity statistics
between the recombining part of the mating-type contig and the other contigs (Table S4A).
339
The study of synonymous divergence (d_S) between the shared orthologs in the M1400 and
M6697 haplotypes suggests that their differentiation is at least 1.5 million years old and that
there is no pattern of evolutionary strata within the MAT-proximal region, i.e. segments with
different levels of differentiation between haplotypes that would indicate stepwise expansion
of recombination suppression away from the mating-type locus. Per-gene synonymous
divergence (d_S) is typically used for detecting gradual expansion of recombination cessation
and for estimating its age, as it is considered a good proxy for the time since recombination
suppression. Indeed, when recombination is suppressed, mutations accumulate independently
in the two non-recombining haplotypes. We plotted the per-gene synonymous divergence (d_S)
between M1400 and M6697 along the M1400 genome, as its mating-type contig is likely
being assembled as a full chromosome (Fig S7A&E; Table S4). Consistent with
recombination suppression, we found significantly higher d_S values in the MAT-proximal
region (computed for 71 genes) than in the other regions of the mating-type contig (pairwise
Wilcoxon test p-value <2e-16 with Bonferroni correction) and other contigs (pairwise
Wilcoxon test p-value <2e-16 with Bonferroni correction). We found no significant

differences between the other regions of the mating-type contig and other contigs (pairwise

Wilcoxon test p-value = 0.23 with Bonferroni correction). The d_S pattern displayed no

indication of gradual expansion of recombination cessation within the MAT-proximal region,

as there was no stair-like pattern. Using synonymous substitution rate estimates across

multiple ascomycetes (from 0.9 x 10^{-9} to 16.7 x 10^{-9} substitutions site per year (Kasuga et al.

2002; Taylor and Berbee 2006) and the mean d_S value (0.0495) across the genes shared

between the MAT-proximal haplotypes, we estimated the age of their divergence to be at least

1.5 Million years (Table S4).

Enrichment in transposable elements in the non-recombining region and inversion

breakpoints

When studying the M1400 and M6697 high-quality assemblies, we detected an enrichment in

transposable elements (TEs) in the MAT-proximal region in the two haplotypes compared to

the rest of the genome (Fig 4B). The percentage of bp occupied by TEs (TE load) was higher

than 50% in the MAT-proximal region, while it was only 9% on average in other regions. TE

load in the MAT proximal region was higher in the M6697 haplotype (76%, MAT-1; MAT-

Prox1) than the M1400 haplotype (68% MAT-2; MAT-Prox2). Class I retrotransposons with

Gypsy (LTR-Ty3) and LARD elements were the most abundant TEs in all genomic regions

(autosomes, MAT-Proximal region and the rest of the mating-type contig), representing >70%

of TE annotations (Fig 5A-B), as shown previously (Demené et al. 2022). The TE load

however varied significantly among TE families and genomic regions (ANOVA; Table S5).

In M6697 for example, TIR elements appeared less abundant in the non-recombining than

recombining regions while LTR-Ty3 elements were more frequent in the MAT-proximal

region (Fig 5B). In M1400, LARD elements were more frequent in the MAT-proximal region

than in recombining regions (Fig 5B).

The presence of transposable elements at the two inversion breakpoints in the two genomes

(Fig 5A) suggests that the inversion may have occurred via non-homologous recombination

mediated by these elements. By dating retrotransposon insertions using between-copy

divergence and within-copy LTR divergence in both M1400 and M6697 genomes (Fig 5C and

D), we found that TEs within the inversion and around the inversion breakpoints were older

than TEs farther from the breakpoints. This is expected in non-recombining regions as

selection is less efficient to purge TE insertions in such regions so that they remain there

longer (Duhamel et al. 2023). The date estimates of LTR element insertions (Fig 5D) further

indicated that the TE accumulation in the MAT-proximal region was as old as a few million

years, confirming the age estimated based on sequence divergence between MAT-proximal

haplotypes.

Gene function in the MAT-proximal region and association to known phenotypes

We found no gene function known to be involved in mating or mating compatibility, nor in

pathogenicity, in the MAT-proximal region (Table S7-Table S8). We nevertheless identified

genes potentially coding for proteins with kinase domain or proteins with homeodomains

(Table S6), that are often involved in the regulation of developmental pathways.

By re-analysing data from a previous study in the light the two haplotypes (Stauber et al.

2021; Stauber et al. 2022), we found no association of the MAT-Proximal haplotypes to

previously described vegetative compatibility groups or to the sensitivity to virus infection,

but more specific studies need to be performed to specifically test hypotheses.

The MAT-proximal region harbours* Starship *elements

In addition to a high density of transposable elements, gene annotations of the MAT-proximal

region in the M1400 and M6697 genome assemblies revealed multiple genes encoding

tyrosine recombinases (genes with DUF3435 domains, that can be *Starship* captains) and gene

content variation (Tables S6-S7), suggesting the presence of *Starships*. By running the Gene

Finder Module of the starfish pipeline designed to identify *Starships* (Gluck-Thaler and

Vogan 2024), we in fact identified the presence of putative *Starship* captains in the M1400

haplotype (MAT-Prox2) and three putative *Starship* captains in the M6697 haplotype (MAT-

Prox1). We detected eight and ten other putative *Starship* captains elsewhere in the M1400

and M6697 genomes, respectively, but the MAT-proximal region appeared enriched in these

elements (Fisher's exact test p-value =0.0004). A phylogenetic analysis with previously

described *Starship* captains (Gluck-Thaler and Vogan 2024) suggested that the captains in the

MAT-Proximal region belonged to the Phoenix *Starship* family, that preferentially target AT-

rich sites (Fig S7 ; (Gluck-Thaler and Vogan 2024)). As the Starfish pipeline performs poorly

in TE-rich regions (Gluck-Thaler and Vogan 2024), we investigated manually the delimitation

of putative starships. Three putative captains shared orthology relationships between the two

haplotypes, one being just at the border of the MAT-proximal region near the mating-type

locus (Figure 3). The high similarity between captains shared by the two haplotypes (>90%

identity), their similar positions and reverse strand orientations suggested that these captains

were present before the differentiation between haplotypes and before the inversion (Figure

5). The genes near these three shared captains are mostly common to the two haplotypes. The

fact that these elements were present in all high-quality assemblies at the same locus

prevented delimiting Starship boundaries and these genes may represent solo Starship

captains or pseudogenes as found in high frequency in Pezizomycotina genomes (Gluck-

Thaler and Vogan 2024). The MAT-Prox2 haplotype carried an additional putative captain,

followed by 18 genes lacking in the MAT-Prox1 haplotype (Figure 5A), suggesting the

presence of a *Starship* element, together with its cargo genes, only in the MAT-Prox2

haplotype. Functional annotation of the putative cargo genes included two genes with

DUF3723 domain and one gene with ferredoxin reductase-type domain, that are often found

in *Starships* (Table S8 (Gluck-Thaler and Vogan 2024). This putative *Starship* was about 44

kb in the M1400 strain.

In addition to the genes only present in the putative Starship specific to the MAT-Prox2

haplotype, the two haplotypes carried additional specific genes, especially the MAT-Prox2

haplotype. Based on automatic gene annotation and reconstruction of orthologous

relationships, we found that, out of 175 groups of single-copy genes in the MAT-proximal

region, 80 (46%) were only present in M1400 (MAT-Prox2) and 24 (14%) were only present

in M6697 (MAT-Prox1). This level or presence/absence gene polymorphism was higher than

in other genomic regions, in which only 3% of genes were only present in one of the two

genomes genome (Fisher's exact test p-value < 2.2e-16). However, not all genes specific to

one haplotype were located close to a putative captain (Figure 3), so that part of the observed

gene presence/absence polymorphism between the two MAT-Proximal haplotypes may also

be due to gene losses due to degeneration or pseudogenisation after *Starship* insertions, to

recombination suppression or to gene gains by other means than *Starships*.

Among the genes specific to M6697 (MAT-Prox1), we found a gene encoding a protein of the

Sirtuin family (IPR003000). A Sirtuin homolog was found in a *Starship* associated with local

thermal climate adaptation in the wheat fungal pathogen *Zymoseptoria tritici* (Tralamazza et

al. 2024).

Gene expression in the MAT-proximal region

We analysed the expression data available for the EP155 strain (Chun et al. 2020), carrying

the MAT-Prox1 haplotype, to test whether some of the genes in the MAT-Proximal region

were upregulated under certain conditions. Out of 127 genes functionally annotated in the

MAT-proximal region of the EP155 genome, 50 genes were supported by RNAseq data

acquired *in vitro* for this strain (Chun et al. 2020), suggesting that the region contains

functionally active genes (Table S6). Among these 50 genes, nine were upregulated during

barrage allorecognition of the EP155 strain with a compatible strain (Belov et al. 2021) and

five were upregulated during infection of the EP155 strain with the hypovirus CHV1 (Chun et

al. 2020) (Table S6), suggesting that the MAT-Proximal region may have a role in fungus-

mycovirus interactions and vegetative incompatibility. Unfortunately, no expression data are

available for a strain with the MAT-Prox2 haplotype, that harbours the additional *Starship*

and the highest number of specific genes and no expression data is available during plant

infection.

Degeneration in the non-recombining region

The MAT-Proximal region was enriched in TEs and depleted in genes (Fig 4 and Fig 5A),

which supports the inference of an old full recombination suppression. We also found overall

higher non-synonymous substitution rate compared to the baseline substitution rates (d_N/d_S

values) between the M1400 and M6697 genomes in the MAT-proximal region (median per

gene $d_N/d_S = 0.80$) than in other contigs ($d_N/d_S = 0.43$; pairwise Wilcoxon test p-value = 1.7e-

07 with Bonferroni correction; Fig S8A; Table S4). This is consistent with relaxed purifying

selection due to recombination suppression in the MAT-proximal region. In addition, 13

genes displayed $d_N/d_S > 1$ between M1400 and M6697, suggesting that they may evolve under

positive selection (Table S9).

Recombination suppression and Starships *in the MAT-Proximal region in other*
populations assessed from high-quality genome assemblies

We analyzed five additional high-quality genome assemblies for strains from the Asian native

range (CL2 and CL4 clusters) and the invasion range (including two North American strains

belonging to the CL1 cluster and one European strain introduced directly from Asia and

belonging to the CL2 cluster), three of which were generated for the present study (Table S3).

Most of the high-quality genomes displayed the same MAT-Proximal chromosomal

arrangement as M1400 (MAT-Prox2, Swiss population and belonging to the CL1 cluster),

suggesting that it is the ancestral state (Fig S9A-D), although the incomplete assemblies may

prevent detecting other rearrangements. The inversion detected in M6697 (MAT-Prox1, Swiss

population and belonging to CL1 cluster) was also present in the EP155 strain (MAT-Prox1,

North America and belonging to CL1 cluster, Crouch et al, 2020; Fig S8D), indicating that

this inversion was already segregating in early invasive populations in North America. In
other genomic regions outside of the MAT-Proximal region, all analyzed high-quality
genomes were colinear to the ESM15 reference genome, except the highly rearranged EP155
genome, as previously reported (Demené et al. 2022).

The MAT-Proximal region in all the high-quality genomes, from Asian native strains and
invasive strains, displayed signs of degeneration, with a high TE load (Fig6; Fig S9),
confirming the ancient age of recombination suppression and its occurrence in the native
range. Study of d_S and d_N/d_S between the MRC10 strain, belonging to the European clonal
lineages of direct Asian origin (CL2 cluster), the M6697, M1400 and the two Asian native
strains further confirmed recombination suppression in the native range and in invasive strains
from different origins (North America or directly from Asia). The d_S and d_N/d_S values were
indeed significantly higher in the MAT-proximal region than other genomic regions (Fig S7
B-E). Mean d_S values in the MAT-proximal region were in the same range as d_S values
between the M6697 and M1400 genomes, suggesting a similar date of recombination
suppression and, therefore, that recombination suppression was ancestral. The MAT-proximal
region appeared smaller in the two Asian strains XIM9508 and ESM15 (<1 Mb) than in
invasive strains (Table S3), but this may be due to the incomplete genome assemblies, and/or
to additional *Starship* presence/absence polymorphism.

We detected the three captains shared between the M6697 and M1400 haplotypes in the
MAT-proximal region of all high-quality genome assemblies, indicating that their insertions
in the MAT-proximal region were old, being already present in the native range. We found
the putative *Starship* specific to M1400 (MAT-Prox2), i.e., the captain and the 18 cargo
genes, in the MAT-proximal region of DUM-005 (MAT-Prox2) and in the haplotypes of
MRC10 and the Japanese strain ESM15 (both genetically close to MAT-Prox2; Figure
S10D1), while it was absent in the MAT-Proximal region of the EP155 strain (MAT-Prox1).
In the XIM9508 Chinese strain (genetically close to MAT-Prox1; Figure S10D1), the captain
and four genes of the *Starship* specific to M1400 were absent in the MAT-proximal region.
However, 14 of its putative cargo genes were present and a very similar captain was found
elsewhere in the genome, suggesting that this *Starship* may have moved away from the MAT-
Proximal region after its insertion there. In the seven high-quality assemblies, the putative
Starship presence/absence polymorphism detected between the 1400 and M6697 genomes
thus seem associated to the MAT-Proximal haplotypes in the invaded range and also in the

native Asian range (Figure S10D), although this would need to be confirmed with additional

high-quality assemblies.

Presence of the two MAT-Proximal haplotypes in other populations, but less differentiated

in the native range

By analyzing available additional Illumina genomes of invasive strains from the first invasion

wave via North America (strains from the US and European clonal lineages), we recovered

the two MAT-proximal haplotypes in balanced frequencies. We indeed observed high

differentiation between two clusters in the MAT-Proximal region, contrasting with a lack of

genetic subdivision in other genomic regions or a completely different genetic subdivision,

corresponding to previously described population genetic structures (Figs S10A and S10C).

This reinforces the view of balancing selection on the MAT-Proximal haplotypes.

In the native range, the MAT-Proximal region also seemed to have undergone a very different

evolutionary history than the rest of the genome (Figure 7). In particular, the neighbor-net

network with multiple strains sequenced previously from the CL2 and CL3 clusters (from the

invaded range and the native range, in South Korean and Japan) and from the CL4 cluster

(from the native range, in China) indicated that sequences from these three clusters were

intermingled in the MAT-proximal region (Fig 7A), while the structure in the rest of the

genome instead corresponded to the different population clusters, i.e., CL1, CL2, CL3 and

CL4 (Fig 7B). However, the two haplotypes were much less differentiated in the native range

and in invasive populations originating directly from Asia than in invasive populations from

the first invasion wave via North America. For example, branch lengths on the neighbor-net

indicate a lower differentiation between haplotypes within CL2 (Fig 7A) than within the

European CL1 or Swiss invasive populations (Fig 2D; Fig S3) and some reticulations were

present between MAT-proximal haplotypes.

We found no association of the MAT-Proximal haplotypes with mating types at the world

scale (Figs 7 and S10). In the S12 European invasive clonal lineage (only MAT-1), we found

a single MAT-Proximal haplotype (MAT-Prox1; Fig S10B), but we detected the two MAT-

Proximal haplotypes in all other populations. Even in two invasive European lineages of

American origin with a predominantly clonal structure (re092 and re103), the strains with a

mating type different from their clone-mates, due to localized introgression from other clonal

lineages (Demené et al. 2019), also had their MAT-Proximal region introgressed (Fig S10C);

this finding reinforces the view of the strong linkage between the mating-type locus and the

MAT-Proximal region, of an advantage of maintaining the two MAT-Proximal haplotypes in

populations, and perhaps of maintaining an association between MAT-proximal haplotypes

and mating types within populations. Three 2019 Swiss strains exhibited a similar

intermediate sequence in the MAT-Proximal region as the other strains from Europe with a

direct Asian origin described above (Fig S10C). This intermediate haplotype thus seemed

associated with the second invasion in Europe, directly from Asia, and carried the same

captains as the MAT-Prox2 haplotype from the M1400 strain. The location as an intermediate

in the neighbor-net network may be due to recombination between the haplotypes, or may just

reflect the fact that the two haplotypes are less differentiated in the native Asian area.

***No evidence of introgression in the MAT-proximal region from* closely related**
Cryphonectria species

The stronger differentiation in the MAT-Proximal region in the Swiss and CL1 invasive

populations than in other populations could also be due to an introgression event from a

closely related species; however, the small genetic distance and the reticulations observed in

the neighbor-net network between the M6697 Swiss reference genome and the Asian

genomes M8444 and XA19 rather indicate that the MAT-Prox1 haplotype in Switzerland

originated from Asian *C. parasitica* populations, likely a population different from those at

the origin of the CL2 invasive cluster (Fig 7).

Gene genealogies also provided no evidence of introgression in the MAT-proximal region

from closely related *Cryphonectria* species. We performed a gene orthology analysis

including three genomes of the closely related species *C. japonica*, C. *carpinicola*, and *C.*

naterciae that were previously sequenced with short read technologies (Stauber et al. 2020).

We retrieved only nine genes from the MAT proximal region that were shared between the

M6697 and M1400 genomes that had orthologs in one of the related species. This may be due

to low quality assemblies in the outgroups (Stauber et al. 2020) and/or to many gene gains in

this region in *C. parasitica* because of *Starships*. We found no gene with a topology

consistent with introgression from another species into the MAT-Proximal region, i.e. with a

MAT-Prox1 or MAT-Prox2 haplotype that would branch with an outgroup allele rather than

with the alternative haplotype of *C. parasitica*. The high load of TEs and gene disruptions in

the two MAT-Proximal haplotypes does not fit with the introgression hypothesis either,

unless the introgression occurred from a species with recombination suppression in the MAT-

Proximal region. The Starships may have moved from a distant species by horizontal gene
transfer together with their cargo genes, but we found no evidence for this by blasting the
captains and cargo genes in public databases.
596
597

Discussion
A large non-recombining region near the mating-type locus, with two highly differentiated
haplotypes maintained polymorphic in the native and introduced ranges
We found strong evidence for the complete cessation of recombination in a region of more
than 1 Mb (1.2 Mb to 1.5 Mb depending on the haplotype), proximal to the mating-type locus
in *C. parasitica*. Indeed, we detected maximal levels of linkage disequilibrium in otherwise
recombining populations and the existence of two differentiated haplotypes, without shared
polymorphism, while even low rates of recombination can homogenize alleles and prevent LD
build-up (Dufresnes et al. 2015). The full cessation of recombination was further supported by
the existence of a previously unknown, large inversion in the center of the non-recombining
region in invasive strains. This is in agreement with previous genetic maps reporting lack of
recombination near the mating-type locus in crosses involving Japanese, North American and
Italian isolates (Kubisiak and Milgroom 2006a). The higher TE load in this region constitutes
additional evidence for full recombination cessation, as the lack of recombination renders
selection against TE accumulation less efficient, in particular due to Muller's ratchet: two
chromosomes with different TE insertions cannot recombine to produce a chromosome free of
TE insertions. TE takes hundreds of thousands of years to accumulate in non-recombining
regions (Duhamel et al. 2023), so that the high TE load in the MAT-Proximal region in *C.*
parasitica indicates old recombination cessation. The particularly high frequency of non-
synonymous substitutions in the MAT-Proximal region worldwide further indicates
degeneration and ancient recombination suppression. The existence of two differentiated
haplotypes appeared less clear in the native range in Asia and in the second-wave invasive
populations originating directly from Asia than in invasive populations from the first
introduction via North America, but the MAT-Proximal region nevertheless appeared to
display a different evolutionary history than the rest of the genome and a high TE load.
623

The estimates for the differentiation between the two MAT-Proximal haplotypes based on
synonymous divergence between shared single-copy orthologs and based on TE insertion
dates were at least 1.5 million years. The suppression of recombination in *C. parasitica* may

be younger than this estimate if one of the two haplotypes was introgressed from a distant

species. The evolution of non-recombining regions by introgression has been reported for

example in ants and butterflies (Jay et al. 2018; Helleu et al. 2022; Stolle et al. 2022).

However, we found little evidence for an introgression: i) the two haplotypes were detected in

native populations and in the CL2 cluster of invasive European populations directly

originating from Asia, even if less differentiated than in the invasive populations from the first

introduction, ii) the MAT-Prox1 haplotype in the CL1 cluster that is the most differentiated

from the MAT-Prox2 haplotype is genetically close to an Asian haplotype, iii) footprints of

recombination suppression are present in the native range in terms of degeneration and

particular genomic structure in the MAT-Proximal region, and iv) we found no signatures of

introgression from closely related species in gene genealogies or by blast in public databases.

In any case, the findings of recombination suppression footprints and of the presence of two

differentiated haplotypes in the two invaded continents, Europe and North America, as well as

in the native range of *C. parasitica*, with intermingled sequences from different Asian

populations in neighbor-net networks, indicate ancient recombination suppression and long-

term polymorphism maintenance in the MAT-Proximal region. These findings point to a

strong balancing selection maintaining two differentiated haplotypes in *C. parasitica.*

In contrast to other known non-recombining regions on sex or mating-type chromosomes

(Bergero and Charlesworth 2009; Hartmann et al. 2021), the region without recombination

was not completely linked to the mating-type locus. Indeed, the occurrence of rare

recombination events between the non-recombining MAT-Proximal region and the mating-

type locus are supported by previous segregation analyses (Kubisiak and Milgroom 2006b)

and the lack of association found here between the MAT-Proximal haplotypes and the mating

types at the worldwide scale, as well as their incomplete association within populations.

Proximal cause of recombination suppression

The large inversion in the MAT-Proximal region may contribute to recombination

suppression. There may also be additional nested inversions and/or translocations in the

MAT-Proximal region as there are breaks of synteny between the inverted fragments with

many transposable element insertions. The chromosomal arrangement of the M1400 MAT-

Prox2 haplotype is likely the ancestral state as it is the most frequent worldwide, although a

more complete sampling in the native range is required to obtain definitive conclusion. The

non-recombining region was defined based on LD pattern and was larger than the inversion.

This may indicate that additional proximal mechanisms suppress recombination. The
inversion may even be a consequence rather than a cause of recombination cessation, as
previously found in other fungal mating-type chromosomes (Grognet et al. 2014; Sun et al.
2017; Branco et al. 2018; Vittorelli et al. 2022), especially as the inversion was not detected
so far in the native range despite the presence of footprints of recombination suppression.
Other proximal causes of recombination cessation may for example be genetic recombination
modifiers, epigenetic marks, histone modifications or lack of recruitment of proteins
responsible for double-strand breaks (Maloisel and Rossignol 1998; Boideau et al. 2022;
Legrand et al. 2024). The lack of rearrangements at the edge of the MAT-proximal region
may allow rare recombination or gene conversion events, which may explain the presence of
two distinct LD blocks with lower LD between them and the lower differentiation at the edges
of the MAT-proximal region. Alternatively, the LD around the inversion may be formed by
less frequent recombination as recombination is often modified around inversion breakpoints
(Pegueroles et al. 2010; Stevison et al. 2011) and the actual size of the non-recombining
region may be smaller than suggested from the LD pattern.

The MAT-proximal haplotypes carry putative* Starship *elements
We detected signatures of *Starship* elements in the non-recombining MAT-proximal region,
with likely three "solo" captains shared between the two haplotypes, but also an additional
captain associated to 18 specific genes, strongly suggesting the presence of a cargo-
mobilizing *Starship*, only present in the MAT-Prox2 haplotype. The difference in gene
content between the two MAT-proximal haplotypes thus likely at least partly result from
specific cargo genes inserted with *Starships,* although there could also be gene losses because
of the less efficient selection due to recombination suppression, and/or additional gene gains.
One of the captains shared between haplotypes was located just nearby the mating-type locus
and at the edge of the MAT-proximal region. This altogether strongly suggest a role of
Starship elements in the formation of the MAT-proximal haplotypes and in their long-term
maintenance, although experiments are required to confirm this hypothesis and elucidate their
role. High-quality assemblies suggested that the presence/absence of the polymorphic
Starship insertion was associated to the MAT-Proximal haplotypes, although this also needs
to be checked in a larger set of strains. The presence of the inversion in the native range also
remains to be investigated. Additional sampling in the native range and high-quality genome
assemblies are required to obtain a more comprehensive view of the history of the MAT-
Proximal region and of the putative *Starship* insertions.

Starships may have been introgressed from another species, as frequent with these elements
(Ropars et al. 2015; Peck et al. 2023), but searches in databases returned no significant blast
results. Such horizontal gene transfers should not have, however, inflated the date estimates of
divergence of recombination suppression in the MAT-Proximal region, as these were
computed based on shared single-copy orthologs between haplotypes and the shared putative
Starship elements seem to correspond to homologous insertions.

The selective forces potentially maintaining the two haplotypes in the MAT-proximal
region
Our findings indicate that selection maintains the two haplotypes polymorphic in the MAT-
Proximal region. There can be several hypotheses regarding the nature of this balancing
selection. The degeneration of the non-recombining region may help maintaining the two
haplotypes by selection due to the sheltering of combinations of deleterious mutations in
repulsion, i.e. thanks to a heterozygote advantage called pseudo-overdominance (Abu-Awad
and Waller 2023). Such pseudo-overdominance advantage may only be a consequence of
recombination suppression, and contribute to its maintenance, or may even be its initial cause
(Branco et al. 2017; Jay et al. 2021; Charlesworth 2023; Jay, D.L. Jeffries, Hartmann, et al.
2024). These hypotheses require the existence of a substantial diploid or dikaryotic phase for
sheltering recessive deleterious alleles in a heterozygous state. In fact, although *C. parasitica*
has a predominant haploid phase, strains can be found as dikaryons heterozygous for the
mating-type locus in some natural populations, and not only as monokaryons (McGuire et al.
2004; McGuire et al. 2005; Stauber et al. 2021; Stauber et al. 2022). The importance and
frequency of heterokaryons in *C. parasitica* in nature would deserve further investigations. It
may be sufficient if recessive deleterious alleles are sheltered in a substantial percentage of
individuals for selecting recombination suppression, and/or if the genes involved are
preferentially expressed during the diploid or dikaryotic phases. This hypothesis of balancing
selection, and in particular the hypothesis of overdominance (i.e. heterozygote advantage),
does not require the same association between the MAT-Proximal haplotypes and mating
types across all populations worldwide, it is sufficient that the MAT-Proximal haplotypes are
strongly associated to mating types within populations. Further sampling will be required to
test whether the MAT-Proximal haplotypes are associated to mating types within local
populations in the CL3 and CL4 clusters and in the native Asian range. It will also be
interesting in future studies to compare the fitness between strains carrying the two MAT-
Proximal haplotypes and dikaryotic strains being homozygous for the MAT-Proximal

haplotype. Such experiments in Sordariales fungi reported a heterozygote advantage (Guyot et
al. 2024). Results from previous studies suggest that strains carrying either the same or
different MAT-Proximal haplotypes can be crossed in *C. parasitica* (Stauber et al. 2021).

It could indeed also be that genes evolve under overdominance in this region for other reasons
than deleterious mutations, so that a partial linkage to a permanently heterozygous locus is
beneficial, perhaps in relationship with the pathogenic lifestyle of the fungus. One could
imagine, for example, that heterozygosity could be advantageous at genes involved in
virulence against the tree host, and especially the novel hosts colonized in invasive ranges, or
in resistance against a parasitic virus known to negatively affect fitness in *C. parasitica* (Choi
and Nuss 1992; Brusini et al. 2017). As a matter of fact, multiple genes present in the MAT-
Proximal region are up-regulated under infection by the hypovirus CHV1 or during the
vegetative incompatibility reaction considered to play a role in the prevention of virus
transmission (Choi et al. 2012; Rigling and Prospero 2018). The specific genes unique to one
or the other MAT-proximal haplotypes, and possibly the cargo genes inserted with the
Starships, may contribute to such a heterozygous advantage.

As an alternative to a heterozygote advantage, the selection maintaining the recombination
suppression and the two haplotypes in the MAT-proximal region could be some kind of
negative-frequency dependent selection of beneficial allelic combinations, possibly linked to a
trench-warfare-like arms race with the host tree, the virus or the microbial community, or to
self-incompatibility, where partial linkage to the mating-type locus would help maintenance
in balanced proportions and therefore would prevent allele loss (Tellier et al. 2014; Jay,
Aubier, et al. 2024). The MAT-proximal region did not include any of the genes previously
identified as controlling vegetative incompatibility in *C. parasitica* but not all self-
incompatibility genes have been identified yet. The MAT-proximal region actually carried
genes upregulated under virus infection or vegetative incompatibility reaction. Such a role in
host-pathogen interactions or vegetative incompatibility would also be consistent with the
selective sweep footprints detected in the MAT-proximal region in a previous study (Stauber
et al. 2021), if there is recurrent positive selection for improving the efficiency of the
pathogen weapons within each of one of the two MAT-proximal haplotypes or new, rare self-
recognition alleles. Such negative-frequency dependent selection, could also explain the
particular population structure in the MAT-proximal region in the native region, with a mix of
sequences from the different genetic clusters, if alleles introgressed between populations are

favored by a positive selection of rare alleles, or with the long-term maintenance of self-
incompatibility alleles, and therefore of ancestral polymorphism. Much higher differentiation
between MAT-Proximal haplotypes in the invasive than the native populations may be due to
a selective sweep of a rare and differentiated MAT-Prox1 allele present in the native range or
having evolved rapidly, by particular demographic effect during the invasion (Moinet et al.
2022), both of which are consistent with the very low genetic diversity in the MAT-Prox1
haplotype in the introduce range. Balancing selection and location adjacent to a non-
recombining region has been reported for loci involved in host resistance in the *Daphnia–*
Pasteuria system (Fredericksen et al. 2023). The MAT-Proximal region may in this case even
include genes involved different traits under balancing selection, and would then constitute a
supergene (Schwander et al. 2014). The MAT-Proximal region also contained, only in the
MAT-Prox1 haplotype, a gene of the Sirtuin family, a homolog of which had previously been
found in a *Starship* associated with local thermal climate adaptation in the wheat fungal
pathogen *Zymoseptoria tritici* _(Tralamazza et al. 2024). Another hypothesis to explain the
balancing selection in the MAT-Proximal region may thus be a heterogeneous selection, with
different genes in the two haplotypes providing contrasted advantages in different conditions
or different phases of the life cycle. The proximity of the mating-type locus, permanently
heterozygous in dikaryotic and diploid stages, may help maintaining balanced frequencies of
the two MAT-proximal haplotypes. Here too, the genes inserted by the *Starships* may play a
role in such balancing selection, especially the genes that are present in a single of the two
MAT-Proximal haplotypes. The other transposable elements insertion polymorphism
observed from the high quality assemblies may also be adaptive (Casacuberta and González
2013; Orteu et al. 2024).

Another hypothesis for explaining such extension of recombination suppression beyond the
mating-type locus is antagonistic selection, i.e. linkage of alleles that would improve fitness
of a MAT-1 gamete while being deleterious in a MAT-2 gamete, or vice-versa. However, full
recombination suppression with the mating-type locus would be expected under such
antagonistic selection, as well as the same association between the MAT-Proximal alleles and
the mating-type in all populations, in contrast to our findings. In addition, we found no
particular predicted function in the MAT-proximal region that could be related to mating
compatibility and there are very little functions, if any, with possible antagonistic functions
between mating types in fungi (Bazzicalupo et al. 2019; Hartmann et al. 2021).

Conclusion

In conclusion, we provide strong evidence for the existence of a non-recombining region partially linked to the mating-type locus in the chestnut blight fungus *C. parasitica*, with two highly differentiated haplotypes, each carrying specific genes, maintained polymorphic by selection. We found footprints of balancing selection in the MAT-proximal region in the first introduction of the pathogen in Europe from North America and a chromosomal inversion. The non-recombining region also displayed footprints of particular evolution in Asia and in the second invasion wave directly from Asia, although the levels of differentiation between haplotypes was lower than in the populations from the first introduction wave. This non-recombining region may underlie important adaptive traits and thereby provide important applications for the control of a devastating tree pathogen. This is supported by the finding of putative *Starships* elements in the MAT-proximal region, i.e., giant mobile elements recently discovered in ascomycete fungi, containing multiple cargo genes (Gluck-Thaler et al. 2022; Urquhart et al. 2024), that can be involved in adaptation. In addition, the high-quality genome assemblies provided here, from the native and invaded ranges, will more generally be useful for studies aiming at understanding the evolution of this invasive and damaging pathogen.

Material and Methods

Strains and genomic data

For population genomic analyses, we studied a collection of 386 monokaryotic *C. parasitica* strains sampled worldwide, from the native and invaded ranges of the pathogen, and sequenced previously with the short-read Illumina technology (strain information are presented in Table S1; (Demené et al. 2019; Stauber et al. 2021; Stauber et al. 2022)). Data were downloaded from NCBI Bioproject numbers PRJNA604575, PRJNA644891 and PRJNA706885. We focused our analyses first on European invasive strains originating from North America. We studied 88 strains belonging to the CL1 genetic cluster, in central and southeastern Europe (Stauber et al. 2021). We excluded the putative clonal genotypes previously identified (Stauber et al. 2021). We also studied 71 strains sampled in the 1990 and 62 strains sampled in 2019 in southern Switzerland (Stauber et al. 2022). Mating-type ratios close to ½ and population structure analyses of the CL1 genetic cluster and the Swiss 1990 population suggest frequent recombination in these populations (Stauber et al. 2021; Stauber et al. 2022). The mating-type ratio was 33% *MAT-1* in the 2019 Swiss population and population structure analyses suggested regular sexual reproduction and recent population

bottleneck. To study the presence of the two haplotypes in the MAT-proximal region more

broadly, we analyzed monokaryotic strains belonging to the CL2, CL3, CL4 genetic clusters,

as well as the S12 European invasive lineage (Stauber et al. 2021) and additional

monokaryotic strains from the US and Europe (with an Asian or North American origin

(Demené et al. 2019)). We excluded heterokaryotic strains, i.e. strains having both MAT-1

and MAT-2 alleles, as the phase was challenging to infer.

For mapping and SNP calling, we first used as reference the 43.9 Mb genome sequence

EP155 v2.0 of the strain EP155 (*MAT-2,* North America, CL1 cluster) available at the Joint

Genome Institute (http://jgi.doe.gov/)(Crouch et al. 2020). For comparative genomics, we

used the published genome of the strain ESM15 (MAT-2, Japan, CL2 cluster; (Demené et al.

2022)) available at DDBJ/ENA/GenBank on the bioproject PRJNA700518 under the

accession JAGDFO000000000. We additionally sequenced *de novo,* with long-read

technologies, five strains from the native and invaded ranges of the pathogen. We sequenced

with PacBio Hifi the genomes of the strains M1400 (MAT-2) and M6697 (MAT-1) sampled

in Gnosca, southern Switzerland (Stauber et al. 2021). Mycelia were stored as glycerol stocks

at -80 C after strain isolation. To produce mycelium for DNA extraction, isolates were

inoculated onto cellophane-covered potato dextrose agar plates (PDA, 39 g/L; BD Becton,

Dickinson and Company, Franklin Lakes, USA) (Hoegger et al. 2000) and incubated for a

minimum of 1 week at 24°C, at a 14 hr light and 10 hr darkness cycle. Mycelium and spores

were harvested by scratching the mycelial mass off the cellophane, transferring it into 2 mL

tubes and freeze-drying it for 24 hr (Stauber et al. 2021). DNA extraction was performed with

the NucleoBond High Molecular Weight DNA kit from Macherey-Nagel, with the mechanical

disruption of about 30 mg of lyophilized mycelium with two beads of 3 mm for 5 min at 30

Hz. Sequencing was outsourced to Novogene, the Netherlands. We additionally sequenced

with Oxford Nanopore MinION technology the strains XIM9508 (China, CL4 cluster, MAT-

1), MRC10 (South Western France introduced directly from Asia, CL2 cluster, MAT-2) and

DUM-005 (USA, CL1 cluster, MAT-2). These isolates had been collected for previous

studies (Milgroom et al. 1996; Liu et al. 2003; Dutech et al. 2012). The protocols for

mycelium production, DNA extraction and sequencing for these three strains were the same

as in (Demené et al. 2022).

Short-read data processing and SNP calling

We used SNP calling datasets of the genomes from monokaryotic strains of the CL1, CL2,

CL3, CL4 genetic clusters, the S12 invasive lineage and the Swiss populations against the

EP155 reference v2.0 genome obtained in (Stauber et al. 2021; Stauber et al. 2022). We

performed mapping and raw SNP calling of short-read data using the M1400 new genome

assembly as a reference as described in (Stauber et al. 2021; Stauber et al. 2022). Briefly, we

trimmed reads with Trimmomatic v0.36 (Bolger et al. 2014) and aligned them with Bowtie 2

v2.3.5.1 (Langmead et al. 2009) and SAMtools v1.9 (Li et al. 2009) to the EP155 v2.0

genome. Raw SNP calling and filtration for quality were conducted with the genome analysis

toolkit GATK v3.8 and v4.0.2.0 (McKenna et al. 2010). We used the filtration parameters

described in (Stauber et al. 2021): QUAL>=100, MQRankSum (lower) >= -2.0, QD:_20.0,

MQRankSum (upper) <=2.0, MQ:_20.0, BaseQRankSum (lower) >=-2.0, ReadPosRankSum

(lower) >=-2.0, ReadPosRankSum (upper) <=2.0, BaseQRankSum (upper) <=2.0. We further

removed SNPs overlapping with transposable elements predicted *de novo* in the EP155 v2.0

genome by (Stauber et al. 2021).

Population genomics analyses

For all population genomics analyses, we excluded SNPs with missing data in more than 10%

of the strains and kept only polymorphic strains with vcftools v0.1.16 (Danecek et al. 2011).

To study linkage disequilibrium, we further excluded rare variants (minor allele frequency

<0.1) with the vcftools (Danecek et al. 2011) option --maf 0.1. We computed LD with the --

hap-r2 option of vcftools (Danecek et al. 2011) for each scaffold and each population

separately. We used the --thin 50000 option of vcftools (Danecek et al. 2011) to sample SNPs

distant of at least 50 kb. We used the R package LDheatmap v1.0-6 (Shin et al. 2006) to plot

LD r^2 values among SNP pairs. To perform principal component analyses (PCAs), we first

used vcftools (Danecek et al. 2011) to convert VCF format files in Plink format. We then used

the Plink v1.90b5.3 (Purcell et al. 2007) –pca command to run PCA analysis. We used the R

package PopGenome v2.7.5 (Pfeifer et al. 2014) to compute nucleotide diversity, Tajima's D

values and the F_{ST} index in 50 kb window overlapping over 10 kbp. Windows containing

fewer than 5 SNPs were removed from the analysis. We used the R package ggplot2 (2_3.5.0)

to plot results. Pairwise Wilcoxon tests were performed in R with Bonferroni or false

discovery rate correction.

We built neighbornet networks using SplitsTree4 v 4.19.2 (Huson 1998). VCF file were

converted to nexus format using PGDSpider v1.0 tool (Lischer and Excoffier 2012). For the

study of other populations, we included only one strain of each haplotype of the invasive

European population but not all CL1 and Swiss strains for network readability; we

nevertheless checked that the results were the same with all CL1 and Swiss strains.

Long-read based assemblies

PacBio Hifi reads of strains M1400 and M6697 were both assembled using canu v1.8 (Koren

et al. 2017) program with a set genome size of 44 Mb. Multiple assembly pipelines were used

for the other strains. Oxford Nanopore MinION reads and Illumina reads of the genomes of

both XIM9508 and MRC10 strains were assembled using HybridSPAdes (Antipov et al.

2016). Assemblies were manually curated and scaffolds were cut when an evidence of a

chimeric connection was detected (i.e. mis-paired short reads) as previously described

(Demené et al. 2022). For DUM005, the assembly was generated by Ra with basic parameters

that uses Oxford Nanopore MinION reads and corrects the assembly with Illumina reads

(https://github.com/lbcb-sci/ra) as it outperformed the HybridSPAdes assembly. As the

HybridSpades assembly of the MRC10 strain suggested absence of collinearity with the

MAT-proximal region M1400 and M6697, we further checked the assembly of the MAT-

proximal region of MRC010 by generating a meta-assembly of this strain. We used the

assembler tool canu v2.2 with an estimated genome size of 42 Mb. We also used Flye v2.9.3-

b1797 (Kolmogorov et al. 2019) with an estimated genome size of 42 Mb, --nano-raw for

reads with an error rate below 20 and a coverage for initial disjointig assembly of 50. Then we

used ragtag (Alonge et al. 2019) to patch the canu assembly with the Flye assembly as query

in a first loop. In a second loop, the canu assembly was patched with the first loop ragtag

assembly. Finally, we polished this second output assembly of ragtag with short reads and the

consensus part of medaka v1.11.3 with no change in the parameters

(https://github.com/nanoporetech/medaka). Assemblies statistics were obtained with quast

v5.1 (Gurevich et al. 2013). We assessed the completeness of each assembly using the

Benchmarking of Universal Single-Copy Orthologue (BUSCO) tools with the Sordariomyceta

ortholog set (sordariomycetes_odb10, 2020-08-05, n = 3817 orthologous groups searched)

(Manni et al. 2021). Gene models were predicted with the Helixer v0.3.1 pipeline (Holst et al.

2023). We also run Helixer pipeline for ESM15 and EP155 strains for the gene orthology

analysis. Statistics of the obtained gene annotation was obtained with the Agat v1.0.0 tool

(Dainat et al. 2020). Transposable elements of all genomes were annotated using

Repeatmasker v4-0-7 (Smit et al. 2013) and the customized library built for *C. parasitica* in

(Demené et al. 2022) contained in the "Curated_TEs_database_ESM015_EP155.fa" file

(available on the "Portail Data INRAE: Chromosomal rearrangements but no change of genes
and transposable elements repertoires in an invasive forest-pathogenic fungus" at
https://doi.org/10.15454/UTIB8U). The class Gipsy invader was renamed LTR-Ty3. We
filtered out TE copies shorter than 100bp. We used the HybridSPAdes preliminary assembly
of MRC10 for gene model annotations. To study gene functions of the EP155 genome, we
used the gene annotation available at http://jgi.doe.gov/ (Crouch et al. 2020). To study support
from RNAseq data, we used RNAseq data from the EP155 strain cultivated *in vitro* available
at Genebank under Project ID number PRJNA588887 and accessions numbers SRR10428542,
SRR10428543, SRR10428544 (Chun et al. 2020). Raw reads were mapped using STAR
v2.7.10a (Dobin et al. 2013) with the settings. --alignIntronMax 1000 --limitBAMsortRAM
1629219965--quantMode GeneCounts. We used the program featureCounts (Liao et al. 2013)
v2.0.6 with the options -p --countReadPairs -M - -B -O --largestOverlap. We considered a
gene to be supported for read count >10. To predict gene functions of the protein predicted by
Helixer, we used the funannotate pipeline with default options. To look for *Starships*, we ran
the Gene Finder Module of Starfish pipeline with default options (Gluck-Thaler and Vogan
2024). To predict gene functions of the protein predicted by Helixer, we used the funannotate
pipeline with default options. To identify putative *Starships*, we ran the Starfish pipeline with
default options (Gluck-Thaler and Vogan 2024). We studied the phylogenetic relationships of
the putative Starships captains by aligning them to the YRsuperfamRefs.faa from the starfish
database (Gluck-Thaler and Vogan 2024). Protein sequences were aligned using Clustal
Omega (Sievers and Higgins 2021)version 1.2.4 allowing for five iterations (--iterations 5).
Gaps in the resulting alignment were trimmed using trimAl v1.4.rev15 and the -gappyout
option. The phylogenetic relationship among proteins was inferred from the trimmed
alignment using FastTree (Price et al. 2009) under the Whelan-And-Goldman 2001 model
after 1,000 bootstraps (-boot 1000 and -wag options). Plots were performed in R v 4.1.2 using
ggtree v3.9.1.
958

959 **Estimation of retrotranposon insertion time and nucleotide divergence time**
960
To get an estimation of the insertion date of the transposable elements present in the MAT-
proximal non-recombining region, we applied two complementary methods. We first used the
divergence between the LTR sequences in retrotransposons, as these LTR sequences at their
edges are identical at the time of TE insertion and then diverge with time. For this, we used a
de-novo prediction of LTR retrotransposons using LTRharvest GenomeTools 1.6.2

(Ellinghaus et al. 2008; Gremme et al. 2013) and LTR_Finder v1.07 (Xu and Wang 2007). To

prepare data for LTRharvest, we first created enhanced suffix for the M1400 and M6697

genome assemblies using the GenomeTools Suffixerator (-dna -suf -lcp -bwt -bck -mirrored

parameters). We ran LTRharvest using two pipelines, designed to identify LTR

retrotransposons with a minimum and maximum LTR length of 100 bp and 7000 bp

respectively and at least 85% identity between two LTR regions, with and without canonical

TGCA motifs, respectively: i) -minlenltr 100 -maxlenltr 7000 -mintsd 4 -maxtsd 6 -similar 85

-vic 10 -seed 20 -motif TGCA -motifmis 1; ii) -minlentltr 100, -maxlenltr 7000, -mintsd 4, -

maxtsd 6, -similar 85, -vic 10, -seed 20. Similarly, we ran LTR_Finder on the M1400 and

M6697 genome assemblies to retrotransposons with both TGCA and non-TGCA motifs and a

minimum and maximum LTR length of 100 bp and 7000 bp respectively and at least 85%

identity between two LTR regions (-D 15000, -d 1000, -l 100, -L 7000, -p 20, -C, -M 0.85).

Finally, we used LTR_retriever v2.8.5 (Ou and Jiang 2018) with default parameters to filter

out false positive LTR candidates identified by LTRharvest and LTR_Finder and get an

estimation of each element insertion date.

As a second method to estimate the insertion date of the transposable elements present in the

MAT-proximal non-recombining region, we used the set of curated consensus sequences from

(Demené et al. 2022) to annotate the inversion sequence or its surroundings. We first used

samtools faidx v1.9 (Li et al. 2009) to extract the sequence corresponding to inversion and

their 50 kb surroundings in M1400 and M6697 genome assemblies. We annotated the

sequences corresponding to the inversions and the concatenated 50 kb surroundings in both

isolates using RepeatMasker version 4.1.5 and rmblast as search engine (v2.10.0) with -no_is

-pa 20 -cutoff 250 -a parameters. Finally, we parsed the RepeatMasker .out file using the

helped script parseRM_merge_interrupted.pl and omitting Simple_repeat and

Low_complexity regions (https://github.com/4ureliek/Parsing-RepeatMasker-Outputs). We

then built a summary of the alignments using the RepeatMasker helper script

buildSummary.pl and calculated sequence divergence using the calcDivergenceFromAlign.pl

script to finally render results with the createRepeatLandscape.pl from the same helper suite.

Comparative genomics analyses

Genome synteny between long reads assemblies were studied using the nucmer v3.1 program

(Marçais et al. 2018). Outputs were plotted with the R programs ggplot2 (Wickham 2009),

genoPlotR (Guy et al. 2010) and RIdeogram (Hao et al. 2020). The dotplot in the putative centromere region of the M1400 mating-type contig was performed using the online megablast alignment tool available at https://blast.ncbi.nlm.nih.gov/(last accessed 13th May 2024).

To study gene disruption, synonymous and non-synonymous divergence (d_S and d_N) and testing introgression in the MAT-proximal region, we build orthology relationships for genes of the genome assemblies of *C. parasitica* strains M1400, M6697, XIM9508, MRC10, DUM-005, ESM15 and EP155 and included three genomes of the closely related species *Cryphonectria japonica* (IF-6), *Cryphonectria carpinicola* (CS3), and *Cryphonectria naterciae* (M3664). Genome, gene annotation and species tree of these closely related species were previously published by (Stauber et al. 2020). Genome data were retrieved from NCBI bioproject number PRJNA644891 and accession IDs JACWRX000000000 for IF-6, JACWRQ000000000 for CS3 and JACWST000000000 for M3664. We run OrthoFinder v2.3.7 (Emms and Kelly 2019) analysis on protein sequences. We used the translatorX v1.1 program (Abascal et al. 2010) with default parameters that use a codon-based approach to align orthologous gene coding sequences. To compute d_S and d_N vaues of one-to-one orthologs between pairs of genome assemblies, we use the yn00 program of PAML (Yang 2007). Estimation of divergence time between haplotypes was performed using computed gene d_S values and the formula Tgenerations = dS/2μ. We used previous estimates of substitution rates in fungi (Kasuga et al. 2002; Taylor and Berbee 2006) and considered that *C. parasitica* undergoes one generation a year (Guerin et al. 2001). We build gene coding sequences trees with the outgroup genomes in the MAT-proximal region using iqtree2 v2.2.2.6 (Minh et al. 2020) with 1000 bootstraps and used the Newick Utilities (https://github.com/tjunier/newick_utils) for displaying phylogenetics tree.

References

Abascal F, Zardoya R, Telford MJ. 2010. TranslatorX: multiple alignment of nucleotide sequences guided by amino acid translations. *Nucleic Acids Res.* 38:W7-13.

Abu-Awad D, Waller D. 2023. Conditions for maintaining and eroding pseudo-overdominance and its contribution to inbreeding depression. *Peer Community J.* [Internet] 3. Available from: https://peercommunityjournal.org/articles/10.24072/pcjournal.224/

Alonge M, Soyk S, Ramakrishnan S, Wang X, Goodwin S, Sedlazeck FJ, Lippman ZB, Schatz MC. 2019. RaGOO: fast and accurate reference-guided scaffolding of draft genomes. *Genome Biol.* 20:224.

Anagnostakis SL. 1987. Chestnut blight: The classical problem of an introduced pathogen.
Mycologia [Internet]. Available from:
https://www.tandfonline.com/doi/abs/10.1080/00275514.1987.12025367

Antipov D, Korobeynikov A, McLean JS, Pevzner PA. 2016. hybridSPAdes: an algorithm for
hybrid assembly of short and long reads. *Bioinformatics* 32:1009–1015.

Bachtrog D. 2013. Y-chromosome evolution: emerging insights into processes of Y-
chromosome degeneration. *Nat. Rev. Genet.* 14:113–124.

Bazzicalupo AL, Carpentier F, Otto SP, Giraud T. 2019. Little evidence of antagonistic
selection in the evolutionary strata of fungal mating-type chromosomes
(*Microbotryum lychnidis-dioicae*). *G3 Genes Genomes Genet.* 9:1987–1998.

Belov AA, Witte TE, Overy DP, Smith ML. 2021. Transcriptome analysis implicates
secondary metabolite production, redox reactions, and programmed cell death during
allorecognition in Cryphonectria parasitica. *G3 GenesGenomesGenetics* 11:jkaa021.

Berdan EL, Blanckaert A, Butlin RK, Flatt T, Slotte T, Wielstra B. 2022. Mutation
accumulation opposes polymorphism: supergenes and the curious case of balanced
lethals. *Philos. Trans. R. Soc. B Biol. Sci.* 377:20210199.

Bergero R, Charlesworth D. 2009. The evolution of restricted recombination in sex
chromosomes. *Trends Ecol. Evol.* 24:94–102.

Boideau F, Richard G, Coriton O, Huteau V, Belser C, Deniot G, Eber F, Falentin C, Ferreira
de Carvalho J, Gilet M, et al. 2022. Epigenomic and structural events preclude
recombination in Brassica napus. *New Phytol.* 234:545–559.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* 30:2114–2120.

Branco S, Badouin H, Rodríguez de la Vega RC, Gouzy J, Carpentier F, Aguileta G, Siguenza
S, Brandenburg J-T, Coelho MA, Hood ME, et al. 2017. Evolutionary strata on young
mating-type chromosomes despite the lack of sexual antagonism. *Proc. Natl. Acad.*
Sci. 114:7067–7072.

Branco S, Carpentier F, Rodríguez de la Vega RC, Badouin H, Snirc A, Prieur SL, Coelho
MA, Vienne DM de, Hartmann FE, Begerow D, et al. 2018. Multiple convergent
supergene evolution events in mating-type chromosomes. *Nat. Commun.* 9:2000.

Brusini J, Wayne ML, Franc A, Robin C. 2017. The impact of parasitism on resource
allocation in a fungal host: the case of *Cryphonectria parasitica* and its mycovirus,
Cryphonectria Hypovirus 1. *Ecol. Evol.* 7:5967–5976.

Bucknell A, Wilson HM, Santos KCG do, Simpfendorfer S, Milgate A, Germain H, Solomon
PS, Bentham A, McDonald MC. 2024. Sanctuary: A Starship transposon facilitating
the movement of the virulence factor ToxA in fungal wheat pathogens.
:2024.03.04.583430. Available from:
https://www.biorxiv.org/content/10.1101/2024.03.04.583430v1

Carpentier F, Rodríguez de la Vega RC, Jay P, Duhamel M, Shykoff JA, Perlin MH, Wallen
RM, Hood ME, Giraud T. 2022. Tempo of degeneration across independently evolved
nonrecombining regions. *Mol. Biol. Evol.* 39:msac060.

Casacuberta E, González J. 2013. The impact of transposable elements in environmental
adaptation. *Mol. Ecol.* 22:1503–1517.

Charlesworth B. 2023. The fitness consequences of genetic divergence between polymorphic
gene arrangements. *Genetics*:iyad218.

Charlesworth D. 2016. The status of supergenes in the 21st century: recombination
suppression in Batesian mimicry and sex chromosomes and other complex
adaptations. *Evol. Appl.* 9:74–90.

Charlesworth D, Vekemans X, Castric V, Glémin S. 2005. Plant self-incompatibility systems:
a molecular evolutionary perspective. *New Phytol.* 168:61–69.

Cheeseman K, Ropars J, Renault P, Dupont J, Gouzy J, Branca A, Abraham A-L, Ceppi M,
Conseiller E, Debuchy R, et al. 2014. Multiple recent horizontal transfers of a large
genomic region in cheese making fungi. *Nat. Commun.* 5:2876.

Choi GH, Dawe AL, Churbanov A, Smith ML, Milgroom MG, Nuss DL. 2012. Molecular
characterization of vegetative incompatibility genes that restrict hypovirus
transmission in the chestnut blight fungus *Cryphonectria parasitica*. *Genetics*
190:113–127.

Choi GH, Nuss DL. 1992. Hypovirulence of chestnut blight fungus conferred by an infectious
viral cDNA. *Science* 257:800–803.

Chun J, Ko Y-H, Kim D-H. 2020. Transcriptome Analysis of Cryphonectria parasitica
Infected With Cryphonectria hypovirus 1 (CHV1) Reveals Distinct Genes Related to
Fungal Metabolites, Virulence, Antiviral RNA-Silencing, and Their Regulation. *Front.*
Microbiol. 11:1711.

Crouch JA, Dawe A, Aerts A, Barry K, Churchill ACL, Grimwood J, Hillman BI, Milgroom
MG, Pangilinan J, Smith M, et al. 2020. Genome sequence of the chestnut blight
fungus Cryphonectria parasitica EP155: A fundamental resource for an archetypical
invasive plant pathogen. *Phytopathology®* 110:1180–1188.

Dainat J, Hereñú D, Pucholt P. 2020. AGAT:Another Gff Analysis Toolkit to handle
annotations in any GTF. *GFF Format*.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter
G, Marth GT, Sherry ST, et al. 2011. The variant call format and VCFtools.
Bioinformatics 27:2156–2158.

Demené A, Laurent B, Cros-Arteil S, Boury C, Dutech C. 2022. Chromosomal
rearrangements with stable repertoires of genes and transposable elements in an
invasive forest-pathogenic fungus. *Peer Community J.* [Internet] 2. Available from:
https://peercommunityjournal.org/articles/10.24072/pcjournal.127/

Demené A, Legrand L, Gouzy J, Debuchy R, Saint-Jean G, Fabreguettes O, Dutech C. 2019.
Whole-genome sequencing reveals recent and frequent genetic recombination between
clonal lineages of Cryphonectria parasitica in western Europe. *Fungal Genet. Biol.*
130:122–133.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,
Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*
29:15–21.

Dufresnes C, Borzée A, Horn A, Stöck M, Ostini M, Sermier R, Wassef J, Litvinchuck SN,
Kosch TA, Waldman B, et al. 2015. Sex-chromosome homomorphy in palearctic tree
frogs results from both turnovers and X-Y recombination. *Mol. Biol. Evol.* 32:2328–
2337.

Duhamel M, Hood ME, Vega RCR de la, Giraud T. 2023. Dynamics of transposable element
accumulation in the non-recombining regions of mating-type chromosomes in anther-
smut fungi. :2022.08.03.502670. Available from:
https://www.biorxiv.org/content/10.1101/2022.08.03.502670v2

Dussert Y, Legrand L, Mazet ID, Couture C, Piron M-C, Serre R-F, Bouchez O, Mestre P,
Toffolatti SL, Giraud T, et al. 2020. Identification of the first oomycete mating-type
locus sequence in the grapevine downy mildew pathogen, *Plasmopara viticola. Curr.*
Biol. 30:3897-3907.e4.

Dutech C, Barrès B, Bridier J, Robin C, Milgroom MG, Ravigné V. 2012. The chestnut blight
fungus world tour: successive introduction events from diverse origins in an invasive
plant fungal pathogen. *Mol. Ecol.* 21:3931–3946.

Dutech C, Fabreguettes O, Capdevielle X, Robin C. 2010. Multiple introductions of divergent
genetic lineages in an invasive fungal pathogen, Cryphonectria parasitica, in France.
Heredity 105:220–228.

Ellinghaus D, Kurtz S, Willhoeft U. 2008. LTRharvest, an efficient and flexible software for
de novo detection of LTR retrotransposons. *BMC Bioinformatics* 9:18.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biol.* 20:238.

Fredericksen M, Fields PD, Pasquier LD, Ricci V, Ebert D. 2023. QTL study reveals
candidate genes underlying host resistance in a Red Queen model system. *PLOS*
Genet. 19:e1010570.

Furman BLS, Metzger DCH, Darolti I, Wright AE, Sandkam BA, Almeida P, Shu JJ, Mank
JE. 2020. Sex chromosome evolution: So many exceptions to the rules. *Genome Biol.*
Evol.

Gluck-Thaler E, Ralston T, Konkel Z, Ocampos CG, Ganeshan VD, Dorrance AE, Niblack
TL, Wood CW, Slot JC, Lopez-Nicora HD, et al. 2022. Giant Starship Elements
Mobilize Accessory Genes in Fungal Genomes. *Mol. Biol. Evol.* 39:msac109.

Gluck-Thaler E, Vogan AA. 2024. Systematic identification of cargo-mobilizing genetic
elements reveals new dimensions of eukaryotic diversity. *Nucleic Acids Res.* 52:5496–
5513.

Gremme G, Steinbiss S, Kurtz S. 2013. GenomeTools: A Comprehensive Software Library
for Efficient Processing of Structured Genome Annotations. *IEEE/ACM Trans.*
Comput. Biol. Bioinform. 10:645–656.

Grognet P, Bidard F, Kuchly C, Tong LCH, Coppin E, Benkhali JA, Couloux A, Wincker P,
Debuchy R, Silar P. 2014. Maintaining two mating types: structure of the mating type
locus and its role in heterokaryosis in *Podospora anserina. Genetics* 197:421–432.

Guerin L, Froidefond G, Xu X-M. 2001. Seasonal patterns of dispersal of ascospores of
Cryphonectria parasitica (chestnut blight). *Plant Pathol.* 50:717–724.

Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for
genome assemblies. *Bioinformatics* 29:1072–1075.

Guy L, Roat Kultima J, Andersson SGE. 2010. genoPlotR: comparative gene and genome
visualization in R. *Bioinformatics* 26:2334–2335.

Guyot L, Chahine E, Lalanne C, Hartmann FE, Giraud T. 2024. Sheltered load in fungal
mating-type chromosomes revealed by fitness experiments. :2024.09.10.612177.
Available from: https://www.biorxiv.org/content/10.1101/2024.09.10.612177v1

Hao Z, Lv D, Ge Y, Shi J, Weijers D, Yu G, Chen J. 2020. RIdeogram: drawing SVG
graphics to visualize and map genome-wide data on the idiograms. *PeerJ Comput. Sci.*
6:e251.

Hartmann FE, Duhamel M, Carpentier F, Hood ME, Foulongne-Oriol M, Silar P, Malagnac F,
Grognet P, Giraud T. 2021. Recombination suppression and evolutionary strata around
mating-type loci in fungi: documenting patterns and understanding evolutionary and
mechanistic causes. *New Phytol.* 229:2470–2491.

Helleu Q, Roux C, Ross KG, Keller L. 2022. Radiation and hybridization underpin the spread
of the fire ant social supergene. *Proc. Natl. Acad. Sci.* 119:e2201040119.

Hoegger PJ, Rigling D, Holdenrieder O, Heiniger U. 2000. Genetic structure of newly
established populations of *Cryphonectria parasitica. Mycol. Res.* 104:1108–1116.

Holst F, Bolger A, Günther C, Maß J, Triesch S, Kindel F, Kiel N, Saadat N, Ebenhöh O,
Usadel B, et al. 2023. Helixer–de novo prediction of primary Eukaryotic gene models
combining deep learning and a hidden markov model. :2023.02.06.527280. Available
from: https://www.biorxiv.org/content/10.1101/2023.02.06.527280v2

Huson DH. 1998. SplitsTree: analyzing and visualizing evolutionary data. *Bioinforma. Oxf.*
Engl. 14:68–73.

Ironside JE. 2010. No amicable divorce? Challenging the notion that sexual antagonism
drives sex chromosome evolution. *BioEssays* 32:718–726.

Jay P, Aubier TG, Joron M. 2024. The interplay of local adaptation and gene flow may lead to
the formation of supergenes. *Mol. Ecol.*:e17297.

Jay P, Chouteau M, Whibley A, Bastide H, Parrinello H, Llaurens V, Joron M. 2021.
Mutation load at a mimicry supergene sheds new light on the evolution of inversion
polymorphisms. *Nat. Genet.* 53:288–293.

Jay P, Jeffries D, Hartmann FE, Véber A, Giraud T. 2024. Why do sex chromosomes
progressively lose recombination? *Trends Genet.* [Internet]. Available from:
https://www.sciencedirect.com/science/article/pii/S0168952524000672

Jay P, Jeffries DL, Hartmann FE, Véber A, Giraud T. 2024. Why do sex chromosomes
progressively lose recombination? *Trends in Genetics*:in press.

Jay P, Tezenas E, Véber A, Giraud T. 2022. Sheltering of deleterious mutations explains the
stepwise extension of recombination suppression on sex chromosomes and other
supergenes. *PLOS Biol.* 20:e3001698.

Jay P, Whibley A, Frézal L, Rodríguez de Cara MÁ, Nowell RW, Mallet J, Dasmahapatra
KK, Joron M. 2018. Supergene evolution triggered by the introgression of a
chromosomal inversion. *Curr. Biol.* 28:1839-1845.e3.

Kasuga T, White TJ, Taylor JW. 2002. Estimation of nucleotide substitution rates in
eurotiomycete fungi. *Mol. Biol. Evol.* 19:2318–2324.

Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, error-prone reads using
repeat graphs. *Nat. Biotechnol.* 37:540–546.

Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu: scalable
and accurate long-read assembly via adaptive k-mer weighting and repeat separation.
Genome Res. 27:722–736.

Kubisiak TL, Milgroom MG. 2006a. Markers linked to vegetative incompatibility (vic) genes
and a region of high heterogeneity and reduced recombination near the mating type
locus (MAT) in *Cryphonectria parasitica*. *Fungal Genet. Biol.* 43:453–463.

Kubisiak TL, Milgroom MG. 2006b. Markers linked to vegetative incompatibility (*vic*) genes
and a region of high heterogeneity and reduced recombination near the mating type
locus (*MAT*) in *Cryphonectria parasitica*. *Fungal Genet. Biol. FG B* 43:453–463.

Küpper C, Stocks M, Risse JE, Remedios N dos, Farrell LL, McRae SB, Morgan TC,
Karlionova N, Pinchuk P, Verkuil YI, et al. 2016. A supergene determines highly
divergent male reproductive morphs in the ruff. *Nat. Genet.* 48:79–83.

Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. *Genome Biol.* 10:R25.

Legrand S, Saifudeen A, Bordelet H, Vernerey J, Guille A, Bignaud A, Thierry A, Acquaviva
L, Gaudin M, Sanchez A, et al. 2024. Absence of chromosome axis protein
recruitment prevents meiotic recombination chromosome-wide in the budding yeast
Lachancea kluyveri. *Proc. Natl. Acad. Sci.* 121:e2312820121.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin
R. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics*
25:2078–2079.

Liao Y, Smyth GK, Shi W. 2013. The Subread aligner: fast, accurate and scalable read
mapping by seed-and-vote. *Nucleic Acids Res.* 41:e108–e108.

Lischer HEL, Excoffier L. 2012. PGDSpider: an automated data conversion tool for
connecting population genetics and genomics programs. *Bioinformatics* 28:298–299.

Liu YC, Linder-Basso D, Hillman BI, Kaneko S, Milgroom MG. 2003. Evidence for
interspecies transmission of viruses in natural populations of filamentous fungi in the
genus Cryphonectria. *Mol. Ecol.* 12:1619–1628.

Lo Y-C, Bruxaux J, Rodríguez de la Vega RC, O'Donnell S, Snirc A, Coton M, Le Piver M,
Le Prieur S, Roueyre D, Dupont J, et al. 2023. Domestication in dry-cured meat
Penicillium fungi: Convergent specific phenotypes and horizontal gene transfers
without strong genetic subdivision. *Evol. Appl.* 16:1637–1660.

Lovat C-A, Donnelly DJ. 2019. Mechanisms and metabolomics of the host–pathogen
interactions between Chestnut (Castanea species) and Chestnut blight (Cryphonectria
parasitica). *For. Pathol.* 49:e12562.

Maloisel L, Rossignol JL. 1998. Suppression of crossing-over by DNA methylation in
Ascobolus. *Genes Dev.* 12:1381–1389.

Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021. BUSCO update: Novel
and streamlined workflows along with broader and deeper phylogenetic coverage for
scoring of eukaryotic, prokaryotic, and viral genomes. *Mol. Biol. Evol.* 38:4647–4654.

Marçais G, Delcher AL, Phillippy AM, Coston R, Salzberg SL, Zimin A. 2018. MUMmer4:
A fast and versatile genome alignment system. *PLOS Comput. Biol.* 14:e1005944.

McGuire IC, Davis JE, Double ML, MacDonald WL, Rauscher JT, McCawley S, Milgroom
MG. 2005. Heterokaryon formation and parasexual recombination between
vegetatively incompatible lineages in a population of the chestnut blight fungus,
Cryphonectria parasitica. *Mol. Ecol.* 14:3657–3669.

McGuire IC, Marra RE, Milgroom MG. 2004. Mating-type heterokaryosis and selfing in
Cryphonectria parasitica. *Fungal Genet. Biol.* 41:521–533.

McGuire IC, Marra RE, Turgeon BG, Milgroom MG. 2001. Analysis of mating-Type genes
in the chestnut blight fungus, *Cryphonectria parasitica*. *Fungal Genet. Biol.* 34:131–
144.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K,
Altshuler D, Gabriel S, Daly M, et al. 2010. The Genome Analysis Toolkit: a
MapReduce framework for analyzing next-generation DNA sequencing data. *Genome*
Res. 20:1297–1303.

Menkis A, Jacobson DJ, Gustafsson T, Johannesson H. 2008. The mating-type chromosome
in the filamentous ascomycete *Neurospora tetrasperma* represents a model for early
evolution of sex chromosomes. *PLOS Genet.* 4:e1000030.

Milgroom MG, Wang K, Zhou Y, Lipari SE, Kaneko S. 1996. Intercontinental population
structure of the chestnut blight fungus, *Cryphonectria parasitica*. *Mycologia* 88:179–
190.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A,
Lanfear R. 2020. IQ-TREE 2: New models and efficient methods for phylogenetic
inference in the genomic era. *Mol. Biol. Evol.* 37:1530–1534.

Moinet A, Schlichta F, Peischl S, Excoffier L. 2022. Strong neutral sweeps occurring during a
population contraction. *Genetics* 220:iyac021.

Orteu A, Kucka M, Gordon IJ, Ng'iru I, van der Heijden ESM, Talavera G, Warren IA,
Collins S, ffrench-Constant RH, Martins DJ, et al. 2024. Transposable element
insertions are associated with Batesian mimicry in the pantropical butterfly
Hypolimnas misippus. *Mol. Biol. Evol.* 41:msae041.

Otto SP, Lenormand T. 2002. Resolving the paradox of sex and recombination. *Nat. Rev.*
Genet. 3:252–261.

Ou S, Jiang N. 2018. LTR_retriever: A highly accurate and sensitive program for
identification of long terminal repeat retrotransposons. *Plant Physiol.* 176:1410–1422.

Peck LD, Llewellyn T, Bennetot B, O'Donnell S, Nowell RW, Ryan MJ, Flood J, Vega RCR
de la, Ropars J, Giraud T, et al. 2023. Horizontal transfers between fungal Fusarium
species contributed to successive outbreaks of coffee wilt disease.
:2023.12.22.572981. Available from:
https://www.biorxiv.org/content/10.1101/2023.12.22.572981v1

Pegueroles C, Ordóñez V, Mestres F, Pascual M. 2010. Recombination and selection in the
maintenance of the adaptive value of inversions. *J. Evol. Biol.* 23:2709–2717.

Pfeifer B, Wittelsbürger U, Ramos-Onsins SE, Lercher MJ. 2014. PopGenome: an efficient
Swiss army knife for population genomic analyses in R. *Mol. Biol. Evol.* 31:1929–
1936.

Price MN, Dehal PS, Arkin AP. 2009. FastTree: Computing Large Minimum Evolution Trees
with Profiles instead of a Distance Matrix. *Mol. Biol. Evol.* 26:1641–1650.

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P,
de Bakker PIW, Daly MJ, et al. 2007. PLINK: A tool set for whole-genome
association and population-based linkage analyses. *Am. J. Hum. Genet.* 81:559–575.

Rigling D, Prospero S. 2018. Cryphonectria parasitica, the causal agent of chestnut blight:
invasion history, population biology and disease control. *Mol. Plant Pathol.* 19:7–20.

Ropars J, Rodríguez de la Vega RC, López-Villavicencio M, Gouzy J, Sallet E, Dumas É,
Lacoste S, Debuchy R, Dupont J, Branca A, et al. 2015. Adaptive horizontal gene
transfers between multiple cheese-associated fungi. *Curr. Biol.* 25:2562–2569.

Schwander T, Libbrecht R, Keller L. 2014. Supergenes and complex phenotypes. *Curr. Biol.*
24:R288–R294.

Shin J-H, Blay S, McNeney B, Graham J, others. 2006. LDheatmap: an R function for
graphical display of pairwise linkage disequilibria between single nucleotide
polymorphisms. *J. Stat. Softw.* 16:1–10.

Sievers F, Higgins DG. 2021. The Clustal Omega Multiple Alignment Package. In: Katoh K,
editor. Multiple Sequence Alignment: Methods and Protocols. New York, NY:
Springer US. p. 3–16. Available from: https://doi.org/10.1007/978-1-0716-1036-7_1

Smit AFA, Hubley R, Green P. 2013. RepeatMasker Open-4.0. Available at:
http://www.repeatmasker.org Accessed: November 30, 2017.

Stauber L, Badet T, Feurtey A, Prospero S, Croll D. 2021. Emergence and diversification of a
highly invasive chestnut pathogen lineage across southeastern Europe.Castric V,
Weigel D, Laine A-L, Castric V, editors. *eLife* 10:e56279.

Stauber L, Croll D, Prospero S. 2022. Temporal changes in pathogen diversity in a perennial
plant–pathogen–hyperparasite system. *Mol. Ecol.* 31:2073–2088.

Stauber L, Prospero S, Croll D. 2020. Comparative genomics analyses of lifestyle transitions
at the origin of an invasive fungal pathogen in the genus *Cryphonectria*. *mSphere*
5:10.1128/msphere.00737-20.

Stevison LS, Hoehn KB, Noor MAF. 2011. Effects of Inversions on Within- and Between-
Species Recombination and Divergence. *Genome Biol. Evol.* 3:830–841.

Stolle E, Pracana R, López-Osorio F, Priebe MK, Hernández GL, Castillo-Carrillo C, Arias
MC, Paris CI, Bollazzi M, Priyam A, et al. 2022. Recurring adaptive introgression of a
supergene variant that determines social organization. *Nat. Commun.* 13:1180.

Sun Y, Svedberg J, Hiltunen M, Corcoran P, Johannesson H. 2017. Large-scale suppression
of recombination predates genomic rearrangements in *Neurospora tetrasperma*. *Nat.*
Commun. 8:1–8.

Taylor JW, Berbee ML. 2006. Dating divergences in the Fungal Tree of Life: review and new
analyses. *Mycologia* 98:838–849.

Tellier A, Moreno-Gámez S, Stephan W. 2014. Speed of adaptation and genomic footprints of
host–parasite coevolution under arms race and trench warfare dynamics. *Evolution*
68:2211–2224.

Tralamazza SM, Gluck-Thaler E, Feurtey A, Croll D. 2024. Copy number variation
introduced by a massive mobile element facilitates global thermal adaptation in a
fungal wheat pathogen. *Nat. Commun.* 15:5728.

Umen JG. 2011. Evolution of sex and mating loci: An expanded view from Volvocine algae.
Curr. Opin. Microbiol. 14:634–641.

Urquhart A, Vogan AA, Gluck-Thaler E. 2024. Starships: a new frontier for fungal biology.
Trends Genet. [Internet] 0. Available from:
https://www.cell.com/trends/genetics/abstract/S0168-9525(24)00183-5

Vittorelli N, Hartmann, Giraud T. 2022. Stepwise recombination suppression around the
mating-type locus in the fungus *Schizothecium tetrasporum* (Ascomycota,
Sordariales).

Wickham H. 2009. ggplot2: elegant graphics for data analysis. Springer Science & Business
Media

Xu Z, Wang H. 2007. LTR_FINDER: an efficient tool for the prediction of full-length LTR
retrotransposons. *Nucleic Acids Res.* 35:W265–W268.

Yan Z, Martin SH, Gotzek D, Arsenault SV, Duchen P, Helleu Q, Riba-Grognuz O, Hunt BG,
Salamin N, Shoemaker D, et al. 2020. Evolution of a supergene that regulates a trans-
species social polymorphism. *Nat. Ecol. Evol.* 4:240–249.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.*
24:1586–1591.

1352

1353

1354

Acknowledgments

This work was supported by the European Research Council (ERC) EvolSexChrom (832352) grant

and a Fondation Louis D grant from the Institut de France to TG, and by the ANR PIA grant # ANR-

20-IDEES-0002 to F.E.H. We acknowledge the GenOuest bioinformatics core facility

(https://www.genouest.org) for providing the computing infrastructure, (GPU) for genome

annotation". We thank Jeanne Ropars, Aaron Vogan and Marie Foulongne for discussions and

suggestions on the analyses. The authors declare that they have no competing interests.

1362

1363 TG and FEH conceptualized the study and acquired funding. TG supervised the study. AL, AS, CD,

1364 AD, SP, DC, LS and TB contributed to data and their availability. AL and AS performed *in vitro*

1365 culture and DNA extraction. FEH, RRDLV, QR, JPV, LS and TB analyzed genomes. FEH and TG

1366 wrote the original draft. All authors edited the manuscript.

1367

1368 The raw data and the new assemblies produced in this study will be published pending scientific

1369 review.

1370

1371 **Tables**

Table 1: Distribution of mating types (*MAT-1* and *MAT-2*) among non-recombining haplotypes of the MAT-proximal region in the European invasive *Cryphonectria parasitica* CL1 genetic cluster and the Swiss populations. MAT-Prox1 and MAT-Prox2 haplotypes were defined based on the clusters of the principal component analysis.

Figures

Figure 1: Linkage disequilibrium (LD) analysis along the contig carrying the mating-type locus in a *Cryphonectria parasitica* European invasive population. LD heatmaps using single nucleotide polymorphisms (SNPs; n=3815) located on the contig carrying the mating-type locus (scaffold_2 of the EP155 genome) in the CL1 genetic cluster (European invasive population introduced from North America); pairs of SNPs with high LD, i.e. r^2 >0.9, correspond to the red triangle. The mating-type locus location is shown with a green triangle and the MAT-proximal region lacking recombination is shown with a red arrow. The two high-LD blocks within the MAT-proximal region are shown with orange arrows. SNPs at the limit of the MAT-proximal region and the two high LD blocks were manually highlighted with red dotted lines and an orange line.

Figure 2: Genetic structure using single nucleotide polymorphisms (SNPs) in the MAT-proximal region lacking recombination and other genomic regions in a *Cryphonectria parasitica* European invasive population. A-C. Principal component analysis (PCA). Two principal components are presented. Percentage of variance explained by each PC is indicated into brackets. Strains are colored according to their mating type (*MAT-1* or *MAT-2*). **D-E.** Neighbor-net network from a SplitsTree analysis. These analyses were performed in the CL1 genetic cluster (European invasive populations introduced from North America) based on: **A and D.** SNPs (n=2,220) located within the MAT-proximal region along the contig carrying the mating-type locus (scaffold_2 of the EP155 genome).; **B.** SNPs (n=1,595) located in other regions along the contig carrying the mating-type locus; **C and E** SNPs (n=11,289) located on other contigs of the EP155 genome. On panels A and D, the two clusters corresponding to the MAT-Prox1 and MAT-Prox2 haplotypes are shown with red circles. The identified haplotype of each strain is indicated in Table S1. The number of strains within the MAT-Prox1 haplotype is indicated on panel A by the letter n. The newly sequenced M1400 and M6697 strains are highlighted with a black rectangle

Figure 3: **Synteny and rearrangements between the two newly sequenced genomes of European invasive strains introduced from North America (M1400 and M6697) in the chromosome carrying the MAT-proximal region lacking recombination in** *Cryphonectria parasitica.* Blue links show colinear regions and orange links show inverted regions in the MAT-proximal region. Grey links show other regions. The mating-type locus is located with a green diamond. The MAT-proximal region defined from LD analyses is indicated with red arrows. Gene density tracks is shown with a color gradient (blue with low density, orange with high density).

Figure 4: **Genetic diversity and divergence between non-recombining haplotypes of the MAT-proximal region along the contig carrying the mating-type locus in a** *Cryphonectria parasitica* **European invasive population.** A-B. GC content (A) and transposable element densities (B) along the contig carrying the mating-type locus (tig_0000001) of the M1400 *C. parasitica* genome. C. Relative divergence (F_{ST}) between strains of the MAT-Prox1 and MAT-Prox2 haplotypes: D-E. Nucleotide diversity within pools of strains for each MAT-Proximal haplotype; F-G-H Tajima's D for all strains pooled and within pools of strains for each MAT-Proximal haplotype; The MAT-proximal region defined from LD analyses and the inversion between M1400 and M6697 genomes are indicated with red and blue arrows respectively. The mating-type locus location is shown with a green triangle. The location of the putative centromere is indicated by a yellow circle. All population statistics were computed for the 1990 Swiss population along the mating-type contig (tig_0000001) of the M1400 genome per 50-kb window overlapping over 10 kbp. Windows containing fewer than 5 SNPs were removed from the analysis.

Figure 5: **Annotation of transposable elements (TEs) and estimates of their insertion time in the MAT-proximal region lacking recombination and other genomic regions in** *Cryphonectria parasitica.* A. Annotation of transposable elements (TEs) and gene density along the MAT-proximal region in M1400 and M6697 genomes. On the left panel (1) the figure shows the genomic region from 7 Mb on tig_0000001 of the M1400 genome and from 2.8 Mb on tig_0000060 of the M6697 genome. The mating-type locus is located with a green triangle. The two high-LD blocks within the MAT-proximal region are shown with orange arrows. Synteny for 10 kb segments with identity > 90 % is shown in red and inversion in blue. Transposons larger than 5 kb are shown in purple for the transposons annotated as LTR-Ty3 by and in green for other transposons. Orthologous genes shared between genomes are

shown in black and unique to each each genome in grey. Genes with DU3435 domains (putative starship captains) are shown in red. On the right panel (2) the figure shows a zoom on the starship present in M1400 and absent in M6697. Transposons larger than 150 pb are shown in orange; orthologous genes shared between genomes are shown in black and unique to one genome in grey and genes with DU3435 domains are shown in red. Synteny for segments with identity > 90 % is shown in red and inversion in blue. B. Relative percentage of each TE annotation (in %) in the non-recombining MAT-proximal region and other recombining regions of M6697 and M1400 genomes. C. Pairwise genetic distance between TE copies within the inversion and around (50 kb) in both M1400 and M6697 genomes. Distribution of the Kimura substitution levels computed using the consensus sequence for the TEs. D. Estimates of the age of intact copies of LTR within and around the inversion.

Figure 6: Transposable elements content and Starship content in high quality genome assemblies from other *Cryphonectria parasitica* populations from the native and the introduced range. A. TE load (percentage of base pairs occupied by transposable elements) in the seven high quality genome assemblies in the MAT proximal region and other recombining regions. **B.** Annotation of transposable elements (TEs) and gene density along the MAT-proximal region in M1400 and the two Asian genomes ESM15 and XIM9508. Orthologous genes shared between M1400 and M6697 genomes are shown in black and unique to M1400 in grey as in Figure 5. Genes of ESM15 and XIM9508 are shown in darkblue? Genes with DU3435 domains (putative starship captains) are shown in red. Transposons larger than 150 pb are shown in orange. Synteny for segments with identity > 90 % is shown in red and inversion in blue.

Figure 7: Genetic structure using single nucleotide polymorphisms (SNPs) in the MAT-proximal region lacking recombination (A) and other genomic regions (B) in other resequenced *Cryphonectria parasitica* populations from the native and the introduced range. Neighbor-net network from a SplitsTree analysis based on: **A** SNPs (n=4,120) located within the MAT-proximal region along the contig carrying the mating-type locus (scaffold_2 of the EP155 genome); **B** SNPs (n= 103,058) located on other contigs of the EP155 genome. Color of empty circles around strain ID indicate the genetic clusters the strains belong to (CL1, CL2, CL3, CL4; n= 33 strains**)**. Dotted circles indicate strains of the native range and plain circles indicate strains of the introduced range. Red and blue dots near strain ID indicate the mating type (*MAT-1* or *MAT-2*, respectively).

	Population	CL1 genetic cluster		Swiss 1990 population		Swiss 2019 population	
	Haplotype	MAT-Prox1	MAT-Prox2	MAT-Prox1	MAT-Prox2	MAT-Prox1	MAT-Prox2
Number of strains in each PCA cluster	total	36	52	20	51	8	54
	MAT-1	30	11	18	17	5	15
	MAT-2	6	41	2	34	3	39
% of strains in each PCA cluster	*MAT-1*	83	21	90	33	63	28
	MAT-2	17	79	10	67	38	72

1474

Mating-type
locus

MAT-proximal

2 high LD blocks
② ①

SNP location
along scaffold_2 of EP155 genome (Mb)

0   1   2   3   4   5

SNPs Position along the heatmap

r² within the
CL1 cluster
(European invasive population)

R² Color Key

0   0.2   0.4   0.6   0.8   1

CL1 genetic cluster (European invasive population)

A

MAT-Prox2

MAT-Prox1

n=36

PC2 (1.7%)

PC1 (95.7%)

B

PC2 (20.3%)

PC1 (23.9%)

C

PC2 (16.6%)

PC1 (26.5%)

ther contigs (EP155 genome)

Mating-type allele:

- MAT−1
- MAT−2

D

MAT-Prox2

MAT-Prox1

MAT-proximal region

E

Other contigs (EP155 genome)

tig00000058   tig00000060

M6697

(MAT-Prox1)

tig00000001

M1400

(MAT-Prox2)

Mating-type
locus   MAT-proximal

2Mbp

Putative centromere

Mating-type locus

MAT-proximal region

Inversion

A GC content (%)

B TE load (%)

C F_{ST} MAT-Prox1/2

D Nucleotide diversity MAT-Prox1

E Nucleotide diversity MAT-Prox2

F Tajima's D all

G Tajima's D MAT-Prox1

H Tajima's D MAT-Prox2

Swiss 1990 population (European invasive population)

M1400 mating-type contig (bp)

A

B

XIM (China)

M1400 (MATProx2)

ESM15 (Japan)

Starship absent in M6697

(1)

MAT-proximal region

Mating-type allele :

 MAT−1

 MAT−2

Genetic cluster :

CL1 (European invasive strains from North America)

CL2 (European invasive strains from Asia and Asian native strains)

CL3 (European invasive strains from Asia and Asian native strains)

CL4 (Asian native strains)

(2)

0.01

other contigs

Genetic cluster :

CL1 (European invasive strains from North America)

CL2 (European invasive strains from Asia and Asian native strains)

CL3 (European invasive strains from Asia and Asian native strains)

CL4 (Asian native strains)