Life-history adaptation under climate warming magnifies the agricultural footprint of a cosmopolitan insect pest

Estelle Burc[1,3†], Camille Girard-Tercieux[1,2,5†], Moa Metz[1,4], Elise Cazaux[1,5], Julian Baur[1],

Mareike Koppik[1,6], Alexandre Rêgo[1], Alex Hart[1] & David Berger[1]*

1. Department of Ecology and Genetics, Program of Animal Ecology. Uppsala University, Norbyvägen 18D, 75236, Uppsala, Sweden.
2. AMAP, Université de Montpellier, CIRAD, CNRS, INRAE, IRD, Montpellier, France
3. Agronomy Institute Rennes-Angers (IARA), Graduate school of agronomy, 35000 Rennes, France
4. Department of Biology, Faculty of Natural Sciences, Norwegian University of Science and Technology, 7491 Trondheim, Norway
5. Ecole nationale supérieure agronomique de Toulouse (ENSAT), Graduate school of agronomy, Toulouse, France.
6. Department of Zoology, Animal Ecology, Martin-Luther University Halle-Wittenberg, Halle (Saale), Germany.

†Both authors contributed equally and are presented in alphabetical order.

*Corresponding author.

david.berger@ebc.uu.se

Keywords: life history, trade-off, temperature, climate change, global warming, agriculture, ectotherm, insect, pest, crops, gene expression.

Abstract

Climate change is affecting the distributions and growth rates of cold-blooded pest species, with dire consequences for agriculture and global food security. To address this challenge, it is essential to forecast how pests will respond to changing climates. However, current projection models typically overlook the potential for climate change adaptation. By unifying life-history theory with thermodynamics based on first principles, we here show that pest species' climate adaptation can cause substantial increases in their agricultural impact via changes in temperature-dependent resource acquisition and allocation strategies. We test these predictions by studying thermal adaptation in life-history traits and gene expression in the cosmopolitan insect pest, *Callosobruchus maculatus*. Five years of adaptation to simulated warming caused an almost two-fold increase in the predicted agricultural footprint of *C. maculatus,* while adaptation to cold temperature had little effect. Consistent with the theoretical predictions, the magnified impact under warming was driven by synergistic increases in larval resource acquisition and adult reproductive potential, resulting in a predicted double blow on agricultural yields. This suggests that adaptation in insect life-history will significantly worsen agricultural loss under future climate change and emphasizes the need for integrating a mechanistic understanding of life-history evolution into ecological forecasts of pests.

Climate change is predicted to greatly impact growth rates and distribution ranges of cold-
blooded pest species[1–7] with potentially severe outcomes for agricultural economics and world
food supply[8–12]. To mitigate these events, forecasting of pests' responses to future climates is
a fundamental task for biologists[13,14]. Current forecasts use deterministic models that rely on
information about species niches and climate data. However, these projections typically do not
incorporate the potential for genetic evolution of climate niches and key life-history traits that
mediate the effect that pests have on their hosts[13,14]. Indeed, the evolution of ecological niches
has been rapid in many cosmopolitan pests and not accounting for this adaptive potential may
therefore provide inaccurate forecasts of their future agricultural impact[1,7,13–17].

Despite its importance, the evolution of pest species' climate niches is not well-
understood[1,5,7,16]. Cold-blooded organisms are constrained by temperature-dependent
biophysical properties of cells[18–22]. On the one hand, metabolism is fundamentally constrained
by cold temperature, predicting that climate warming may increase growth rates[23,24] and
thereby worsen pest impact in many parts of the world[2]. On the other hand, hot temperatures
can jeopardize cellular homeostasis[25–27], leading to fertility decline[28,29] and increased
mortality[22,30]. These thermodynamic constraints are expected to limit the distribution ranges of
ectotherms. However, typical signatures of pest species are their broad climate niches coupled
with fast growth rates[1,16]. One mechanism that can allow organisms to occupy broad climate
niches is up-regulation of molecular chaperones that aid cellular homeostasis under harsh
conditions[20,31]. Such responses are energetically costly and therefore need to be compensated
by either increased resource intake[5,19], or by diverting resources away from reproduction in
favour of cellular maintenance[20,31]. Such alternative strategies in pest species' life-history are
predicted to impact both their invasion potential and the footprint left on their hosts[1,14,17,32–34],
and thus, can have important consequences for pest management[1,3,35].

However, exactly how life-history evolution unfolds under climate change remains
unclear[1,16,33]. Moreover, for adaptation to occur, genetic variation is a prerequisite.
Geographically isolated populations can differ greatly in the abundance of both segregating
and fixed genetic variation due to differences in their evolutionary past, which may affect future
adaptive potential[36–40]. Such historical contingencies may be of particular importance in insect
pests, as geographically isolated populations often originate from human-mediated long
distance dispersal events of only a few individual genetic founders. Evolution under climate
change may therefore proceed along different trajectories and rates across a species
distribution range, even when climates change in parallel.

Here we explored the agricultural impact of life-history evolution under climate change by
employing experimental evolution in the seed beetle, *Callosobruchus maculatus*, an
economically important cosmopolitan pest on legume seeds, and a model organism for life-
history research[41]. Life-history evolution over 5 years in response to warming caused a
substantial increase in the agricultural footprint of *C. maculatus* via trait-specific responses, an
effect not seen in populations adapting to cold conditions. This increase was also contingent
on the genetic make-up of the founding populations. These findings suggest that equipping
species distribution models with genetically explicit details accounting for life-history evolution
can lead to a substantial improvement in predictive accuracy of pest impact under climate
change, even over relatively short time scales.

Results

The agricultural impact of thermal adaptation via compensatory resource acquisition
To formalize predictions for how thermal adaptation of insect pests will impact agriculture, we
combined a classic life-history framework of energy acquisition and allocation trade-offs[42–45]
with the thermodynamic laws that govern enzyme kinetics[20,46]. Warm temperatures increase
growth rates of ectotherms[19,23] because any biological rate is at some level governed by
enzymatic reaction rates, k, that scale predictably with temperature according to[47,48]:

$$k \propto Ae^{-Ea/RT} \qquad \text{(Eq. 1)}$$

where Ea is the activation energy required for a reaction to occur, R is the universal gas
constant, T is temperature in Kelvin, and A is a species and rate specific constant that may
itself evolve to accommodate effects of temperature[49,50] (**Fig. 1A**). Complementing equation 1
with the power law describing how productivity scales with body mass, M, results in the well-
known prediction of body mass and temperature-dependent population growth rate based on
the metabolic theory of ecology[23,24]: $r \propto M^b k$, where b is the allometric exponent, often taken
to be 0.75 based on the physics of distribution networks in plants and animals[23,51]. However,
excessively hot temperatures lead to increased molecular failure rates[30] due to a range of
detrimental processes that also scale exponentially with temperature, including protein
unfolding[25,26], metabolic expenditure[52,53], production of reactive oxygen species[20], and oxygen
depletion[54] (**Fig. 1A**). Thus, mortality rate can be described by equation 1 with its own
activation energy (Ea) and rate-specific constant (A) (e.g.[30,53,55]). As follows, population growth
rate is the result of temperature-dependent productivity and mortality rate[1,2] (**Fig. 1B**).

While there is variation in biological rates across organisms[49,56,57], inherent enzyme properties

governing the temperature-dependence of reaction rate kinetics (Ea in eq. 1) evolve slowly[50,58].

Instead, thermal adaptation often involves rapid evolution of compensatory stress responses

(A in eq. 1)[20,31,49,59]. Such responses involve upregulation of molecular chaperones, such as

heat-shock proteins[20,60,61], that maintain cellular homeostasis at stressful temperatures but are

costly to produce[31,60,62]. Here we therefore placed the predictions of equation 1 into a life-

history framework incorporating acquisition and allocation trade-offs[42,43,62]. First, assume that

body reserves, M, are accumulated by host feeding at a cost, c, so that survival to maturity

equals: $(1-c)^M$. Further assume that body reserves can be allocated with fraction p to increase

enzymatic reaction rate, and with fraction q to decrease molecular failure rates, leaving $M(1-$

$p-q)$ for reproduction, B, according to:

$$B \propto [M(1-p-q)]^b \; ; \quad A_B \propto pM \; ; \quad A_m \propto 1/qM \qquad \text{(Eq. 2)}$$

where A_B and A_m are the rate-specific constant for productivity and mortality, respectively.

Population growth rate, r', is then:

$$r'(T) \propto (1-c)^M * B * A_B\, e^{-\frac{Ea_B}{RT}} * (1 - A_m\, e^{-\frac{Ea_m}{RT}}) \qquad \text{(Eq. 3)}$$

We can find the optimal strategy of resource acquisition (increasing M at cost c) and allocation

(of M to p and q) at different temperatures by numerically solving for the combination of M, p

and q that maximize r'. The solution to this multidimensional trade-off depends on the specific

form of equations 1 and 2 and parameters that govern temperature sensitivity (i.e. Ea, A and

their implementation); mechanistic details which are empirically not well-defined. Indeed,

equation 1 represents a phenomenological, rather than a mechanistic, relationship and studies

have highlighted that the coefficients Ea_B and Ea_m can themselves be temperature-dependent

(e.g. [30]), emphasizing the need for further experimental work quantifying these relationships.

Nevertheless, the qualitative predictions from this heuristic model are robust (see also [53]). We

provide a parameterized example in **Figure 1** grounded in knowledge of how temperature

affects protein catalysis and stability[25,27,58,63] (details in **Suppl. 1**). First, and as pointed out

previously[2], warm temperatures increase growth rates and agricultural impact by reducing

thermodynamic constraints on reaction rates. Second, when thermodynamic constraints

jeopardize survival, hot temperatures favour increased resource acquisition and more body

reserves (M) that can be allocated to maintenance[53] (**Fig. S1A, Fig. 1B**). When increased

resource acquisition comes at low cost, this life-history strategy is particularly favourable and

results in increased thermal niche breadth and population growth rates (**Fig. 1C**). Third, this
has an even greater agricultural impact, given by the product of population growth rate (*r'*) and
the per capita host consumption (assumed proportional to *M)* (**Fig. 1D**). In fact, the strategy to
increase resource acquisition at stressful temperatures causes populations experiencing
temperatures slightly hotter than those which maximize growth rate to have the greatest
agricultural impact (**Fig. S1**).
162
Experimental evolution of life-history traits and agricultural impact
To evaluate the prediction that trait and temperature-specific responses in pest species' life-
history can influence their agricultural impact under climate change, we compared twelve
experimental evolution lines of the seed beetle *C. maculatus*; six adapted to 35°C (*Hot*) and
six adapted to 23°C (*Cold*). All lines were reared on seeds from their preferred and
economically important host, black-eyed bean (*Vigna unguiculata)* (**Fig. 2**). The lines were
derived from three geographically isolated founding populations (*Ancestors*, kept at 29°C)
originating from Brazil, California (USA) and Yemen[64], allowing us to assess the importance of
geographic differences and evolutionary history for future responses to climate change.
172
Molecular evidence for a temperature-dependent trade-off between reproduction and
maintenance
To provide evidence for the central assumption of a trade-off between investment in
reproduction and cellular maintenance under thermal stress (Eq. 2), we analysed data on gene
expression from two separate experiments on *C. maculatus*. These datasets contained genes
that were significantly differentially expressed in female abdomens in response to heat shock,
and between virgin and reproducing females. We identified 1269 reproduction-related genes
and 765 heat stress genes. We found 137 genes present in both datasets, which is significantly
more than expected by chance (exp: 83.4 genes, $\chi^2 = 37.5$, df = 1, p < 0.001, **Fig. 3A**).
Moreover, only 3 of the 137 genes showed concordant responses to mating and heat-shock
(all three decreased in expression), whereas 134 genes showed antagonistic responses
(binomial test: P < 0.001, **Fig. 3B**), providing strong evidence for a trade-off between
reproduction and the heat stress response. Out of these genes, 21 were up-regulated in
response to reproduction and down-regulated in response to heat. These genes included
common ribosomal genes that are crucial for protein translation, signifying reproductive
investment. The 113 antagonistic genes that were down-regulated in response to mating and
up-regulated in response to heat stress included common insect heat-shock proteins (**Suppl.**
Table 2a). GO enrichment analysis showed that the antagonistic genes that overlapped both
gene sets were enriched for biological process involved in protein folding and degradation of

unfolded proteins, which are essential components of cellular maintenance under thermal stress (**Fig. 3C, Suppl. Table 2b**).

To confirm that the revealed trade-off also was at play in the thermal conditions used during experimental evolution, we analysed gene expression data from all Cold and Hot lines, as well as their Ancestors, reared in a common garden design including 23, 29 and 35°C assay temperature. Each evolution line was represented by a single mRNA library per temperature, based on a pool of 10 abdomens from mated females. The ancestors were represented by two such replicates. To evaluate investment into reproduction versus maintenance, we calculated a score for each of these mRNA libraries along the axis describing the trade-off based on the expression of the 134 antagonistic genes (**Fig. 3B**). We then analysed differences between assay temperatures and evolution regimes using nested ANOVAS (**Suppl. Table 2c**). Beetles reared at 35°C showing a strong change in allocation towards cellular maintenance and away from reproduction compared to beetles reared at the other two temperatures (assay temperature: F = 263.0, P < 0.001, **Fig. 3D**). There were no significant differences between evolution regimes overall (F = 0.12, P = 0.89), although the differences between the evolution regimes and their ancestors showed a dependence on geographic origin (interaction: F = 11.0, P = 0.002). Qualitatively similar results were obtained when analysing all 765 heat stress genes and 1269 reproduction genes (**Suppl. Tables 2d, e**). While statistical power was low due to the small number of mRNA libraries, this implies that there has been little evolutionary change in resource allocation overall (i.e. q in Eq. 2). This result further predicts that if hot-adapted lines have evolved a means to deal with the increased cellular stress of hot temperature (35°C), they must have done so mainly via increased resource acquisition (M in Eq. 2), which should be apparent as an evolved increase in adult body mass.

Thermal adaptation in life-history traits

We assessed the predicted agricultural impact of life-history evolution by assaying three classic traits; juvenile development time, female adult body mass, and lifetime reproductive success (LRS: number of offspring produced), and four rate-dependent traits assayed over the first 16h of female reproduction; early fecundity, weight loss, water loss and metabolic rate (**Fig. 4**). This also allowed us to evaluate if Hot lines had adapted to their temperature regime by increased energy acquisition (M; captured by adult body mass).

To characterize trait-values at the start of experimental evolution, we assayed the three ancestors reared at 23, 29 and 35°C. This allowed us to quantify both the direction and magnitude of adaptation during experimental evolution, as well as its subsequent agricultural impact (*see below*). Temperature affected the expression of all seven life-history traits. For

development time, body mass, early fecundity and LRS, there were also main effects of the geographic origin of the ancestors. Body mass, early fecundity and LRS also showed a significant interaction between geographic origin and assay temperature, indicating differences in thermal plasticity between ancestors (**Fig. 4A-G, Suppl. 3**).

To assess temperature-dependent life-history evolution we assayed the 12 evolved lines when reared in a common garden experiment including 23°C and 35°C. All traits except metabolic rate and weight loss showed significant differentiation between the Hot and Cold regime. In particular, the Hot regime had indeed evolved larger adult body mass and higher reproductive output (**Fig. 4A-G**), in accordance with model predictions (**Fig. 1C**). All traits except metabolic rate and LRS showed a significant interaction between regime and assay temperature, demonstrating thermal adaptation *sensu stricto* (**Fig. 4B-G**). Several alternative estimates of laboratory fitness also showed clear signs of temperature-specific adaptation, with Hot lines outperforming Cold lines at 35°C, while there were only small differences at 23°C (**Suppl. 4**). In addition, for metabolic there was an interaction between evolution regime, assay temperature and geographic origin, indicating that thermal adaptation was contingent on founding genetic variation (**Fig. 4B, E, Suppl. 3**). Analysing all traits together using a non-parametric MANOVA[65] confirmed main effects of assay temperature (F = 205.5, P < 0.001), evolution regime (F = 57.1, P < 0.001) and the interaction between evolution regime and assay temperature (F = 6.8, P < 0.001).

Thermal adaptation in life-history shapes the agricultural footprint

To predict how the observed life-history evolution may affect agricultural yields, we estimated the relationship between life-history traits and the amount of host consumption by *C. maculatus.* We reared Cold, Hot and Ancestral lines at 23, 29 and 35°C. At each assay temperature, we let females lay eggs in a petri-dish containing ad libitum black-eyed beans. The number of eggs laid, the number of individuals surviving to pupation, the mass of adult beetles, and the seed mass consumed, was calculated from each dish.

The Hot regime consumed significantly more of the host than the Cold regime ($F_{1, 5.97}$ = 22.8, P = 0.003), demonstrating that the agricultural footprint indeed had evolved. The difference between the Hot and Cold regime tended to be greater at 35°C, although the two-way interaction between regime and assay temperature was marginally non-significant ($F_{2, 11.9}$ = 3.25, P = 0.075, **Fig. 5A**). To assess the importance of life-history traits in affecting food consumption, we performed analyses while sequentially including information on the number of eggs laid, juvenile survival, and body mass so that the final model predicted the amount of host consumed as a function of the cumulative beetle mass produced. Adding life-history traits

to the models reduced the differences between evolution regimes and in the final analysis including all traits we no longer found a significant effect of regime ($F_{1, 5.73} = 0.60$, $P = 0.47$), demonstrating that life-history adaptation alone seems responsible for the evolved differences in host consumption. Indeed, a model containing only the main effects of eggs laid, adult body mass and juvenile survival to pupation explained 93% ($P < 0.001$) of all the observed variance in bean consumption across the entire experiment (**Fig. 5A**). The amount of host mass lost was 3-4 times greater than the mass of beetle produced, which is similar to what has been reported in a previous study[66], underlining the severe impact that *C. maculatus* can have on host crops (**Suppl. 5**).

We predicted the agricultural impact of *C. maculatus* at cold and hot temperature based on the observed life-history evolution by calculating the agricultural footprint as: $\varphi_1 = C_T \omega M / d_1$, where C_T is the temperature-specific constant converting beetle mass produced to host mass consumed (**Fig. 5A**), ω is lifetime offspring production of a single female (**Fig. 4A**), M is body mass at adult emergence (**Fig. 4B**) and d_1 is mean development time in days (**Fig. 4C**). Hence, this measure of the footprint captures the magnitude of agricultural loss per time unit as a function of an organism's life-history strategy[67]. To assess the sensitivity of our predictions to both estimation error and choice of life-history variables we calculated a second independent estimate of the footprint, based on measurements of a different set of individuals: $\varphi_2 = \psi C_E / d_2$, where ψ is the number of eggs laid by a single female over her first 16h of reproduction (**Fig. 4D**), C_E is the amount of host consumed per laid egg, and d_2 is the mean development time in days of the fastest developing individual in each petri-dish (both variables estimated from the host consumption experiment presented in **Fig. 5A**). These two metrics were highly correlated ($r = 0.88$, $t_{31} = 10.4$, $P < 0.001$, **Fig. 5B**) suggesting high predictive accuracy.

To provide a generalisable metric of the agricultural impact of life-history evolution under climate change, we scaled the predicted footprint for all lines relative to their respective ancestor reared at 29°C (**Fig. 5C**). This metric allowed us to assess both the direct effect of temperature on the footprint in the ancestors, and how evolution, relative to this direct effect of temperature, impacted the footprint. Cold (23°C) and hot (35°C) assay temperature reduced the footprint by 35-45% and 0-25% respectively, relative to benign 29°C in the ancestors (**Fig. 5C**). However, evolution had a fundamental impact on how temperature affected the footprint. While cold-adapted lines showed very little difference to ancestral lines at 23°C (Δfootprint = 7.6%; CI: 2.0 – 15%), Hot-adapted lines left almost twice as large a footprint as the ancestral lines at 35°C (Δfootprint = 80%; CI: 70 – 97%). Moreover, these effects were contingent on the geographic origin of the founders. At 23°C, cold-adapted lines from California left a slightly larger footprint compared to the ancestor (Δfootprint = 29%; CI: 13 – 42%) while the footprint

had not evolved on the Brazil (Δfootprint = -3.1%; CI: -11 – 6.3%) and Yemen (Δfootprint = 1.0%; CI: -11 – 11%) background. At 35°C, the footprint evolved substantially on all backgrounds, but the increase relative to the ancestor tended to be greater for Brazil (Δfootprint = 84%; CI: 64 – 110%) and California (Δfootprint = 93%; CI: 71 – 120%) compared to Yemen (Δfootprint = 62%; CI: 44 – 87%) (**Fig. 5C**).

Discussion

Climate change is predicted to fundamentally impact growth rates and distribution ranges of ectotherms[5,30,37,68–71], many of which are pests on economically important crops[1,2,7,16]. Current predictions of increased agricultural impacts of pests typically rely on deterministic statistical projections of how temperature affects ectotherm metabolism and population growth, but do not incorporate evolution[13,14,67,72]. However, rapid evolutionary responses are characteristic of many pest species[1,5,7,16]. Here we therefore asked to what extent the incorporation of evolution can alter current projections of climate-induced changes in pest impact. We show three features that emphasize that future projections need to account for thermal adaptation in pest species' life-history. First, adaptation affected the agricultural footprint left by *C. maculatus,* but the effect was dependent on temperature; with little or no effect of cold adaptation while hot-adapted populations caused an almost doubling of the footprint relative to ancestors (**Fig. 5**). Second, these effects depended on the ancestral founder of the adapting populations, suggesting that geographic differences in segregating genetic variation in life-history traits may influence future agricultural loss under climate warming. Third, attempts to incorporate niche evolution in species' distribution models may need to do so by relatively simple means in the absence of empirical data that allow parametrisation and evaluation of more complex mechanistic models[13]. However, our study echoes concerns that such predictions may be off the mark because the mechanisms and type of traits that mediate niche evolution have central influence on pest impact[14,17,34,35].

Indeed, the observed temperature-dependent effect of evolution on the agricultural footprint can be understood by considering trait-specific responses. Even though cold-adapted populations did show evolved responses in traits such as development time, weight loss and water loss, these traits are predicted to have small to no effect on the agricultural footprint compared to lifetime reproductive success, early fecundity and adult body mass, that evolved more readily in hot-adapted populations (**Fig. 4**). The responses of hot-adapted populations are in line with two predictions from biophysical theory. First, warmer temperatures are predicted to ease thermodynamic constraints on metabolic rate[52], which can increase selection

for larger size via effects on fecundity[73,74]. Second, evolution at stressful hot temperature may favour compensatory feeding and sequestering of resources devoted to maintaining cellular homeostasis[5,19,20,31,42,53,62] (**Fig. 1, Suppl. 1**). Indeed, increased larval feeding and larger adult body mass in *C. maculatus* is associated with extended adult lifespan and greater fecundity under aphagous conditions corresponding to laboratory and grain storage environments (e.g.[75,76]). These effects are likely magnified at hot temperatures that increase metabolism and water loss ([23,77] and **Fig. 4**), and the need for upregulation of molecular chaperones to secure cellular homeostasis ([20,60,61] and **Fig. 3**). In accordance with this hypothesis, previous studies on *C. maculatus* also found that evolution at hot temperature increased body mass[78,79], an adaptive genetic response that paralleled the observed thermal plasticity in body mass[78]. To further support this hypothesis, we used the estimates of female life-history to calculate selection differentials on body size at hot and cold assay temperature. For all three estimates, selection was stronger at the hot temperature (**Suppl. 6**). Thus, the evolved increase in the agricultural footprint at hot temperature observed here is likely mediated by two synergistic effects of life-history adaptation, *i)* increased larval feeding enabling larger size at maturity, and *ii)* the positive effects of a large adult size on survival and reproduction at hot and aphagous conditions.

These observations agree broadly with the model prediction that stressful temperature can favour increased host consumption and energy allocation to maintenance to mitigate thermal constraints on growth and reproduction (**Fig. 1, Suppl. 1**). Note, however, that this prediction does not necessarily equate to the evolution of a large size at maturity, as resources might be spent on higher metabolic expenditure and maintenance at warm temperature prior to the organism reaching adult maturity, and our simple model makes no explicit predictions in this regard (see also[80,81]). Moreover, increased energy intake necessitates access to host plants. If food availability becomes limiting under climate warming, this mechanism could instead severely limit thermal tolerance and population growth rates of heat exposed ectotherms[53,55]. Thus, while host abundance is already acknowledged as a key factor regulating pest outbreaks[1,7,67,82], the interdependency between heat stress and the benefits of energy acquisition may magnify the effects of host accessibility on pest outbreaks under future climate warming (**Fig. 1C, D**).

Early findings suggested that most ectotherms follow the "temperature-size rule"[83,84], stating that body size shows plastic reductions with increasing temperature. However, later results have shown that these patterns are far from general. In particular, terrestrial ectotherms, the group to which many insect pests belong, show no consistent plastic response to temperature, and genetic adaptation along latitude shows a trend towards larger genetic body sizes at

warmer latitudes[85], which is consistent with the evolved response observed here. Moreover,

thermal responses in body size tend to vary even within and between closely related species[71],

as for *C. maculatus*[78,86] and other Chrysomelid beetles[87]. Understanding whether taxonomic

variation represent constraints rooted in thermodynamics or adaptive plasticity may offer

insights into the future routes of evolution under climate warming but has proven a major

challenge for the field of eco-physiological adaptation[5,19,71,80,84]. Moreover, as discussed above,

the effect of heat on energy expenditure suggests that the temperature scaling of adult body

mass may not follow a one-to-one relationship with that of host consumption. Controlled

experimental approaches as used here may provide further clues to what extent patterns of

thermal plasticity are indicative of future genetic responses to climate and how the evolution

of classic life-history traits affects host consumption. Such general information could be used

to increase accuracy of future projections for species where thermal plasticity can be accessed,

but where evolution is hard to study directly.

In contrast to the measured life-history traits, feeding efficiency was not strongly affected by

assay temperature and did not evolve in our experiment, suggesting that differences in the

agricultural footprint of insect pests may be accurately predicted from life-history variables

alone. Yet, our study also highlights the important role of founding genetic variation in dictating

life-history adaptation. This suggests that evolution under climate change may proceed along

different trajectories across a species distribution range, even when regional climates change

in parallel. When such divergent evolutionary responses involve key life-history traits, as

observed here, this could lead to regional differences in pest impact[1,2,35].

The impact of insect pests on agriculture is paramount[11,67], with severe repercussions in human

societies; 2.6 billion people (26% of the world's population) are currently directly dependent on

agriculture for a secure living[88]. In African countries alone, the financial loss from pests was

estimated to at least 1 billion USD for 2019, with agriculture accounting for 99% of this loss in

central regions[9,10]. Clearly, the redistribution of species has already caused massive

ramifications for the Earth's ecosystems[89], and the fear is now that future climate warming will

allow insect pests to colonize new habitats[1,2,4,7,11,12]. To mitigate these effects, management

strategies need to be equipped with accurate predictive models[13,14,72,90–92], as well as an

understanding of the limits to their accuracy[38,93–95]. It is our hope that our study can contribute

towards this development by *i)* providing general insights into the repeatability and mode of

thermal adaptation in life-history of insect pests, which can inform on the general accuracy of

ecological forecasting of species responses to climate change, and *ii)* by emphasizing the need

for incorporating a mechanistic and genetic basis of thermal niche evolution into eco-

evolutionary forecasts, on which improved management decision can be based[6,13,17,35,72,90,96,97].

Methods

Experimental evolution

C. maculatus is a pest on legume seeds and is common in tropical and sub-tropical regions[98]. Females lay eggs on the surface of seeds. After the egg hatches, the larva burrows into the seed to feed until it pupates. Adults emerge from seeds after ca. three weeks at standard temperatures around 29°C and are reproductively mature almost instantly and do not require food or water to reproduce[64,78,99]. These characteristics have likely contributed to *C. maculatus* being a cosmopolitan pest that invades seed storages[12,41]. Indeed, *C. maculatus* has become a model organism for research in both evolutionary ecology and pest management because laboratory and grain storage environments are very similar.

The experimental evolution protocol exposed six lines to 35°C (*Hot* regime) and six lines to 23°C (*Cold* regime), for 85 and 130 generations, respectively (difference due to effects of temperature on generation time). The lines were derived from three ancestors that were collected from different geographic regions (Brazil, Yemen and California (USA)), with two replicate lines per thermal regime created per region. The three founders were kept at ancestral 29°C throughout experimental evolution (**Fig. 2A**). A humidity of 50-55% was used throughout this study. Ancestors were reared, both before and during experimental evolution, by placing 300-400 newly emerged adults on 150 ml black-eyed beans, whereas evolution lines were reared by placing 600 adults on 250 ml black eyed beans[64], resulting in similar densities of adult beetles per bean. To reduce direct selection on development time, all lines were transferred to a new jar with fresh beans around the peak of emergence of adult beetles at each generation. This occurred approximately three days after the first beetle emerged at for ancestors and the Hot regime, and five days for the Cold regime.

Gene expression

To provide evidence for a temperature-dependent trade-off between reproduction and maintenance, we leveraged transcriptomic data from three separate experiments. The first two experiments estimated gene expression responses in female abdomens to heat shock and mating, respectively, in a separate lab population collected from Lomé, Togo[100]. The third dataset estimated expression in female abdomens in response to rearing temperature in our experimental evolution lines.

To identify heat stress responsive genes, we compared reproductively active control and heat
shocked females. Individuals were collected as virgins and kept individually. On day 2, females
were heat shocked in individual 60 mm petri dishes at 55° C for 20 minutes, which is stressful
and reduces female fertility[101,102]. After a 2.5 h at 29° C and 55% rh, all females were mated to
males. Females were then kept in individual 60 mm dishes until being flash frozen in liquid
nitrogen 2-3 h, 6-7 h or 24-25 h after mating. The experiment was carried out on three
consecutive days. Each replicate sample consisted of 5 pooled female abdomens from the
same treatment, time-point and day, resulting in a total of seven replicate samples per time-
point for the control treatment and five replicate samples per time-point for the heat shock
treatment. Preliminary analyses showed that there were only three differentially expressed
genes at the last time-point, which therefore was removed in the final analyses (although
results did not qualitatively change if including them), resulting in 24 RNA libraries.

To identify reproductive genes, we compared virgin and mated females 24 h after a single
mating. We mated females to males from 8 different genetic lines. The lines had undergone
53 generations of experimental evolution under one of three mating regimes manipulating the
relative strength of natural and sexual selection[102]. Accordingly, we had 8 samples of mated
females, with 2 or 3 samples per male mating regime. Additionally, we had 3 samples of virgin
control females. Flash freezing of all females occurred 24 h after mating, with females kept
singly and allowed to lay eggs on fresh beans during the 24h period. The final 11 RNA libraries
each consisted of 12 female abdomens.

To quantify expression of reproductive and heat stress responsive genes (identified in the first
two datasets) in our experimental evolution lines and their ancestors, we reared all lines at 23,
29 and 35°C. All lines were first propagated for one generation at 29°C to remove potential
difference between evolution regimes stemming from temperature-induced (non-genetic)
parental effects. Once adults emerged at respective rearing temperature, 5 males and 5
females were placed together in a 90mm petri dish with access to fresh beans. Beetles were
allowed to reproduce for 23h (for 29 and 35°C) or 46h (for 23°C) after which females were
flash frozen. The longer time allowed for reproduction at cold assay temperature was given to
make a more direct comparison across females from different temperatures; reproductive
output (Fig. 4B), metabolism (Fig. 4E) and weight loss (Fig. 4F) is about twice as fast at 35°C
compared to 23°C. Two petri dishes of females were pooled into one sample (i.e. 10 female
abdomens pooled in total) for extractions and further analysis, resulting in one sample per line
and temperature for the Hot and Cold regime. Two samples were taken per temperature for
each of the three ancestors. This resulted in a total of 54 RNA libraries.

RNA from all samples was extracted using the Qiagen RNeasy Mini kit with beta-mercapto-
Ethanol added to the lysis buffer and an on-column DNase treatment with the Qiagen RNase-
free DNase kit. Tissue lysis was done in a bead mill with two stainless steel beads at 28 Hz for
90 s and RNA was eluted in two times 30-50 µl water. Some samples went through an
additional clean-up step with the Qiagen RNeasy Mini Kit after the RNA extraction. Next
generation sequencing of the samples was done at the SNP&SEQ Technology Platform in
Uppsala. Libraries were prepared using the TruSeq stranded mRNA library preparation kit with
polyA selection. Libraries were sequenced in one flowcell as paired-end 150 bp reads on a
NovaSeq 600 system.

Raw reads were quality checked with FastQC 0.11.9[103] and MultiQC 1.11[104] and afterwards
cleaned up using Trimmomatic 0.39[105]. Adapters were trimmed from the raw reads and ends
below an average accuracy of 99% over a five base pair sliding window were clipped. Finally,
reads shorter than 20 bp were discarded. Resulting read pairs were mapped to the
Callosobruchus maculatus genome[106] using HISAT2 2.2.1[107] with default settings for stranded
libraries and sensitivity set to "--very-sensitive". Number of reads per gene was then
determined using HTSeq 2.0.2[108] with default settings for stranded libraries. This resulted in
10-27M (heat stress), 13-30M (reproduction) and 9-35M (rearing temperature during
experimental evolution) uniquely mapped reads per library going into the final downstream
analyses.

Read count data was pre-processed and analysed in R 4.3.1[109] using edgeR[110,111]. For the
heat shock response, only genes with at least 3 counts per million in at least 5 samples were
further analysed, resulting in 10246 genes for final analysis. For the mating response, only
genes with at least 1 count per million in at least 2 samples were further analysed, resulting in
11640 genes for final analysis. The last approach was also used for the genes expressed in
the evolution regimes, resulting in 11292 genes for analysis. Counts were normalized using
the 'Trimmed Mean of M-values' method via edgeR's NormLibSizes() function. Count data for
the heat shock and mating response was further analysed with edgeR[110,111] and
limma[112] using linear models on the normalized log2 transformed counts per million to
determine differentially expressed genes. All analyses used a false discovery cut-off of 5 %.

For reproductive genes, effect sizes (i.e. Log-fold changes) were averaged across the three
male selection regimes. However, we only considered genes that were significantly
differentially expressed in all three separate contrasts between virgin females and females
mated to males from the three mating regimes. For heat stress genes, effect sizes and
significance was calculated by averaging across the two time-points. Significant genes were

further subject to gene ontology analysis, using the R package GOstats (2.66.0)[113] and the

HyperGTest function. Results were visualised using clusterProfiler (4.10)[114].

To provide a powerful, yet, unbiased estimate of temperature-dependent allocation between

reproduction (mating response) and maintenance (heat stress response) in the experimental

evolution lines, we focused on the 134 genes that displayed significant antagonistic differential

expression patterns in response to heat shock and mating (Fig. 3B). We first multiplied the

estimated log fold changes of the antagonistic genes for both mating and heat shock responses

by the vector of normalized read counts of the same 134 antagonistic genes found for each of

the 54 RNA libraries from the experimental evolution lines. The resulting values were then

summed to compute a score for the heat stress and reproduction response separately. As

expected, the scores for heat stress and reproduction, based on only the antagonistic genes,

were tightly negatively correlated (r = -0.999) across all combinations of evolved lines and

assay temperatures and fell along a singular linear trade-off axis (Supplementary Fig. 2). The

response along the allocation trade-off for each sample was taken by projecting samples along

this axis (quantified as the first eigenvector of the data). We repeated the same analyses using

all differentially expressed genes in response to heat shock (765) and mating (1269) with

qualitatively similar results (Suppl. Tables 2d, e).

Life-history traits

We quantified thermal adaptation in female life-history by measuring three core traits: lifetime

reproductive success (LRS), juvenile development time and adult body mass, and four rate-

dependent traits: early fecundity, weight loss, water loss, and mass-specific metabolic rate (ml

CO_2/mass/min) over the first 16h of female reproduction. All life-history traits were collected at

generations 40 for cold-adapted lines and 60 for hot-adapted lines, in a large common garden

experiment including the two assay temperatures corresponding to the experimental evolution

treatments (23°C and 35°C). Ancestral lines were scored in the same experimental conditions

with the addition of the ancestral 29°C assay temperature, but on a later occasion following

ca. 125 generations of experimental evolution. Note that the ancestors had been kept at the

ancestral conditions, to which they had already adapted for more than 300 generations prior

to the start of experimental evolution. It can therefore be assumed that the measured trait

values correspond well with the trait values at the start of experimental evolution. To control

for potential differences in the separate experiments on ancestors and evolved lines stemming

from unknown sources, we reared an independent laboratory adapted reference population in

both experiments. This indicated that differences in rearing had affected the life-history traits

scored over the first 16h of reproduction. We therefore standardized the traits scored during

respirometry of the three founding ancestors by this estimated amount (adult mass: increased

by 6.4%, metabolic rate: reduced by 12%; early fecundity: reduced by 18%; water loss:

reduced by 15%, and weight loss: reduced by 25%) in order not to erroneously assign these

differences to evolutionary divergence between ancestors and evolved lines. Note that this

was done averaged across the three assay temperatures and geographic origins. Therefore,

our approach to provide more accurate measures of evolutionary divergence between evolved

lines and ancestors did not affect the estimated temperature-dependence of adaptation or the

importance of geographic differences.

Before assays of life-history traits, non-genetic parental effects were removed by moving F0

grandparents of the assayed individuals into a common temperature of 29°C to lay eggs. The

emerging beetles in the next (parental) F1 generation were allowed to mate and lay eggs on

beans provided *ad libitum*. Following 48 hours of egg laying, the beans were split and assigned

to one of the two (for ancestors, three) assay temperatures. The emerging adult F2 offspring

were phenotyped for their life-history (**Fig. 4**). Newly emerged (0-48 hours old) virgin females

were mated to males by placing three males and females together in a petri dish over night at

the assay temperature. In the following morning, the three females were weighed for their body

mass and then placed together inside a glass vial filled with black eyed beans to be measured

for their metabolic rate, water loss and early fecundity at their designated assay temperature.

The glass vials were placed in a Sable Systems (Las Vegas, NV, USA) high-throughput

respirometry system. Briefly, the respirometry was set up in stop-flow mode, and CO^2

production and water-loss was measured for up to 23 vials on a given experimental day. The

first vial was left empty and served as a baseline to control for any drift of the gas analysers

during each session. Vials were measured over 17 cycles, each of a length of 57.5 minutes.

Mean metabolic rate and water loss for each vial was calculated across cycles 2-17, with the

readings from the first cycle discarded (as it contains human-produced water and CO_2). After

respirometry, females were weighed again to record their weight loss and beans with eggs

were isolated and counted to record early fecundity. In total we followed 386 triads of females

for the evolved lines and another 115 triads from their ancestors.

From the same rearing we measured egg-to-adult development time for two technical

replicates per line and assay temperature, each consisting of 40-120 individuals. We

calculated a mean development time per technical replicate and used this in analysis. We also

collected virgin males and females and placed three males and three females together in a

petri dish with ad libitum beans to record lifetime reproductive output (LRS) at each assay

temperature. In total we recorded LRS for 258 couple triplets for evolved lines, and another

115 couple triplets for the ancestors. These data were complemented with additional data from

both evolved and ancestral lines reared in a common garden design in two consecutive years

(corresponding to generation 120/135 for ancestors, 115/130 for hot-adapted lines, and 80/90

for cold-adapted lines). In these rearings, a single male and female were put together in a petri

dish with ad libitum host seeds, with otherwise the same conditions. For ancestors we scored

396 couples, and for evolved lines 789 couples, across both experimental years. LRS was

analysed per female, hence we divided all offspring counts from female triads by three before

analysis.

We carried out all statistical analyses using the statistical and programming software R (v.

3.6.1). All seven traits were first analysed separately using linear mixed effect models available

in the lme4 package[115]. When analysing differences between hot- and cold-adapted lines,

evolution regime, geographic origin and assay temperature were added as fully crossed fixed

effects. Line replicate crossed with assay temperature and nested within geographic origin

were added as random terms to assure correct level of replication when estimating significance

of fixed terms including evolution regime. For analyses on metabolic rate, water loss and

weight loss, the date of the respirometry run was added as an additional random effect.

Metabolic rate and weight loss were corrected for body mass of the measured female triad by

taking an average of the weight measures before and after respirometry and adding it as a

covariate in analysis. For water loss measured from evolved lines, measures from vial 2 (the

first vial in the measuring sequence containing beetles) and measurements from three entire

experimental days were discarded following an outlier analysis as these measures were

magnitudes greater than other measures and clearly represented water vapour from other

sources than the beetles. For LRS, experiment (triads and two separate rearings of single

couples) and its interaction with assay temperature were added as additional fixed effects to

account for possible block effects. The data from ancestral lines were analyzed in similar

fashion but without the fixed effect of evolution regime and random effect of population

replicate.

P-values were calculated using the car package and type II sums of squares with the Kenward-

Roger approximation for the degrees of freedom. To analyse evolutionary responses while

taking all traits into consideration at once, we also performed a non-parametric MANOVA using

residual randomization[65] with the same structure for fixed and random effects (**Fig. 4H**).

The agricultural footprint

We set up both evolved and ancestral lines in a common garden experiment including the 23,

29 and 35°C assay temperature, following approximately 85, 120 and 125 generations of

experimental evolution for cold-adapted, hot-adapted and ancestral lines, respectively. We

removed parental effects by moving all lines to 29°C two generations prior to the start of the

experiment. Newly laid F2 eggs were split among the three assay temperatures and resulting virgin adults were collected. Three males and three females were placed together in petri-dishes with ad libitum *V. unguiculata* seeds for 24h at 29°C and 35°C, and 34h at 23°C (as reproductive rate is slower at cold temperature; **Fig. 4B**). Three dishes were prepared per line and assay temperature. Note that the experiment was aimed at linking life-history variation to host consumption, and the sample size was low for detecting more fine-scaled differences between evolution regimes due to the imprecise measures of fecundity (6-9 females per line and temperature measured over 24-34h), which were much better estimated in the previous experiments (~80 females per line and temperature across the entire adult stage).

To quantify the amount of host seed consumed by beetles, each petri dish with beans was weighed prior to the addition of beetles, to record a starting weight of each assay. After beetles had emerged and been removed, assays were placed in standard conditions for 6 weeks during which they were weighed on several occasions to record the weight of the infested beans and to check for potential inconsistencies and time-dependencies in estimates of bean consumption. However, the measures of the beans' weight loss following infestation were highly repeatable (r^2 = 94%) and we chose the final (fourth) measure to calculate our estimate of host consumption. To account for potential weight loss of beans that was independent of the beetle infestation, we also set up 5 assays without beetles at each temperature and measured them in the exact same way. The host consumption was then corrected based on the weight changes in these control assays; this correction was very small and only corresponded to 1-2% of the total bean weight. For each assay, we also recorded the number of hatched eggs laid on beans (i.e. fecundity), the number of those that resulted in the adult pupating (i.e. juvenile survival), and the mean body mass of emerged adult beetles.

Differences in food consumption were first analyzed using linear mixed effect models in the lme4 package with the same structure for fixed and random terms as described for life-history traits. We excluded the ancestors from formal analysis since these had only half the samples compared to the evolution regimes, but we included them in **Fig. 5A** to show the direction of evolution of host consumption. We sequentially added information on fecundity, juvenile survival, and mean body mass for each assay to explore how life-history traits accounted for host consumption. We finally fitted a linear model including only the three traits (removing the fixed terms of assay temperature, evolution regime and geographic background) to estimate how much of the variation in host consumption that could be predicted by life-history alone.

Motivated by the observation that the life-history traits accounted for more than 93% of the observed variation in host consumption (see **Fig. 5A** and[66]), we calculated the two independent

measures of the agricultural footprint (see main text) from our measured life-history traits in the main experiment, as this data was estimated with high accuracy for all lines. To provide 95% confidence limits and P-values, we performed parametric bootstrap on all calculations using the means and standard errors for each trait per line and assay temperature derived from the univariate mixed models (**Fig. 4A-G, Suppl. 3**).

Acknowledgements

We thank Johanna Liljestrand-Rönn, Elina Immonen and Göran Arnqvist for help in the lab as well as for helpful input on our study design and results. Sequencing was performed by the SNP&SEQ Technology Platform in Uppsala. The facility is part of the National Genomics Infrastructure (NGI) Sweden and Science for Life Laboratory. The SNP&SEQ Platform is also supported by the Swedish Research Council and the Knut and Alice Wallenberg Foundation. This work was supported by grant no. 2019-05023 from the Swedish Research Council (VR), grant no. 2022-01117 from Formas, and grant no. CTS22:2101 from Carl Tryggers Stiftelse, to DB.

References

1. Lehmann, P. *et al.* Complex responses of global insect pests to climate warming. *Front. Ecol. Environ.* **18**, 141–150 (2020).
2. Deutsch, C. A. *et al.* Increase in crop losses to insect pests in a warming climate. *Science* **361**, 916–919 (2018).
3. Harvey, J. A. *et al.* Scientists' warning on climate change and insects. *Ecol. Monogr.* **93**, e1553 (2023).
4. Daly, E. Z. *et al.* A synthesis of biological invasion hypotheses associated with the introduction– naturalisation–invasion continuum. *Oikos* **n/a**, e09645.
5. González-Tokman, D. *et al.* Insect responses to heat: physiological mechanisms, evolution and ecological implications in a warming world. *Biol. Rev.* **95**, 802–821 (2020).
6. Diamond, S. E. Contemporary climate-driven range shifts: Putting evolution back on the table. *Funct. Ecol.* **32**, 1652–1665 (2018).
7. Harvey, J. A., Heinen, R., Gols, R. & Thakur, M. P. Climate change-mediated temperature extremes and insects: From outbreaks to breakdowns. *Glob. Change Biol.* **26**, 6685–6701 (2020).
8. Simberloff, D. *et al.* Impacts of biological invasions: what's what and the way forward. *Trends Ecol. Evol.* **28**, 58–66 (2013).
9. Diagne, C. *et al.* High and rising economic costs of biological invasions worldwide. *Nature* **592**, 571–576 (2021).
10. Diagne, C. *et al.* The economic costs of biological invasions in Africa: a growing but neglected threat? *NeoBiota* **67**, 11–51 (2021).
11. Change, C. R. P. on C. & Security, A. and F. *CGIAR Research Program on Climate Change, Agriculture and Food Security Annual Report 2020*. https://cgspace.cgiar.org/handle/10568/114818 (2021).

12. Adler, C. *et al.* Changes in the distribution and pest risk of stored product insects in Europe due
to global warming: Need for pan-European pest monitoring and improved food-safety. *J. Stored*
Prod. Res. **97**, 101977 (2022).
13. Urban, M. C. *et al.* Improving the forecast for biodiversity under climate change. *Science* **353**,
aad8466 (2016).
14. Macfadyen, S., McDonald, G. & Hill, M. P. From species distributions to climate change
adaptation: Knowledge gaps in managing invertebrate pests in broad-acre grain crops. *Agric.*
Ecosyst. Environ. **253**, 208–219 (2018).
15. Galante, P. J. *et al.* The challenge of modeling niches and distributions for data-poor species: a
comprehensive approach to model complexity. *Ecography* **41**, 726–736 (2018).
16. Hoffmann, A. A. Rapid adaptation of invertebrate pests to climatic stress? *Curr. Opin. Insect Sci.*
21, 7–13 (2017).
17. Helmuth, B., Kingsolver, J. G. & Carrington, E. BIOPHYSICS, PHYSIOLOGICAL ECOLOGY, AND
CLIMATE CHANGE: Does Mechanism Matter? *Annu. Rev. Physiol.* **67**, 177–201 (2005).
18. Huey, R. B. & Kingsolver, J. G. Evolution of thermal sensitivity of ectotherm performance. *Trends*
Ecol. Evol. **4**, 131–135 (1989).
19. Angilletta, M. J. *Thermal Adaptation: A Theoretical and Empirical Synthesis*. (Oxford University
Press, Oxford ; New York, 2009).
20. Hochachka, P. W. & Somero, G. N. *Biochemical Adaptation: Mechanism and Process in*
Physiological Evolution. (Oxford University Press, 2002).
21. Berger, D., Stångberg, J., Baur, J. & Walters, R. J. Elevated temperature increases genome-wide
selection on de novo mutations. *Proc. R. Soc. B Biol. Sci.* **288**, 20203094 (2021).
22. Kellermann, V., van Heerwaarden, B., Sgrò, C. M. & Hoffmann, A. A. Fundamental evolutionary
limits in ecological traits drive Drosophila species distributions. *Science* **325**, 1244–1246 (2009).
23. Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M. & West, G. B. Toward a Metabolic Theory of
Ecology. *Ecology* **85**, 1771–1789 (2004).
24. Savage, V. M., Gillooly, J. F., Brown, J. H., West, G. B. & Charnov, E. L. Effects of Body Size and
Temperature on Population Growth. *Am. Nat.* **163**, 429–441 (2004).
25. Bloom, J. D. *et al.* Thermodynamic prediction of protein neutrality. *Proc. Natl. Acad. Sci. U. S. A.*
102, 606–611 (2005).
26. Agozzino, L. & Dill, K. A. Protein evolution speed depends on its stability and abundance and on
chaperone concentrations. *Proc. Natl. Acad. Sci.* **115**, 9092–9097 (2018).
27. Chen, P. & Shakhnovich, E. I. Thermal adaptation of viruses and bacteria. *Biophys. J.* **98**, 1109–
1118 (2010).
28. Parratt, S. R. *et al.* Temperatures that sterilize males better match global species distributions
than lethal temperatures. *Nat. Clim. Change* 1–4 (2021) doi:10.1038/s41558-021-01047-0.
29. Walsh, B. S. *et al.* The Impact of Climate Change on Fertility. *Trends Ecol. Evol.* **34**, 249–259
(2019).
30. Jørgensen, L. B., Ørsted, M., Malte, H., Wang, T. & Overgaard, J. Extreme escalation of heat
failure rates in ectotherms with global warming. *Nature* **611**, 93–98 (2022).
31. Feder, M. E., Bennett, A. F. & Huey, R. B. EVOLUTIONARY PHYSIOLOGY1. 29 (2000).
32. Skoracka, A. *et al.* Effective specialist or jack of all trades? Experimental evolution of a crop pest
in fluctuating and stable environments. *Evol. Appl.* **15**, 1639–1652 (2022).
33. Alexandridis, N. *et al.* Archetype models upscale understanding of natural pest control response
to land-use change. *Ecol. Appl.* **n/a**, e2696.
34. Towards an interactive, process-based approach to understanding range shifts: developmental
and environmental dependencies matter. doi:10.1111/ecog.03975.
35. Segoli, M. *et al.* Trait-based approaches to predicting biological control success: challenges and
prospects. *Trends Ecol. Evol.* **38**, 802–811 (2023).
36. Barrett, R. & Schluter, D. Adaptation from standing genetic variation. *Trends Ecol. Evol.* **23**, 38–44
(2008).

37. Bennett, J. M. *et al.* The evolution of critical thermal limits of life on Earth. *Nat. Commun.* **12**,
1198 (2021).
38. Figueiredo, L., Krauss, J., Steffan-Dewenter, I. & Sarmento Cabral, J. Understanding extinction
debts: spatio–temporal scales, mechanisms and a roadmap for future research. *Ecography* **42**,
1973–1990 (2019).
39. Chen, Y. *et al.* The combination of genomic offset and niche modelling provides insights into
climate change-driven vulnerability. *Nat. Commun.* **13**, 4821 (2022).
40. Nadeau, C. P., Urban, M. C. & Bridle, J. R. Climates Past, Present, and Yet-to-Come Shape Climate
Change Vulnerabilities. *Trends Ecol. Evol.* **32**, 786–800 (2017).
41. Tuda, M., Kagoshima, K., Toquenaga, Y. & Arnqvist, G. Global Genetic Differentiation in a
Cosmopolitan Pest of Stored Beans: Effects of Geography, Host-Plant Usage and Anthropogenic
Factors. *PLOS ONE* **9**, e106268 (2014).
42. Jong, G. de & Noordwijk, A. J. van. Acquisition and Allocation of Resources: Genetic (CO)
Variances, Selection, and Life Histories. *Am. Nat.* **139**, 749–770 (1992).
43. Houle, D. Genetic Covariance of Fitness Correlates: What Genetic Correlations are Made of and
Why it Matters. *Evolution* **45**, 630 (1991).
44. Roff, D. *Evolution Of Life Histories: Theory and Analysis*. (Springer Science & Business Media,
1993).
45. Stearns, S. C. THE EVOLUTION OF LIFE HISTORIES.
46. Cornish-Bowden, A. Kinetics, Enzymes. in *Encyclopedia of Bioprocess Technology* (John Wiley &
Sons, Inc., 2002). doi:10.1002/0471250589.ebt123.
47. Evans, M. G. & Polanyi, M. Some applications of the transition state method to the calculation of
reaction velocities, especially in solution. *Trans. Faraday Soc.* **31**, 875–894 (1935).
48. Eyring, H. The Activated Complex in Chemical Reactions. *J. Chem. Phys.* **3**, 107–115 (1935).
49. Clarke, A. Is there a Universal Temperature Dependence of metabolism? *Funct. Ecol.* **18**, 252–256
(2004).
50. Dell, A. I., Pawar, S. & Savage, V. M. Systematic variation in the temperature dependence of
physiological and ecological traits. *Proc. Natl. Acad. Sci.* **108**, 10591–10596 (2011).
51. Gillooly, J. F., Charnov, E. L., West, G. B., Savage, V. M. & Brown, J. H. Effects of size and
temperature on developmental time. *Nature* **417**, 70–73 (2002).
52. Gillooly, J. F., Brown, J. H., West, G. B., Savage, V. M. & Charnov, E. L. Effects of Size and
Temperature on Metabolic Rate. *Science* **293**, 2248–2251 (2001).
53. Huey, R. B. & Kingsolver, J. G. Climate Warming, Resource Availability, and the Metabolic
Meltdown of Ectotherms. *Am. Nat.* **194**, E140–E150 (2019).
54. Dahlke, F. T., Wohlrab, S., Butzin, M. & Pörtner, H.-O. Thermal bottlenecks in the life cycle define
climate vulnerability of fish. *Science* **369**, 65–70 (2020).
55. Thomas, M. K. *et al.* Temperature–nutrient interactions exacerbate sensitivity to warming in
phytoplankton. *Glob. Change Biol.* **23**, 3269–3280 (2017).
56. Angilletta, M. J. *Thermal Adaptation: A Theoretical and Empirical Synthesis*. (OUP Oxford, 2009).
57. Huey, R. B. & Kingsolver, J. G. Variation in universal temperature dependence of biological rates.
Proc. Natl. Acad. Sci. **108**, 10377–10378 (2011).
58. Dill, K. A., Ghosh, K. & Schmit, J. D. Physical limits of cells and proteomes. *Proc. Natl. Acad. Sci.*
108, 17876–17882 (2011).
59. Clarke, A. Temperature and the Metabolic Theory of Ecology. *Funct. Ecol.* **20**, 405–412 (2006).
60. Taipale, M., Jarosz, D. F. & Lindquist, S. HSP90 at the hub of protein homeostasis: emerging
mechanistic insights. *Nat. Rev. Mol. Cell Biol.* **11**, 515–528 (2010).
61. Mayer, M. P. & Bukau, B. Hsp70 chaperones: Cellular functions and molecular mechanism. *Cell.*
Mol. Life Sci. **62**, 670–684 (2005).
62. Kingsolver, J. G. & Woods, H. A. Beyond Thermal Performance Curves: Modeling Time-
Dependent Effects of Thermal Stress on Ectotherm Growth Rates. *Am. Nat.* **187**, 283–294 (2016).
63. Echave, J. & Wilke, C. O. Biophysical models of protein evolution: understanding the patterns of
evolutionary sequence divergence. *Annu. Rev. Biophys.* **46**, 85–103 (2017).

64. Immonen, E., Berger, D., Sayadi, A., Liljestrand-Rönn, J. & Arnqvist, G. An experimental test of
temperature-dependent selection on mitochondrial haplotypes in Callosobruchus maculatus
seed beetles. *Ecol. Evol.* **10**, 11387–11398 (2020).

65. Collyer, M. L. & Adams, D. C. RRPP: An r package for fitting linear models to high-dimensional
data using residual randomization. *Methods Ecol. Evol.* **9**, 1772–1779 (2018).

66. Umeozor, O. C. Effect of the Infection of Callosobruchus maculatus (Fab.) on the Weight Loss of
Stored Cowpea (Vigna unguiculata (L.) Walp). *O C.*

67. Dent, D. & Binks, R. H. *Insect Pest Management, 3rd Edition*. (CABI, 2020).

68. Urban, M. C. Accelerating extinction risk from climate change. *Science* **348**, 571–573 (2015).

69. Wagner, D. L. Insect Declines in the Anthropocene. *Annu. Rev. Entomol.* **65**, 457–480 (2020).

70. Deutsch, C. A. *et al.* Impacts of climate warming on terrestrial ectotherms across latitude. *Proc.*
Natl. Acad. Sci. **105**, 6668–6672 (2008).

71. Chown, S. L. & Gaston, K. J. Body size variation in insects: a macroecological perspective. *Biol.*
Rev. **85**, 139–169 (2010).

72. Urban, M. C. *et al.* Coding for Life: Designing a Platform for Projecting and Protecting Global
Biodiversity. *BioScience* **72**, 91–104 (2022).

73. Angilletta, M. J., Huey, R. B. & Frazier, M. R. Thermodynamic Effects on Organismal Performance:
Is Hotter Better? *Physiol. Biochem. Zool.* **83**, 197–206 (2010).

74. Berger, D., Walters, R. & Gotthard, K. What limits insect fecundity? Body size- and temperature-
dependent egg maturation and oviposition in a butterfly. *Funct. Ecol.* **22**, 523–529 (2008).

75. Berger, D. *et al.* Intralocus Sexual Conflict and the Tragedy of the Commons in Seed Beetles. *Am.*
Nat. **188**, E98–E112 (2016).

76. Berger, D., Berg, E. C., Widegren, W., Arnqvist, G. & Maklakov, A. A. Multivariate intralocus
sexual conflict in seed beetles: SEXUAL CONFLICT AND LIFE-HISTORY PLEIOTROPY. *Evolution* **68**,
3457–3469 (2014).

77. Mallard, F., Nolte, V., Tobler, R., Kapun, M. & Schlötterer, C. A simple genetic basis of adaptation
to a novel thermal environment results in complex metabolic rewiring in Drosophila. *Genome*
Biol. **19**, 119 (2018).

78. Rogell, B. *et al.* Sex-dependent evolution of life-history traits following adaptation to climate
warming. *Funct. Ecol.* **28**, 469–478 (2014).

79. Ivimey-Cook, E. R., Piani, C., Hung, W.-T. & Berg, E. C. Genetic background and thermal regime
influence adaptation to novel environment in the seed beetle, Callosobruchus maculatus. *J. Evol.*
Biol. **37**, 1–13 (2024).

80. Angilletta, Jr., , Michael J. & Dunham, A. E. The Temperature-Size Rule in Ectotherms: Simple
Evolutionary Explanations May Not Be General. *Am. Nat.* **162**, 332–342 (2003).

81. Zuo, W., Moses, M. E., West, G. B., Hou, C. & Brown, J. H. A general model for effects of
temperature on ectotherm ontogenetic growth and development. *Proc. R. Soc. B Biol. Sci.* **279**,
1840–1846 (2011).

82. Awmack, C. S. & Leather, S. R. Host Plant Quality and Fecundity in Herbivorous Insects. *Annu.*
Rev. Entomol. **47**, 817–844 (2002).

83. Atkinson, D. & Sibly, R. M. Why are organisms usually bigger in colder environments? Making
sense of a life history puzzle. *Trends Ecol. Evol.* **12**, 235–239 (1997).

84. Blanckenhorn, W. U. Bergmann and Converse Bergmann Latitudinal Clines in Arthropods: Two
Ends of a Continuum? *Integr. Comp. Biol.* **44**, 413–424 (2004).

85. Horne, C. R., Hirst, Andrew. G. & Atkinson, D. Temperature-size responses match latitudinal-size
clines in arthropods, revealing critical differences between aquatic and terrestrial species. *Ecol.*
Lett. **18**, 327–335 (2015).

86. Stillwell, R. C., Wallin, W. G., Hitchcock, L. J. & Fox, C. W. Phenotypic plasticity in a complex
world: interactive effects of food and temperature on fitness components of a seed beetle.
Oecologia **153**, 309–321 (2007).

87.   Garcia-Robledo, C., Baer, C. S., Lippert, K. & Sarathy, V. Evolutionary history, not ecogeographic
867   rules, explains size variation of tropical insects along elevational gradients. *Funct. Ecol.* **34**, 2513–
868   2523 (2020).
869   88.   Martin. Goal 2: Zero Hunger. *United Nations Sustainable Development*
870   https://www.un.org/sustainabledevelopment/hunger/.
871   89.   Climate Change 2022: Impacts, Adaptation and Vulnerability.
872   https://www.ipcc.ch/report/ar6/wg2/.
873   90.   Cabral, J. S. *et al. The Road to Integrate Climate Change Effects on Land-Use Change in Regional*
874   *Biodiversity Models*. https://www.authorea.com/users/409019/articles/558105-the-road-to-
875   integrate-climate-change-effects-on-land-use-change-in-regional-biodiversity-
876   models?commit=1a1c610450c40c57b8bc51276fcbcc05242acfa5 (2022)
877   doi:10.22541/au.164608831.19029067/v1.
878   91.   Bocedi, G. *et al.* RangeShifter 2.0: an extended and enhanced platform for modelling spatial eco-
879   evolutionary dynamics and species' responses to environmental changes. *Ecography* **44**, 1453–
880   1462 (2021).
881   92.   Hagen, O. *et al.* gen3sis: A general engine for eco-evolutionary simulations of the processes that
882   shape Earth's biodiversity. *PLOS Biol.* **19**, e3001340 (2021).
883   93.   Dormann, C. F. *et al.* Correlation and process in species distribution models: bridging a
884   dichotomy. *J. Biogeogr.* **39**, 2119–2131 (2012).
885   94.   Mouquet, N. *et al.* REVIEW: Predictive ecology in a changing world. *J. Appl. Ecol.* **52**, 1293–1310
886   (2015).
887   95.   Zurell, D. *et al.* Benchmarking novel approaches for modelling species range dynamics. *Glob.*
888   *Change Biol.* **22**, 2651–2664 (2016).
889   96.   Anderson, R., Bayer, P. E. & Edwards, D. Climate change and the need for agricultural adaptation.
890   *Curr. Opin. Plant Biol.* **56**, 197–202 (2020).
891   97.   Garcia-Costoya, G., Williams, C. E., Faske, T. M., Moorman, J. D. & Logan, M. L. Evolutionary
892   constraints mediate extinction risk under climate change. *Ecol. Lett.* **26**, 529–539 (2023).
893   98.   Beck, C. W. A Handbook on Bean Beetles,. 15.
894   99.   Fox, C. W. Multiple Mating, Lifetime Fecundity and Female Mortality of the Bruchid Beetle,
895   Callosobruchus maculatus (Coleoptera: Bruchidae). *Funct. Ecol.* **7**, 203–208 (1993).
896   100.   Berger, D. *et al.* INTRALOCUS SEXUAL CONFLICT AND ENVIRONMENTAL STRESS: SEX,
897   GENES, AND CONFLICT IN STRESSFUL ENVIRONMENTS. *Evolution* n/a-n/a (2014)
898   doi:10.1111/evo.12439.
899   101.   Baur, J., Jagusch, D., Michalak, P., Koppik, M. & Berger, D. The mating system affects
900   the temperature sensitivity of male and female fertility. *Funct. Ecol.* **36**, 92–106 (2022).
901   102.   Baur, J., Zwoinska, M., Koppik, M., Snook, R. R. & Berger, D. Heat stress reveals a
902   fertility debt owing to postcopulatory sexual selection. *Evol. Lett.* qrad007 (2023)
903   doi:10.1093/evlett/qrad007.
904   103.   Babraham Bioinformatics - FastQC A Quality Control tool for High Throughput
905   Sequence Data. https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.
906   104.   Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: summarize analysis results
907   for multiple tools and samples in a single report. *Bioinforma. Oxf. Engl.* **32**, 3047–3048 (2016).
908   105.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
909   sequence data. *Bioinformatics* **30**, 2114–2120 (2014).
910   106.   Arnqvist, G. *et al.* A chromosome-level assembly of the seed beetle Callosobruchus
911   maculatus genome with annotation of its repetitive elements. *G3 GenesGenomesGenetics*
912   jkad266 (2023) doi:10.1093/g3journal/jkad266.
913   107.   Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome
914   alignment and genotyping with HISAT2 and HISAT-genotype. *Nat. Biotechnol.* **37**, 907–915
915   (2019).
916   108.   Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with high-
917   throughput sequencing data. *Bioinforma. Oxf. Engl.* **31**, 166–169 (2015).

109.   R Core Team. *R: A Language and Environment for Statistical Computing*. (R Foundation
for Statistical Computing, Vienna, Austria, 2020).

110.   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140
(2010).

111.   McCarthy, D. J., Chen, Y. & Smyth, G. K. Differential expression analysis of multifactor
RNA-Seq experiments with respect to biological variation. *Nucleic Acids Res.* **40**, 4288–4297
(2012).

112.   Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing
and microarray studies. *Nucleic Acids Res.* **43**, e47 (2015).

113.   Falcon, S. & Gentleman, R. Using GOstats to test gene lists for GO term association.
Bioinforma. Oxf. Engl. **23**, 257–258 (2007).

114.   Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics
data. *The Innovation* **2**, 100141 (2021).

115.   Bates, D. Mixed models in R using the lme4 package Part 4: Inference based on
profiled deviance. 23.

116.   Jackai, L. E. N. & Daoust, R. A. Insect Pests of Cowpeas. *Annu. Rev. Entomol.* **31**, 95–
119 (1986).

Figure 1: Thermal adaptation via compensatory energy acquisition can increase pest species' impact on agriculture.

Shown are predictions based on protein kinetics (details in Supplementary 1). **A**) Temperature increases enzymatic reaction rates[20,51] (pink lines) while too hot temperatures lead to (reversable) protein unfolding (grey lines), leaving fewer enzymes available to do work[25,26]. Reduced costs of feeding (broken lines) leads to more body reserves acquired (*M*), of which a fraction (*p*) can be devoted to increased production of catalytic enzymes that speed up constrained reaction rates at cold temperature[20], and another fraction (*q*) that can be devoted to molecular chaperones that stabilize proteins at hot temperature[31]. **B**) The resulting growth rate is the product of the remaining body reserves that can be devoted to growth and reproduction, *M* (1-*p*-*q*), the reaction rate, and the proportion of properly folded protein. **C**) The resulting growth rates for populations adapting to cold (23°C; blue) and hot (35°C; red) temperature via compensatory responses (increased *M*, and allocation to *p* and *q*) are shown for high (full lines) and low (broken lines) costs of feeding. While the cold-adapted population outcompetes the hot-adapted population at cold temperature, growth rates are greater at warm temperatures and for the hot-adapted population due to thermodynamic constraints on reaction rates[23,51]. Horizontal lines on the right-hand y-axis show *M* at the optimal strategy; lower costs of feeding favour increased energy acquisition, which improves thermal performance and increases thermal niche breadth. In the depicted scenario, increased energy acquisition is particularly beneficial at hot temperature that puts increased demands on cellular maintenance to ensure protein stability. **D)** The possibility to evolve thermal performance via compensatory energy acquisition leads to an even greater agricultural impact (the product of *M* and population growth rate; compare difference between full and hatched lines in panels **C** and **D**)**.** All panels show responses relative to the maximum value for ancestors reared at benign 29°C at high costs of feeding (black full line in panel B).

Figure 2: Experimental evolution of thermal niches and life history in seed beetles.
Dryland legumes are nutritionally rich, drought tolerant and resilient to harsh weather[11,116] and provide subsistence for more than half a billion people in the driest regions of the world[11]. However, these crops suffer substantial losses from seed beetles[116]. To investigate how genetic adaptation to climate change might shape the impact of seed beetles on legume crops, we performed large-scale phenotyping of long-term experimental evolution lines of *C. maculatus*. **A**) Populations from three geographic locations (Brazil, California and Yemen) were sampled and allowed to adapt to common laboratory conditions for ca. 300 generations before experimental evolution. Four replicate populations were created per genetic background and split between the two evolution regimes (23°C and 35°C), while the original ancestor remained at the benign temperature (29°C). Experimental evolution proceeded over 90 and 130 generations for the Cold (blue) and Hot (red) adapted populations. During this time, the ancestors went through 130 generations of evolution. **B**) To provide molecular evidence for a temperature-dependent trade-off between maintenance and reproduction, gene expression was measured in both evolved and ancestral lines after 80-125 generations of evolution. **C**) After 45 (Cold adapted) and 60 (Hot adapted) generations, life-history adaptation was assessed in a common garden experiment. This data was complemented in generations 90 (Cold lines) and 130 (Hot Lines and Ancestors). Life-history traits were measured in the three ancestors after 120 generations of evolution. **D**) Following 80-125 generations of experimental evolution, the host consumption of all lines and ancestors was scored in a common garden to assess how life-history evolution under simulated climate change had shaped the agricultural footprint left by *C. maculatus*.

Figure 3. Molecular evidence for a temperature-dependent trade-off between maintenance and reproduction.

A) Differentially expressed genes in response to heat shock (HS) and mating (R); 137 genes responded to both treatments, which was more than expected by chance. **B**) Of these genes, all but three showed antagonistic responses to the two treatments, signifying a trade-off between investment in reproduction and maintenance under heat stress. **C**) GO term enrichments for genes upregulated (+) or downregulated (-) in response to mating (R) and heat stress (HS), as well as for the overlapping antagonistic genes. **D)** The expression of the 134 antagonistic genes in the experimental evolution lines and ancestors indicate a clear increase in allocation towards cellular maintenance at the cost of reduced reproduction when beetles were reared at 35°C (the thermal regime of hot-adapted lines). However, there were no consistent differences in expression between evolution regimes.

1003

1004

1005

Figure 4. Thermal adaptation in life-history traits.
Single traits in (**A-G**) for Hot (red) and Cold (blue) adapted lines, as well as their Ancestors
(black). Development time is presented as a rate (1/development time in days) for consistency
with other traits. Means ± 1SE. **H**) Ordination using principal component analysis showing that
multivariate divergence occurs along both a temperature-sensitive axis (PC1) and along a
general life-history axis describing overall variation in body mass and LRS (PC2), with more
divergence from the ancestor in Hot-adapted lines.
1013

Figure 5: Temperature-dependent evolution of the agricultural footprint.
A) The relationship between beetle mass produced and host bean consumed. Each assay is represented by an individual datapoint. The slopes of the dashed lines correspond to the temperature-specific constant (C_T) giving the amount of host consumed per beetle produced (23°C: 3.17, 29°C: 3.66, 35°C: 3.55). **B**) The two independent metrics used to predict the agricultural footprint. **C**) The predicted footprint of each line (using the first metric) relative to respective ancestor raised at 29°C (grey hatched line). Means ± 95% CIs. The effect of life-history adaptation at cold temperature is small, but life-history adaptation at hot temperature almost doubles the agricultural footprint relative to the ancestral impact.