Life-history adaptation under climate warming magnifies the agricultural footprint of a cosmopolitan insect pest

Estelle Burc[1,3†], Camille Girard-Tercieux[1,2,5†], Moa Metz[1,4], Elise Cazaux[1,5], Julian Baur[1],

Mareike Koppik[1,6], Alexandre Rêgo[1], Alex Hart[1] & David Berger[1]*

1. Department of Ecology and Genetics, Program of Animal Ecology. Uppsala University, Norbyvägen 18D, 75236, Uppsala, Sweden.
2. AMAP, Université de Montpellier, CIRAD, CNRS, INRAE, IRD, Montpellier, France
3. Agronomy Institute Rennes-Angers (IARA), Graduate school of agronomy, 35000 Rennes, France
4. Department of Biology, Faculty of Natural Sciences, Norwegian University of Science and Technology, 7491 Trondheim, Norway
5. Ecole nationale supérieure agronomique de Toulouse (ENSAT), Graduate school of agronomy, Toulouse, France.
6. Department of Zoology, Animal Ecology, Martin-Luther University Halle-Wittenberg, Halle (Saale), Germany.

†Both authors contributed equally and are presented in alphabetical order.

*Corresponding author.

david.berger@ebc.uu.se

Keywords: life history, temperature, climate change, global warming, agriculture, insect, pest, Callosobruchus maculatus, Vigna unguiculata, cowpea.

Abstract

Climate change is affecting population growth rates of ectothermic pests with potentially dire consequences for agriculture, but how rapid genetic adaptation impacts these dynamics remains unclear. To address this challenge, we predicted how climate change adaptation in life-history traits of insect pests may affect future agricultural yields by unifying thermodynamics based on first principles with classic life-history theory. Our model predicts that warming temperatures favour changes in resource allocation decisions coupled with increased larval host consumption, resulting in a predicted double-blow on agricultural yields under future climate change. We find support for these predictions by studying thermal adaptation in life-history traits and underlying gene expression in the wide-spread insect pest, *Callosobruchus maculatus*, with five years of life-history evolution under experimental warming causing an almost two-fold increase in its predicted agricultural footprint. These results emphasize the need for integrating a mechanistic understanding of life-history evolution into forecasts of pest impact.

Introduction

Climate change is predicted to impact growth rates and distribution ranges of ectothermic pests[1-7] with potentially severe outcomes for agricultural economics and world food supply[8-12]. To mitigate these events, forecasting of pests' responses to future climates is an important task for biologists[13,14]. Current forecasts use deterministic models that rely on information about species niches and climate data. However, these projections typically do not incorporate the potential for genetic evolution of climate niches and key life-history traits that mediate the effect that pests have on their hosts[13,14]. Indeed, the evolution of ecological niches has been rapid in many cosmopolitan pests and not accounting for this adaptive potential may therefore provide inaccurate forecasts of their future agricultural impact[1,7,13-17].

Ectothermic organisms are constrained by temperature-dependent biophysical properties of cells[18-22]. On the one hand, metabolism is constrained by cold temperature that slows down both juvenile development and adult reproduction[19], predicting that climate warming may lead to faster population growth rates via plastic increases in these rate-dependent life-history traits[23,24] and thereby worsen pest impact in many parts of the world[2]. On the other hand, acute hot temperatures can jeopardize cellular homeostasis[25-27], leading to rapid fertility decline[28,29] and increased mortality[22,30], which predicts that climate warming may reduce insect growth rates in the world's hottest regions[2,31]. These thermodynamic constraints are thus expected to have a strong influence on ectotherm pests facing climate warming, but how genetic adaptation to temperature may influence long-term outcomes remains unclear. For example, theory predicts that thermal adaptation should be governed by trade-offs and that shifts in physiology allowing superior performance at one temperature extreme should lead to reduced performance at the other[18,20,32,33]. However, when thermal adaptation occurs it often does so seemingly without any apparent trade-offs, with some genotypes displaying superior performance across the entire thermal range studied[19,33-35].

Indeed, a typical signature of insect pests is their broad climate niches coupled with fast growth rates[1,16]. One mechanism that can allow organisms to occupy broad climate niches is up-regulation of molecular chaperones that aid cellular homeostasis under harsh conditions[20,36]. Such responses are energetically costly and therefore need to be compensated by either allocation of resources away from reproduction in favour of cellular maintenance[20,36], or by acquiring more resources overall by increased food intake[5,19]. Notably, the first strategy, involving allocation trade-offs, is predicted to increase survival but reduce insect reproductive rates at stressful temperatures, whereas the second strategy, to increase resource acquisition, permits both sustained adult survival and reproductive output but is only possible if larvae can increase their host consumption. This makes the prediction that the effect of climate warming on insect growth rates may critically depend on the access to host plants[37]. Moreover, applying this reasoning to pest species suggests that the rich host

82 abundance supplied by agricultural crops might make the strategy to increase energy
83 acquisition particularly favourable, which could result in increased per capita host
84 consumption by pests facing stressful temperatures. These different routes to deal with
85 stressful temperatures can thus have important consequences for predicting the agricultural
86 impact of insects under climate change[1,14,17,38–40].
87
Despite of its importance, however, how insect life-histories are optimized under climate
change remains unclear[1,16,39]. Moreover, for adaptation to occur, genetic variation is a
prerequisite. Geographically isolated populations can differ greatly in the abundance of both
segregating and fixed genetic variation due to differences in their evolutionary past, which
may affect future adaptive potential[41–45]. Such historical contingencies may be of particular
importance in pests species, as geographically isolated populations often originate from
human-mediated long distance dispersal events of only a few individual genetic founders.
Evolution under climate change may therefore proceed along different trajectories and rates
across a species distribution range, even when climates change in parallel.
97
Here we explored how thermal adaptation in life-history traits of insect pests affect their
population growth rates and host consumption. We first constructed a model of thermal
adaptation in insect life-history by combining theory on thermodynamics[46,47] and acquisition
and allocation trade-offs[48,49]. The model predicts that warming temperatures not only
increase the potential for fast growth rates, but also benefit the evolution of increased host
consumption in larvae, which allows sustained survival and reproduction in adults, causing a
double-blow to agricultural yields. We then employed experimental evolution in the seed
beetle, *Callosobruchus maculatus*, an economically important cosmopolitan pest on legume
seeds[50]. Following 5 years of experimental evolution at either hot or cold temperature, we
assayed several life-history traits predicted to affect agricultural loss, including development
rate, metabolic rate, body size and adult offspring production. In accordance with model
predictions, life-history evolution in response to hot temperature caused an almost two-fold
increase in the agricultural footprint of *C. maculatus* that was mediated by increases in larval
host consumption, in turn leading to increased adult body mass and reproductive output. This
effect of life-history adaptation was contingent on the genetic make-up of the founding
populations and was not seen in populations adapting to cold temperature. These findings
suggest that equipping species distribution models with genetically explicit details accounting
for life-history evolution can lead to a substantial improvement in predictive accuracy of pest
impact under climate change, even over relatively short time scales.
117

Results

The agricultural impact of thermal adaptation via compensatory resource acquisition

To formalize predictions for how thermal adaptation in insect life-history will impact
agriculture, we combined the classic theoretical framework of energy acquisition and
allocation trade-offs[48,49,51,52] with the thermodynamic laws that govern enzyme kinetics[20,53].
Warm temperatures increase productivity rates of ectotherms[19,23] because biological rates
are at some level governed by enzymatic reaction rates, k_B, that scale with temperature
according to[46,47]:
126

$$k_B \propto A_B e^{-Ea_B/RT}$$
(Eq. 1)

128
where Ea_B is the activation energy required for the reaction to occur, R is the universal gas
constant, T is temperature in Kelvin, and A_B is a species and rate specific constant that may
itself evolve to accommodate effects of temperature[23,54,55] (Fig. 1A). Complementing equation
(1) with the power law describing how productivity scales with body mass, m, results in the
well-known prediction of mass and temperature-dependent population growth based on the
metabolic theory of ecology[23,24]: $r \propto m^b k_B$, where b is the allometric exponent, often taken to
be 0.75 based on the physics of distribution networks in plants and animals[23,56] (but see[57–59]).

However, excessively hot temperatures lead to increased molecular failure rates[30] due to a
range of detrimental processes that also scale with temperature, including protein
unfolding[25,26], metabolic expenditure[37,60], production of reactive oxygen species[20], and
oxygen depletion[61]. Several studies have highlighted that activation energies (*Ea*) can
themselves be temperature-dependent, which is particularly apparent for molecular failure
rates, that show a disproportionate increase at hot temperature[18–20,30,62] (Fig. 1A). To account
for this increase, mortality rate can be described by equation (1) with its own, temperature-
dependent, activation energy, $Ea_D(T)$[21,26,30,63,64] and rate-specific constant (A_D):

$$k_D \propto A_D e^{-Ea_D(T)/RT}$$ (Eq. 2)

As follows, population growth rate is the result of temperature-dependent productivity and
mortality rate[1,2] (Fig. 1B). Inherent enzyme properties governing the temperature-
dependence of reaction rate kinetics have been argued to evolve more slowly than thermal
adaptation via evolution of compensatory stress responses[20,36,55,65–67]. Such compensatory
responses involve upregulation of molecular chaperones, such as heat-shock proteins[20,68,69],
that maintain cellular homeostasis at stressful temperatures but are costly to produce[36,68,70].
Here we therefore integrated the thermodynamic predictions of equations (1) and (2) into a
life-history framework incorporating energy acquisition and allocation trade-offs[48,49,70]. First,
assume that energy reserves, *M*, are accumulated by larval host feeding at a cost, *c*, so that
survival to maturity equals: $(1-c)^M$. Further assume that accumulated energy reserves can be
allocated with fraction *p* to increase enzymatic reaction rates, and with fraction *q* to decrease
molecular failure rates, leaving $M(1-p-q)$ resources that can be devoted to productivity, *B*,
(i.e. growth and reproduction) according to:

$$B \propto [M(1-p-q)]^b \; ; \quad A_B \propto pM \; ; \quad A_D \propto 1/qM$$ (Eq. 3)

where A_B and A_D are the rate-specific constants for productivity and mortality, respectively.
Note that we here chose to model effects of energy acquisition and allocation via *A*, and not
Ea. This choice has both a practical and biological motivation as parametrization is
straightforward and seems justified; a proportional change in allocation to *A* leads to a
proportional change in reaction rates, but we acknowledge that our approach does not
capture the full range of possibilities for how energy allocation and acquisition trade-offs can
affect thermal performance[20,65,71–73]. Population growth rate, r', is then:

$$r'(T) \propto (1-c)^M * B * A_B \, e^{-\frac{Ea_B}{RT}} * \frac{1}{1 + A_D \, e^{-\frac{Ea_D(T)}{RT}}}$$ (Eq. 4)

We can find the optimal strategy of juvenile resource acquisition (increasing *M* at cost *c*) and
allocation (of *M* to *p* and *q*) at different temperatures by numerically solving for the
combination of *M*, *p* and *q* that maximize r'. The solution to this multidimensional trade-off
depends on the specific form of equations (1-3) and parameters that govern temperature
sensitivity (i.e. *Ea, A* and their implementation); mechanistic details which are empirically not
well-defined. Indeed, equations (1-2) represent phenomenological rather than mechanistic
relationships, emphasizing the need for further experimental work quantifying these
relationships.

Nevertheless, the qualitative predictions from this heuristic model are robust (see also [37]).
We provide a parameterized example in Figure 1 grounded in knowledge of how temperature
affects protein catalysis and stability[21,25,27,63,64,67]. Note that, because our focus is on climate
warming, our example models the effects of temperature on reaction rates and the
detrimental effects of acute hot temperatures via loss of protein stability, but does not model
lethal effects of acute cold temperatures (details in Supplementary 1). First, and as pointed

out previously[2], warm temperatures increase growth rates and agricultural impact via
thermodynamic effects on enzyme reaction rates. Second, hot temperatures that jeopardize
survival favour increased larval resource acquisition and more energy reserves (*M*) that can
be allocated to maintenance[37] (Supplementary Figure S1A, Fig. 1B). When increased
resource acquisition comes at low cost, this life-history strategy is particularly favourable and
results in increased thermal niche breadth and higher population growth rates (Fig. 1C).
Third, this has an even greater agricultural impact, given by the product of population growth
rate (*r*) and per capita host consumption (assumed proportional to *M*) (Fig. 1D). In fact, the
strategy to increase resource acquisition at hot temperature causes populations experiencing
temperatures slightly higher than those which maximize growth rate to have the greatest
agricultural impact (Supplementary Figure S1).
200
Experimental evolution of life-history traits and agricultural impact
To evaluate the prediction that trait and temperature-specific responses in pest species' life-
history can influence their agricultural impact under climate change, we compared twelve
experimental evolution lines of the seed beetle *C. maculatus*; a capital breeding
holometabolous insect that when kept in aphagous conditions acquires all its resources for
adult reproduction and survival in the larval stage[74]. Six lines were adapted to 35°C (*Hot*) and
six lines adapted to 23°C (*Cold*). All lines were reared on seeds from their preferred and
economically important host, black-eyed bean (*Vigna unguiculata*) (Fig. 2). The lines were
derived from three geographically isolated founding populations (*Ancestors*, kept at 29°C)
originating from Brazil, California (USA) and Yemen[75], allowing us to assess the importance
of geographic differences and evolutionary history for future responses to climate change.
212
Molecular evidence for a temperature-dependent trade-off between reproduction and
maintenance
To provide evidence for the model assumption of a trade-off between investment in
reproduction and cellular maintenance under thermal stress (Equation 3), we analysed data
on gene expression from two separate experiments on *C. maculatus*. These datasets
contained genes that were significantly differentially expressed in female abdomens in
response to heat shock, and between virgin and reproducing females. We identified 1269
reproduction-related genes and 765 heat stress genes. We found 137 genes present in both
datasets, which is significantly more than expected by chance (exp: 83.4 genes, χ^2 = 37.5, df
= 1, p < 0.001, Fig. 3A). Moreover, only 3 of the 137 genes showed concordant responses to
mating and heat-shock (all three decreased in expression), whereas 134 genes showed
antagonistic responses (binomial test: P < 0.001, Fig. 3B), providing strong evidence for a
trade-off between reproduction and the heat stress response. Out of these genes, 21 were
up-regulated in response to reproduction and down-regulated in response to heat. These
genes included common ribosomal genes that are crucial for protein translation, signifying
reproductive investment. The 113 antagonistic genes that were down-regulated in response
to mating and up-regulated in response to heat stress included common insect heat-shock
proteins (Supplementary Table 2a). GO enrichment analysis showed that the antagonistic
genes that overlapped both gene sets were enriched for biological process involved in
protein folding and degradation of unfolded proteins, which are essential components of
cellular maintenance under thermal stress (Fig. 3C, Supplementary Table 2b).
234
To confirm that the revealed trade-off also was at play in the thermal conditions used during
experimental evolution, we analysed gene expression data from all Cold and Hot lines, as
well as their Ancestors, reared in a common garden design including 23, 29 and 35°C assay
temperature. Each evolution line was represented by a single mRNA library per temperature,
based on a pool of 10 abdomens from mated young females of the same age as the females
phenotyped for life-history traits (see Fig. 4, further below). The ancestors were represented
by two such replicates. To evaluate investment into reproduction versus maintenance, we
calculated a score for each of these mRNA libraries along the axis describing the trade-off
based on the expression of the 134 antagonistic genes (Fig. 3B). We then analysed

differences between assay temperatures and evolution regimes using nested ANOVAS
(Supplementary Table 2c). Beetles reared at 35°C showing a strong change in allocation
towards cellular maintenance and away from reproduction compared to beetles reared at the
other two temperatures (assay temperature: F = 263.0, P < 0.001, Fig. 3D). There were no
significant differences between evolution regimes overall (F = 0.12, P = 0.89), although the
differences between the evolution regimes and their ancestors showed a dependence on
geographic origin (interaction: F = 11.0, P = 0.002). Qualitatively similar results were
obtained if restricting the analysis to the 113 genes that were upregulated in response to
heat stress (lower right quadrant of Fig. 3b) or when analysing all 765 heat stress genes and
1269 reproduction genes (Supplementary Figures S2 and S3, Supplementary Tables 2d, f,
and g). When restricting the analysis to the 21 overlapping genes that were upregulated in
response to mating (upper left quadrant of Fig. 3b), however, ancestors showed overall
higher expression than both the cold and hot regime (Supplementary Figure S3,
Supplementary Table 2e), hinting at a potential decrease during experimental evolution in
some aspects of reproduction early in life. These analyses nevertheless imply that there has
been little temperature-specific adaptation in resource allocation overall (i.e. q in Equation 3).
This result further predicts that if hot-adapted lines have evolved a means to deal with the
increased cellular stress of hot temperature (35°C), they must have done so mainly via
increased resource acquisition (M in Equation 3).
263
Thermal adaptation in life-history traits
We assessed the predicted agricultural impact of life-history evolution by assaying three
classic traits; juvenile development time, female adult body mass, and lifetime reproductive
success (LRS: number of offspring produced), and four rate-dependent traits assayed over
the first 16h of female reproduction; early fecundity, weight loss, water loss and metabolic
rate (Fig. 4). This also allowed us to evaluate if Hot lines had adapted to their temperature
regime by increased energy acquisition ($M;$ approximated by adult body mass at eclosion).
271
To characterize trait-values at the start of experimental evolution, we assayed the three
ancestors reared at 23, 29 and 35°C. This allowed us to quantify both the direction and
magnitude of adaptation during experimental evolution, as well as its subsequent agricultural
impact (*see below*). Temperature affected the expression of all seven life-history traits. For
development time, body mass, early fecundity and LRS, there were also main effects of the
geographic origin of the ancestors. Body mass, early fecundity and LRS also showed a
significant interaction between geographic origin and assay temperature, indicating
differences in thermal plasticity between ancestors (Fig. 4A-G, Supplementary 3).
280
To assess temperature-dependent life-history evolution we assayed the 12 evolved lines
when reared in a common garden experiment including 23°C and 35°C. All traits except
metabolic rate and weight loss showed significant differentiation between the Hot and Cold
regime. In particular, the Hot regime had indeed evolved larger adult body mass and higher
reproductive output (Fig. 4A-G), in accordance with model predictions (Fig. 1C). All traits
except metabolic rate and LRS showed a significant interaction between regime and assay
temperature, demonstrating thermal adaptation *sensu stricto* (Fig. 4B-G). Several alternative
estimates of laboratory fitness also showed clear signs of temperature-specific adaptation,
with Hot lines outperforming Cold lines at 35°C, while there were only small differences at
23°C (Supplementary Figure S4). In addition, for metabolic there was an interaction between
evolution regime, assay temperature and geographic origin, indicating that thermal
adaptation was contingent on founding genetic variation (Fig. 4B, E, Supplementary 3).
Analysing all traits together using a non-parametric MANOVA[76] confirmed main effects of
assay temperature (F = 205.5, P < 0.001), evolution regime (F = 57.1, P < 0.001) and the
interaction between evolution regime and assay temperature (F = 6.8, P < 0.001).
296
297 *Thermal adaptation in life-history shapes the agricultural footprint*

To predict how the observed life-history evolution may affect agricultural yields, we estimated the relationship between life-history traits and the amount of host consumption by *C. maculatus.* We reared Cold, Hot and Ancestral lines at 23, 29 and 35°C. At each assay temperature, we let females lay eggs in a petri-dish containing ad libitum black-eyed beans. The number of eggs laid, the number of individuals surviving to pupation, the mass of adult beetles, and the seed mass consumed, was calculated from each dish.

The Hot regime consumed significantly more of the host than the Cold regime ($F_{1, 5.97} = 22.8$, $P = 0.003$), demonstrating that the agricultural footprint indeed had evolved. The difference between the Hot and Cold regime tended to be greater at 35°C, although the two-way interaction between regime and assay temperature was marginally non-significant ($F_{2, 11.9} = 3.25$, $P = 0.075$, Fig. 5A). To assess the importance of life-history traits in affecting food consumption, we performed analyses while sequentially including information on the number of eggs laid, juvenile survival, and body mass so that the final model predicted the amount of host consumed as a function of the cumulative beetle mass produced. Adding life-history traits to the models reduced the differences between evolution regimes and in the final analysis including all traits we no longer found a significant effect of regime ($F_{1, 5.73} = 0.60$, $P = 0.47$), demonstrating that life-history adaptation alone seems responsible for the evolved differences in host consumption. Indeed, a model containing only the main effects of eggs laid, adult body mass and juvenile survival to pupation explained 93% ($P < 0.001$) of all the observed variance in bean consumption across the entire experiment (Fig. 5A). The amount of host mass lost was 3-4 times greater than the mass of beetle produced, which is similar to what has been reported in a previous study[77], underlining the severe impact that *C. maculatus* can have on host crops (Supplementary 5).

We predicted the agricultural impact of *C. maculatus* at cold and hot temperature based on the observed life-history evolution by calculating the agricultural footprint as: $\varphi1 = C_T\omega M/d_1$, where C_T is the temperature-specific constant converting beetle mass produced to host mass consumed (Fig. 5A), ω is lifetime offspring production of a single female (Fig. 4A), M is body mass at adult emergence (Fig. 4B) and d_1 is mean development time in days (Fig. 4C). Hence, this measure of the footprint captures the magnitude of agricultural loss per time unit as a function of an organism's life-history strategy[78]. To assess the sensitivity of our predictions to both estimation error and choice of life-history variables we calculated a second independent estimate of the footprint, based on measurements of a different set of individuals: $\varphi2 = \psi C_E/d_2$, where ψ is the number of eggs laid by a single female over her first 16h of reproduction (Fig. 4D), C_E is the amount of host consumed per laid egg, and d_2 is the mean development time in days of the fastest developing individual in each petri-dish (both variables estimated from the host consumption experiment presented in Fig. 5A). These two metrics were highly correlated ($r = 0.88$, $t_{31} = 10.4$, $P < 0.001$, Fig. 5B) suggesting high predictive accuracy.

To provide a generalisable metric of the agricultural impact of life-history evolution under climate change, we scaled the predicted footprint for all lines relative to their respective ancestor reared at 29°C (Fig. 5C). This metric allowed us to assess both the direct effect of temperature (i.e. thermal plasticity) on the footprint in the ancestors, and how evolution, relative to this direct effect of temperature, impacted the footprint. Cold (23°C) and hot (35°C) assay temperature reduced the footprint by 35-45% and 0-25% respectively, relative to benign 29°C in the ancestors (Fig. 5C). However, evolution had a fundamental impact on how temperature affected the footprint. While cold-adapted lines showed very little difference to ancestral lines at 23°C (Δfootprint = 7.6%; CI: 2.0 – 15%), Hot-adapted lines left almost twice as large a footprint as the ancestral lines at 35°C (Δfootprint = 80%; CI: 70 – 97%). Moreover, these effects were contingent on the geographic origin of the founders. At 23°C, cold-adapted lines from California left a slightly larger footprint compared to the ancestor (Δfootprint = 29%; CI: 13 – 42%) while the footprint had not evolved on the Brazil (Δfootprint = -3.1%; CI: -11 – 6.3%) and Yemen (Δfootprint = 1.0%; CI: -11 – 11%) background. At

35°C, the footprint evolved substantially on all backgrounds, but the increase relative to the
ancestor tended to be greater for Brazil (Δfootprint = 84%; CI: 64 – 110%) and California
(Δfootprint = 93%; CI: 71 – 120%) compared to Yemen (Δfootprint = 62%; CI: 44 – 87%)
(Fig. 5C).

These results indicate that life-history evolution can cause significant changes to the
agricultural impact of insect pests over a relatively short period of time. Due to the
thermodynamic effects on generation time[56], our evolved regimes had undergone different
numbers of generations at the time of assaying (60 for hot vs. 45 generations for cold lines
for traits reported in Fig. 4, after 5 years of evolution). Reported effects should correspond
well with predictions for populations having continuous generations adapting at cooling and
warming temperatures, respectively. However, for seasonal populations that are restricted in
the number of generations per year by other ecological factors than temperature, per-
generation estimates may be more relevant. Assuming that trait-evolution proceeded at a
constant (linear) pace in our experiment, the average change in the agricultural footprint due
to adaptation at hot temperature (relative to ancestral populations) would correspond to a
1.3% (1.2-1.6) increase per generation, with the estimate for cold adaptation being 0.2%
(0.05-0.35).

Discussion
Climate change is predicted to fundamentally impact growth rates and distribution ranges of
ectotherms[5,30,31,42,79–81], many of which are pests on economically important crops[1,2,7,16].
Current predictions of increased agricultural impacts of pests typically rely on deterministic
statistical projections of how temperature affects ectotherm metabolism and population
growth, but do not incorporate evolution[13,14,78,82]. However, rapid evolutionary responses are
characteristic of many pest species[1,5,7,16]. Here we therefore asked to what extent the
incorporation of evolution can alter current projections of climate-induced changes in pest
impact. We show three features that emphasize that future projections need to account for
thermal adaptation in pest species' life-history. First, adaptation affected the agricultural
footprint left by *C. maculatus,* but the effect was dependent on temperature; with little or no
effect of cold adaptation while hot-adapted populations caused an almost doubling of the
footprint relative to ancestors (Fig. 5). Second, these effects depended on the ancestral
founder of the adapting populations, suggesting that geographic differences in segregating
genetic variation in life-history traits may influence future agricultural loss under climate
warming. Third, attempts to incorporate niche evolution in species' distribution models may
need to do so by relatively simple means in the absence of empirical data that allow
parametrisation and evaluation of more complex mechanistic models[13]. However, our study
echoes concerns that such predictions may be off the mark because the mechanisms and
type of traits that mediate niche evolution have a strong influence on pest impact[14,17,40,83].

Indeed, the observed temperature-dependent effect of evolution on the agricultural footprint
can be understood by considering trait-specific responses. Even though cold-adapted
populations did show evolved responses in traits such as weight loss and water loss, these
traits are predicted to have small to no effect on the agricultural footprint. Cold populations
also showed genetic responses in development rate and metabolism (the latter dependent
on genetic background) that could impact agricultural loss[1,2], but these genetic changes were
very modest compared to the responses observed for lifetime reproductive success, early
fecundity and adult body mass, that evolved more readily in hot-adapted populations (Fig. 4).
The responses of hot-adapted populations are in line with two predictions from biophysical
theory. First, warmer temperatures are predicted to ease thermodynamic constraints on
metabolic rate[60], which can benefit larger size via stronger selection on fecundity[84–86].
Second, evolution at stressful hot temperature may favour compensatory feeding and
sequestering of resources devoted to maintaining cellular homeostasis[5,19,20,36,37,49,70] (Fig. 1,
Supplementary 1). Indeed, increased larval feeding and larger adult body mass in *C.*

maculatus is associated with extended adult lifespan and greater fecundity under aphagous conditions corresponding to laboratory and grain storage environments (e.g.[87,88]). These effects are likely magnified at hot temperatures that increase metabolism and water loss ([23,89] and Fig. 4), and the need for upregulation of molecular chaperones to secure cellular homeostasis ([20,68,69] and Fig. 3). In accordance with this hypothesis, previous studies on *C. maculatus* also found that evolution at hot temperature increased body mass[90,91], an adaptive genetic response that paralleled the observed thermal plasticity in body mass[90]. Indeed, also for our populations used here, even though body size is largest when beetles are raised at 23°C, beetles reared at 35°C are larger than when raised at benign 29°C (Supplementary Figure S5). To further evaluate the hypothesis that larger sizes is beneficial at hot temperature in *C. maculatus*, we used the estimates of female life-history to calculate selection differentials on body size at hot and cold assay temperature. For all three estimates, selection on size was stronger at the hot temperature (Supplementary 6). Thus, the evolved increase in the agricultural footprint at hot temperature observed here is likely mediated by two synergistic effects of life-history adaptation, *i)* increased larval feeding enabling larger size at maturity, and *ii)* the positive effects of a large adult size on survival and reproduction at hot and aphagous conditions.

These observations agree broadly with the model prediction that stressful temperature can favour increased host consumption and energy allocation to maintenance to mitigate thermal constraints on growth and reproduction (Fig. 1, Supplementary 1). Note, however, that this prediction does not necessarily equate to the evolution of a large size at maturity, as resources might be spent on higher metabolic expenditure and maintenance at warm temperature prior to the organism reaching adult maturity, and our simple model makes no explicit predictions in this regard (see also[92,93]). Moreover, increased energy intake necessitates access to host plants. If food availability becomes limiting under climate warming, this mechanism could instead severely limit thermal tolerance and population growth rates of heat exposed ectotherms[37,94]. Thus, while host abundance is already acknowledged as a key factor regulating pest outbreaks[1,7,78,95], the interdependency between heat stress and the benefits of energy acquisition may magnify the effects of host accessibility on pest outbreaks under future climate warming (Fig. 1C, D). Manipulating temperature and larval densities simultaneously and examining the outcome for both population growth and host consumption could test this prediction explicitly and further evaluate our model's predictions.

Early findings suggested that most ectotherms follow the "temperature-size rule"[96,97], stating that body size shows plastic reductions with increasing temperature. However, later results have shown that these patterns are far from general. In particular, terrestrial ectotherms, the group to which many insect pests belong, show no consistent plastic response to temperature, and genetic adaptation along latitude shows a trend towards larger genetic body sizes at warmer latitudes[98], which is consistent with the evolved response observed here. Moreover, thermal responses in body size tend to vary even within and between closely related species[81], as for *C. maculatus*[90,99] and other Chrysomelid beetles[100]. Understanding whether taxonomic variation represent constraints rooted in thermodynamics or adaptive plasticity may offer insights into the future routes of evolution under climate warming but has proven a major challenge for the field of eco-physiological adaptation[5,19,81,92,97]. Moreover, as discussed above, the effect of heat on energy expenditure suggests that the temperature scaling of adult body mass may not follow a one-to-one relationship with that of host consumption. Controlled experimental approaches as used here may provide further clues to what extent patterns of thermal plasticity are indicative of future genetic responses to climate and how the evolution of classic life-history traits affects host consumption. Such general information could be used to increase accuracy of future projections for species where thermal plasticity can be accessed, but where evolution is hard to study directly.

In contrast to the measured life-history traits, feeding efficiency was not strongly affected by assay temperature and did not evolve in our experiment, suggesting that differences in the agricultural footprint of insect pests may be accurately predicted from life-history variables alone. Yet, our study also highlights the important role of founding genetic variation in dictating life-history adaptation. This suggests that evolution under climate change may proceed along different trajectories across a species distribution range, even when regional climates change in parallel. When such divergent evolutionary responses involve key life-history traits, as observed here, this could lead to regional differences in pest impact[1,2,83]. In addition, while our assays of offspring production and host consumption capture variation stemming from temperature effects on larval survival and male fertility, our study is focused on female life-history traits. Future studies might also want to consider the role of male life-history, fertility, and male-female mating interactions, which can be affected by temperature and have strong effects on population regulation[29,101–106], as also shown for *C. maculatus*[107–110].

The impact of insect pests on agriculture is paramount[11,78], with severe repercussions in human societies; 2.6 billion people (26% of the world's population) are currently directly dependent on agriculture for a secure living[111]. In African countries alone, the financial loss from pests was estimated to at least 1 billion USD for 2019, with agriculture accounting for 99% of this loss in central regions[9,10]. Clearly, the redistribution of species has already caused massive ramifications for the Earth's ecosystems[112], and the fear is now that future climate warming will allow insect pests to colonize new habitats[1,2,4,7,11,12]. To mitigate these effects, management strategies need to be equipped with accurate predictive models[13,14,82,113–115], as well as an understanding of the limits to their accuracy[43,116–118]. It is our hope that our study can contribute towards this development by *i)* providing general insights into the repeatability and mode of thermal adaptation in life-history of insect pests, which can inform on the general accuracy of ecological forecasting of species responses to climate change, and *ii)* by emphasizing the need for incorporating a mechanistic and genetic basis of thermal niche evolution into eco-evolutionary forecasts, on which improved management decision can be based[6,13,17,82,83,113,119,120].

Methods

Experimental evolution

C. maculatus is a pest on legume seeds and is common in tropical and sub-tropical regions[121]. Females lay eggs on the surface of seeds. After the egg hatches, the larva burrows into the seed to feed until it pupates. Adults emerge from seeds after ca. three weeks at standard temperatures around 29°C and are reproductively mature almost instantly and do not require food or water to reproduce[74,75,90]. These characteristics have likely contributed to *C. maculatus* being a cosmopolitan pest that invades seed storages[12,50]. Indeed, *C. maculatus* has become a model organism for research in both evolutionary ecology and pest management because laboratory and grain storage environments are very similar.

The experimental evolution protocol exposed six lines to 35°C (*Hot* regime) and six lines to 23°C (*Cold* regime), for 85 and 130 generations, respectively (difference due to effects of temperature on generation time). The lines were derived from three ancestors that were collected from different geographic regions (Brazil, Yemen and California (USA)), with two replicate lines per thermal regime created per region. Each line was created by mating 150 females with 150 males from the respective ancestral founding population. Population size was immediately expanded to ~3000 beetles in the next generation and beetles were split in groups of 600 to create four population replicates; two placed at hot, and two placed at cold

evolution temperature per ancestor[75]. The three founders were kept at ancestral 29°C
throughout experimental evolution (Fig. 2A). A humidity of 50-55% was used throughout this
study. Ancestors were reared, both before and during experimental evolution, by placing
300-400 newly emerged adults on 150 ml black-eyed beans, whereas evolution lines were
reared by placing 600 adults on 250 ml black eyed beans, resulting in similar densities of
adult beetles per bean. To reduce direct selection on development time, all lines were
transferred to a new jar with fresh beans around the peak of emergence of adult beetles at
each generation. This occurred approximately three days after the first beetle emerged for
ancestors and the Hot regime, and five days for the Cold regime.

526
Gene expression
To provide evidence for a temperature-dependent trade-off between reproduction and
maintenance, we leveraged transcriptomic data from three separate experiments. The first
two experiments estimated gene expression responses in female abdomens to heat shock
and mating, respectively, in a separate lab population collected from Lomé, Togo[122]. The
third dataset estimated expression in female abdomens in response to rearing temperature in
our experimental evolution lines. All beetles were reared at standard laboratory conditions of
29°C and 55% relative humidity unless otherwise stated.

535
To identify heat stress responsive genes, we compared reproductively active control and
heat shocked females. Individuals were collected as virgins and kept individually. On the next
day, females were heat shocked in individual 60 mm petri dishes at 55° C for 20 minutes,
which is stressful and reduces female fertility while still representing a thermal stress
occurring at the sampling site of the population[108,109]. After a 2.5 h recovery at 29° C and 55%
rh, all females were mated to males. Females were then kept in individual 60 mm dishes until
being flash frozen in liquid nitrogen 2-3 h, 6-7 h or 24-25 h after mating. The experiment was
repeated on three consecutive days (blocks). Each replicate sample consisted of 5 pooled
female abdomens from the same treatment, time-point and block. For sequencing, we
selected a total of seven replicate samples per time-point for the control treatment and five
replicate samples per time-point for the heat shock treatment. Preliminary analyses showed
that there were only three differentially expressed genes at the last time-point, which
therefore was removed in the final analyses (although results did not qualitatively change if
including them), resulting in 24 RNA libraries for analysis.

550
To identify reproductive genes, we compared virgin and mated females 24 h after a single
mating. We mated females to males from 8 different genetic lines. The lines had undergone
53 generations of experimental evolution under one of three mating regimes manipulating the
relative strength of natural and sexual selection[109]. Accordingly, we had 8 samples of mated
females, with 2 or 3 samples per male mating regime. Additionally, we had 3 samples of
virgin control females. Flash freezing of all females occurred 24 h after mating, with females
kept singly and allowed to lay eggs on fresh beans during the 24h period. The final 11 RNA
libraries each consisted of 12 female abdomens.

559
To quantify expression of reproductive and heat stress responsive genes (identified in the
first two datasets) in our experimental evolution lines and their ancestors, we reared all lines
at 23, 29 and 35°C. All lines were first propagated for one generation at 29°C to remove
potential difference between evolution regimes stemming from temperature-induced (non-
genetic) parental effects. Once adults emerged at respective rearing temperature, 5 males
and 5 females were placed together in a 90mm petri dish with access to fresh beans. Beetles
were allowed to reproduce for 23h (for 29 and 35°C) or 46h (for 23°C) after which females
were flash frozen. The longer time allowed for reproduction at cold assay temperature was
given to make a more direct comparison across females from different temperatures;
reproductive output (Fig. 4B), metabolism (Fig. 4E) and weight loss (Fig. 4F) is about twice
as fast at 35°C compared to 23°C. Two petri dishes of females were pooled into one sample

(i.e. 10 female abdomens pooled in total) for extractions and further analysis, resulting in one
sample per line and temperature for the Hot and Cold regime. Two samples were taken per
temperature for each of the three ancestors. This resulted in a total of 54 RNA libraries.
574
RNA from all samples was extracted using the Qiagen RNeasy Mini kit with beta-mercapto-
Ethanol added to the lysis buffer and an on-column DNase treatment with the Qiagen
RNase-free DNase kit. Tissue lysis was done in a bead mill with two stainless steel beads at
28 Hz for 90 s and RNA was eluted in two times 30-50 µl water. Some samples went through
an additional clean-up step to increase RNA concentrations with the Qiagen RNeasy Mini Kit
after the RNA extraction. Next generation sequencing of the samples was done at the
SNP&SEQ Technology Platform in Uppsala. Libraries were prepared using the TruSeq
stranded mRNA library preparation kit with polyA selection. Libraries were sequenced in one
flowcell as paired-end 150 bp reads on a NovaSeq 6000 system.
584
Raw reads were quality checked with FastQC 0.11.9[123] and MultiQC 1.11[124] and afterwards
cleaned up using Trimmomatic 0.39[125]. Adapters were trimmed from the raw reads and ends
below an average accuracy of 99% over a five base pair sliding window were clipped. Finally,
reads shorter than 20 bp were discarded. Resulting read pairs were mapped to the
Callosobruchus maculatus genome[126] using HISAT2 2.2.1[127] with default settings for
stranded libraries and sensitivity set to "--very-sensitive". Number of reads per gene was then
determined using HTSeq 2.0.2[128] with default settings for stranded libraries. This resulted in
10-27M (heat stress), 13-30M (reproduction) and 9-35M (rearing temperature during
experimental evolution) uniquely mapped reads per library going into the final downstream
analyses.
595
Read count data was pre-processed and analysed in R 4.3.1[129] using edgeR[130,131]. For the
heat shock response, only genes with at least 3 counts per million in at least 5 samples were
further analysed, resulting in 10246 genes for final analysis. For the mating response, only
genes with at least 1 count per million in at least 2 samples were further analysed, resulting
in 11640 genes for final analysis. The last approach was also used for the genes expressed
in the evolution regimes, resulting in 11292 genes for analysis. Counts were normalized
using the 'Trimmed Mean of M-values' method via edgeR's NormLibSizes() function. Count
data for the heat shock and mating response was further analysed with edgeR[130,131] and
limma[132] using linear models on the normalized log2 transformed counts per million to
determine differentially expressed genes. All analyses used a false discovery cut-off of 5 %.
606
For reproductive genes, effect sizes (i.e. log-fold changes) were averaged across the three
male selection regimes. However, we only considered genes that were significantly
differentially expressed in all three separate contrasts between virgin females and females
mated to males from the three mating regimes. For heat stress genes, effect sizes and
significance was calculated by averaging across the two time-points. Significant genes were
further subject to gene ontology analysis, using the R package GOstats (2.66.0)[133] and the
HyperGTest function. Results were visualised using clusterProfiler (4.10)[134].
614
To provide a powerful, yet, unbiased estimate of temperature-dependent allocation between
reproduction (mating response) and maintenance (heat stress response) in the experimental
evolution lines, we focused on the 134 genes that displayed significant antagonistic
differential expression patterns in response to heat shock and mating (Fig. 3B). We first
multiplied the estimated log fold changes of the antagonistic genes for both mating and heat
shock responses by the vector of normalized read counts of the same 134 antagonistic
genes found for each of the 54 RNA libraries from the experimental evolution lines. The
resulting values were then summed to compute a score for the heat stress and reproduction
response separately. As expected, the scores for heat stress and reproduction, based on
only the antagonistic genes, were tightly negatively correlated ($r = -0.999$) across all
combinations of evolved lines and assay temperatures and fell along a singular linear trade-

off axis (Supplementary Figure S3). The response along the allocation trade-off for each
sample was taken by projecting samples along this axis (quantified as the first eigenvector of
the data). We repeated the same analyses using all differentially expressed genes in
response to heat shock (765) and mating (1269) with qualitatively similar results.
630
Life-history traits
We quantified thermal adaptation in female life-history by measuring three core traits: lifetime
reproductive success (LRS), juvenile development time and adult body mass, and four rate-
dependent traits: early fecundity, weight loss, water loss, and metabolic rate over the first
16h of female reproduction. All life-history traits were collected at generations 40 for cold-
adapted lines and 60 for hot-adapted lines, in a large common garden experiment including
the two assay temperatures corresponding to the experimental evolution treatments (23°C
and 35°C). Ancestral lines were scored in the same experimental conditions with the addition
of the ancestral 29°C assay temperature, but on a later occasion following ca. 125
generations of experimental evolution. Note that the ancestors had been kept at the
ancestral conditions to which they had already adapted for more than 300 generations prior
to the start of experimental evolution. It can therefore be assumed that the measured trait
values will not have changed much due to selection during experimental evolution. Further,
while we cannot completely exclude a role of genetic drift in the measured life history traits,
the censuses size of 300-400 adult beetles in the ancestors corresponds to an effective
population size of roughly 200[107], and the measured traits are likely under relatively strong
(stabilizing) selection. Hence, we expect any effect of drift to be small and that the measured
trait values should correspond reasonably well with those at the start of experimental
evolution. To control for potential differences in the separate experiments on ancestors and
evolved lines stemming from unknown sources, we reared an independent laboratory
adapted reference population in both experiments. This indicated that differences in rearing
had affected the life-history traits scored over the first 16h of reproduction. We therefore
standardized the traits scored during respirometry of the three founding ancestors by this
estimated amount (adult mass: increased by 6.4%, metabolic rate: reduced by 12%; early
fecundity: reduced by 18%; water loss: reduced by 15%, and weight loss: reduced by 25%)
in order not to erroneously assign these differences to evolutionary divergence between
ancestors and evolved lines. Note that this was done averaged across the three assay
temperatures and geographic origins. Therefore, our approach to provide more accurate
measures of evolutionary divergence between evolved lines and ancestors did not affect the
estimated temperature-dependence of adaptation or the importance of geographic
differences.
662
Before assays of life-history traits, non-genetic parental effects were removed by moving F0
grandparents of the assayed individuals into a common temperature of 29°C to lay eggs. The
emerging beetles in the next (parental) F1 generation were allowed to mate and lay eggs on
beans provided *ad libitum*. Following 48 hours of egg laying, the beans were split and
assigned to one of the two (for ancestors, three) assay temperatures. The emerging adult F2
offspring were phenotyped for their life-history (Fig. 4). Newly emerged (0-48 hours old)
virgin females were mated to males by placing three males and females together in a petri
dish over night at the assay temperature. In the following morning, the three females were
weighed for their body mass and then placed together inside a glass vial filled with black
eyed beans (ca. 20-25) to be measured for their metabolic rate, water loss and early
fecundity at their designated assay temperature. We note here that our goal was to measure
metabolism in conditions that correspond to the most common state of adult seed beetles
and most relevant ecological setting to compare the evolved populations. *C. maculatus* mate
on the first day of eclosion, and females from our lines start laying eggs almost immediately
at high adult densities. Further, early laid eggs are likely to be more important for individual
fitness than eggs laid at older age due to the larval competition that subsequently occurs
among offspring inside seeds. We thus measured metabolic rate during the first day of mated
adult life for female trios. We nevertheless note that our measures of metabolism may not

directly correlate with commonly used measures of resting metabolic rate from single
individuals kept in isolation.
683
The glass vials (diameter = 2 cm, length = 4 cm) were integrated into a high-throughput
respirometry system and housed inside a Sanyo MIR-153 incubator (set to an assay
temperature of either 23, 29 or 35°C, with lights turned on). Briefly, the respirometry was set
up in stop-flow mode, with incoming air controlled at a flow rate of 50ml/min by an SS-4
pump (Sable Systems, Las Vegas, NV, USA) and a Model 840 mass control valve (Sierra
Instruments, Monterey, CA). CO^2 production and water-loss were measured for up to 23 vials
on a given experimental day using a LiCor 7000 infrared CO_2 gas-analyser (Lincoln, NE) and
a RH-300 water vapor pressure meter (Sable Systems), respectively. The system was
connected via three RM-8 (eight-channel) multiplexors (Sable Systems). The first vial was
left empty and served as a baseline to control for any drift of the gas analysers during each
session and was measured at both the start and end of each cycle. Vials were measured
over 17 cycles, each of a length of 57.5 minutes. The LiCor gas analyser was span calibrated
using 1000ppm NO_2 and CO_2 in the morning before each trial began. Mean metabolic rate
and water loss for each vial was calculated across cycles 2-17, with the readings from the
first cycle discarded (as it contains human-produced water and CO_2). Respirometry data was
retrieved using ExpeData Pro v1.5.6 (Sable Systems). After respirometry, females were
weighed again to record their weight loss and beans with eggs were isolated and counted to
record early fecundity. In total we followed 386 triads of females for the evolved lines and
another 115 triads from their ancestors.
703
From the same rearing we measured egg-to-adult development time for two technical
replicates per line and assay temperature, each consisting of 40-120 individuals. We
calculated a mean development time per technical replicate and used this in analysis. We
also collected virgin males and females and placed three males and three females together
in a petri dish with ad libitum beans to record lifetime reproductive output (LRS) at each
assay temperature. In total we recorded LRS for 258 couple triplets for evolved lines, and
another 115 couple triplets for the ancestors. These data were complemented with additional
data from both evolved and ancestral lines reared in a common garden design in two
consecutive years (corresponding to generation 120/135 for ancestors, 115/130 for hot-
adapted lines, and 80/90 for cold-adapted lines). In these rearings, a single male and female
were put together in a petri dish with ad libitum host seeds, with otherwise the same
conditions. For ancestors we scored 396 couples, and for evolved lines 789 couples, across
both experimental years. LRS was analysed per female, hence we divided all offspring
counts from female triads by three before analysis.
718
We carried out all statistical analyses using the statistical and programming software R (v.
3.6.1). All seven traits were first analysed separately using linear mixed effect models
available in the lme4 package[135]. When analysing differences between hot- and cold-adapted
lines, evolution regime, geographic origin and assay temperature were added as fully
crossed fixed effects. Line replicate crossed with assay temperature and nested within
geographic origin were added as random terms to assure correct level of replication when
estimating significance of fixed terms including evolution regime. For analyses on metabolic
rate, water loss and weight loss, the date of the respirometry run and the measurement vial
ID were added as additional random effects. Metabolic rate and weight loss were corrected
for body mass of the measured female triad by taking an average of the weight measures
before and after respirometry and adding it as a covariate in analysis. We first fitted individual
slopes for the logged weight covariate (with logged CO_2 as response) for the hot and cold
evolution regime. As these estimates were very similar (slope cold = 0.95, slope hot = 1.0)
and not different statistically (P = 0.7), we estimated a shared slope across evolution regimes
for the final model. This slope was equal to 0.98 and not significantly different from 1, as
seen for other studies on *C. maculatus*[87,136,137]. We therefore chose to graphically present
mass-specific metabolic rate as produced CO_2 in ml per minute and milligram of beetle (Fig.

4E). For water loss measured from evolved lines, measures from vial 2 (the first vial in the measuring sequence containing beetles) and measurements from three entire experimental days were discarded following an outlier analysis as these measures were magnitudes greater than other measures and clearly represented water vapour from other sources than the beetles. For LRS, experiment (triads and two separate rearings of single couples) and its interaction with assay temperature were added as additional fixed effects to account for possible block effects. The data from ancestral lines were analysed in similar fashion but without the fixed effect of evolution regime and random effect of population replicate.

P-values were calculated using the car package and type II sums of squares with the Kenward-Roger approximation for the degrees of freedom. To analyse evolutionary responses while taking all traits into consideration at once, we also performed a non-parametric MANOVA using residual randomization[76] with the same structure for fixed effects but where observations were based on line means, as all traits were not recorded in the same individuals (Fig. 4H).

The agricultural footprint
We set up both evolved and ancestral lines in a common garden experiment including the 23, 29 and 35°C assay temperature, following approximately 85, 120 and 125 generations of experimental evolution for cold-adapted, hot-adapted and ancestral lines, respectively. We removed parental effects by moving all lines to 29°C two generations prior to the start of the experiment. Newly laid F2 eggs were split among the three assay temperatures and resulting virgin adults were collected. Three males and three females were placed together in petri-dishes with ad libitum *V. unguiculata* seeds for 24h at 29°C and 35°C, and 34h at 23°C (as reproductive rate is slower at cold temperature; Fig. 4B). Three dishes were prepared per line and assay temperature. Note that the experiment was aimed at linking life-history variation to host consumption, and the sample size was low for detecting more fine-scaled differences between evolution regimes due to the imprecise measures of fecundity (6-9 females per line and temperature measured over 24-34h), which were much better estimated in the previous experiments (~80 females per line and temperature across the entire adult stage).

To quantify the amount of host seed consumed by beetles, each petri dish with beans was weighed prior to the addition of beetles, to record a starting weight of each assay. After beetles had emerged and been removed, assays were placed in standard conditions for 6 weeks during which they were weighed on several occasions to record the weight of the infested beans and to check for potential inconsistencies and time-dependencies in estimates of bean consumption. However, the measures of the beans' weight loss following infestation were highly repeatable ($r^2 = 94\%$) and we chose the final (fourth) measure to calculate our estimate of host consumption. To account for potential weight loss of beans that was independent of the beetle infestation, we also set up 5 assays without beetles at each temperature and measured them in the exact same way. The host consumption was then corrected based on the weight changes in these control assays; this correction was very small and only corresponded to 1-2% of the total bean weight. For each assay, we also recorded the number of hatched eggs laid on beans (i.e. fecundity), the number of those that resulted in the adult pupating (i.e. juvenile survival), and the mean body mass of emerged adult beetles.

Differences in food consumption were first analyzed using linear mixed effect models in the lme4 package with the same structure for fixed and random terms as described for life-history traits. We excluded the ancestors from formal analysis since these had only half the samples compared to the evolution regimes, but we included them in Fig. 5A to show the direction of evolution of host consumption. We sequentially added information on fecundity, juvenile survival, and mean body mass for each assay to explore how life-history traits accounted for host consumption. We finally fitted a linear model including only the three traits

791 (removing the fixed terms of assay temperature, evolution regime and geographic
792 background) to estimate how much of the variation in host consumption that could be
793 predicted by life-history alone.
794
Motivated by the observation that the life-history traits accounted for more than 93% of the
observed variation in host consumption (see Fig. 5A and[77]), we calculated the two
independent measures of the agricultural footprint (see main text) from our measured life-
history traits in the main experiment, as this data was estimated with high accuracy for all
lines. To provide 95% confidence limits and P-values, we performed parametric bootstrap on
all calculations using the means and standard errors for each trait per line and assay
temperature, derived from the univariate mixed models (Fig. 4A-G, Supplementary 3).
802
803
804
805

Data Availability
807 All data on life-history traits and host consumption, along with R code for all analyses, are
808 available at Figshare [https://doi.org/10.6084/m9.figshare.26048389].
809
The gene expression data is under embargo at:
Mating response:
https://dataview.ncbi.nlm.nih.gov/object/PRJNA1119937
Heat stress:
https://dataview.ncbi.nlm.nih.gov/object/PRJNA1120212
Evolution and Thermal Plasticity:
https://dataview.ncbi.nlm.nih.gov/object/PRJNA1121056
817
818

Code Availability
820 The R code underlying the model predictions in Figure 1 and Supplementary 1 is available as
821 supplementary material.
822
823

References

1.   Lehmann, P. *et al.* Complex responses of global insect pests to climate warming. *Front. Ecol.*
Environ. **18**, 141–150 (2020).
2.   Deutsch, C. A. *et al.* Increase in crop losses to insect pests in a warming climate. *Science* **361**,
916–919 (2018).
3.   Harvey, J. A. *et al.* Scientists' warning on climate change and insects. *Ecol. Monogr.* **93**, e1553
(2023).
4.   Daly, E. Z. *et al.* A synthesis of biological invasion hypotheses associated with the introduction–
naturalisation–invasion continuum. *Oikos* **n/a**, e09645.
5.   González-Tokman, D. *et al.* Insect responses to heat: physiological mechanisms, evolution and
ecological implications in a warming world. *Biol. Rev.* **95**, 802–821 (2020).
6.   Diamond, S. E. Contemporary climate-driven range shifts: Putting evolution back on the table.
Funct. Ecol. **32**, 1652–1665 (2018).
7.   Harvey, J. A., Heinen, R., Gols, R. & Thakur, M. P. Climate change-mediated temperature
extremes and insects: From outbreaks to breakdowns. *Glob. Change Biol.* **26**, 6685–6701 (2020).
8.   Simberloff, D. *et al.* Impacts of biological invasions: what's what and the way forward. *Trends*
Ecol. Evol. **28**, 58–66 (2013).

9.   Diagne, C. *et al.* High and rising economic costs of biological invasions worldwide. *Nature* **592**,
571–576 (2021).
10. Diagne, C. *et al.* The economic costs of biological invasions in Africa: a growing but neglected
threat? *NeoBiota* **67**, 11–51 (2021).
11. Change, C. R. P. on C. & Security, A. and F. *CGIAR Research Program on Climate Change,*
Agriculture and Food Security Annual Report 2020.
https://cgspace.cgiar.org/handle/10568/114818 (2021).
12. Adler, C. *et al.* Changes in the distribution and pest risk of stored product insects in Europe due
to global warming: Need for pan-European pest monitoring and improved food-safety. *J. Stored*
Prod. Res. **97**, 101977 (2022).
13. Urban, M. C. *et al.* Improving the forecast for biodiversity under climate change. *Science* **353**,
aad8466 (2016).
14. Macfadyen, S., McDonald, G. & Hill, M. P. From species distributions to climate change
adaptation: Knowledge gaps in managing invertebrate pests in broad-acre grain crops. *Agric.*
Ecosyst. Environ. **253**, 208–219 (2018).
15. Galante, P. J. *et al.* The challenge of modeling niches and distributions for data-poor species: a
comprehensive approach to model complexity. *Ecography* **41**, 726–736 (2018).
16. Hoffmann, A. A. Rapid adaptation of invertebrate pests to climatic stress? *Curr. Opin. Insect Sci.*
21, 7–13 (2017).
17. Helmuth, B., Kingsolver, J. G. & Carrington, E. BIOPHYSICS, PHYSIOLOGICAL ECOLOGY, AND
CLIMATE CHANGE: Does Mechanism Matter? *Annu. Rev. Physiol.* **67**, 177–201 (2005).
18. Huey, R. B. & Kingsolver, J. G. Evolution of thermal sensitivity of ectotherm performance. *Trends*
Ecol. Evol. **4**, 131–135 (1989).
19. Angilletta, M. J. *Thermal Adaptation: A Theoretical and Empirical Synthesis.* (Oxford University
Press, Oxford⁠; New York, 2009).
20. Hochachka, P. W. & Somero, G. N. *Biochemical Adaptation: Mechanism and Process in*
Physiological Evolution. (Oxford University Press, 2002).
21. Berger, D., Stångberg, J., Baur, J. & Walters, R. J. Elevated temperature increases genome-wide
selection on de novo mutations. *Proc. R. Soc. B Biol. Sci.* **288**, 20203094 (2021).
22. Kellermann, V., van Heerwaarden, B., Sgrò, C. M. & Hoffmann, A. A. Fundamental evolutionary
limits in ecological traits drive Drosophila species distributions. *Science* **325**, 1244–1246 (2009).
23. Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M. & West, G. B. Toward a Metabolic Theory of
Ecology. *Ecology* **85**, 1771–1789 (2004).
24. Savage, V. M., Gillooly, J. F., Brown, J. H., West, G. B. & Charnov, E. L. Effects of Body Size and
Temperature on Population Growth. *Am. Nat.* **163**, 429–441 (2004).
25. Bloom, J. D. *et al.* Thermodynamic prediction of protein neutrality. *Proc. Natl. Acad. Sci. U. S. A.*
102, 606–611 (2005).
26. Agozzino, L & Dill, K. A. Protein evolution speed depends on its stability and abundance and on
chaperone concentrations. *Proc. Natl. Acad. Sci.* **115**, 9092–9097 (2018).
27. Chen, P. & Shakhnovich, E. I. Thermal adaptation of viruses and bacteria. *Biophys. J.* **98**, 1109–
1118 (2010).
28. Parratt, S. R. *et al.* Temperatures that sterilize males better match global species distributions
than lethal temperatures. *Nat. Clim. Change* 1–4 (2021) doi:10.1038/s41558-021-01047-0.
29. Walsh, B. S. *et al.* The Impact of Climate Change on Fertility. *Trends Ecol. Evol.* **34**, 249–259
(2019).
30. Jørgensen, L. B., Ørsted, M., Malte, H., Wang, T. & Overgaard, J. Extreme escalation of heat
failure rates in ectotherms with global warming. *Nature* **611**, 93–98 (2022).
31. Deutsch, C. A. *et al.* Impacts of climate warming on terrestrial ectotherms across latitude. *Proc.*
Natl. Acad. Sci. **105**, 6668–6672 (2008).
32. Levins, R. *Evolution in Changing Environments: Some Theoretical Explorations.* (Princeton
University Press, 1968).

33.   Kassen, R. The experimental evolution of specialists, generalists, and the maintenance of
diversity: Experimental evolution in variable environments. *J. Evol. Biol.* **15**, 173–190 (2002).

34.   Conover, D. O. & Schultz, E. T. Phenotypic similarity and the evolutionary significance of
countergradient variation. *Trends Ecol. Evol.* **10**, 248–252 (1995).

35.   Murren, C. J. *et al.* Evolutionary Change in Continuous Reaction Norms. *Am. Nat.* **183**, 453–467
(2014).

36.   Feder, M. E., Bennett, A. F. & Huey, R. B. EVOLUTIONARY PHYSIOLOGY1. 29 (2000).

37.   Huey, R. B. & Kingsolver, J. G. Climate Warming, Resource Availability, and the Metabolic
Meltdown of Ectotherms. *Am. Nat.* **194**, E140–E150 (2019).

38.   Skoracka, A. *et al.* Effective specialist or jack of all trades? Experimental evolution of a crop pest
in fluctuating and stable environments. *Evol. Appl.* **15**, 1639–1652 (2022).

39.   Alexandridis, N. *et al.* Archetype models upscale understanding of natural pest control response
to land-use change. *Ecol. Appl.* **n/a**, e2696.

40.   Towards an interactive, process-based approach to understanding range shifts: developmental
and environmental dependencies matter. doi:10.1111/ecog.03975.

41.   Barrett, R. & Schluter, D. Adaptation from standing genetic variation. *Trends Ecol. Evol.* **23**, 38–44
(2008).

42.   Bennett, J. M. *et al.* The evolution of critical thermal limits of life on Earth. *Nat. Commun.* **12**,
1198 (2021).

43.   Figueiredo, L., Krauss, J., Steffan-Dewenter, I. & Sarmento Cabral, J. Understanding extinction
debts: spatio–temporal scales, mechanisms and a roadmap for future research. *Ecography* **42**,
1973–1990 (2019).

44.   Chen, Y. *et al.* The combination of genomic offset and niche modelling provides insights into
climate change-driven vulnerability. *Nat. Commun.* **13**, 4821 (2022).

45.   Nadeau, C. P., Urban, M. C. & Bridle, J. R. Climates Past, Present, and Yet-to-Come Shape Climate
Change Vulnerabilities. *Trends Ecol. Evol.* **32**, 786–800 (2017).

46.   Eyring, H. The Activated Complex in Chemical Reactions. *J. Chem. Phys.* **3**, 107–115 (1935).

47.   Evans, M. G. & Polanyi, M. Some applications of the transition state method to the calculation of
reaction velocities, especially in solution. *Trans. Faraday Soc.* **31**, 875–894 (1935).

48.   Houle, D. Genetic Covariance of Fitness Correlates: What Genetic Correlations are Made of and
Why it Matters. *Evolution* **45**, 630 (1991).

49.   Jong, G. de & Noordwijk, A. J. van. Acquisition and Allocation of Resources: Genetic (CO)
Variances, Selection, and Life Histories. *Am. Nat.* **139**, 749–770 (1992).

50.   Tuda, M., Kagoshima, K., Toquenaga, Y. & Arnqvist, G. Global Genetic Differentiation in a
Cosmopolitan Pest of Stored Beans: Effects of Geography, Host-Plant Usage and Anthropogenic
Factors. *PLOS ONE* **9**, e106268 (2014).

51.   Roff, D. *Evolution Of Life Histories: Theory and Analysis*. (Springer Science & Business Media,
1993).

52.   Stearns, S. C. THE EVOLUTION OF LIFE HISTORIES.

53.   Cornish-Bowden, A. Kinetics, Enzymes. in *Encyclopedia of Bioprocess Technology* (John Wiley &
Sons, Inc., 2002). doi:10.1002/0471250589.ebt123.

54.   Clarke, A. Is there a Universal Temperature Dependence of metabolism? *Funct. Ecol.* **18**, 252–256
(2004).

55.   Gillooly, J. F. *et al.* Response to Clarke and Fraser: Effects of Temperature on Metabolic Rate.
Funct. Ecol. **20**, 400–404 (2006).

56.   Gillooly, J. F., Charnov, E. L., West, G. B., Savage, V. M. & Brown, J. H. Effects of size and
temperature on developmental time. *Nature* **417**, 70–73 (2002).

57.   Glazier, D. S. Beyond the▯: variation in the intra- and interspecific scaling of metabolic rate in
animals. *Biol. Rev.* **80**, 611 (2005).

58.   Berger, D. *et al.* Intraspecific variation in body size and the rate of reproduction in female insects
- adaptive allometry or biophysical constraint? *J. Anim. Ecol.* **81**, 1244–1258 (2012).

59. Kozlowski, J. & Konarzewski, M. Is West, Brown and Enquist's model of allometric scaling
mathematically correct and biologically relevant? *Funct. Ecol.* **18**, 283–289 (2004).
60. Gillooly, J. F., Brown, J. H., West, G. B., Savage, V. M. & Charnov, E. L. Effects of Size and
Temperature on Metabolic Rate. *Science* **293**, 2248–2251 (2001).
61. Dahlke, F. T., Wohlrab, S., Butzin, M. & Pörtner, H.-O. Thermal bottlenecks in the life cycle define
climate vulnerability of fish. *Science* **369**, 65–70 (2020).
62. Martin, T. L. & Huey, R. B. Why "Suboptimal" Is Optimal: Jensen's Inequality and Ectotherm
Thermal Preferences. *Am. Nat.* **171**, E102–E118 (2008).
63. Echave, J. & Wilke, C. O. Biophysical models of protein evolution: understanding the patterns of
evolutionary sequence divergence. *Annu. Rev. Biophys.* **46**, 85–103 (2017).
64. Bershtein, S., Serohijos, A. W. & Shakhnovich, E. I. Bridging the physical scales in evolutionary
biology: from protein sequence space to fitness of organisms and populations. *Curr. Opin. Struct.*
Biol. **42**, 31–40 (2017).
65. Glazier, D. S. Is metabolic rate a universal 'pacemaker' for biological processes? *Biol. Rev.* **90**,
377–407 (2015).
66. Addo-Bediako, A., Chown, S. L. & Gaston, K. J. Thermal tolerance, climatic variability and latitude.
Proc. R. Soc. Lond. B Biol. Sci. **267**, 739–745 (2000).
67. Dill, K. A., Ghosh, K. & Schmit, J. D. Physical limits of cells and proteomes. *Proc. Natl. Acad. Sci.*
108, 17876–17882 (2011).
68. Taipale, M., Jarosz, D. F. & Lindquist, S. HSP90 at the hub of protein homeostasis: emerging
mechanistic insights. *Nat. Rev. Mol. Cell Biol.* **11**, 515–528 (2010).
69. Mayer, M. P. & Bukau, B. Hsp70 chaperones: Cellular functions and molecular mechanism. *Cell.*
Mol. Life Sci. **62**, 670–684 (2005).
70. Kingsolver, J. G. & Woods, H. A. Beyond Thermal Performance Curves: Modeling Time-
Dependent Effects of Thermal Stress on Ectotherm Growth Rates. *Am. Nat.* **187**, 283–294 (2016).
71. Clarke, A. Temperature and the Metabolic Theory of Ecology. *Funct. Ecol.* **20**, 405–412 (2006).
72. Schulte, P. M. The effects of temperature on aerobic metabolism: towards a mechanistic
understanding of the responses of ectotherms to a changing environment. *J. Exp. Biol.* **218**,
1856–1866 (2015).
73. Chown, S. L. *et al.* Adapting to climate change: a perspective from evolutionary physiology. *Clim.*
Res. **43**, 3–15 (2010).
74. Fox, C. W. Multiple Mating, Lifetime Fecundity and Female Mortality of the Bruchid Beetle,
Callosobruchus maculatus (Coleoptera: Bruchidae). *Funct. Ecol.* **7**, 203–208 (1993).
75. Immonen, E., Berger, D., Sayadi, A., Liljestrand-Rönn, J. & Arnqvist, G. An experimental test of
temperature-dependent selection on mitochondrial haplotypes in Callosobruchus maculatus
seed beetles. *Ecol. Evol.* **10**, 11387–11398 (2020).
76. Collyer, M. L. & Adams, D. C. RRPP: An r package for fitting linear models to high-dimensional
data using residual randomization. *Methods Ecol. Evol.* **9**, 1772–1779 (2018).
77. Umeozor, O. C. Effect of the Infection of Callosobruchus maculatus (Fab.) on the Weight Loss of
Stored Cowpea (Vigna unguiculata (L.) Walp). *O C.*
78. Dent, D. & Binks, R. H. *Insect Pest Management, 3rd Edition.* (CABI, 2020).
79. Urban, M. C. Accelerating extinction risk from climate change. *Science* **348**, 571–573 (2015).
80. Wagner, D. L. Insect Declines in the Anthropocene. *Annu. Rev. Entomol.* **65**, 457–480 (2020).
81. Chown, S. L. & Gaston, K. J. Body size variation in insects: a macroecological perspective. *Biol.*
Rev. **85**, 139–169 (2010).
82. Urban, M. C. *et al.* Coding for Life: Designing a Platform for Projecting and Protecting Global
Biodiversity. *BioScience* **72**, 91–104 (2022).
83. Segoli, M. *et al.* Trait-based approaches to predicting biological control success: challenges and
prospects. *Trends Ecol. Evol.* **38**, 802–811 (2023).
84. Angilletta, M. J., Huey, R. B. & Frazier, M. R. Thermodynamic Effects on Organismal Performance:
Is Hotter Better? *Physiol. Biochem. Zool.* **83**, 197–206 (2010).

85. Berger, D., Walters, R. & Gotthard, K. What limits insect fecundity? Body size- and temperature-
dependent egg maturation and oviposition in a butterfly. *Funct. Ecol.* **22**, 523–529 (2008).

86. Frazier, M. R., Huey, R. B. & Berrigan, D. Thermodynamics Constrains the Evolution of Insect
Population Growth Rates: "Warmer Is Better". *Am. Nat.* **168**, 512–520 (2006).

87. Berger, D. *et al.* Intralocus Sexual Conflict and the Tragedy of the Commons in Seed Beetles. *Am.*
Nat. **188**, E98–E112 (2016).

88. Berger, D., Berg, E. C., Widegren, W., Arnqvist, G. & Maklakov, A. A. Multivariate intralocus
sexual conflict in seed beetles: SEXUAL CONFLICT AND LIFE-HISTORY PLEIOTROPY. *Evolution* **68**,
3457–3469 (2014).

89. Mallard, F., Nolte, V., Tobler, R., Kapun, M. & Schlötterer, C. A simple genetic basis of adaptation
to a novel thermal environment results in complex metabolic rewiring in Drosophila. *Genome*
Biol. **19**, 119 (2018).

90. Rogell, B. *et al.* Sex-dependent evolution of life-history traits following adaptation to climate
warming. *Funct. Ecol.* **28**, 469–478 (2014).

91. Ivimey-Cook, E. R., Piani, C., Hung, W.-T. & Berg, E. C. Genetic background and thermal regime
influence adaptation to novel environment in the seed beetle, Callosobruchus maculatus. *J. Evol.*
Biol. **37**, 1–13 (2024).

92. Angilletta, Jr., , Michael J. & Dunham, A. E. The Temperature-Size Rule in Ectotherms: Simple
Evolutionary Explanations May Not Be General. *Am. Nat.* **162**, 332–342 (2003).

93. Zuo, W., Moses, M. E., West, G. B., Hou, C. & Brown, J. H. A general model for effects of
temperature on ectotherm ontogenetic growth and development. *Proc. R. Soc. B Biol. Sci.* **279**,
1840–1846 (2011).

94. Thomas, M. K. *et al.* Temperature–nutrient interactions exacerbate sensitivity to warming in
phytoplankton. *Glob. Change Biol.* **23**, 3269–3280 (2017).

95. Awmack, C. S. & Leather, S. R. Host Plant Quality and Fecundity in Herbivorous Insects. *Annu.*
Rev. Entomol. **47**, 817–844 (2002).

96. Atkinson, D. & Sibly, R. M. Why are organisms usually bigger in colder environments? Making
sense of a life history puzzle. *Trends Ecol. Evol.* **12**, 235–239 (1997).

97. Blanckenhorn, W. U. Bergmann and Converse Bergmann Latitudinal Clines in Arthropods: Two
Ends of a Continuum? *Integr. Comp. Biol.* **44**, 413–424 (2004).

98. Horne, C. R., Hirst, Andrew. G. & Atkinson, D. Temperature-size responses match latitudinal-size
clines in arthropods, revealing critical differences between aquatic and terrestrial species. *Ecol.*
Lett. **18**, 327–335 (2015).

99. Stillwell, R. C., Wallin, W. G., Hitchcock, L. J. & Fox, C. W. Phenotypic plasticity in a complex
world: interactive effects of food and temperature on fitness components of a seed beetle.
Oecologia **153**, 309–321 (2007).

100.   Garcia-Robledo, C., Baer, C. S., Lippert, K. & Sarathy, V. Evolutionary history, not
ecogeographic rules, explains size variation of tropical insects along elevational gradients. *Funct.*
Ecol. **34**, 2513–2523 (2020).

101.   Whitlock, M. C. & Agrawal, A. F. PURGING THE GENOME WITH SEXUAL SELECTION:
REDUCING MUTATION LOAD THROUGH SELECTION ON MALES. *Evolution* **63**, 569–582 (2009).

102.   Arnqvist, G. & Rowe, L. *Sexual Conflict*. (Princeton University Press, 2005).

103.   Kokko, H. & Brooks, R. Sexy to die for? Sexual selection and the risk of extinction. *Ann.*
Zool. Fenn. **40**, 207–219 (2003).

104.   García-Roa, R., Garcia-Gonzalez, F., Noble, D. W. A. & Carazo, P. Temperature as a
modulator of sexual selection. *Biol. Rev.* **n/a**,.

105.   Parrett, J. M., Mann, D. J., Chung, A. Y. C., Slade, E. M. & Knell, R. J. Sexual selection
predicts the persistence of populations within altered environments. *Ecol. Lett.* **0**,.

106.   Plesnar-Bielak, A., Skrzynecka, A. M., Prokop, Z. M. & Radwan, J. Mating system affects
population performance and extinction risk under environmental challenge. *Proc. R. Soc. B Biol.*
Sci. **279**, 4661–4667 (2012).

107.   Martinossi-Allibert, I., Thilliez, E., Arnqvist, G. & Berger, D. Sexual selection,
environmental robustness, and evolutionary demography of maladapted populations: A test
using experimental evolution in seed beetles. *Evol. Appl.* **12**, 1371–1384 (2019).

108.   Baur, J., Jagusch, D., Michalak, P., Koppik, M. & Berger, D. The mating system affects
the temperature sensitivity of male and female fertility. *Funct. Ecol.* **36**, 92–106 (2022).

109.   Baur, J., Zwoinska, M., Koppik, M., Snook, R. R. & Berger, D. Heat stress reveals a
fertility debt owing to postcopulatory sexual selection. *Evol. Lett.* qrad007 (2023)
doi:10.1093/evlett/qrad007.

110.   Berger, D. & Liljestrand-Rönn, J. Environmental complexity mitigates the demographic
impact of sexual selection. *Ecol. Lett.* **27**, e14355 (2024).

111.   Martin. Goal 2: Zero Hunger. *United Nations Sustainable Development*
https://www.un.org/sustainabledevelopment/hunger/.

112.   Climate Change 2022: Impacts, Adaptation and Vulnerability.
https://www.ipcc.ch/report/ar6/wg2/.

113.   Cabral, J. S. *et al. The Road to Integrate Climate Change Effects on Land-Use Change in*
Regional Biodiversity Models. https://www.authorea.com/users/409019/articles/558105-the-
road-to-integrate-climate-change-effects-on-land-use-change-in-regional-biodiversity-
models?commit=1a1c610450c40c57b8bc51276fcbcc05242acfa5 (2022)
doi:10.22541/au.164608831.19029067/v1.

114.   Bocedi, G. *et al.* RangeShifter 2.0: an extended and enhanced platform for modelling
spatial eco-evolutionary dynamics and species' responses to environmental changes. *Ecography*
44, 1453–1462 (2021).

115.   Hagen, O. *et al.* gen3sis: A general engine for eco-evolutionary simulations of the
processes that shape Earth's biodiversity. *PLOS Biol.* **19**, e3001340 (2021).

116.   Dormann, C. F. *et al.* Correlation and process in species distribution models: bridging a
dichotomy. *J. Biogeogr.* **39**, 2119–2131 (2012).

117.   Mouquet, N. *et al.* REVIEW: Predictive ecology in a changing world. *J. Appl. Ecol.* **52**,
1293–1310 (2015).

118.   Zurell, D. *et al.* Benchmarking novel approaches for modelling species range dynamics.
Glob. Change Biol. **22**, 2651–2664 (2016).

119.   Anderson, R., Bayer, P. E. & Edwards, D. Climate change and the need for agricultural
adaptation. *Curr. Opin. Plant Biol.* **56**, 197–202 (2020).

120.   Garcia-Costoya, G., Williams, C. E., Faske, T. M., Moorman, J. D. & Logan, M. L.
Evolutionary constraints mediate extinction risk under climate change. *Ecol. Lett.* **26**, 529–539
(2023).

121.   Beck, C. W. A Handbook on Bean Beetles,. 15.

122.   Berger, D. *et al.* INTRALOCUS SEXUAL CONFLICT AND ENVIRONMENTAL STRESS: SEX,
GENES, AND CONFLICT IN STRESSFUL ENVIRONMENTS. *Evolution* n/a-n/a (2014)
doi:10.1111/evo.12439.

123.   Babraham Bioinformatics - FastQC A Quality Control tool for High Throughput
Sequence Data. https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

124.   Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: summarize analysis results
for multiple tools and samples in a single report. *Bioinforma. Oxf. Engl.* **32**, 3047–3048 (2016).

125.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

126.   Arnqvist, G. *et al.* A chromosome-level assembly of the seed beetle Callosobruchus
maculatus genome with annotation of its repetitive elements. *G3 GenesGenomesGenetics*
jkad266 (2023) doi:10.1093/g3journal/jkad266.

127.   Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nat. Biotechnol.* **37**, 907–915
(2019).

128. Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with high-
throughput sequencing data. *Bioinforma. Oxf. Engl.* **31**, 166–169 (2015).
129. R Core Team. *R: A Language and Environment for Statistical Computing.* (R Foundation
for Statistical Computing, Vienna, Austria, 2020).
130. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140
(2010).
131. McCarthy, D. J., Chen, Y. & Smyth, G. K. Differential expression analysis of multifactor
RNA-Seq experiments with respect to biological variation. *Nucleic Acids Res.* **40**, 4288–4297
(2012).
132. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing
and microarray studies. *Nucleic Acids Res.* **43**, e47 (2015).
133. Falcon, S. & Gentleman, R. Using GOstats to test gene lists for GO term association.
Bioinforma. Oxf. Engl. **23**, 257–258 (2007).
134. Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics
data. *The Innovation* **2**, 100141 (2021).
135. Bates, D. Mixed models in R using the lme4 package Part 4: Inference based on
profiled deviance. 23.
136. Immonen, E., Rönn, J., Watson, C., Berger, D. & Arnqvist, G. Complex mitonuclear
interactions and metabolic costs of mating in male seed beetles. *J. Evol. Biol.* **29**, 360–370 (2016).
137. Arnqvist, G., Rönn, J., Watson, C., Goenaga, J. & Immonen, E. Concerted evolution of
metabolic rate, economics of mating, ecology, and pace of life across seed beetles. *Proc. Natl.*
Acad. Sci. **119**, e2205564119 (2022).
138. Jackai, L. E. N. & Daoust, R. A. Insect Pests of Cowpeas. *Annu. Rev. Entomol.* **31**, 95–
119 (1986).

1121
1122
1123

Acknowledgements

We thank Johanna Liljestrand-Rönn for help in the lab, and Elina Immonen and Göran
Arnqvist for helpful input on our study design and results. Sequencing was performed by the
SNP&SEQ Technology Platform in Uppsala. The facility is part of the National Genomics
Infrastructure (NGI) Sweden and Science for Life Laboratory. The SNP&SEQ Platform is also
supported by the Swedish Research Council and the Knut and Alice Wallenberg Foundation.
This work was supported by grant no. 2019-05023 from the Swedish Research Council (VR),
grant no. 2022-01117 from Formas, and grant no. CTS22:2101 from Carl Tryggers Stiftelse,
to DB.

1133
1134

Author Contributions

1136 EB, CGT, MM, EC and JB planned, coordinated and performed the collection of all
1137 phenotypic data. MK and EB planned and performed the collection of gene expression data.
1138 AR and AH analyzed all gene expression data. DB designed the study, analyzed the
1139 phenotypic data, constructed the model, and wrote the manuscript. All authors read and
1140 commented on the first manuscript draft and approved the submission.

1141
1142

Competing Interests

1144 The authors declare no competing interests.

1145

Figure 1: Thermal adaptation via compensatory energy acquisition can increase pest species' impact on agriculture.

Shown are predictions based on protein kinetics (details in Supplementary 1). **A)** Temperature increases enzymatic reaction rates[20,56] (pink lines) while too hot temperatures lead to (reversable) protein unfolding (grey lines), leaving fewer enzymes available to do work[25,26]. Reduced costs of feeding (broken lines) leads to more energy reserves acquired (M), of which a fraction (p) can be devoted to increased production of catalytic enzymes that speed up constrained reaction rates at cold temperature[20], and another fraction (q) that can be devoted to molecular chaperones that stabilize proteins at hot temperature[36]. **B)** The resulting growth rate is the product of the remaining reserves that can be devoted to growth and reproduction, M (1-p-q), the reaction rate, and the proportion of properly folded protein. **C)** The resulting growth rates for populations adapting to cold (23°C; blue) and hot (35°C; red) temperature via compensatory responses (increased M, and allocation to p and q) are shown for high (full lines) and low (broken lines) costs of feeding. While the cold-adapted population outcompetes the hot-adapted population at cold temperature, growth rates are greater at warm temperatures and for the hot-adapted population due to thermodynamic constraints on reaction rates[23,56]. Horizontal lines on the right-hand y-axis show M at the optimal strategy; lower costs of feeding favour increased energy acquisition, which improves thermal performance and increases thermal niche breadth. In the depicted scenario, increased energy acquisition is particularly beneficial at hot temperature that puts increased demands on cellular maintenance to ensure protein stability. **D)** The possibility to evolve thermal performance via compensatory energy acquisition leads to an even greater agricultural impact (the product of M and population growth rate; compare difference between full and hatched lines in panels **C** and **D**). All panels show responses relative to the maximum value for ancestors reared at benign 29°C at high costs of feeding (black full line in panel B).

1175
Figure 2: Experimental evolution of thermal niches and life history in seed beetles.
Dryland legumes are nutritionally rich, drought tolerant and resilient to harsh weather[11,138]
and provide subsistence for more than half a billion people in the driest regions of the
world[11]. However, these crops suffer substantial losses from seed beetles[138]. To investigate
how genetic adaptation to climate change might shape the impact of seed beetles on legume
crops, we performed large-scale phenotyping of long-term experimental evolution lines of *C.*
maculatus. **A**) Populations from three geographic locations (Brazil, California and Yemen)
were sampled and allowed to adapt to common laboratory conditions for ca. 300 generations
before experimental evolution. Four replicate populations were created per genetic
background and split between the two evolution regimes (23°C and 35°C), while the original
ancestor remained at the benign temperature (29°C). Experimental evolution proceeded over
90 and 130 generations for the Cold (blue) and Hot (red) adapted populations. During this
time, the ancestors went through 130 generations of evolution. **B**) To provide molecular
evidence for a temperature-dependent trade-off between maintenance and reproduction,
gene expression was measured in both evolved and ancestral lines after 80-125 generations
of evolution. **C**) After 40 (Cold adapted) and 60 (Hot adapted) generations, life-history
adaptation was assessed in a common garden experiment. This data was complemented in
generations 90 (Cold lines) and 130 (Hot Lines and Ancestors). Life-history traits were
measured in the three ancestors after 120 generations of evolution. **D**) Following 80-125
generations of experimental evolution, the host consumption of all lines and ancestors was
scored in a common garden to assess how life-history evolution under simulated climate
change had shaped the agricultural footprint left by *C. maculatus*.

1198
1199
1200
Figure 3. Molecular evidence for a temperature-dependent trade-off between
maintenance and reproduction.
A) Differentially expressed genes in response to heat shock (HS) and mating (R); 137 genes
responded to both treatments, which was more than expected by chance. **B**) Of these genes,
all but three showed antagonistic responses to the two treatments, signifying a trade-off
between investment in reproduction and maintenance under heat stress. **C)** GO term
enrichments for genes upregulated (+) or downregulated (-) in response to mating (R) and
heat stress (HS), as well as for the overlapping antagonistic genes. **D)** The expression of the
134 antagonistic genes in the experimental evolution lines and ancestors indicate a clear
increase in allocation towards cellular maintenance at the cost of reduced reproduction when
beetles were reared at 35°C (the thermal regime of hot-adapted lines). However, there were
no consistent differences in expression between evolution regimes.
1213

1214

1215
1216
Figure 4. Thermal adaptation in life-history traits.
Single traits in (**A-G**) for Hot (red) and Cold (blue) adapted lines, as well as their Ancestors
(black). Development time is presented as a rate (1/development time in days) for
consistency with other traits. Means ± 1SE. **H**) Ordination using principal component analysis
showing that multivariate divergence occurs along both a temperature-sensitive axis (PC1)
and along a general life-history axis describing overall variation in body mass and LRS
(PC2), with more divergence from the ancestor in Hot-adapted lines.
1224

Figure 5: Temperature-dependent evolution of the agricultural footprint.
A) The relationship between beetle mass produced and host bean consumed. Each assay is represented by an individual datapoint. The slopes of the dashed lines correspond to the temperature-specific constant (C_T) giving the amount of host consumed per beetle produced (23°C: 3.17, 29°C: 3.66, 35°C: 3.55). **B**) The two independent metrics used to predict the agricultural footprint. **C**) The predicted footprint of each line (using the first metric) relative to respective ancestor raised at 29°C (grey hatched line). Means ± 95% CIs. The effect of life-history adaptation at cold temperature is small, but life-history adaptation at hot temperature almost doubles the agricultural footprint relative to the ancestral impact.