1 **Partitioning changes in ecosystem productivity by effects of species interactions in**

2 **biodiversity experiments**

4 Jing Tao[1], Charles A. Nock[2], Eric B. Searle[3], Shongming Huang[4], Rongzhou Man[3*], Hua Yang[5*],
5 Grégoire T. Freschet[6], Cyrille Violle[7], and Ji Zheng[8]
6

8 [1]Jilin Provincial Academy of Forestry Sciences, Changchun, China taojing8116@126.com
9
10 [2]Department of Renewable Resources, Faculty of Agriculture, Life, and Environmental Sciences,
11 University of Alberta, Edmonton, Alberta, Canada, T6G 2H1 nock@ualberta.ca
12
13 [3]Ontario Ministry of Natural Resources and Forestry, Ontario Forest Research Institute, 1235
14 Queen Street East, Sault Ste. Marie, ON, P6A 2E5 Canada eric.searle@ontario.ca;
15 rongzhou.man@ontario.ca; wayne.bell@ontario.ca
16
17 [4]Forestry Division, Department of Agriculture and Forestry, Government of Alberta, Edmonton,
18 Alberta, Canada T6H 5T6 shongming.huang@gov.ab.ca
19
20 [5]College of Forestry, Beijing Forestry University, China huayang8747@163.com
21
22 [6]Station d'Ecologie Théorique et Expérimentale, CNRS, 2 route du CNRS, 09200 Moulis,
23 France; gregoire.freschet@sete.cnrs.fr
24
25 [7]CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France; cyrille.violle@cefe.cnrs.fr
26
27 [8]School of Agriculture and Biology, and Shanghai Urban Forest Ecosystem Research Station of
28 National Forestry and Grassland Administration, Shanghai Jiao Tong University, China
29 zhengjiji456@sjtu.edu.cn
30

31
[*]Corresponding authors: rongzhou.man@ontario.ca; huayang8747@163.com

Abstract

Species interactions affect ecosystem productivity. Positive interactions (resource partitioning

and facilitation) increase productivity while negative interactions (species interference) decrease

productivity relative to the null expectations defined by monoculture yields. Effects of

competitive interactions (resource competition) can be either positive or negative. Distinguishing

effects of species interactions is therefore difficult, if not impossible, with current biodiversity

experiments involving mixtures and full density monocultures.

To partition changes in ecosystem productivity by effects of species interactions, we

modify null expectations with competitive growth responses, i.e., proportional changes in

individual size (biomass or volume) expected in mixture based on species differences in growth

and competitive ability. We use partial density (species density in mixture) monocultures and the

competitive exclusion principle to determine maximum competitive growth responses and full

density monoculture yields to measure species ability to achieve maximum competitive growth

responses in mixture. Deviations of observed yields from competitive expectations represent the

effects of positive/negative species interactions, while the differences between competitive and

null expectations reflect the effects of competitive interactions.

We demonstrate the effectiveness of our competitive partitioning model in distinguishing

effects of species interactions using both simulated and experimental species mixtures. Our

competitive partitioning model enables meaningful assessments of species interactions at both

species and community levels and helps disentangle underlying mechanisms of species

interactions responsible for changes in ecosystem productivity and identify species mixtures that

maximize positive effects.

Keywords: competitive interaction, null expectation, competitive expectation, biodiversity effect,

competitive partitioning

Species interactions and ecosystem productivity

Detecting changes in ecosystem productivity with biodiversity and understanding how these changes are affected by mechanisms of species interactions has been a central focus in ecology for decades (Loreau, 2000; Loreau and Hector, 2019; Mahaut et al., 2020). However, effective methods are lacking (Drake, 2003; Forrester and Pretzsch, 2015; Jaillard et al., 2018; Barry et al., 2019; Loreau and Hector, 2019; Bourrat et al., 2023). The method of additive partitioning was developed from the covariance equation to quantify changes in ecosystem productivity by comparing species performance in mixture with those expected from their monocultures using a replacement design (de Wit, 1960; Loreau and Hector, 2001; Barry et al., 2019). The resultant changes in ecosystem productivity (commonly referred to as net biodiversity effects) are mathematically decomposed into additive components, namely the complementarity effect (CE) measuring community average response and the selection effect (SE) measuring interspecies differences (Loreau and Hector, 2001; Forrester and Pretzsch, 2015; Loreau and Hector, 2019). CE ad SE are widely used to examine mechanisms of species interactions responsible for biodiversity effects, i.e., CE for positive interactions and SE for competitive interactions (Loreau and Hector, 2001; Petchey, 2003; Polley et al., 2003; Montès et al., 2008; Barry et al., 2019; Hagan et al., 2021; Feng et al., 2022), despite that neither of the additive components can be linked to specific effects of species interactions (Petchey, 2003; Hooper et al., 2005; Montès et al., 2008; Loreau et al., 2012; Forrester and Pretzsch, 2015; Loreau and Hector, 2019; Hagan et al., 2021; Bourrat et al., 2023).

Among the interspecific interactions in plants, positive interactions (resource partitioning and facilitation or collectively referred to as complementarity) increase productivity and negative interactions (species interference or interference competition) decrease productivity relative to

the null expectations, i.e., species yields expected in mixture based on composition (relative

yield) and full density monoculture yields (Loreau, 2000; Loreau and et al., 2005; Barry et al.,

2019). Uncertainty comes with effects of competitive interactions (i.e., resource competition)

that are positive for more competitive species and negative for less competitive species

(Aschehoug et al., 2016; Barry et al., 2019). For example, when a highly competitive and less

competitive species are mixed, more resources flow to the more competitive species. The yield

gain of more competitive species from increases in resource availability (resources per capita)

often exceeds the yield loss of less competitive species from decreases in resource availability

(Huston, 1997; Tilman et al., 1997; Wardle, 1999; Montazeaud et al., 2018; Pillai and Gouhier,

2019), resulting in positive deviation from the total expected yield of both species (i.e., positive

biodiversity effect). Competitive interactions can also result in negative biodiversity effects if the

less competitive species suffers more than the more competitive species benefits.

The positive effects of competitive dominance on ecosystem productivity can be

illustrated with species mixture data. In the situations of both simulated and experimental

mixtures with two species, the partial density monoculture yield of the more competitive species

exceeded the total expected yield of both species in 32 of the 35 mixtures (*Table 1*), meaning

that a higher productivity expected for a given species mixture under the null hypothesis can be

generally achieved by removing less competitive species from the mixture (*Table S1; Table S2*).

In other words, the positive biodiversity effects of mixtures detected through the null expectation

can be obtained from the monocultures of more competitive species at partial (lower) densities.

These positive biodiversity effects, however, result from competition-induced changes in

resource availability, i.e., the yield gain of more competitive species from all resources in the

mixture exceeds the total yield loss expected for the less competitive species, and are not related

to effects of positive species interactions. Competitive interactions are the predominant type of

interspecific relationships in plants and occur in all mixtures where constituent species are

competitively different (Goldberg and Barton, 1992; Pillai and Gouhier, 2019). Reporting

positive biodiversity effects, without indication of possible contributions from competitive

interactions, can lead to unrealistic expectations for effects of positive interactions.

Given the limits of the additive partitioning in deciphering mechanisms of changes in

ecosystem productivity (Loreau and Hector, 2001; Petchey, 2003; Forrester and Pretzsch, 2015),

some alternative methods have been explored to identify underlying mechanisms. This includes

the use of metrics such as relative yield totals (Hector, 1998; Roscher and Schumacher, 2016)

and transgressive overyielding or superyielding (Drake, 2003; Forrester and Pretzsch, 2015;

Špaèková and Lepš, 2001), the multiplicative partitioning or statistical analyses to separate

effects of species interactions and species composition (Kirwan et al., 2009; Jaillard et al., 2018,

2021; Chen et al., 2020), the identification of relative species influences via measurement of

competitive interactions (Montès et al., 2008; Mahaut et al., 2020; Brun et al., 2022), or

influences of species diversity, composition, and density on net biodiversity effects (Polley et al.,

2003; Roscher and Schumacher, 2016; Jaillard et al., 2018; Mahaut et al., 2020; Tatsumi and

Loreau, 2023). However, although these methods may indicate the presence of strong positive or

competitive interactions or connections of the additive components with niche and fitness

differences (Carroll et al., 2011; Turnbull et al., 2013; Godoy et al., 2020), they are not capable

of quantifying changes in ecosystem productivity by species interactions at species or

community level.

Here, we present the competitive partitioning model, a new framework for assessing

effects of species interactions on ecosystem productivity based on null expectations and

competitive growth responses, the proportional changes of individual size (biomass or volume)

from full density monocultures to mixtures expected from species differences in growth and

competitive ability (*Figure 1*). Current biodiversity experiments are generally established based

on a replacement design with equal density across mixtures and monocultures (Sackville

Hamilton, 1994) and do not provide data for estimating competitive growth responses. Thus, we

applied this new methodological approach to simulated data generated with the GYPSY model

(Huang et al., 2009) for trembling aspen (*Populus tremuloides* Michx.) and white spruce (*Picea*

glauca (Moench) Voss) tree mixtures and greenhouse experimental data from grassland mixtures

(Mahaut et al., 2020) (*Table 1*). We used partial density (species density in mixture)

monocultures and the competitive exclusion principle to determine maximum competitive

growth responses and full density monoculture yields to measure species ability to compete for

resources and achieve maximum competitive growth responses in mixture. Our objectives are to

(1) demonstrate the power of this new framework in distinguishing effects of species interactions

responsible for changes in ecosystem productivity, (2) illustrate how the mechanisms of changes

in ecosystem productivity differ between the assessments with competitive and additive

partitioning models, and (3) assess the effects of competitive interactions on ecosystem

productivity relative to those of other species interactions.

Competitive expectation

Under the null hypothesis, species in mixture are assumed to be competitively equivalent (i.e.,

absence of interspecific interactions) and species growth (or any other function) stays the same in

mixture as in monocultures (Loreau and Hector, 2001). Deviations of observed yields (biomass

or volume) from total null expectation therefore produce net biodiversity effects or changes in

ecosystem productivity (Loreau and Hector, 2001; Barry *et al.*, 2019).

$$\sum Y_{Oi} - \sum Y_{NEi} = \sum RY_{Oi}M_{if} - \sum RY_{Ei}M_{if}$$

$$= \sum \Delta RY_i M_{if} \qquad\qquad (1)$$

Thus, the net biodiversity effect is the difference between total observed yields $\sum Y_{Oi}$ and total

null expectation $\sum Y_{NEi}$, calculated from species composition (or relative yield) in mixture RY_{Ei}

and full density monoculture yield M_{if}. Mathematically, the net biodiversity effect is an

arithmetic mean of full density monoculture yields weighted by changes in species relative yield

ΔRY_I, and results from all species interactions including positive (positive biodiversity effects),

competitive (positive or negative biodiversity effects), and negative (negative biodiversity effects)

interactions. Net biodiversity effects can be mathematically decomposed into an arithmetic mean

(a product of mean monoculture yield by mean relative yield change) representing community

average response (CE) and a covariance (between monoculture yield and relative yield change)

representing interspecies differences (SE) (Loreau and Hector, 2001).

To separate effects of competitive interactions from those of other species interactions,

we would need the hypothesis that constituent species share an identical niche but differ in

growth and competitive ability (i.e., absence of positive/negative interactions). Competitive

growth responses, i.e., proportional changes in individual size (biomass or volume) expected

from full density monocultures to mixtures under this competitive hypothesis, can be determined

from species differences in growth and competitive ability (*Figure 1*). The magnitudes of

competitive growth responses depend on: (1) species resource needs in full density monocultures;

and (2) changes in resource availability (resources per capita) from full density monocultures to

full density mixtures that are related to species composition and relative ability to compete for

resources in mixture.

To estimate competitive growth response, we first determine the maximum competitive

growth response MG_i that species can reach in a scenario of competitive exclusion (Murray, 1986)

where a more competitive species completely dominates and grows like a partial density

monoculture, while a less competitive species is competitively eliminated.

$$MG_i = \frac{w_{ip}}{w_{if}} - 1, \text{ for more competitive species}$$

$$= -1, \text{ for less competitive species} \qquad (2)$$

Here, w_{ip} stands for individual size (biomass or volume) in partial density (species density in

mixture) monoculture and w_{if} is for individual size in full density monoculture; both parameters

are calculated from species monoculture yields and initial densities in partial/full density

monocultures. A more competitive species gains $\frac{1}{RY_{Ei}} - 1$ in resource availability and $\frac{w_{ip}}{w_{if}} - 1$ in

individual size, while a less competitive species loses 100% resource availability and size in

mixture relative to full density monoculture (***Figure 1***). This simplified competitive dominance

applies to mixtures where constituent species differ considerably in competitive ability,

especially when competition for light is intense (Grime, 1973; Lieffers et al., 1993). In most

cases, however, competitive exclusion would not occur in biodiversity experiments that are

controlled in density, composition, duration, and scale. More competitive species are negatively

affected by interspecific competition relative to their partial density monocultures (Holland and

DeAngelis, 2009), while less competitive species are competitively suppressed relative to their

null expectations, but not eliminated (Aarssen, 1989; Connolly et al., 2001; Montazeaud et al.,

2018; Mahaut et al., 2020).

Next, we use relative competitive ability RC_i to represent species ability to compete for

resources and achieve their maximum competitive growth response in mixture (Aarssen, 1989).

RC_i is defined by the deviation of species competitive ability H_i from community average $\overline{H}_{i\notin}$

(excluding species i) relative to community maximum H_m. Competitive ability is species ability

to exploit resources and can be assessed by growth attributes such as full density monoculture

yields in biomass or volume (Gaudet and Keddy, 1988; Aarssen, 1989; Grace, 1990; Aschehoug

et al., 2016; Montazeaud et al., 2018).

199                  $$RC_i = \frac{|H_i - \overline{H}_{i\notin}|}{H_m}$$ (3)

200   Greater competitive advantages would lead to increasingly positive competitive growth

201   responses and larger competitive disadvantages would lead to increasingly negative competitive

202   growth responses in mixture relative to full density monocultures.

203                  $$Y_{CEi} = Y_{NEi}[1 + MG_i RC_i]$$ (4)

Species competitive expectation Y_{CEi} is the yield based on null expectation Y_{NEi} and competitive

growth response $MG_i RC_i$ derived from species differences in growth and competitive ability

(*Figure 1*). The term $MG_i RC_i$ would be positive for more competitive species $MG_i > 0$, negative

for less competitive species $MG_i < 0$, and zero when species are at community average or

competitively equivalent $RC_i = 0$. At individual species level, the deviation of observed yield

Y_{Oi} from the competitive expectation Y_{CEi} can be positive (representing relaxation of interspecific

competition from competitive expectation and therefore dominance of positive interactions),

neutral (representing similar intra- and interspecific interactions or offset of positive and negative

interactions), or negative (representing dominance of negative interactions). The differences

between competitive and null expectations are competitive yield changes resulting from

competitive interactions. Isolating facilitative effect from those of other positive interactions (i.e.,

resource partitioning) is also possible for individual species by comparing species' observed

yield in mixture to their partial density monoculture yield, adapting the principle of additive

design for biodiversity–ecosystem productivity experiments (Sackville Hamilton, 1994). At the

community level, the difference in total yield between competitive expectations $\sum Y_{CEi}$ and null

expectations $\sum Y_{NEi}$ of all species can be attributed to the competitive effect, whereas the

difference between observed yields $\sum Y_{Oi}$ and competitive expectations $\sum Y_{CEi}$ can be attributed to

the dominance of positive effects when the difference >0 or to the dominance of negative effects

when the difference <0 (***Figure 1***).

Case studies

Simulated aspen and spruce mixtures

Mixed forests composed of trembling aspen (*Populus tremuloides* Michx.) and white spruce

(*Picea glauca* (Moench) Voss) are widely distributed in North America (Man and Lieffers, 1999).

The two species, occurring in pure and mixed stands of varying compositions in natural

conditions, generally have different canopy structure, rooting depth, leaf phenology, and

requirements for light, moisture, and nutrients so are considered different in niche (Man and

Lieffers, 1999).

We used a growth and yield simulation system GYPSY, developed for mixed species

forests in western Canada (Huang et al., 2009), to generate data required for determining

competitive expectation, i.e., mixtures and monocultures at partial and full densities. We were

interested in knowing how effects of species interactions may vary with stand age and

composition. We chose four ages: 20, 40, 60, and 80 years, to represent young, middle-aged, pre-

mature, and mature stands, and five relative yields from nearly pure aspen (90% aspen), aspen-

dominated (70% aspen), equal mixed (50% aspen), spruce-dominated (30% aspen), to nearly

pure spruce (10% aspen) mixtures at average densities on fully stocked medium productivity

sites (Peterson and Peterson, 1992) (*Table 1*). As GYPSY is developed from extensive field

observations (Huang et al., 2009), the simulated data represent average yields in natural

monocultures and mixtures of various composition and ages that may result from direct or

indirect interactions between the two species (Man and Lieffers, 1999; Barry et al., 2019).

Species competitive ability was approximated with stand volume in full density monocultures at

site index 17 m and 14 m (height), respectively, for dominant and codominant aspen and spruce

at 50 years of total age on medium productivity sites (Alberta Forest Service, 1985; Huang et al.,

1994).

The partitioning of net biodiversity effects with competitive partitioning model indicated

that the relative contribution of competitive interactions to net biodiversity effects decreased with

age (*Figure 2a*), whereas community positive effects occurred generally in pre-mature and

mature stands and more so in equal aspen and spruce mixtures (*Figure 2b*). Evidence for a

facilitative effect (i.e., species observed yield in mixture > partial density monoculture yield) was

only found on spruce in nearly pure mature spruce mixture (*Table 1*). Community negative

effects occurred in young and middled-age mixtures where aspen did poorer than expected under

the competitive expectation (*Table S1*). Averaged across all ages and compositions, the net

biodiversity effect was 14.3 m^3 ha^{-1} or 14% of the average full density monoculture yield. On a

relative basis, 57% of the value (8.2 m^3 ha^{-1}) was attributable to positive effects, 65% (9.3 m^3 ha^{-}

^{1}) to competitive effects and -23% (-3.2 m^3 ha^{-1}) to negative effects, compared to 63% and 37%

for CE and SE with additive partitioning (*Figure 2c; Figure 2d*).

Experimental grassland mixtures

Mahaut et al. (2020) conducted an experimental study in pots with six grassland species in full

and half-density monocultures, as well as in every combination of two, three and six species in

mixture. The studied species vary in productivity and functional traits. We used aboveground

biomass data from full and half-density monocultures, as well as all combinations of two species

in full density mixtures (***Table 1***). Species competitive ability was approximated with species

biomass in full density monocultures.

Competitive partitioning suggested a weak facilitative effect in two mixtures that were

associated with *Lotus corniculatus*, one of the two legume species studied (***Table 1; Table S2***).

Community negative effects occurred in two mixtures with *Sanguisorba minor* and between the

two legumes, due to negative effects of *Sanguisorba minor* on other species and *Lotus*

corniculatus on *Trifolium repens* (***Table S2***). Across all 15 two-species mixtures, the average of

net biodiversity effects on aboveground biomass was 3.9 g/pot or 49% of the full density

monoculture yield averaged across all species. On a relative basis, 60% (2.3 g) of the value was

attributable to positive effects, 45% (1.7 g) to competitive effects, and -5% (-0.2 g) to negative

effects, compared to 84% and 16% for CE and SE with additive partitioning (***Table 2***).

Comparison of additive and competitive partitioning

The competitive partitioning model provides a conceptual framework that helps establish the

linkage between changes in ecosystem productivity and effects of species interactions. Under

this framework, in both simulated tree and experimental grassland mixtures, net biodiversity

effects were generally positive, resulting from both positive and competitive interactions. The

positive competitive effects result from greater competitive yield responses of more competitive

species (Huston, 1997; Tilman et al., 1997; Wardle, 1999; Montazeaud et al., 2018), i.e., the

yield gain of more competitive species from increases in resource availability exceeds the yield

loss of less competitive species from decrease in resource availability, due to different

competitive growth responses and full density monoculture yields (***Table S1; Table S2***). The

greater competitive yield response of more competitive trembling aspen in young and middle-

aged mixtures resulted in net biodiversity effects that were predominantly from competitive

interactions (***Figure 2a; Figure 2b; Table S1***). A facilitative effect was rarely detected, only on

spruce in nearly pure spruce mature mixture, possibly due to improved nutrient availability by

aspen (Peterson and Peterson, 1992; Man and Lieffers, 1999) and in mixtures with legume *Lotus*

corniculatus in grassland mixtures, likely due to improved nitrogen availability (Mahaut et al.,

2020). The strong negative effect of mixing on aspen (i.e., observed yields < competitive

expectations, see ***Table S1***) in young and middled-age mixtures suggests that trembling aspen

originating naturally in high density does not benefit from mixing with white spruce. This is in

contrast with white spruce where the deviations of observed yields from competitive

expectations were generally positive, likely due to its shade tolerance and positive effects of

aspen on nutrient availability (Peterson and Peterson, 1992; Man and Lieffers, 1999).

The net biodiversity effects reveal overall community yield changes from what would be

expected from full density monoculture yields (Loreau and Hector, 2001; Loreau and Hector,

2019). However, the underlying mechanisms of species interactions for detected changes are

uncertain, due to effects of competitive interactions that can be positive or negative. In both

simulated mixed forests and experimental grassland mixtures, the partial density monoculture

yield of the more competitive species exceeded the total expected yield of both species in nearly

all mixtures (see ***Table S1*** and ***Table S2***), resulting in positive biodiversity effects when positive

species interactions are not involved. Such positive biodiversity effects may help understand

changes in ecosystem productivity detected through the null expectation (Carroll et al., 2011;

Godoy et al., 2020) but do not support the general expectation that positive changes in ecosystem

productivity are associated with positive species interactions (Loreau, 2000; Drake, 2003;

Hooper et al., 2005; Barry et al., 2019) or crop mixing in agriculture and forestry where interests

are positive effects (Man & Lieffers, 1999; Forrester & Pretzsch, 2015; Montazeaud et al., 2018).

Thus, partitioning net biodiversity effects by effects of species interactions (i.e., positive,

competitive, negative effects) allows for estimating relative importance of mechanisms of

species interactions responsible for changes in ecosystem productivity. This enables ecologists to

determine which communities are truly benefitting from diversity in the form of positive

interactions and which result from interspecies differences in growth and competitive ability. In

the aspen and spruce mixtures examined, community positive effects occurred largely in pre-

mature and mature mixtures (>=60 years), even though net biodiversity effects and CE were

positive at nearly all stages of stand development (*Figure 2*). In grassland mixtures, two

mixtures (i.e., *Dactylis glomerata - Sanguisorba minor* and *Lotus corniculatus - Trifolium repens*)

had positive net biodiversity effects that resulted entirely from competitive interactions (*Table 2;*

Table S2). These positive net biodiversity effects may indicate evidence for positive changes in

ecosystem productivity when these changes were actually driven by differential outcomes of

interspecific competition.

We demonstrate with simulated and experimental species mixtures that competitive

interaction is a major source of the net biodiversity effects and affects the additive components

of both CE and SE. In the scenario of partial density monocultures emulating a simplified

scenario of competitive exclusion where positive net biodiversity effects resulted entirely from

competitive interactions, CE was generally negative in tree mixtures where relative yield

changes were smaller with more competitive species (negative mean relative yield change) but

positive in grassland mixtures where relative yield changes were greater with more competitive

species (positive mean relative yield change) (*Table S1; Table S2*). In aspen-spruce mixtures, SE

was either larger (age 80) or smaller (age 20, 40, and 60) than competitive effect. In the

grassland communities, SE was similar to competitive effect in 3 of the 15 mixtures (*Bromus*

erectus - Lotus corniculatus, *Dactylis glomerata - Lotus corniculatus*, and *Lotus corniculatus -*

Sanguisorba minor) but substantially smaller in the remaining mixtures (*Table 2*). Even in the

extreme cases where SE was nearly zero (*Bromus erectus - Sanguisorba minor* and *Bromus*

erectus - Trifolium repens), competitive effect was still relatively strong, suggesting that a

common assumption to attribute positive net biodiversity effects to positive species interactions

when SE is close to zero (Loreau and Hector, 2001; Feng et al., 2022) is not justified.

Competitive interactions amplify interspecies differences and therefore SE, while species

interference on more competitive species (*Bromus erectus - Sanguisorba minor*) and greater

positive species interactions on less competitive species (*Bromus erectus - Trifolium repens*)

(*Table S2*) reduce SE. Our findings, along with some recent suggestions (Carroll et al., 2011;

Turnbull et al., 2013; Godoy et al., 2020; Bourrat et al., 2023), do not support some common

exercises that attribute positive/negative CEs to positive/negative interactions (Loreau and

Hector, 2001; Petchey, 2003; Fargione et al., 2007; Barry et al., 2019; Hagan et al., 2021; Feng

et al., 2022) and positive SEs to competitive interactions (Loreau, 2000; Loreau and Hector,

2001; Montazeaud et al., 2018).

Several assumptions are adopted with the development of competitive partitioning model.

First, we assumed that more productive species are more competitive, which was mostly true

with the species mixtures examined in terms of species yields in mixture relative to their full

density monoculture yields (*Table 1*). There were exceptions with Dactylis:Plantago and

Plantago:Trifolium in grassland mixtures where less productive species Plantago based on full

density monoculture yields performed better in mixtures, which may suggest positive effects on

Plantago and negative effects on the other species. However, species yields in partial versus full

density monocultures indicate that these exceptions resulted from the stagnation of more

productive Plantago in full (high) density monocultures, due to restricted rooting volumes and

supplies of water and nutrients in pots (Chung et al., 2021). The release of growth stagnation in

partial density monocultures and mixtures affects not only the mechanisms of changes in

ecosystem productivity assessed with competitive and additive partitioning models but also the

magnitudes of the changes in ecosystem productivity detected through the null expectation

(Loreau and Hector, 2001). In this case, the use of partial density monoculture yields in

assessment of species competitive ability are probably more appropriate, which would increase

the values of competitive effects and reduce those of positive effects based on the competitive

expectations. Partial density monocultures help detect competition-related yield changes, which

would be otherwise unknown. Second, we used partial density monocultures to determine

maximum competitive growth responses in a simplified competitive exclusion where more

competitive species are not affected and grow like partial density monocultures, while less

competitive species are competitively eliminated. We assumed a linear relationship between

competitive growth responses and species relative competitive ability (*Figure 1*), which should

be a good approximation (Gaudet and Keddy, 1988), but may not be the most appropriate, e.g.,

light availability and plant height (Brüllhardt et al., 2020). Linear models can overestimate or

underestimate competitive expectations of all species and affect individual species assessments

but would have limited impacts on community level assessments, due to different directions of

competitive growth responses among species.

The competitive partitioning model incorporates effects of competitive interactions into

the conventional null expectation and assists: (1) understanding the mechanisms of species

interactions driving positive biodiversity-productivity relationships (Cardinale et al., 2012; Isbell

et al., 2017) by relative contributions of positive interactions from more species of diverse niche

(Tilman et al., 1997; Hooper et al., 2005) and competitive interactions from greater yield

responses of more competitive and more productive species to changes in resource availability in

mixture (Aarssen, 1997; Huston, 1997; Tilman et al., 1997; Pillai and Gouhier, 2019), (2)

examining the mechanisms of key species in influencing ecosystem productivity by roles of

competitive interactions due to their higher productivity and competitive advantages (Loreau and

Hector, 2001; Mahaut et al., 2020), (3) meaningful comparisons of changes in ecosystem

productivity (i.e., relative yield totals) across different ecosystems based on competitive

expectations, and (4) redefining changes in ecosystem productivity such that positive changes

result from positive interactions and negative changes from negative interactions (Loreau, 2000;

Drake, 2003; Hooper et al., 2005). Under this competitive partitioning framework, null

expectations are either raised for more competitive species with increased resource availability or

lowered for less competitive species with decreased resource availability in mixture relative to

full density monocultures. The magnitudes of competitive growth responses are derived from

species relative competitive ability in mixture and their maximum competitive growth responses

determined from full and partial density monocultures under a simplified scenario of competitive

exclusion. Current biodiversity experiments do not have partial density monocultures and

therefore do not provide estimates of competitive growth responses that would occur in mixture

due to interspecies differences in growth and competitive ability. However, density-size/yield

relationship is one of the most extensively studied areas in ecology (Watkinson, 1980; Weiner

and Freckleton, 2010). Previous research can help determine maximum competitive growth

responses through density-size/yield relationships (Huang et al., 2009). This means that the

competitive partitioning model can be used to investigate changes in ecosystem productivity by

effects of species interactions in current biodiversity experiments established with replacement

design (Loreau and Hector, 2001; Fargione et al., 2007).

Conclusions

Competitive interactions are the major source of biodiversity effects and affect the additive

components of both CE and SE. The interpretations of CEs and SEs with specific effects of

species interactions lack mathematical or ecological justifications. For example, attributing

positive biodiversity effects solely to effects of positive interactions, as commonly seen in the

literature (Loreau and Hector, 2001; Hooper et al., 2005; Fargione et al., 2007; Barry et al., 2019;

Feng et al., 2022), assumes that competitive interactions are productivity-neutral (SE close to

zero) (Loreau and Hector, 2001; Feng et al., 2022). This assumption is true if ecosystem

productivity is defined by total resources pool (Tilman et al., 2014) but generally not true if

mixture productivity is defined by species monoculture yields as with the additive partitioning

(Loreau and Hector, 2001), due to the predominance of competitive interactions in interspecific

relationships (Goldberg and Barton, 1992) and positive effects of competitive interactions on

ecosystem productivity based on the null expectation.

The competitive partitioning model is based on ecological theories of interspecific

relationships, competitive ability, and competitive exclusion principle and experimental designs

of both replacement and additive series. This new partitioning approach enables meaningful

assessments of species interactions at both species and community levels and provides detailed

insights into the mechanisms of species interactions that drive changes in ecosystem productivity.

We believe that our framework, which is admittedly perfectible, is one promising avenue to

determine effects of species interactions responsible for changes in ecosystem productivity, an

approach that is long sought after in biodiversity–ecosystem productivity research (Loreau and

Hector, 2019; Mahaut et al., 2020).

Acknowledgements

431 We thank Gordon Kayahara and Bill Parker of Ontario Ministry of Natural Resources and

432 Forestry, and Dan Kneeshaw of Université du Québec à Montréal for helpful criticism and

433 comments on previous versions of this article, and Phil Comeau of University of Alberta for

434 advice on growth and yield models.

Author contributions

437 R.M. and H.Y. designed the research; J.T., S.H. and R.M. generated growth and yield data with

438 the GYPSY; and J.T., C.A.N., E.B.S., S.H., R.M., H.Y., G.T.F., C.V., and J.Z. contributed to the

439 interpretations of the results and to the writing of the paper.

References

Aarssen LW. 1997. High productivity in grassland ecosystems: effected by species diversity or

productive species? *Oikos* **80**:183–184.

Aarssen LW. 1989. Competitive ability and species coexistence: a 'plant's-eye' view. *Oikos*

56:386401.

Aschehoug ET, Brooker R, Atwater DZ, Maron JL, Callaway RM. 2016. The mechanisms and

consequences of interspecifc competition among plants. *Annual Review of Ecology,*

Evolution, and Systematics **47**:263–81.

Alberta Forest Service. 1985. *Alberta Phase 3 forest inventory - yield tables for unmanaged*

stands. Edmonton, Alberta: Alberta Forest Service.

Barry KE, Mommer L, Ruijven J, Wirth C, Wright AJ, Bai Y, …Weigelt A. 2019. The future of

complementarity: disentangling causes from consequences. *Trends in Ecology and*

Evolution **34**:167–180.

Bourrat B, Godsoe W, Pillai P, Gouhier TC, Ulrich W, Gotelli NJ, van Veelen M. 2023. What is

the price of using the Price equation in ecology? *Oikos* **2023**:e10024.

Brüllhardt M, Rotach P, Schleppi P, Bugmann H. 2020. Vertical light transmission profiles in

structured mixed deciduous forest canopies assessed by UAV-based hemispherical

photography and photogrammetric vegetation height models. *Agricultural and Forest*

Meteorology **281**:107843.

Brun P, Violle C, Mouillot D, Mouquet N, Enquist BJ, Munoz F, … Thuiller W. 2022. Plant

community impact on productivity: Trait diversity or key (stone) species effects? *Ecology*

letters **25**:913–925.

Cardinale BJ, Duffy JE, Gonzalez A, Hooper DU, Perrings C, Venail P, … Naeem S. 2012.

Biodiversity loss and its impact on humanity. *Nature* **486**:59–67.

Chen Y, Huang Y, Niklaus PA, Castro-Izaguirre N, Clark AT, Bruelheide H, … Schmid B. 2020.

Directed species loss reduces community productivity in a subtropical forest biodiversity

experiment. *Nature Ecology & Evolution* **4**:550–559.

Carroll IT, Cardinale BJ, Nisbet RM. 2011. Niche and fitness differences relate the maintenance

of diversity to ecosystem function. *Ecology* **92**:1157–1165.

Chung PW, Livesley SJ, Rayner JP, Farrell C. 2021. Rooting volume impacts growth, coverage

and thermal tolerance of green façade climbing plants. *Land* **10**:1281.

Connolly J, Wayne P, Bazzaz FA, Gaines AESD. 2001. Interspecific competition in plants: How

well do current methods answer fundamental questions? *The American Naturalist*

157:107–125.

de Wit CT. 1960. On competition. *Verslagen Landbouwkundige Onderzoekigen* **66***:1–82.

Drake JM. 2003. Why does grassland productivity increase with species richness?

Disentangling species richness and composition with tests for overyielding and

superyielding in biodiversity experiments. *Proceedings of the Royal Society of London.*

Series B **270**:1713–1719.

Fargione J, Tilman D, Dybzinski R, Lambers JHR, Clark C, Harpole WS, … Loreau M. 2007.

From selection to complementarity: shifts in the causes of biodiversity–productivity

relationships in a long-term biodiversity experiment. *Proceedings of the Royal Society of*

London. Series B **274***:871–876.

Feng Y, Schmid B, Loreau M, Forrester DI, Fei S, Zhu J, … Fang J. 2022. Multispecies forest

plantations outyield monocultures across a broad range of conditions. *Science* **376***:*865–

868.

Forrester DI, Pretzsch H. 2015. Tamm Review: On the strength of evidence when comparing

ecosystem functions of mixtures with monocultures. *Forest Ecology and Management*

356:41–53.

Gaudet CL, Keddy PA. 1988. A comparative approach to predicting competitive ability from

plant traits. *Nature* **334***:*242–243.

Godoy O, Gómez-Aparicio L, Matías L, Pérez-Ramos IM, Allan E. 2020. An excess of niche

differences maximizes ecosystem functioning. *Nature Communications* **11**:4180.

Goldberg DE, Barton AM. 1992. Patterns and consequences of interspecific competition in

natural communities: a review of field experiments with plants. *The American Naturalist*

139*:*771–801.

Grace JB. 1990. On the relationship between plant traits and competitive ability. *InPerspectives*

on plant competition. Academic Press, New York, p. 51–66.

Grime J.P. 1973. Competitive exclusion in herbaceous vegetation. *Nature* **242**:344–347.

Hagan JG, Vanschoenwinkel B, Gamfeldt L. 2021. We should not necessarily expect positive

relationships between biodiversity and ecosystem functioning in observational field data.

Ecology Letters **24**:2537–2548.

Hector A. 1998. The effect of diversity on productivity: detecting the role of species

complementarity. *Oikos* **82***:*597–599.

Holland NJ, DeAngelis DL. 2009. Consumer-resource theory predicts dynamic transitions

between outcomes of interspecific interactions. *Ecology Letters* **12***:*357–1366.

Hooper DU, Chapin III FS, Ewel JJ, Hector A, Inchausti P, Lavorel S, … Wardle DA. 2005.

Effects of biodiversity on ecosystem functioning: a consensus of current knowledge.

Ecological Monograph **75**:3–35.

Huang S, Titus SJ, Lakusta TW. 1994. *Ecologically based site index curves and tables for major*

Alberta tree species. Forest Management Division, Land and Forest Resources, Alberta

Environmental Protection, Rep. T/307.

Huang S, Meng SX, Yang Y. 2009. *Growth and yield simulation system (GYPSY) for natural*

and post-harvest stands in Alberta. Edmonton, Alberta: Alberta Sustainable Resource

Development.

Huston MA. 1997. Hidden treatments in ecological experiments: re-evaluating the ecosystem

function of biodiversity. *Oecologia* **110**:449–460.

Isbell F, Gonzalez A, Loreau M, Cowles J, Díaz S, Hector A, … Larigauderie A. 2017. Linking

the influence and dependence of people on biodiversity across scales. *Nature* **546**:65–72.

Jaillard B, Richon C, Deleporte P, Loreau M, Violle C. 2018. An *a posteriori* species clustering

for quantifying the effects of species interactions on ecosystem functioning. *Methods in*

Ecology and Evolution **9**:704–715.

Jaillard B, Deleporte P, Isbell F, Loreau M, Violle C. 2021. Consistent functional clusters

explain the effects of biodiversity on ecosystem productivity in a long-term experiment.

Ecology **102**:e03441.

Kirwan L, Connolly J, Finn JA, Brophy C, Lüscher A, Nyfeler D, Sebastia MT. 2009.

Diversity–interaction modeling: estimating contributions of species identities and

interactions to ecosystem function. *Ecology* **90**:2032–2038.

Lieffers VJ, Macdonald SE, Hogg EH. 1993. Ecology of and control strategies for

Calamagrostis canadensis in boreal forest sites. *Canadian Journal of Forest Research*

23*:*2070-2077.

Loreau M. 2000. Biodiversity and ecosystem functioning: recent theoretical advances. *Oikos* **91**:

3–17.

Loreau M, Hector A. 2001. Partitioning selection and complementarity in biodiversity

experiments. *Nature* **412***:*72–76.

Loreau M, Sapijanskas J, Isbell F, Hector A. 2012. Niche and fitness differences relate the

maintenance of diversity to ecosystem function: comment. *Ecology* **93**:1482–1487.

Loreau M, Hector A. 2019. Not even wrong: Comment by Loreau and Hector. *Ecology*

100:e02794.

Mahaut L, Fort F, Violle C, Freschet GT. 2020. Multiple facets of diversity effects on plant

productivity: Species richness, functional diversity, species identity and intraspecific

competition. *Functional Ecology* **34**:287–298.

Man R, Lieffers VJ. 1999. Are mixtures of aspen and white spruce more productive than single

species stands? *Forestry Chronicle* **75**:505–513.

Montazeaud G, Violle C, Fréville H, Luquet D, Ahmadi N, … Fort F. 2018. Crop mixtures:

does niche complementarity hold for belowground resources? An experimental test using

rice genotypic pairs. *Plant and Soil* **424**:187–202.

Montès N, Maestre FT, Ballini C, Baldy V, Gauquelin T, Planquette M, … Perret JB. 2008. On

the relative importance of the effects of selection and complementarity as drivers of

diversity-productivity relationships in Mediterranean shrublands. *Oikos* **117**:1345–1350.

Murray Jr BG. 1986. The structure of theory, and the role of competition in community dynamics.

Oikos **46**:145-158.

Petchey OL. 2003. Integrating methods that investigate how complementarity influences

ecosystem functioning. *Oikos* **101***:323–330.

Peterson EB, Peterson NM. 1992. *Ecology, management, and use of aspen and balsam poplar*

in the prairie provinces. Edmonton, Alberta: Forestry Canada.

Pillai P, Gouhier TC. 2019 Not even wrong: The spurious measurement of biodiversity's effects

on ecosystem functioning. *Ecology* **100**:e02645

Polley HW, Wilsey BJ, Dernerk JD. 2003. Do species evenness and plant density influence the

magnitude of selection and complementarity effects in annual plant species mixtures?

Ecology Letters **6**:248–256.

Roscher C, Schumacher J. 2016. Positive diversity effects on productivity in mixtures of arable

weed species as related to density-size relationships. *Journal of Plant Ecology* **9**:792–804.

Sackville Hamilton NR. 1994. Replacement and additive designs for plant competition studies.

Journal of Applied Ecology **31**:599–603.

Špaèková I, Lepš J. 2001. Procedure for separating the selection effect from other effects in

diversity–productivity relationships. *Ecology Letters* **4***:585–594.

Tatsumi S, Loreau M. 2023. Partitioning the biodiversity effects on productivity into density and

size components. *Ecology Letters* **26**:ele.14300

Tilman D, Lehman CL, Thomson KT. 1997. Plant diversity and ecosystem productivity:

theoretical considerations. *Proceedings of the National Academy of Sciences, USA*

94:1857–1861.

Tilman D, Isbell F, Cowles JM. 2014. Biodiversity and ecosystem functioning. *Annual Review of*

Ecology, Evolution, and Systematics **45***:*471–93.

Turnbull LA, Levine JM., Loreau M, Hector A. 2013. Coexistence, niches and biodiversity

effects on ecosystem functioning. *Ecology Letters* **16**:116–127.

Wardle DA. 1999. Is ''sampling effect'' a problem for experiments investigating biodiversity–

ecosystem function relationships? *Oikos* **87**:403–407.

Watkinson AR. 1980. Density-dependence in single-species populations of plants. *Journal of*

Theoretical Biology **83**:345–357.

Weiner J, Freckleton RP. 2010. Constant final yield. *Annual Review of Ecology, Evolution, and*

Systematics **41**:173–192.

Figure 1. Under the competitive hypothesis, more competitive species gain size (biomass or
volume) and less competitive species lose size in mixture relative to their full density
monocultures. The magnitude of competitive growth responses (i.e., proportional changes in
individual size) increases with relative competitive ability (proportional deviations of species
competitive ability from community average) from a minimum of 0 (at community average) to a
maximum of $W_p/W_f - 1$ for more competitive species and -1 for less competitive species in
competitive exclusion. The changes may not be linear, greater near community average and
smaller with deviations of competitive ability (Gaudet & Keddy, 1988) or recourse availability
(e.g., light, see Brüllhardt *et al.* 2020) away from community averages. W_f represents for
individual size in full density monocultures and W_p represents for individual size in partial
density monocultures.

Figure 2. Partitioning net biodiversity effects (changes in stand volume in mixtures relative to monocultures) with competitive (a and b) and additive (c and d) partitioning models for 20- to 80-year-old mixed trembling aspen and white spruce with varying species compositions from nearly pure aspen (90% aspen) to nearly pure spruce (90% spruce) on average medium productivity sites in western Canada. The growth and yield data (***Table 1***) are generated with GYPSY (Huang et al., 2009) and calculations of biodiversity components with this figure are detailed in ***Table S1***.

604 **Table 1.** Simulated and experimental species mixtures. BPM (biodiversity effects of partial density monocultures) are calculated from
605 the differences between the partial density monoculture yield of more competitive species and total yield expected from species
606 relative yields and full density monoculture yields (see competitive exclusion in *Table S1* and *Table S2* for detailed calculations).

Species mixture (species1:species2)	Age	Total density	Mixture composition		Full density monoculture yields		Observed yields in mixtures		Partial density monoculture yields		
			Species1	Species2	Species1	Species2	Species1	Species2	Species1	Species2	BPM
Simulated mixed trembling aspen and white spruce in western Canada generated with GYPSY model (Huang et al., 2009) (density and stand volume yield, stems and m^3 per hectare)											
Populus:Picea	20 years	11,000	0.9	0.1	23.7	0.7	22.3	0.1	23.3	0.5	1.9
Populus:Picea	20 years	11,000	0.7	0.3	23.7	0.7	19.0	0.2	22.5	0.6	5.7
Populus:Picea	20 years	11,000	0.5	0.5	23.7	0.7	15.1	0.3	21.4	0.6	9.2
Populus:Picea	20 years	11,000	0.3	0.7	23.7	0.7	10.6	0.4	19.8	0.7	12.2
Populus:Picea	20 years	11,000	0.1	0.9	23.7	0.7	5.0	0.5	16.9	0.7	13.9
Populus:Picea	40 years	3,600	0.9	0.1	110.4	27.2	104.5	3.4	108.3	15.5	6.3
Populus:Picea	40 years	3,600	0.7	0.3	110.4	27.2	90.3	7.6	103.5	20.7	18.1
Populus:Picea	40 years	3,600	0.5	0.5	110.4	27.2	72.4	12.1	97.0	23.4	28.2
Populus:Picea	40 years	3,600	0.3	0.7	110.4	27.2	49.6	17.5	87.4	25.3	35.2
Populus:Picea	40 years	3,600	0.1	0.9	110.4	27.2	20.6	23.9	70.6	26.7	35.1
Populus:Picea	60 years	1,800	0.9	0.1	189.7	88.3	180.5	9.6	186.1	47.0	6.6
Populus:Picea	60 years	1,800	0.7	0.3	189.7	88.3	158.1	22.4	177.3	64.3	18.0
Populus:Picea	60 years	1,800	0.5	0.5	189.7	88.3	128.1	38.1	164.8	73.7	25.8
Populus:Picea	60 years	1,800	0.3	0.7	189.7	88.3	85.4	58.5	144.8	80.5	26.1
Populus:Picea	60 years	1,800	0.1	0.9	189.7	88.3	28.6	83.0	109.2	85.9	10.8
Populus:Picea	80 years	1,100	0.9	0.1	240.2	143.1	229.6	13.7	235.6	73.3	5.1
Populus:Picea	80 years	1,100	0.7	0.3	240.2	143.1	203.7	33.0	224.1	102.0	13.1
Populus:Picea	80 years	1,100	0.5	0.5	240.2	143.1	167.3	58.2	207.6	117.8	15.9
Populus:Picea	80 years	1,100	0.3	0.7	240.2	143.1	110.3	94.0	179.1	129.4	6.8
Populus:Picea	80 years	1,100	0.1	0.9	240.2	143.1	26.2	139.6	124.5	138.9	-28.4

Experimental grassland mixtures (density and aboveground biomass yield, stems and grams per pot) (Mahaut et al., 2020)											
Bromus: Dactylis	13 weeks	6	0.5	0.5	5.5	9.8	2.6	8.6	5.3	11.5	3.8
Bromus: Lotus	13 weeks	6	0.5	0.5	5.5	6.3	3.8	3.2	5.3	4.7	-1.2
Bromus:Plantago	13 weeks	6	0.5	0.5	5.5	8.6	1.9	10.9	5.3	16.1	9.0
Bromus:Sanguisorba	13 weeks	6	0.5	0.5	5.5	4.4	2.9	2.2	5.3	3.6	0.3
Bromus:Trifolium	13 weeks	6	0.5	0.5	5.5	13.3	3.8	9.3	5.3	14.2	4.8
Dactylis:Lotus	13 weeks	6	0.5	0.5	9.8	6.3	11.8	2.0	11.5	4.7	3.4
Dactylis:Plantago	13 weeks	6	0.5	0.5	9.8	8.6	4.4	9.5	11.5	16.1	2.3
Dactylis:Sanguisorba	13 weeks	6	0.5	0.5	9.8	4.4	6.0	1.9	11.5	3.6	4.4
Dactylis:Trifolium	13 weeks	6	0.5	0.5	9.8	13.3	7.5	7.7	11.5	14.2	2.6
Lotus:Plantago	13 weeks	6	0.5	0.5	6.3	8.6	1.0	16.2	4.7	16.1	8.6
Lotus:Sanguisorba	13 weeks	6	0.5	0.5	6.3	4.4	1.9	2.6	4.7	3.6	-0.6
Lotus:Trifolium	13 weeks	6	0.5	0.5	6.3	13.3	2.4	9.2	4.7	14.2	4.4
Plantago:Sanguisorba	13 weeks	6	0.5	0.5	8.6	4.4	11.6	1.1	16.1	3.6	9.6
Plantago:Trifolium	13 weeks	6	0.5	0.5	8.6	13.3	10.2	6.9	16.1	14.2	3.3
Sanguisorba:Trifolium	13 weeks	6	0.5	0.5	4.4	13.3	1.9	13.4	3.6	14.2	5.3

607 Species codes: Populus - *Populus tremuloides,* Picea - *Picea glauca*, Bromus - *Bromus erectus,* Dactylis - *Dactylis glomerata*, Lotus -
608 *Lotus corniculatus*, Plantago - *Plantago lanceolata*, Sanguisorba - *Sanguisorba minor*, and Trifolium - *Trifolium repens*.

610   **Table 2.** Partitioning net biodiversity effects (changes in aboveground biomass, g) with

611   competitive and additive partitioning models using experimental data (***Table 1***) from a grassland

612   diversity–productivity study (Mahaut et al., 2020). Calculations are detailed in ***Table S2***.

Species mixture	Competitive partitioning		Additive partitioning	
0.50Species1: 0.50Species2	Positive effect	Competitive/ negative effect	Complementarity effect (CE)	Selection effect (SE)
Bromus: Dactylis	1.8	1.7	2.7	0.9
Bromus: Lotus	1.2	-0.1	1.1	-0.1
Bromus:Plantago	2.5	3.2	4.3	1.4
Bromus:Sanguisorba	0.1	0.1	0.1	0.0
Bromus:Trifolium	0.9	2.8	3.6	0.0
Dactylis:Lotus	4.5	1.2	4.2	1.6
Dactylis:Plantago	4.3	0.3	5.1	-0.4
Dactylis:Sanguisorba	0.0	0.8	0.3	0.5
Dactylis:Trifolium	3.0	0.7	4.0	-0.3
Lotus:Plantago	7.6	2.3	7.9	1.9
Lotus:Sanguisorba	0.0	-0.9	-0.6	-0.3
Lotus:Trifolium	0.0	1.8	0.8	1.1
Plantago:Sanguisorba	1.5	4.7	3.9	2.3
Plantago:Trifolium	5.0	1.2	7.8	-1.6
Sanguisorba:Trifolium	2.9	3.6	3.9	2.6
Average	2.3	Competitive: 1.7 Negative: -0.2	3.3	0.6

613   Species codes: See ***Table 1***.