Carbohydrate Hydrolytic Activity, Antibiotic Resistance and Stress Tolerance of *Lacticaseibacillus paracasei* BCRC-16100 and *Lacticaseibacillus paracasei* ZFM54 for Probiotics Using Genomic and Biochemical Approaches

[1,2,3,4]**Peter James Icalia Gann***, [1,6]Jimmbeth Zenila P. Fabia, [1,3]Althea Gay B. Pagurayan, [1,6]Ma. Joy Theresa Agcaoili, [6]Ryan James Pascual, [1,6]Suerte M. Baranda, [1,6]Arc Josam J. Racho, [1,6]Marvielyn P. Olivar, [1,5]Jayson F. Cariaga, [3,5]Alvin Domingo, [6]Dionisio S. Bucao, and [1,3]Shirley C. Agrupis

[1]Genomics and Genetic Engineering Laboratory, College of Medicine, Mariano Marcos State University, City of Batac, Ilocos Norte, Philippines; [2]Department of Biochemistry and Molecular Biology, College of Medicine, Mariano Marcos State University, City of Batac, Ilocos Norte, Philippines; [3]Department of Biology, College of Arts and Sciences, Mariano Marcos State University, City of Batac, Ilocos Norte, Philippines; [4]Department of Agricultural Sciences, College of Agriculture, Food, and Sustainable Development, Mariano Marcos State University, City of Batac, Ilocos Norte, Philippines; [5]National Bioenergy Research and Innovation Center, Mariano Marcos State University, City of Batac, Ilocos Norte; and [6]Research Directorate, Mariano Marcos State University, City of Batac, Ilocos Norte, Philippines

*Corresponding Author
Email: pjicalia@gmail.com

Data Availability Statement: Whole genome sequences are available upon request from RJP and other data including gene expression, microbiological and biochemical assays from PJIG.

Funding Statement: The funding is from the Commission on Higher Education in the Philippines grant number LAKAS2022-005.

Conflict of Interest: The authors have declared no conflict of interest.

ABSTRACT

Probiotics are live microorganisms infused in food products or beverages that offer several health benefits in the body including acceleration of nutrient digestion and improvement of gut microbiota. In addition to the potential health benefits, it is also essential to ensure the safety of probiotics incorporated in food. Here, we present evidence of the ability of two probiotic isolates, *Lacticaseibacillus paracasei* BCRC-16100 and *Lacticaseibacillus paracasei* ZFM54, in the (1) enhancement of carbohydrate digestion, (2) tolerance to a range of drug and food processing conditions, and (3) antibiotic resistance and finally propose corresponding molecular mechanisms. Approaches include predictions using whole genome sequence (WGS) analysis, carbohydrate hydrolytic activity, and antibiotic resistance. WGS predictions suggested the ability of the two isolates to increase digestion of macronutrients particularly starch and antibacterial activity against opportunistic pathogens. The ability to facilitate carbohydrate hydrolysis was confirmed through a biochemical assay where the two isolates when grown alongside with starch causes release of glucose. However, the two isolates did not show antibacterial activities against pathogenic *S. aureus*, *S. epidermidis*, *E. coli*, *E. faecalis*, and *P. aeruginosa*. In terms of safety, WGS revealed the presence of antibiotic resistance (AbR) genes with different mechanisms, permease, and efflux pump, contributing to multidrug resistance in both isolates which were validated by their resistance against six (6) antibiotics including gentamicin, clindamycin, erythromycin, streptomycin, ofloxacin and vancomycin at 24 hours but susceptibility at 48 hours was observed in five (5) antibiotics except vancomycin. Furthermore, promoter and transposable element analysis showed the most active and highly transferable AbR genes in each of the isolates. Altogether, we show the potential of two probiotic isolates to be incorporated in food products for improved carbohydrate digestion and the need to address the removal of active and mobile antibiotic resistance genes that may compromise safety.

Keywords: probiotics; nutrient digestion*;* antibiotic resistance; antibacterial activity; whole genome sequence

Significance Statement: Probiotic isolates from nipa sap, *Lacticaseibacillus paracasei* BCRC-16100 and *Lacticaseibacillus paracasei* ZFM54, can potentially improve digestibility of carbohydrates and are tolerant to certain conditions common in drug development and food processing, however, the genome of the two isolates show antibiotic resistance genes that require understanding of the mechanisms to employ safety strategies prior to use for product development.

INTRODUCTION

In the last ten years, there has been a significant progress in probiotic research with many studies suggesting the essential role of probiotics in maintaining human health. Probiotics is increasingly gaining a huge popularity and share in the market due to potential health benefits such as improved bioavailability of macronutrients, changing the diversity of the gut microbiota and treatment of several diseases. Microbes under the genera *Pediococcus, Lactococcus, Enterococcus, Streptococcus, Propionibacterium*, and *Bacillus* are generally considered as probiotics provided that they qualify screenings for function and safety (de Brito Alves *et al*, 2016 and Hamad *et al*, 2022).

Several studies have shown that probiotics can improve digestion of carbohydrates such as lactose (Roobab *et al*, 2020) where lactose intolerant patients with supplemented with *lactobacilli* or *bifidobacteria* in their diet showed improved bioavailability and digestion of lactose (Lin *et al*, 1991; Kim *et al*, 1983; Lin *et al*, 1998; and Vitellio *et al*, 2019). Moreover, probiotics can also facilitate digestion of resistant starch such as fiber. When fiber passes through the stomach and small intestine undigested, the presence of *Bifidobacteria* in the large intestine can catalyze its hydrolysis (De Vuyst *et al*, 2021; Turroni *et al*, 2018; Rivière *et al*, 2016). There are also evidences on the enhanced protein digestion and absorption with probiotics which includes multiple mechanisms such as production of proteases, regulation of microflora to favor microbes with peptidase activities, and improved absorption of peptides and amino acids through accelerated transport in the intestine. The feasibility of lactobacilli derived proteases have been shown to catalyze hydrolysis of complex polypeptide chains (Jisha *et al*, 2013). Some of the most common protein hydrolytic enzymes in microbes are aspartic proteases, cysteine proteases, metalloproteases, and serine proteases (Rao *et al*, 1998). These findings indicate the potential of probiotics to facilitate utilization of nutrients from food sources with low digestibility which consequently improves digestion and overall nutritional health.

In addition to the digestive benefits, probiotics can also alter the gut microbiome favoring groups of microbial flora that can improve health in three ways including (1) modifying the internal host environment, (2) change in the composition of the gut microbiome, and (3) inhibition of pathogen proliferation. First, a shift in the diversity of the gut microflora also changes the host gut internal environment through increase in the amount of specific metabolic compounds. Metabolites from probiotics including organic acids, short-chain fatty acids, teichoic acids, peptidoglycans, plasmalogens, neurotransmitters, biosurfactants, amino acids, and flavonoid-derived compounds such as desaminotyrosine, equol daidzein, noratirriol, terpenoids, and phenolic compounds has been found to cause positive effects on consumer health (Erginkaya *et al*, 2022; Nataraj *et al*, 2020; and Stiles *et al*, 1996). Second, observation animal models suggests that composition of the gut microbiome changes with probiotic supplementation (Zhao *et al*, 2022). A wide range of studies have shown that a perturbed microbial community in the gut can be restored using probiotics as it has been associated with competitive exclusion of pathogen binding (Gareau *et al*, 2010; Dowarah *et al,* 2016; and Hooper *et al*, 2001). This ability of lactic acid bacteria in modulating the gut microbiota is due to the production of antimicrobial peptides (AMPs) such as bacteriocins which function as a natural bacterial immune defense system that can be combined with the corresponding receptor of both narrow and broad spectrum microbiota, and other metabolites such as lactic acid, which are harmful to several pathogenic strains of microorganisms (Yang *et al*, 2014; Servin AL, 2004; Konings *et al*, 2000; and Cotter *et al*, 2005), thus, bacteria which produce lactic acid as end product of carbohydrate fermentation are generally recognized as safe (GRAS) by Food and Drug Administration (FDA). Scant nutrients in the environment trigger the production of bacteriocins for competition of space and resources, excluding potential pathogens in gut microbiome, thus altering the composition of intestinal microbiota. Lastly, alteration of the gut microbiome and modification of the host internal environment with the consumption of probiotics consequently treats gastrointestinal diseases in controlled clinical trials (Preidis and Versalovic, 2009 and Thomas and Versalovic, 2010) and improves disease susceptibility (Frank *et al*, 2011).

Probiotics may provide multiple health benefits but it is equally essential that microbes used for these purposes are non-pathogenic, non-toxic, and versatile to remain stable and viable for long periods of storage and harsh conditions for incorporation in both functional food and dietary supplements and drugs (Yang *et al*, 2014), and generally safe for the intended use and meets standards for purity, identity, and potency. Scrutiny of pathogenicity, physiological and metabolic activities, and intrinsic properties are required. Comprehensive assessment of probiotic strain intended to be utilized in foods and human supplements, despite the variation in regulatory requirements in each country must be highly considered (Roe *et al*, 2022) through ways including (1) assessment of physiological activities, (2) strain identity, and (3) antibiotic resistance. First, the industrial production of probiotic strains should be based on the properties of the strains involved and their ability to withstand stress during processing and storage (Fossi *et al*, 2017). Second, unambiguous identification of the probiotic strains of interest should be considered and whole genome sequence (WGS) is used for this purpose. Lastly, only probiotics strains which do not have antibiotic resistant (AbR) genes should be selected for use in food and supplements (Ouwehand *et al*, 2016). Phenotypic assessment of antibiotic resistant genes comprised of testing the probiotic strain to a set of clinically important antibiotics, and determination of colony forming units for each antibiotic exposure with respect to the original log of the strain culture. Annotated genome sequence of lactic acid bacteria is utilized to inspect genetic elements responsible for metabolic activities, physiological activities, hemolytic activity, toxins, and antibiotic resistance. The horizontal transfer capability of antibiotic resistance is further determined by gene identification either on a plasmid or in near proximity to mobile elements such as to transposases. The vancomycin-resistance phenotype present in several strains of *Lactobacillus* is one of the most used examples of intrinsic AbR which is attributed to properties of the cell wall preventing the binding of the antibiotic (Gueimonde *et al*, 2013). Based on EFSA in 2012, any functional antibiotic resistance genes found within the genome of probiotic strain should be characterized as intrinsic or transmissible, which will further determine the risk of potential spread, which is a major public-health concern nowadays (Jindal *et al*, 2015). Countries with modern regulatory structures utilize advanced technology in establishing probiotic safety profiles, with greater emphasis on whole genome sequencing, and relying less on *in-vivo* testing.

Candidate probiotic strains are ubiquitous in nature. They emanate from human and animal origins such as gastrointestinal tract and breast milk, various food biotopes like fermented food products and dairy products, as well as from different parts of plants (Zommitti *et al*, 2020). Abundant sources of potential probiotic *Lactobacillus* strains were isolated from fermented food product of plant origin (Zielinska *et al*, 2015). This broad range of inexhaustible sources provide a challenge for the lactic acid bacteria strains to adapt to various environments, and these abilities vary significantly among species and even at a strain level (Morales *et al*, 2020), hence a significant variation in their genomic and metabolomic profiles as well. This variation arises through processes such as genetic mutation, horizontal gene transfer, leading to variations in traits like phenotypic, metabolism, and antibiotic resistance. Additionally, environmental factors such as but not limited to nutrient availability, temperature, pH, and competition with other microorganisms exert selective pressures shaping bacterial populations, further contributing to the diversity in bacterial characteristics across various sources (Nguyen *et al*, 2021).

It is evident that probiotics is becoming increasingly popular to consumers as it offers multiple health benefits. This trend brings in the need to isolate probiotics from different sources, ensure safety and evaluate functional use when used for supplementation. Here, we elucidate the potential benefit of two probiotic isolates from nipa sap in improving digestion of carbohydrates and proteins. Moreover, we explain bottlenecks on safety with emphasis on antibiotic resistance and the underlying biomolecular mechanisms. Finally, we provide insights to work around the known limitations for safety of the two isolates.

RESULTS

Phenotypic and Genomic Characteristics and Identification of Isolates

There were two lactic acid bacteria isolated from nipa sap which were both subjected to phenotypic characterization and identification (**Figure 1a-c**). Both bacterial isolates were gram-positive with the shape of a bacilli. The two isolates showed negative reaction to 3% H_2O_2 and exhibited γ hemolysis as there is no formation of clear, greenish, or opaque zones around the colonies (**Figure 1a**). These phenotypes have been identified to be common in lactic acid bacterial isolates (Yasmin *et al*, 2020).

Through this, it is inferred that these putative lactic acid bacteria are safe for further utilization as probiotics that can be incorporated in drugs or functional food. The colonies of the two isolates were translucent, round, shiny and convex (**Figure 1b**) which are all consistent to the characteristics of a lactic acid bacteria described from other studies (Siregar *et al*, 2014; Manzoor *et al*, 2016; Mahulette *et al*, 2018; Rahayu & Qurbaniah, 2019; Nasution *et al*, 2020; and Nasri *et al*, 2021). Using 16s rRNA, the isolates were identified as *Lacticaseibacillus paracasei* BCRC-16100 and *Lacticaseibacillus paracasei* ZFM54 with 100% homologies from the database sequences with accession numbers NZ_CP086132 and NZ_CP032637, respectively (**Figure 1c**). Despite genotypic similarities, alignment of genome sequences showed variability among the isolates and database hits (**Supporting Table 11**). Visualization of the whole genome of *Lacticaseibacillus paracasei* BCRC-16100 shows 2,885 contigs (3,029,123 bp) and *Lacticaseibacillus paracasei* ZFM54 shows 2,945 contigs (3,015,887 bp). In addition to the difference in genome length, the two isolates also showed variability in coding sequences on forward strand, genomic scaffolds, coding sequences on reverse strand, GC content, and GC skew (**Figure 2a-b**).

Figure 1. Phenotypic characteristics and identification of lactic acid bacterial (LAB) isolates from nipa sap, *Lacticaseibacillus paracasei* BCRC-16100 and *Lacticaseibacillus paracasei* ZFM54. (**a**) Tests for safety in the two isolates indicating gram positive, hemolysin negative, catalase negative for both, (**b**) observable physical appearance and (**c**) identity using 16s rRNA. Gram Stain Test (Purple cell morphology = gram positive, pink cell morphology = gram negative); Hemolysin Test (yellow to translucent inhibition = β-hemolysis, translucent with greenish inhibition = α-hemolysis, no clearing zones = γ-hemolysis; Catalase Test (presence of bubbles = positive, absence of bubbles – negative).

Figure 2. Whole genome sequence (WGS) assembly and analysis of *Lacticaseibacillus paracasei* BCRC-16100 and *Lacticaseibacillus paracasei* ZFM54. (**a**) Visualization of regions of interest in the whole genome of *Lacticaseibacillus paracasei* BCRC-16100 showing 2,885 contigs (3,029,123 bp) and (**b**) Visualization of regions of interest in the whole genome of *Lacticaseibacillus paracasei* ZFM54 showing 2,945 contigs (3,015,887 bp). Circles from the outside to the center illustrate the following characteristics: (1) coding sequences on forward strand, (5) genomic scaffolds, (2) coding sequences on reverse strand, (3) GC content, and (4) GC skew. Genetic map is generated using ProkSee v1.0.0a6.

Tolerance in Different Conditions

The tolerance of *Lacticaseibacillus paracasei* BCRC-16100 and *Lacticaseibacillus paracasei* ZFM54 were tested in ranges of pH (3-7), alcohol concentration (5-10%), and temperature (25-50°C) in 24-hour incubation (**Figure 3a-c**). **Figure 3a** shows that *Lacticaseibacillus paracasei* ZFM54 grows in a wider range from 35-45°C with 45°C as the optimal temperature. The optimal temperature for growth *Lacticaseibacillus paracasei* BCRC-16100 is the same but there is a significant decline in growth for any deviation in the temperature (±1°C). However, it has been previously reported that the growth for different *Lactobacillus* spp. (*rhamnosus, paracasei, reuteri, plantarum* and *pentosus*) is optimal at 37 °C (Śliżewska and Chlebicz-Wójcik, 2020). This can be explained in the differences in the temperature points observed where the previous study observed growth in 4, 20, 30, 37, 44, and 55°C. In the different concentrations of alcohol tested (5, 10, 15 and 20%), growth for the two isolates were the same suggesting the ability of the isolates to survive when incorporated in functional food products or drugs with alcohol content ranging from 5-20% (**Figure 3b**). For both probiotic isolates, the growth at pH 5 and 7 were significantly higher than pH 3 and 4 (**Figure 3c**). This is consistent to the result of a previous study indicating that optimal pH for the growth of *Lactobacillus* spp. (*rhamnosus, paracasei, reuteri, plantarum* and *pentosus*) is 5.5 to 6.5 which may vary depending on the strain (Śliżewska and Chlebicz-Wójcik, 2020).

The differences in the frequency of the stress tolerance genes and variability in the type and number of transcription factors (TFs) binding to cis-regulatory elements offers an explanation in differential or similar growth response of the two isolates in a particular stressful condition (**Figure 4a-c**). The growth of *Lacticaseibacillus paracasei* ZFM54 is less influenced by temperature and higher frequency of TFs that may potentially bind to its promoter region to a thermoregulatory gene *dnaK* can play a significant role in this tolerance. A previous study has shown that in known thermotolerant *Bacillus pumilus*, *dnaK* is highly

expressed (Kumar *et al*, 2013). Although it cannot be ruled out that there might be differences in *dnaK* of different species and strains and other genes present in the genome of the two isolates play synergistic or antagonistic roles in tolerance to certain temperatures.

Figure 3. Tolerance of lactic acid bacterial isolates from nipa sap, *Lacticaseibacillus paracasei* BCRC-16100 and *Lacticaseibacillus paracasei* ZFM54. *Lacticaseibacillus paracasei* ZFM54 has wider range of tolerance across (**a**) temperature and (**b**) presence of alcohol while (c) *Lacticaseibacillus paracasei* BCRC-16100 performs better in a range of pH. Means within each bar (comparison within the same strain) having the same letter are not significantly different in Tukey's multiple comparison at $\alpha = 0.05$. and significant for condition as source of variation at $p \leq 0.01$. Error bars represent standard error within 3 biological replicates of each type of condition.

Figure 4. Promoter analysis of genes associated to (**a**) thermoregulation, (**b**) alcohol tolerance, and (**c**) pH response in *Lacticaseibacillus paracasei* BCRC-16100 and *Lacticaseibacillus paracasei* ZFM54. There is differential frequency in the number of tolerance genes and transcription factors binding to each gene responsible for stress response. Colors in the bars indicate the number of each type of transcription factors (TF) binding to a stress tolerance gene. Complete list of stress tolerance genes with the cis-regulatory elements where TFs bind is shown in Supporting Figure 11.

Antibiotic Resistance

The antibiotic resistance of the two probiotic isolates showed similar pattern in six (6) types of antibiotics both at 24- and 48-hours post-treatment (**Figure 5a-b**). In 24 hours, *Lacticaseibacillus paracasei* BCRC-16100 and *Lacticaseibacillus paracasei* ZFM54, showed complete resistance to clindamycin, vancomycin, gentamicin, ofloxacin, erythromycin, and streptomycin. However, at 48 hours post-treatment, five antibiotics (clindamycin, gentamycin, ofloxacin, erythromycin, and streptomycin) showed inhibition in the growth of two isolates except for vancomycin (**Figure 5a**). Differences in the zone of inhibition (mm) were significantly different (p=0.0005, α=0.05) in various antibiotics for each of the two isolates (**Figure 5b**). Higher zone of inhibition was observed in erythromycin while no inhibition was observed in vancomycin for the two isolates. The result on disk diffusion is corroborated by the colony forming unit quantity (CFU/ml) when the two isolates were independently grown in liquid broth infused with antibiotics for 48 hrs. The CFU for both isolates were significantly higher in vancomycin (BCRC-16100=6.330413773, and ZFM54=5.252853031) suggesting resistance in this antibiotic that also support results of the disk diffusion analysis. These analyses indicate differential resistance of *Lacticaseibacillus paracasei* BCRC-16100 and *Lacticaseibacillus paracasei* ZFM54 depending on the type of antibiotic and time of exposure. Moreover, it also suggests the preponderance of antibiotic resistance genes containing active cis-regulatory elements in these isolates promoting survivability in antibacterial agents.

Figure 5. Higher antibiotic resistance in *Lacticaseibacillus paracasei* ZFM54 compared to *Lacticaseibacillus paracasei* BCRC-16100 suggested by higher growth in six (6) types of antibiotics and expression of antibiotic resistant (AbR) genes. (**a**) disk diffusion assay, zone of inhibition in antibiotic disks and (**b**) and colony forming units (CFU) of the two isolates in different antibiotics at 12 hrs. From top clockwise in (a) the antibiotics are clindamycin, vancomycin, gentamycin, ofloxacin, erythromycin, and streptomycin. Means within each bar (comparison within the same strain) having the same letter are not significantly different in Tukey's multiple comparison at α = 0.05. and significant for type of antibiotic as source of variation at $p ≤ 0.01$. Error bars represent standard error within 3 biological replicates of each type of condition.

Genomic Analysis of Antibiotic Resistant (AbR) Genes

AbR genes are present in both probiotic isolates which differ in terms of frequency, activity, and genomic, and amino acid structure (**Figure 6a-d**). There are five (*bmr3, stp, lmrA, emrY, yheI*) and nine (*stp, yheI, yheH, bmr3, tetA_1, lmrA, tetA_2, tetO, marR*) AbR genes in *Lacticaseibacillus paracasei* BCRC-16100 and *Lacticaseibacillus paracasei* ZFM54, respectively (**Figure 6a**). Briefly, there are two antibiotic resistant mechanisms (transcriptional regulation and transporter activity) in *Lacticaseibacillus paracasei* BCRC-16100 while there are three in *Lacticaseibacillus paracasei* ZFM54 suggesting that the later can have a wider range of resistance to antibiotics (transcriptional regulation, transporter activity, and efflux system). In *Lacticaseibacillus paracasei* BCRC-16100, the antibiotic resistant genes are mostly responsible in encoding transporter protein subunits (*bmr3, stp, emrY, yheI*) and one is a transcriptional repressor (*lmrA*). The two isolates share four AbR genes namely *bmr3, stp, lmrA, yheI* and unique AbR genes in *Lacticaseibacillus paracasei* ZFM54 include *yheH, tetA_1, tetA_2, tetO,* and *marR* where three are efflux pumps (*tetA_1, tetA_2,* and *tetO*), one encodes a transcriptional regulator (*marR*) and another that encodes a transporter protein subunit (*yheH*).

Although four AbR genes are shared between *Lacticaseibacillus paracasei* BCRC-16100 and *Lacticaseibacillus paracasei* ZFM54, analysis of the homology of these four genes relative to each other including the rest of AbR genes indicate that there are differences. Among the four shared AbR genes in the two isolates, *yheI* and *stp* are the most diverged as suggested by the distance in their cluster or clades in the phylogenetic analysis (**Figure 6b**). Given the overlapping and diversity of similar AbR genes in the two isolates, genes responsible for antibiotic resistance were tested for differential gene expression analysis in the presence in vancomycin (**Figure 6c**), an antibiotic where both isolates showed resistance (**Figure 5a**). For both isolates, AbR genes encoding transcriptional regulators (*lmrA and marR*) are upregulated in

vancomycin treatment. This may suggest a similar mechanism for vancomycin resistance in *Enterococci* where vancomycin activates *VanR* which then cascades to the activation of other AbR genes (Guffey and Loll, 2021). Other AbR genes that are significantly upregulated in vancomycin are *emrY* and *yheI* which are encoding for transporters protein subunits and *tetA_1* encoding for an efflux pump. Among the AbR genes, only *emrY* has the highest probability to be transposed laterally due to the presence of transposable elements in the -54bp (5') and +637bp (3') of the gene (**Figure 6d**).

Figure 6. Promoter, phylogeny, gene expression, and transposable elements of antibiotic resistant (AbR) genes in *Lacticaseibacillus paracasei* BCRC-16100 and *Lacticaseibacillus paracasei* ZFM54. (**a**) AbR genes and frequency of each type of transcription factor (TF) binding in the promoter region (1000bp) in every gene. More resistant genes are found in the genome of *Lacticaseibacillus paracasei* ZFM54 and the frequency of TFs that can bind to each of the AbR genes promoter region vary. (**b**) Phylogeny of AbR genes showing differential clustering of some similar genes (*stp* and *bmr3*) shared by the two isolates. (**c**) Expression of AbR genes in the two isolates. (**d**) Transposable elements located in the upstream (5') and downstream (3') regions (500bp) of AbR genes. Colors in the bars indicate the number of each type of TF binding to a stress tolerance gene. Complete list of AbR genes with the cis-regulatory elements where TFs bind is shown in Supporting Figure 10. Relative expression is against the internal control (*16s rRNA*) is shown. Transcription Start Site (TSS). Error bars represent standard error within 3 biological replicates.

Carbohydrate Hydrolytic Activity

The two isolates were analyzed for ability to hydrolyze carbohydrates and TF binding to the promoter and expression of genes responsible for hydrolysis (**Figure 7**). In a starch assay, both isolates showed hydrolytic activity qualitatively (**Figure 7a**) and quantitatively (**Figure 7b**). There were multiple genes found in the WGS which maybe responsible for the carbohydrate hydrolytic activity having 8 and 11, in *Lacticaseibacillus paracasei* BCRC-16100 and *Lacticaseibacillus paracasei* ZFM54, respectively (**Figure 7c**). Notably, promoter analysis showed more TF binding to *bga*, *lacG* and *bglA* suggesting the importance of the three genes in carbohydrate analysis. However, gene expression analysis revealed that during carbohydrate hydrolysis, *rafA*, *malL*, and *bglC* are the genes that are upregulated. It can be argued that despite the presence of more cis regulatory elements in some carbohydrate hydrolysis genes, the observed lower expression can be explained as a result of positional effect of where the TF bind which was previously described to influence gene activity (ENCODE Project Consortium, 2012).

Figure 7. Carbohydrate hydrolytic activity of *Lacticaseibacillus paracasei* BCRC-16100 and *Lacticaseibacillus paracasei* ZFM54. (**a**) Amylase activities of BCRC-16100 and ZFM54 presented by zone of clearance, (**b**) the quantitative zone of clearance expressed in mm, (**c**) promoter analysis of carbohydrate hydrolysis genes, and (**d**) expression of amylases genes. Colors in the bars indicate the number of each type of transcription factors (TF) binding to a stress tolerance gene. Complete list of AbR genes with the cis-regulatory elements where TFs bind is shown in Supporting Figure 11. Relative expression is against the internal control (*16s rRNA*) is shown. Error bars represent standard error within 3 biological replicates.

DISCUSSION

Probiotics has been widely used for numerous health benefits such as improved gut microflora, enhance digestion of nutrients and reduction in the risk of various diseases (Lin, 2003 and Salminen *et al*, 2005). Despite the known positive health effects, it is also essential to ensure the (1) flexibility of probiotic isolates in different conditions to allow incorporation in drugs and functional food, (2) absence of transposable antibiotic resistance genes, (3) and bioactivity of any proposed benefit such as carbohydrate digestion. Probiotics can be isolated from animal and dairy sources where novel or similar genotypes obtained from a different source can have different characteristics (Abdou *et al*, 2018; Karami *et al*, 2017; and Haghshenas *et al*, 2017). However, efforts to isolate probiotics from plant sources are very limited (Menendez and Garcia-Fraile, 2017).

There are multiple indigenous plants in northern Philippines that can be instrumental as sources of probiotics due to their abundance in the region such as nipa. Two strains of the same species of probiotics can be isolated from this source namely *Lacticaseibacillus paracasei* BCRC-16100 and *Lacticaseibacillus paracasei* ZFM54. The genus of the two strains is classified as probiotics but there is very little to know understanding on their health benefits, tolerance to stress and safety. Moreover, the species of the two strains are also recognized as probiotics only known for ameliorating allergic airway, antimicrobial activity, enhancement of intestinal microbiota and stress modulator, among others (Lin *et al*, 2023; Hill *et al*, 2018; and Jones, 2017). The intrinsic heterogeneity across strains of *Lacticaseibacillus paracasei* makes the species a strong group for the selection of probiotics isolates for health benefits and development of drugs or functional food for the abovementioned purpose (Balzaretti et al, 2017). Although in recent years much effort has been made to study this, the answer is not conclusive, and there remains much to be elucidated.

As described above, the health-promoting traits from the species *Lacticaseibacillus paracasei* as probiotics has not been extensively explored and the strains isolated from nipa sap, *Lacticaseibacillus paracasei* BCRC-16100 and *Lacticaseibacillus paracasei* ZFM54, has not been characterized for its contribution to overall health, compatibility for drug and functional food development and absence of transposable antibiotic resistance elements.

To assess the potential of *Lacticaseibacillus paracasei* BCRC-16100 and *Lacticaseibacillus paracasei* ZFM54 as probiotic for commercial use, we carried out whole genome sequence, gene expression and biochemical analyses. There are several genes that confer thermotolerance in both isolates which allows growth in a wide range of temperature where *dnaK* and *dnaJ* are likely responsible for this trait these genes show the highest potential to be regulated due to the association of promoter regions to multiple transcription factors (TFs) (**Figure 4a**). Both *dnaK* and *dnaJ* belong to a chaperone system in microbes known to confer wider tolerance to temperatures (Kumar *et al*, 2013 and Kim *et al*, 2021). The two isolates grow optimally at pH 5-7, however, promoter analysis of genes responsible for this trait only suggested *clpB* as may be responsible in differential growth in pH. The gene *clpB* is highly conserved in bacteria to provide tolerance to oxidative stress, nutrient starvation and low pH (Meibom *et al*, 2008; Glaza *et al*, 2020; and Tripathi *et al*, 2020). In addition to flexibility to stressful conditions, we also explored AbR genes in each strain with the corresponding molecular mechanisms and transposability. The two strains were highly resistant to vancomycin and gene expression analysis suggests four genes conferring this resistance namely *lmrA, marR, emrY* and *yheI* and mechanisms of resistance include efflux, transcriptional control and regulation of transport (**Figure 6**). Moreover, these AbR genes possess transposable elements in the upstream (5') and downstream (3') region. This is a consideration in using these strains as probiotics to avoid potential lateral transfer of AbR genes to pathogenic microbes (Sieber *et al*, 2017). Lastly, high rates of available genes that encode carbohydrate active enzymes are detected in the genome data, which correlated the enzymatic hydrolytic activities evident *in-vitro*. Among the identified genes, *rafA*, *malL*, and *bglC*, are most likely to be responsible to the carbohydrate hydrolytic activity of *L. paracasei* BCRC-16100 and *L. paracasei* ZFM54 based on the promoter analysis of genes (**Figure 7**). Bacteria that colonize the

intestine collectively possesses a large repertoire of degradative enzymes and metabolic capabilities. To promote the development of human gastrointestinal nutrition and health is to regulate the host mucosal and balance intestinal microflora.

In conclusion, *Lacticaseibacillus paracasei* BCRC-16100 and *Lacticaseibacillus paracasei* ZFM54 can arguably be utilized as probiotics that can be incorporated in drugs and functional food to promote digestion of carbohydrates in food products with low digestibility with careful consideration of the antibiotic resistance in these strains.

EXPERIMENTAL PROCEDURES

Isolation and Screening of LAB. The de Man Rogosa Sharpe agar (Sigma-Aldrich Canada Co.) supplemented with 1% CaCO$_3$ was used as a selective medium for the cultivation and isolation of the two probiotic isolates from nipa sap. One hundred microliter of the serially diluted sample was aseptically inoculated in MRSA plate using spread plate method. The plates were incubated for 24 hrs at 37°C. Individual colonies with colony morphology of a putative lactic acid bacterium grown overnight was further subjected to standard purification of bacterial cultures. Purified putative LAB undergone biochemical test consists of staining test using gram staining solutions (Sigma-Aldrich 111885) and catalase test using 3% H2O2. Moreover, hemolysin test using Sheep Blood Agar (Remel TSA w/ 5% Sheep Blood) was conducted with gram positive and catalase negative putative LAB. Isolates which exhibited γ-hemolysis were stored and sent for molecular identification.

Identification of Isolates by Capillary Sequencing. Genomic DNA (gDNA) of S1 and S2 was extracted using the Quick-DNA fungal/Bacterial Miniprep Kit (Zymo Research, USA) according to manufacturer's protocol. PCR amplicons were subjected to purification using AMPure XP beads (Cat. No. 163881). One microliter of the purified PCR amplicons was loaded into 1% agarose gel run at 120 V for 45 min, with Invitrogen 1kb Plus DNA Ladder. Capillary sequencing involved the incorporation of fluorescently labeled chain terminator ddNTPs. The reaction components include amplicons, corresponding primers, and the ABI BigDye® Terminator v3.1 Cycle Sequencing Kit (Cat No. 4337455). The cycling parameters on the thermal cycler were as follows: pre-hold at 4 °C; 96 °C for 1 min; 25 cycles at 96°C for 10 s, 50°C for 5 s, 62°C for 4 min; and hold at 4 °C. Ethanol precipitation was performed to remove unincorporated ddNTPs, excess primers, and primer dimers. Capillary electrophoresis was carried out on the ABI 3730xl DNA Analyzer using a 50cm 96-capillary array, POP7 Polymer (Cat No. 4393714), and 3730xl Data Collection Software v3.1. Base calling was performed using the Sequencing Analysis Software v5.4.

Whole Genome Sequencing. Library preparation was performed using the TruSeq DNA Nano Kit (Illumina, USA), and sequencing was conducted using an Illumina MiSeq instrument and a paired end read format of 2 x 150bp for 300 cycles at the Philippine Genome Center, Quezon City, Philippines.

Antibiotic Resistance. The probiotic isolates were subjected to antibiotic susceptibility using commonly used antibiotics (BD BBL Sensi-Disc) from various classes with various pharmacological actions, namely, natural penicillins, glycopeptides, aminoglycosides, macrolides, lincosamides, and fluorquinolones. Kirby-Bauer technique (disc-diffusion) was employed where one hundred microliters of bacterial culture was inoculated in the plate containing sterile Mueller Hinton Agar (TM Media, India) and discs of Vancomycin (30μg), Clindamycin (2μg), Gentamycin (10μg), Ofloxacin (5μg), Erythromycin (15μg), and Streptomycin (300μg) were used as antibiotic wafers. Zone of Inhibition (ZOI) was measured after 48 hrs using digital caliper. Computed ZOI were categorized according to the accepted inhibition ranges of the three levels of susceptibility namely sensitive, intermediate, and resistant. Moreover, viable cells of the probiotic isolates suspended in Mueller Hinton Broth (TM Media, India) containing the antibiotic with known concentration was quantified to correlate the semi-quantitative data obtained from disc-diffusion method.

Carbohydrate Hydrolysis. *Lacticaseibacillus paracasei* BCRC-16100 and *Lacticaseibacillus paracasei* ZFM54 were qualitatively identified through the Seliwanoffs' assay using a solution of 0.05% resorcinol (Sigma-Aldrich, Missouri, United States) in 3 N HCl. The experimental control was 1 mL of the test sample (from broth with cultured sample) and 1 mL of distilled water as a blank, then 2mL of Seliwanoffs' reagent were added to controls with thorough mixing and were placed in a water bath for 1 minute. The formation of the cherry, red-colored complex indicated a positive result for ketose production.

Prediction of Promoter Elements. Bioinformatical tools such as BPROM and PRODORIC were used to predict the promoter elements involved in the expression of genes associated with antibiotic resistance, stress tolerance and hydrolytic activity of *Lacticaseibacillus paracasei* BCRC-16100 and *Lacticaseibacillus paracasei* ZFM54. The upstream regions of these genes were extracted from their whole genome sequences (WGS) and subjected to analysis in the BPROM website where results show possible – 10 and –35 boxes of predicted promoters, their positions in the submitted sequence along with possible transcription factors. These transcription factors were further analyzed in the PRODORIC website's virtual footprint in which their potential binding site with the highest relative score is shown in **Supporting Table 10-11**.

Gene Expression and Polymerase Chain Reaction. For gene expression, total RNA was isolated using Trizol (Invitrogen Inc.) and quantified using Nano-drop 2000 (Thermo-Fisher Inc). Two micrograms of total RNA were treated with RQ1-RNAse free DNase (Thermofisher Inc.), and one microgram of the DNase-treated RNA was used for cDNA synthesis using PrimeScript RT reagent kit (Takara Bio, CA, USA). The expression analysis was performed using TB green Premix Ex Taq II (Takara Bio, CA, USA) on Bio-Rad CFX 96 C1000 with following conditions: 95°C for 30 sec and 40 cycles of 95°C for 5 sec + 60°C for 30 sec. The product specificity was verified by the melt curve analysis. The Ct values of all the genes were normalized against *16s rRNA* fused protein as the reference gene. Primer sequences used for gene expression are in **Supporting Table 1**. The polymerase chain reaction (PCR) was performed on genomic DNA to amplify *16s rRNA* using Emerald Amp MAX PCR Master Mix (Takara Bio) using the primers presented in **Supporting Table 1**.

Data Analysis. Experiments subjected to statistical analysis were replicated three times. For bioactivity on macronutrient hydrolysis and other phenotypic characteristics between the two isolates, means were compared using studentized t-Test. Antibiotic resistance assays experiments were laid out in completely randomized design. Data were transformed with arcsine and was subjected to one-way ANOVA. Differences in the mean antibiotic resistance in different antibacterial drugs were determined using Tukey's multiple comparison test. Statistical analyses were performed in SAS statistical software (version 9.4, SAS Institute Inc.) and results are presented in **Supporting Tables 2-9**.

REFERENCES

Roobab U., Batool Z., Manzoor M. F., Shabbir M. A., Khan M. R., Aadil R. M. (2020). Sources, formulations, advanced delivery and health benefits of probiotics. *Curr. Opin. Food Sci.* 32, 17–28. doi: 10.1016/j.cofs.2020.01.003

Lin, M. Y., Savaiano, D., & Harlander, S. (1991). Influence of nonfermented dairy products containing bacterial starter cultures on lactose maldigestion in humans. *Journal of dairy science*, *74*(1), 87–95. https://doi.org/10.3168/jds.S0022-0302(91)78147-2

Kim, H. S., & Gilliland, S. E. (1983). Lactobacillus acidophilus as a dietary adjunct for milk to aid lactose digestion in humans. *Journal of dairy science*, *66*(5), 959–966. https://doi.org/10.3168/jds.S0022-0302(83)81887-6

Lin, M. Y., Yen, C. L., & Chen, S. H. (1998). Management of lactose maldigestion by consuming milk containing lactobacilli. *Digestive diseases and sciences*, *43*(1), 133–137. https://doi.org/10.1023/a:1018840507952

Vitellio, P., Celano, G., Bonfrate, L., Gobbetti, M., Portincasa, P., & De Angelis, M. (2019). Effects of *Bifidobacterium longum* and *Lactobacillus rhamnosus* on Gut Microbiota in Patients with Lactose Intolerance and Persisting Functional Gastrointestinal Symptoms: A Randomised, Double-Blind, Cross-Over Study. *Nutrients*, *11*(4), 886. https://doi.org/10.3390/nu11040886

Jisha, V.N., Smitha, R.B., Pradeep, S., Sreedevi, S., Unni, K.N., Sajith, S., Priji, P., Josh, M.S., & Benjamin, S. (2013). Versatility of microbial proteases. *Advances in Enzyme Research, 2013*, 39-51.

Rao, M. B., Tanksale, A. M., Ghatge M. S., and Deshpande V. V. (1998). Molecular and biotechnological aspects of microbial proteases. *Microbiol. Mol. Biol R* 62, 597–635. doi: 10.1128/MMBR.62.3.597-635.1998

De Vuyst, L., & Leroy, F. (2011). Cross-feeding between bifidobacteria and butyrate-producing colon bacteria explains bifdobacterial competitiveness, butyrate production, and gas production. *International journal of food microbiology*, *149*(1), 73–80. https://doi.org/10.1016/j.ijfoodmicro.2011.03.003

Turroni, F., Milani, C., Duranti, S., Mahony, J., van Sinderen, D., & Ventura, M. (2018). Glycan Utilization and Cross-Feeding Activities by Bifidobacteria. *Trends in microbiology*, *26*(4), 339–350. https://doi.org/10.1016/j.tim.2017.10.001

Rivière, A., Selak, M., Lantin, D., Leroy, F., & De Vuyst, L. (2016). Bifidobacteria and Butyrate-Producing Colon Bacteria: Importance and Strategies for Their Stimulation in the Human Gut. *Frontiers in microbiology*, *7*, 979. https://doi.org/10.3389/fmicb.2016.00979

de Brito Alves, J. L., de Sousa, V. P., Cavalcanti Neto, M. P., Magnani, M., Braga, V. A., da Costa-Silva, J. H., Leandro, C. G., Vidal, H., & Pirola, L. (2016). New Insights on the Use of Dietary Polyphenols or Probiotics for the Management of Arterial Hypertension. *Frontiers in physiology*, *7*, 448. https://doi.org/10.3389/fphys.2016.00448

Hamad, G., Ombarak, R. A., Eskander, M., Mehany, T., Anees, F. R., Elfayoumy, R. A., Sabrien A. O., José M. L., Abou-Alella, S.A. (2022). Detection and inhibition of *Clostridium botulinum* in some Egyptian fish products by probiotics cell-free supernatants as bio-preservation agents. *LWT* 163:113603. doi: 10.1016/j.lwt.2022.113603

Stiles M. E. (1996). Biopreservation by lactic acid bacteria. *Antonie van Leeuwenhoek*, *70*(2-4), 331–345. https://doi.org/10.1007/BF00395940

Nataraj, B. H., Ali, S. A., Behare, P. V., & Yadav, H. (2020). Postbiotics-parabiotics: the new horizons in microbial biotherapy and functional foods. *Microbial cell factories*, *19*(1), 168. https://doi.org/10.1186/s12934-020-01426-w

Erginkaya Z., Konuray-Altun G. (2022). Potential Biotherapeutic Properties of Lactic Acid Bacteria in Foods. *Food Biosci.* 46:101544. doi: 10.1016/j.fbio.2022.101544.

Zhao, N., Shu, Y., Jian, C., Zhou, Z., Bao, H., Li, X., Cheng, X., Zhao, Y., Jin, S., & Shu, X. (2022). *Lactobacillus* Ameliorates SD-Induced Stress Responses and Gut Dysbiosis by Increasing the Absorption of Gut-Derived GABA in Rhesus Monkeys. *Frontiers in immunology*, *13*, 915393. https://doi.org/10.3389/fimmu.2022.915393

Preidis, G. A., & Versalovic, J. (2009). Targeting the human microbiome with antibiotics, probiotics, and prebiotics: gastroenterology enters the metagenomics era. *Gastroenterology*, *136*(6), 2015–2031. https://doi.org/10.1053/j.gastro.2009.01.072

Thomas, C. M., & Versalovic, J. (2010). Probiotics-host communication: Modulation of signaling pathways in the intestine. *Gut microbes*, *1*(3), 148–163. https://doi.org/10.4161/gmic.1.3.11712

Frank, D. N., Zhu, W., Sartor, R. B., & Li, E. (2011). Investigating the biological and clinical significance of human dysbioses. *Trends in microbiology*, *19*(9), 427–434. https://doi.org/10.1016/j.tim.2011.06.005

Śliżewska, K., & Chlebicz-Wójcik, A. (2020). Growth Kinetics of Probiotic *Lactobacillus* Strains in the Alternative, Cost-Efficient Semi-Solid Fermentation Medium. *Biology*, *9*(12), 423. https://doi.org/10.3390/biology9120423

Siregar, M. T. P., Kusdiyantini, E., & Rukmi, M. I. R. (2014). Isolasi Dan Karakterisasi Bakteri Asam Laktat Pada Pangan Fermentasi Mandai. Jurnal Akademika Biologi, 3(2), 40–48. Retrieved from https://ejournal3.undip.ac.id/index.php/biologi/ article/view/19443

Manzoor, A., Ul-haq, I., Baig, S., Qazi, J. I., & Seratlic, S. (2016). Efficacy of Locally Isolated Lactic Acid Bacteria Against Antibiotic-Resistant Uropathogens. Jundishapur journal of microbiology, 9(1), 1–11. https://doi.org/10.5812/jjm.18952

Mahulette, F., Mubarik, N. R., Suwanto, A., & Widanarni. (2018). Microbiological and Physicochemical Characteristics of Inasua Traditional Fish Fermented from Maluku Islands. Biosaintifika: Journal of Biology & Biology Education, 10(2), 298–305. http://dx.doi.org/10.15294/biosaintifika.v10i2.135 37

Rahayu, H. M., & Qurbaniah, M. (2019). Selection of Tempoyak Lactic Acid Bacteria As Candidate Strain for Yoghurt Starter Culture. Biosaintifika. 11(111), 39–46. http://dx.doi.org/10.15294/biosaintifika.v11i1.167 69

Nasution, M. H. B., Ramadhani, S., & Fachrial, E. (2020). Isolation, Characterization and Antibacterial Activities of Lactic Acid Bacteria Isolated From Batak's Special Food "Dali Ni Horbo". Jurnal Natur Indonesia, 18(1), 1. https://doi.org/10.31258/jnat.18.1.1-11

Nasri, Harahap, U., Silalahi, J., & Satria, D. (2021). Antibacterial activity of lactic acid bacteria isolated from dengke naniura of carp (Cyprinus carpio) against diarrhea-causing pathogenic bacteria. Biodiversitas, 22(8), 3098–3104. https://doi.org/10.13057/biodiv/d220802

Yasmin, I., Saeed, M., Khan, W. A., Khaliq, A., Chughtai, M. F. J., Iqbal, R., Tehseen, S., Naz, S., Liaqat, A., Mehmood, T., Ahsan, S., & Tanweer, S. (2020). In vitro Probiotic Potential and Safety Evaluation (Hemolytic, Cytotoxic Activity) of *Bifidobacterium* Strains Isolated from Raw Camel Milk. *Microorganisms*, *8*(3), 354. https://doi.org/10.3390/microorganisms8030354

Gareau, M. G., Sherman, P. M., & Walker, W. A. (2010). Probiotics and the gut microbiota in intestinal health and disease. *Nature Reviews. Gastroenterology & Hepatology,* 7(9), 503-514. https://doi.org/10.1038/nrgastro.2010.117

Dowarah, R., Verma, A. K., & Agarwal, N. (2017). The use of Lactobacillus as an alternative of antibiotic growth promoters in pigs: A review. *Animal Nutrition*, 3(1), 1-6. https://doi.org/10.1016/j.aninu.2016.11.002

Hooper, L. V., & Gordon, J. I. (2001). Commensal host-bacterial relationships in the gut. Science, 292(5519), 1115-1118. https://doi.org/10.1126/science.1058709

Yang, S. C., Lin, C. H., Sung, C. T., & Fang, J. Y. (2014). Antibacterial activities of bacteriocins: application in foods and pharmaceuticals. *Frontiers in Microbiology*, 5, 241. https://doi.org/10.3389/fmicb.2014.00241

Servin, A. L. (2004). Antagonistic activities of lactobacilli and bifidobacteria against microbial pathogens. *FEMS Microbiology Reviews*, 28(4), 405-440. https://doi.org/10.1016/j.femsre.2004.01.003

Konings, W. N., Kok, J., Kuipers, O. P., & Poolman, B. (2000). Lactic acid bacteria: the bugs of the new millennium. *Current Opinion in Microbiology*, 3(3), 276-282. https://doi.org/10.1016/s1369-5274(00)00089-8

Cotter, P. D., Hill, C., & Ross, R. P. (2005). Bacteriocins: developing innate immunity for food. *Nature Reviews. Microbiology*, 3(10), 777-788. https://doi.org/10.1038/nrmicro1273

Preidis, G. A., & Versalovic, J. (2009). Targeting the human microbiome with antibiotics, probiotics, and prebiotics: gastroenterology enters the metagenomics era. *Gastroenterology*, 136(6), 2015-2031. https://doi.org/10.1053/j.gastro.2009.01.072

Thomas, C. M., & Versalovic, J. (2010). Probiotics-host communication: Modulation of signaling pathways in the intestine. *Gut Microbes*, 1(3), 148-163. https://doi.org/10.4161/gmic.1.3.11712

Frank, D. N., Zhu, W., Sartor, R. B., & Li, E. (2011). Investigating the biological and clinical significance of human dysbioses. *Trends in Microbiology*, 19(9), 427-434. https://doi.org/10.1016/j.tim.2011.06.005

Roe, A. L., Boyte, M. E., Elkins, C. A., Goldman, V. S., Heimbach, J., Madden, E., & Smith, A. (2022). Considerations for determining safety of probiotics: A USP perspective. *Regulatory Toxicology and Pharmacology*, 136, 105266. https://doi.org/10.1016/j.yrtph.2022.105266

Fossi, B. T., & Ndjouenkeu, R. (2017). Probiotic potential of thermotolerant lactic acid bacteria isolated from "Gari" a cassava-based African fermented food. *Journal of Applied Biology & Biotechnology*, 5(04), 001-005.

Ouwehand, A. C., Forssten, S., Hibberd, A. A., Lyra, A., & Stahl, B. (2016). Probiotic approach to prevent antibiotic resistance. *Annals of Medicine*, 48(4), 246-255. https://doi.org/10.3109/07853890.2016.1161232

Gueimonde, M., Sánchez, B., G de Los Reyes-Gavilán, C., & Margolles, A. (2013). Antibiotic resistance in probiotic bacteria. *Frontiers in Microbiology*, 4, 202. https://doi.org/10.3389/fmicb.2013.00202

European Food Safety Authority. (2012). Guidance on the assessment of bacterial susceptibility to antimicrobials of human and veterinary importance. *EFSA Journal*, 10(6), 2740. https://doi.org/10.2903/j.efsa.2012.2740

Jindal, A. K., Pandya, K., & Khan, I. D. (2015). Antimicrobial resistance: A public health challenge. *Medical Journal Armed Forces India*, 71(2), 178-181. https://doi.org/10.1016/j.mjafi.2014.04.011

Zommiti, M., Feuilloley, M. G. J., & Connil, N. (2020). Update of Probiotics in Human World: A Nonstop Source of Benefactions till the End of Time. *Microorganisms*, 8(12), 1907. https://doi.org/10.3390/microorganisms8121907

Zielińska, D., Rzepkowska, A., Radawska, A., & Zieliński, K. (2015). In vitro screening of selected probiotic properties of Lactobacillus strains isolated from traditional fermented cabbage and cucumber. *Current Microbiology*, 70(2), 183-194. https://doi.org/10.1007/s00284-014-0699-0

Linares-Morales, J. R., Cuellar-Nevárez, G. E., Rivera-Chavira, B. E., Gutiérrez-Méndez, N., Pérez-Vega, S. B., & Nevárez-Moorillón, G. V. (2020). Selection of Lactic Acid Bacteria Isolated from Fresh Fruits and Vegetables Based on Their Antimicrobial and Enzymatic Activities. *Foods*, 9(10). https://doi.org/10.3390/foods9101399

Nguyen, J., Lara-Gutiérrez, J., & Stocker, R. (2021). Environmental fluctuations and their effects on microbial communities, populations and individuals. *FEMS Microbiology Reviews*, 45(4). https://doi.org/10.1093/femsre/fuaa068

Guffey, A. A., & Loll, P. J. (2021). Regulation of Resistance in Vancomycin-Resistant Enterococci: The VanRS Two-Component System. *Microorganisms*, *9*(10), 2026. https://doi.org/10.3390/microorganisms9102026

Kumar, M., Prasanna, R., Lone, S., Padaria, J. C., & Saxena, A. K. (2013). Cloning and expression of dnaK gene from Bacillus pumilus of hot water spring origin. *Applied & translational genomics*, *3*(1), 14–20. https://doi.org/10.1016/j.atg.2013.10.001

Lin D. C. (2003). Probiotics as functional foods. *Nutrition in clinical practice : official publication of the American Society for Parenteral and Enteral Nutrition*, *18*(6), 497–506. https://doi.org/10.1177/0115426503018006497

Salminen, S. J., Gueimonde, M., & Isolauri, E. (2005). Probiotics that modify disease risk. *The Journal of nutrition*, *135*(5), 1294–1298. https://doi.org/10.1093/jn/135.5.1294

Abdou, A. M., Hedia, R. H., Omara, S. T., Mahmoud, M. A. E., Kandil, M. M., & Bakry, M. A. (2018). Interspecies comparison of probiotics isolated from different animals. *Veterinary world*, *11*(2), 227–230. https://doi.org/10.14202/vetworld.2018.227-230

Karami, S., Roayaei, M., Hamzavi, H., Bahmani, M., Hassanzad-Azar, H., Leila, M., & Rafieian-Kopaei, M. (2017). Isolation and identification of probiotic *Lactobacillus* from local dairy and evaluating their antagonistic effect on pathogens. *International journal of pharmaceutical investigation*, *7*(3), 137–141. https://doi.org/10.4103/jphi.JPHI_8_17

Haghshenas, B., Nami, Y., Almasi, A., Abdullah, N., Radiah, D., Rosli, R., Barzegari, A., & Khosroushahi, A. Y. (2017). Isolation and characterization of probiotics from dairies. *Iranian journal of microbiology*, *9*(4), 234–243.

Menendez, E., & Garcia-Fraile, P. (2017). Plant probiotic bacteria: solutions to feed the world. *AIMS microbiology*, *3*(3), 502–524. https://doi.org/10.3934/microbiol.2017.3.502

Lin, E. K., Chang, W. W., Jhong, J. H., Tsai, W. H., Chou, C. H., & Wang, I. J. (2023). *Lacticaseibacillus paracasei* GM-080 Ameliorates Allergic Airway Inflammation in Children with Allergic Rhinitis: From an Animal Model to a Double-Blind, Randomized, Placebo-Controlled Trial. *Cells*, *12*(5), 768. https://doi.org/10.3390/cells12050768

Hill, D., Sugrue, I., Tobin, C., Hill, C., Stanton, C., & Ross, R. P. (2018). The *Lactobacillus casei* Group: History and Health Related Applications. *Frontiers in microbiology*, *9*, 2107. https://doi.org/10.3389/fmicb.2018.02107

Jones R.M. (2017). The Microbiota in Gastrointestinal Pathophysiology: Implications for Human Health, Prebiotics, Probiotics, and Dysbiosis. *The Use of Lactobacillus casei and Lactobacillus paracasei in Clinical Trials for the Improvement of Human Health*; Academic Press; Cambridge, MA, USA: pp. 99–108.

Balzaretti, S., Taverniti, V., Guglielmetti, S., Fiore, W., Minuzzo, M., Ngo, H. N., Ngere, J. B., Sadiq, S., Humphreys, P. N., & Laws, A. P. (2017). A Novel Rhamnose-Rich Hetero-exopolysaccharide Isolated from Lactobacillus paracasei DG Activates THP-1 Human Monocytic Cells. *Applied and environmental microbiology*, *83*(3), e02702-16. https://doi.org/10.1128/AEM.02702-16

Kim, J. S., Liu, L., & Vázquez-Torres, A. (2021). The DnaK/DnaJ Chaperone System Enables RNA Polymerase-DksA Complex Formation in Salmonella Experiencing Oxidative Stress. *mBio*, *12*(3), e03443-20. https://doi.org/10.1128/mBio.03443-20

Meibom K. L., Dubail I., Dupuis M., Barel M., Lenco J., Stulik J., et al. (2008). The heat-shock protein ClpB of *Francisella tularensis* is involved in stress tolerance and is required for multiplication in target organs of infected mice. *Mol. Microbiol.* 67 1384–1401. 10.1111/j.1365-2958.2008.06139.x

Glaza P., Ranaweera C. B., Shiva S., Roy A., Geisbrecht B. V., Schoenen F. J., et al. (2020). Repurposing p97 inhibitors for chemical modulation of the bacterial ClpB/DnaK bi-chaperone system. *J. Biol. Chem.* 296:100079. 10.1074/jbc.RA120.015413

Tripathi P., Singh L. K., Kumari S., Hakiem O. R., Batra J. K. (2020). ClpB is an essential stress regulator of *Mycobacterium tuberculosis* and endows survival advantage to dormant bacilli. *Int. J. Med. Microbiol.* 310:151402. 10.1016/j.ijmm.2020.151402

Sieber, K. B., Bromley, R. E., & Dunning Hotopp, J. C. (2017). Lateral gene transfer between prokaryotes and eukaryotes. *Experimental cell research*, *358*(2), 421–426. https://doi.org/10.1016/j.yexcr.2017.02.009

ENCODE Project Consortium (2012). An integrated encyclopedia of DNA elements in the human genome. *Nature*, *489*(7414), 57–74. https://doi.org/10.1038/nature11247

ACCESSION NUMBERS: Antimicrobial Resistance Genes: *bmr3* (IFVACORR3RYYD9F), *stp* (6FL26G1YG-JQNVC), *lmrA* (WINBPF2SHX11TZ1), *emrY* (J481GWNCZRGXWUT), *yheI* (OJ5BGFKMGZOHUGG), *yheH* (JZLJH8Cl_GBL7X3), *tetA_1* (386040461), *tetO* (FOFYWHHBJ59NNME), *marR* (BENSSXVNWHFX88W); Alcohol tolerance: *yjmD* (9W56F4TCKFOLMA1), *clpB* (CVUASSMOQSZSHEN), *adha* (TGAE3M-9ODZ14KZ), *katA* (WJGVHOFP_ASOOTW); pH tolerance: *phoP* (KB935FL1AFT6JHC), *clpB* (CVUASSMOQSZSHEN); and Thermotolerance: *clpB* (CVUASSMOQSZSHEN), *dnaJ* (HWAVQ5V5PEOWE73), *dnaK* (Z7ZALBCSB3HQBMO), *grpE* (N7OKVC2YHKHXBG0), *hsp15* (FTT_0739C), *hslR* (BSU_00590)

ACKNOWLEDGMENTS: This project was supported by the Commission on Higher Education (CHED) in the Philippines with grant number LAKAS2022-005. We thank Dr. Policarpio Joves for providing a microbiology laboratory.

AUTHOR CONTRIBUTIONS: PJIG conceptualized, designed and directed the project and analyzed the data. JC and AD isolated the probiotics from nipa sap. RJP and AGBP performed the genome assembly and analysis. MTJA and JZPF identified the samples. SMB, AJR, MPO performed the bioactivity assays. SCA and DSB supervised the isolation and characterization activities. PJIG, JZPF, and AGBP wrote the paper.