Shewanella is a putative producer of polyunsaturated fatty acids in the gut soil of the composting earthworm *Eisenia fetida*

Jan-Philipp Wittlinger[1], Natalia Castejón[2], Bela Hausmann[3,4], David Berry[1,3‡], Stephanie L. Schnorr[1‡*]

1. Centre for Microbiology and Environmental Systems Science, Division of Microbial Ecology, University of Vienna, Vienna Austria

2. Department of Food Chemistry and Toxicology, Faculty of Chemistry, University of Vienna, Waehringer Str. 38, 1090 Vienna, Austria

3. Joint Microbiome Facility of the Medical University of Vienna and the University of Vienna, Vienna, Austria

4. Division of Clinical Microbiology, Department of Laboratory Medicine, Medical University of Vienna, Vienna, Austria

‡ These authors share senior authorship

* Corresponding author:

Stephanie L. Schnorr

Djerassiplatz 1, 1030 University of Vienna

stephanie.schnorr@univie.ac.at

Abstract (max 250)

Polyunsaturated fatty acids (PUFAs) play a crucial role in aiding bacteria to adapt to extreme and stressful environments. While there is a well-established understanding of their production, accrual, and transfer within marine ecosystems, knowledge about terrestrial environments remains limited. Investigation of the intestinal microbiome of earthworms has illuminated the presence of PUFAs presumably of microbial origin, which contrasts with the surrounding soil.

To comprehensively study this phenomenon, a multi-faceted approach was employed, combining fatty acid analysis with amplicon sequencing of the PfaA-KS domain of the anaerobic fatty acid synthase gene (*pfa*), as well as the 16S rRNA and 18S rRNA genes. This methodology was applied to scrutinize the gut microbiome of *Eisenia fetida*, its compost-based dietary source, and the resultant castings.

This study unveiled a distinct gut soil ecosystem from input compost and output castings in fatty acid profile as well as type and abundance of organisms. 16S sequencing provided insights into the microbial composition, showing increased relative abundance of certain Pseudomonadota, including *Shewanellaceae*, and Planctomycetota, including *Gemmataceae*> within the gut microbiome compared to input bulk soil compost, while Actinomycetota and Bacillota were relatively enriched compared to the casted feces. Sequencing of the PfaA-KS domain revealed ASVs belonging primarily to Shewanella. Intriguingly, the 20C PUFAs were identified only in gut-soil samples, though PfaA-KS sequence abundance was highest in output castings. Overall, the results indicate that *Shewanella* in particular can explain PUFA enrichment in the gut environment as a result of the *pfa* gene presence detected via PfaA-KS sequence data.

Importance (max 150)

Prior research has demonstrated that earthworm microbiomes can potentially harbor PUFAs that are not found within their residing soil environment. Moreover, distinct indicator species have been pinpointed for various earthworm genera. Nevertheless, none of these studies have integrated metataxonomic and fatty acid analysis to explore the origin of PUFA synthesis in any earthworm species, with the objective of identifying the specific organisms and locations responsible for this production. This study suggests that earthworms accumulate PUFAs produced from bacteria, especially *Shewanella,* activated through the gut ecosystem.

Introduction

Fatty acids (FAs) are a key component of various lipid types and play important biological roles in life, such as serving as an energy resource, participating in signaling pathways, and stabilizing cell membranes (1–3). They are incorporated into lipid structures through ester bonds and can be classified into different groups based on their carbon chain length. Further they can be classified on their degree of saturation and the location and orientation of double bonds (4). Poly-unsaturated fatty acids (PUFAs), especially omega-3 and omega-6 PUFAs, are essential fatty acids for all higher organisms and have diverse functions, including building the lipid bilayer of plasma membranes (5–8). The degree of saturation of the lipid bilayer determines its permeability and fluidity, which can influence cell integrity and metabolism for energy production, growth and replication, particularly in response to environmental stressors like pressure, temperature, or oxidative stress (2, 9, 10). Omega-3 ($\omega 3$) fatty acids have anti-inflammatory functions and oppose the pro-inflammatory effects mediated by omega-6 ($\omega 6$) fatty acids. The omega-3 and omega-6 PUFAs, especially eicosapentaenoic acid (EPA), arachidonic acid (ARA), and docosahexaenoic acid (DHA), are physiologically important for broad precursors for eicosanoid secondary metabolites (8, 11). The production of eicosanoids is critical for invertebrate microfauna such as insects and annelids to modulate physiological processes and immunity (12, 13).

De novo PUFA synthesis by microorganisms

PUFAs are synthesized by two known pathways. The aerobic pathway involves elongation and oxygenic desaturation steps which are catalyzed by specific desaturases and elongases until the final PUFA length and unsaturation is reached (14). The anaerobic pathway, synthesizes PUFAs via a multifunctional mega-enzyme and double bonds are not introduced through oxygenic desaturation but rather by a PUFA synthase (15). This process, which was fully elucidated in 2001 (16), is encoded by a type I polyketide-like fatty acid synthase consisting of canonical fatty acid synthase domains (15), situated among either three genes (protozoa and myxobacteria) or five genes (marine bacteria). While the aerobic pathway is used by higher animals, lower plants, and eukaryotic microorganisms, the anaerobic pathway is limited to microorganisms such as fungi, algae, and bacteria (17).

The final fatty acid product that can be produced depends on the types of elongases and desaturases encoded in the genome and the selective conditions of their activity. A description of the enzymatic pathway and evolutionary considerations of organismal production patterns have been covered in detail, readers are especially referred to Kabeya et al. 2018 and Twining et al. 2021 (18, 19). To summarize, there is widespread presence of aerobic PUFA synthase genes among invertebrate taxa, including members of Nematoda, Cnidaria, Rotifera, Mollusca, Arthropoda, and Annelida. These genes putatively confer de novo synthesis capacity, and their functional characterization gives further evidence that their presence is part of an evolutionarily conserved adaptation to meet physiological needs for certain long-chain PUFAs. Even so, major groups of invertebrate producers experience a fitness cost when dietarily preformed sources of PUFAs are absent, and exhibit dramatic reduction in growth and reproduction (19, 20), suggesting that outsourcing PUFA production to primary producers is preferred. Earthworms, as part of Annelida, may contain the enzymatic repertoire in their genome to

produce long-chain PUFAs (18), though this has not been specifically explored in Lumbricidae.
Terrestrial and freshwater vertebrates are physiologically constrained by the limited availability of long-
chain PUFAs, and cope by either adjust dietary foraging behaviors, or through adaptive radiations to
increase synthesis capacity encoded in the genome (19, 21). For example, mammals lack Δ12- and
Δ15-desaturases, preventing the potential to synthesize linoleic and alpha-linoleic acid, which are
precursors to longer-chain FAs (22). Therefore, EPA, ARA and DHA are considered essential in the
mammalian diet. However, PUFAs accumulate at higher trophic levels, starting from microalgae and
bacteria as primary producers (23, 24). Marine bacteria and microalgae use PUFAs in membrane
lipids to maintain sufficient membrane fluidity in high-pressure and low-temperature environments (2,
9).

Linking microbial activity to the food web

As PUFAs are important to human and animal development, physiology, and health, it is crucial to
better understand PUFA producers and their role in the food web. This knowledge is particularly
valuable to the biochemical engineering industry, as it aims to develop biosynthetic methods for PUFA
production (25). Additionally, there is a significant goal to increase PUFA yield from natural
microorganisms, reducing dependence on marine resources and promoting sustainable production
(26). However, the supply of PUFAs is linked with aquatic organisms, and there is little evidence or
expectation for a de-novo PUFA producer in terrestrial habitats, leaving open the question as to how
these habitats are adequately supplied. Therefore, understanding how PUFAs are produced in
terrestrial ecosystems and supplied to terrestrial food webs has cross-disciplinary interest from fields
such as human evolution, ecology, and microbiology. The bacterial contribution to this effect has
received relatively far less attention than have plant and eukaryotic producer systems, yet there is
promising genetic and functional evidence that soil bacteria are capable PUFA producers (27). One
unique study focused on the earthworm *Lumbricus terrestris*, and revealed increasing concentrations
of PUFAs, including ARA, EPA, and DHA, from the surrounding soil to the intestinal contents –
referred to hereafter as "gut-soil" – of *L. terrestris* (28). The authors concluded that a variety of unique
fatty acids, including the long-chain PUFAs, are produced by microorganism inside of the earthworm
gut, and that these lipids are then transferred from the intestine to the luminal cells and then finally into
muscle tissue. The long-chain PUFAs were hypothesized to be used for energy needs of the *L.*
terrestris. Another study analyzed the fatty acid composition of gut-soil from the epigeic composting
earthworm, *Eisenia fetida*, in a comparison between bulk soil containing the earthworms, and soil
without (29). The study reported a lower total fatty acid content in soil with *E. fetida*, along with
reduced microbial biomass and diversity. The identified fatty acids ranged from twelve to twenty
carbon atoms in length, but no unsaturation was detected among the twenty-carbon chain varieties.
Overall, these studies report that earthworms modify soil properties and the microbial communities by
selectively filtering microbial biomass, and additionally that earthworms likely benefit from the
absorption of bacterial metabolites, including certain lipids.

Efforts have been dedicated not only to the exploration of fatty acid production in earthworms but also
to the analysis of microbial compositions within diverse earthworm species towards understanding

their role in composting and soil revitalization. One study examined the gut microbiome of earthworm
species from three genera: *Aporrectodea, Allolobophora*, and *Lumbricus*, which occupy two different
ecotypes. The *Lumbricus* are vertically burrowing "anecic" worms whereas the *Aporrectodea* and
Allolobophora are horizontally burrowing "endogeic" worms (30). By utilizing 16S rRNA amplicon
sequencing, it was discovered that Pseudomonadota was the primary phylum present in both
earthworm ecotypes. Furthermore, the genus *Arthrobacter* and the class of Gammaproteobacteria
were indicator taxa for *Lumbricus*, while at higher taxonomic levels the Acidobacter and
Alphaproteobacteria were indicative of *Aporrectodea* (30). Further distinctions between ecotypes
included presence of bacteriovore protists, as well as richness of bacterial taxa, which is considered to
reflect the variation in feeding behavior of these earthworms. Another study focused on the top-soil
and leaf-litter dwelling "epigeic" *E. fetida* and assessed the microbial composition of its microbiome
when fed with different compost types, including brewers spent grain, cow manure, and a 50/50
mixture of both (31). Analysis of the earthworm castings (feces) revealed that, regardless of the
treatment, the dominant phyla remained Proteobacteria and Bacteroidetes. However, a significant
increase in Firmicutes and Actinobacteria was observed when *E. fetida* was fed with either 100% or
50% brewers spent grain. Similar results have been reported when *E. fetida* was fed vegetable waste
(32). In sum, the effects of feeding behavior and the impact of soil conditions have been well studied in
how they can modify composition of the earthworm gut-soil microbiota. These works lend important
insights about how the earthworm gut-soil profile is distinct but related to the microbial composition of
bulk soil. Variations between different types of earthworms appear dependent upon their ecotype
rather than species.

These previous works suggest that the gut-soil microbiome of earthworms may contain PUFAs that
are not initially present in the soil that they inhabit, and so, are not taken-up from primary consumption.
However, the combined study of the gut microbiota and the lipid metabolites from the same sample set
is needed to resolve the source of PUFA production and to define the microbial organisms involved in
this production. We address this gap with an analysis of the fatty acid and microbial compositions of
gut-soil, bulk soil, and earthworm castings of *E. fetida*. We hypothesize that bacteria in terrestrial
environments, including those in the guts of terrestrial animals, may be capable of producing PUFAs,
but only under certain conditions or with specific microbial neighbors. To test this hypothesis, the gut
soil content of *E. fetida* was analyzed in three parts. First, the fatty acid composition was analyzed
using gas-chromatography mass spectrometry (GC-MS). Second, the prokaryotic and eukaryotic
constituents were profiled using 16S and 18S rRNA gene amplicon sequencing. Finally, amplicon
sequencing was performed on the initiator ketoacyl-synthase (KS) domain on the polyunsaturated fatty
acid gene (*pfaA*) (33) in order to identify microorganisms with the genetic potential to de-novo produce
PUFAs using the iterative type-1 polyketide-like synthase system. This work uses the earthworm and
its gut microbiota as a model for terrestrial ecosystem dynamics to identify potential PUFA producers
and to help us understand the origin of PUFAs in terrestrial ecosystems and the diversity of
microorganisms that may provide them.

Results and Discussion

We analyzed the total fatty acid content and microbiome profiles of three sample types: pre-compost

(PC), vermicompost (VC), and the gut soil (GS) of the earthworm *E. fetida.* PC is a mixture of horse

manure and thermophilic compost used as feed and applied once per week, while VC is worm castings

(feces) that is normally collected via sifting and used as an organic fertilizer. The sampling was

conducted over two weeks, with samples collected on three specifically spaced days within each week.

PC was only sampled on the first day of each weekly "feeding" cycle, as it is laid freshly onto the

existing *E. fetida* compost housing once a week.

Earthworm gut-soil harbors a distinct fatty acid profile enriched in PUFAs

The fatty acid profile of samples collected in week one and week two were compared to identify any

batch effects between sampling weeks based on sample type or sampling days (as shown in Materials

and Methods). Permutational multivariate analysis of variance (PerMANOVA) showed a significant

difference (p = 0.016) between the sampling weeks, but this only explained 0.5% of the variation, while

most of the variation (87.6%, p = 0.001) was explained by the sample type (Figure 1). We then

examined whether there were any significant differences between the respective days of sampling for

each sample type using the Kolmogorov-Smirnov test. No significant differences were identified for all

sample types, therefore, the data from week one and week two were merged for all subsequent fatty

acid composition analysis.

According to the analysis of the fatty acid composition of PC presented in Table 1, palmitic acid (16:0)

was found to be the most abundant fatty acid, constituting on average over 30% of the detected fatty

acids, followed by stearic acid (18:0) and oleic acid (18:1 *cis-ω9*) at 13% and 14%, respectively. The

only PUFA detected in PC was linoleic acid (18:2 *ω6*), which accounted for more than 6% of the total

fatty acid content, and likely derives from bulk plant material in the PC mixture. Linoleic acid is known

to be a precursor for the synthesis of the omega-6 series of fatty acids, particularly leading to ARA,

whereas α-linolenic acid, which is the precursor to the omega-3 series, including EPA, and DHA, was

not detected.

One curious finding was the presence of *trans*-fatty acids. Although bacteria do not usually produce

trans-fatty acids, they can produce them under specific circumstances, which has been demonstrated

in *Vibrio* and *Pseudomonas*. In *Vibrio*, a temperature shift caused cis/trans isomerization, while in

Pseudomonas, toxic compounds such as organic hydrocarbons resulted in isomerization (34–37). PC

partly consists of thermophilic compost which has been heat processed through fermentative activity,

and this temperature increase could potentially have caused *cis/trans* isomerization from bacterially

originated FAs.

[Table 1: Fatty acid composition of pre-compost, vermicompost, and gut-soil samples]

The percentage of PUFAs in GS was found to range from 17 to 20% of all FAs across the three

sampling days. ARA (20:4 *ω6*) and EPA (20:5 *ω3*) were identified as two notable fatty acids,

comprising ~7-9% and 4.5-5.5% respectively of the detected fatty acids. Other major fatty acids

included lauric acid (12:0, ~4-6%), elaidic acid (the trans isomer of oleic acid,18:1 *trans-ω9*, ~4-6%)

and gondoic acid (20:1, ~6-9%). No significant differences in the relative abundance of fatty acids were

observed between sampling days (p > 0.05), except for lauric acid (12:0) (two-way ANOVA with post-
hoc Tukey, day 1 – day 3: $p = 0.0280$), arachidonic acid (20:4 ω6) (two-way ANOVA with post-hoc
Tukey, day 1 – day 2: $p = 0.0451$), the overall SFA content (two-way ANOVA with post-hoc Tukey, day
1 – day 2: $p = 0.0002$, day 2 – day 3: $p = 0.0002$) and the overall PUFA content (two-way ANOVA with
post-hoc Tukey, day 1 – day 2: $p = 0.0002$). PUFAs were found to have a lower relative content on day
one of sampling compared to day two. Lauric acid content increased steadily from day one to day
three, but only the difference between day one and three was found to be significant. This may be due
to the feeding cycle, which occurs on day one of the week, and therefore the second sampling point
captures the peak nutrient availability and metabolic activity, which then falls off by the end of the
week. The high percentage of unidentified fatty acids (n.i.), which accounted for approximately 23% of
the total content, were compounds either structurally similar to fatty acids but unidentifiable with the
standard fatty acid methyl-ester (FAME) mix, or compounds with low identification scores in the
national institute of standards and technology (NIST) mass spectral library and thus grouped together
as "not identified".

Only six fatty acids were identified in VC, all of which were the same as those found in PC samples.
Among these, linoleic acid (18:2 ω6) was the only PUFA, accounting for less than 4% of fatty acids.
Palmitic acid (16:0) was one of the major fatty acids found in VC, accounting for 28-31% of the total
content, along with stearic acid (18:0), which accounted for approximately 15%. Palmitic acid showed
significant differences over the one-week sampling period (two-way ANOVA with post-hoc Tukey, p =
0.0321). We detected oleamide, or oleic acid amide, in all samples, which likely originated from the
compost added to PC, since oleamide has been detected in mixtures containing starch, sunflower oil,
and soy protein (38). These results show that PUFAs are uniquely concentrated in GS samples
compared to the starting compost and the resulting vermicompost (Figure 1).

[Figure 1]

Figure 1. Principal component analysis (PCA) of the fatty acid content extracted from pre-compost
(yellow triangle), gut soil (red dot) and vermicompost (blue square) after center-log-ratio
transformation (CLR) with each loading arrow representing one fatty acid identified.
229

These results are largely consistent with the findings of Sampedro et al. (2006), who reported elevated
levels of PUFAs including ARA, EPA, and DHA in the gut of *L. terrestris* compared to the surrounding
bulk soil (28). In our study with *E. fetida*, the PC is the bulk soil that serves as substrate gut microbiota
metabolism, the products of which are available for host uptake. Since we did not find PUFAs in PC
(incoming) nor VC (outgoing) samples, then PUFAs found in the GS, either from microbial activity or
from the shedding of earthworm epithelial cells, appears to be selectively retained in the earthworm
and its microbiome for physiological needs. The diverse range of fatty acids that we recovered in the
GS may be indicative of high metabolic activity stimulated in the gut.

Metataxonomic amplicon analysis

To identify potential microbial producers of ARA and EPA in the gut soil samples, we used 16S and
18S rRNA gene sequencing alongside amplicon sequencing of the PfaA-KS domain using primers

previously developed to target the PUFA synthase complex (described in the next section) (33). By

using high-level taxonomic binning of the functional gene sequences and then examining members of

these taxonomic bins among the sample types, potential producers of ARA and EPA could be

identified. To account for effects of low biomass and interindividual variation, samples of GS from

individual *E. fetida* (denoted IG or MG) were also compared with the pooled GS samples (see Table

S1 for sample metadata).

The V4 hypervariable region of the 16S rRNA was targeted using the 515F/806R universal primers

with slight modification, and the TAReuk454FWD1/TAReukREV3mod universal primers were used to

target the eukaryotic 18S rRNA gene for amplification and sequencing on an Illumina MiSeq (see

Materials and Methods for primer sequences). After decontamination of 16S rRNA sequences, an

average of 8116 (±4193 SD) reads were obtained for GS, 5606 (±4124 SD) for IG, 8296 (±5664 SD)

for PC, and 7648 (±4295 SD) for VC. A total of 4238 ASVs were identified across all the samples. The

18S sequencing resulted in an average of 30753 (±15703 SD) reads for GS, 25977 (±10907 SD) for

IG, 18221 (±12272 SD) for PC and 20034 (±14504 SD) for VC after decontamination with a total of 886

identified ASVs (see Tables S2 & S3 for 16S and 18S ASV tables).

<u>Prokaryote taxonomic sequence profiles</u>

Chao1 and Shannon metrics were used to analyse alpha-diversity of the microbiome, based on a

rarefied ASV table with 5000 reads for 16S rRNA analysis and 10000 reads for 18S rRNA analysis

(Figure S1 rarefaction curves). No significant difference was found in richness or diversity between GS

and IG samples. However, gut soils (GS & IG) showed significant variation compared to compost

samples (PC & VC), with VC displaying significantly higher measures of richness compared to both gut

soils (Wilcoxon, GS-VC: p = 3.166e-06 and IG-VC: p = 1.615e-07; Figure 2a). Similar variation was

observed in Shannon diversity metrics, with significant differences identified between both gut soil

samples compared to VC (Wilcoxon, GS-VC: p = 0.001636 and IG-VC: p = 9.578e-06; Figure 2b). This

suggests that VC and PC have a higher evenness and contain more ASVs that were only identified

once or twice compared to GS samples. Beta-diversity analysis was conducted using the unrarefied

ASV table, but was normalized using a variance stabilizing transformation with DESeq2. Ordination

analysis using Euclidean distance on center-log-ratio transformed data indicated distinct sample

clusters for PC and VC, while IG and GS clustered together (Figure 2c).

[Figure 2]

Figure 2. Alpha- and beta-diversity analysis comparisons using the 16S rRNA data for gut soil (GS,
red), individual gut soil (IG, green), Pre-compost (PC, yellow) and Vermicompost (VC, blue). (**A**, **B**)
Boxplots for Chao1 and Shannon alpha-diversity metrics with significant differences shown for
different sample types with pair wise comparisons between groups by Wilcoxon test (p > 0.05 (ns), p
≤ 0.05 (*), p ≤ 0.01 (**), p ≤ 0.001 (***), p ≤ 0.0001 (****)). (**C**) Ordination of ASVs using Euclidean
distance metrics shows separation of sample types. (**D**) Ward's clustering of rarefied ASVs binned at
family- or lowest assigned taxa level and filtered for 2% abundance in at least 5% of samples.

Based on the analysis of alpha- and beta-diversity, it was found that there were no significant

differences in richness or diversity between the pooled and individual GS samples, and that they

clustered together. In contrast, the PC, VC, and gut soil samples were found to each have distinct

microbial community compositions. Figure 2c shows clustering of class-level bins and the relatively
enriched or depleted taxa across samples, and reveals that GS samples largely cluster together with
the exception of a few IG samples, while the compost samples form a separate cluster. This suggests
that either the earthworm gut community is truly an endemic microbiome, or that the gut environment
provides highly selective conditions that promote transient growth and activity in a stable
reproduceable manner (39).

The relative abundance of major phyla and families in each sample type provides additional evidence
of different microbial community compositions per sample type. Phyla that contribute at least 5% and
families that contribute at least 2% to the total relative abundance are compared for relative
abundance. To test for significant differences in relative abundance, a pairwise Wilcoxon analysis with
post Benjamini-Hochberg was conducted between each sample type (Figure 3, and see Figure S2
pairwise family level corr). Comparing IG and GS samples revealed significant differences across
sample types at the phylum, class, and family level, however, these differences were primarily driven
by only a few family-level groups within the primary segregating phyla. Specifically, significant
differences were found in three of 51 families in Gammaproteobacteria, five of 61 families within
Firmicutes, and seven of 37 families within Bacteroidota for gut sample comparisons. On the other
hand, pronounced differences were observed in the comparisons between the gut soils and composts,
where most families showed significant changes in abundance. In general, there was significant
variation in relative abundances when gut soils were compared with composts (Figure S3a *relative*
abundance barplots). While the significance was less for IG than for GS in most cases, the trend of the
mean values was the same, supporting that the features seen in pooled samples can be generalized to
individual specimens, and are not driven by stochastic or random features from outliers.

Notably, Actinobacteriota and Firmicutes display a higher relative abundance in gut soil samples
compared to the compost samples, whereas Bacteroidota show a higher relative abundance in the
compost samples compared to the gut soils.

[Figure 3]

Figure 3. Pair wise comparisons of each sample type (GS), individual gut soil (IG), pre-compost (PC)
and vermicompost (VC) by Wilcoxon test with post Benjamini-Hochberg showing adjusted p-values for
each comparison (p > 0.05 (ns), p ≤ 0.05 (*), p ≤ 0.01 (**), p ≤ 0.001 (***), p ≤ 0.0001 (****)). Blue color
scale indicates that the mean of sample A is lower than in sample B. Red color scale indicates that the
mean of sample A is greater than the mean of sample B.

The pair wise comparison of sample types supported the findings from the taxonomic summary
regarding Actinobacteriota, Bacteroidota, and Bacillota (formerly Firmicutes). Specifically,
Bacteroidota, Gammaproteobacteria, and Chloroflexi had a higher relative abundance in both compost
samples compared to each of the gut soil samples. This was also true for Alphaproteobacteria when
comparing the gut soil samples to VC. Verrucomicrobiota had a higher relative abundance in the gut
soils compared to PC, but still lower than in VC. No differences were found in Planctomycetota for all
comparisons and in Alphaproteobacteria when comparing the gut soils to PC.

In previous works, the major phyla identified in *E. fetida* when fed with different types of compost were
Proteobacteria and Bacteroidota. However, since the compost used in our study differed from that

used in the analysis by Budroni et al. (2020), slight differences in the microbial composition of GS are
expected (31). This is supported by previous reports that show changes in microbial composition
based on the compost composition, as well as other studies that varied the feeding compost (40). A
study by Sapkota et al. (2020) also identified Proteobacteria as a major phylum in their analysis for the
earthworm genera *Aporrectodea* and *Lumbricus* (30). Although Proteobacteria and Bacteroidota are
still present in GS from our study, they do not account for most of the microbial composition in this
sample type. Their lower abundance could be explained by the nutrient-rich PC substrate, applied to
the earthworm population as feed and housing matrix. Previous research has shown that
Proteobacteria and Bacteroidota are particularly increased in abundance when *E. fetida* is fed with a
nutrient-poor substrate, whereas a nutrient-rich substrate results in higher microbial diversity and a
higher abundance of Firmicutes and Actinobacteria (40).

<u>PfaA-KS for assessment of bacterial PUFA producers</u>
To understand the potential that PUFA lipids concentrating in the gut of the earthworm are derived
from bacteria, we used a previously developed universal primer targeting the beta-ketoacyl synthase
(KS) region in the single-copy pfaA gene, which amplifies an approximately 502 bp region spanning
the PfaA-KS N- and C-terminals (33). The KS domains from PKS types tend to cluster as a
monophyletic group, and can approximate phylogeny since horizontal gene transfers (HGT) of PUFA
synthase complex somewhat mimic patterns of vertical inheritance (41, 42). In addition, the KS
sequence phylogeny is distinctive of the type of PKS complex in which they occur, which makes it a
suitable candidate marker for targeting the genes involved in synthesis of a specific metabolite. We
sequenced the amplified KS products using an Illumina MiSeq, resulting in 53 samples with
successfully generated, filtered, and PfaA-KS-assigned reads, and final average sequence counts
segregating per sample type: 16 (±27 SD) for GS derived reads, 315 (±526 SD) for PC, and 1230
(±1459 SD) for VC. A total of 65 PfaA-KS ASVs were derived using the DADA2 workflow on merged
reads, leading to the identification of 6 resolved unique bacterial strains within 14 taxonomic bins
across 5 phyla (Table S5). Neither microeukaryotes nor fungi were found among the annotations,
although some of these organisms harbor an iterative type I PKS (T1PKS) gene complex and KS in a
neighboring clade to bacterial iterative T1PKS (15, 42, 43).

Little is known about the distribution of PUFA-synthase genes and production potential in
microorganisms outside of marine contexts, with only very recent investigations beginning to probe the
terrestrial biospheres (17, 44). To date, we are aware of only one primer set that has been developed
for specifically targeting the **pfa** T1PKS KS domain, which was modeled from the sequences of marine
Pseudomonadota, and especially those belonging to γ-Proteobacteria. Since these bacteria appear to
have propagated T1PKS genes via HGT (42), a single primer pair may work universally within this
phylum to capture many **pfa** KS sequences. However, this specificity towards marine
Gammproteobacteria likely hinders recovery of bacteria in diverse phyla deriving from terrestrial soil
and freshwater ecosystems such as actinobacteria and cyanobacteria respectively, where T1PKS
appear to have evolved rather from a common ancestor. Indeed, certain soil taxa with known PUFA
production competence were surprisingly not found among our annotated sequences, such as some
strains within myxobacteria, Actinomycetes, and Desulfobacteria (27, 45–47). Despite this limitation, a

surprising number of sequences and diversity were recovered from the compost and earthworm gut
soil sample material that we report here. We recovered PfaA-KS sequences that had >95% nucleotide
homology to those found among marine bacterial reference isolates using BLASTn query on the full
nucleotide collection (nt). For sequences that were not well resolved using the NCBI RefSeq nt
database, a BLASTx translated amino acid sequence was mapped against the non-redundant (nr)
database, with most hits achieving >95% homology (see Tables S6 and S7 for a full summary of
BLAST hit results).

<u>Phylogenetic inference among recovered sequences</u>
In order to understand the relationship of the sequences we captured from earthworm gut and compost
soil samples, we constructed a phylogenetic inference tree using our 65 bacterial PfaA-KS ASVs plus
the 446 PfaA-KS sequences that were published in the original publication for creation of the PfaA-KS
primer set (33) (Figure 4; Table S8). The relevant difference with our present study is use of the PfaA-
KS primers on terrestrial soil-derived samples, whereas prior research has been restricted to their
application on marine water and sediment samples. Using the *Escherichia coli fab*F gene as an
outgroup (representative of a KS subtype involved in the canonical type II FAS system), we see that
our sequences surprisingly form clusters among shallow branches of the tree which match their
taxonomic assignments, rather than their environmental source. However, known evolutionary
relationships that are based on whole-genomes are not recapitulated, since more distantly related taxa
(such as Pseudomonadota and Bacteroidota) form larger clusters that break up established
evolutionary lineages, such as between members of Gammproteobacteria.

[Figure 4 Tree]

Figure 4. Phylogenetic inference of 511 PfaA-KS amplicon sequences from this study (earthworm) and
prior work investigating marine bacterial sources of PUFA synthesis. The fabF gene from E. coli was
used as outgroup and root. Stacked barplots show the number of samples that contain the ASV
sequence and are colored by the sample type.

There is a clear separation between sequences found within earthworm GS samples and those of the
compost samples (denoted by barplots along the outer ring of the tree). All GS sequences are
assigned to *Shewanella* and cluster among a fairly undifferentiated lineage that include other *Vibrio*
assigned sequences, and which stem from a bifurcation that separates 1) facultatively anaerobic cold-
water-marine Gammaproteobacteria (*Vibrio*, *Colwellia*, *Moritella*, *Psychromonas*) and aerobic
saprophytic or predatory Bacteroidota (*Flammeovirga*, Saprospirace); from 2) subsurface anoxic
dwelling taxa (Legionellales, Candidatus Hydrogenedentes), anaerobic fermenters (*Anaerolineae*) and
terrestrial aerobic chemoheterotrophs (Armatimonadota) (48–50). The PfaA-KS sequences found
within compost samples were spread more widely across the tree among organisms such as *Gemmata*
(Planctomycetota), Cloacimonadota, and other Anaerolineales that are often found among carbon-rich
sediments such as peatbogs, waste-water treatments, or the rhizosphere. To date, the activity and
product of these putatively competent PUFA synthase complexes remains unconfirmed (47). The
remaining one-third of the tree is wholly dominated by a basal cluster of sequences assigned to
Deltaproteobacteria or its sublineage, SAR324, which are distinguished from the other taxonomically
and phylogenetically diverse branchings. This suggests that metabolic niche participation has broadly
determined the structure of the tree, which separates primarily between marine

chemolithoheterotrophs (sulfate reducers) (51) and diverse chemoorganoheterotrophs
(carbon/nitrogen utilizers through predatory or saprophytic processes) (48, 52). This topology further
supports the observation that T1PKS are prone to HGT events (42), and that HGT propagation of the
pfa complex helps to explain its environmental and phylogenetically widespread occurrence and lack of
strict coherence among species or sub-species level presence/absence patterns (47).

PfaA-KS sequence distribution and abundance across sample types
To gain insight about which members of the microbial community that are concentrated in earthworm
gut-soil and which may be contributing to production of PUFA, we conducted in-depth investigation on
the ASV table abundances, which was first transformed to center-log-ratio (CLR) values (53). An
extremely sparse compositional table rendered normalization practices such as rarefaction and relative
abundance inadequate to standardize sequence abundances across samples. We confirmed that there
is no apparent bias in the amount of genetic data recovered from the samples based on sample type,
starting amount, or concentration, supporting the conclusion that the observed sparsity of the PfaA-KS
ASV table is biologically real (see Table S1 for sample metadata; see Figure S4 for associations
between PfaA-KS sequence recovery and sample metadata). Surprisingly, in contrast to the metabolite
lipid profiles generated from the same sample material, the recovered PfaA-KS sequences were most
diverse and concentrate among compost samples (PC and VC) rather than the GS samples (Figure
5a). The PfaA-KS sequences were sparsely distributed at very low abundance in GS samples, both
pooled and individual (Figure 5b). While the ASV taxonomic assignments suggest that just a handful of
unique taxa contribute to the total sequence variation, the sequence variants themselves are quite
numerous even within the same putative bacterial strain, and are strongly segregated between GS and
compost samples (Figure 5c). No single ASV is shared across all three sample types. This can be
most clearly seen when ASVs are aggregated at genus level, or otherwise their lowest taxonomic bin,
and then hierarchically clustered according to Ward's sum of squares method (Figure 5b). The GS
samples are composed entirely of *Shewanella* assigned sequences while the compost samples
contain a mixture of the other seven environmental taxa bins. This suggests that only *Shewanella* are
remaining in close association to the earthworm host tissue rather than passing through as transients
(indicated by the strains found in VC but not in GS), and perhaps actually are co-residents of the
earthworm gut (54).

[Figure 5]

Visualization of the sparse distribution and low abundance of 65 bacterial PfaA-KS ASVs by sample
type based on center-log-ratio transformed ASV table (**A**); and hierarchical clustering of samples
based on ASVs binned at lowest taxonomic assignment (**B**). Number of unique sequences shown
across 5 phyla assignments shows clear dominance by Pseudomonadota, and primarily from the
genus *Shewanella* (n=48) (**C**).

PfaA-KS taxa within the microbiome profile
To understand the prevalence of putative PUFA-producing taxa among the overall prokaryote
microbiome community, the taxonomic bins assigned to the PfaA-KS sequences were sought among
the 16S rRNA dataset, and the matching ASVs from the abundance tables were subset for further
analysis (Figure S6 *prok.pfa abundance barplots*). The subset of PfaA-KS taxa from the 16S data

(hereon referred to as **prok.pfa**) were normalized by the total number of ASVs, and then transformed
using CLR to compare with the PfaA-KS sequence amplicon dataset (hereon referred to as **pfa**). First,
Procrustes analysis was used to infer the correspondence of the ASV level ordinations of the **prok.pfa**
and **pfa** data using Euclidean distance. Using the function "protest" with 999 permutations, the
datasets were found to have modestly significant correlation (m12 squared = 0.758, corr. = 0.492, p =
0.001), and their ordinations show similar separation of sample types along the first and second
principal components (Figure 6a). Next, the ASVs counts were binned at genus level (or lowest
taxonomic assignment), resulting in seven PfaA-KS taxonomic assignments: Anaerolineales,
Armatimonadota, *Gemmataceae*, Hydrogenedentes, *Moritella*, Phycisphaera, and *Shewanella*. The
correspondence of these binned taxa vectors within the **pfa** and **prok.pfa** abundance tables was
calculated using a Mantel test on the Euclidean distance matrices, yielding again low correlation
(Spearman's rho = 0.194, p = 0.007, Figure S7 *mantel test*), suggesting that the abundance of the
sequences captured using the PfaA-KS primers cannot be related to the microbiome profile
reconstructed from 16S rRNA amplification. Correlation and visualization of the abundance distribution
of each taxonomic bin shows that only for *Shewanella* abundance is there coherence between the
PfaA-KS and 16S rRNA abundances (Pearson's rho: PC = 0.465, GS = 0.662; Figure 6b & 6c; Figure
S8 *TaxaLinearAssoc-pfaclrM2;* & Figure S9 *Histograms*), lending more evidence to the idea that
Shewanella are maintaining a stable community within the earthworm gut, and are not just transient
environmental taxa.

[Figure 6]

Figure 6. Comparison of ASV abundance tables for the PfaA-KS amplicons (**pfa**), and the subsetted
PfaA-KS assigned taxa from the 16S rRNA dataframe (**prok.pfa**). Ordination and procrustes analysis
in the transformed dataframes (**A**) indicate modest correlation, shown by the sum of squares (m12²)
and correlation coefficient. The principal component plots of the CLR-transformed dataframes
demonstrate this correlation from some interspersion of GS and VC samples, and with PC samples
clustering separately. The abundance values of the binned taxa for **pfa** and **prok.pfa** show the
distribution of taxa by sample (**B**). The abundance of *Shewanella* taxa were correspondent between
datasets (**C**), shown separated by sample type, as there are two distinct linear relationships for PC
and GS samples.
473
Using effect-size as a measure of explanatory power for the difference in ASV abundance in the
prok.pfa table also found Shewanella-assigned ASVs to be highly indicative of differences between
vermicompost (earthworm castings), and pre-compost (compost soil applied as feed but not containing
earthworms) (Tables S9 & S10; Figure S10 *aldex2 results*). The dissociation between **pfa** and
prok.pfa abundances for the remaining taxa is most likely because the taxa that contain a putative
PUFA gene complex are extremely low-abundant members of the total microbial community
(especially in soil or compost), and so are negatively biased during 16S rRNA primer annealing and
amplification. Additionally, the 16S rRNA sequence diversity of such rare taxa is likely not well
represented among reference databases, leading to underestimation of their presence and abundance
in complex community reconstructions using culture-independent techniques.

<u>Eukaryote taxonomic assessment for other soil microbial or microfaunal PUFA producers</u>
Earthworms ingest a variety of non-bacterial microbiota from the soil and can even be selective
foragers. Phospholipid fatty acid (PLFA) biomarkers indicate the concentration of microeukaryotes and

fungi within the earthworm gut relative to the bulk soil (55), and a robust immune response has evolved
to entrap and expel parasitic nematodes (56, 57). Prior research on the lipid content of earthworm GS
attributed the accumulation of PUFA in GS and somatic tissue to the metabolic activity of
microeukaryotes such as fungi and protozoa. This was in accordance with the understanding that
certain FAs are lipid biomarkers for the presence of organismal groups, on account of the FA
composition of their phospholipid membrane (58). According to the pattern of observations, algae,
yeast and other fungi are typically associated with synthesis of long-chain polyunsaturated acyl-
moieties, including 20C+ fatty acids and sphingolipids, from both denovo and precursor 18C fatty acids
(18, 28, 58–61). Common terrestrial bacteria on the other hand are attributed with medium-chain,
straight or branched, odd-numbered-saturated or even-numbered-monounsaturated fatty acids (58,
62), and so they have not been considered candidate producers for PUFAs found in a terrestrial
sediment context. Examples of soil-living microeukaryote organisms with notable concentrations of
PUFA include invertebrate microfauna such as nematodes (Chromadorea), and collembola (Ellipura),
fungi such as *Mortierellaceae*, and protozoan microalgae such as Labyrinthulomycetes. However, the
origin of earthworm gut PUFAs from a microeukaryotic source has not been fully investigated.
Therefore, we used 18S rRNA amplification to catalogue the presence of putative PUFA producing
eukaryotes, and to determine whether their pattern of distribution across sample types may be a factor
towards explaining PUFA accumulation in the earthworm gut environment. Eukaryote ASVs that
mapped to the Oligochaetes subclass (aquatic and terrestrial worms) were first removed to eliminate
the host signal that otherwise obliterated our ability to see representative taxa. For GS samples, this
unfortunately resulted in a loss of >90% of reads. Filtered samples were then rarefied to 1000 reads
and ASVs were binned at the "class/order/family" level for visualization and assessment (Figure 7,
Figure S3 *Eukaryote taxa assigns barplot*). Of the resulting 46 taxonomic bins achieving at least 1%
abundance in 5% of samples, four are identified as containing known PUFA producers: Chromadorea
(nematodes), Ellipura (collembola), *Mortierellaceae* (soil fungi), and *Thraustochytriaceae* (saprotrophic
protists).

[Figure 7]

Figure 7. Binned taxa filtered for those achieving 1% abundance in 5% of samples in the rarefied ASV
table plotted by Ward's clustering of scaled abundance shows a distinct division in relative enrichment
of microeukaryotes characterized by the 18S rRNA gene. Compost samples are generally enriched
across a larger number of taxonomic bins.

The Chromadorea and Ellipura percent relative abundance in GS samples were approximately 1.88%
and 0.02% of rarefied reads respectively, while significantly greater proportions of these two organisms
were found in VC samples (castings) (Wilcoxon: p<0001). *Mortierellaceae* were most prevalent in PC
(7.55%) while <1% in VC and GS, and *Thraustochytriaceae* were <1% for all sample types (Table S4).
Nematodes contain all desaturase and elongase enzymes to denovo produce up to 20C length FAs,
including 20:5ω3 EPA and 20:4n6 ARA, which were the uniquely enriched PUFAs found among our
GS samples (61). Therefore, we cannot exclude that the prevalence of nematodes passing through the
earthworm intestinal tract and expelled with the castings may explain the detection of PUFAs in the
GS. However, since PC samples also contain a smaller but detectable amount of nematode 18S rRNA
signal, and yet are devoid of any detectable PUFA, then it is unclear whether the nematodes in the

earthworm composting system where we sampled are sequestering PUFA. Previous studies have
shown that *Caenorhabditis elegans* may contain anywhere from 1% to 20% of total lipid PUFA,
depending on growth temperature (63). These are largely due to an abundance of 18C FAs, namely a
signature peak at 18:1ω7 (*cis*-vaccenic acid) in all lipid fractions – phosphatidylcholine,
phatidylethanolamine – but to a lesser degree also EPA and ARA (59, 61, 63, 64). However, none of
our sample peaks were annotated for 18:1ω7, which would be expected if nematode lipids (or
otherwise aerobic bacteria) were contributing to the pool of measured fatty acids in our samples.
Furthermore, earthworms respond to nematode infestations that occur in the coelomic cavity and near
the nephridia by encapsulating the tiny worms in cysts called "brown bodies" and destroying them
using reactive oxygen species (57). Alternatively, the earthworm itself may be able to produce PUFAs,
as recent genomic analysis has revealed genes encoding "methyl"-end desaturase homologs to the
Δ12 (ω6) and Δ15 (ω3) desaturases in Oligochaeta (18). Little is known about the ability of
invertebrates to produce long-chain PUFA, especially the amount and rate that this might occur.
Evidence from genomic analysis attests to wide-spread occurrence of elongation and desaturation
enzymes, but only a handful of earlier publications have indicated some limited de novo production
such as in Collembola (springtails), *Daphnia* (water fleas) and nematodes (20, 59, 61). Still, Arthropoda
exhibit poor fecundity and decline in fitness when their dietary supply of pre-formed PUFA are absent
or restricted, despite encoding the competent enzymes for de novo production (20, 59). In earthworms,
the increasing concentration of PUFAs from bulk soil to gut soil, and finally peaking in muscle tissue,
which was interpreted as trophic transfer from soil microorganisms to the earthworm (28), may well
work in the opposite direction. The earthworm may be producing PUFA within its own tissue, perhaps
as precursors to eicosanoids or for cellular membrane permeability, and as the epithelial cells shed
into the gut, then PUFAs appear to concentrate in gut soil. However, in this prior study the entire
unbroken intestinal track was surgically isolated from body, and it challenges any notion that somatic
cell metabolites should be leaked into the gut soil, and rather strengthens the original interpretation. To
definitively rule out earthworm-derived PUFAs, further work should isolate metabolic activity using
approaches such as stable isotope probing with heavy isotopes of carbon and hydrogen.

Conclusions

In sum, we found that the earthworm gut soil ecosystem is distinct from the input compost and output
castings according to the type and abundance of organisms present, as well as in the fatty acid
metabolite profiles. Amplicon sequencing of the KS functional domain on the *pfaA* gene as part of the
PUFA synthase complex illuminated the sparse and phylogenetically narrow distribution of putative
PUFA producing bacterial taxa, especially the ASVs from GS samples, which all unambiguously
belonged to strains of *Shewanella*. Most surprising was that while the 20C PUFAs were found among
the total lipid pool of the GS samples and absent in both compost sample types, the PfaA-KS
sequence abundance pattern was exactly the contrary. Samples in which the largest number and
diversity of PfaA-KS sequences were recovered were VC, which are the castings or feces of the
earthworms. These samples contained taxa also found among the GS samples which were implicated

as PUFA-competent based on PfaA-KS presence, but VC samples also contained other PfaA-KS-
positive taxa known to be wide-spread free-living organisms. Based on 16S and 18S rRNA taxonomic
profiling, the GS samples diverged from the compost samples based on presence and abundance of
ASVs, suggestive of a unique ecosystem that is maintained by the conditions of the earthworm gut.
Eukaryotic origin of the PUFAs is possible but not well supported, since organisms such as nematodes
and collembola failed to show other signature 18C fatty acid biomarkers, and were not well-
represented among the amplicon sequence pool, especially of the GS samples. The overall picture
appears to be that earthworms may aggregate the ARA and EPA PUFAs stemming from bacterial
metabolism that is activated via conditions in the intestinal tract, and from *Shewanella* taxa that
uniquely belong to the earthworm gut ecosystem, rather than other transient environmental bacteria.

<u>Material and Methods</u>

Materials

Method	Reagent
Lipid extraction	- Chloroform
	- Methanol
	- dH$_2$0
	- Sodium chloride (NaCl)
Basic derivatization	- Hexane
	- Potassium hydroxide (KOH)
	- Methanol
	- Sodium Bisulphate (NaHSO$_4$)
Acid derivatization	- Sulfuric acid (H$_2$SO$_4$)
	- Methanol
	- Hexane
	- Sodium bicarbonate (NaHCO$_3$)
	- Sodium sulphate anhydrous (Na$_2$SO$_4$)
Nucleic acid extraction: ZymoBIOMICS™ MagBead DNA/RNA Kit	- ZymoBIOMICS ™ MagBinding Beads
	- DNA/RNA Shield ™
	- DNA/RNA Lysis Buffer
	- DNA/RNA Prep Buffer
	- MagBead DNA/RNA Wash 1
	- MagBead DNA/RNA Wash 2
	- DNase/RNase-Free Water
	- DNase I 3 (lyophilized) 250 U
	- DNA Digestion Buffer 4 ml 4 ml
	- Proteinase K 3 (lyophilized) & Storage Buffer

582 **Equipment**

Equipment	Company
CentriVap Benchtop Vacuum concentrator	Labconco
HP-5ms Ultra Inert Column	Aglient
7890A GC system	Aglient
Fastprep-96	MP
MiSeq System	Illumina

Experimental design and sample collection

Adult *Eisenia fetida* were collected from Vermigrand GmbH in Absdorf, Austria together with different
compost samples, designated Pre-Compost and Vermicompost. Pre-Compost (PC) is compost
consisting of horse manure and thermophilic compost, it is laid freshly onto the existing compost every
week to feed the worms. Vermicompost (VC) is worm castings and is used as an organic fertilizer.
Samples were collected on 3 specific days within a one-week period, for 2 separate weeks. This
sampling scheme was chosen to capture the possible effects within a week of composting, since new
Pre-compost is laid onto existing soil once a week.

Collected worms were kept in their natural soil composition until the dissection, taking place not more
than two hours later. For dissection, the worms were put into a petri dish and cleaned from residual
soil before being euthanized in hot water (70°C) for 20s. Worms were then dissected aseptically under

a stereomicroscope to recover soil from the intestinal tract (called "Gut soil" or GS). Gut soil from ten
to fifteen worms was collected to recover at least 275 mg wet gut soil per replicate. To collect castings
of *E. fetida*, worms were cleaned externally with deionized water and then separated into individual
culture boxes with ventilation for 48 hours. The castings produced the worms were collected, weighed,
and then pooled to recover at least 275 mg wet weight. Samples were frozen immediately after
collection, freeze dried on the next day, and stored at -20°C until further use.

Every sample was given a unique identifier for tracking the sample during the analysis as shown

Composition of the unique identifier for each sample with the first two letters accounting for the sample type followed by indicators for sampling week and day as well as its replicate

below:

Lipid extraction

The Folch extraction method was carried out following the procedure described by Folch et al. (65): 70
mg dry sample (PC, VC and GS) was extracted with 1.4 mL of chloroform:methanol (2:1) vortexing
for two minutes. The mixture was centrifuged at 4000 rpm for 10 min before collecting the
supernatant. The extraction process was carried out three times on the same sample and pooled. The
collected organic layers were purified washing with 1 mL dH_2O and a spatula tip of NaCl. The mixture
was vortexed for one minute and centrifuged at 4000 rpm for 10 minutes. The chloroform layer that
contains the extracted lipids was recovered and evaporated in a Labconco CentriVap Benchtop
Vacuum concentrator under reduced pressure at 10 °C. The lipid content was determined
gravimetrically and calculated as weight percentage of dry biomass. Lipid extracts obtained were
directly used for derivatization.

<u>Basic derivatization</u>
Basic derivatization was carried out by resuspending 10 mg lipid extracts in 100 µL of hexane. After
vortexing, to completely dissolve the extract, 50 µL of 2 M KOH solution in methanol was added and
vortexed for one minute. The sample was incubated for five minutes at room temperature. 125 mg of
sodium bisulphate was added to the sample, vortexed and centrifuged at 4000 rpm for five minutes.
100 µL of supernatant was collected, mixed with 400 µL of hexane and filtered through a 0.22 µm
PTFE filter. The filter was washed with an additional 100 µL of Hexane.

 <u>GC-MS method</u>

FAMEs were separated using a HP-5ms Ultra Inert Column (30 m x 250 μm x 0.25 μm) (Agilent, CA,

USA). 1 μL sample was injected in spitless mode and an injector temperature of 280 °C. The initial

oven temperature was set at 150 °C for 1 min and the temperature was gradually raised to 220 °C

at 3 °C/min with a final increase to 300 °C for 3 min. Helium was used as carrier gas at a constant

column flow rate of 2.52 mL/min. The GC to MS interface temperature was fixed at 280 °C and an

electron ionization system was set on the MS in scan mode. The mass range evaluated was 50–600

m/z, where MS quad and source temperatures were maintained at 150 °C and 230 °C respectively. To

search and identify each fatty acid, the NIST-MS Library (2.2) was used. It was also used to measure

the relative percentage of each compound and relative peak areas of the total ionic chromatogram

were used. Additionally, a commercial standard (Supelco 37 Component FAME Mix cat.#47885) was

used to identify peaks of the fatty acids by comparing retention times to those of known fatty acids in

the commercial standard.

<u>Statistical Analysis of identified fatty acids</u>

All extractions were performed in biological triplicate for each time point within a week. The results

were expressed as mean ± standard deviation while the effect of the extraction method and the

extraction conditions on lipid recovery was analyzed by using two-way ANOVA followed by Tukey

post-hoc test (statistically significance at $p < 0.05$). Statistical analysis was performed with GraphPad

Prism 8 (8.0.1) (GraphPad Software, San Diego, CA, USA). To test differences between the two

sampling weeks and to discriminate differences in fatty acid composition between samples, the data

were transformed by using center-log ratio and then evaluated using a Kolmogorov-Smirnov t-test and

Euclidean distance-based ordination respectively. Between-group differences were assessed by

permutational ANOVA by explaining community distances based by sampling week interacting with

sampling day and sample type: (Response = adonis2 (community data ~ sampling week * sampling

day * sample type)). Every analysis was performed with R version 4.1.1 using the "vegan" package

(v2.6-4) (66).

Nucleic acids extraction and sequencing

DNA and RNA extraction was carried out by using the ZymoBIOMICS[TM] MagBead DNA/RNA Kit (67).

Samples were weighed at roughly equivalent starting biomass (approximately 20 mg dry weight when

sample quantity allowed) into prefilled tube strips (8 x 12) containing 750 μl DNA/RNA Shield[TM]

(ZymoBIOMICS) and 40-400 μm glass beads (Macherey-Nagel) and arranged on a 96-well rack.

Homogenization was conducted on an MP Fastprep-96 with one minute on maximum speed and five

minutes rest, which was repeated five times. DNA and RNA were purified from the same input material

following the kit protocol, yielding two separate eluates for DNA and RNA respectively. The

concentrations were measured with the Qubit 1x dsDNA HS Assay Kit and RNA HS Assay Kit. The

detailed protocol for amplicon generation and sequencing are published by Pjevac et al. (68). In brief,

a two-step PCR was performed for amplicon generation and to add linkers as well as barcodes.

Amplicons were generated for the V4 region of the 16S (515F: GTG YCA GCM GCC GCG GTA A;

806R: GGA CTA CNV GGG TWT CTA AT) and 18S (TAReuk454FWD1: CCA GCA SCY GCG GTA

ATT CC; TAR eukREV3mod: ACT TTC GTT CTT GAT YRA TGA) rRNA genes as well as the PfaA-
KS domain region of the *pfa* gene (PfaA-KS-Fw: TGG GAA GAR AWT TCC C; PfaA-KS-Rv: GTR
CCN GTR CNG CTT C), part of the Pfa-Synthase complex (33). Linker sequences from primer to
barcode were forward GCT ATG CGC GAG CTG C and reverse TAG CGC ACA CCT GGT A. Cycling
conditions for the 16S and 18S rRNA amplicons were: initial denature 94° C for 3 mins, and then
denaturing at 95° C for 30 s, annealing at 55° C for 30 s, and extension at 72° C for 60 s. Cycling
conditions for the PfaA-KS amplicons were: initial denature at 94° C for 3 min, and then denaturing at
94° C for 45 s, annealing at 50.3° C for 30 s, and extension at 72° C for 90 s. The first PCR for all-
three primer sets was done with 30 cycles and 0,25 µM primer concentration while the second PCR
was done with 7 cycles for the 16S and 18S rRNA amplicons and 15 cycles for the PfaA-KS amplicons
using 0,8 µM primer concentration. Sequencing was performed by the Joint Microbiome Facility (JMF)
on an Illumina MiSeq using 2x300 base pair sequencing (69).

Analysis of amplicon sequencing data

In-house generated data

Input data were filtered for PhiX contamination with BBDuk (70). Demultiplexing was performed with
the python package demultiplex allowing 1 mismatch for barcodes and 2 mismatches for linkers (71).
Primers were verified with the python package demultiplex allowing 2 and 2 mismatches for forward
and reverse primers, respectively. Barcodes, linkers, and primers were trimmed off using BBDuk with
47 and 48 bases being left-trimmed for F.1/R.2 and F.2/R.1, respectively allowing a minimum length of
220 base pairs. Around 80% of reads across all samples passed all trimming, filtering and denoising
criteria.

Amplicon Sequence Variants (ASV) determination

ASVs for 16S and 18S sequences were generated by in-house sequencing facility using a standard
workflow with DADA2 (72). The PfaA-KS sequences were handled separately according to the
following steps: (i) Forward and reverse orientated sequences were kept separated based on primer
(F.1/F.2 and R.1/R.2), and merged using PEAR (73) with minimum overlap (-o) of 5, minimum trim
length (-t) of 300, maximum assembly length (-m) of 600, minimum quality score (-q) of 30, and
maximum uncalled bases allowed (-u) of 0; (ii) The forward-orientated FASTQ reads starting with the
reverse primer (R: merged from R1/R2) were reverse complemented, and then all merged sequences
were concatenated per sample and used for ASV calling DADA2; (iii) Filtering was conducted based
on quality profile plots: length truncated to 475 nt, maxN=0, maxEE=5, truncQ=3, rm.phix=TRUE. After
merging and filtering, 11 samples were discarded (01, 16, 24, 27, 40, 41, 42, 50, 52, 63, and 64)
because no reads passed filtering; (iv) The remaining 54 samples were used for error modelling and
ASV calling, yielding 66 unique biological PfaA-KS sequence ASVs. The sample ASV table and ASV
sequences were exported for further analyses.

Taxonomy Assignments

SSU rRNA ASVs were classified using SINA version 1.6.1 and the SILVA database SSU Ref NR 99
release 138.1 (74, 75). The PfaA-KS sequences were classified using BLASTn and BLASTx against
the nucleotide (nt) and non-redundant protein (nr) databases respectively and the top hit extracted for

putative assignment (Table S5). The results from BLASTn were primarily used unless the identity and

query coverage fell below 80%, in which the BLASTx results were consulted. Sixteen samples

remained with ambiguous assignments ("uncultured organism") and for these the BLASTx (nr) search

was narrowed to Bacterial references only, which resolved the assignments, with a minimum percent

identity of 81% and full query coverage (see Tables S6 & S7 for all BLAST output). Top-hits for each

amplicon variant were used for the final assignment, and the full taxonomy lineage was retrieved from

the Genome Taxonomy Database, release 202 metadata (76–79).

<u>Alignment and tree-building</u>

The PfaA-KS sequence variants from the original study (33) were accessed from the NCBI GenBank

PopSet 303162115 (446 total entries) and reannotated following the same procedures with BLASTn

and BLASTx as for our in-house generated PfaA-KS ASVs (Table S8). These sequences were

combined with our in-house generated PfaA-KS ASVs. The *Escherichia coli fabF* gene was used as

an outgroup to root the phylogeny. Sequences were aligned with MAFFT (v7.520) using the --localpair

option and --maxiterate 1000 (80, 81). Phylogenetic inference was made using IQ-TREE (multicore

version 1.6.12) (82–85), first using model selection, and then using the best-fit model with the following

parameters: -m TIM2e+R5 -bb 1000 -nt 4 -redo. The resulting tree was visualized using iTOL (86).

<u>Analysis and statistics</u>

All analysis on the ASV abundance tables were handled in R version 4.1.1 including the following

packages: {decontam} (v1.14.0) was used for removing contaminants (87); sampling depth was

normalised with {DESeq2} (v1.36.0) using a variance stabilizing transformation (88); compositional

transformations using center-log-ratio (CLR) and estimation of effect sizes were handled with {propr},

{vegan}, and {ALDEx2} (53, 89–92). An effect size >1 was considered as explanatory. The base R

{stats} was used for clustering samples as well as pair-wise comparisons with the Wilcoxon test.

Additionally, to reduce the rate of type-I errors, multiple testing corrections were done with the

Benjamini-Hochberg method. False discovery rate (FDR) < 0.05 was considered as statistically

significant (93). {phyloseq} (v1.40.0) was used for alpha- and beta-diversity calculation while

rarefaction for alpha diversity calculation was done with {vegan} (2.6-4) (89, 94). Heatplots were

generated with {made4} (v1.68.0) (95), {massageR} and {gplots} (96, 97); dataframe transformation

and reformatting were handled with {reshape2}, {dplyr} (v1.0.10), {tidyr} (v1.2.1) (98, 99), and plotting

conducted with {ggplot2} using {wesanderson} color palettes (100, 101).

References

1.　Eckhart S, Jörg H. 2004. Microbial Type I Fatty Acid Synthases (FAS): Major Players in a
Network of Cellular FAS Systems. Microbiol Mol Biol Rev 68:501–517.

2.　Parsons JB, Rock CO. 2013. Bacterial lipids: Metabolism and membrane homeostasis. Prog
Lipid Res 52:249–276.

3.　White SW, Zheng J, Zhang Y-M, Rock CO. 2005. THE STRUCTURAL BIOLOGY OF TYPE II
FATTY ACID BIOSYNTHESIS. Annu Rev Biochem 74:791–831.

4.　Huang C. 2001. Mixed-Chain Phospholipids: Structures and Chain-Melting Behavior. Lipids
36:1077–1097.

5.　Harayama T, Shimizu T. 2020. Roles of polyunsaturated fatty acids, from mediators to
membranes. J Lipid Res 61:1150–1160.

6.　Hau HH, Gralnick JA. 2007. Ecology and Biotechnology of the Genus Shewanella. Annu Rev
Microbiol 61:237–258.

7.　Valentine RC, Valentine DL. 2004. Omega-3 fatty acids in cellular membranes: a unified
concept. Prog Lipid Res 43:383–402.

8.　Calder PC. 2015. Marine omega-3 fatty acids and inflammatory processes: Effects,
mechanisms and clinical relevance. Biochim Biophys Acta - Mol Cell Biol Lipids 1851:469–484.

9.　Yoshida K, Hashimoto M, Hori R, Adachi T, Okuyama H, Orikasa Y, Nagamine T, Shimizu S,
Ueno A, Morita N. 2016. Bacterial Long-Chain Polyunsaturated Fatty Acids: Their Biosynthetic
Genes, Functions, and Practical Use. Mar Drugs 14:94.

10.　Zhang Y-M, Rock CO. 2008. Membrane lipid homeostasis in bacteria. Nat Rev Microbiol
6:222–233.

11.　Harwood JL. 2019. Algae: Critical Sources of Very Long-Chain Polyunsaturated Fatty Acids.
Biomolecules https://doi.org/10.3390/biom9110708.

12.　Stanley D, Kim Y. 2020. Why most insects have very low proportions of C20 polyunsaturated
fatty acids: The oxidative stress hypothesis. Arch Insect Biochem Physiol 103:e21622.

13.　Schlotz N, Roulin A, Ebert D, Martin-Creuzburg D. 2016. Combined effects of dietary
polyunsaturated fatty acids and parasite exposure on eicosanoid-related gene expression in an
invertebrate model. Comp Biochem Physiol Part A Mol Integr Physiol 201:115–123.

14.　Guillou H, Zadravec D, Martin PGP, Jacobsson A. 2010. The key roles of elongases and
desaturases in mammalian fatty acid metabolism: Insights from transgenic mice. Prog Lipid
Res 49:186–199.

15.　Herbst DA, Townsend CA, Maier T. 2018. The architectures of iterative type I PKS and FAS.

Nat Prod Rep 35:1046–1069.

16. Metz JG, Roessler P, Facciotti D, Levering C, Dittrich F, Lassner M, Valentine R, Lardizabal K,
Domergue F, Yamada A, Yazawa K, Knauf V, Browse J. 2001. Production of Polyunsaturated
Fatty Acids by Polyketide Synthases in Both Prokaryotes and Eukaryotes. Science (80-)
293:290–293.

17. Singh H, Creamer K, Chase A, Klau L, Sheila P, Jensen P. 2023. Metagenomic data reveals
type I polyketide synthase distributions across biomes. mSystems 8:e00012-23.

18. Kabeya N, Fonseca MM, Ferrier DEK, Navarro JC, Bay LK, Francis DS, Tocher DR, Castro
LFC, Monroig Ó. 2018. Genes for de novo biosynthesis of omega-3 polyunsaturated fatty acids
are widespread in animals. Sci Adv 4:eaar6849.

19. Twining CW, Bernhardt JR, Derry AM, Hudson CM, Ishikawa A, Kabeya N, Kainz MJ, Kitano J,
Kowarik C, Ladd SN, Leal MC, Scharnweber K, Shipley JR, Matthews B. 2021. The
evolutionary ecology of fatty-acid variation: Implications for consumer adaptation and
diversification. Ecol Lett 24:1709–1731.

20. Pilecky M, Závorka L, Arts MT, Kainz MJ. 2021. Omega-3 PUFA profoundly affect neural,
physiological, and behavioural competences – implications for systemic changes in trophic
interactions. Biol Rev 96:2127–2145.

21. Ishikawa A, Kabeya N, Ikeya K, Kakioka R, Cech JN, Osada N, Leal MC, Inoue J, Kume M,
Toyoda A, Tezuka A, Nagano AJ, Yamasaki YY, Suzuki Y, Kokita T, Takahashi H, Lucek K,
Marques D, Takehana Y, Naruse K, Mori S, Monroig O, Ladd N, Schubert CJ, Matthews B,
Peichel CL, Seehausen O, Yoshizaki G, Kitano J. 2019. A key metabolic gene for recurrent
freshwater colonization and radiation in fishes. Science (80-) 364:886–889.

22. Lee JM, Lee H, Kang S, Park WJ. 2016. Fatty Acid Desaturases, Polyunsaturated Fatty Acid
Regulation, and Biotechnological Advances. Nutrients https://doi.org/10.3390/nu8010023.

23. Carvalho CCCR, Caramujo MJ. 2012. Lipids of prokaryotic origin at the base of marine food
webs. Mar Drugs 10:2698–2714.

24. Mühlroth A, Li K, Røkke G, Winge P, Olsen Y, Hohmann-Marriott MF, Vadstein O, Bones AM.
2013. Pathways of lipid metabolism in marine algae, co-expression network, bottlenecks and
candidate genes for enhanced production of EPA and DHA in species of Chromista. Mar Drugs
11:4662–4697.

25. Ochsenreither K, Glück C, Stressler T, Fischer L, Syldatk C. 2016. Production Strategies and
Applications of Microbial Single Cell Oils   . Front Microbiol .

26. Tocher DR, Betancor MB, Sprague M, Olsen RE, Napier JA. 2019. Omega-3 Long-Chain
Polyunsaturated Fatty Acids, EPA and DHA: Bridging the Gap between Supply and Demand.
Nutrients https://doi.org/10.3390/nu11010089.

27. Garcia R, Pistorius D, Stadler M, Müller R. 2011. Fatty acid-related phylogeny of myxobacteria

as an approach to discover polyunsaturated omega-3/6 fatty acids. J Bacteriol 193:1930–1942.

28.   Sampedro L, Jeannotte R, Whalen J. 2006. Trophic transfer of fatty acids from gut microbiota
801   to the earthworm Lumbricus terrestris. Soil Biol Biochem 38:2188–2198.

802   29.   Gómez-Brandón M, Aira M, Lores M, Domínguez J. 2011. Changes in microbial community
803   structure and function during vermicomposting of pig slurry. Bioresour Technol 102:4171–4178.

804   30.   Sapkota R, Santos S, Farias P, Krogh PH, Winding A. 2020. Insights into the earthworm gut
805   multi-kingdom microbial communities. Sci Total Environ 727:138301.

806   31.   Budroni M, Mannazzu I, Zara S, Saba S, Pais A, Zara G. 2020. Composition and functional
807   profiling of the microbiota in the casts of Eisenia fetida during vermicomposting of brewers'
808   spent grains. Biotechnol reports (Amsterdam, Netherlands) 25:e00439.

809   32.   Huang K, Li F, Wei Y, Chen X, Fu X. 2013. Changes of bacterial and fungal community
810   compositions during vermicomposting of vegetable wastes by Eisenia foetida. Bioresour
811   Technol 150:235–241.

812   33.   Shulse CN, Allen EE. 2011. Diversity and distribution of microbial long-chain fatty acid
813   biosynthetic genes in the marine environment. Environ Microbiol 13:684–695.

814   34.   Pinkart HC, White DC. 1997. Phospholipid biosynthesis and solvent tolerance in Pseudomonas
815   putida strains. J Bacteriol 179:4219–4226.

816   35.   Okuyama H, Okajima N, Sasaki S, Higashi S, Murata N. 1991. The cis/trans isomerization of
817   the double bond of a fatty acid as a strategy for adaptation to changes in ambient temperature
818   in the psychrophilic bacterium, Vibrio sp. strain ABE-1. Biochim Biophys Acta - Lipids Lipid
819   Metab 1084:13–20.

820   36.   Heipieper HJ, Meinhardt F, Segura A. The cisâ☐☐trans isomerase of unsaturated fatty acids in
821   Pseudomonas and Vibrio☐: biochemistry, molecular biology and physiological function of a
822   unique stress adaptive mechanism. FEMS Microbiol Lett 229:1–7.

823   37.   Chatgilialoglu C, Ferreri C, Melchiorre M, Sansone A, Torreggiani A. 2014. Lipid geometrical
824   isomerism: from chemistry to biology and diagnostics. Chem Rev 114:255–284.

825   38.   Kolb M, Wichmann H, Schröder U. 2019. GC/MS-screening analyses of valuable products in
826   the aqueous phase from microwave-assisted hydrothermal processing of Lemna minor.
827   Sustain Chem Pharm 13:100165.

828   39.   Zeibich L, Schmidt O, Drake HL. 2019. Fermenters in the earthworm gut: do transients matter?
829   FEMS Microbiol Ecol 95.

830   40.   Liu D, Lian B, Wu C, Guo P. 2018. A comparative study of gut microbiota profiles of
831   earthworms fed in three different substrates. Symbiosis 74:21–29.

832   41.   Ginolhac A, Jarrin C, Robe P, Perrière G, Vogel TM, Simonet P, Nalin R. 2005. Type I
833   Polyketide Synthases May Have Evolved Through Horizontal Gene Transfer. J Mol Evol

834  60:716–725.

835 42. Jenke-Kodama H, Sandmann A, Müller R, Dittmann E. 2005. Evolutionary Implications of
836  Bacterial Polyketide Synthases. Mol Biol Evol 22:2027–2039.

837 43. Quilodrán B, Cortinez G, Bravo A, Silva D. 2020. Characterization and comparison of lipid and
838  PUFA production by native thraustochytrid strains using complex carbon sources. Heliyon
839  6:e05404.

840 44. Schnorr SL, Berry D. 2022. Lipid synthesis at the trophic base as the source for energy
841  management to build complex structures. Curr Opin Biotechnol 73:364–373.

842 45. Gemperlein K, Rachid S, Garcia R, Wenzel S, Muller R. 2014. Polyunsaturated fatty acid
843  biosynthesis in myxobacteria: different PUFA synthases and their product diversity. Chem Sci
844  5:1733–1741.

845 46. Tice H, Mayilraj S, Sims D, Lapidus A, Nolan M, Lucas S, Glavina Del Rio T, Copeland A,
846  Cheng J-F, Meincke L, Bruce D, Goodwin L, Pitluck S, Ivanova N, Mavromatis K, Ovchinnikova
847  G, Pati A, Chen A, Palaniappan K, Land M, Hauser L, Chang Y-J, Jeffries CD, Detter JC,
848  Brettin T, Rohde M, Göker M, Bristow J, Eisen JA, Markowitz V, Hugenholtz P, Kyrpides NC,
849  Klenk H-P, Chen F. 2010. Complete genome sequence of Nakamurella multipartita type strain
850  (Y-104T). Stand Genomic Sci 2:168–175.

851 47. Shulse CN, Allen EE. 2011. Widespread occurrence of secondary lipid biosynthesis potential in
852  microbial lineages. PLoS One 6:e20146.

853 48. Xia Y, Wang Y, Wang Y, Chin FYL, Zhang T. 2016. Cellular adhesiveness and cellulolytic
854  capacity in Anaerolineae revealed by omics-based genome interpretation. Biotechnol Biofuels
855  9:111.

856 49. Graells T, Ishak H, Larsson M, Guy L. 2018. The all-intracellular order Legionellales is
857  unexpectedly diverse, globally distributed and lowly abundant. FEMS Microbiol Ecol 94:fiy185.

858 50. Momper L, Aronson HS, Amend JP. 2018. Genomic Description of 'Candidatus
859  Abyssubacteria,' a Novel Subsurface Lineage Within the Candidate Phylum Hydrogenedentes
860  . Front Microbiol .

861 51. Boeuf D, Eppley JM, Mende DR, Malmstrom RR, Woyke T, DeLong EF. 2021.
862  Metapangenomics reveals depth-dependent shifts in metabolic potential for the ubiquitous
863  marine bacterial SAR324 lineage. Microbiome 9:172.

864 52. Kondrotaite Z, Valk LC, Petriglieri F, Singleton C, Nierychlo M, Dueholm MKD, Nielsen PH.
865  2022. Diversity and Ecophysiology of the Genus OLB8 and Other Abundant Uncultured
866  Saprospiraceae Genera in Global Wastewater Treatment Systems. Front Microbiol 13:917553.

867 53. Quinn T, Richardson MF, Lovell D, Crowley T. 2017. propr: An R-package for Identifying
868  Proportionally Abundant Features Using Compositional Data Analysis. Sci Rep 7.

869   54.   Hong, Wook S, Kim IS, Lee JS, Sub CK. 2011. Culture-Based and Denaturing Gradient Gel
870   Electrophoresis Analysis of the Bacterial Community Structure from the Intestinal Tracts of
871   Earthworms(Eisenia fetida). J Microbiol Biotechnol2011/09/28. 21:885–892.

872   55.   Sampedro L, Whalen JK. 2007. Changes in the fatty acid profiles through the digestive tract of
873   the earthworm Lumbricus terrestris L. Appl Soil Ecol 35:226–236.

874   56.   Poinar GO, Hess RT. 1977. Immune Responses in the Earthworm, Aporrectodea trapezoides
875   (Annelida), Against Rhabditis pellio (Nematoda) BT - Comparative Pathobiology: Volume 3
876   Invertebrate Immune Responses, p. 69–84. In Bulla, LA, Cheng, TC (eds.), . Springer US,
877   Boston, MA.

878   57.   Geier B, Oetjen J, Ruthensteiner B, Polikarpov M, Gruber-Vodicka HR, Liebeke M. 2021.
879   Connecting structure and function from organisms to molecules in small-animal symbioses
880   through chemo-histo-tomography. Proc Natl Acad Sci 118:e2023773118.

881   58.   Vestal JR, White DC. 1989. Lipid Analysis in Microbial Ecology. Bioscience 39:535–541.

882   59.   Menzel R, Geweiler D, Sass A, Simsek D, Ruess L. 2018. Nematodes as Important Source for
883   Omega-3 Long-Chain Fatty Acids in the Soil Food Web and the Impact in Nutrition for Higher
884   Trophic Levels. Front Ecol Evol 6.

885   60.   Allemann MN, Allen EE. 2018. Chapter One - Characterization and Application of Marine
886   Microbial Omega-3 Polyunsaturated Fatty Acid Synthesis, p. 3–32. In Moore, BSBT-M in E
887   (ed.), Marine Enzymes and Specialized Metabolism - Part B. Academic Press.

888   61.   Watts JL, Browse J. 2002. Genetic dissection of polyunsaturated fatty acid synthesis in
889   Caenorhabditis elegans. Proc Natl Acad Sci 99:5854–5859.

890   62.   Delong EF, Yayanos AA. 1986. Biochemical function and ecological significance of novel
891   bacterial lipids in deep-sea procaryotes. Appl Environ Microbiol 51:730–737.

892   63.   Tanaka T, Ikita K, Ashida T, Motoyama Y, Yamaguchi Y, Satouchi K. 1996. Effects of growth
893   temperature on the fatty acid composition of the free-living nematodeCaenorhabditis elegans.
894   Lipids 31:1173–1178.

895   64.   Brock TJ, Browse J, Watts JL. 2007. Fatty Acid Desaturation and the Regulation of Adiposity in
896   Caenorhabditis elegans. Genetics 176:865–875.

897   65.   FOLCH J, LEES M, SLOANE STANLEY GH. 1957. A simple method for the isolation and
898   purification of total lipides from animal tissues. J Biol Chem 226:497–509.

899   66.   2022. Community Ecology Package [R package vegan version 2.6-4]. Comprehensive R
900   Archive Network (CRAN).

901   67.   RESEARCH   Z.   ZymoBIOMICS   MagBead   DNA/RNA.
902   https://www.zymoresearch.com/products/zymobiomics-magbead-dna-rna.

68.   Pjevac P, Hausmann B, Schwarz J, Kohl G, Herbold CW, Loy A, Berry D. 2021. An Economical

and Flexible Dual Barcoding, Two-Step PCR Approach for Highly Multiplexed Amplicon
Sequencing. Front Microbiol 12:669776.

69.   MiSeq Reagent Kit v3 | Longest read lengths on any Illumina sequencer.
https://www.illumina.com/products/by-type/sequencing-kits/cluster-gen-sequencing-
reagents/miseq-reagent-kit-v3.html.

70.   SourceForge. BBMap. https://sourceforge.net/projects/bbmap/.

71.   GitHub.   GitHub   -   jfjlaros/demultiplex: Versatile   FASTA/FASTQ   demultiplexer.
https://github.com/jfjlaros/demultiplex.

72.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes S. 2016. DADA2: High-
resolution sample inference from Illumina amplicon data. Nat Methods 13:581–587.

73.   Zhang J, Kobert K, Flouri T, Stamatakis A. 2014. PEAR: a fast and accurate Illumina Paired-
End reAd mergeR. Bioinformatics 30:614–620.

74.   Pruesse E, Peplies J, Glöckner FO. 2012. SINA: accurate high-throughput multiple sequence
alignment of ribosomal RNA genes. Bioinformatics 28:1823–1829.

75.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glockner F. 2013.
The SILVA ribosomal RNA gene database project: improved data processing and web-based
tools. Nucleic Acids Res 41:D590–D596.

76.   Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil P-A, Hugenholtz P.
2018. A standardized bacterial taxonomy based on genome phylogeny substantially revises the
tree of life. Nat Biotechnol 36:996–1004.

77.   Parks DH, Chuvochina M, Chaumeil P-A, Rinke C, Mussig AJ, Hugenholtz P. 2020. A complete
domain-to-species taxonomy for Bacteria and Archaea. Nat Biotechnol 38:1079–1086.

78.   Rinke C, Chuvochina M, Mussig AJ, Chaumeil P-A, Davín AA, Waite DW, Whitman WB, Parks
DH, Hugenholtz P. 2021. A standardized archaeal taxonomy for the Genome Taxonomy
Database. Nat Microbiol 6:946–959.

79.   Parks DH, Chuvochina M, Rinke C, Mussig AJ, Chaumeil P-A, Hugenholtz P. 2022. GTDB: an
ongoing census of bacterial and archaeal diversity through a phylogenetically consistent, rank
normalized and complete genome-based taxonomy. Nucleic Acids Res 50:D785–D794.

80.   Katoh K, Misawa K, Kuma K, Miyata T. 2002. MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform. Nucleic Acids Res 30:3059–3066.

81.   Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol Biol Evol 30:772–780.

82.   Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: Improving the
Ultrafast Bootstrap Approximation. Mol Biol Evol 35:518–522.

83.   Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017. ModelFinder:
939   fast model selection for accurate phylogenetic estimates. Nat Methods 14:587–589.

940   84.   Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear
941   R. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the
942   Genomic Era. Mol Biol Evol 37:1530–1534.

943   85.   Minh BQ, Nguyen MAT, von Haeseler A. 2013. Ultrafast Approximation for Phylogenetic
944   Bootstrap. Mol Biol Evol 30:1188–1195.

945   86.   Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree
946   display and annotation. Nucleic Acids Res 49:W293–W296.

947   87.   Davis NM, Proctor DM, Holmes SP, Relman DA, Callahan BJ. 2018. Simple statistical
948   identification and removal of contaminant sequences in marker-gene and metagenomics data.
949   Microbiome 6:226.

950   88.   Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
951   RNA-seq data with DESeq2. Genome Biol 15:550.

952   89.   Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, Simpson GL, Solymos
953   P, Stevens MHH, Wagner H. 2013. vegan: Community Ecology Package. 2.0-8.

954   90.   Fernandes A, Macklaim J, Linn T, Reid G, Gloor G. 2013. ANOVA-Like Differential Gene
955   Expression Analysis of Single-Organism and Meta-RNA-Seq. PLoS ONE2 8:e67019.

956   91.   Fernandes A, Reid J, Macklaim J, McMurrough A, Edgell R, Gloor G. 2014. Unifying the
957   analysis of high-throughput sequencing datasets: characterizing RNA-seq, 16S rRNA gene
958   sequencing and selective growth experiments by compositional data analysis. Microbiome 2.

959   92.   Gloor G, Macklaim J, Fernandes A. 2016. Displaying Variation in Large Datasets: a Visual
960   Summary   of   Effect   Sizes.   J   Comput   Graph   Stat
961   https://doi.org/http://dx.doi.org/10.1080/10618600.2015.1131161.

962   93.   Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful
963   Approach to Multiple Testing. J R Stat Soc Ser B 57:289–300.

964   94.   McMurdie PJ, Holmes S. 2013. phyloseq: an R package for reproducible interactive analysis
965   and graphics of microbiome census data. PLoS One 8:e61217.

966   95.   Culhane AC, Thioulouse J, Perrière G, Higgins DG. 2005. MADE4: an R package for
967   multivariate analysis of gene expression data. Bioinformatics 21:2789–90.

968   96.   Stanstrup J. 2017. massageR: massageR. 0.7.2.

969   97.   Warnes GR, Bolker B, Bonebakker L, Gentleman R, Huber W, Liaw A, Lumley T, Maechler M,
970   Magnusson A, Moeller S, Schwartz M, Venables B. 2022. gplots: Various R Programming
971   Tools for Plotting Data. 3.1.3.

972   98.   Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, Grolemund G, Hayes A,
973   Henry L, Hester J, Kuhn M, Pedersen T, Miller E, Bache S, Müller K, Ooms J, Robinson D,
974   Seidel D, Spinu V, Takahashi K, Vaughan D, Wilke C, Woo K, Yutani H. 2019. Welcome to the
975   Tidyverse. J Open Source Softw 4:1686.

976   99.   Wickham H. 2016. Flexibly Reshape Data: A Reboot of the Reshape Package. CRAN.

977   100.   Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

978   101.   Ram K, Wickham H. 2018. wesanderson: A Wes Anderson Palette Generator. 0.3.6.

Figure 1. Principal component analysis (PCA) of the fatty acid content extracted from pre-compost (yellow triangle), gut soil (red dot) and vermicompost (blue square) after center-log-ratio transformation (CLR) with each loading arrow representing one fatty acid identified.

Figure 2. Alpha- and beta-diversity analysis comparisons for gut soil (GS, red), individual gut soil (IG, green), Pre-compost (PC, yellow) and Vermicompost (VC, blue). (**A**, **B**) Boxplots for Chao1 and Shannon alpha-diversity metrics with significant differences shown for different sample types with pair wise comparisons between groups by Wilcoxon test (p > 0.05 (ns), p ≤ 0.05 (*), p ≤ 0.01 (**), p ≤ 0.001 (***), p ≤ 0.0001 (****)). (**C**) Ordination of ASVs using Euclidean distance metrics shows separation of sample types. (D) Ward's clustering of rarefied ASVs binned at family- or lowest assigned taxa level and filtered for 2% abundance in at least 5% of samples.

Figure 3. Pair wise comparisons of each sample type (GS), individual gut soil (IG), Pre-compost (PC) and Vermicompost (VC) by Wilcoxon test with post Benjamini-Hochberg showing adjusted p-values for each comparison (p > 0.05 (ns), p ≤ 0.05 (*), p ≤ 0.01 (**), p ≤ 0.001 (***), p ≤ 0.0001 (****)). Blue color scale indicates that the mean of sample A is lower that in sample B. Red color scale indicates that the mean of sample A is greater than the mean of sample B.

Figure 4. Phylogenetic inference of 511 PfaA-KS amplicon sequences from this study (earthworm) and prior work investigating marine bacterial sources of PUFA synthesis. The fabF gene from *E. coli* was used as outgroup and root. Stacked barplots show the number of samples that contain the ASV sequence and are colored by the sample type.

Figure 5. Visualization of the sparse distribution and low abundance of 65 bacterial PfaA-KS ASVs by sample type based on center-log-ratio transformed ASV table (**A**); and hierarchical clustering of samples based on ASVs binned at lowest taxonomic assignment (**B**). Number of unique sequences shown across 5 phyla assignments shows clear dominance by Pseudomonadota, and primarily from the genus *Shewanella* (n=48) (**C**).

Figure 6. Comparison of ASV abundance tables for the PfaA-KS amplicons (**pfa**), and the subsetted PfaA-KS assigned taxa from the 16S rRNA dataframe (**prok.pfa**). Ordination and procrustes analysis on the transformed dataframes (**A**) indicates modest correlation, shown by the sum of squares (m12²) and correlation coefficient. The principal component plots of the CLR-transformed dataframes demonstrate this correlation from some interspersion of GS and VC samples, and with PC samples clustering separately. The abundance values of the binned taxa for **pfa** and **prok.pfa** show the distribution of taxa by sample (**B**). The abundance of *Shewanella* taxa were correspondent between datasets (**C**), shown separated by sample type, as there are two distinct linear relationships for PC and GS samples.

Figure 7. Binned taxa filtered for those achieving 1% abundance in 5% of samples in the rarefied ASV table plotted by Ward's clustering of scaled abundance shows a distinct division in relative enrichment of microeukaryotes. Compost samples PC and VC (yellow and blue respectively) are generally enriched across a larger number of taxonomic bins relative to GS samples (red).