Genes in Humans and Mice: Insights from Deep learning of 777K Bulk Transcriptomes

Zheng Su[1#], Mingyan Fang[2,3#], Andrei Smolnikov[1], Fatemeh Vafaee[1*] & Marcel E. Dinger[1,4*] & Emily C. Oates[1*]

[1] School of Biotechnology and Biomolecular Sciences, Faculty of Science, The University of New South Wales, Sydney, NSW 2052, Australia.

[2] BGI Research, Shenzhen 518083, China.

[3] BGI Australia, Herston, Queensland, Australia.

[4] School of Life and Environmental Sciences, Faculty of Science, University of Sydney, Sydney, NSW 2006, Australia.

[#] These authors contributed equally to this work.

[*] These corresponding authors contributed equally to this work, e-mails: f.vafaee@unsw.edu.au; marcel.dinger@sydney.edu.au; e.oates@unsw.edu.au

Abstract

Mice are widely used as animal models in biomedical research, favored for their small size, ease of breeding, and anatomical and physiological similarities to humans[1,2]. However, discrepancies between mouse gene experimental results and the actual behavior of human genes are not uncommon, despite their shared DNA sequence similarity[3-8]. This suggests that DNA sequence similarity does not always reliably predict functional similarity. On the other hand, RNA-level gene expression could offer additional information about gene function[9,10]. In this study, we undertook characterization and inter-species comparison of human and mouse genes by applying innovative deep learning methodologies to a large dataset of 410K human and 366K mouse bulk RNA-seq samples. This was achieved by using gene representations from our Transformer-based GeneRAIN model[11,12]. These gene representations aggregate information from large gene expression datasets, and provide insights beyond DNA sequence similarity. We identified 2,407 human-mouse homologous genes with high DNA similarity but distinct RNA characteristics, and showed that these genes are more likely to have differing disease/phenotype associations between the two species. Additionally, we found 3,070 homologous genes with low similarity at both the DNA and RNA levels, suggesting the highest risk of discrepancies in study results between the two species. We propose that this approach will support future decision making around whether the mouse will be an appropriate model for studying specific human genes, and whether the results of specific mouse gene studies are likely to be recapitulated in humans. Our methodological innovations offer valuable lessons for future deep learning applications in cross-species omics data. The interspecies gene relationship findings from our study also contribute valuable insights into the gene biology and evolution of the two species.

Introduction

Mouse models are extensively used in various human biomedical research scenarios, including the study of disease mechanisms, validation of gene functions, and drug development[1,2,4,6,13]. The sequencing of both human and mouse genomes has dramatically increased our knowledge of gene relationships between these two species. It has enabled us to identify mouse homologous genes that could represent human genes using DNA sequence similarity information[14-16].

However, mouse genes selected based on DNA sequence similarity do not always mirror the function of human genes. Studies have shown discrepancies between outcomes in mouse models and human clinical settings[3-8]. These inconsistencies might arise from cellular, organ-level, or environmental differences between humans and mice. They might also stem from the limitations of using DNA sequences to infer gene function similarities. DNA provides a static blueprint of biology but may not fully capture the complexity of gene function, as genes typically function via the RNAs and proteins they encode[17]. RNAs may therefore offer an additional layer of information about gene function.

Although multiple mouse gene expression studies have been conducted, they have often relied on techniques such as dimension reduction, phylogenetic clustering, co-expression analysis, and differential expression analysis[16,18-33]. These traditional methods face challenges in achieving a comprehensive comparison of genes at the RNA level, as they can be susceptible to batch effects, biased by small sample sizes, and constrained by the limited availability of samples from matched biological conditions[33,34]. Given the dynamic and complex nature of gene expression, which varies across genders, ages, tissues, and conditions, a thorough characterization at the RNA level necessitates integrating data from diverse biological contexts and a large collection of samples.

In this study, our goal was to characterize the similarities and differences between human and mouse genes at the RNA level. This was achieved by integrating information from 410K human and 366K mouse bulk RNA-seq samples. Our approach leveraged advanced deep learning methods, including state-of-the-art Transformer models[11], self-supervised techniques[35-37], and the development of embedding alignment methodology.

Results

Representing genes across species using RNA information

In our study, we started by creating representations of protein-coding genes, by integrating gene expression information from 410,850 human and 366,750 mouse bulk RNA-seq samples from the ARCHS4 database[38]. These samples were from a wide range of tissues, genders, ages, and biological conditions. Our gene representations were created using the ARCHS4 RNA-seq datasets to train our state-of-the-art gene representation GeneRAIN model[12], a Transformer-based[11] Generative Pre-Training (GPT) model[37]. The gene representations were 200-number vector embeddings from the model. Prior study revealed that the gene embeddings generated by GeneRAIN model could effectively represent genes by encoding a comprehensive set of biological information[12]. This includes information about gene-associated diseases/phenotypes, protein interactions, transcription factors, biological pathways,

gene ontology, associated cell types and more. Throughout this text, we will use the
terms 'gene embedding' and 'gene representation' interchangeably.

In the study, we performed intensive methodological development and evaluation to
represent genes of the two species in the same space. Here, a 'space' is a
mathematical framework where similar numerical values in the gene representations
represent similar biological attributes. Conversely, if genes are represented in
separate spaces, the same numerical values can represent different attributes.
Traditional single-species models will represent genes of two species in distinct
spaces, making comparison challenging. Thus, for this study, developing methods to
represent genes from two species in the same space was crucial to make them
comparable.

We started with a mixed training approach, inspired by joint training techniques used
in aligning cross-lingual word embeddings[39]. In this approach, human samples were
randomly mixed with mouse samples to create a training dataset for a single
GeneRAIN model (Fig. 1a). In the model, 18,757 human genes and 19,163 mouse
genes had their own gene embeddings but shared other model parameters.
Throughout the training, although the model progressively 'pulled' the gene
embeddings of the two species closer (Extended Data Fig. 1a and 2, observe the
percentage of explained variance on the PCA plot x-axis), they were not in a same
space (Fig. 1d), indicating the need for further methodological enhancements.

Subsequently, we adopted a 'shared embeddings' approach to further align gene
embeddings, using the information of 16,983 one-to-one orthologous genes from the
MGI database[40]. As a significant portion of these orthologous genes should share
similar functions across the two species[41], they should have very similar gene
representations. Thus, we forced the 5,000 randomly selected one-to-one orthologous
gene pairs to have the same gene embeddings (Fig. 1b). We hoped to use these 5,000
gene pairs as anchors to further pull the representations of other genes closer together.
Note that only the representations of these other genes would be used for later
analyses. Three different models, each applying shared embeddings to a different set
of orthologs, were trained to make all genes have usable embeddings (Methods). Our
analysis indicated that, although this method did bring the gene embeddings closer
together, they remained in relatively distinct spaces (Fig. 1e).

Despite being in distinct spaces, we investigated whether the gene embeddings
already encoded information about inter-species gene relationships. This was done
by examining if orthologous genes from the two species had similar representations.
It was found that, for 2,459 of the 16,983 human gene examined, their mouse
orthologs were the nearest embedding neighbor among all mouse genes. Additionally,
for 6,007 human genes, their mouse orthologs were among the ten nearest embedding
neighbors (Fig. 1g). This suggests that the gene embeddings had indeed captured
some inter-species gene relationship information.

We further explored representation alignment strategies by testing the advanced
unsupervised and supervised embedding alignment methods from MUSE[42] (Fig. 1c),
on the embeddings from mixed training. These methods adopt approaches of
contrastive learning and Procrustes alignment[43], and find a single transformation
matrix to align all the genes. Assessment revealed that unsupervised method had
limited efficacy (1,932 orthologs being the nearest, Fig. 1g). However, the supervised

method successfully made 4,216 human orthologs the nearest neighbor, and 8,140
human orthologs ranked among the top 10 closest embeddings (Fig. 1f, g).

Additionally, to evaluate how the result can be impacted by different alignment
techniques, we assessed the correlation of embedding similarities from different
approaches, focusing on one-to-one orthologous gene pairs. Strong correlation was
observed between supervised alignment and unsupervised alignment approaches,
especially in the high similarity orthologs (Spearman correlation rho = 0.89, p < 1e-20,
Fig. 1h). The high agreement with the unsupervised approach indicates low risk of
overfitting in the supervised approach. We also compared similarities between shared
embeddings and supervised alignment approaches. Despite lower overall similarities
from the shared embedding approach, strong correlation was observed (Spearman
correlation rho = 0.74, p < 1e-20, Extended data Fig. 3), indicating gene relationship
results were robust to the used embedding alignment techniques.

To further optimize and evaluate embedding alignment methodologies, we performed
multiple experiments, as detailed in Supplementary Information. Briefly, it was
revealed that combining shared embeddings with supervised alignment methods
slightly further enhanced the performance. Our 'positive control' experiment
demonstrated the high effectiveness of our embedding alignment approach in
reconstructing inter-species gene relationships. The experiment indicated that it was
not due to methodology limitations that fewer than 5,000 of 16,983 mouse genes were
the nearest embedding neighbors of their human orthologs. Additionally, a 'negative
control' experiment confirmed that our approach did not arbitrarily align genes in the
supervision set regardless of their embeddings. We also showed the alignment results
were not limited by training epochs or number of orthologous gene pairs for
supervision, and they were robust to Transformer architectures. Moreover, mixed
training proved beneficial for embedding alignment.

To mitigate bias from a single approach, we averaged the gene pair similarities from
shared embeddings plus supervised alignment approach and supervised alignment-
only approach. We used the averaged similarities for subsequent analyses. Our
comprehensive optimization and evaluation of embedding alignment methodologies
created a solid, reliable foundation for future analysis and application of these gene
representations.

RNA embeddings and DNA attributes in homologous genes

With human and mouse gene representations learned from expression data in a
shared space, we analyzed inter-species gene similarities. We observed varying
degrees of representation similarity (termed as RNA similarity or RNA simi.) among
human-mouse homologous gene pairs. To facilitate comparison, for each pair, we
determined the rank of the mouse gene based on its similarity to the corresponding
human gene, compared to all mouse genes. We then categorized these homologs into
two groups based on their ranks: a high similarity group comprising the top 40% of
homologs (n = 7,249, rank <= 3) and a low similarity group including the bottom 40%
(n = 7,103, rank >= 21). We excluded the middle 20% to avoid ambiguity in the
similarity property.

We explored the DNA attributes of two homolog groups. We examined DNA sequence
similarities, which we termed as DNA similarity or DNA simi., in coding regions, as well

as 1kb upstream and downstream regions of transcription sites through pairwise
transcript level alignment. Our analysis revealed that homologs with high RNA
embedding similarity generally exhibited high DNA similarity in all these regions,
though the overall DNA similarity was lower in the non-coding regions (Methods, Fig.
2a-c, Extended Data Fig. 6a-c and Supplementary Table 1-4). We also noted that
homologs with high RNA similarity have higher GC content (Fig. 2d and Extended
Data Fig. 7a-e), indicating stronger guanine-cytosine base pairs and greater molecule
stability[44]. Furthermore, high RNA similarity homologs tended to have slightly longer
coding sequences (Fig. 2e and Extended Data Fig. 6d).

RNA similarity and phenotype associations

Discrepancies between human and mouse gene study results could arise because
human and mouse homologs do not share the same disease or phenotype
associations. We explored whether gene representations derived from RNA
information could better explain phenotype association than DNA sequence similarity.
For this purpose, homologs were further categorized into high (top 40%, based on
coding region DNA alignment scores) and low (bottom 40%) DNA similarity groups.
We identified homologs with high similarity in both DNA and RNA (n = 4,694), in only
RNA (n = 1,107), in only DNA (n = 2,407), and in neither (n = 3,070).

For this investigation, we used mouse/human orthology phenotype annotations from
the MGI database[40] (Methods). Analysis revealed that homologs with high similarity in
both RNA and DNA had the largest number of shared associated phenotypes. This
was followed by homologs with high RNA but low DNA similarity, then those with low
RNA but high DNA similarity, and finally, homologs with low similarity in both (Fig. 3a
and Supplementary Table 2). This suggests RNA similarity provides information not
always captured by DNA similarity. Moreover, homologs with low DNA similarity can
have more shared associated phenotypes than those with high DNA similarity, if the
former have higher RNA similarity. A similar pattern was observed when analyzing the
percentage of homologs with shared associated phenotypes (Extended Data Fig. 8a).

To mitigate bias from using a single phenotype annotation dataset, we analyzed
another dataset, specifically the 'mouse models of human diseases' annotations from
the MGI database. This dataset includes information about human diseases, their
mouse models, and the associated genes. Given the low number of shared associated
diseases among the homologous genes (Extended Data Fig. 8b), we focused on
comparing proportion of each gene group with shared disease association(s). Results
indicated that homologs with high RNA similarity have largest proportion of shared
association(s). In contrast, homologs with low RNA similarity showed a smaller
proportion, even if their DNA similarity is high (Fig. 3b and Supplementary Table 2).

We extended our analysis using the third phenotype association annotations,
specifically the Alliance of Genome Resources database[45]. As association type and
evidence type information is available in this annotation dataset, we filtered the
association annotations to retain only those with well-supported evidence (Methods).
This analysis reaffirmed similar patterns of shared disease associations (Fig. 3c,
Extended Data Fig. 8c and Supplementary Table 2), reinforcing the notion that RNA
similarity offers additional, important information not captured by DNA sequence
similarity.

We then compared the performance of our embedding similarities for phenotype
association with an existing, state-of-the-art human-mouse gene conservation metric
called LECIF[33]. LECIF assesses human-mouse genomic functional conservation by
integrating epigenomic, transcription factor binding, and transcriptomic data using
contrastive learning neural networks. It incorporates data on over 8,000 human and
3,000 mouse features from consortia like ENCODE[46], Roadmap Epigenomics[47], and
FANTOM[48]. Our results showed that our representation similarities outperformed
LECIF in predicting shared phenotype and disease associations of human-mouse
homologs. (Fig. 3d-f, Extended Data Fig. 8d-f). This suggests that our approach,
although relying solely on transcriptome data, can achieve superior performance
compared to models that incorporate multi-omics data.

Given the predictive ability of our RNA embedding similarities, this approach could
potentially help explain discrepancies observed in mouse model studies. For example,
transgenic mouse models of certain genetic disorders may only partially replicate
human disease phenotypes, or therapeutics targeting some genes or their encoding
proteins may show promise in mice but fail to result in a positive clinical outcome in
humans. Examples include discrepancies observed in studies of SOD1 (similarity rank
171; rank ≥ 21 was considered as low similarity), C9orf72 (rank 89) for Amyotrophic
Lateral Sclerosis[49,50], PSEN2 (rank 587) for Alzheimer's Disease[51,52], PARK7
(encodes DJ-1, rank 105) for Parkinson's disease[8,53-55], and IL4 (rank 528), NOS2
(rank 50), and PTAFR (rank 33) for asthma[8,56]. These genes have low RNA similarities
in mice, suggesting that the correlation of these genes with other genes and possibly
their functions have diverged between mice and humans. This divergence could
contribute to the discrepancies observed between the human and mouse studies.

Modelling lncRNAs and pseudogenes across species

Finally, we expanded our analysis to include 20,184 long noncoding RNA (lncRNA)
genes and 8,439 pseudogenes alongside protein-coding genes in the model
(Extended Data Fig. 1e). We employed mixed training and supervised alignment for
the expanded geneset. We found that the similarities obtained were highly correlated
with those from the protein-coding only model (Fig. 4a), indicating consistent
performance when modelling more genes.

Next, we classified the genes into seven groups: protein-coding genes, lncRNAs, and
five pseudogene groups (Fig. 4b). For each group, we identified the most similar
mouse gene for each human gene based on gene embedding. To avoid bias,
orthologs used for supervision in supervised alignment were excluded when assessing
similarity. The analysis revealed that protein-coding genes had the highest similarity
between the two species (Fig. 4b), suggesting a conservation of their characteristics
at the RNA level. Processed pseudogenes and lncRNAs followed, and the other
pseudogene groups had the lowest similarity, indicating more expression-level
variability between species. These RNA similarity insights could be valuable for
understanding gene conservation at the RNA level, and potentially provide new
information about gene functions and attributes. Furthermore, they could assist in
identifying appropriate mouse representative genes for human lncRNAs and
pseudogenes in biomedical research.

Discussion

In this study, we developed a novel deep learning approach to characterize and
compare human and mouse genes at the RNA level, using a large bulk RNA-seq
dataset. Our thorough optimization and evaluation of gene embedding alignment
significantly improves the methodology and offers valuable lessons for future studies.
Our RNA-level representations offer insights beyond DNA sequence similarity, aiding
in selecting appropriate animal models for gene studies and improving our
understanding of genes in humans and mice.

Aligning gene embeddings from different species into the same space opens up the
possibility of using advanced deep learning technology for inter-species comparisons.
This approach could potentially be extended to aligning different biological conditions
and even different omics data. Our experiments demonstrate that mixed training,
shared embeddings, and supervised alignment techniques can improve alignment
quality. Importantly, our study found that alignment results were robust to the
alignment techniques used, making this a powerful approach even when some
aligning techniques may not be applicable in certain future scenarios.

The inter-species DNA and RNA similarities of genes do not fully overlap. Some
homologous gene pairs exhibited low DNA similarity but high RNA similarity,
suggesting that despite differences in DNA sequences, they still have similar
expression relationships with other genes. On the other hand, some gene pairs had
similar DNA sequences but low RNA similarity, suggesting that their functions may
have diverged. Therefore, we suggest exercising caution when selecting animal
models or interpreting results from mouse models for studying genes with low RNA
similarity, particularly those with low similarity in both DNA and RNA.

It's important to note that our RNA similarities reflect the average across tissues and
biological conditions. In certain applications, it may be beneficial to use the RNA
similarity for a specific tissue or a specific biological condition. While not explicitly
tested, it is theoretically feasible to obtain tissue- or condition-specific similarities by
adding corresponding expression embeddings to the gene embeddings, as this is the
operation adopted by the Transformer model during input, and our GeneRAIN study
provided preliminary results supporting this method[12]. Although our similarities are the
averaged similarities, they still provide valuable information about gene function.
Furthermore, our averaged similarities can be helpful for studying the evolution of the
two species and potentially enhance our understanding of their gene biology.

References

Peters, L. L. *et al.* The mouse as a model for human biology: a resource guide for complex trait analysis. *Nature Reviews Genetics* **8**, 58-69, doi:10.1038/nrg2025 (2007).

Bryda, E. C. The Mighty Mouse: the impact of rodents on advances in biomedical research. *Mo Med* **110**, 207-211 (2013).

Seok, J. *et al.* Genomic responses in mouse models poorly mimic human inflammatory diseases. *PNAS* **110**, 3507-3512, doi:doi:10.1073/pnas.1222878110 (2013).

Cummings, J. Lessons Learned from Alzheimer Disease: Clinical Trials with Negative Outcomes. *Clin Transl Sci* **11**, 147-152, doi:10.1111/cts.12491 (2018).

Stortz, J. A. *et al.* Murine Models of Sepsis and Trauma: Can We Bridge the Gap? *ILAR Journal* **58**, 90-105, doi:10.1093/ilar/ilx007 (2017).

McGonigle, P. & Ruggeri, B. Animal models of human disease: challenges in enabling translation. *Biochem Pharmacol* **87**, 162-171, doi:10.1016/j.bcp.2013.08.006 (2014).

Leenaars, C. H. C. *et al.* Animal to human translation: a systematic scoping review of reported concordance rates. *Journal of Translational Medicine* **17**, 223, doi:10.1186/s12967-019-1976-2 (2019).

Marshall, L. J., Bailey, J., Cassotta, M., Herrmann, K. & Pistollato, F. Poor Translatability of Biomedical Research Using Animals — A Narrative Review. *Alternatives to Laboratory Animals* **51**, 102-135, doi:10.1177/02611929231157756 (2023).

Zhang, W. *et al.* The functional landscape of mouse gene expression. *J Biol* **3**, 21, doi:10.1186/jbiol16 (2004).

Normand, R. *et al.* Found In Translation: a machine learning model for mouse-to-human inference. *Nat Methods* **15**, 1067-1073, doi:10.1038/s41592-018-0214-9 (2018).

Vaswani, A. *et al.* Attention is All you Need. *Advances in Neural Information Processing Systems* **30** (2017).

Zheng, S. *et al.* Multifaceted Representation of Genes via Deep Learning of Gene Expression Networks. *bioRxiv*, 2024.2003.2007.583777, doi:10.1101/2024.03.07.583777 (2024).

Hickman, D. L., Johnson, J., Vemulapalli, T. H., Crisler, J. R. & Shepherd, R. Commonly used animal models. *Principles of animal research for graduate and undergraduate students*, 117 (2017).

Chinwalla, A. T. *et al.* Initial sequencing and comparative analysis of the mouse genome. *Nature* **420**, 520-562, doi:10.1038/nature01262 (2002).

Schneider, V. A. *et al.* Evaluation of GRCh38 and de novo haploid genome assemblies demonstrates the enduring quality of the reference assembly. *Genome Res* **27**, 849-864, doi:10.1101/gr.213611.116 (2017).

Baldarelli, R. M. *et al.* The mouse Gene Expression Database (GXD): 2021 update. *Nucleic Acids Research* **49**, D924-D931, doi:10.1093/nar/gkaa914 (2020).

An integrated encyclopedia of DNA elements in the human genome. *Nature* **489**, 57-74, doi:10.1038/nature11247 (2012).

Schaum, N. *et al.* Single-cell transcriptomics of 20 mouse organs creates a Tabula Muris. *Nature* **562**, 367-372, doi:10.1038/s41586-018-0590-4 (2018).

19   Yu, Y. *et al.* A rat RNA-Seq transcriptomic BodyMap across 11 organs and 4
developmental stages. *Nat Commun* **5**, 3230, doi:10.1038/ncomms4230 (2014).

20   Abascal, F. *et al.* Expanded encyclopaedias of DNA elements in the human and
mouse genomes. *Nature* **583**, 699-710, doi:10.1038/s41586-020-2493-4 (2020).

21   Söllner, J. F. *et al.* An RNA-Seq atlas of gene expression in mouse and rat normal
tissues. *Scientific Data* **4**, 170185, doi:10.1038/sdata.2017.185 (2017).

22   Hu, Y. *et al.* Omic horizon expression: a database of gene expression based on RNA
sequencing data. *BMC Genomics* **24**, 674, doi:10.1186/s12864-023-09781-9 (2023).

23   Leach, S. M. *et al.* Human and Mouse Transcriptome Profiling Identifies Cross-
Species Homology in Pulmonary and Lymph Node Mononuclear Phagocytes. *Cell*
Rep **33**, 108337, doi:10.1016/j.celrep.2020.108337 (2020).

24   Mortazavi, A., Williams, B. A., McCue, K., Schaeffer, L. & Wold, B. Mapping and
quantifying mammalian transcriptomes by RNA-Seq. *Nat Methods* **5**, 621-628,
doi:10.1038/nmeth.1226 (2008).

25   Brawand, D. *et al.* The evolution of gene expression levels in mammalian organs.
Nature **478**, 343-348, doi:10.1038/nature10532 (2011).

26   Merkin, J., Russell, C., Chen, P. & Burge, C. B. Evolutionary dynamics of gene and
isoform regulation in Mammalian tissues. *Science* **338**, 1593-1599,
doi:10.1126/science.1228186 (2012).

27   Lin, S. *et al.* Comparison of the transcriptional landscapes between human and mouse
tissues. *Proc Natl Acad Sci U S A* **111**, 17224-17229, doi:10.1073/pnas.1413624111
(2014).

28   Monaco, G., van Dam, S., Casal Novo Ribeiro, J. L., Larbi, A. & de Magalhães, J. P.
A comparison of human and mouse gene co-expression networks reveals conservation
and divergence at the tissue, pathway and disease levels. *BMC Evolutionary Biology*
15, 259, doi:10.1186/s12862-015-0534-7 (2015).

29   Pervouchine, D. D. *et al.* Enhanced transcriptome maps from multiple mouse tissues
reveal evolutionary constraint in gene expression. *Nat Commun* **6**, 5903,
doi:10.1038/ncomms6903 (2015).

30   Sudmant, P. H., Alexis, M. S. & Burge, C. B. Meta-analysis of RNA-seq expression
data across species, tissues and studies. *Genome Biol* **16**, 287, doi:10.1186/s13059-
015-0853-4 (2015).

31   Yue, F. *et al.* A comparative encyclopedia of DNA elements in the mouse genome.
Nature **515**, 355-364, doi:10.1038/nature13992 (2014).

32   Breschi, A. *et al.* Gene-specific patterns of expression variation across organs and
species. *Genome Biol* **17**, 151, doi:10.1186/s13059-016-1008-y (2016).

33   Kwon, S. B. & Ernst, J. Learning a genome-wide score of human-mouse conservation
at the functional genomics level. *Nat Commun* **12**, 2495, doi:10.1038/s41467-021-
22653-8 (2021).

34   Breschi, A., Gingeras, T. R. & Guigó, R. Comparative transcriptomics in human and
mouse. *Nat Rev Genet* **18**, 425-440, doi:10.1038/nrg.2017.19 (2017).

35   Radford, A., Metz, L. & Chintala, S. Unsupervised Representation Learning with
Deep Convolutional Generative Adversarial Networks. *arXiv*,
doi:10.48550/arXiv.1511.06434 (2015).

36   Devlin, J., Chang, M.-W., Lee, K. & Toutanova, K. BERT: Pre-training of Deep
Bidirectional Transformers for Language Understanding. *arXiv*,
doi:10.48550/arXiv.1810.04805 (2018).

37   Radford, A., Narasimhan, K., Salimans, T. & Sutskever, I. Improving language
understanding by generative pre-training. (2018).

400	38	Lachmann, A. *et al.* Massive mining of publicly available RNA-seq data from human and mouse. *Nature Communications* **9**, 1366, doi:10.1038/s41467-018-03751-6 (2018).
403	39	Wang*, Z. *et al.* Cross-lingual Alignment vs Joint Training: A Comparative Study and A Simple Unified Framework. *OpenReview* (2019).
405	40	Blake, J. A. *et al.* Mouse Genome Database (MGD): Knowledgebase for mouse–human comparative biology. *Nucleic acids research* **49**, D981-D987 (2021).
407	41	Gabaldón, T. & Koonin, E. V. Functional and evolutionary implications of gene orthology. *Nature Reviews Genetics* **14**, 360-366, doi:10.1038/nrg3456 (2013).
409	42	Conneau, A., Lample, G., Ranzato, M. A., Denoyer, L. & Jégou, H. Word Translation Without Parallel Data. *arXiv*, doi:10.48550/arXiv.1710.04087 (2017).
411	43	Schönemann, P. H. A generalized solution of the orthogonal procrustes problem. *Psychometrika* **31**, 1–10, doi:10.1007/bf02289451 (1966).
413	44	Nakano, S. & Sugimoto, N. Roles of the amino group of purine bases in the thermodynamic stability of DNA base pairing. *Molecules* **19**, 11613-11627, doi:10.3390/molecules190811613 (2014).
416	45	Bult, C. J. & Sternberg, P. W. The alliance of genome resources: transforming comparative genomics. *Mamm Genome* **34**, 531-544, doi:10.1007/s00335-023-10015-2 (2023).
419	46	Frankish, A. *et al.* GENCODE 2021. *Nucleic Acids Res* **49**, D916-d923, doi:10.1093/nar/gkaa1087 (2021).
421	47	Kundaje, A. *et al.* Integrative analysis of 111 reference human epigenomes. *Nature* **518**, 317-330, doi:10.1038/nature14248 (2015).
423	48	Ramilowski, J. A. *et al.* Functional annotation of human long noncoding RNAs via molecular phenotyping. *Genome Res* **30**, 1060-1072, doi:10.1101/gr.254219.119 (2020).
426	49	Petrov, D., Mansfield, C., Moussy, A. & Hermine, O. ALS Clinical Trials Review: 20 Years of Failure. Are We Any Closer to Registering a New Treatment? *Front Aging Neurosci* **9**, 68, doi:10.3389/fnagi.2017.00068 (2017).
429	50	Mejzini, R. *et al.* ALS genetics, mechanisms, and therapeutics: Where are we now? *Front. Neurosci.* **13** (2019).
431	51	Donoviel, D. B. *et al.* Mice lacking both presenilin genes exhibit early embryonic patterning defects. *Genes Dev* **13**, 2801-2810, doi:10.1101/gad.13.21.2801 (1999).
433	52	Herreman, A. *et al.* Presenilin 2 deficiency causes a mild pulmonary phenotype and no changes in amyloid precursor protein processing but enhances the embryonic lethal phenotype of presenilin 1 deficiency. *Proceedings of the National Academy of Sciences of the United States of America* **96**, 11872-11877, doi:10.1073/pnas.96.21.11872 (1999).
438	53	Gibson, R., Dalvi, S. P. & Dalvi, P. S. DJ-1 and Parkinson's disease.
439	54	Hijioka, M., Inden, M., Yanagisawa, D. & Kitamura, Y. DJ-1/PARK7: A new therapeutic target for neurodegenerative disorders. *Biological and Pharmaceutical Bulletin* **40**, 548-552, doi:10.1248/bpb.b16-01006 (2017).
442	55	Bonifati, V. *et al.* Mutations in the DJ-1 gene associated with autosomal recessive early-onset parkinsonism. *Science* **299**, 256-259, doi:10.1126/science.1077209 (2003).
445	56	Nials, A. T. & Uddin, S. Mouse models of allergic asthma: acute and chronic allergen challenge. *Dis Model Mech* **1**, 213-220, doi:10.1242/dmm.000323 (2008).

Figure and Legends

Fig. 1 | Alignment of cross-species gene embeddings. a, Schematic of mixed training: Human RNA-seq samples were randomly mixed with mouse samples before being fed into a single Transformer model. Four samples are illustrated as examples. Lower boxes in each line denote gene symbols, while upper boxes indicate embedding indices. The colors represent gene expression level. Genes were ordered from highest to lowest expression prior to model input. Upper- and lower-case gene symbols represent orthologs in the two species. Different embeddings were used for genes in two species. **b,** Schematic of mixed training plus shared embedding strategy. Some randomly selected one-to-one orthologs from both species were forced to share identical embeddings (denoted by blue boxes). **c,** Supervised alignment approach. One-to-one orthologous gene pairs (connected by dashed lines) were used to supervise the alignment of gene embeddings from two species. **d to f,** PCA plots depicting the distribution of gene embeddings from three different alignment strategies, with each dot symbolizing a gene. **g,** Distribution of embedding similarity ranks for orthologs**.** For each human gene, the embedding similarity rank of its ortholog (where available) among all mouse genes was determined, and this histogram displays the

distributions of these ranks from three different embedding alignment strategies. **h,**
Correlation of orthologous gene similarities from the supervised alignment and
unsupervised alignment approaches. This scatter plot visualizes the Spearman
correlation of cosine similarities of one-to-one orthologous gene embeddings, from
supervised alignment and unsupervised alignment approaches. The dashed line
represents the line of equality (y=x).
473

474
Fig. 2 I DNA characteristics of homologs grouped by embedding similarity. a-c,
Boxplots depicting DNA alignment scores (Methods) for **(a)** coding sequences, **(b)** 1kb
upstream regions of transcription start sites, and **(c)** 1kb downstream regions of
transcription termination sites in homologs with high embedding similarity (n = 7,249)
and low embedding similarity (n = 7,103). Random gene pairs (n = 29,998) are
included for comparison. DNA alignment scores are determined by the highest
pairwise alignment similarity across transcripts for each gene pair, reflecting the extent
of DNA sequence conservation. **d,** GC content in the human genes from three gene
pair groups. **f,** Average coding sequence length in the human genes from three gene
pair groups. In all boxplots: center line, median; box limits, upper and lower quartiles;
whiskers, 1.5× interquartile range; outliers, points. All comparisons were evaluated
using a two-sided Wilcoxon rank-sum test. *** $p < 0.00001$, ** $p < 0.001$, * $p < 0.05$,
'ns' not significant ($p >= 0.05$).
488

Fig. 3 | Embedding similarity and phenotype association analysis. a, Empirical
cumulative distribution function (ECDF) plot illustrating the number of shared
phenotype associations for homologs, categorized by DNA similarity and embedding
similarity learned from RNA data. The horizontal axis represents the number of shared
associated phenotypes for each homolog gene pairs, while the vertical axis denotes
the fraction of homologs with a greater number of shared associated phenotypes.
Lines higher on the plot indicate more shared associated phenotypes. All group
comparisons had p-values less than 1e-14, except for the second vs. third group (p-
value = 0.0039), two-sided Kolmogorov-Smirnov test. **b,** Barplot displaying the
percentages of homologs with shared associated diseases, according to the human
disease mouse model annotations from MGI database. **c,** Barplot showing the
percentages of homologs with shared associated diseases, according to filtered
disease association annotations from the Alliance of Genome Resources database
(Methods). Error bars represent the standard errors in both barplots. In barplots, two-
sample z-test for proportions was used for comparisons. *** $p < 0.00001$, ** $p < 0.001$,
* $p < 0.05$, 'ns' denotes not significant ($p \geq 0.05$). **d-f,** F1 Score vs. threshold curves
comparing the performance of our RNA similarities and LECIF scores in predicting
whether homologous gene pairs have shared association(s), using annotations from
the database as indicated in the title. For LECIF, the average score within the genic
region was used for each gene. The curves visualize how the F1 score (harmonic
mean of precision and recall) varies across different classification thresholds. The p-
values, obtained from the Wilcoxon signed-rank test, compare the F1 scores of the
two metrics across 100 quantile thresholds.

513
514
515
516

517
Fig. 4 l Embedding similarities from the coding genes, lncRNAs and pseudogene
model. a, Spearman correlation of one-to-one orthologous gene embedding
similarities from the protein-coding gene-only model and the model including coding
genes, lncRNAs and pseudogenes. The dashed line represents the line of equality
(y=x). **b,** Distribution of embedding similarity for the closest gene across different gene
groups. Gene embeddings from the model of coding genes, lncRNAs, and
pseudogenes were used. In each gene group, for each human gene, the plot displays
the similarity with its closest mouse gene on the vertical axis. Note that genes used
for supervision in the supervised embedding alignment were excluded from the
protein-coding gene group in the plot. Boxplot details: center line indicates the median;
box limits represent the upper and lower quartiles; whiskers extend to 1.5× the
interquartile range; points beyond the whiskers are outliers.
In boxplots, two-sided Wilcoxon rank-sum test was used for comparisons. *** p <
0.00001, ** p < 0.001, * p < 0.05, 'ns' denotes not significant (p ≥ 0.05).
532
533

Methods

Data preprocessing

Gene-level human and mouse bulk RNA-seq data were retrieved from the ARCHS4 database (version 2.2)[38]. Data collection and processing adhered to ethical guidelines, under the supervision and approval of the University of New South Wales (UNSW) Human Research Ethics Committee. The data processing followed the methodology described in our previous GeneRAIN study[12]. Briefly, sample selection was based on stringent criteria: total gene expression read counts exceeding one million, a minimum of 2,000 genes exhibiting non-zero expression, and a single-cell probability below 0.5 (as estimated by the database) to eliminate potential single-cell samples. These criteria ensured the inclusion of high-quality samples.

The samples underwent library size normalization to have a uniform total read count of 10 million per sample. Subsequently, read counts were log-transformed (base 10), and only genes with a mean normalized expression above 0.1 were retained. This selection process resulted in a final dataset comprising 18,762 human protein-coding genes, 13,006 human lncRNA genes, 4,956 human pseudogenes, 19,072 mouse protein-coding genes, 7,178 mouse lncRNA genes, and 3,599 mouse pseudogenes.

Expression data discretization followed the procedure outlined in our previous GeneRAIN study. Gene expressions across all samples were categorized into 2,000 bins according to their expression ranks, using the Binning-By-Gene normalization method. This approach ensured that each bin, except the very first, contained equal number of samples. Before being fed into the Transformer GPT[37] or BERT[36] models, gene expressions in each sample were ordered from highest to lowest based on their bin assignments. The models received only the top 2048 genes, with attention paying to any genes in the inputs, including those with zero expression. In the BERT model specifically, 15% of genes exhibiting non-zero expression were randomly selected to have their identities (gene symbols) masked. The model was trained to reconstruct the identities of those genes.

Gene annotations

DNA sequence data was downloaded from Table Browser of UCSC Genome Browser[57]. For genic region annotation and gene transcript annotation, the GENCODE database was used[46]. Specifically, for humans, the GRCh38 assembly and GENCODE V44 were used. For mice, GRCm39 and GENCODE VM33 were used. Gene type annotations, such as annotations of being protein coding genes, lncRNA genes, etc., were primarily extracted from the GENCODE database. For genes whose information was not available in GENCODE database, gene type annotations from the metadata in the ARCHS4 data file was used. The information for mapping transcript IDs to gene names was obtained from Ensembl database Release 111 (https://ftp.ensembl.org/pub/)[58].

Information regarding human-mouse homologs and orthologs was obtained from the file 'HOM_MouseHumanSequence.rpt' in the MGI database[40], downloaded in January 2024. One-to-one homologs identified in this file were used as one-to-one orthologs.

Model hyperparameters

The configuration and parameters of the model were consistent with those detailed in our previous GeneRAIN GPT study. Briefly, each gene and its expression value were

represented in a 200-dimensional space. In both the BERT and GPT models, gene
embedding and expression embedding were added together. The architecture of the
model consisted of six layers, with each layer containing four attention heads and each
head had 32 dimensions. The AdamW optimizer was utilized for model
optimization[59,60], starting with a base learning rate of 0.00001 and allowing it to peak
at 0.0001. To optimize memory utilization and manage batch size effectively, gradient
accumulation steps were set at 5. The batch size was 12 for coding genes only model,
and batch size was set to 8 for models with coding gene, lncRNAs and pseudogenes
due to memory constraint. The dataset was divided into training (90%) and validation
(10%) sets.

Gene embedding alignment

In the mixed training approach, human samples were randomly mixed with mouse
samples to form the training dataset (Fig. 1a). Each mouse gene symbol was prefixed
with "m_" to distinguish it from human gene symbols. While human genes had gene
embeddings different from those of mouse genes, both shared the same expression
embeddings. Each training instance was either a single human sample or a single
mouse sample.

For the shared embedding training, three separate models were trained. Within each
model, 5,000 one-to-one orthologous gene pairs were randomly selected, and each
selected orthologous gene pair, consisting of a human gene and a mouse gene, was
assigned the same gene embedding. Importantly, the selection of 5,000 orthologous
gene pairs for each model was non-overlapping, ensuring that none of those gene
pairs were selected in more than one model. After training, we only used information
from the genes that did not have shared embeddings. The cosine similarity between
these genes was computed using their respective embeddings from each model. We
then averaged the similarities from the three models, and used them as the final
similarity values from this embedding alignment approach.

For unsupervised alignment, the program 'unsupervised.py' implemented by the
MUSE python package was used (https://github.com/facebookresearch/MUSE)[42]. The
training was conducted for 5 epochs. The function for selecting embeddings of the k
most frequent words for discrimination was disabled, and other parameters were set
to their default values.

In the case of supervised alignment, the program 'supervised.py' implemented by the
MUSE python package was employed (https://github.com/facebookresearch/MUSE).
A random selection of 95% (n=16134) of one-to-one orthologous gene pairs
constituted the training set, while the remaining 5% (n=849) served as the evaluation
set. Five refinement iterations were performed. All other parameters were maintained
at their default settings.

For the shared embedding plus supervised alignment approach, three shared
embedding models were first trained as previously described. Subsequently,
supervised alignment was applied to the genes without shared embeddings in each
model. We calculated the average gene-pair embedding similarities across the three
models, and used them as the final similarities from this approach.

Evaluation of embedding alignment

In the pseudo non-human 'positive control' experiment, we created a pseudo non-human species. The 410K human RNA-seq samples were divided into two halves, each comprising 205K samples. The first 205K samples were labeled as real human samples, and the remaining 205K samples were assigned to the pseudo non-human species. New gene symbols were 'invented' for each gene in the pseudo non-human species, and the mapping relationship with the actual human gene symbols was documented. That relationship was later used for embedding alignment supervision and evaluation of the alignment performance.

Different embeddings were assigned to the genes in the real human species and the genes in the pseudo non-human species. This resulted in embeddings for 18,757 genes from the real human species and another set of embeddings for 18,757 'invented' genes from the pseudo non-human species. The real human and pseudo non-human species underwent mixed training. Post mixed training, the documented mapping relationship between the actual human gene symbols and the 'invented' genes from the pseudo non-human species was used for supervision to align the embeddings from these two 'species'. Subsequently, we assessed the extent to which the mapping relationships were reconstructed by the alignment.

In the shuffled embeddings 'negative control' experiment, real human samples and real mouse samples first underwent mixed training as normal. Then prior to supervised alignment, the order of embeddings within each species was randomly shuffled, thereby disrupting the gene-embedding correspondence. Supervised alignment, utilizing the one-to-one orthologous gene pairs for supervision, was conducted on these shuffled embeddings. Subsequently, we assessed the extent to which the one-to-one orthologous relationships were reconstructed by the alignment.

For evaluating embedding alignment methodologies and experiments, the ranks of ortholog embedding similarity were examined. The process was performed as follows: for each human gene with a corresponding one-to-one mouse ortholog, its cosine similarity with every mouse gene was calculated using gene embeddings. Subsequently, all mouse genes were ranked based on these similarities, from highest to lowest. The rank of the corresponding mouse ortholog for the human gene was recorded. This procedure was repeated for every human gene (if a one-to-one mouse ortholog is available). The ranks for all the human genes were then used for assessing the closeness of embeddings between orthologous genes, and for evaluating the performance of the models or alignment methods.

Additionally, the ranking process could be reversed, starting with each mouse gene and examining the ranks of its human orthologs. Evaluations indicated that rankings in both directions are highly correlated, as detailed in Extended Data Fig. 5e, f.

DNA alignment

For the DNA alignment, the Biopython package was employed, specifically using PairwiseAligner function from Bio.Align module[61]. Global alignment was performed at the transcript level for each gene. Transcripts with a length of less than 100 base pairs (bp) were excluded from the analysis. For our pairwise alignments, following scoring system was used, the match score was set to 2, penalizing mismatches and gap openings with scores of -1 and -2, respectively, and a gap extension penalty of -1. The calculated scores for a pairwise alignment were normalized by the average length of

the two sequences being compared, to facilitate the comparison of alignment scores
across different gene pairs.

For aligning sequences from two genes, such as Gene A from humans and Gene B
from mice, following approached was used. For each transcript in Gene A, we
identified the best possible matched transcripts by finding the transcript of Gene B that
result in highest alignment. Subsequently, the maximum of scores from all transcripts
in Gene A were used as the DNA alignment score between Gene A and Gene B. While
the mean of scores from all transcripts in Gene A were used as the average transcript
alignment score. This approach provided a measure of similarity, taking into account
the splicing diversity of genes.

Phenotype association analysis

For our human and mouse gene phenotype association analysis, we used the
'HMD_HumanPhenotype.rpt' file from the the 'Mouse/Human Orthology with
Phenotype Annotations' section in the Mouse Genome Informatics (MGI) database
(https://www.informatics.jax.org/downloads/reports/index.html)[40]. From this file, we
extracted the number of high-level mammalian phenotypes associated with each gene
pair. The extracted numbers were then used for comparative analysis and
visualization.

For human disease mouse model association analysis, we obtained the annotation
data from the 'MGI_DiseaseGeneModel.rpt' file, which located in the 'Mouse models
of Human Disease by Human gene' section in the MGI database. This file provided
information on mouse models corresponding to human diseases. For each gene pair,
we counted the number of their associated 'DO term names', for comparative analysis
and visualization.

For Alliance disease association analysis, we obtained the annotation dataset from
the Alliance of Genome Resources website, Version: 6.0.0
(https://www.alliancegenome.org/downloads)[45]. Specifically, we used two
compressed tab-separated values (tsv) files: 'DISEASE-ALLIANCE_HUMAN.tsv.gz'
for human genes and 'DISEASE-ALLIANCE_MGI.tsv.gz' for mouse genes. This
dataset provided comprehensive information on gene-disease associations for both
human and mouse genes. We extracted gene symbols and associated disease names
from these files using columns 'DBObjectSymbol' and 'DOtermName', respectively.
Subsequently, as association type and evidence type information is available in this
dataset, we performed filtering to include only well-supported association records.

For human genes, we included association records with 'AssociationType' being
'is_implicated_in' or 'is_marker_for', and excluded those being
'implicated_via_orthology', 'biomarker_via_orthology', or 'is_not_implicated_in'.
Additionally, we applied filtering to the association evidence. We selected only records
where the 'EvidenceCodeName' included 'inference by association of genotype from
phenotype used in manual assertion', 'direct assay evidence used in manual assertion',
and 'mutant phenotype evidence used in manual assertion', excluding those supported
only by automatic assertions.

Similarly, for mouse genes, we selected records with 'DBobjectType' being 'gene'
(excluded 'allele' and 'affected_genomic_model') and 'AssociationType' being
'is_implicated_in' or 'is_marker_for'. We also applied filtering to the

'EvidenceCodeName', retaining only those being 'author statement supported by
traceable reference'. These filters provided us a dataset of high-confidence
association records, enabling us to perform association and comparative analyses on
homologs with different DNA and RNA similarities.

Comparison with LECIF scores

To obtain LECIF scores[33], we downloaded the bigwig format[62] file
'hg38.LECIFv1.1.bw' from https://github.com/ernstlab/LECIF and converted it to wig
format using the UCSC bigWigToWig tool[57]. Since LECIF scores are assigned to 50-
bp bins, we calculated gene-level LECIF scores to facilitate phenotype and disease
association comparisons. This was done by averaging LECIF scores for all positions
within each gene's genic region. We also calculated a separate set of scores using
only positions within coding regions. Genomic coordinates for both genic and coding
regions were obtained from GENCODE V44 annotation[46]. We then compared the
ability of both versions of calculated LECIF scores and our embedding similarities in
predicting whether homologous gene pairs shared phenotype(s) or disease
association(s).

Data availability

The supervised aligned gene embeddings derived from the protein-coding genes-only
GPT model, as well as from the model of coding genes, lncRNAs + pseudogenes, are
available at https://zenodo.org/records/10866876. These embeddings allow for
straightforward calculation of pairwise similarity. The 37,926 by 37,926 pairwise
similarity matrix, produced by averaging the similarities from the supervised alignment
approach and those from shared embeddings + supervised alignment approach, is
also available at the same link. Provided files are made available under the Creative
Commons Attribution Non Commercial 4.0 International License. All data is free to use
for non-commercial purposes.

Code availability

740 The source code used for training the models, aligning cross-species gene
741 embeddings, and performing all the analyses to generate the results in the study, is
742 available in GitHub repository https://github.com/suzheng/GeneRAIN_HM. Provided
743 code is made available under the Creative Commons Attribution Non Commercial 4.0
744 International License. All code is free to use for non-commercial purposes.

Method references

Raney, B. J. *et al.* The UCSC Genome Browser database: 2024 update. *Nucleic Acids Res* **52**, D1082-d1088, doi:10.1093/nar/gkad987 (2024).

Martin, F. J. *et al.* Ensembl 2023. *Nucleic Acids Research* **51**, D933-D941, doi:10.1093/nar/gkac958 (2022).

Loshchilov, I. & Hutter, F. Decoupled Weight Decay Regularization. *arXiv*, doi:10.48550/arXiv.1711.05101 (2017).

Kingma, D. P. & Ba, J. Adam: A Method for Stochastic Optimization. *arXiv*, doi:10.48550/arXiv.1412.6980 (2014).

Cock, P. J. A. *et al.* Biopython: freely available Python tools for computational molecular biology and bioinformatics. *Bioinformatics* **25**, 1422 (2009).

Kent, W. J., Zweig, A. S., Barber, G., Hinrichs, A. S. & Karolchik, D. BigWig and BigBed: enabling browsing of large distributed datasets. *Bioinformatics* **26**, 2204-2207, doi:10.1093/bioinformatics/btq351 (2010).

Acknowledgements

We would like to thank the Research Technology Services (ResTech) at the University of New South Wales for their help with computational resources, with a special thank you to Martin Thompson for his support and assistance. We also extend our thanks to the National Computational Infrastructure (NCI) for their support. This research was undertaken with the assistance of resources and services from the National Computational Infrastructure (NCI Australia), which is supported by the Australian Government through the National Collaborative Research Infrastructure Strategy (NCRIS). Z.S. acknowledges funding support from The University of New South Wales (University Postgraduate Award scholarship).

Author contributions

Z.S. and M.F conceptualized and designed the research, Z.S. conducted the data analysis. E.O., M.D. and F.V. provided project supervision and strategic guidance on research methodologies and analysis. Z.S. and M.F. generated visualizations, and led the writing of the manuscript. All authors contributed to the intellectual content, provided critical feedback, participated in revising the manuscript, and approved the final version for publication.

Competing interest declaration

F.V. declares commercial association with OmniOmics.AI Pty Ltd.

Additional information

Extended Data

Extended Data Fig. 1 l Training and validation loss of across epochs. This figure illustrates the training and validation loss over epochs for various models.

Extended Data Fig. 2 | PCA plots of gene embedding distributions over training
epochs. The plots illustrate the distribution of gene embeddings derived from the
mixed training GPT model at various training epochs. Each dot in the plots represents
a gene. Please note the percentage of explained variance on the x-axis.

Extended Data Fig. 3 | Embedding similarities from the shared embedding and
supervised alignment approaches. This scatter plot visualizes the Spearman

799
Extended Data Fig. 4 | Evaluation of embedding alignment strategies. a,
Distribution of embedding similarity ranks for one-to-one orthologs from supervised
alignment only and shared embedding plus supervised alignment approaches, with
statistical significance evaluated by a two-sided Kolmogorov-Smirnov test. **b,**
Spearman correlation of one-to-one orthologous gene embedding similarities from
these two alignment approaches. The dashed line represents the line of equality (y=x).
c, Distribution of embedding similarity ranks for one-to-one orthologs from the pseudo
non-human species in the 'positive control' experiment. **d,** Distribution of embedding
similarity ranks for orthologs from the shuffled embedding 'negative control'
experiment. **e,** Efficiency of supervised alignment with varying fractions of orthologous
gene pairs used for supervision, illustrated by the percentage of human genes with
their mouse orthologs ranking as the closest gene (blue line), and within the top 10
genes based on embedding similarity. **f,** Efficiency of supervised alignment across
different numbers of training epochs.

815
Extended Data Fig. 5 | Evaluation of embedding from different models. a,
Distribution of embedding similarity ranks for one-to-one orthologs from GPT and
BERT models, with mixed training and supervised alignment were used for both
models. **b,** Spearman correlation of one-to-one orthologous gene embedding
similarities from the GPT and BERT models. **c,** Distribution of embedding similarity
ranks for one-to-one orthologs from models with and without mixed training. GPT
model and supervised alignment were applied in both scenarios. **d,** Spearman
correlation of one-to-one orthologous gene embedding similarities from the models
with and without mixed training. **e,** Comparative analysis of human ortholog ranks in
two directions. The rank of each human gene's ortholog within all mouse genes was
calculated using embeddings from the shared embedding plus supervised alignment
approach. Similarly, the rank of each mouse gene's ortholog within all human genes
was also determined. This histogram visualizes the distribution of these two sets of

ranks. **f,** A scatter plot showing the correlation between these two sets of ranks,
assessed using Spearman's rho and p-value. In all barplots, statistical significance
was evaluated by a two-sided Kolmogorov-Smirnov test. In all scatter plot, each dot
symbolizing a orthologous gene pair. The dashed line represents the line of equality
(y=x).
834

835
Extended Data Fig. 6 l DNA characteristics of homologs grouped by embedding
similarity. a-c, Boxplots depicting mean transcript alignment scores (Methods) for **(a)**
coding sequences, **(b)** 1kb upstream regions of transcription start sites, and **(c)** 1kb
downstream regions of transcription termination sites in homologs with high
embedding similarity (n = 7,249) and low embedding similarity (n = 7,103). Mean
transcript alignment scores are calculated as the average pairwise alignment similarity
across the human transcripts for each gene pair (Methods), offering a complementary
perspective on DNA sequence conservation. Random gene pairs are included for
comparison. **d,** Average coding sequence length in the human genes from three gene
pair groups. In all boxplots: center line, median; box limits, upper and lower quartiles;
whiskers, 1.5× interquartile range; outliers, points. All comparisons were evaluated
using a two-sided Wilcoxon rank-sum test. *** $p < 0.00001$, ** $p < 0.001$, * $p < 0.05$,
'ns' not significant ($p >= 0.05$).
849

Extended Data Fig. 7 l GC content of homologs grouped by embedding similarity.
a, Boxplots visualize the GC content of the coding sequences of mouse genes from
homologs with high embedding similarity (n = 7,249) and low embedding similarity (n
= 7,103). Random gene pairs are included for comparative analysis. **b-e,** Analysis of
GC content in regulatory regions. With **b** for the 1kb upstream of transcription start
sites in human genes; **c** for the 1kb upstream of transcription start sites in mouse
genes; **d** for the 1kb downstream of transcription termination sites in human genes;
and **e** for the 1kb downstream of transcription termination sites in mouse genes. Each
analysis compares the two homolog groups and a random gene pair group. In all
boxplots: center line, median; box limits, upper and lower quartiles; whiskers, 1.5×
interquartile range; outliers, points. All comparisons were evaluated using a two-sided
Wilcoxon rank-sum test. *** *p* < 0.00001, ** *p* < 0.001, * *p* < 0.05, 'ns' not significant
(*p* >= 0.05).

865
Extended Data Fig. 8 | Embedding similarity and phenotype association analysis.
a Barplot displaying the percentages of homologs with shared associated phenotype
associations, based on ortholog phenotype association annotations from MGI
database. Homologs were categorized by DNA similarity and embedding similarity
learned from RNA data. Error bars represent the standard errors. **b,** Boxplot for
number of shared associated diseases across homolog groups, according to the
human disease mouse model annotations from MGI. **c,** Barplot showing number of
shared associated diseases across homolog groups, according to filtered disease
association annotations from the Alliance of Genome Resources database (Methods).
In all boxplots: center line, median; box limits, upper and lower quartiles; whiskers,
1.5× interquartile range; outliers, points. In barplots, two-sample z-test for proportions
was used for comparisons, while in boxplots, two-sided Wilcoxon rank-sum test was
used. *** $p < 0.00001$, ** $p < 0.001$, * $p < 0.05$, 'ns' not significant ($p >= 0.05$). **d-f,** F1
Score vs. threshold curves comparing the performance of our RNA similarities and
LECIF scores in predicting whether homologous gene pairs have shared
association(s), using annotations from the database as indicated in the title. For LECIF,
in contrast to Fig. 3, the average score within the coding region rather than genic
region was used for each gene. The curves visualize how the F1 score (harmonic
mean of precision and recall) varies across different classification thresholds. The p-
values, obtained from the Wilcoxon signed-rank test, compare the F1 scores of the
two metrics across 100 quantile thresholds.
887
888