Title: Trait distribution in Geometrid moth assemblages does not vary along a western Himalayan elevational gradient

Author: Pritha Dey

Author details: Wildlife Institute of India, Chandrabani, Dehradun-248001, India

Current address: LUOMUS-Finnish Museum of Natural History, Pohjoinen Rautatiekatu 13, 00100 Helsinki, Finland

 dey.pritha126@gmail.com

Abstract:

Elevation gradients pose challenges for flying insects due to shifts in temperature and air density, impacting their thermoregulation and flight abilities. Understanding how these challenges influence morphological traits across elevational gradients can provide insights into species distribution mechanisms. Geometrid moths, a model group for trait-based analysis among insects, show no significant variation in body size despite declining species diversity along a 1500m elevation span in the western Himalayas. Investigating 697 specimens of 120 Geometrid moth species, I assessed species diversity, turnover, and various traits related to body size and flight capabilities. Results indicate decreasing species diversity and increasing turnover with elevation. Although body length, wing loading, and maneuverability decrease with elevation, intraspecific variability remains insignificant, and trait overlap is high across elevations, suggesting no significant trait partitioning and niche differentiation among species. This supports the argument that species with similar ecological roles persist through turnover, maintaining consistency in trait values across assemblages. These findings align with previous inconsistencies in trait-environment relationships along elevational gradients in Geometrid moths, spanning neotropical and temperate regions. Additionally, the study reiterates the

importance of multivariate trait-based approaches in understanding species responses to

environmental changes within diverse clades, overcoming taxonomic challenges.

Introduction:

The inhabitant fauna of mountainous regions, including flying insects, encounters varying

physical parameters such as temperature and air density along elevation gradients (Dillon et

al., 2006). This range of physical environments induces notable shifts in species demands,

leading to observable variations in biological assemblages along the gradients.

At the nexus of trait-environment relationships are individual and species-level dynamics

within communities, spanning various spatial scales. It reveals how organisms, with distinctive

trait values, adapt to diverse environmental conditions (Gaston et al., 2009). The resultant trait

distribution among coexisting species sheds light on the process of community assembly

(Kearney et al., 2010; Lamanna et al., 2014). In low elevations, where warm temperatures and

ample resources diminish environmental constraints, species diversify into different niches.

Consequently, a lower overall trait overlap and narrower trait width facilitate the coexistence

of diverse species. Whereas at higher elevations with higher environmental filters, species

diversity decreases, and there is a higher overlap of trait space (Read et al., 2018).

Associated with the rules governing trait-based assemblages is the concept of species packing

within niche space, encompassing trait overlap (Kearney et al., 2010; Lamanna et al., 2014).

Whether trait overlap decreases or increases hinges on the interplay between niche packing and

the displacement of functional trait values over evolutionary time (Macarthur & Levins, 1967;

Ricklefs & O'Rourke, 1975). Nonetheless, the total niche occupied by an assemblage is

anticipated to increase with growing species richness, aligning with MacArthur's model

(MacArthur, 1965), where a 'niche' is defined as an n-dimensional hypervolume comprising

biotic and abiotic factors (Blonder et al., 2014). In the tropical Himalayan Mountain system,

bat communities in the western Himalaya and bird communities in the eastern Himalaya exhibit

higher niche overlap at lower elevations with greater species richness (Schumm et al., 2020;

Chakravarty et al., 2023). However, the applicability of these patterns to insect taxa across

elevational gradients and geographical regions remains unclear.

While Bergmann's rule established that species in colder regions should be larger for

thermoregulation (Bergmann, 1848), there exists a lack of consensus on this rule among

insects, exemplified by varied body size patterns across elevation and latitude gradients (Brehm

& Fiedler, 2004; Shelomi, 2012). For arthropods, particularly insects, the decrease in mean

temperature does not necessarily translate into analogous changes in body temperature

(Stevenson, 1985). The regulation of body temperatures is contingent upon factors such as

habitat, species shape, size, and behavior, all of which govern how species respond to

temperature fluctuations (Porter & Gates, 1969; Bakken, 1976; Samejima & Tsubaki, 2010).

The aforementioned variability in body size among insects reflects the intricate interplay

between ecological, physiological, and climatic factors. For instance, bees exhibit a decrease

in body size with increasing elevation (Osorio-Canadas et al., 2022.), contrasting with an

increase across latitudes (Gérard et al., 2018). Dragonflies and damselflies are generally larger

in temperate regions than in the tropics (Svensson et al., 2023). Geometrid moths, a model

group for many such studies, display diverse patterns, with some studies indicating a positive

relationship between body size and elevation in tropical and temperate elevational gradients

(Beck et al., 2016; Brehm et al., 2018), while others reveal no change in body size along an

elevational gradient in the neotropics (Brehm & Fiedler, 2004). Further complicating matters,

noctuid moths show larger body sizes at higher elevations in temperate regions (Heidrich et al.,

2021).

Predominantly, studies of trait-environment relationships in insects have centered on body size,

given its direct link to thermal sensitivity and other physiological and ecological traits (Chown

& Gaston, 2010). However, the influence of body size allometry extends beyond thermal

sensitivity, impacting flight-related morphology such as wing loading, and maneuverability

which directly predicts power requirements during flight in insects (Grula et al., 2021). The

morphological traits associated with flight in insects respond to the aerodynamic necessities

dictated by their specific environments (Bartholomew & Casey, 1977).

Insects at higher elevations are predicted to exhibit reduced flight capabilities due to alterations

in the aerodynamic forces generated by their wings (Dudley, 2002). To counteract the reduced

air density at higher elevations, insects employ strategies such as longer wings relative to body

size and greater wing area (Dudley, 2002; Dillon et al., 2006). Geometridae moths, the primary

focus of this study, exemplify this phenomenon with their large wing areas compared to body

mass (lower wing loading), facilitating energetically effective flight even at low temperatures

(Heinrich, 1974; Casey & Joos, 1983).

The Geometridae family, known for its diversity, serves as a model for ecological studies (Beck

et al., 2016; Brehm & Fiedler, 2004; Holm et al., 2019; Holm et al., 2019; Seifert et al., 2022)

has received comparatively little attention in the context of testing macroecological patterns in

the paleotropical montane systems. This study seeks to address this gap by examining

Geometrid moth communities across an elevation gradient of 1500-3000 m in the western

Himalaya, India. The study area, situated on the fringes of the Kedarnath Wildlife Sanctuary,

typifies a western Himalayan gradient with stark habitat changes over a short spatial scale.

The overarching objective of this study is to investigate how individual traits, variability of

traits among species, and niche overlap of Geometrid moth assemblages shift across an

elevational gradient, considering changing species diversity. Given the inconsistent findings

regarding the impact of elevation on body size in prior studies (Brehm & Fiedler, 2004; Beck

et al., 2016; Heidrich et al., 2021), the current research aims to ascertain whether these patterns

extend to the broader trait-space for insects in the Himalayan gradient. The hypothesis posits

that despite changes in species diversity, the body size and traits related to flight capabilities

will not differ significantly across the elevation. Therefore, no significant alterations will occur

in the overall trait-space at the assemblage level, and niche differentiation across the elevation

Materials and methods:

Study Area:

The study was done in 2018-2019 around the fringes of the Kedarnath Wildlife Sanctuary (KWLS)

(30°25′–30°41′N, 78° 55′–79°22′E) located in the Chamoli-Rudraprayag district in the state of

Uttarakhand. It is one of the most prominent protected areas in the western Himalaya (Map1),

home to the endangered Himalayan Musk deer (*Moschus chrysogaster*). KWLS lies in the upper

catchment area of the rivers Alakananda and Mandakini, which form the major tributaries of the

river Ganga, covering an area of 975 sq. km. The elevation of the protected area ranges from 1160-

7068 masl. Habitats range from mixed oak forests in the south to lush alpine meadows. The

combination of human pressure, pristine forest areas, and a large altitudinal range make it an ideal

site for biogeographic studies. The forest types are dominated by oak (*Quercus leucotrichophora,*

Quercus floribunda, Quercus glauca, and Quercus semecarpiflora).

Field sampling:

The study area was stratified based on elevation & vegetation types to sample the moth

diversity along the gradients. Sampling was done every 200m along the elevation from 1500m

up to 3100m. I studied the changes across elevation, considering it as a factor in three

catergories or classes. I categorized the sampling sites based on similar elevation and

vegetation types as follows: Khalla, Mandal, Gondi (below 2000m: Low Elevation); Jatholi,

Ansuya, Kanchula Kharak (2000-2500m: Mid Elevation); Bulkhan and Shokharak (>2500m:

High Elevation). At each site, 3-night sampling was done (details provided in the Table 1 of

the Supplementary material) at different times of the night to maximize species capture in that

area, as different species are active at different times of the night. The sampling sites are shown

in Map 1. On each sampling date, two LepiLED (Brehm, 2017) light traps were operated for

3-4 hours after dusk, depending on the weather conditions. Logistic constraints do not permit

all-night sampling. Sampling was done for a total ~140 hours from March to May in 2018. The

minimum distance between the two light traps was 50 m, with lamps not in line of sight of

neighbouring sites, so that cross-site sampling does not occur. The moths attracted to the trap

were photographed and representative samples from each species were collected as vouchers.

Collection was done with the aim of minimum three samples for each species, for

morphological measurements. The specimens were collected only from March to May in 2018,

as per the permit from the Forest Department. The specimens were then identified till species,

as much as possible. The voucher specimens have been submitted to the Research Collections

Facility at the National Centre for Biological Sciences, Bangalore, India.

Morphological trait measurement:

All voucher specimens were curated and photographed. We measured morphological traits on

the scaled photographs of 697 specimens of 120 morphospecies for morphometry using the

software Image J (Schneider et al., 2012). The measured traits were body length, wingspan,

forewing length, hindwing length, forewing area, hindwing area. I also calculated wing loading

[2*body length/ (forewing area + hindwing area)], and the ratio of hindwing area: to forewing

area which is roughly termed maneuverability (Arrizabalaga-Escudero et al., 2019). These

traits were measured because they are related to the body size and flight capability of moths,

which might govern their occurrence and sustenance across elevational gradients.

Analyses:

Species diversity

To show sample completeness, I calculated the rarefied sample coverage at each of the

elevation classes. I pooled the data collected from each light trapping night in each elevation

class and calculated the rarefied species diversity extrapolated to the number of individuals,

across the elevation classes. Both these analyses were done using the 'iNEXT' R package

(Hsieh et al., 2016).

For beta diversity across the elevation, I calculated the overall beta-diversity (Sørensen

pairwise dissimilarity), and species spatial turnover (Simpson's pairwise dissimilarity), all

measured as average dissimilarity of the observed units to their group centroid (Anderson et

al., 2006).

Variation in the individual trait value, intraspecific trait variability and niche overlap

For the subsequent analyses, the outliers in range of individual trait values were omitted, to

arrive at more robust patterns**.** The outliers were few and not abundant species and hence would

not affect the overall pattern. (details in Figure 4 and Table 2 of the Supplemenatary Material)

To look into intraspecific trait variation across the elevational gradient, I compared the

homogeneity of dispersion (Anderson et al., 2006) for the assemblages in each elevation

category. I used the *betadisper* function of the R package vegan (Oksanen et al., 2009) to

calculate dispersion, which tests whether the variance or dispersion of two or more groups is

significantly different or not based on Euclidean distance. This was followed by a permutest

for the homogeneity of multivariate dispersion. An ANOVA was then performed to identify

significance in the groups and a post hoc test to identify significant differences between specific

elevation categories. These beta values refer to a species' position along a gradient or

intraspecific trait variation defined by assemblage-level mean trait values (Ackerly &

Cornwell, 2007).

I used two approaches to examine the response of community-trait values of Geometrid moths

across the elevation classes. First, I investigated the response using community abundance-

weighted mean trait value (CWM; Lavorel et al., 2008), a widely used method to calculate

community trait values weighted by abundance of species in that community. Then I looked at

the response of the mean trait values for each of the traits across the elevation. The change in

both the community means with elevation was quantified using ordinary least squares

regression. It is useful to look at both the approaches as the relationships between community-

weighted mean (CWM) traits and elevation, is supposed to reflect selection towards locally

optimal traits, but it is limited by the large amount of interspecific trait variation typically found

within ecological communities (Muscarella & Uriarte, 2016).

To look at the niche overlap among species across the elevational gradient, body length, wing

loading, and the ratio of hindwing area: forewing area (Maneuverability) were chosen as the

least correlated variable. For this, I used O-statistic (Read et al., 2018) to generate a single

community-level (pooling the assemblages across the elevation) overlap index across multiple

traits by fitting non-parametric kernel density functions to the distribution of trait values within

an elevation (Mouillot et al., 2005; Geange et al., 2011; Read et al., 2018). The median pairwise

overlap was calculated by first determining the overlap of traits in each species pair, and then

taking the median overlap of each species pair in a community. This median overlap value is

called the O-statistic (O for overlap)

The R package *Ostats* (Read et al. 2022) was used to calculate O-statistic, the multivariate O-

statistic estimation in this package relies on functions imported from the package *hypervolume*

(Blonder et al., 2018). To test whether the degree of niche overlap among species in each

elevation category is greater than or less than expected by chance, I evaluated the O-statistic

against a null model, where the median of the pairwise hypervolume overlaps are weighted

against the abundances of the species pairs in each community or Standardized Effect Size (z-

scores).

Results:

Species diversity

Species diversity showed lower diversity in the high elevation (Fig.1b). The Euclidean distance

to centroid for Sørensen pairwise dissimilarity among the elevation categories were as follows:

Low-Mid (0.314); Low-High (0.489) and Mid-High (0.34); similarly, the Simpson

dissimilarity index (spatial turnover) was: Low-Mid (0.05); Low-High (0.14) and Mid-High

(0.12) (Fig 1 c and d). This means, the species belonging to the low elevation were quite

different (with higher spatial turnover) from the high elevation, than between Low and Mid,

and Mid and High elevation classes.

Variation in the individual trait value, intraspecific trait variability and niche overlap

The average distance (Euclidean) to centroid which gives a measure of the dispersion of traits

at the assemblage level, was higher in the lowest elevation (87.94) as compared to mid (78.63)

and high (76.14) elevations. The distances are not significantly different from each other

(ANOVA, $p>0.05$), which indicates that the intraspecific trait variation does not change across

the elevation gradient (Fig.2)

The ordinary least squares regression did not show any significant effect of elevation on the

Community-weighted means of traits across the elevation classes: Forewing Length ($p=0.57$),

Hindwing Length ($p=0.534$), Body Length ($p=0.31$), Wingspan ($p=0.14$), Forewing Area

($p=0.41$), Hindwing Area ($p=0.32$), Wing loading ($p=0.14$), Maneuverabilty ($p=0.51$) (Fig.3).

For the mean trait values, Forewing Length ($p=0.03$), Hindwing Length ($p=0.02$), and

Hindwing area ($p=0.03$) showed a significant decline at high elevation. Body Length ($p=0.19$),

Wingspan ($p=0.26$), Forewing Area ($p=0.23$), Wing Loading ($p=0.24$), Maneuverability

(p=0.19) did not show any significant change with elevation (Figure 2 in Supplementary

Material). There is high variability and a weak pattern in the CWM across elevation, which

makes it evident that the patterns are sensitive to the abundance of species. The species of same

trait values are abundant across the elevation. The density plots of individual traits added in the

Supplementary Material (Figure 3) would make it clearer.

There is a high level of niche overlap among the species of the Geometrid moth communities

across the elevation, as shown by high (0.84) O-statistic index of the multivariate trait space

(Fig. 4). This indicates that for each trait the observed niche overlap did not differ from that

which would be expected due to chance.

Discussion:

The findings on Geometrid moth assemblages along a 1500m gradient in the western Himalaya

provides valuable insights into the dynamics of individual traits, trait variability within the

assemblages, and niche overlap across elevational gradients in response to changing species

diversity. The results show a decrease in species diversity along the elevational gradient and

that there is consistent species turnover from the low to the high elevations. However, elevation

does not significantly impact morphological traits related to thermal sensitivity (body length)

and flight capability (wing-loading and maneuverability) at the assemblage level in Geometrid

moths.

Additionally, the study reveals the absence of significant intraspecific trait variation and

significant niche differentiation among species across the elevation. This sheds light on the

assemblage-level trait variability within Geometrid moth , revealing that ecological similarity

remains high across elevational gradients. Supporting these findings, evidence from the

Paleotropical region further underscores the lack of consistent trends in individual traits studied

in Geometrid moths and other insect taxa across diverse elevation ranges in neotropic and

palearctic regions (Brehm & Fiedler, 2004b; Peters et al., 2016; Beck et al., 2016; Classen et

al., 2017).

Species diversity across elevation

Here, it is evident that trait-level patterns may not directly correlate with the occurrence,

diversity, and abundance of species. Geometrid moths across mountains either show a mid-

elevation peak or a decreasing trend with elevation, with temperature having minimal role in

their species richness patterns (Beck et al., 2017). It is noteworthy that other evolutionary

hypotheses of species diversity patterns, like phylogenetic history (McCain, 2009), climatic

variation (Colwell et al., 2008) or biotic interactions (Novotny et al., 2006) could be system -

specific.

In this study, species from the Ennominae family from the genera *Alcis, Arichanna, Abraxas,*

Petelia were quite abundant in the low and mid elevations, whereas *Dalima truncataria*,

Blepharoctenucha virescens were few. *Psyra crypta* was more abundant towards higher

elevations. There were singletons from the Geometrinae family like *Jodis, Tanaorhinus cf.*

reciprocata. The turnover of species from the low to high elevations seemed to be influenced

by higher abundance of species belonging to the Larentiinae subfamily, specifically from the

genera *Perizoma*, *Photoscotosia* known to specialize on herbaceous plants (Skou, 1986). These

findings are similar to the patterns of Geometrid moths in neotropical and palearctic mountain

gradients (Brehm & Fiedler, 2003; Brehm et al., 2013). Thus the results point towards effect

of microclimate variables (Hof et al., 2011) on the species diversity and turnover. Most likely,

host-plant availability is a strong factor in determining species distribution.

It is already known, that the flight morphology of Geometrid moths makes them 'poor' fliers,

such that their occurrence is tightly linked to their host-plant availability (Heidrich et al., 2021).

Further evidence of host-plant relationships and their long-term effect on species distributions,

could support this claim. However, the host-plant relationships for moths remain insufficiently

understood in from this part of the globe.

Morphological trait variation across elevation

Elevation is adiabatically linked to temperature, and diverse moth families respond differently

to the changing temperature due to their distinct thermal sensitivities (Bladon et al., 2020;

Heidrich et al., 2021). Thermoregulation, a critical process controlled by the flight muscles in

insects (Heinrich, 1974), is influenced by the flight characteristics unique to each taxon.

Geometrid moths, characterized by larger wings relative to their body size, exhibit lower wing

loading compared to other moth families, capable of flight over a wide range of temperatures

(Casey & Joos, 1983). This trait allows them to engage in immediate flight without the need

for warming up-characteristically referred to as 'cool moth' in Heinrich (2013). Consequently,

it is not surprising that Geometrid moths demonstrate a lack of significant elevation-induced

effects on flight-related traits.

Moreover, the overall morphological trait space (niche dimension) of Geometrid moths along

the elevational gradient demonstrates little variation, likely due to the prevalence of species

with similar traits at each elevation. However, it is crucial to acknowledge that the intraspecific

variation of other physiological trade-offs, such as stress tolerance, resource use, and life

history strategies along the biotic and abiotic axes, may unveil actual niche segregation among

species (Mouillot et al., 2005; Wang et al., 2022). This interpretation exceeds the scope of this

paper and requires further investigation. The consistent overall assemblage-level trait overlap

in this study, further suggests that, despite decreasing species diversity, trait similarity and

functional redundancy among species persist, contributing to a stable trait distribution pattern.

Species having similar 'function' not only emphasize enhanced temporal stability of

communities (Pillar et al., 2013), but also provide a buffer against rapid species extinction on

the community's 'function' (Naeem, 1998). They are less likely to lose an entire functional

group due to stochastic processes only (Biggs et al., 2020). Also, following community

assembly rules, species of similar trait inhabit the same habitat, allowing them to 'function'

under specific abiotic conditions (Díaz et al., 1998; Weiher et al., 1998). Notably, the only

other research on trait-level patterns in moths across elevational gradients in the Himalaya,

found significant trait variations and internal filters influencing community assembly of

Hawkmoths (Mungee & Athreya, 2021). However, this study did not specifically focus on

examining the mechanisms of community assembly concerning changing elevation and related

abiotic factors.

Limitations and future possibilities

Heidrich et al. 2021 demonstrated a notable elevation-related increase in body size with

elevation by incorporating sites from lower elevations into their analysis. Extending this

approach to include additional sites from the species-rich lower elevations (<1500m) could

substantially impact the observed patterns in our study. However, locating continuous gradients

in deeper valleys from the foothills to over 3000m in this part of the Himalaya poses a

significant challenge.

The sampling in this study was restricted to the spring-summer season due to logistic

constraints. Reports from other western Himalayan elevational gradients, (Sanyal et al., 2017)

and Dey (2019) indicate variations in species composition across seasons. Consequently, it

would be worthwhile to investigate how the observed patterns hold across seasons.

Additionally, examining how life history traits related to larval host plant use (i.e., diet breadth

and host-plant growth form) and seasonal life cycle (i.e., overwintering stage and caterpillar

phenology) are linked to adult body size and flight morphology in species-rich herbivore insect

clades, Geometrids and alike, is crucial.

For species-rich taxa, trait- based conclusions are useful to overcome the taxonomic

impediment (Wong et al., 2019). In the current context of increasing demand for monitoring

insect populations, understanding traits that render species vulnerable to environmental

alterations caused by human activities provides the foundation for predicting community-level

response. Future studies should adopt a long-term perspective and a comprehensively examine

potential shifts in the observed consistent trait-space. Such investigations may reveal range

shifts in species due to changing thermal regimes, particularly in the fragile Himalayan

ecosystem.

Conclusion

The findings of the study indicate a decline in species diversity and an increase in species

turnover along elevational gradients. Notably, the elevation does not exert a significant

influence on the body size and flight capability of Geometrid moths. Therefore, the results

challenge prevailing notions about association between elevation and insect morphology.

Additionally, the study supports the idea that species with similar ecological functions are

sustained through the high turnover of species sharing similar trait values across assemblages.

In the case of Geometrid moths, the study underscores the insignificance of trait-environment

relationships, as evidenced by previous research (Beck et al., 2016; Heidrich et al., 2018; Xing

et al., 2018). Here, a clear need is identified to transition from a predominantly univariate

approach, focused on single-trait responses along individual environmental gradients, to a

multivariate approach that realistically considers the responses at species level across various

environmental gradients typical of most human-dominated landscapes.

The study's implications for the vulnerable Himalayan Mountain system, particularly in the

face of accelerated warming, emphasize the critical importance of comprehending species and

trait distribution patterns for the formulation of effective conservation strategies.

Acknowledgements: This work would not have been possible without funding from the
Rufford Foundation, UK. It was an independent endeavour, and I am immensely thankful to
the Dean, Wildlife Institute of India, Dr. V.P. Uniyal and the Uttarakhand Forest Department
(Letter no.2921/5-6(WII) dated 17 April 2018) for permitting me to carry out this work on my
own. I would like to thank Prabhat Singh Bisht, my local field assistant, and all the volunteers
who helped me during the fieldwork, namely: Meghavi Purohit, Kunjan Joshi, Bhargav
Bhadresvara, and Mohit. I extend my sincere thanks to Anza Simon and Rama Narayanan for
helping with the morphometry of the specimens, which were pinned and curated by Tarun
Karmakar. Lastly and most importantly, I am immensely thankful to Saskya Van Nouhuys,
Mayank Kohli, and Rohit Chakravarty for their valuable time, support, and discussion
regarding the analysis and writing of the manuscript. The manuscript was written and the data
was analyzed while I was funded by Dr. D.S. Kothari postdoctoral fellowship (2020-2023),
though this fellowship was not used in conducting the research for this paper.

Data availability statement: The data that support the findings of this study are available from
the corresponding author upon reasonable request. Upon the acceptance of the manuscript, the
data will be made accessible in a designated data depository.

References:

Ackerly DD, Cornwell WK. A trait-based approach to community assembly: partitioning of
species trait values into within- and among-community components. Ecology Letters
2007;10: 135–145.

Anderson MJ, Ellingsen KE, McArdle BH. Multivariate dispersion as a measure of beta
diversity. Ecology Letters 2006; 9: 683–693.

Arrizabalaga-Escudero A, Merckx T, et al., Trait-based functional dietary analysis provides a better insight into the foraging ecology of bats. Journal of Animal Ecology 2019; 88: 1587–1600.

Bakken GS. A heat transfer analysis of animals: Unifying concepts and the application of metabolism chamber data to field ecology. Journal of Theoretical Biology 1976; 60: 337–384.

Bartholomew GA, Casey TM. Body temperature and oxygen consumption during rest and activity in relation to body size in some tropical beetles. Journal of Thermal Biology 1977; 2: 173–176.

Beck J, Liedtke HC, Widler S, et al., Patterns or mechanisms? Bergmann's and Rapoport's rule in moths along an elevational gradient. Community Ecology 2016; 17: 137–148.

Beck J, McCain CM, Axmacher JC, et al., Elevational species richness gradients in a hyperdiverse insect taxon: a global meta-study on geometrid moths. Global Ecology and Biogeography 2017; 26: 412–424.

Biggs CR, Yeager LA, Bolser DG, et al., Does functional redundancy affect ecological stability and resilience? A review and meta-analysis. Ecosphere 2020; 11: e03184.

Bladon AJ, Lewis M, Bladon EK, et al., How butterflies keep their cool: Physical and ecological traits influence thermoregulatory ability and population trends. Journal of Animal Ecology 2020; 89: 2440–2450.

Blonder B, Lamanna C, Violle C, Enquist BJ. The n-dimensional hypervolume. Global Ecology and Biogeography 2014; 23: 595–609.

Blonder B, Morrow CB, Maitner B, et al., New approaches for delineating n-dimensional

hypervolumes. Methods in Ecology and Evolution 2018; 9: 305–319.

Brehm G. A new LED lamp for the collection of nocturnal Lepidoptera and a spectral

comparison of light-trapping lamps. Nota Lepidopterologica 2017; 40: 87–108.

Brehm G, Fiedler K. Faunal composition of geometrid moths changes with altitude in an

Andean montane rain forest. Journal of Biogeography 2003; 30: 431–440.

Brehm G, Fiedler K. Bergmann's rule does not apply to geometrid moths along an elevational

gradient in an Andean montane rain forest. Global Ecology and Biogeography 2004; 13: 7–

14.

Brehm G, Strutzenberger P, Fiedler K. Phylogenetic diversity of geometrid moths decreases

with elevation in the tropical Andes. Ecography 2013; 36: 1247–1253.

Brehm G, Zeuss D, Colwell RK. Moth body size increases with elevation along a complete

tropical elevational gradient for two hyperdiverse clades. Ecography 2018; 42(4): 632-642.

Casey TM, Joos BA. Morphometrics, Conductance, Thoracic Temperature, and Flight

Energetics of Noctuid and Geometrid Moths. Physiological Zoology 1983; 56: 160–173.

Chakravarty R, Radchuk V, Managave S, Voigt CC. Increasing species richness along

elevational gradients is associated with niche packing in bat assemblages. Journal of Animal

Ecology 2023; 92: 863–874.

Chown SL, Gaston KJ. Body size variation in insects: a macroecological perspective.

Biological Reviews 2010; 85: 139–169.

Classen A, Steffan-Dewenter I, Kindeketa WJ, et al., Integrating intraspecific variation in

community ecology unifies theories on body size shifts along climatic gradients. Functional

Ecology 2017; 31: 768–777.

Colwell RK, Brehm G, Cardelús CL, et al., Global warming, elevational range shifts, and

lowland biotic attrition in the wet tropics. science 2008; 322: 258–261.

Díaz S, Cabido M, Casanoves F. Plant Functional Traits and Environmental Filters at a

Regional Scale. Journal of Vegetation Science 2008; 9: 113–122.

Dillon ME, Frazier MR, Dudley R. Into thin air: Physiology and evolution of alpine insects.

Integrative and Comparative Biology 2006; 46: 49–61.

Dudley R. Mechanisms and Implications of Animal Flight Maneuverability. Integrative and

Comparative Biology 2002; 42: 135–140.

Gaston KJ, Chown SL, Calosi P, et al., Macrophysiology: A Conceptual Reunification. The

American Naturalist 2009; 174: 595–612.

Geange SW, Pledger S, Burns KC, Shima JS. A unified analysis of niche overlap

incorporating data of different types. Methods in Ecology and Evolution 2011; 2: 175–184.

Gérard M, Vanderplanck M, Franzen M, et al., Patterns of size variation in bees at a

continental scale: does Bergmann's rule apply? Oikos 2018; 127: 1095–1103.

Grula CC, Rinehart JP, Greenlee KJ, Bowsher JH. Body size allometry impacts flight-related

morphology and metabolic rates in the solitary bee Megachile rotundata. Journal of Insect

Physiology 2021; 133: 104275.

Heidrich L, Friess N, Fiedler K, et al., The dark side of Lepidoptera: Colour lightness of

geometrid moths decreases with increasing latitude. Global Ecology and Biogeography 2018;

27: 407–416.

Heidrich L, Pinkert S, Brandl R, et al., Noctuid and geometrid moth assemblages show

divergent elevational gradients in body size and color lightness. Ecography 2021; 44: 1169-

1179.

Heinrich B. Thermoregulation in Endothermic Insects: Body temperature is closely attuned to

activity and energy supplies. Science 1974; 185: 747–756.

Heinrich B. The Hot-Blooded Insects: Strategies and Mechanisms of Thermoregulation.

Springer Science & Business Media 2013.

Hof C, Levinsky I, Araújo MB, Rahbek C. Rethinking species' ability to cope with rapid

climate change. Global Change Biology 2011; 17: 2987–2990.

Holm S, Javoiš J, Kaasik A, et al., Size-related life-history traits in geometrid moths: a

comparison of a temperate and a tropical community. Ecological Entomology 2019a; 44:

711–716.

Holm S, Javoiš J, Molleman F, et al., No Indication of High Host-Plant Specificity in

Afrotropical Geometrid Moths. Journal of Insect Science 2019b; 19: 1.

Hsieh TC, Ma KH, Chao A. iNEXT: an R package for rarefaction and extrapolation of

species diversity (Hill numbers). Methods in Ecology and Evolution 2016; 7: 1451–1456.

Kearney M, Simpson SJ, Raubenheimer D, Helmuth B. Modelling the ecological niche from

functional traits. Philosophical Transactions of the Royal Society B: Biological Sciences

2010; 365: 3469–3483.

Lamanna C, Blonder B, Violle C, et al., Functional trait space and the latitudinal diversity

gradient. Proceedings of the National Academy of Sciences 2014; 111: 13745–13750.

Macarthur RH. Patterns of Species Diversity. Biological Reviews 1965; 40: 510–533.

Macarthur R, Levins R. The Limiting Similarity, Convergence, and Divergence of Coexisting

Species. The American Naturalist 1967; 101: 377–385.

McCain CM. Global analysis of bird elevational diversity. Global Ecology and Biogeography

2009; 18: 346–360.

Mouillot D, Stubbs W, Faure M, et al., Niche overlap estimates based on quantitative

functional traits: a new family of non-parametric indices. Oecologia 2005; 145: 345–353.

Mungee M, Athreya R. Intraspecific trait variability and community assembly in hawkmoths

(Lepidoptera: Sphingidae) across an elevational gradient in the eastern Himalayas, India.

Ecology and Evolution 2021; 11: 2471–2487.

Muscarella R, Uriarte M. Do community-weighted mean functional traits reflect optimal

strategies? Proceedings of the Royal Society B: Biological Sciences 2016; 283: 20152434.

Naeem S. Species Redundancy and Ecosystem Reliability. Conservation Biology 1998; 12:

39–45.

Novotny V, Drozd P, Miller SE, et al., Why are there so many species of herbivorous insects

in tropical rainforests? Science (New York, N.Y.) 2006; 313: 1115–1118.

Oksanen J, Kindt R, Legendre P, et al., 2009. The vegan Package.

Peters MK, Peisker J, Steffan-Dewenter I, Hoiss B. Morphological traits are linked to the

cold performance and distribution of bees along elevational gradients. Journal of

Biogeography 2016; 43: 2040–2049.

Pillar VD, Blanco CC, Müller SC, et al., Functional redundancy and stability in plant

communities. Journal of Vegetation Science 2013; 24: 963–974.

Porter WP, Gates DM. Thermodynamic Equilibria of Animals with Environment. Ecological

Monographs 1969; 39: 227–244.

Read QD, Grady JM, Zarnetske PL, et al., Among-species overlap in rodent body size

distributions predicts species richness along a temperature gradient. Ecography 2018; 41:

1718–1727.

Ricklefs RE, O'Rourke K. Aspect Diversity in Moths: A Temperate-Tropical Comparison.

Evolution 1975; 29: 313–324.

Samejima Y, Tsubaki Y. Body Temperature and Body Size Affect Flight Performance in a

Damselfly. Behavioral Ecology and Sociobiology 2010; 64: 685–692.

Sanyal AK, Dey P, Uniyal VP, et al., Geometridae Stephens, 1829 from different altitudes in

Western Himalayan Protected Areas of Uttarakhand, India.(Lepidoptera: Geometridae).

SHILAP Revista de Lepidopterología 2017; 45: 143-163.

Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis.

Nature Methods 2012; 9: 671–675.

Schumm M, White AE, Supriya K, Price TD. Ecological Limits as the Driver of Bird Species

Richness Patterns along the East Himalayan Elevational Gradient. The American Naturalist

2020; 195: 802–817.

Seifert CL, Strutzenberger P, Hausmann A, Fiedler K. Dietary specialization mirrors

Rapoport's rule in European geometrid moths. Global Ecology and Biogeography 2022; 31:

1161–1171.

Shelomi M. Where Are We Now? Bergmann's Rule Sensu Lato in Insects. The American

Naturalist 2012; 180: 511–519.

Skou P. The Geometroid Moths of North Europe (Lepidoptera: Drepanidae and

Geometridae). Brill Archive, 1986.

Stevenson RD. Body Size and Limits to the Daily Range of Body Temperature in Terrestrial

Ectotherms. The American Naturalist 1985; 125: 102–117.

Svensson EI, Gómez-Llano M, Waller JT. Out of the tropics: Macroevolutionary size trends

in an old insect order are shaped by temperature and predators. Journal of Biogeography

2023; 50: 489–502.

Wang XZ, Sun SW, Sedio BE, et al., Niche differentiation along multiple functional-trait

dimensions contributes to high local diversity of Euphorbiaceae in a tropical tree assemblage.

Journal of Ecology 2022; 110: 2731–2744.

Weiher E, Clarke GDP, Keddy PA. Community Assembly Rules, Morphological Dispersion,

and the Coexistence of Plant Species. Oikos 1998; 81: 309–322.

Wong MKL, Guénard B, Lewis OT. Trait-based ecology of terrestrial arthropods. Biological

Reviews 2019; 94: 999–1022.

Xing S, Bonebrake TC, Ashton LA, et al., Colors of night: climate–morphology relationships

of geometrid moths along spatial gradients in southwestern China. Oecologia 2018; 188:

537–546.

Map 1: Map showing the sampling locations in and around the buffer zone of Kedarnath Wildlife Sanctuary, which is located in northern side of the mountainous state of Uttarakhand (highlighted on the map of India on the left side)

Figure 1: Rarefied curves for a) Sample coverage and b) Species diversity patterns extrapolated across the number of individuals; c) overall beta diversity measured as Sorensen

pairwise dissimilarity and d) the spatial turnover of the species measured as Simpson pairwise dissimilarity across the low, mid and high elevation assemblages

Figure 2: The figure shows the dispersion of traits as a measure of intraspecific trait variability in each of the assemblages along the elevation and all the elevations together (last panel). The beta-dispersions refer to a species' position along a gradient or intraspecific trait variation defined by community-level mean trait values (shown in red)

Figure 3: The figure shows the community weighted means (mean ± SE) of each of the traits across the low, mid and high-elevation assemblages.

Figure 4: The figure shows the degree of overlap (O-statistic) of the multivariate trait-space, based on body length (BL), maneuverability (MV), and wing loading (WL) of the Geometrid moth assemblages in low, mid, and high elevations