Title: Trait distribution in Geometrid moth assemblages does not vary along a western Himalayan elevational gradient

Running Title: Trait Patterns of Himalayan Geometrids

Author: Pritha Dey

Author details:

Affiliation during the grant period: Wildlife Institute of India, Chandrabani, Dehradun-248001

Present address: National Centre for Biological Sciences, Tata Institute of Fundamental Research, Bellary Road, Bangalore-560065

Abstract:

Elevation gradients pose challenges for flying insects due to temperature and air density shifts, impacting their thermoregulation, flight abilities and subsequently species distribution patterns. Geometrid moths are a model group for studying these traits among insects. This study examines how trait-patterns and niche differentiation of Geometrid moth assemblages change with elevation. Investigating 697 specimens of 120 Geometrid moth species, I assessed species diversity, turnover, and various traits related to body size and flight capabilities along a 1500m elevation span in the western Himalayas. Results indicate a decline in species diversity along the elevational gradients, accompanied by consistent turnover from lower to higher elevations. Despite this pattern, elevation did not significantly influence morphological traits related to thermal sensitivity (body size) and flight capability (wing loading and manoeuvrability) at the assemblage level in Geometrid moths. Furthermore, there was no significant niche differentiation among species across elevations. This supports the argument that species with similar ecological roles persist through turnover, maintaining consistency in trait values across assemblages. These findings from the paleotropics align with patterns observed in Geometrid moths across neotropical and temperate regions, highlighting the importance of trait-based approaches in understanding species responses to environmental changes.

Summary Statement: This study examines how Geometrid moth assemblages in the western Himalayas respond to elevational gradients, revealing consistent turnover, stable traits, and no significant niche differentiation, emphasizing trait-based species adaptation

Introduction:

Mountain ecosystems are increasingly threatened by climate change, posing a risk to cold-adapted and endemic species that depend on these regions (Laiolo *et al.,* 2018; Trew &

Maclean, 2021). These unique ecosystems have drastic environmental changes over small

spatial scales (Rogora *et al.,* 2018) affecting species distributions across gradients. To survive

these conditions, species need to adapt physiologically, which is often reflected their

morphological traits (Ryding *et al.*, 2021; Warrington & Waterman, 2023). Morphological

traits are, therefore, crucial for our understanding of species assembly (Hoiss *et al.*, 2012;

Chichorro *et al.*, 2022). The importance of morphological traits is enhanced while studying

assemblages of species rich clades such as insects, where species identification is challenging

(Yates *et al.,* 2014)

Insects adapt to abiotic factors such as elevation, temperature, and air density indirectly

through interactions with hostplants, competitor species, parasitoids and predators

(Hodkinson, 2005), or directly through traits like body size, wing size and colouration (Xing

et al., 2018; Bladon *et al.*, 2020; Clusella-Trullas & Nielsen, 2020). Body size (often

evaluated as wingspan), in particular influences thermoregulation, and is a key focus in

studies on patterns of traits in insects across elevation and latitudes (reviewed in Chown &

Gaston, 2010). For example, bees decrease in size with elevation (Osorio-Canadas *et al*.,

2022), but increase across latitudes (Gérard *et al.,* 2018). Dragonflies and damselflies are

larger in temperate regions than in the tropics (Svensson *et al.*, 2023). Noctuid moths show

larger body sizes at higher elevations in temperate regions (Heidrich *et al.*, 2021). However,

there is significant variability among taxa. For instance, Geometrid moths, show diverse

patterns: some studies indicate a positive relationship between body size and elevation (Beck

et al., 2016; Brehm *et al.,* 2018), while others reveal no change (Brehm & Fiedler, 2004).

Additionally, body size directly impacts flight-related traits (Grula *et al.,* 2021), with larger

insects having higher wing loading and lower manoevrability (Shyy *et al.,* 2016). At higher

elevations, lower temperatures favour species with smaller body and reduced wing loading

(therefore proportionally long wings) (Dudley, 2002) which is advantageous in low air

density (Dillon & Dudley, 2004; Dillon *et al.*, 2006; Grula *et al.*, 2021).

Morphological traits are critical in defining a species' niche in an assemblage (Violle &

Jiang, 2009) and inferences can be made regarding community assembly by measuring

changes in the niche space and trait overlap across ecological gradients. At lower elevations,

warm temperature and abundant resources reduce constraints, leading to high diversity and

niche specialization with lower trait overlap (Read *et al.*, 2018), as seen in Orthopteran

(König *et al.,* 2024) and bee communities (Hoiss *et al.,* 2012). Conversely, higher elevations

impose stricter environmental constraints, resulting in lower diversity and higher trait

overlap, consistent with the 'niche packing' model (Macarthur,1965; Pigot *et al.,* 2016;

Pellissier *et al.,* 2018). The degree of trait overlap is shaped by niche differentiation and the

displacement of functional trait values over evolutionary time (Macarthur & Levins, 1967;

Ricklefs & O'Rourke, 1975). However, distinguishing traits affected by biotic versus abiotic

environmental filters, in trait-based approaches, can be challenging (Violle & Jiang, 2009).

Analyzing the traits weighed by the relative abundance of the species best reflect the

adaptation of traits to the environment (Garnier *et al.,* 2004; Shipley *et al.,* 2006).

Community Assembly Theory (HilleRisLambers *et al.,* 2012; Kraft *et al.,* 2015) predicts that

assemblages that are shaped by interspecific competition show overdispersed trait spaces

whereas those shaped by environmental adaptations are underdispersed, given the relatedness

of the species in the assemblage (Weiher & Keddy, 1995)

In this context, I Firstly wanted to understand 1) how does species diversity and turnover

change across the elevation? If there is a change in species diversity across elevation, 2) do

body size and flight related traits show a predictable relationship with elevation?, and 3) do

trait dispersion and the overall trait overlap (niche differentiation) at assemblage level vary

with changing species diversity along the elevation? I predict species diversity to decrease

with high turnover with increasing elevation. I hypothesize that body size and flight traits are expected to change with elevation to achieve optimal thermoregulation and flight, leading to reduced body size, reduced wing loading and higher manoevrability. Following species assembly rules, higher elevations are expected to show trait under dispersion, higher trait overlap, with less niche differentiation due to decreased species diversity. While these patterns have been observed in bird and bat communities (Schumm *et al.,* 2020; Chakravarty *et al.,* 2023), in the tropical Himalayan mountain system, it remains unexplored if similar patterns apply to insects as well.

I studied Geometrid moth assemblages across a 1500m elevation gradient on the fringes of Kedarnath Wildlife Sanctuary. Despite the extensive use of Geometrid moths in ecological studies (Brehm & Fiedler, 2004; Davis *et al.,* 2013; Beck *et al.,* 2016; Holm *et al.,* 2019a, 2019b; Choi *et al.,* 2022; Seifert *et al.*, 2022; Foerster *et al.,* 2024) so far no study has considered body size and flight-related traits together to understand the effect of elevation on the overall trait space and the niche differentiation. This trait-based approach aims to fill this gap, enhancing our ability to address ecological questions, particularly given the unclear phylogenetic positions of Geometrid moths from this region in global studies.

Furthermore, research on macroecological patterns within paleotropical montane systems is limited. While previous studies from the western Himalayan region, Sanyal et al. (2017) and Dey (2019) have investigated species diversity patterns across elevational gradients, they did not consider trait-based patterns. Trait-based research on moths from this region focused on less diverse groups like Hawkmoths, highlighting significant trait variations and internal filters influencing community assembly (Mungee & Athreya 2021). This study specifically focused on, on understanding how elevation and related abiotic factors influence trait-level patterns among highly diverse Geometrid moths.

Results:

Species diversity

Out of 119, there were 56 species level identifications, 56 morphospecies identified only till genus, one was identified till subfamily, three morphospecies were identified till tribe and three morphospecies remain unidentified (Table S3). Figure 1b illustrates a decrease in species diversity at higher elevations. Rarefaction curves facilitate the comparison of diversity across various communities or habitats. By scrutinizing the shapes and positions of these curves (Figure 1a and 1b), the disparities in species diversity, sampling completeness, and overall species diversity among sites are clear. A steep slope indicates rapid accumulation of new species with additional sampling effort, suggesting high species richness and/or incomplete sampling. In contrast, a shallow slope indicates that most species have already been sampled, and further sampling is unlikely to yield many new species.

For beta diversity, the resulting box-plot shows the distribution of βnested and βturnover values across the different pairs of sites or groups in your data (Fig 1c and Fig 1d). This visualization helps to understand the relative contributions of nestedness and turnover to the overall beta diversity pattern. The Euclidean distance to centroid for Sørensen pairwise dissimilarity among the elevation categories were as follows: Low-Mid (0.314); Low-High (0.489) and Mid-High (0.34); similarly, the Simpson dissimilarity index (spatial turnover) was: Low-Mid (0.05); Low-High (0.14) and Mid-High (0.12) (Fig 1 c and d). Though there is indication of dissimilarity in species composition and turnover of species between the low and high elevation sites, the Euclidean distances are not significantly different from each other (Sørensen pairwise dissimilarity: ANOVA, df=2, p>0.05; Spatial turnover: ANOVA, df=2, p>0.05)

Variation in the individual trait values, trait dispersion and trait space overlap (niche

differentiation)

There was no significant effect of elevation on the community-weighted means of the

individual trait values (Fig.2) (model summary provided in Table S2). The pairwise

comparisons of the raw means of the individual traits across the elevation classes also did not

show any significant difference (details in Figure S3). There is high variability and a weak

pattern in the CWM across elevation, which makes it evident that the patterns are sensitive to

the abundance of species. The species of same trait values are abundant across the elevation.

The density plots of individual traits would make it clearer, where kernel density estimate is

used to show the distribution of each of the traits, indicating the mean values (Figure S4).

Trait dispersion did not differ significantly among the subfamilies in the low (df=3, p=0.149),

mid (df=3, p=0.252), and in high elevation (df=3, p=0.2). Similarly, no significant

differences were found in trait dispersion among assemblages across the elevation

(df=2,p=0.3) (Fig.3).

There is also a high level of niche overlap among the species of the Geometrid moth

communities across the elevation, as shown by high (0.84) O-statistic index of the

multivariate trait space (Fig. 4). This indicates that for each trait the observed niche overlap

did not differ from that which would be expected due to chance.

Discussion:

The findings offer important insights into the dynamics of Geometrid moth assemblages

along a 1500m gradient in the western Himalaya. The results show a decrease in species

diversity with increasing elevation, along with consistent species turnover from the low to the

high elevations. Despite this, elevation does not significantly impact morphological traits

related to thermal sensitivity (body size) and flight capability (wing loading and

manoevrability) at the assemblage level. This suggests that Geometrid moths maintain trait

consistency across different elevations. Additionally, the study reveals no significant niche

differentiation among species across the elevation.

Species diversity across elevation

The results indicate that trait-level patterns may not directly correlate with the species

diversity, and relative abundance. For instance, species from subfamily Ennominae, such as

from the genera *Alcis* and *Abraxas*, were abundant at low and mid elevations, likely due to its

polyphagous diet (Choi *et al.,* 2022), while species like *Dalima truncataria* were scarce.

Conversely, *Psyra crypta* was more prevalent at higher elevations. The turnover from low to

high elevations is influenced by the higher abundance of species from Larentiinae subfamily,

particularly those specializing in herbaceous plants, reflecting similar patterns observed in

other mountain gradients (Skou, 1986; Brehm & Fiedler, 2003; Brehm *et al.,* 2013). Further,

local host-plant availability, diet breadth, and breeding strategies likely affects species

diversity (Heidrich *et al.*, 2021; Seifert *et al.*, 2022). Moreover, the consistent presence and

high abundance of some species across elevations might also indicate density compensation

mechanisms due to competition and predator relaxation (Longino & Colwell 2011; Choi *et al.*

2022)

Morphological trait variation across elevation

Body size significantly affects how insects respond to temperature changes, influencing their

distribution and thermal tolerance (Hodkinson *et al.,* 1999; Hodkinson, 2005). However,

Geometrid moths do not show significant body size changes with elevation (Bladon *et al.,*

2020; Heidrich *et al.,* 2021), likely due to their large wings and low wing loading, which

allow them to fly across varying temperatures without extensive warm-up (Casey & Joos

1983; Heinrich, 2013). This wide thermal tolerance may contribute to their stable body size

and flight traits across elevations. Similar adaptation patterns have been observed in other

taxa, such as the African dung beetles (Gaston & Chown, 1999). The high abundance of

certain common species also played a role in maintaining stability, as these species tend to

have traits well-suited to the prevailing environmental conditions (Cingolani *et al.,* 2007)

The uniformity in trait dispersion among Geometrid subfamilies suggests that similar

selective pressures or trait convergence are at play, despite the taxonomic diversity. Although

the impact of phylogenetic constraints remains unclear due to limited taxonomic and

phylogenetic knowledge of moths from this region, the minimal variation in morphological

traits along elevation is likely attributed to the abundance of species with similar traits and

evolutionary relatedness (Griffith *et al,*2023.). Traits that are conserved through shared

evolutionary history are apparent in many phytophagous insects. For example, closely related

herbivore species often feed on the same or similar plant taxa (Nylin *et al.,* 2014;

Strutzenberger *et al.*, 2017) and exhibit comparable levels of dietary specialization (Slove &

Janz, 2011; Hardy *et al.,* 2016) and body size (Ribeiro & Freitas, 2011). Such similarity in

traits across communities, indicate a lower degree of specialization and a higher prevalence

of generalist species. As a result, this increased trait similarity may reduce ecosystem

resilience, affect community functioning, and impair ecosystem services (Lavorel & Garnier,

2002; Olden, 2006)

More definitive conclusions about the mechanisms driving the lack of variation in body size

and flight traits in insects, and how these influence species distributions, are hindered by

insufficient life-history, ecological, and abundance data collected over extended periods and

with higher temporal resolution, from the Himalayan region.

Conclusion

The study provides valuable insights into how individual traits, trait variability within the assemblages, and niche differentiation respond to changing species diversity across elevational gradients. With evidence from the Paleotropical region, the results align with the lack of consistent trends in individual traits studied in Geometrid moths and other insect taxa across diverse elevation ranges in neotropics and palearctic regions (Brehm & Fiedler, 2004; Peters *et al.*, 2016; Beck *et al.,* 2016; Classen *et al.,* 2017).

No significant variation in trait-patterns with changing species diversity, challenge prevailing notions about the association between elevation and insect morphology. It highlights the possible role of functional redundancy (Pillar *et al.*, 2013) or biotic homogenization (Olden, 2006)in maintaining trait distributions across varied elevations and underscores the need for a nuanced species-specific understanding of trait-environment relationships.

However, the limitations, including restricted sampling to the spring-summer season and the exclusion of lower elevation sites, suggest a need for further research. Seasonal variations and additional lower elevation sites could provide a comprehensive understanding of trait-environment dynamics. Previous studies from other western Himalayan elevational gradients, (Sanyal *et al.*, 2017) and (Dey, 2019) indicate variations in species composition across seasons, highlighting the need for such an investigation. Further, integrating phylogenetic information is essential when examining species traits, their determinants, and their relationships.

Future research could explore how life history traits, such as larval host plant use (i.e., diet breadth and host-plant growth form) and seasonal life cycle (i.e., overwintering stage and caterpillar phenology); and factors like predator density are linked to adult body size and flight morphology. Such causal relationships would be crucial for developing effective conservation strategies, especially as climate change alters environmental conditions. Long-

term studies investigating potential shifts in trait distributions, can provide valuable insights

into how species respond to changing thermal regimes and help tailor conservation efforts to

protect the unique biodiversity of the Himalayan ecosystem.

Materials and methods:

Study Area:

The study was done in 2018 around the fringes of the Kedarnath Wildlife Sanctuary (KWLS)

(30°25′–30°41′N, 78° 55′–79°22′E) located in the Chamoli-Rudraprayag district in the state

of Uttarakhand (Map 1). KWLS covers an area of 975 sq. km and lies in the upper catchment

area of the rivers Alakananda and Mandakini, the major tributaries of the river Ganga. The

elevation of the protected area ranges from 1160-7068 meters above sea level, encompassing

diverse habitats from moist temperate forests, mixed oak forests to lush alpine meadows. The

combination of human pressure, pristine forest areas, and a large altitudinal range make it an

ideal site for biogeographic studies. The forest types are dominated by oak (*Quercus*

leucotrichophora, Quercus floribunda, Quercus glauca, and Quercus semecarpiflora). The

area receives annual rainfall of about 3000 mm with maximum contribution during monsoon

(June to September). During the winter months (December to February) heavy snowfall takes

place in the alpine and temperate regions (Thakur *et al.*, 2011).

Field sampling:

The study area was stratified based on elevation and vegetation types to sample the moth

diversity along the gradients. Sampling was done every 200m along the elevation from

1500m up to 3100m. For analyses, I categorized the sampling sites based on similar elevation

and vegetation types as follows: Khalla, Mandal, Gondi (below 2000m: Low Elevation);

Jatholi, Ansuya, Kanchula Kharak (2000-2500m: Mid Elevation); Bulkhan and Shokharak

(>2500m: High Elevation). The vegetation types ranged from the Himalayan Moist

Temperate Forests dominated by *Quercus* spp.(Low elevation) and mixed Oak Forests (Mid

elevation) and; sub-alpine forests with *Rhododendron* spp- Alpine meadows with (High

elevation) (Champion and Seth, 1968). The low, medium, and high elevations were therefore

used as factors in statistical analyses. Moreover, spacing elevation bands out in this manner

also reduces the spatial autocorrelation across our sampling sites.

At each site, 3-night sampling was done (details provided in Table S1 and the sampling sites

are shown in Map 1). On each sampling date, two LepiLED (Brehm, 2017) light traps were

operated for 3-4 hours after dusk, depending on the weather conditions (Estimated trap hours-

Trap 1: 1900-2200; Trap 2: 2000-2300 hrs) which is the peak time of activity of Geometrid

moths (Brehm & Fiedler, 2003; and also from personal observations during reconnaissance

survey). Also, logistic constraints do not permit all-night sampling. Sampling was done for a

total ~140 hours from March to May (Spring-Summer season) in 2018, during which the

average temperatures during sampling were, 16.2 °C, 11°C and 7.5°C at the low, mid and

high elevation zones respectively.

The minimum distance between the two light traps was 50 m, with lamps not in line of sight

of neighbouring sites, so that cross-site sampling does not occur. Geometrid moths attracted

to traps were counted, and representative samples from each species were collected for

morphometric measurements and as vouchers. While many species were represented by only

one specimen (singletons), an average of three specimens were collected whenever possible

for more reliable morphological analysis. All species were included for morphometric

measurements, analyses, and diversity assessment. The specimens were collected only from

March to May in 2018, as per the permit from the Forest Department. The specimens were

then identified till species, as much as possible. The voucher specimens have been submitted

to the Research Collections Facility at the National Centre for Biological Sciences,

Bangalore, India.

Morphological trait measurement:

All voucher specimens were curated and photographed. I measured morphological traits on

the scaled photographs of 697 specimens for morphometry using the software Image J

(Schneider, Rasband & Eliceiri, 2012). As different linear measurements widely used proxies

for body size in Geometrid moths, (Brehm *et al.,* 2019; Foerster *et al., 2024*)). I measured the

body length, wingspan (the width between the apices of the forewings, with the trailing wing

edge perpendicular the body), forewing length (measured from the body to the apex of the

forewing) and hindwing length. Further, forewing area, hindwing area, were used to calculate

Wing loading [body length/ 2*(forewing area R+ hindwing area R)] with body length as a

proxy for body mass and total wing area is twice the area of right forewing and hindwing

combined (Shi *et al.*, 2015), and the ratio of hindwing area to forewing area which is roughly

termed manoeuvrability (Arrizabalaga☐Escudero *et al,* 2019), (Figure S2) which is directly

related to the flight capability of moths. These measured traits are thus predicted to govern

the occurrence and sustenance of geometrid moths across elevational gradients.

Analyses:

Species diversity

The elevation bands for the sites selected for sampling in my study area were separated by

roughly 200 m each. In order to perform meaningful statistics using elevation as a continuous

variable, the bare minimum would have been three sites at every 200 m. This was,

unfortunately, not possible given the topography of the landscape. Therefore, I pooled the

sampling sites into low (1500-1600m; Mandal, Khalla, Gondi), medium (1700-2000m), and

high (2600-3000 m). The low, medium, and high elevations were therefore used as factors. I

used the 'iNEXT' R package (Hsieh *et al.,* 2016) to calculate rarefied species diversity

extrapolated to the number of individuals across different elevation classes. This approach

allows to visualize species diversity as a function of sampling effort or the number of

individuals sampled. By rarefying the data, the species diversity estimates can be

standardized across elevation classes, enabling a comparison of species richness among sites

with different sampling intensities. To show sample completeness, I similarly calculated the

rarefied sample coverage at each of the elevation classes.

To understand the effect of elevation on the species composition in the assemblages, I

calculated the overall beta-diversity (Sørensen pairwise dissimilarity), and species spatial

turnover (Simpson's pairwise dissimilarity), all measured as average dissimilarity of the

observed units to their group centroid (Anderson *et al.*, 2006). This analysis was done with

the 'betapart' package in R (Baselga & Orme, 2012), which decomposes the total beta

diversity into two components, Nestedness (βnested): this component represents the variation

in community composition due to species loss or gain across sites that are nested within

others. It reflects the turnover of species due to differences in site richness; Turnover

(βturnover): this component represents the variation in community composition due to

species replacement between sites. It reflects the substitution of species among sites.

Variation in the individual trait values, trait dispersion and trait space overlap (niche

differentiation)

For understanding the effect of elevation on individual traits, I used two approaches. First, I

investigated the response using community abundance-weighted mean trait value (CWM;

Lavorel *et al.,* 2008), a widely used method to calculate community trait values weighted by

abundance of species in that community. The change in CWM with elevation was quantified

using ordinary least squares regression, where each trait was used as a predictor variable in a

separate model with Elevation (factor) as the response variable. Then the raw mean of the

individual trait values were compared by pairwise comparisons. It is useful to look at both

the approaches as the relationships between community-weighted mean (CWM) traits and

elevation, is supposed to reflect selection towards locally optimal traits (Violle & Jiang,

2009), but it is limited by the large amount of interspecific trait variation typically found

within ecological communities (Muscarella & Uriarte, 2016).

To understand the trait dispersion at the subfamily-level and at the species assemblage level

across the elevational gradient, I compared the homogeneity of dispersion (Anderson *et al.,*

2006) for the assemblages in each elevation category. I used the *betadisper* function of the R

package vegan (Oksanen *et al.* 2009) to calculate dispersion, which tests whether the variance

or dispersion of two or more groups is significantly different or not based on Euclidean

distance. This was followed by a *permutest* for the homogeneity of multivariate dispersion.

Additionally, I have used Principal Coordinates Analysis (PCoA), to visualize pairwise

distances in a two-dimensional plot followed by PERMANOVA.

To understand the overall trait overlap (niche differentiation) among species assemblages

across the elevational gradient, I used O-statistic (Read *et al.,* 2018) to generate a single

community-level (pooling the assemblages across the elevation) overlap index across

multiple traits by fitting non-parametric kernel density functions to the distribution of trait

values within an elevation (Mouillot *et al.,* 2005; Geange *et al.,* 2011; Read *et al.,* 2018).

For this, body length, wing loading, and Manoeuvrability were chosen as the least correlated

independent variable, to avoid multicollinearity. For this, The median pairwise overlap was

calculated by first determining the overlap of traits in each species pair, and then taking the

median overlap of each species pair in a community. This median overlap value is called the

O-statistic (O for overlap)

The R package *Ostats* (Read *et al.,* 2022) was used to calculate O-statistic, the multivariate O-statistic estimation in this package relies on functions imported from the package *hypervolume* (Blonder *et al.,* 2018). To test whether the degree of niche overlap among species in each elevation category is greater than or less than expected by chance, I evaluated the O-statistic against a null model, where the median of the pairwise hypervolume overlaps are weighted against the abundances of the species pairs in each community or Standardized Effect Size (z☐scores).

Acknowledgements: This work would not have been possible without funding from the Rufford Foundation, UK. It was an independent endeavour, and I am immensely thankful to the Dean, Wildlife Institute of India, Dr. V.P. Uniyal and the Uttarakhand Forest Department (Letter no.2921/5-6(WII) dated 17 April 2018) for permitting me to carry out this work on my own. I would like to thank Prabhat Singh Bisht, my local field assistant, and all the volunteers who helped me during the fieldwork, namely: Meghavi Purohit, Kunjan Joshi, Bhargav Bhadresvara, and Mohit. I extend my sincere thanks to Anza Simon and Rama Narayanan for helping with the morphometry of the specimens, which were pinned and curated by Tarun Karmakar. Lastly and most importantly, I am immensely thankful to Saskya Van Nouhuys, Mayank Kohli, and Rohit Chakravarty for their valuable time, support, and discussion regarding the analysis and writing of the manuscript. The manuscript was written and the data was analyzed while I was funded by Dr. D.S. Kothari postdoctoral fellowship (2020-2023), though this fellowship was not used in conducting the research for this paper.

Competing interests: There are no competing interests

Funding: The study was funded by the Rufford Foundation small grants programme, UK

Data availability statement: The data that support the findings of this study are available

from the corresponding author upon reasonable request. Upon the acceptance of the

manuscript, the data will be made accessible in a designated data depository.

References:

Anderson MJ, Ellingsen KE, et al., Multivariate dispersion as a measure of beta diversity, Ecology
Letters 2006;9:683–693.

Arrizabalaga-Escudero A, Merckx T, García-Baquero G et al., Trait-based functional dietary analysis
provides a better insight into the foraging ecology of bats, Journal of Animal Ecology 2019;88:1587–
1600.

Baselga A, Orme CDL. betapart: an R package for the study of beta diversity, Methods in Ecology and
Evolution 2012;3:808–812.

Beck J, Liedtke HC, Widler S et al., Patterns or mechanisms? Bergmann's and Rapoport's rule in
moths along an elevational gradient, Community Ecology 2016;17:137–148.

Bladon AJ, Lewis M, Bladon EK et al., How butterflies keep their cool: Physical and ecological traits
influence thermoregulatory ability and population trends, Journal of Animal Ecology 2020;89:2440–
2450.

Blonder B, Morrow CB, Maitner B et al., New approaches for delineating n-dimensional
hypervolumes, Methods in Ecology and Evolution 2018;9:305–319.

Brehm G. A new LED lamp for the collection of nocturnal Lepidoptera and a spectral comparison of
light-trapping lamps, Nota Lepidopterologica 2017;40:87–108.

Brehm G, Fiedler K. Faunal composition of geometrid moths changes with altitude in an Andean
montane rain forest, Journal of Biogeography 2003;30:431–440.

Brehm G, Fiedler K. Bergmann's rule does not apply to geometrid moths along an elevational
gradient in an Andean montane rain forest, Global Ecology and Biogeography 2004;13:7–14.

Brehm G, Strutzenberger P, et al., Phylogenetic diversity of geometrid moths decreases with
elevation in the tropical Andes, Ecography 2013;36:1247–1253.

Brehm G, Zeuss D, et al., Moth body size increases with elevation along a complete tropical
elevational gradient for two hyperdiverse clades, Ecography 2019;42:632–642.

Casey TM, Joos BA. Morphometrics, Conductance, Thoracic Temperature, and Flight Energetics of
Noctuid and Geometrid Moths, Physiological Zoology 1983;56:160–173.

Chakravarty R, Radchuk V, Managave S et al., Increasing species richness along elevational gradients
is associated with niche packing in bat assemblages, Journal of Animal Ecology 2023;92:863–874.

Chichorro F, Urbano F, Teixeira D et al., Trait-based prediction of extinction risk across terrestrial
taxa, Biological Conservation 2022;274:109738.

Choi S-W, An J-S, Lee J-Y et al., Spatial and temporal changes in moth assemblages along an altitudinal gradient, Jeju-do island, Scientific Reports 2022;12:20534.

Chown SL, Gaston KJ. Body size variation in insects: a macroecological perspective, Biological Reviews 2010;85:139–169.

Cingolani AM, Cabido M, Gurvich DE et al., Filtering processes in the assembly of plant communities: Are species presence and abundance driven by the same traits? Journal of Vegetation Science 2007;18:911–920.

Classen A, Steffan-Dewenter I, Kindeketa WJ et al., Integrating intraspecific variation in community ecology unifies theories on body size shifts along climatic gradients, Functional Ecology 2017;31:768–777.

Clusella-Trullas S, Nielsen M. The evolution of insect body coloration under changing climates, Current Opinion in Insect Science 2020;41:25–32.

Davis RB, Õunap E, Javoiš J et al.,Degree of specialization is related to body size in herbivorous insects: a phylogenetic confirmation, Evolution 2013;67:583–589.

Dey P. Diversity assessment and molecular characterization of Geometridae moths (Lepidoptera: Heterocera) from Nanda Devi Biosphere Reserve, Uttarakhand, India. PhD thesis. Saurashtra University, 2019

Dillon ME, Dudley R. Allometry of maximum vertical force production during hovering flight of neotropical orchid bees (Apidae: Euglossini), Journal of Experimental Biology 2004;207:417–425.

Dillon ME, Frazier MR, et al.,. Into thin air: Physiology and evolution of alpine insects, Integrative and Comparative Biology 2006;46:49–61.

Dudley R. Mechanisms and Implications of Animal Flight Maneuverability, Integrative and Comparative Biology 2002;42:135–140.

Foerster SÍ, Javoiš J, Holm S et al., Predicting insect body masses based on linear measurements: a phylogenetic case study on geometrid moths, Biological Journal of the Linnean Society 2024;141:71–86.

Garnier E, Cortez J, Billès G et al. Plant functional markers capture ecosystem properties during secondary succession. Ecology 2004;85:2630–2637.

Gaston KJ, Chown SL. Elevation and Climatic Tolerance: A Test Using Dung Beetles. Oikos 1999;86:584–590.

Geange SW, Pledger S, Burns KC et al., A unified analysis of niche overlap incorporating data of different types, Methods in Ecology and Evolution 2011;2:175–184.

Gérard M, Vanderplanck M, Franzen M et al., Patterns of size variation in bees at a continental scale: does Bergmann's rule apply? Oikos 2018;127:1095–1103.

Griffith DM, Byrd KB, Anderegg LDL et al., Capturing patterns of evolutionary relatedness with reflectance spectra to model and monitor biodiversity, Proceedings of National Academy of Science US 2023, 120:e2215533120.

Grula CC, Rinehart JP, Greenlee KJ et al., Body size allometry impacts flight-related morphology and
metabolic rates in the solitary bee Megachile rotundata, Journal of Insect Physiology
2021;133:104275.

Hardy NB, Peterson DA, et al., Nonadaptive radiation: Pervasive diet specialization by drift in scale
insects? Evolution 2016;70:2421–2428.

Heidrich L, Pinkert S, Brandl R et al., Noctuid and geometrid moth assemblages show divergent
elevational gradients in body size and color lightness, Ecography 2021;44:1169–1179.

Heinrich B. The Hot-Blooded Insects: Strategies and Mechanisms of Thermoregulation, Springer
Science & Business Media, 2013.

HilleRisLambers J, Adler PB, Harpole WS et al., Rethinking Community Assembly through the Lens of
Coexistence Theory, Annual Review of Ecology, Evolution, and Systematics 2012;43:227–248.

Hodkinson ID. Terrestrial insects along elevation gradients: species and community responses to
altitude, Biological Reviews 2005;80:489–513.

Hodkinson ID, Bird J, Miles JE et al., Climatic signals in the life histories of insects: the distribution
and abundance of heather psyllids (Strophingia spp.) in the UK, Functional Ecology 1999;13:83–95.

Hoiss B, Krauss J, Potts SG et al. Altitude acts as an environmental filter on phylogenetic
composition, traits and diversity in bee communities. Proceedings of the Royal Society B: Biological
Sciences 2012;279:4447–4456.

Holm S, Javoiš J, Kaasik A et al., Size-related life-history traits in geometrid moths: a comparison of a
temperate and a tropical community, Ecological Entomology 2019a;44:711–716.

Holm S, Javoiš J, Molleman F et al., No Indication of High Host-Plant Specificity in Afrotropical
Geometrid Moths, Journal of Insect Science 2019b;1-16.

Hsieh TC, Ma K, et al., iNEXT: an R package for rarefaction and extrapolation of species diversity (Hill
numbers), Methods in ecology and evolution 2016; 7(12): 1451-1456

König S, Krauss J, Classen A et al., Micro- and macroclimate interactively shape diversity, niches and
traits of Orthoptera communities along elevational gradients, Diversity and Distributions
2024;30:e13810.

Kraft NJB, Adler PB, Godoy O et al., Community assembly, coexistence and the environmental
filtering metaphor, Functional Ecology 2015;29:592–599.

Laiolo P, Pato J, et al., Ecological and evolutionary drivers of the elevational gradient of diversity.
Ecology Letters 2018;21:1022–1032.

Lavorel S, Garnier E. Predicting changes in community composition and ecosystem functioning from
plant traits: revisiting the Holy Grail, Functional Ecology 2002;16:545–556.

Longino JT, Colwell RK. Density compensation, species composition, and richness of ants on a
neotropical elevational gradient, Ecosphere 2011;2:art29.

Macarthur R, Levins R. The Limiting Similarity, Convergence, and Divergence of Coexisting Species,
The American Naturalist 1967;101:377–385.

Macarthur RH. Patterns of Species Diversity, Biological Reviews 1965;40:510–533.

Mouillot D, Stubbs W, Faure M et al., Niche overlap estimates based on quantitative functional
traits: a new family of non-parametric indices, Oecologia 2005;145:345–353.

Mungee M, Athreya R. Intraspecific trait variability and community assembly in hawkmoths
(Lepidoptera: Sphingidae) across an elevational gradient in the eastern Himalayas, India, Ecology and
Evolution 2021;11:2471–2487.

Muscarella R, Uriarte M. Do community-weighted mean functional traits reflect optimal strategies?
Proceedings of the Royal Society B 2016;283:20152434.

Nylin S, Slove J, et al., Host plant utilization, host range oscillations and diversification in nymphalid
butterflies: a phylogenetic investigation, Evolution 2014;68:105–124.

Oksanen J, Kindt R, Legendre P et al., The vegan Package. 2009.

Olden JD. Biotic homogenization: a new research agenda for conservation biogeography, Journal of
Biogeography 2006;33:2027–2039.

Osorio-Canadas S, Flores- Hernández N, Sánchez Ortiz T et. al., Changes in bee functional traits at
community and intrasepcific levels along an elevational gradient in a Mexical-type scrubland,
Oecologia 2022,1: 145-148

Pellissier V, Barnagaud J-Y, Kissling WD et al., Niche packing and expansion account for species
richness–productivity relationships in global bird assemblages, Global Ecology and Biogeography
2018;27:604–615.

Peters MK, Peisker J, Steffan-Dewenter I et al., Morphological traits are linked to the cold
performance and distribution of bees along elevational gradients, Journal of Biogeography
2016;43:2040–2049.

Pigot AL, Trisos CH, et al., Functional traits reveal the expansion and packing of ecological niche
space underlying an elevational diversity gradient in passerine birds, Proceedings of the Royal
Society B: Biological Sciences 2016;283:20152013.

Pillar VD, Blanco CC, Müller SC et al., Functional redundancy and stability in plant communities,
Journal of Vegetation Science 2013;24:963–974.

Read QD, Grady JM, Zarnetske PL et al., Among-species overlap in rodent body size distributions
predicts species richness along a temperature gradient, Ecography 2018;41:1718–1727.

Read QD, Yue A, Fluck IE et al., The Ostats package. 2022.

Ribeiro DB, Freitas AVL. Large-sized insects show stronger seasonality than small-sized ones: a case
study of fruit-feeding butterflies, Biological Journal of the Linnean Society 2011;104:820–827.

Ricklefs RE, O'Rourke K. Aspect Diversity in Moths: A Temperate-Tropical Comparison, Evolution
1975;29:313–324.

Rogora M, Frate L, Carranza ML et al., Assessment of climate change effects on mountain
ecosystems through a cross-site analysis in the Alps and Apennines, Science of The Total
Environment 2018;624:1429–1442.

Ryding S, Klaassen M, Tattersall GJ et al., Shape-shifting: changing animal morphologies as a
response to climatic warming, Trends in Ecology & Evolution 2021;36:1036–1048.

Sanyal AK, Dey P, Uniyal VP et al., Geometridae Stephens, 1829 from different altitudes in Western
Himalayan Protected Areas of Uttarakhand, India.(Lepidoptera: Geometridae), SHILAP Revista de
Lepidopterología 2017;45: 143-63.

Schneider CA, Rasband WS, et al., NIH Image to ImageJ: 25 years of image analysis, Nature Methods
2012;9:671–675.

Schumm M, White AE, Supriya K et al., Ecological Limits as the Driver of Bird Species Richness
Patterns along the East Himalayan Elevational Gradient, The American Naturalist 2020;195:802–817.

Seifert CL, Strutzenberger P, et al., Ecological specialisation and range size determine intraspecific
body size variation in a speciose clade of insect herbivores, Oikos 2022;n/a:e09338.

Shi J, Chen F, et al., Differences in Wing Morphometrics of Lymantria dispar (Lepidoptera: Erebidae)
Between Populations That Vary in Female Flight Capability, Annals of the Entomological Society
America 2015;108:528–535.

Shipley B, Vile D, et al., From Plant Traits to Plant Communities: A Statistical Mechanistic Approach
to Biodiversity, Science 2006;314:812–814.

Shyy W, Kang C, Chirarattananon P et al., Aerodynamics, sensing and control of insect-scale flapping-
wing flight, Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences
2016;472:20150712.

Skou P. The Geometroid Moths of North Europe (Lepidoptera: Drepanidae and Geometridae), Brill
Archive, 1986.

Slove J, Janz N. The Relationship between Diet Breadth and Geographic Range Size in the Butterfly
Subfamily Nymphalinae – A Study of Global Scale, PLOS ONE 2011;6:e16057.

Strutzenberger P, Brehm G, Gottsberger B et al. Diversification rates, host plant shifts and an
updated molecular phylogeny of Andean Eois moths (Lepidoptera: Geometridae), PLOS ONE
2017;12:e0188430.

Svensson EI, Gómez-Llano M, et al.,. Out of the tropics: Macroevolutionary size trends in an old
insect order are shaped by temperature and predators, Journal of Biogeography 2023;50:489–502.

Thakur AK, Singh G, Singh S et al., Impact of Pastoral Practices on Forest Cover and Regeneration in
the Outer Fringes of Kedarnath Wildlife Sanctuary, Western Himalaya, Journal of Indian Society of
Remote Sensing 2011;39:127–134.

Trew BT, Maclean IMD. Vulnerability of global biodiversity hotspots to climate change, Global
Ecology and Biogeography 2021;30:768–783.

Violle C, Jiang L. Towards a trait-based quantification of species niche, Journal of Plant Ecology
2009;2:87–93.

Warrington MH, Waterman J. Temperature-associated morphological changes in an African arid-
zone ground squirrel. Goheen J (ed.).,Journal of Mammalogy 2023;104:410–420.

Weiher E, Keddy PA. Assembly Rules, Null Models, and Trait Dispersion: New Questions from Old
Patterns, Oikos 1995;74:159–164.

Xing S, Bonebrake TC, Ashton LA et al., Colors of night: climate–morphology relationships of
geometrid moths along spatial gradients in southwestern China, Oecologia 2018;188:537–546.

Yates ML, Andrew NR, Binns M et al., Morphological traits: predictable responses to macrohabitats
across a 300 km scale, PeerJ 2014;2:e271.

Figure Legends:

Map 1: Map showing the sampling locations in and around the buffer zone of Kedarnath

Wildlife Sanctuary, which is located in northern side of the mountainous state of Uttarakhand

(highlighted on the map of India on the left side)

Figure 1: Rarefied curves for a) Sample coverage and b) Species diversity patterns

extrapolated across the number of individuals; c) overall beta diversity measured as Sorensen

pairwise dissimilarity and d) the spatial turnover of the species measured as Simpson

pairwise dissimilarity across the low, mid and high elevation assemblages

Figure 2: The figure shows the community weighted means (mean ± SE) of each of the traits

across the low, mid and high-elevation assemblages.

Figure 3: The figure shows the comparison of subfamily-level trait dispersion in a) Low, b)

Mid and c) High elevation classes; and overall comparison of trait dispersion at the d)

assemblage level. The red-black shapes denote the centroid for the respective level of

comparison.

Figure 4: The figure shows the degree of overlap (O-statistic) of the multivariate trait-space,

based on body length (BL), maneuverability (MV), and wing loading (WL) of the Geometrid

moth assemblages in low, mid, and high elevations

1: (overlap = 0.84)

- High
- Low
- Mid