Two distinct integrin binding sites on MMP9 drive cancer invasion by mediating integrin

membrane trafficking & stabilization

Sarbajeet Dutta[1], Soumili Sarkar[1], Simran Tolani[1], Md. Asrafuddoza Hazari[1], Shamik Sen[1*]

[1]Dept. of Biosciences & Bioengineering, IIT Bombay, Mumbai 400076

*Correspondence: shamiks@iitb.ac.in

Abstract

Matrix stiffening has been established to drive cancer progression through increased activity
of matrix metalloproteases (MMPs) which degrade the matrix creating paths for migration.
However, the non-proteolytic functions of MMPs in cancer invasion remain relatively less
understood. Here we have probed the importance of proteolytic and non-proteolytic
functions of MMP9, which exhibits robust stiffness dependent expression and secretion in
highly invasive cancer cells. We show that while MMP9 sustains spreading and 2D migration
non-proteolytically by stabilizing focal adhesions, MMP9 proteolytic activity is essential for
3D invasion. We then establish the function of two distinct integrin β1 (ITG β1) binding sites
on MMP9, with the hemopexin domain mediating co-packaging and co-transport of ITG
β1/MMP9 to the cell periphery, and the RGD domain stabilizing ITG β1 on the cell membrane
prior to matrix degradation. Together, our results illustrate how MMP9 optimizes cancer
invasion by spatiotemporally integrating matrix remodeling with adhesion formation.

22
23
24

25   **Keywords:** MMP9, invasion, integrin β1 (ITGβ1), matrix degradation, non-proteolytic

26   function

Introduction

The extracellular matrix (ECM) constitutes the acellular component of all tissues [1], [2], [3]. Cells residing within the tissue deposit a variety of protein/non-protein components to form this complex 3D structure. Although ECM homeostasis is strictly regulated in normal tissues, this homeostasis is compromised in cancerous tissues [4]. Excess deposition of fibrillar collagen and crosslinking by enzymes such as lysyl oxidase leads to matrix stiffening in multiple epithelial cancers including breast cancer and pancreatic cancer [5], [6]. Altered cell-matrix signaling is associated with malignant transformation leading to epithelial-to-mesenchymal transition (EMT) [7] through increased secretion of matrix degrading enzymes such as matrix metalloproteinases (MMPs). Indeed, expression of multiple MMPs including MMP1, 2, 3, 7, 9, 13, and 14 are closely associated with cancer invasion and worse prognosis [8], [9], [10], [11], [12].

The multidomain structural architecture of MMPs with different substrate specificities is well documented in the literature. While MMP1, 8, 13, and 18 mostly act on fibrillar collagen, MMP2 and 9 can also degrade non-fibrillar collagen and gelatin [13][14]. Despite the structural similarities in their catalytic domain, pro-peptide, and signal peptide, individual MMPs have distinct structural traits that define their functional properties. While most membrane bound MMPs have a transmembrane/cytoplasmic domain (MMP14, 15, 16, and 24), MMP17 and 25 have GPI anchoring to the membrane. Except for MMP7 and 26, all MMPs have hemopexin repeats; in addition, MMP2/9 possess fibronectin type II repeats in their catalytic domains, which can purportedly bind to gelatin, collagen types I and IV, and laminin [14].

The proteolytic activity of MMPs mediated by their catalytic domain has been studied extensively and is well established. For example, overexpression of MMP3 and 14 has been shown to induce breast cancer and melanoma [15], [16], whereas MMP2 and 14 promote cell invasion by cleaving laminin 5 [17], [18]. MMP3 and 7 degrade E-cadherin in MCF7 and MDCK cells, causing cancer cells to separate from the tissue [19]. A recent study has implicated MMP9 in tumor progression in triple-negative breast cancer [20]. MMP9 expression increases during the early stages of melanoma development [21]. In the animal model, inhibiting MMP9 lowers lung metastases [22].

Beyond their proteolytic function, a growing body of literature has identified the importance of non-proteolytic activity of MMPs mediated by their hemopexin, transmembrane, and cytoplasmic domains, via interactions with diverse cell surface proteins that aid in invasion and cancer growth. According to emerging evidence in the literature, MMP9 and 14 can localize on the cell membrane via a variety of mechanisms. MMP14 interacts with CD44, a hyaluronan receptor, and promotes invasion by releasing CD44 from the cell membrane in pancreatic and breast cancer [23]. MMP14 has also been shown to engage integrin β1 (ITGβ1) via its transmembrane/cytoplasmic domain and activate MAPK in mouse mammary epithelial cells leading to branching morphogenesis [24]. The interaction of CD44 with MMP9 via the hemopexin domain leads to enhanced invasion and angiogenesis in mouse mammary carcinoma and human melanoma cells [25]. In melanoma cells, MMP9 has also been shown to interact with ITGβ1 via its hemopexin domain, leading to the formation of cell-substrate adhesions [26], [27].

While insightful, these studies have not parsed the proteolytic and non-proteolytic functions of MMP9 in cancer invasion. We address this question by first showing that among several MMPs, MMP9 exhibits strong stiffness dependent expression and secretion in highly invasive cancer cells. Cell rounding and loss of cell motility upon inhibition of MMP proteolytic activity are attributed to defects in adhesion stability [28]. While similar effects were observed upon MMP9 knockdown, restoration of cell spreading and 2D motility in knockdown cells re-expressing catalytically inactive MMP9 through stabilization of adhesions, but no restoration of invasion in 3D collagen matrices, suggests that MMP9 non-proteolytically regulates adhesion formation, with matrix degradation essential for 3D invasion. We then show that two ITGβ1 binding sites, a RGD motif in the catalytic domain and a previously identified peptide sequence in the hemopexin domain [29], collectively regulate adhesion formation. While MMP9/ ITGβ1 association via the hemopexin domain is necessary for integrin trafficking to the membrane and MMP9 secretion, the RGD motif is responsible for stabilizing ITGβ1 on the cell membrane. Collectively, our study illustrates how invasiveness is optimally sustained by MMP9 through a combination of path generation via proteolytic remodeling and stabilization of integrins at the leading edge.

Materials & Methods

TCGA Data analysis: The data from The Cancer Genome Atlas (TCGA) was obtained through curation of the UCSC Xena Browser, an online resource accessible at https://xenabrowser.net/. The analysis focused exclusively on the TCGA breast cancer (BRCA) and TCGA lung cancer (LUNG) datasets, among the various types of cancer data available. The gene expression levels of MMP1, MMP2, MMP9, and MMP14 were determined by RNA sequencing (Illumina HiSeq) in both cancerous and normal tissue samples. Subsequently, the data obtained was organized and subjected to analysis. A total of 1097 individuals diagnosed with breast cancer and 114 healthy individuals were included in the breast cancer analysis. Similarly, the lung cancer analysis comprised 1017 individuals diagnosed with lung cancer and 110 healthy individuals.

Cell culture & reagents: In our study, the following human cell lines were used: HT1080 invasive fibrosarcoma cell line, MDA-MB-231 invasive breast cancer cell line, A549 lung carcinoma cell line, MCF-7 breast cancer cell line, MCF-10A breast epithelial cell line, and Normal Human Dermal Fibroblasts (HDF) cells isolated from the healthy human tissue sample. The cancer cells utilized in this study were procured from the NCCS Cell Repository in Pune. The HEK293T cells utilized for lentivirus generation were generously provided by the laboratory of Dr. Prakriti Tayalia. Cells were cultured in Dulbecco's Modified Eagle Medium (DMEM) (Cat # 12100046, Gibco) supplemented with 10% fetal bovine serum (FBS) (Cat. # 10270106, Gibco) and 1% antibiotic (Cat. # A002, HiMedia) under 5% CO_2 conditions. For drug treatment experiments, GM6001 (Cat. # 2983, Tocris Bioscience) and SB-3CT (Cat. # 6088, Tocris Bioscience) were used at a working concentration of 10 µM [30] and ARP100 (Cat. # 2621, Tocris Bioscience) was used at a working concentration 50nM [31].

Cloning and generation of stable cell lines: MMP9 full-length cDNA was cloned from HT1080 cDNA to create expression plasmids using gene-specific cloning primers M9_F_KpnI and M9_R_NotI (Supplementary Table 1). The amplified product was first cloned into pcDNA3 (Addgene plasmid# 13031), followed by MMP9-fused eGFP that was subcloned into the pLenti-puro vector (Addgene plasmid# 39481) to generate the MMP9 full-length expression plasmid (M9FL) using primer pairs M9_F_EcoRI and eGFP_R_XbaI. To generate a catalytic inactive mutant (Δcat), the primer pairs M9_E4A_F and M9_E4A_R were used to introduce a point mutation of glutamic acid (E) to alanine (A) at amino acid position 402 (E402A) in M9FL.

RGE, ΔB1, and ΔB1Δcat plasmids were obtained by amplifying M9FL and M9Δcat plasmids

with the primer pairs M9_RGE_F and R and M9_B1_F and R, respectively (Supplementary

Table 1).

Stable MMP9 knockdown cell lines were generated using two independent shRNA sequences

(CCACAACATCACCTATTGGAT and CAGTTTCCATTCATCTTCCAA; Sigma Aldrich). The transduced

cells were selected using puromycin (Catalogue number CMS8861, HiMedia). Re-expression

of M9FL and other MMP9 cDNAs was done using lentiviral transduction, produced by

HEK293T cells according to the standard protocol [32]. Cells were transduced with lentiviral

particles in the presence of polybrene (8 g/mL, Cat# TR-1003-G, Merck); after 48 hours of

transduction, positive cells were sorted using fluorescent-assisted cell sorting (FACS).

Cell Experiments on Polyacrylamide (PA) gels: PA gels were prepared as per previously

established lab protocols to generate 0.5 kPa soft gels mimicking the stiffness of normal breast

tissue and 5 kPa malignant tissue [30]. Gels were functionalized with 0.1M sulfo-succinimidyl

6-(4'-azido-2'-nitrophenylamino) hexanoate (Sulfo-SANPAH) (Cat# 22589, Thermo Scientific)

dissolved in HEPES before overnight incubation in collagen type-I solution (Catalogue #

A1048301, Thermo Scientific) at 4^0C to achieve a theoretical coating density of 10 μg/cm^2. For

experiments, cells were seeded sparsely (2000 cells/cm^2) and cultured for 24 hours before

imaging and motility experiments; cells were seeded at high density (20000 cells/cm^2) for RT-

PCR, western blotting, and gelatin zymography experiments.

RT-PCR, western blotting, zymography and Co-IP: Total RNA was isolated using TRIzol (Cat#

15596026, Invitrogen) reagent, followed by cDNA synthesis using High-capacity cDNA

synthesis KIT (Cat# 4368814, Thermofisher). Next, the cDNA was used in qPCR using

PowerUp^{TM} SYBR^{TM} Green Master Mix (Cat# A25742, Applied Biosystem) and gene-specific

primers (Supplementary Table 1) using Quantstudio^{TM} 5. Cyclophilin A (CycA) gene was used

as an internal housekeeping gene. Gene fold change was calculated using the $\Delta\Delta C_t$ technique

[33].

For western blotting, cell lysates were prepared using RIPA buffer (Sigma Aldrich Cat. No.

R0278) with 1% protease and phosphatase inhibitors. Equal amounts of denatured proteins

were loaded into 10% PA-gels and run for 2 hours at 90V. After protein transfer onto methanol-

activated PVDF membranes (Cat# 1620177, Bio-Rad), membranes were blocked using a 1%

gelatin solution for 1 hour at RT. The following primary antibodies were used: mouse anti-

GAPDH antibody (Cat# AC033, ABclonal), rabbit anti-MMP9 antibody (Cat# MA5-45511,

Invitrogen), rabbit anti-ITGβ1 antibody (Cat# PA578028, Invitrogen), and rabbit anti-pMLC

antibody (Cat# 3674, Cell Signaling). The following secondary HRP-conjugated antibodies were

used: anti-rabbit IgG (Cat. No. 31460, Invitrogen) and HRP-conjugated anti-mouse IgG (Cat.

No. 31430, Invitrogen). Finally, the blots were developed in a ChemiDoc^{TM}MP Gel Imaging

System (Biorad) with ECL substrate solution (Cat. No. K-12045-D20, Advansta). The Fiji-ImageJ

software was used to perform densitometric analysis.

Gelatin zymography was performed using cell-secreted conditioned media (CM) to investigate

the secretion profile of MMP2/9 by different cell types as described previously [30]. Briefly,

the CM collected 24 hours after culture, was run in 10% PA gels containing 0.1% gelatin type

B (Cat# 83740, SRL Chemical) using a non-denaturing sample loading buffer. The samples were

run at a voltage of 130V and at 4^0C until the dye front exited the bottom of the gels. After

rinsing with renaturing buffer (2.5% TritonX-100 in distilled water) at RT, the gels were

incubated in developing buffer (50 mM Tris-base, 50 mM Tris-HCL, 0.2 mM NaCl, 5 mM CaCl2

and pH adjusted to 7.4) for 24 hours at 37^0C. Next, after incubation for 1 hour at RT with

staining solution (40% methanol, 10% acetic acid, 0.1% Coomassie Brilliant Blue in distilled

water), the gels were rinsed with detaining solution (40% methanol, 10% acetic acid in distilled

water) until the degradation bands became visible. Gel images were captured using Bio rad's

ChemiDoc^{TM}MP Gel Imaging System.

The Protein-G CoIP kit (IP50, Sigma Aldrich) was used for coimmunoprecipitation (CoIP)

experiments. After lysing cells in non-denaturing lysis buffer (20 mM Tris HCl at pH 8, 137 mM

NaCl, 10% glycerol, 1% NP-40, and 2 mM EDTA), cell lysates were incubated overnight with

either the anti-ITGβ1 antibody (Cat# PA578028, Invitrogen) or the control IgG antibody (Cat#

CR1, Sino Biological Inc.) in an IP spin column at 40 °C. Next, 30μL of Sepharose beads were

added and incubated again overnight at 40 °C. After spin, the beads were washed 6-7 times

with IP buffer. Finally, the bound protein was eluted by incubating the beads with 1x Laemmli

buffer at 95^0C for 15 minutes. SDS-PAGE was used to load 20 μL of eluted protein and equal

amounts of whole cell lysate. Western blotting was performed using antibodies against MMP9

(Cat# MA5-45511, Invitrogen) and GAPDH (Cat# AC033, ABclonal).

Immunocytochemistry (ICC) & Microscopy: For staining, cells were fixed with 4%

paraformaldehyde (PFA) (Cat# 158127, Sigma Aldrich) for 15 minutes at room temperature

(RT), washed three times with DPBS, and permeabilized with 0.25% TritonX-100 (Cat# 64518,

SRL Chemicals) for 10 minutes at RT. After blocking with 5% BSA (Cat# 1650500501730, Genei)

for 1hr at RT, cells were incubated overnight at 40C with one or more of the following

antibodies: rabbit anti-pMLC (ser19) (Cat# 3675, Cell Signaling), rabbit anti-paxillin (Cat#

PA534910, Invitrogen), rabbit anti-ITGβ1 (Cat# PA578028, Invitrogen), and rabbit anti-MMP9

(Cat# MA5-45511, Invitrogen). In addition, cells were incubated with Wheat Germ Agglutinin

(WGA) conjugated with Alexa Flour (AF) 647 (Cat# W32466, Invitrogen) to stain cell glycocalyx

and phalloidin conjugated with Alexa Flour (AF) 555 (Cat# A34055, Invitrogen) to stain F-actin

stress fibre. The next day, after washing with DPBS, cells were incubated with one of the

following secondary antibodies at RT for 2 hours: anti-rabbit Alexa Flour-488 conjugated

antibody (Cat# A-11008, Invitrogen), anti-rabbit Alexa Flour-555 conjugated antibody (Cat# A-

21428), anti-rabbit Alexa Flour-647 conjugated antibody (Cat# A-31573, Invitrogen), and anti-

mouse. The coverslips were treated with DAPI (Cat# D9542, Sigma Aldrich) for 15 minutes at

RT after three DPBS washes. After mounting coverslips, samples were imaged using an

Olympus IX83 inverted fluorescence microscope and/or an LSM 780 Carl Zeiss confocal

microscope at 60x magnification. Images were analyzed with Fiji-ImageJ. Filament-Sensor

software [34] was used to measure F-actin stress fibre length and count.

To conduct the Proximity Ligation Assay (PLA) in order to identify the interactions between

ITGβ1 and CD44, the Duolink® In Situ Red Starter Kit Mouse/Rabbit (DUO92101, Sigma-

Aldrich) was used, following the manufacturer's instructions. In short, the cells were fixed with

4% PFA for 10 minutes, washed twice with PBS followed by permeabilization with 0.25%

Triton-X100 solution for 15 minutes. After washing with PBS twice, the cells were blocked for

one hour at 37^0C in a humid chamber using the supplied blocking solution in the kit. Next, an

overnight incubation at 4^0C was carried out for the primary antibodies against anti-MMP9 and

anti-integrin β1. After that, the cells were washed twice using 1x washing solution A.

Subsequently, one-hour incubation of the Duolink PLUS/MINUS secondary antibody in a

humid chamber at 37^0C was performed. After washing twice with solution A In a humid 37^0C

chamber, a ligation reaction was carried out for 30 minutes using the Duolink ligation regent.

This was followed by an amplification step using Duolink amplification reagents for two hours

in the dark humid 37^0C chamber. Wash solution B was used for the last washing, and Duolink

DAPI-containing mounting media was used for the mounting process. Images were captured

in an Olympus IX83 inverted fluorescence microscope at 60x magnification.

Focal adhesion (FA) dynamics were assessed by paxillin-mRFP overexpressing cells seeded at

low density on glass bottomed dishes (Cat# D35-C4-20-1.5N, CellVis). Time-lapse images were

collected using a spinning disc confocal microscope (Yokogawa Electric Corporation, CSU-X1)

at 15s intervals for a duration of 10min. After subtracting the background in Fiji-ImageJ,

adhesion lifetimes were estimated using the open access Focal Adhesion Analysis Server

(FAAS) [35].

For studying the localization dynamics of MMP9 and ITGβ1, GFP-fused MMP9-expressing cells

were transfected with ITGβ1-mRFP cDNA. These cells were seeded on glass-bottomed dishes

at low density for imaging. Time-lapse images were collected on a spinning disc confocal

microscope at 5s intervals for 10 min at low laser power to prevent bleaching. Analysis was

performed using Fiji-ImageJ. After subtracting the background, ITGβ1 foci were tracked, and

diffusion speed was quantified using manual tracking. Additionally, the ratio of the ITGβ1 foci

moving towards the cell migration front to the foci internalized was quantified. For kymograph

experiments, cells were sparsely seeded on collagen-coated 24 well plates, and images of live

cells were acquired at 5 s intervals for 15 minutes using a phase-contrast microscope (Olympus

IX83, 20x objective). Protrusion and retraction rates were quantified using the previously

described protocols [36].

Single cell biophysics measurements: For 2D cell motility experiments, a total of 15,000 cells

were seeded onto each 12mm polyacrylamide (PA) gel coverslips. After allowing cells to settle

for 12 hours, images were captured at 20-minute intervals over 12 hours using an Olympus

IX83 inverted microscope equipped with a 10x objective. For 3D collagen invasion

experiments, cells were encapsulated within 1.2 mg/ml 3D collagen matrices as done

elsewhere [30], [37]. Timelapse images were recorded at 20 min intervals for 12 hours. This

was accomplished using an Olympus IX83 inverted microscope equipped with a 10x objective

lens. The velocity of cells in both 2D motility and 3D invasion was measured using the manual

tracking plug-in of the Fiji-ImageJ software. Subsequently, an Excel macro, as detailed in a

separate publication [38], was utilized.

For gel compaction assay, 5×10^4 cells were copolymerized with 1.2mg/mL collagen (Cat#

354236, Corning) to a total volume of 150μL in 48 well plates. After 24hr, the polymerized gels

were released from the sides using a needle. After 48hr, gel images were captured in an

Olympus IX83 phase contrast microscope at 4x magnification. The quantification of the gel

area was done in Fiji-ImageJ software using the following equation:

$$\% \, of \, compaction = \frac{Well \, Area - final \, gel \, area}{Well \, Area} \, x \, 100$$

Atomic force microscopy (AFM) was used to measure the cortical stiffness of cells. Pyramidal

AFM probes (10 kHz, Cat # TR400PB, Olympus) with a nominal spring constant of 30pN/nm

were calibrated thermally to obtain the precise value of the spring constant. Cells were grown

at low density on stiff PA gels (5 kPa) on 12mm coverslips for 12-16 hours. Indentation was

done in the cell cortex area to measure the force curve, and the data was fitted in the Hertz

model till 500nm.

$$F = \frac{3}{4} \, X \, E \, \frac{\tan(\alpha)\delta^2}{(1 - \vartheta^2)}$$

Here, F = indentation force, E = elastic modulus of the material, α = half tip angle, δ =

indentation depth, ν = Poisson ratio of the material (assumed to be 0.45 for the cell).

Adhesion tests using AFM were carried out to detect RGD-ITGβ1 interactions. Cells were

grown on 5 kPa stiff PA-gels for 12-16h. The spherical probe (67 kHz, Novascan) was coated

with 10μg/mL poly-L-lysin solution for 12 hours and treated with 0.5% glutaraldehyde for 4

hours. Ultrapure water is used three times to wash the probes to remove surplus solutions.

Next, the probe was treated with 10μg/mL RGD peptide for 2 hours. The probe was rinsed

with ultrapure water to eliminate unbound RGD. During the experiment, the functionalized

probe was held on the cell surface for 10s to generate RGD integrin bonds, then retracted at

4μm/s. In this step, we recorded the retraction curve and estimated the adhesion force using

Igor Pro and MATLAB.

Statistical Analysis: The Kolmogorov-Smirnov normality test was used to examine data

distribution. A parametric or nonparametric statistical test was done based on the outcome.

For parametric data, one-way ANOVA was employed for statistical analysis, and Tukey's test

was utilized to compare means. For non-parametric data, the Mann-Whitney test was used.

Prism 9 (GraphPad Software, LLC) was used for statistical analysis and generating graphs, with

P<0.05 considered statistically significant.

Results

ECM stiffness differentially modulates MMP expression and activity in cancer cells

Pan-cancer analysis of the TCGA database using the UCSC Xena browser (Fig. 1A) revealed

elevated expression of MMP1, 9, and 14 in primary and secondary tumor sites compared to

solid benign tumors, with MMP2 transcript levels remaining unchanged (Fig. 1B). Similar trend

observed in breast and lung cancers establishes a strong positive correlation between

overexpression of MMP1, 9, and 14 with tumorigenicity and malignancy (Fig. 1C, D).

Since matrix stiffening is implicated in cancer progression [30], stiffness dependent expression

profiling of MMP1/MMP2/MMP9/MMP14 was carried out on soft (0.5 kPa) and stiff (5 kPa)

polyacrylamide (PA) gels mimicking normal and metastatic tissue stiffness (Fig. 1F) [30].

Experiments were performed using normal human dermal fibroblasts (HDFs), non-cancerous

MCF10A breast epithelial cells, non-invasive ER^+PR^+ MCF7 breast cancer cells, moderately

invasive A549 lung adenocarcinoma cells, and highly invasive MDA-MB-231 triple-negative

breast cancer (TNBC) cells and HT1080 fibrosarcoma cells (hereafter referred to as MDA and

HT cells, respectively). While MMP1, MMP2, and MMP14 levels were comparable across

HDFs, MDA, and HT cells, more than 100-fold MMP9 overexpression was observed in MDA

and HT cells (Fig. 1F). In addition, MDA and HT cells also exhibited ~2-fold stiffness dependent

MMP9 upregulation on 5 kPa gels compared to 0.5 kPa gels. Consistent with these

observations, the western blot of cell lysates revealed elevated expression of MMP9 and the

mesenchymal marker integrin-β1 in all the cells cultured on 5 kPa gels (Fig. 1G, H and Supp.

Fig. 1A). Stiffness-mediated MMP2 upregulation was observed in MCF-10A and MDA only.

Assaying MMP2/MMP9 secretion using cell-secreted conditioned media (CM), revealed

significant stiffness dependent MMP9 activity in gelatin zymogram from highly invasive MDA

and HT cells only, with no stiffness dependent alteration in MMP2 activity (Fig. 1I, J and Supp.

Fig. 1B). Collectively, these results suggest that high baseline MMP9 expression and stiffness

dependent MMP9 modulation are hallmarks of highly metastatic cancer cells.

Soluble MMPs drive cell spreading and migration by regulating focal adhesions

To probe the role of soluble MMPs (i.e., MMP2 and MMP9) in mediating stiffness dependent

invasive phenotype, we measured cell spreading and motility of A549, MDA, and HT cells on

soft (0.5 kPa) and stiff (5 kPa) PA gels in the presence and absence of the pan-MMP inhibitor

GM6001 (hereafter referred to as GM) and the soluble MMP2/9 inhibitor SB3-CT (hereafter

referred to as SB3). DMSO-treated and untreated cells served as controls. While both GM and

SB3-treated cells exhibited a dramatic drop in cell spreading and increased cell rounding on

stiff PA gels (Fig. 2A, B and Supp. Fig. 2C), no significant changes were observed on soft PA gels

(Supp. Fig. 2A, B and C). In line with these observations, cell motility was markedly reduced

on stiff gels (Fig. 2C, D, Supp. Fig. 2E and Supp. Movie 2) with no changes on the soft gels

(Supp. Figs. 2D, F and Supp. Movie 1). Similarly, when encapsulated in 3D collagen gels, cell

invasiveness was inhibited in GM and SB3 treated cells (Fig. 2E, F, Supp. Fig. 2G, and Supp.

Movie 3).

Defects in migration can be attributed to reduced cytoskeletal forces and/or perturbed

adhesion formation and dynamics. Both GM and SB3 treatment led to prominent cell

softening in A549, MDA, and HT cells (Fig. 2G, H and Supp. Fig. 2I), and ~50% drop in the extent

of gel compaction (Fig. 2I, J and Supp. Fig. 2H) indicative of a compromised cytoskeleton, as

evident from a marked reduction in the number of F-actin stress fibres/cell (Fig. 2K, L and

Supp. Fig. 2J). Since focal adhesions stabilize the cytoskeleton, we next probed adhesion size

distribution and adhesion stability in GM/SB3 treated cells. Visualization of paxillin-stained

focal adhesions revealed a dramatic drop in the average number of focal adhesions in drug-

treated cells (Fig. 2K, M and Supp. Fig. 2J). Furthermore, time-lapse images of cells stably

transfected with GFP-paxillin revealed a marked reduction in focal adhesion (FA) lifetime upon

SB3 treatment (Fig. 2N, O, Supp. Fig. 2K and Supp. Movie 4). Reduction in FA stability was

associated with a drop in the secretion of MMP2 and MMP9 and a reduction in ITGβ1

expression (Fig. 2P) in HT1080 cells. Together, these results suggest that cell secreted soluble

MMPs mediate cell spreading and migration by modulating adhesion stability and turnover.

Both proteolytic and non-proteolytic functions of MMP9 are essential for sustaining cancer invasion

In contrast to SB3, treating MDA and HT cells with ARP100 at concentrations that inhibited only MMP2 [31], did not impact cell spreading and motility (Supp. Fig. 2L, M and Supp. Movie 5), suggesting that defects in cell spreading and motility observed in SB3-treated cells is attributed to MMP9. To test the role of MMP9 directly, we established shCTL (i.e., cells transfected with scrambled shRNA) and shKD1/shKD2 knockdown cells (i.e., MMP9 was knocked down using two distinct shRNAs) in HT cells. Knockdown cells were re-transfected with MMP9 full-length cDNA to generate M9FL1 and M9FL2 cells for recovery experiments. For probing non-proteolytic functions of MMP9, knockdown cells were also transfected with catalytically inactive MMP9 (E402A) cDNA to generate ΔCAT cells (Fig. 3A, Supp. Fig. 3A and B) [39]. While comparable levels of MMP9 were detected in conditioned media (CM) secreted by M9FL1/M9FL2 cells, degradation bands in gelatin zymograms were detected in shCTL and M9FL cells only (Fig. 3B).

In contrast to shCTL cells which were well spread and polarized, shKD1/shKD2 cells exhibited reduced spreading with rounded morphologies similar to SB3 treated cells (Fig. 3C and D). Remarkably, cell spreading was rescued not only in M9FL1/M9FL2 cells but also in ΔCAT cells. In line with these observations, reduction in cell motility on 5 kPa gels observed in shKD1/shKD2 cells, was rescued in both M9FL1/M9FL2 and ΔCAT cells (Fig. 3E, F and Supp. Movie 6). However, when entrapped in 3D collagen gels, in contrast to shCTL and M9FL cells which migrated at comparable speeds, both shKD1/shKD2 and ΔCAT cells were unable to migrate (Fig. 3G, H and Supp. Movie 7). These results highlight the importance of MMP9 catalytic activity in sustaining invasion through dense matrices.

To probe the non-proteolytic functions of MMP9, cytoskeletal organization and focal adhesion dynamics were compared across the different conditions. Reduction in stress fibre formation in shKD1/shKD2 cells (Fig. 3I, J) was associated with reduced actomyosin contractility (assessed using pMLC staining) (Supp. Fig. 3C and D) and gel compaction (Fig. 3M and N), and increased cell softening (Fig. 3K, L). However, re-expression of full length MMP9 or MMP9 ΔCAT restored cytoskeletal organization, contractility, and cell stiffness to levels comparable to shCTL cells.

Cell migration involves cycles of protrusion and adhesion stabilization at the cell front. Since

kymograph analysis revealed perturbed protrusion dynamics in shKD1/shKD2 cells only, but

not in ΔCAT cells (Supp. Fig. 3E-G), we hypothesized MMP9 ΔCAT is fully capable of stabilizing

adhesions. Indeed, in direct contrast to shKD1/shKD2 cells where the frequency of focal

adhesions and their stability were significantly reduced, both focal adhesion number and

lifetime were comparable in M9FL and ΔCAT cells (Fig. 3O-R and Supp. Movie 8). Together,

these results suggest that proteolytic and non-proteolytic functions of MMP9 collectively

sustain cancer invasion, with MMP9 stabilizing adhesions in a non-proteolytic manner and

proteolytic remodeling of the matrix.

MMP9-integrin β1 (ITGβ1) interaction via two distinct sites is essential for sustaining cancer

invasion

How does MMP9 stabilize adhesions? In a previous study, hemopexin domains of MMP14

have been shown to mediate binding with integrin β1 (ITGβ1) and activate MAPK [40].

Another study identified a 17-amino acid motif in blade IV of the hemopexin domain that

mediates integrin binding [29] (Fig. 4A). MMP9 catalytic domain is also known to harbour

three fibronectin type II like repeats [41], [42]. Bioinformatics analysis revealed the presence

of only one RGD motif in the catalytic domain of MMP9 (Fig. 4A). Based on the background

literature, and our bioinformatics analysis, we speculated MMP9 might regulate focal

adhesions non-proteolytically through its interactions with ITGβ1 (Fig. 4A). To test our

hypothesis, we generated two mutations at these sites: an RGD-to-RGE substitution and a

deletion of 17 amino acids in the PEX4 domain (Δβ1) (Fig. 4B). This was done in both M9FL

and ΔCAT cells, with these cell lines referred to as RGE, Δβ1 and Δβ1ΔCAT cells, respectively.

Thus, while matrix proteolysis is expected to remain unchanged in M9FL, RGE and Δβ1 cells,

integrin binding is expected to be perturbed in RGE, Δβ1 and Δβ1ΔCAT cells (Fig. 4B). Co-

immunoprecipitation (CoIP) experiments confirmed that interaction between MMP9 and

ITGβ1 was eliminated in Δβ1 and Δβ1ΔCAT cells, but not in M9FL, ΔCAT and RGE cells (Fig. 4C).

Proximity ligation assay (PLA) enabled us to visualize how MMP9/ITGβ1 association was

spatially distributed (Fig. 4D). While co-localization of MMP9 (red) with PLA signal (green) was

prominently observed in M9FL, ΔCAT and RGE cells at the cell periphery (marked by white

dotted circles), this was markedly reduced and/or absent in Δβ1 and Δβ1ΔCAT cells, as evident

from quantification of the integrated density of the PLA signal (Fig. 4E, F).

To next assess if MMP9/ ITGβ1 association influenced MMP9 secretion, gelatin zymography

experiments were performed using cell secreted conditioned media (CM) from the five cell

lines. As expected, MMP9 band was prominently detected in M9FL cells only (Fig. 4G). While

the absence of MMP9 band in ΔCAT cells was expected, MMP9 band intensities were

substantially reduced in both RGE and Δβ1 cells. Consistent with this, no MMP9 bands were

detected in western blotting of CM secreted by RGE, Δβ1 and Δβ1ΔCAT cells, suggesting that

MMP9/ ITGβ1 interaction is required for MMP9 secretion.

To assess the functional importance of MMP9/ ITGβ1 association, cell spreading, motility and

invasion experiments were performed as described above. Loss of MMP9/ ITGβ1 association

in RGE, Δβ1 and Δβ1Δcat cells led to pronounced reduction in cell spreading and increase in

cell rounding (Fig. 4H, I, Supp. Fig. 4A). However, the addition of CM from M9FL cells led to

near complete recovery in cell spreading and polarized morphology. RGE, Δβ1 and Δβ1Δcat

cells, which exhibited minimal spreading, also exhibited defective motility, which was rescued

upon the addition of M9FL CM (Fig. 4J, K, Supp. Fig. 4C and Supp. Movie 9 and 10). Similar

results were obtained when cells were entrapped in 3D collagen gels and their invasiveness

was tracked, i.e., RGE, Δβ1 and Δβ1Δcat cells were unable to invade along with Δcat, but

invasiveness was rescued by adding M9FL CM (Fig. 4L, M, Supp. Fig. 4D and Supp. Movie 11

and 12). Taken together, these results suggest that MMP9/ ITGβ1 association is essential for

sustaining cancer invasion.

MMP9-integrin association via PEX4 domain mediates ITGβ1 membrane trafficking while

RGD domain of MMP9 mediates ITGβ1 membrane stabilization

To understand the mechanisms underlying defective spreading and motility of RGE, Δβ1, and

Δβ1Δcat cells, we first assessed if adhesion formation/dynamics and/or cytoskeletal

organization were negatively impacted. Paxillin and stress fiber staining showed that

disruption of MMP9/ITGβ1 association led to a reduction in both the number focal adhesions

as well as the number of stress fibers in RGE, Δβ1, and Δβ1Δcat cells (Fig. 5A, B, Supp. Fig. 5A

and B). Timelapse imaging of paxillin-eGFP transfected cells revealed adhesion stability was

reduced in RGE, Δβ1 and Δβ1Δcat cells compared to M9FL/Δcat cells (Fig. 5C, D, Supp. Movie

13).

Smaller adhesion formation and reduced adhesion stability can be attributed to reduced

availability of integrins at the cell membrane and/or their membrane stability. To test this

possibility, we performed adhesion experiments using AFM as detailed elsewhere [30]. In

brief, cells were indented with RGD functionalized spherical AFM tips; upon indentation, the

tip position was held fixed for 10 seconds to allow for adhesions to form, and then retracted

(Fig. 5E). Bonds of adhesion formed between the cell surface and the tip, if any, had to be

broken for the tip to retract, leading to a characteristic peak in the retraction curve and

corresponds to the force required to break the adhesions. As evident from the representative

retraction curves (Fig. 5E), compared to M9FL/Δcat cells, the maximum adhesion force was

reduced by nearly 50% in RGE, Δβ1 and Δβ1Δcat cells (Fig. 5F), indicative of reduced ITGβ1

localization and/or stability at the cell membrane. Consistent with these observations,

MMP9/ITGβ1 dual staining revealed MMP9/ITGβ1 co-localization at the migration front in

M9FL and Δcat cells (Fig. 5G, Supp. Fig. 5C). In comparison, MMP9/ITGβ1 colocalization at the

cell periphery was reduced in RGE cells, and completely lost in Δβ1/Δβ1Δcat cells, where both

MMP9 and ITG β1 exhibited cytoplasmic localization.

Both MMP9 and ITGβ1 are trafficked to the cell membrane via vesicles [43], [44], [45]. Since

MMP9 secretion was significantly reduced in Δβ1/Δβ1Δcat cells, we hypothesized that altered

localization of both MMP9 and ITGβ1 in Δβ1/Δβ1Δcat cells is attributed to defective vesicular

trafficking, with MMP9 and ITGβ1 association ensuring the are being co-transported. To test

this, we performed three colour timelapse imaging wherein M9FL, Δcat, RGE, Δβ1 and

Δβ1Δcat cells (tagged in green) were transfected with ITGβ1 mRFP, and vesicles were tagged

using bio-tracker dye (Blue) (Fig. 5H and Supp. Fig. 5D). Indeed, in M9FL, Δcat and RGE cells,

movement of co-localized dots towards the cell periphery is indicative of co-transport of

MMP9 and ITGβ1 in the same vesicles (Supp. Movie. 14-18). In contrast, in Δβ1 and Δβ1Δcat

cells, co-transport was near completely abolished. Quantification of speed of individual ITGβ1

foci revealed comparable rates of transport in M9FL, Δcat and RGE cells, and substantially

reduced transport in Δβ1 and Δβ1Δcat cells (Fig. 5I).

While AFM experiments revealed drop in peak forces in RGE cells, no differences in ITGβ1

transport was observed in these cells. To test if this apparent anomaly can be explained by

altered ITGβ1 stability, we tracked the ratio (M/N) of the number of foci going towards the

membrane (M) to those being internalized (N). While this ratio was around 2 for both M9FL

and Δcat cells, this reduced to ~1 for RGE cells (Fig. 5J), suggesting that in spite of being

transported to the membrane, ITGβ1 is not stabilized at the cell membrane in RGE cells.

Collectively, these results highlight two distinct roles for the two ITGβ1 binding sites on MMP9,

with the 17 amino acid residue essential for ITGβ1 transport and the RGD domain essential

for stabilizing ITGβ1 at the cell membrane.

Discussion

MMP9 expression is associated with increased cancer cell invasion and metastasis [46]. While

the proteolytic activity of MMP9 involved in matrix remodeling is well established, an

increasing body of literature has demonstrated its non-proteolytic functions play an equally

crucial role. Here, we have established how the combination of proteolytic and non-

proteolytic functions of MMP9 together render cancer invasiveness via its association with

ITGβ1. Based on our findings, we propose a mechanism whereby MMP9/ITGβ1 association is

essential for their co-transport to the leading edge, ITGβ1 stabilization at the cell membrane,

MMP9 secretion and MMP9 mediated matrix degradation thereby creating migration paths

(Fig. 6).

The stiffness of the extracellular matrix has a significant impact on cell spreading, motility,

cytoskeletal organization, and MMP secretion [30], [47], [48], [49]. Among the secreted

MMPs, MMP2 and MMP9, increased cancer invasiveness was associated with a stiffness

dependent increase in MMP9 secretion, with MMP inhibition inducing cell rounding, and cell

and nuclear softening [50], [51]. By screening both normal and cancer cells of varying

invasiveness, here we show that stiffness modulates MMP9 expression transcriptionally in

A549, MDA-MB-231 and HT-1080 invasive cancer cells only. However, even in these invasive

cells, stiffness dependent modulation of MMP1, MMP2 and MMP14 was not observed. Our results suggest that MMP9 mediates faster motility on stiffer substrates by stabilising adhesions as both pharmacological inhibition of MMPs as well as MMP9 knockdown led to reduction in adhesion size and stability, and destabilization of the cell cytoskeleton [52], [53], [54].

Previous literature has shown that interaction between the MMP9 PEX (PEX9) domain and ITGβ1 is essential for formation of cell-matrix adhesions [26], [27], [55]. Building on this work, another study showed that a 17 amino acid stretch from the fourth blade of the PEX9 domain stabilizes MMP9/ITGβ1 association leading to loss of adhesions. Consistent with these studies, by deleting this stretch of amino acids, we show here that reduced spreading, migration, and invasion of Δβ1 cells is attributed to loss of focal adhesion formation. While the presence of fibronectin type II repeats in MMP9 has been documented before, their role has not been elucidated. For the first time, we identify the presence of an RGD motif in these repeats, which when mutated to RGE, leads to loss of adhesions. Thus, our study establishes that both integrin binding sites on MMP9 are essential for adhesion formation. This MMP9/ITGβ1 association can explain the restoration of the cell spreading, adhesion, and 2D motility in catalytic inactive MMP9 (Δcat) on 2D substrates. But the defects in cell invasion of Δcat cells inside 3D collagen illustrates the necessity of matrix degradation in 3D invasion.

Vesicular transport of MMP9 towards lamellipodia has been reported in HMEC-1 cell invasion previously [56]. Here we show that the 17 amino acid sequence mediates co-trafficking of MMP9 and ITGβ1 to the migration front. Co-trafficking of MMP9 and ITGβ1 ensures that MMPs are secreted at the vicinity of existing adhesions thereby leading to localized matrix degradation and adhesion strengthening. Defective MMP9 secretion in Δβ1 and Δβ1Δcat cells suggests that adhesion maturation and matrix remodeling are closely coupled, with MMPs being released only after integrins are stabilized on the membrane. Reduced MMP9 secretion observed in SB3-CT cells may be attributed to disruption of MMP9/ITGβ1 association. RGD peptide is widely used for integrin blocking experiments. The combination of AFM experiments and time-lapse imaging suggests that RGD domain of MMP9 FNII domain anchors integrins at the cell membrane.

Recent research has demonstrated that MDA-MB-231 cells secrete MMP9-enriched exosomes, which can diffuse and degrade the surrounding matrix [57]. Since exosome bio-

genesis involves the double invagination of the plasma membrane to form intraluminal bodies (ILB) inside a single multivesicular body (MVB) [58], it is likely that MMP9 is present both in the MVBs which merge with the cell membrane, and is also packaged in the exosomes which get secreted. While the MVB-associated MMPs stabilize ITGβ1 on the cell membrane via the RGD domain, upon matrix degradation mediated by exosomal MMPs, membrane-stabilized ITGβ1 forms firm adhesions with the underlying matrix. However, the details of this mechanism is yet to be established.

In conclusion, our data illustrates the importance of MMP9/ITGβ1 association in sustaining cancer invasion via ITGβ1/MMP9 co-trafficking and ITGβ1 stabilization at the cell membrane and subsequent MMP mediated matrix degradation.

Acknowledgements

We acknowledge IIT Bombay for providing Bio-AFM, FACS and Confocal Microscopy facilities.

Author Contributions

Conceptualization: S.D., S.Sen; Methodology: S.D., S.S.; Formal analysis: S.D., S.T., M.A.H; Investigation: S.D., S.Sen.; Data curation: S.D.; Writing - original draft: S.D., S.Sen; Writing - review & editing: S.Sen; Supervision: S.Sen; Project administration: S.Sen; Funding acquisition: S.Sen

Competing interests

The authors declare no competing or financial interests.

Funding

S. Sen acknowledges financial support from the Department of Science and Technology, Ministry of Science and Technology, India (Grant # DST/SJF/LSA-01/2016-17) and STARS, Ministry of Education (Grant # MoE-STARS/STARS-2/2023-0229). S.D. was supported by the fellowship from the Department of Biotechnology, Ministry of Science and Technology, India (Grant # DBT/2017/IIT-B/850).

References

[1] R. P. Mecham, 'Overview of extracellular matrix', *Curr Protoc Cell Biol*, no. SUPPL.57, 2012, doi: 10.1002/0471143030.cb1001s57.

[2] C. Frantz, K. M. Stewart, and V. M. Weaver, 'The extracellular matrix at a glance', *J Cell Sci*, vol. 123, no. 24, pp. 4195–4200, Dec. 2010, doi: 10.1242/jcs.023820.

[3] Richard O. Hynes, 'Extracellular matrix: not just pretty fibrils', *Science*, vol. 326, no. 5957. pp. 1212–1216, Nov. 27, 2009. doi: 10.1126/science.1176488.

[4] J. D. Humphrey, E. R. Dufresne, and M. A. Schwartz, 'Mechanotransduction and extracellular matrix homeostasis', *Nat Rev Mol Cell Biol*, vol. 15, no. 12, pp. 802–812, Dec. 2014, doi: 10.1038/nrm3896.

[5] K. R. Levental, 'Matrix Crosslinking Forces Tumor Progression by Enhancing Integrin signaling', *Cell*, vol. 139, 2010, doi: 10.1016/j.cell.2009.10.027.

[6] Q. Xiao and G. Ge, 'Lysyl Oxidase, Extracellular Matrix Remodeling and Cancer Metastasis', *Cancer Microenvironment*, vol. 5, no. 3, pp. 261–273, Dec. 2012, doi: 10.1007/s12307-012-0105-z.

[7] A. H. Mekhdjian *et al.*, 'Integrin-mediated traction force enhances paxillin molecular associations and adhesion dynamics that increase the invasiveness of tumor cells into a three-dimensional extracellular matrix', *Mol Biol Cell*, vol. 28, no. 11, pp. 1467–1488, Jun. 2017, doi: 10.1091/mbc.e16-09-0654.

[8] A. LOCHTER, M. D. STERNLICHT, Z. WERB, and M. J. BISSELL, 'The Significance of Matrix Metalloproteinases during Early Stages of Tumor Progression [a]', *Ann N Y Acad Sci*, vol. 857, no. 1, pp. 180–193, Oct. 1998, doi: 10.1111/j.1749-6632.1998.tb10116.x.

[9]   E. Kerkelä and U. Saarialho-Kere, 'Matrix metalloproteinases in tumor progression: focus on
basal and squamous cell skin cancer', *Exp Dermatol*, vol. 12, no. 2, pp. 109–125, Apr. 2003,
doi: 10.1034/j.1600-0625.2003.120201.x.

[10] O. R. F. Mook, W. M. Frederiks, and C. J. F. Van Noorden, 'The role of gelatinases in colorectal
cancer progression and metastasis', *Biochimica et Biophysica Acta (BBA) - Reviews on Cancer*,
vol. 1705, no. 2, pp. 69–89, Dec. 2004, doi: 10.1016/j.bbcan.2004.09.006.

[11] R. A. Wagenaar-Miller, L. Gorden, and L. M. Matrisian, 'Matrix metalloproteinases in
colorectal cancer: Is it worth talking about?', *Cancer and Metastasis Reviews*, vol. 23, no. 1/2,
pp. 119–135, Jan. 2004, doi: 10.1023/A:1025819214508.

[12] M. Björklund and E. Koivunen, 'Gelatinase-mediated migration and invasion of cancer cells',
Biochimica et Biophysica Acta (BBA) - Reviews on Cancer, vol. 1755, no. 1, pp. 37–69, May
2005, doi: 10.1016/j.bbcan.2005.03.001.

[13] S. Mondal, N. Adhikari, S. Banerjee, S. A. Amin, and T. Jha, 'Matrix metalloproteinase-9 (MMP-
9) and its inhibitors in cancer: A minireview', *Eur J Med Chem*, vol. 194, p. 112260, May 2020,
doi: 10.1016/j.ejmech.2020.112260.

[14] M. Egeblad and Z. Werb, 'New functions for the matrix metalloproteinases in cancer
progression', *Nat. Rev. Cancer*, vol. 2, 2002, doi: 10.1038/nrc745.

[15] A. Lochter, 'Matrix metalloproteinase stromelysin-1 triggers a cascade of molecular
alterations that leads to stable epithelial-to-mesenchymal conversion and a premalignant
phenotype in mammary epithelial cells', *J. Cell Biol.*, vol. 139, 1997.

[16] M. Ahonen, A. H. Baker, and V. M. Kahari, 'Adenovirus-mediated gene delivery of tissue
inhibitor of metalloproteinases-3 inhibits invasion and induces apoptosis in melanoma cells',
Cancer Res., vol. 58, 1998.

[17] N. Koshikawa, G. Giannelli, V. Cirulli, K. Miyazaki, and V. Quaranta, 'Role of cell surface
metalloprotease MT1-MMP in epithelial cell migration over laminin-5', *J. Cell Biol.*, vol. 148,
2000.

[18] G. Giannelli, J. Falk-Marzillier, O. Schiraldi, W. G. Stetler-Stevenson, and V. Quaranta,
'Induction of Cell Migration by Matrix Metalloprotease-2 Cleavage of Laminin-5', *Science
(1979)*, vol. 277, no. 5323, pp. 225–228, Jul. 1997, doi: 10.1126/science.277.5323.225.

[19] V. Noe, 'Release of an invasion promoter E-cadherin fragment by matrilysin and stromelysin-
1', *J. Cell Sci.*, vol. 114, 2001.

[20] C. Mehner, A. Hockla, E. Miller, S. Ran, D. C. Radisky, and E. S. Radisky, 'Tumor cell-produced
matrix metalloproteinase 9 (MMP-9) drives malignant progression and metastasis of basal-like
triple negative breast cancer', *Oncotarget*, vol. 5, no. 9, pp. 2736–2749, May 2014, doi:
10.18632/oncotarget.1932.

[21] S. Napoli *et al.*, 'Functional Roles of Matrix Metalloproteinases and Their Inhibitors in
Melanoma', *Cells*, vol. 9, no. 5, p. 1151, May 2020, doi: 10.3390/cells9051151.

[22] J. Hua and R. J. Muschel, 'Inhibition of matrix metalloproteinase 9 expression by a ribozyme
blocks metastasis in a rat sarcoma model system', *Cancer Res.*, vol. 56, 1996.

[23] M. Kajita, 'Membrane-type 1 matrix metalloproteinase cleaves CD44 and promotes cell
migration', *J. Cell Biol.*, vol. 153, 2001.

[24] H. Nakahara, 'Transmembrane/cytoplasmic domain-mediated membrane type 1-matrix
metalloprotease docking to invadopodia is required for cell invasion', *Proc. Natl Acad. Sci.*
USA, vol. 94, 1997.

[25] Q. Yu and I. Stamenkovic, 'Localization of matrix metalloproteinase 9 to the cell surface
provides a mechanism for CD44-mediated tumor invasion', *Genes Dev*, vol. 13, no. 1, pp. 35–
48, Jan. 1999, doi: 10.1101/gad.13.1.35.

[26] J. Redondo-Muñoz *et al.*, 'α4β1 integrin and 190-kDa CD44v constitute a cell surface docking
complex for gelatinase B/MMP-9 in chronic leukemic but not in normal B cells', *Blood*, vol.
112, no. 1, pp. 169–178, Jul. 2008, doi: 10.1182/blood-2007-08-109249.

[27] V. M. Alford *et al.*, 'Targeting the Hemopexin-like Domain of Latent Matrix Metalloproteinase-
9 (proMMP-9) with a Small Molecule Inhibitor Prevents the Formation of Focal Adhesion
Junctions', *ACS Chem Biol*, vol. 12, no. 11, pp. 2788–2803, 2017, doi:
10.1021/acschembio.7b00758.

[28] S. I. Fraley *et al.*, 'A distinctive role for focal adhesion proteins in three-dimensional cell
motility', *Nat Cell Biol*, vol. 12, no. 6, pp. 598–604, Jun. 2010, doi: 10.1038/ncb2062.

[29] E. Ugarte-Berzal *et al.*, 'A 17-residue Sequence from the Matrix Metalloproteinase-9 (MMP-9)
Hemopexin Domain Binds α4β1 Integrin and Inhibits MMP-9-induced Functions in Chronic
Lymphocytic Leukemia B Cells', *Journal of Biological Chemistry*, vol. 287, no. 33, pp. 27601–
27613, Aug. 2012, doi: 10.1074/jbc.M112.354670.

[30] A. Das, M. Monteiro, A. Barai, S. Kumar, and S. Sen, 'MMP proteolytic activity regulates cancer
invasiveness by modulating integrins', *Sci Rep*, 2017, doi: 10.1038/s41598-017-14340-w.

[31] A. Rossello *et al.*, 'New N-arylsulfonyl-N-alkoxyaminoacetohydroxamic acids as selective
inhibitors of gelatinase A (MMP-2)', *Bioorg Med Chem*, vol. 12, no. 9, pp. 2441–2450, May
2004, doi: 10.1016/j.bmc.2004.01.047.

[32] G. Tiscornia, O. Singer, and I. M. Verma, 'Production and purification of lentiviral vectors', *Nat*
Protoc, vol. 1, no. 1, pp. 241–245, Jun. 2006, doi: 10.1038/nprot.2006.37.

[33] T. D. Schmittgen and K. J. Livak, 'Analyzing real-time PCR data by the comparative CT method',
Nat Protoc, vol. 3, no. 6, pp. 1101–1108, Jun. 2008, doi: 10.1038/nprot.2008.73.

[34] B. Eltzner, C. Wollnik, C. Gottschlich, S. Huckemann, and F. Rehfeldt, 'The Filament Sensor for
Near Real-Time Detection of Cytoskeletal Fiber Structures', *PLoS One*, vol. 10, no. 5, p.
e0126346, May 2015, doi: 10.1371/journal.pone.0126346.

[35] M. E. Berginski and S. M. Gomez, 'The Focal Adhesion Analysis Server: a web tool for
analyzing focal adhesion dynamics', *F1000Res*, Mar. 2013, doi: 10.3410/f1000research.2-
68.v1.

[36] A. D. Doyle, M. L. Kutys, M. A. Conti, K. Matsumoto, R. S. Adelstein, and K. M. Yamada,
'Microenvironmental control of cell migration: Myosin IIA is required for efficient migration in
fibrillar environments through control of cell adhesion dynamics', *J Cell Sci*, Jan. 2012, doi:
10.1242/jcs.098806.

[37]   A. Barai, A. Mukherjee, A. Das, N. Saxena, and S. Sen, 'α-Actinin-4 drives invasiveness by
675   regulating myosin IIB expression and myosin IIA localization', *J Cell Sci*, vol. 134, no. 23, Dec.
676   2021, doi: 10.1242/jcs.258581.

677   [38]   R. Gorelik and A. Gautreau, 'Quantitative and unbiased analysis of directional persistence in
678   cell migration', *Nat Protoc*, vol. 9, no. 8, pp. 1931–1943, Aug. 2014, doi:
679   10.1038/nprot.2014.131.

680   [39]   P. Michaluk *et al.*, 'Influence of matrix metalloproteinase MMP-9 on dendritic spine
681   morphology', *J Cell Sci*, vol. 124, no. 19, pp. 3369–3380, Oct. 2011, doi: 10.1242/jcs.090852.

682   [40]   H. Mori, 'Transmembrane/cytoplasmic, rather than catalytic, domains of Mmp14 signal to
683   MAPK activation and mammary branching morphogenesis via binding to integrin β1',
684   *Development*, vol. 352, 2013, doi: 10.1242/dev.084236.

685   [41]   H. NAGASE, R. VISSE, and G. MURPHY, 'Structure and function of matrix metalloproteinases
686   and TIMPs', *Cardiovasc Res*, vol. 69, no. 3, pp. 562–573, Feb. 2006, doi:
687   10.1016/j.cardiores.2005.12.002.

688   [42]   C. Tallant, A. Marrero, and F. X. Gomis-Rüth, 'Matrix metalloproteinases: Fold and function of
689   their catalytic domains', *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*, vol.
690   1803, no. 1, pp. 20–28, Jan. 2010, doi: 10.1016/j.bbamcr.2009.04.003.

691   [43]   A. Jacob *et al.*, 'Rab40b regulates MMP2 and MMP9 trafficking during invadopodia formation
692   and breast cancer cell invasion', *J Cell Sci*, Jan. 2013, doi: 10.1242/jcs.126573.

693   [44]   C. M. Regen and A. F. Horwitz, 'Dynamics of beta 1 integrin-mediated adhesive contacts in
694   motile fibroblasts.', *J Cell Biol*, vol. 119, no. 5, pp. 1347–1359, Dec. 1992, doi:
695   10.1083/jcb.119.5.1347.

696   [45]   R. E. Bridgewater, J. C. Norman, and P. T. Caswell, 'Integrin trafficking at a glance', *J Cell Sci*, vol.
697   125, no. 16, pp. 3695–3701, Aug. 2012, doi: 10.1242/jcs.095810.

698   [46]   K. Nabeshima, T. Inoue, Y. Shimao, and T. Sameshima, 'Matrix metalloproteinases in tumor
699   invasion: Role for cell migration', *Pathol Int*, vol. 52, no. 4, pp. 255–264, Apr. 2002, doi:
700   10.1046/j.1440-1827.2002.01343.x.

701   [47]   A. Parekh and A. M. Weaver, 'Regulation of cancer invasiveness by the physical extracellular
702   matrix environment', *Cell Adh Migr*, vol. 3, no. 3, pp. 288–292, Jul. 2009, doi:
703   10.4161/cam.3.3.8888.

704   [48]   F. Kai, H. Laklai, and V. M. Weaver, 'Force Matters: Biomechanical Regulation of Cell Invasion
705   and Migration in Disease', *Trends Cell Biol*, vol. 26, no. 7, pp. 486–497, Jul. 2016, doi:
706   10.1016/j.tcb.2016.03.007.

707   [49]   T. A. Ulrich, E. M. de Juan Pardo, and S. Kumar, 'The Mechanical Rigidity of the Extracellular
708   Matrix Regulates the Structure, Motility, and Proliferation of Glioma Cells', *Cancer Res*, vol. 69,
709   no. 10, pp. 4167–4174, May 2009, doi: 10.1158/0008-5472.CAN-08-4859.

710   [50]   M. Ehrbar *et al.*, 'Elucidating the Role of Matrix Stiffness in 3D Cell Migration and
711   Remodeling', *Biophys J*, vol. 100, no. 2, pp. 284–293, Jan. 2011, doi:
712   10.1016/j.bpj.2010.11.082.

713   [51]   A. Das, A. Barai, M. Monteiro, S. Kumar, and S. Sen, 'Nuclear softening is essential for
714   protease-independent migration', *Matrix Biology*, vol. 82, pp. 4–19, Sep. 2019, doi:
715   10.1016/j.matbio.2019.01.001.

716   [52]   R. J. Pelham and Y. Wang, 'Cell locomotion and focal adhesions are regulated by
717   substrate flexibility', *Proceedings of the National Academy of Sciences*, vol. 94, no. 25, pp.
718   13661–13665, Dec. 1997, doi: 10.1073/pnas.94.25.13661.

719   [53]   D. J. Webb, J. T. Parsons, and A. F. Horwitz, 'Adhesion assembly, disassembly and turnover in
720   migrating cells – over and over and over again', *Nat Cell Biol*, vol. 4, no. 4, pp. E97–E100, Apr.
721   2002, doi: 10.1038/ncb0402-e97.

722   [54]   D. Kim and D. Wirtz, 'Focal adhesion size uniquely predicts cell migration', *The FASEB Journal*,
723   vol. 27, no. 4, pp. 1351–1361, Apr. 2013, doi: 10.1096/fj.12-220160.

724   [55]   J. Redondo-Muñoz *et al.*, 'Matrix Metalloproteinase-9 Promotes Chronic Lymphocytic
725   Leukemia B Cell Survival through Its Hemopexin Domain', *Cancer Cell*, vol. 17, no. 2, pp. 160–
726   172, Feb. 2010, doi: 10.1016/j.ccr.2009.12.044.

727   [56]   I. Abécassis, B. Olofsson, M. Schmid, G. Zalcman, and A. Karniguian, 'RhoA induces MMP-9
728   expression at CD44 lamellipodial focal complexes and promotes HMEC-1 cell invasion', *Exp
729   Cell Res*, vol. 291, no. 2, pp. 363–376, Dec. 2003, doi: 10.1016/j.yexcr.2003.08.006.

730   [57]   S. Patwardhan, P. Mahadik, O. Shetty, and S. Sen, 'ECM stiffness-tuned exosomes drive breast
731   cancer motility through thrombospondin-1', *Biomaterials*, vol. 279, p. 121185, Dec. 2021, doi:
732   10.1016/j.biomaterials.2021.121185.

733   [58]   R. Kalluri and V. S. LeBleu, 'The biology **,** function **,** and biomedical applications of exosomes',
734   *Science (1979)*, vol. 367, no. 6478, Feb. 2020, doi: 10.1126/science.aau6977.

Figure Legends

Figure 1: ECM stiffness differentially modulates MMP expression and activity in cancer cells.

(A) Curing TCGA patient datasets and processing to assess the expression profile of major

MMPs (MMP1, 2, 9, and 14) using UCSC Xena browser. (B) The mRNA expression profiles of

MMP1, 2, 9, and 14 in ~110 normal tissue samples, ~1100 tissues from primary malignant

sites, and ~8 tissues from secondary malignant sites. (C, D) The expression profiles of MMP1,

2, 9, and 14 in ~115 normal and ~1100 malignant tissue samples in breast cancer patients

and lung cancer patients. (E) Schematic for assessing stiffness dependent MMP expression

and secretion. Cells were grown on 0.5 kPa soft and 5 kPa stiff polyacrylamide gels. (F) Relative

fold change of gene expression of MMP1, 2, 9, and 14 by quantitative PCR in human dermal

fibroblasts (HDFs), MCF10A normal mammary epithelial cells, MCF7 ERPR+ breast cancer cells,

A549 lung adhenocarcinoma cells, MDA-MB-231 (MDA) triple negative breast cancer cells,

and HT1080 (HT) fibrosarcoma cells on both 0.5 and 5 kPa PA-gels. The fold change was

calculated for the expression of HDFs on 0.5 kPa gels. (G) Western blot analysis of protein

expression of MMP2, 9, and ITGβ1 in the cells mentioned above on 0.5 and 5 kPa PA-gels.

GAPDH was used as a loading control. (H) The fold change of the gene expression of A549,

MDA-MB-231, and HT1080 cells on 5 kPa PA gels normalized with GAPDH expression is

represented as a bar diagram analyzed from three independent experiments ($mean \pm SEM$).

(I) Gelatin zymogram of cell-secreted conditioned media (CM) from soft and stiff substrates to

probe the secretion profiles of MMP9 and MMP2. (J) The relative fold change of the secreted

MMP2/9 of A549, MDA-MB-231, and HT1080 on 5 kPa PA gels is represented as the bar

diagram normalized to on 0.5 kPa PA gels. The analysis was performed three independent

times and densitometric analysis of the band intensity was calculated for statistical

comparisons ($mean \pm SEM$). For all sub-figures, statistical significance was assessed using

One-Way ANOVA or nonparametric test. ($* p < 0.05, ns: not\ significant$).

Figure 2: Soluble MMPs drive cell spreading and migration by regulating focal adhesions. (A, B) Representative Lectin (green)/F-actin (red)/DAPI (blue) stained images of untreated and drug treated HT1080 cells on 5 kPa PA gels and quantification of cell spreading area of untreated, control (DMSO) and drug (GM6001 and SB3-CT) treated A549, MDA-MB-231 and

HT1080 cells ($\approx 80 - 100$ cells, $N = 3$). (C, D) Representative migration trajectories of control

and drug treated HT1080 cells on 5 kPa PA gels, and quantification of cell speed ($\approx 50 - 80$

cells per condition, $N = 3$). (E, F) Representative invasion trajectories of control and drug

treated HT1080 cells entrapped inside 3D collagen gels and quantification of cell speed ($\approx$

$30 - 50$ cells per condition, $N = 3$). (G, H) Representative force vs. indentation curves of

control and drug treated HT1080 cells on 5 kPa PA gel. The data were fitted to obtain estimates

of cell cortical stiffness ($\approx 40 - 50$ cells per condition, $N = 3$). (I, J) Representative images of

HT1080 cells/Collagen I gel mixture cultured for 48 hr in the presence of DMSO/GM6001/SB3-

CT and quantification of gel compaction ($\approx 3 - 4$ gels per condition, $N = 3$). White/yellow

dotted circles represent the initial/final gel areas. (K-M) Representative images of F-actin

stress fibre (green) and paxillin (red) stained HT1080 cells treated with DMSO, GM6001 and

SB3-CT on 5 kPa PA gels, and quantification of stress fibers and focal adhesion size ($\approx 30 - 50$

cells per condition, $N = 3$). The yellow lines represent processed stress fibres. (N, O)

Representative lifetime heatmaps and quantification of focal adhesion lifetime of HT1080 cells

treated with DMSO and SB3-CT ($\approx 7 - 10$ cells per condition, $N = 3$). (P) The gelatin

zymogram of conditioned media (CM) and western blot of ITGβ1 in DMSO, GM6001 and SB3-

CT treated HT1080 cells. Densitometric analysis of the bands were performed ($N = 3$). GAPDH

was used as a loading control. For all sub-figures, statistical significance was assessed using

One-Way ANOVA or nonparametric test. ($* p < 0.05, ** p < 0.01, *** p < 0.001, **** p <$

$0.0001, ns$: not significant.

Figure 3: Both proteolytic and non-proteolytic functions of MMP9 are essential for sustaining cancer invasion. (A) Schematic of generation of MMP9 knockdown cells (shKD1/2) and M9FL/Δcat expressing cell lines using lentiviral mediated shRNA transduction. (B) Gelatin zymogram of CM and western blotting of CM/ whole cell lysate from shCTL, shKD1/2, M9FL1/2 and Δcat cells. (C, D) Representative lectin (green), phalloidin (red) and DAPI (blue) stained images of HT1080 shCTL, shKD1/2, M9FL1/2 and Δcat images of cells on stiff 5 kPa PA gels, and quantification of cell spread area and circularity ($\approx 50 - 80$ cells per condition, $N = 3$).

(E, F) Representative 2D migration trajectories of HT1080 shCTL, shKD, M9FL and Δcat cells on

5 kPa PA gels and quantification of cell motility ($\approx 50 - 80$ cells per condition, $N = 3$). (G, H)

3D invasion trajectories of HT1080 shCTL, shKD, M9FL and Δcat cells inside 3D Col and

quantification of cell speed ($\approx 40 - 50$ cells per condition, $N = 3$). (I, J) Representative

images of F-actin (green) of HT1080 shCTL, shKD, M9FL and Δcat cells, and quantification of

stress fibers ($\approx 50 - 60$ cells per condition, $N = 3$). Yellow lines indicate individual stress

fibres. (K, L) Representative force-indentation of M9FL, shKD and Δcat cells and quantification

of the cortical stiffness of these cells ($\approx 50 - 80$ cells per condition, $N = 3$). (M, N)

Representative images of 3D gel compaction of M9FL, shKD and Δcat cells and quantification

of gel compaction ($\approx 3 - 4$ gels per condition from $N = 3$). White/yellow dotted circles

represent the initial/final gel areas. (O, P) Representative paxillin (red) and F-actin (green)

stained shCTL, shKD, M9FL and Δcat cells and quantification of the number of focal adhesions

per cell ($\approx 30 - 50$ cells per condition, $N = 3$). (Q, R) Representative Focal adhesion lifetime

heatmaps and quantification of adhesion lifetime in M9FL, shKD and Δcat cells ($\approx 15 - 20$

cells per condition, $N = 3$). For all sub-figures, statistical significance was assessed using One-

Way ANOVA or nonparametric test ($* p < 0.05, ** p < 0.01, *** p < 0.001, **** p <$

$0.0001, ns$: not significant).

Figure 4: MMP9- ITGβ1 interaction via two distinct sites is essential for sustaining cancer

invasion. (A) Domain organization of MMP9 showing two putative integrin bindings sites, the

RGD sequence in the catalytic domain and a 17 aa sequence in blade IV of the hemopexin

domain. To assess the importance of these two sites, two mutants RGE and Δβ1 were

generated. (B) Proposed proteolytic and nonproteolytic functions of different mutated MMP9 constructs, where green/red circles represent active/inactive functions. (C) Co-immunoprecipitation for the verification of the perturbation of association between different MMP9 constructs (M9FL, Δcat, RGE, Δβ1, and Δβ1Δcat) and ITGβ1. (D) The working principle of proximity ligation assay (PLA) using Duolink Probe. (E, F) Spatial localization of MMP9-ITGβ1 association determined by PLA (green) and MMP9 (red) signal in M9FL, Δcat, RGE, Δβ1, and Δβ1Δcat cells and quantification of PLA integrated density ($\approx 60 - 80$ cells per condition, $N = 3$). IgG control and only secondary antibody-treated conditions served as negative controls. (G) MMP9 secretion and expression in M9FL, Δcat, RGE, Δβ1, and Δβ1Δcat cells assessed using gelatin zymography and western blotting of CM/whole cell lysate. GAPDH served as loading control. (H, I) Representative lectin (red)/F-actin (green)/DAPI (blue) stained images of M9FL, Δcat, RGE, Δβ1, and Δβ1Δcat cells and quantification of cell spread area ($\approx 70 - 80$ cells per condition, $N = 3$). (J-M) Representative 2D migration and 3D invasion trajectories of M9FL, Δcat, RGE, Δβ1, and Δβ1Δcat cells on 5 kPa PA gel and 3D collagen respectively and quantification of cell speed ($\approx 60 - 80$ cells per condition, $N = 3$). For all sub-figures, statistical significance was assessed using One-Way ANOVA or nonparametric test ($* p < 0.05, ** p < 0.01, *** p < 0.001, **** p < 0.0001$).

D) Focal adhesion lifetime heatmaps of M9FL, Δcat, RGE, Δβ1, and Δβ1Δcat cells and its

quantification ($\approx 12 - 15$ cells per condition, $N = 3$). (E, F) Representative AFM retraction

curves of M9FL, Δcat, RGE, Δβ1, and Δβ1Δcat cells obtained by indenting cells with RGD-

coated spherical tips, and retracting after 10s of dwell time. The maximum adhesion force was

quantified for the different cell lines ($\approx 50 - 80$ cells per condition, $N = 3$). (G)

Representative images of ITGβ1 (red), MMP9 (green), and DAPI (blue) stained M9FL, RGE and

Δβ1 cells, with intensity along the Line profiles drawn in the direction of cell migration

(magenta and yellow circles depict the migration front and cell rear respectively). (H)

Trajectories of ITGβ1 (red) foci in M9FL, RGE and Δβ1 cells along the direction of migration

(white arrow). Merged images of MMP9 (green) and ITGβ1 foci (red) are also shown. Time

stamp images of MMP9 (green), ITGβ1 foci (red) and bio-tracker (blue) of the indicated dotted

box (inset) are represented. (I) Quantification of trafficking speed of ITGβ1 foci in the different

cell lines ($\approx 20 - 30$ taken from $5 - 7$ cells per condition, $N = 3$). (J) Quantification of the

ratio of the number of the ITGβ1 foci going towards the membrane (M) to the number of foci

being internalized (N) at a given time ($\approx 15 - 30$ taken from $5 - 6$ cells per condition, $N = 3$).

3). For all sub-figures, statistical significance was assessed using One-Way ANOVA or

nonparametric test ($* p < 0.05, ** p < 0.01, *** p < 0.001, **** p < 0.0001$).

Figure 6: Proposed model of MMP9 mediated ITGβ1 co-trafficking and stabilization on the

cell membrane. Both MMP9 and ITGβ1 are co-trafficked via intracellular vesicles to the cell

membrane. While hemopexin domain of MMP9 is responsible for co-packaging of

MMP9/ITGβ1 in vesicles and

their co-transport, RGD domain mediates membrane surface stabilization of ITGβ1. Upon

MMP9 mediated matrix degradation, focal adhesions are stabilized leading to forward

migration.