Decoding Missense Variants by Incorporating Phase Separation via Machine Learning

Mofan Feng[1,2+], Xiaoxi Wei[1+], Xi Zheng[1,2], Liangjie Liu[1,2], Lin Lin[1], Manying Xia[1], Guang He[1,2*], Yi Shi[1,2*], Qing Lu[3*]

1 Bio-X Institutes, Key Laboratory for the Genetics of Developmental and Neuropsychiatric Disorders, Shanghai Jiao Tong University, Shanghai, China.

2 The Collaborative Innovation Center for Brain Science, and Brain Science and Technology Research Center, Shanghai Jiao Tong University, Shanghai, China.

3 Department of Otorhinolaryngology-Head and Neck Surgery, Chongqing General Hospital, Chongqing, China. Bio-X Institutes, Key Laboratory for the Genetics of Developmental and Neuropsychiatric Disorders, Shanghai Jiao Tong University, Shanghai, China. Ear Institute, Shanghai Jiao Tong University School of Medicine, Shanghai, China. Shanghai Key Laboratory of Translational Medicine on Ear and Nose Diseases, Shanghai, China.

+ These authors contributed equally as co-first authors.

* Corresponding author at: Bio-X Institutes, Shanghai Jiao Tong University,

1954 Huashan Road, Shanghai 200030, P.R.China. Tel & Fax: +86 (0) 21

62822491 Email:

Guang He: heguang@sjtu.edu.cn,

Yi Shi: yishi@sjtu.edu.cn

Qing Lu: luqing67@sjtu.edu.cn

Abstract

Computational models have made significant progress in predicting the effect of protein variants. However, deciphering numerous variants of unknown significance (VUS) located within intrinsically disordered regions (IDRs) remains challenging. To address this issue, we introduced phase separation (PS), which is tightly linked to IDRs, into the investigation of missense variants. Phase separation is vital for multiple physiological processes. By leveraging missense variants that alter phase separation propensity, we developed a machine learning approach named PSMutPred to predict the impact of missense mutations on phase separation. PSMutPred demonstrated robust performance in predicting missense variants that affect natural phase separation. In vitro experimental findings further underscore its validity. By applying PSMutPred on over 522,000 ClinVar missense variants, it significantly contributes to decoding the pathogenesis of disease variants, especially those in IDRs. Our work provides unique insights into the understanding of a vast number of VUSs in IDRs, thereby expediting clinical interpretation and diagnosis of disease variants.

Keywords: phase separation, missense mutation, variant effect prediction, machine learning, intrinsically disordered regions

Introduction

Approximately 25% of disease-associated missense variants are located in intrinsically disordered regions (IDRs)[1], present in approximately 63% of human proteins. However, studying variants in IDRs is challenging due to the lack of a fixed tertiary structure and limited evolutionary conservation[2], as only a small proportion (around 15%) of IDRs exhibit high conservation and high pLDDT scores (AlphaFold2[3]), which denote prediction confidence[4]. Numerous IDR variants are therefore annotated as variants of unknown significance (VUSs), making it difficult to evaluate and predict

their functional impact on diseases.

IDRs, particularly low-complexity IDRs, are crucial for the formation of membrane-

less biomolecular condensates through phase separation, a complex and not fully

understood physicochemical process in which molecules segregate into a distinct fluid

phase[5,6]. Phase separation leads to the formation of membrane-less organelles called

condensates which play essential roles in various biological processes. An increasing

number of proteins are recognized for their roles via the formation of phase separation

condensates. These condensates include the nucleolus and transcription factories in the

nucleus[7], stress granules[8], and protein densities at neuronal synapses[9] and inner-ear hair

cells[10-12].

Accumulating studies in IDRs reveal that missense variants in IDRs can perturb

protein functions by altering phase separation[13-15]. Missense mutation impacts phase

separation by altering specific residues, such as key residues for polar interaction[16], pi-

contact[17,18] or other multivalent interaction[19-21]. These mutations can affect the IDR

conformations, intra-molecular interactions, and inter-molecular protein-protein

interactions, leading to abnormal condensate formations[7]. This can cause either loss[22,23]

or gain[13,14,24] of natural phase separation, leading to misplacement and related gain or

loss-of-function outcomes. Notably, the gain of unnecessary phase separation can

contribute to disease phenotypes like neurodegenerative disorders such as Alzheimer's

and Parkinson's diseases [25-27]. These pioneering works have confirmed the association

between phase separation and disease variants in several cases.

Existing missense variant prediction algorithms typically rely on protein structural

features or evolutionary features derived from multiple sequence alignments (MSA).

Using such features, prediction algorithms of disease variants have made significant

progress in predicting protein clinical relevance[28-32], bridging the variant interpretation

gap left by experiments due to the cost and labor constraints. However, for numerous

disease variants located in IDRs which often lack a fixed structure[7] and show poor

evolutionary conservation[2], encoding variants using these traditional features becomes

unsuitable, making accurate prediction challenging. To address this challenge, we

propose to employ phase separation as a promising feature for improving the prediction

of IDR disease variants, given the role of numerous IDRs in phase separation associated

with various diseases.

Machine learning algorithms have shown remarkable performance in predicting the

propensity of proteins to undergo phase separation. Several sequence-related features

have been identified that made phase separation prediction effective[33-39] including pi-

pi[35], cation-pi interactions[36,37,40], electrostatic interactions[39,41], hydrophobic

interactions[38,40], and the valency and patterning of the low-complexity region[8,36,42].

Protein-protein interactions (PPIs) and post-translational modifications (PTMs) were

also found to be promising features recently[43-45]. Missense mutations can cause

pathogenic changes by altering phase separation. However, current sequence-based

phase separation prediction algorithms are trained on a limited set of known phase-

separating proteins, their ability to predict the impact of missense mutations on phase

separation remains unclear and has not been systematically evaluated.

To enhance comprehension of the correlation between IDR missense mutations and

diseases, we approached the issue from the phase separation standpoint and devised

features to represent the underlying phase separation alterations resulting from the

mutation. Using experimentally validated missense mutations that alter phase

separation propensity in proteins naturally undergoing this process, we constructed a

computational tool named PSMutPred to predict the effect of missense mutation on

phase separation. We demonstrated that missense mutations that impact the normal

phase separation propensity can be predicted, and the direction of the shifts in the PS

threshold caused by these mutations are also predictable.

We next explored whether PS can be integrated into the pathogenicity prediction of

disease variants. By analyzing PSMutPred scores for 520,000 + missense variants, we

observed a positive correlation between the variant's tendency to impact phase

separation and its pathogenicity, especially in proteins prone to phase separation. We

also observed that in proteins related to neurodegenerative diseases, disease variants

that were predicted to enhance phase separation were proportionally more prevalent

compared to those that might weaken phase separation. By integrating PSMutPred

scores and other PS-related features into representative unsupervised pathogenicity

prediction methods for missense variants, such as EVE[31] and ESM1b[46], we observed

significant improvement in prediction accuracy, especially for variants within low-

conservation IDRs, with an approximately 10% increase in AUPR. These findings

demonstrate that PSMutPred not only can serve as a tool for predicting the impact of

mutations on phase separation, but also can provide informative encoding for mutation

impacts on IDRs. Additionally, phase separation offers fresh perspectives and

opportunities for pre-screening studies on the pathogenicity of numerous VUSs.

Results

Collection of Phase Separation (PS)-related missense variants

To investigate relationships between missense mutations and protein phase

separation (PS) properties, we reviewed missense variants with altered PS propensity

that were documented in PhaSepDB[47] and LLPSDB[48,49] databases. To minimize the

noise effect caused by multiple mutations within a single sequence, we narrowed our

selection to mutation records with a limited number of mutations in the sequence that

alter the normal PS threshold (annotated as 'Impact' mutations). Examples include

P22L in *Ape1*, which solidifies semi-liquid *Ape1* Droplets[50], and S48E in *TDP-43,*

which disrupts PS[51]. We limited our analysis to variants influencing proteins'

spontaneous PS, excluding partner-dependent PS.

Our compilation yielded a list of 307 experimentally validated 'Impact' mutation

records from 70 proteins (Fig. S1A), including 79 that strengthened the PS properties

(annotated as 'Strengthen') and 228 that weakened or disabled them (annotated as

'Weaken/Disable'). The PScore[35] and PhaSePred[43] scores indicate that the proteins

from which these missense variants originate are predominantly proteins undergoing

PS spontaneously, as the PS propensity scores predicted for these proteins are

significantly higher compared to those of the human proteome (Fig. 2A).

Properties of variants impacting phase separation

We observed that mutations impacting PS (annotated as 'Impact' mutations) are

predominantly located in intrinsically disordered regions (IDRs) rather than structured

domain areas (Domains) (Fig. 2B). Existing variant effect prediction models primarily

rely on evolutionary features[28,30,31,52] and structural features[32,53-55]. However, these
features are inadequate for representing variants in IDRs and their effects on PS. We
found that current advanced pathogenicity prediction models, such as AlphaMissense[32]
and EVE[31], exhibit higher uncertainty of predictions for variants in IDRs than in
Domains (Fig. S1D). Moreover, they both fail to distinguish 'Impact' mutations from
random background mutations (Fig. S1C). This underscores the urgent need for
developing specialized models to predict the effects of mutations on PS.

Among these 'Impact' mutations, those involving serine, tyrosine, arginine, lysine
and glutamine are most prevalent (Fig. 2C, Fig. S5A). This aligns with prior findings
that tyrosine, lysine and arginine govern the saturation concentration of PS, while
glutamine and serine promote it[37,56-58].

Interestingly, our analysis revealed that 'Impact' mutations tend to occur near domain
boundaries. Specifically, the amino acid distance from each mutation site to its closest
domain boundary was calculated. We noted that, whether within Domains or IDRs,
'Impact' mutations are significantly closer to these boundaries compared to random
mutation sites (P <= 0.0001, Fig. 2D and Fig. S5B). Furthermore, we observed that
pathogenic mutants predicted by AlphaMissense[32] also demonstrated shorter distances
to domain boundaries compared to those predicted as benign in both Domains and IDRs
(P<=0.0001, Fig. S1E).

In addition, we assessed pi-contact values at mutation sites using the prediction
function provided by PScore[35], as pi-pi interactions are vital in determining PS. For 6
out of the 8 predicted pi-contact values, 'Impact' mutations have higher predicted
values than random mutation sites (Fig. 2E). This suggests that mutations affecting PS
are likely to occur at sites with high pi-contact frequencies.

Next, we compared the changes in amino acid properties before and after mutation
between the 'Strengthen' and 'Weaken/Disable' groups. We observed that missense
mutations that strengthen PS propensity are prone to have higher amino acid mass (P =
0.0097), hydrophobicity (P = 0.0024), and decreased polarity (P < 0.0001) compared to
'Weaken/Disable' mutations (Fig. 2F and Fig. S1F). Notably, the increase in
hydrophobicity supports the previous discovery that hydrophobicity serves as an

important PS driver [38,40].

Together, these findings show the distinct properties of missense mutations impacting

PS and the varying characteristics between mutations that either strengthen or weaken

PS. Based on these properties, we have developed tools to predict the impact of

mutations on PS, which will be discussed in the following section.

PSMutPred for predicting the impact of missense mutation on phase separation tendency

To predict the impact of missense mutations on protein PS properties, we developed

PSMutPred composed of two machine learning (ML) approaches. The first approach

termed the 'Impact Prediction' (IP) task trained ML models to predict missense

mutations that impact PS. In the second approach termed the 'Strengthen/Weaken

Prediction' (SP) task, ML models were trained to predict the direction of the shifts in

the normal cellular PS threshold induced by 'Impact' mutations (Methods). The process

is depicted in Figure 1, indicated by the green section (Fig. 1).

To encode such changes caused by missense mutations for quantification and model

learning, we considered the physicochemical properties of both the wild-type amino

acid (AA) and mutant AA at each mutation site, as well as the properties of amino acids

within IDRs. We applied pi-pi contact frequency[35] to encode the mutation site's

underlying significance for PS. Additionally, features such as IUPred[59] score and

mutation distance to domain boundary were used to encode the position of these

mutations. Each mutation sample was converted into a 39-dimension feature vector

(Methods). Neither MSA features nor structural features widely adopted in existing

mutation-related machine learning prediction[28-31,52,53,60] were considered due to the

nature of IDRs.

Due to the limited availability of variants that have no PS effect, for each of the 70

proteins, we randomly generated 500 single amino acid (AA) variants, resulting in a

total of 35,000 random 'Background' mutations, which were further used as negative

samples (i.e., 'Background' mutations) for analyses (Methods). The dataset of missense

mutations was divided into a cross-validation dataset (246 'Impact' samples and 23,500

'Background' samples from 47 proteins) and an independent test set (61 'Impact'

samples and 11,500 'Background' samples from 23 proteins) (Methods). We trained

prediction models including Logistic Regression (LR), Random Forest (RF), and

Support Vector Regression (SVR) for both tasks to explore the discriminative power of

predicting missense mutations' effect on PS.

Performance evaluation of PSMutPred

Although there is currently no computational algorithm for missense mutations on

predicting the effect on PS, we attempted to determine if existing PS prediction methods

are capable of discerning alterations in PS propensity caused by missense mutations.

Five high-performing PS methods were selected, including DeePhase[39], PSAP[61],

PScore[35], catGRANULE[38], and FuzDrop[62]. We found that the prediction score

differences of 'Impact' mutations were higher than those of 'Background' mutations,

by comparing the absolute differences of prediction scores (Fig. 3A). However, the

AUROCs were unsatisfactory (Fig 3B). Except for FuzDrop[62], none of the methods

could differentiate between 'Strengthen' mutations and 'Weaken' mutations (Fig. S2 C

and D), as their prediction scores did not accurately reflect the increase or decrease of

PS propensity caused by mutations.

To test whether PSMutPred-IP can identify mutations that impact PS, we

implemented leave-one-source-out cross-validation (LOSO CV) (Method). LOSO is

used to evaluate the models' predictive capabilities for variants from unseen proteins,

ensuring their generalizability. AUROCs and AUPRs on LOSO (Fig. 3 C and D)

showed the accuracy of our models, especially IP-SVR and IP-RF. Evaluation results

on the independent test set showed that our models have stable prediction performance

when facing mutations from different PS protein categories (Fig. 3 C and D).

To ensure that the superior performance of the algorithm was not caused by the

distribution bias of 'Impact' mutations, which tend to be located in IDRs (Fig. 2 B and

D), we conducted an additional LOSO CV and an independent validation for the IP task.

Here we generated 'Background' samples maintaining the same IDRs: Domains ratio

as observed in the 'Impact' samples. LOSO CV results and independent validation

results (Fig. 3 E and F) were as promising as those from the previous dataset (Fig. 3 C

and D). We also created another 'Background' dataset by aligning the amino acid

substitution frequencies with those in the impact dataset. This was done to test whether

the predictive power was due to the high ratio of specific amino acid types in 'Impact'

samples (Fig. 3 G and H).

Moreover, a small proportion of 'Impact' mutations are multi-point mutations (Fig.

S1B), the proposed models still exhibited efficient predictive power and outperformed

representative PS models when analyzing the results without considering multi-point

mutations (Fig. S2 A and B). We also assess the performance of PSMutPred-IP by only

including naturally occurring mutations in the database, the model allowed efficient

predictive power on these mutations (Fig. S5 D and E).

We next investigated the performance of PSMutPred-SP models. The significant

discriminative power to distinguish between 'Strengthen' samples and 'Weaken'

mutation samples from unseen proteins under LOSO CV (SP-LR, $P < 0.0001$) (Fig. S2

E and F) and AUROCs on the independent test set (Fig. S2G) indicated that SP-LR, SP-

RF models can identify the direction of the shifts in PS caused by missense mutations.

Feature importance indices were calculated for random forest models of both tasks

and grouped into feature types to discover potentially key features (Fig. S2 G and H).

In both tasks, the pi-contact frequency of the mutation site ranked first, suggesting that

the pi-pi interaction at the mutation's specific location significantly influences the effect

of mutations on protein phase separation. Beyond Pi-contact, no single feature was

identified as a dominating one, suggesting the non-linear nature among features and a

multi-factor causal relationship between features and the PS outcome.

Overall, PSMutPred models not only predict missense mutations that impact PS but

also predict the direction of the PS-threshold shift. Our final model generates

predictions made by the IP-SVR, IP-RF, IP-LR, SP-LR, and SP-RF models as well as

their corresponding rank scores (Methods). These models can be employed as effective

tools for assessing the tendency of missense mutations to affect PS, enhancing the

interpretation of disease variants' pathogenicity.

Experimental validation on the novel PS-related mutations identified by PSMutPred

Aberrant phase separation (PS) or aggregation processes have been implicated in the pathogenesis of various diseases, including neurodegenerative disorders, autism[7], and hearing loss[10-12]. In this study, we employed PSMutPred to predict the PS impact ('Impact' prediction and 'Strengthen/Weaken' prediction) of missense variants from genes associated with PS-related diseases. Specifically, we selected *EPS8*, known for its association with deafness, and analyzed the PS impact of its 'Uncertain' missense mutations from the ClinVar[63,64] database to assess the accuracy of our prediction model.

Epidermal growth factor receptor pathway substrate 8 (EPS8) is a multifunctional protein involved in cell mitosis and differentiation[65-68], in capping proteins through side-binding, in bundling of actin filaments[69-71] as well as in the elongation of actin in hair cell stereocilia[65]. Prior research has shown that EPS8 localizes to the tips of stereocilia and contributes to the formation of PS-mediated condensates at the stereocilia tip complex[11,72]. These findings suggest that EPS8 has the capacity for self-phase separation and to interact with other molecules.

To validate our predictions, we selected missense mutations, including R265C, D586G, and K676R, from the 8 candidate mutations. These candidates were predicted by PSMutPred-IP to impact PS, either by strengthening or weakening it, with rank scores above 0.5 across all models (IP-SVR, IP-RF, and IP-LR). Among these candidates, D586G had the highest scores for both SP-LR and SP-RF, suggesting a stronger propensity to enhance PS, while K676R scored the lowest for both metrics implying an impairment of PS capacity. As negative controls, we selected R702W and E728V from the 15 candidate mutations (all with IP rank scores below 0.5 for IP-LR, IP-SVR, and IP-RF; at least one had a score below 0.1).

For experimental validation, we overexpressed the wild-type and mutants of mouse *Eps8* (which is highly conserved with human *EPS8*, as shown in Fig. S3D) fused with a GFP tag in HEK293 cells. Observations made using Olympus fluorescence microscopy highlighted distinct changes in puncta formation quantity to evaluate PS

capacity. Specifically, R265C and D585G exhibited a notable increase in the number of

puncta, while K675R showed a significant reduction (Fig. 4A and B, Fig. S3 A and B).

Notably, D586G (equivalent to D585G in mice) demonstrated enhanced PS capacity,

aligning with its high SP-RF and SP-LR scores. In contrast, K676R (equivalent to

K675R in mice) showed diminished PS ability, supported by its low SP-RF and SP-LR

scores. To confirm that the observed puncta are indeed a result of liquid-liquid phase

separation (LLPS), we also conducted a fluorescence recovery experiment after a

photobleaching (FRAP) experiment to validate the dynamic and rapid formation of

droplets observed of both wild-type and mutant (D585G) of mouse EPS8 in HEK293

cells (Fig. S4 H-J).

Eps8 comprises six domains, encompassing both well-structured regions and

intrinsically disordered regions (IDRs) (Fig. S3C). D585 in mice (equivalent to D586

in humans) is situated in the SH3 domain. It forms stable hydrogen bonds with the

sidechains or backbones of K536 and K538 in mice (equivalent to K537 and K539 in

humans), maintaining structural stability through either a β-sheet-loop conformation in

mice or β-sheet-β-sheet conformation in humans (Fig. 4 D and E, Fig. S3 E and F).

Additionally, D585 and its neighboring residues, as well as interaction residues, exhibit

high conservation. However, the substitution of glycine for aspartate disrupts the

formation of these hydrogen bonds, leading to a destabilized structure. This change

might enhance PS by increasing flexibility and modifying the β-sheet-loop structure.

Conversely, K675 in mice (equivalent to K676 in humans) is located in an IDR at the

C-terminus of EPS8 (Fig. 4C, Fig. S3D). Sequence alignment underscores the high

conservation of K675 and its adjacent residues (Fig. 4F, Fig. S3G). Lysine's

characteristic long and hydrophobic side chain increases the likelihood of extending

and 'capturing' polar residues from self-proteins or other proteins. This substitution

may alter these interactions, potentially impairing PS.

In summary, our study demonstrates that the changes in PS propensity, as predicted

by our model, accurately correspond to a distinct number of puncta formed in live cells.

This serves to underscore the utility and accuracy of PSMutPred.

Phase separation effect prediction of disease variants by PSMutPred

Phase separation turns out to be a general mechanism for protein condensate assembly forming as membrane-less organelles in various physiological processes[9,73-82]. Mutations that change phase separation (PS) are likely causes of disease[13,22,23,83]. To exam this theory, we divided missense variants corresponding to 8,611 human genes from ClinVar[63,64] (522,016 variants in total, downloaded in 2022.12) into PS-prone and low-PS-prone propensity groups for between-group analyses (Methods).

By comparing the PS-prone group defined by 83 PS ClinVar proteins with the low-PS-prone group (other proteins), we found that in the PS-prone group, the PSMutPred-IP scores of variants (n = 1,451) were skewed to pathogenicity compared to those from low-PS-prone group (n = 86,291) (Fig. 5A). To make the sizes of the two groups more comparable and improve the efficiency of the comparison, we also defined the PS-prone group by collecting PS proteins predicted by PScore[35] and a meta-predictor PhaSePred[43] (1,276 proteins, 30,889 variants) and grouped variants from other proteins as low-PS-prone group (56,853 variants). The between-group analysis showed consistent results (Fig. 5B). However, when conducting the between-group analysis using representative PS prediction methods trained on phase-separation proteins, the outcomes for DeePhase[39] and catGRANULE[38] displayed inconsistencies across the analyses (Fig. S6 A and B). For the other three methods, which include PSAP[61], PScore[35], and FuzDrop[62], the PS-prone group did not demonstrate significantly higher Pearson correlation coefficients than the low-PS-prone group (Fig. S6 A and B).

IDRs and structured domains are both crucial for phase separation[6]. When analyzing the PS-prone group defined by algorithms, we found that PSMutPred-IP, along with other sequence-based PS metrics (except PSAP), showed a significant positive correlation with the pathogenicity of variants in both IDRs (n = 15,427) and domains (n = 15,427) (Fig. 5C and S6C).

Additionally, we isolated variants likely in IDRs (unmapped by PfamScan[84,85], with IUPred3[59]score > 0.5) from the PS-prone group and directly used PhaSePred-IP scores

to predict pathogenicity ((Likely) pathogenic as 1 and (Likely) benign as 0). Separate

analyses were conducted for the PS-prone group as defined by PS proteins and by

algorithms (Fig. 5 E and F). The resulting AUROC scores indicated that PSMutPred-IP

can identify disease variants that lead to PS alterations.

Studies indicate that neurodegenerative lesions may be associated with excessive

PS[25,86]. Within the predicted PS-prone group, 19 proteins were identified to be highly

associated with neurodegenerative disease (Methods). SP prediction scores for the

variants (n = 252) of these proteins showed different patterns compared to other variants

(n = 19,266). Specifically, among these 252 variants, those predicted to strengthen PS

were more inclined towards pathogenicity compared to those predicted to weaken PS.

This observation was reflected by the positive Pearson coefficient values for both SP-

LR and SP-RF models, contrasting with the negative coefficient values for variants

from other PS proteins (Fig. 5D). catGRANULE[38] and FuzDrop[62] also demonstrate a

higher Pearson correlation coefficient than the overall pattern (Fig. S6D).

Additionally, we observed that mutations within IDRs of PS proteins are more likely

to affect PS than those in proteins less likely to undergo PS. This conclusion was based

on evaluating the proportion of variants of 'Uncertain Significance' (VUSs) predicted

to impact PS for each protein (see Methods). Using Kolmogorov's D statistic to

compare these proportions, we observed that the PS-prone group had a higher incidence

of PS-affecting variants than the low-PS-prone group, as shown in Fig. 5G. We found

that only for PSAP[61] and PScore[35], PS-prone proteins have a higher proportion of

missense mutations predicted to 'Impact' PS (Fig. S6E).

In summary, our comprehensive analyses revealed a notable clustering of pathogenic

missense mutations with an impact on PS propensity in proteins with inherently higher

PS propensity. Additionally, our findings indicate that compared to mutations that might

weaken PS, gain-of-PS mutations tend to aggregate specifically in disease mutations

associated with neurodegenerative-related genes. These results not only provide

valuable insights into the relationship between PS and disease but also underscore the

reliability and validity of PSMutPred.

Introducing a new feature for pathogenicity prediction of disease variants

Current variant interpretation methods heavily rely on evolutionary features[28,30,31,52,87] generated by multiple sequence alignment (MSA). As IDRs, especially those with poor evolutionary conservation[2], challenge the effectiveness of traditional features[7]. Using two representative pathogenicity prediction models, including EVE[31] and ESM1b[46,88], we aim to test whether PS-related features can address the IDR gaps left by evolutionary features in pathogenicity prediction (Fig. 1, orange section). EVE predicts pathogenic variants by thoroughly leveraging MSA information, demonstrating that this feature alone can predict the impact of most known disease mutations, proving the dominant role of the evolutionary feature in this field. ESM1b utilizes large language models (LLM) to learn protein information across species, modeling the space of known protein sequences selected throughout evolution, and can thus be considered an advanced representation of evolutionary features.

We first selected EVE and applied a straightforward approach, that combined the unsupervised EVE score with a simple feature group, including PSMutPred scores as the variant feature (Methods). We then trained three models including Random Forest (RF), Support Vector Regression (SVR), and Logistic Regression (LR), using ClinVar[64] significances as labels. Their performances were evaluated using both blocked 3-fold cross-validation and an independent test set (Methods). We observed that all three combined models demonstrated improved pathogenicity prediction, with RF showing the best performance in terms of AUROC and AUPR scores (Fig. S4 A, B, and Table. S1). We subsequently selected the RF model as the combined model for further analysis, based on validation on the independent test set.

Given that IDRs typically exhibit poorer evolutionary conservation than Domains[7], it is unsurprising that we found EVE to be less effective in predicting IDR variants compared to Domain variants (Fig. 6 A and B, Table. S1). As expected, the combined model led to a more pronounced improvement in identifying IDR disease variants (n = 5,656) than those within Domains (n = 9,738). Specifically, the combined model

showed a 4.3% improvement in AUROC and a 7.6% improvement in AUPR for IDR

variants, compared to a 2.6% AUROC improvement and only a 1.7% AUPR rise for

Domain variants (Fig. 6 A and B, Table. S1). The Mann-whitely test further indicated a

significant improvement in the prediction of both pathogenic variants as well as benign

variants in IDRs (Fig. 6 C and D). Additionally, we consider that IDRs include a small

subset of potentially conditionally folded IDRs with high evolutionary conservation[4],

characterized by high AlphaFold2[3]-predicted confidence scores (pLDDT scores). For

the 5,656 IDR mutations in the test set, we mapped their pLDDT scores and categorized

them into a high pLDDT group (pLDDT >= 0.7) and a low pLDDT group (pLDDT <

0.5). We found a more pronounced improvement in AUPR for IDR variants with low

pLDDT scores compared to those with high pLDDT scores (9.8% compared to 3.9%

AUPR improvement, Fig. 6E). This indicates that current PS-related features can

supplement the evolutionary feature's weakness in IDRs, especially for low

conservation IDRs.

To investigate the role of PSMutPred in pathogenicity prediction, we examined the

effect of including PSMutPred scores within various feature combinations, and their

performance on the test set was quantified by AUROC and AUPR (Fig. S4C). The

comparative analysis highlighted a significant enhancement in the prediction accuracy

when PSMutPred scores were incorporated. Next, we analyzed ClinVar mutations

predicted by EVE with opposite outcomes (pathogenic/likely pathogenic mutations

with EVE scores < 0.5 and benign/likely benign mutations with EVE scores ≥ 0.5) to

evaluate the PSMutPred's discriminative power on these mutations. We focused on

mutations within disordered regions of potential phase-separating proteins (n = 600,

Fig. S4D). PSMutPred scores for false negatives of EVE are significantly higher than

those for false positives of EVE, with P-values of 1.3e-4, 9.7e-5 and 9e-3 for IP-RF, IP-

SVR and IP-LR, respectively, as evaluated by the Mann-Whitney test. This shows that

PSMutPred can capture information of mutants where evolutionary features failed.

Subsequently, similar to the process of appending features to EVE, we combined PS-

related features with the ESM1b score, which can be considered an advanced

representation of evolutionary features. We mapped ESM1b scores to 140,320 ClinVar

variants and tested the new combined RF model using a blocked 5-fold cross-validation

(Methods). The combined model also achieved improved prediction accuracy (Fig. 6F),

especially for variants located in IDRs with low pLDDT scores (Fig. 6G). This

consistent result indicates that PS-related features can address the weaknesses of

evolutionary features in predicting IDR variants, particularly those in IDRs with low

conservation. Using the PS features, combined with ESM1b scores, we predicted

pathogenicity scores for 1,015,769 ClinVar variants of uncertain significance

(Methods). Among them, 527,524 are IDR variants (Fig. 6H), 9.3% were predicted

pathogenic, and 78.4% were predicted benign (Fig. 6I).

We chose EVE, ESM1b due to their unsupervised nature, which can offer an unbiased

baseline, making testing with our features straightforward. These findings reveal that

the PS-related features including variants' impact on PS serve as a valuable encoding

for IDR mutations and can be integrated into pathogenicity prediction models in the

future to provide a better interpretation of pathogenicity variants.

451 **Discussion**

Missense mutations in IDRs are overlooked compared to those in folded regions.

Pathogenicity prediction models, such as AlphaMissense[32], typically rely on

evolutionary information and protein structural features. However, since IDRs often

lack a consistent structure and show limited evolutionary conservation, using these

traditional features to encode variants in IDRs turns out to be unsuitable, making

interpreting the numerous VUSs in IDRs difficult[2]. To address the challenge, we turn

to phase separation (PS), a widely acknowledged property of IDRs linked to various

diseases. As recent studies emphasize, mutations within IDRs can disrupt regular PS,

which can lead to disease[22,23], highlighting its potential as a predictive feature.

Although a comprehensive understanding of the specific mechanisms by which

missense mutations alter PS remains elusive[7], we considered differences in

physicochemical properties, viewing them as features that quantify mutation-induced

changes. Additionally, we used the predicted pi-pi contact frequency[35] at the mutation

site as a position-specific encoding. These considerations were used to develop models

predicting the effect of missense mutation on PS. Separate models were constructed to

predict the 'Impact' and 'Strengthen/Weaken' effects of missense variants. The

reliability of the model predictions was validated through prediction results on variants

from unseen proteins and further experimental validations. Moreover, subsequent

analyses conducted on a larger-scale dataset further affirmed the accuracy and

effectiveness of our models.

It is crucial to note that, for single or a few key amino acids, changes at the hot spots

involved in charged or multivalent interactions can impact phase separation. These key

residues vary among different proteins, with some being specific residues responsible

for post-translational modifications (PTMs)[89,90], polar interaction[16] or pi-contact[17,18]

within low complexity domains (LCD), while others are residues located at domain

surfaces or boundaries facilitating multivalent interactions[19,20], or contributing to cross-

beta structures within LCD[21]. Given this, we developed machine learning models using

both randomly generated mutations and experimentally validated missense mutations

that alter PS propensity. These mutations mainly involve single amino acid mutations,

aiming to uncover the unique characteristics of 'hot spots' mutations that alter phase

separation. The discriminative power of our algorithms indicates an inherent pattern

among missense variants impacting PS, allowing for their prediction.

We currently focus on predicting single-site mutations because single amino acid

substitutions play a more critical role in diagnosing Mendelian diseases and are more

prominently featured in current disease datasets, such as those in ClinVar[64], compared

to multi-site mutations. Multi-site mutations, which we are unsure how to perfectly

encode, are currently treated as supplementary to our dataset to help the model learn

information about amino acid properties or segments that are important for phase

separation. The model trained in this manner has shown a good ability to predict the

performance of a single amino acid mutation that alters phase separation. Our final

model only predicts the effect of single amino acid mutation at present. Going forward,

as the dataset expands, different approaches to processing multi-site mutation samples

can be explored and the development of a more comprehensive multi-site mutation

prediction approach can be employed to improve the accuracy and adaptability of
PSMutPred.

Through analysis of PSMutPred scores using ClinVar[64] variants, we discovered that
proteins prone to PS have a higher proportion of disease variants altering PS, compared
to proteins less likely to undergo this process. Among these mutations, those likely to
enhance PS were proportionally more prevalent in neurodegenerative-disease-related
genes than those that weaken PS, which supports the current discoveries[25,86]. Therefore,
we can confidently assert that PSMutPred scores serve as valuable encodings for
assessing the potential impact of missense mutations on PS, making them applicable to
a broader range of research.

We further combined PS-related features, including PSMutPred scores, with
evolutionary scores[31] as well as ESM1b[46] to predict pathogenicity and assessed the
performance using ClinVar labels. PSMutPred targets the 'IDR gap' by evaluating
variants in IDRs by their potential effects on phase separation. Consequently, this
approach showed a more comprehensive interpretation of pathogenicity, especially in
IDRs, our intuitive and straightforward collaborative approach offers a more
comprehensive interpretation of pathogenicity across both domains and IDRs. It further
demonstrates that the impact of phase separation features effectively complements the
current focus on pathogenicity interpretations, which predominantly consider ordered
regions. We didn't build a new model from scratch; instead, we highlighted the
contribution of PS to improving the accuracy of pathogenicity predictions, offering a
fresh perspective in this field. Looking ahead, leveraging diverse data such as weak
labels from population frequency and variants from HGMD[91] and gnomAD[92] may
further refine these predictions. Additionally, advanced machine learning methods,
including deep learning, hold the potential to boost accuracy.

However, except IDRs, some folded domains also play important roles in driving
phase separation, such as multivalent tandem structured domains[6]. Our feature
encoding metrics currently favor mutations that impact IDRs rather than those
impacting folded domains. Due to the limited training data available and to avoid
overfitting, we chose simpler feature encoding and traditional machine learning models

instead of developing a complex encoding method to comprehensively capture the

effects of mutations in structured domain areas. In subsequent evaluations of

pathogenicity, we also assessed the pathogenicity of mutations within domains. The

results show that the improvement in predictive accuracy for mutations in IDRs was

significantly greater than for those in domain areas. Including more high-quality,

experimentally validated data, especially regarding the effects of missense mutations

on structured domains, and integrating more comprehensive features could both refine

the prediction of mutation impact on PS.

Ultimately, our study not only presents methods to predict the effects of missense

mutations on PS but also contributes to the prediction and improved understanding of

Variants of Uncertain Significance (VUS) occurring in IDRs, enhancing the diagnostic

accuracy for rare genetic disorders.

Methods

Data acquisition

Experimentally validated missense mutations that impact phase separation (PS) and

their corresponding experimental sequences were curated from the PhaSepDB[47] and

LLPSDB[48,49] databases up to 2022.11. We focused on entries that documented

individual proteins undergoing phase separation, rather than multi-protein co-phase

separation; no additional filtering criteria were applied. In the corresponding literature

of each entry, proteins involved in these entries were reported to undergo phase

separation, along with the mutations that resulted in changes to their phase separation

propensities. We retained single amino acid mutation samples as well as multi amino

acid mutations with a number of mutation sites less than or equal to five. We got 214

single amino acid mutation samples, 43 two-mutation-sites-samples, 26 three-mutation-

sites-samples, 18 four-mutation-sites-samples and 6 five-mutation-sites-samples (Fig.

S1B). We finally obtained 307 samples corresponding to 70 proteins including 79

'Strengthen' samples that strengthened PS, and 228 'Weaken/Disable' samples that

weakened or diminished PS.

As a limited number of experimentally validated mutations that do not affect PS are

available for reference, we created 'Background' samples by generating random single

amino acid (AA) missense mutations with the wild-type sequences for each of the 70

proteins. Mutations already included in the 'Impact' samples were excluded. For each

gene, we generated 500 random mutations, leading to a total of 35,000 mutations.

It should be noted that this does not imply that background mutations cannot alter

PS under any conditions, instead, it allowed us to learn missense mutations that impact

PS.

Statistical analysis of phase separation-related missense mutations

We used PfamScan[84,85] to predict the structural domain regions, abbreviated as

'Domains', within the experimental sequences. Any segment of the sequence not

identified as a domain by PfamScan was annotated as an IDR. Multi-amino acid (AA)

mutation samples were broken down into single AA mutations for subsequent statistical

analysis.

We estimated the pi-contact frequency at the mutation site using the pi-pi interaction

prediction function in PScore[35]. These values correspond to 8 sp^2 groups. The

categorization process divides pi-contacts based on: 1) short-range (4 sequence

separation) vs. long-range (>4), 2) sidechain vs. backbone, and 3) absolute predicted

frequency vs. relative difference from sp2 groups with the same identity respectively

(See PScore[35] for more details).

To analyze the changes in the amino acid properties pre and post-mutation, we

calculated the index differences for properties including hydrophobicity[93], polarity[94],

and mass[95]. The Kolmogorov–Smirnov test was then applied to compare these index

differences between the 'Strengthen' and 'Weaken/Disable' mutation groups.

Development of PSMutPred

Training and testing dataset

We divided the mutation samples into a cross-validation dataset (47 human proteins)

and an independent test set (23 non-human proteins). 'Non-human' in this context

specifically refers to experimental protein sequences originating from species other

than humans. Specifically, we classified the proteins based on their HGNC gene names.
The 47 human proteins were those found in the Human Uniprot database, while the
remaining 23 proteins, which were not found in the Human Uniprot database, were
labeled as non-human proteins.

In the case of the 'Impact Prediction' (IP) models ('IP task' in short), we grouped
'Strengthen' and 'Weaken/Disable' mutations into one group and set their labels to 1,
while the labels of 'Background' mutations were set to 0. For the IP task, we got 246
'Impact' mutation samples, and 23,500 'Background' samples in the cross-validation
dataset, 61 'Impact' samples, and 11,500 'Background' samples in the independent test
set. In the case of the 'Strengthen/Weaken Prediction' (SP) models ('SP task' in short),
labels of 'Strengthen' mutations were set to 1, and labels of 'Weaken/Disable' mutations
were set to 0. For the SP task, we got 174 'Weaken/Disable' samples and 72 'Strengthen'
samples in the cross-validation dataset; 7 'Strengthen' samples, and 54
'Weaken/Disable' samples in the independent test set.

Machine learning features

We used a set of simple features (39 dimensions) to encode each mutation entry. The
two tasks share a common set of feature encoding. For multi-point mutations, certain
features were adjusted to account for the presence of multiple mutation sites. The details
of the features for each sample are described below:

IDR-related features of the mutation site(s) (5 dimensions): The structural domains
of the experimental sequences corresponding to the sample were predicted using
PfamScan[84]. Segments not predicted as Domains were designated as IDRs. 1) One
binary value indicating whether the site is located in IDRs. For multi-point mutation,
this value is set to 0 if any of the mutation sites are located within Domains; 2) One
value quantifies the residue distance of the mutation site to its nearest domain boundary.
For multi-point mutation, this value reflects the average distance of all mutation sites;
3) One value representing the likelihood of the mutation site being within IDRs
predicted by IUPred3[59], For multi-point mutations, we use the average of the predicted
values for all mutation sites; 4) One value representing the average of the full sequence
IUPred3[59] scores; 5) One binary value indicating whether the protein sequence has

IDRs or not.

The predicted pi-contact frequency at the mutation site (8 dimensions): We derived 8

predicted values for pi-pi interactions at the mutation site from the wild-type sequence

using the pi-pi interaction prediction function in PScore[35] (See PScore[35] for more

details). In the case of multi-point mutation, we computed the 8 values as the mean

scores across all mutation positions.

Physicochemical feature encoding for the mutation site (26 dimensions): We targeted

the changes in AA characteristics at the mutation site whenever possible by employing

five physicochemical indices that were previously used for protein feature encoding[96-

99]. We selected hydrophobicity[93], polarity[100], volumes of side chains (VSC) of amino

acids[101], solvent-accessible surface area (SASA)[102], and net charge index (NCI) of side

chains of amino acids[96,103]. For each index, we computed four feature values: 1) Index

value for the wild-type AA at the mutation site. For multi-point mutation, we used the

mean value; 2) Numerical difference in index value between mutant and wild-type AA,

we summed the difference values for multi-point mutations; 3) Average index values

for all AAs in the wild-type sequence' IDRs (defined as areas outside of Domains

predicted by PfamScan[84,85]), and 4) the difference between the index value of mutant

AA and the average index value (from point 3), for multi-point mutation, we

accumulated the difference values for each mutant AA. Additionally, six binary values

(6 dimensions) were used to capture the presence of three properties (positively charged,

negatively charged, and hybridized) in both the mutant AA and wild-type AA. For

multi-point mutations, an element is set to 1 if at least one AA exhibits the respective

property.

Machine learning algorithm and model performance evaluation

For both the IP task and the SP task, to evaluate the generalizability of our models on

variants from unseen proteins, we implemented a blind test called 'leave-one-source-

out cross-validation' (LOSO CV). In this approach, for each validation iteration, we

held out variants from a single protein from the total set of proteins (variants from cross-

validation dataset; 47 proteins). We iteratively held out all variants (both positive and

negative samples) associated with a single protein as the validation dataset from the

total set of cross-validation proteins, while variants from the other proteins in the

dataset were used for model training, and the trained model predicted the values for the

left-out samples. After cycling through all proteins in the dataset, the prediction results

of the validation dataset corresponding to each protein were combined, and metrics

including AUC and AUPR were applied to assess predictive performance.

Especially for the IP task, a dataset balancing process was applied before the LOSO

CV was initiated. First, before initiating the LOSO CV process (i.e., before traversing

samples corresponding to different proteins), we randomly selected a subset of negative

samples. This subset, drawn randomly from the total pool of 35,000 mutations (500 for

each protein), was twice the size of the positive sample set. These selected negative

samples were then combined with the positive samples to form a new sub-dataset. Next,

the LOSO CV procedure was initiated on this new sub-dataset, where we sequentially

traverse different proteins, using the mutation samples corresponding to a single protein

as the test set and the remaining for training the model. This whole process is repeated

100 times for robustness.

We added representative PS predictors including DeePhase[39], PSAP[61], PScore[35],

catGRANULE[38] as well as FuzDrop[62] for comparison in the IP task. For FuzDrop[62], we

applied the residue-level PS scores for calculation. Expecting greater score changes for

'Impact' mutations than 'Background' mutations, we calculated the absolute difference

in scores before and after each mutation to represent the predicted values.

We applied the Python Scikit learn (sklearn) package to develop machine learning

models. Data was normalized by the MinMaxScaler from sklearn before fitting the

models. We trained models including a RandomForestClassifier (with parameters:

class_weight = 'balanced', and max_depth = 10), an SVR (with default parameters),

and a LogisticRregression (with parameters: class_weight = 'balanced', kernel =

'liblinear', and penalty = '11') for both the IP and the SP task. We calculated the area

under the curve of the receiving operating characteristics (AUROC) with the sklearn

roc_auc_score, and roc_curve metric to evaluate the performance. Additionally, the

precision and recall were calculated with sklearn precision_recall_curve with the area

under   the   curve   of   the   precision-recall   curve   (PRC)   (AUPR)   using

precision_recall_curve. The discriminative power of models was evaluated using the

Mann-Whitely test.

The cross-validation dataset and independent test set were combined to train the final

PSMutPred models for both tasks. According to the models' performance, we produced

IP-RF, IP-SVR, and IP-LR scores and SP-RF, SP-LR scores. To prevent the overuse of

negative samples in the IP task, for each model, 10 different subsets of samples were

randomly sampled from the 'Background' mutations with twice the size of the collected

'Impact' mutations to train 10 sub-models. The averaged prediction scores of the 10

trained models as well as the ranks for these scores in all ClinVar[63,64] variants were used

as the final prediction scores for each IP-RF, IP-SVR, and IP-LR model. These models

were used to predict PSMutPred scores for each missense variant in ClinVar.

Between-group analysis

Missense mutation data was collected from ClinVar up to 2022.12, which contains

522,016 variants corresponding to 8,611 human proteins. Of these, 39,166 are

pathogenic or likely pathogenic, 48,576 are benign or likely benign and the remainder

434,274 are 'Uncertain Significance'.

After excluding the training data points for PSMutPred, we divided missense variants

corresponding to 8,611 proteins into a PS-prone group and a low-PS propensity group

in two ways: 1. By grouping variants from experiment-verified PS protein to the PS-

prone group and the rest to the low-PS-prone group. Specifically, we referenced 155

human PS proteins (59 PS-Self proteins, 96 PS-Part proteins) from PhaSepDB[43,47],

among which 83 were found in ClinVar[63]. This resulted in 1,451 variants in the PS-

prone group, leaving 86,291 variants for the low-PS-prone group. 2. Variants from

predicted PS proteins, based on either a PScore[35] higher than 4 or a PhaSePred[43] rank

(PdPS-10fea_rnk) higher than 0.9, were allocated to the PS-prone group. This resulted

in 30,889 variants from 1,276 proteins in the PS-prone group and leaving 56,853

variants for the low-PS-prone group.

We first performed analysis excluding variants with 'Uncertain significance'. For

each group, the Pearson correlation score was computed using the Python Scipy

package by comparing the PSMutPred-IP ranks of its variants against their respective

ClinVar pathogenicity labels (pathogenic or likely pathogenic coded as 1 and benign or

likely benign coded as 0). The ranks of the absolute score difference of DeePhase[39],

PSAP[61], PScore[35], catGRANULE[38] as well as FuzDrop[62], were also used to compute

the Pearson correlation.

The significance of each computed Pearson value is annotated with corresponding P-

values, indicating the robustness of the correlation within each group. We compared the

coefficient values between the PS-prone group and the low-PS-prone group. We also

computed and compared the coefficient values between IDR variants (n = 15,427) and

Domain variants (n = 15,462) within the PS-prone group defined by predictions (a

variant was classified by Domain variants if identified in a structured domain by

PfamScan[84,85] and as an IDR variant if it is unmapped).

In addition, we divided the PS-prone group into a neurodegenerative disease (ND)

group and those that are not into a non-ND group. To do that, we collected disease genes

related to common neurodegenerative diseases including Alzheimer's disease,

Amyotrophic lateral sclerosis, Frontotemporal dementia, Huntington's disease,

Multiple Sclerosis, and Parkinson's disease from DisGeNet[104]. After removing

duplicates, we screened genes with a gene-disease association score (GDA score)

higher than 0.5, resulting in a subset of 91 genes, 19 of which were both matched in

ClinVar genes and the PS-prone group (363 variants, 252 being predicted 'Impact'

mutation by PSMutPred-IP, having a combined IP-LR, IP-SVR, and IP-RF rank score

sum above 1.5). We compared the coefficient values of PSMutPred-SP between variants

corresponding to the ND genes and those that were not within the predicted PS-prone

group. The ranks of the score difference of various sequence-based PS methods, were

also used to compute the Pearson correlation.

We then performed between-group analysis to test whether proteins that have a higher

propensity to undergo PS have a higher proportion of variants that are predicted to

impact PS. For each PSMutPred-IP model (IP-RF, IP-LR, IP-SVR), we calculated the

proportion of IDR variants of 'Uncertain Significance' (VUSs) predicted as 'Impact'

(PSMutPred-IP rank score > 0.8) for each protein. The proportion values of the PS-

prone group and the low-PS-prone group were compared. We compared the PS-prone group (1,276 proteins) with the low-PS-prone group (7,335 proteins) defined by algorithm prediction and compared the PS-prone group (83 proteins) with the low-PS-prone group (8,528 proteins) defined by experimental-verified PS proteins. Sequence-based PS methods were also added to the comparison, and for each method, a variant is predicted to be an 'Impact' variant if its corresponding absolute score difference rank is above 0.8.

Analysis of phase separation feature's performance in refining pathogenicity prediction

Dataset acquisition

We collected 'EVE_scores_ASM' scores from EVE[31], which include scoring for all possible missense variants corresponding to 3,219 disease-associated genes. Among them, 47,870 ClinVar[63] variants (by annotations up to 2021.4.) were used to evaluate the EVE model, 30,758 pathogenic or likely pathogenic were marked as positive samples and 17,112 'Benign' or 'Likely benign' variants were marked as negative samples. We applied these samples for model cross-validation (cross-validation set). To construct an independent test dataset, we screened new variants (ClinVar[63] annotations up to 2022.12) that were not used to evaluate the EVE model and removed variants in the same amino acid position seen in the cross-validation set to avoid data leakage. This resulted in 15,394 variants including 8,709 pathogenic or likely pathogenic variants and 6,685 benign or likely benign variants.

We collected ESM1b[46] scores, which encompass the scoring of all possible missense variants for 42,286 proteins and their isoforms. In total, scores for 140,321 variants were mapped to ClinVar[63]. These variants were then used as a cross-validation dataset to evaluate the combination of ESM1b and PS features.

Pathogenicity prediction combining PS-related features

To test whether the integration of PS-related properties can improve the accuracy of pathogenicity prediction. We built a model based on the following features: PSMutPred-IP model scores predicting the propensity of the missense mutation to

impact PS (3 dim, predicted values of IP-LR, IP-RF, and IP-SVR); PSMutPred-SP

model scores predicting the direction of the shift in the normal PS threshold induced by

the mutation (2 dim, SP-LR, SP-RF); PScore[35] representing the PS tendency of the

wild-type protein (1 dim); IUPred[59] score of the mutation site representing the

probability of the site to locate in IDRs (1 dim); binary encoding whether the mutation

is in a Domain predicted by PfamScan[84,85] (1 dim); the residue distance of the mutation

site to the nearest IDR (1 dim); one-hot encoding for wild-type and mutant aa (40 dim);

EVE score or ESM1b score of the variant (1 dim). We trained models including a

RandomForestClassifier (with parameters: n_estimators = 200, class_weight =

'balanced', and max_depth = 15), an SVR (with default parameters), and a

LogisticRregression (with parameters: class_weight = 'balanced', kernel = 'liblinear',

and penalty = 'l1'). The dataset was standardized using the MinMaxScaler from Scikit

Learn before fitting each model. Based on the performance, we selected the

RandomForestClassifier as the optimal model.

Model performance benchmarks

To evaluate the models on the cross-validation set, we applied a blocked n-fold cross-

validation where variants from the same gene were strictly assigned to the same group.

To evaluate the models on the independent test set, the models were first trained on the

cross-validation set and then evaluated using the independent test set.

Pathogenicity prediction

The data points were combined to train a final pathogenicity predictor (140,321

variants; PS feat. + ESM1b predictor). And to transform the continuous pathogenicity

scores into 'Pathogenic', 'Likely benign', and 'Uncertain' categories, we determined

0.5 as the initial threshold to distinguish between 'Likely pathogenic' and 'Likely

benign' based on the test set's F1 score. To define the uncertainty of prediction, we

considered applying an offset around this threshold. For instance, with a 0.1 offset,

predictions below 0.4 are categorized as 'Likely benign', while those above 0.6 are

classified as 'Likely pathogenic'. We observed that starting from the initial 0.5

threshold, the AUROC and Accuracy improved as the offset increased (Fig. S4F).

Experimental Materials

Plasmid preparation and cell culture

The full-length coding sequences of mouse *Eps8* (NM_007945.4) and its mutations were PCR amplified and cloned into the pEGFP-C3 vector. These recombinant plasmids were then transiently transfected into HEK293 cells using the Lipofectamine 3000 Kit (Invitrogen), with 1-2 μg of plasmid used per well in a 12-well plate (Costar, Corning) for each transfection. Before fixation, HEK293 cells were cultured for 16-24 hours in DMEM medium (Gibco) supplemented with 10% FBS (Gibco) and 1% Penicillin-Streptomycin Solution (Gibco), under 5% CO_2 conditions. Cells were washed 3 times with PBS for 3 minutes each, then fixed in 4% PFA for 20 minutes at room temperature. Detailed validation information and references for these products can be found on the respective manufacturers' websites.

Image analysis and quantification

HEK293 cells were visualized using an IX73 Inverted Fluorescence Microscope (Olympus) equipped with a 60× 1.42 NA Plan Apochromat oil objective. Images were captured with an Iris 9 sCMOS camera (Teledyne Photometrics) and the setup was controlled by the cellSens imaging system (Olympus). After subtracting the background and setting an identical fluorescence threshold, ImageJ software was applied to quantify the number of puncta per cell, with modules including 'Analysis Particles', 'Find Maxima', and 'Set Measurement'. The statistical analysis applied a two-tailed Student's t-test for experimental comparisons of puncta numbers using GraphPad Prism 9. Each assay was performed at least three times.

Protein structure analysis of EPS8

We performed the sequence alignment of EPS8 with the ClustalX tool. The structures of human and mouse EPS8 were predicted using AlphaFold2[3,105]. The interaction among residues was analyzed using the crystal structure of the SH3 domain of EPS8 (PDB: 7TZK). The protein structure analysis was performed using PyMol version 2.5.0.

Data Availability

Variants used in this study are collected from PhaSepDB (http://db.phasep.pro), LLPSDB (v2.0), and ClinVar. All data supporting the findings of this study are available within the article and its Supplementary Information files or from the corresponding author upon reasonable request. Source data are provided with this paper.

Code Availability

PSMutPred is freely available at https://codeocean.com/capsule/5744011/tree.

References

1.	Vacic, V. & Iakoucheva, L.M. Disease mutations in disordered regions--exception to the rule? *Mol Biosyst* **8**, 27-32 (2012).
2.	Colak, R. *et al.* Distinct types of disorder in the human proteome: functional implications for alternative splicing. *PLoS Comput Biol* **9**, e1003030 (2013).
3.	Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-+ (2021).
4.	Alderson, T.R., Pritisanac, I., Kolaric, D., Moses, A.M. & Forman-Kay, J.D. Systematic identification of conditionally folded intrinsically disordered regions by AlphaFold2. *Proc Natl Acad Sci U S A* **120**, e2304302120 (2023).
5.	Alberti, S. Phase separation in biology. *Curr Biol* **27**, R1097-r1102 (2017).
6.	Gao, Y., Li, X., Li, P. & Lin, Y. A brief guideline for studies of phase-separated biomolecular condensates. *Nat Chem Biol* **18**, 1307-1318 (2022).
7.	Tsang, B., Pritisanac, I., Scherer, S.W., Moses, A.M. & Forman-Kay, J.D. Phase Separation as a Missing Mechanism for Interpretation of Disease Mutations. *Cell* **183**, 1742-1756 (2020).
8.	Gomes, E. & Shorter, J. The molecular language of membraneless organelles. *J Biol Chem* **294**, 7115-7127 (2019).
9.	Banani, S.F., Lee, H.O., Hyman, A.A. & Rosen, M.K. Biomolecular condensates: organizers of cellular biochemistry. *Nat Rev Mol Cell Biol* **18**, 285-298 (2017).
10.	Wang, H. *et al.* Temporal and spatial assembly of inner ear hair cell ankle link condensate through phase separation. *Nat Commun* **14**, 1657 (2023).
11.	Lin, L. *et al.* Phase separation-mediated condensation of Whirlin-Myo15-Eps8 stereocilia tip complex. *Cell Rep* **34**, 108770 (2021).
12.	He, Y., Li, J. & Zhang, M. Myosin VII, USH1C, and ANKS4B or USH1G Together Form Condensed Molecular Assembly via Liquid-Liquid Phase Separation. *Cell Rep* **29**, 974-986.e4 (2019).
13.	Molliex, A. *et al.* Phase separation by low complexity domains promotes stress granule assembly and drives pathological fibrillization. *Cell* **163**, 123-33 (2015).
14.	Murakami, T. *et al.* ALS/FTD Mutation-Induced Phase Transition of FUS Liquid Droplets and

Reversible Hydrogels into Irreversible Hydrogels Impairs RNP Granule Function. *Neuron* **88**,
678-90 (2015).

15.　Gopal, P.P., Nirschl, J.J., Klinman, E. & Holzbaur, E.L. Amyotrophic lateral sclerosis-linked
mutations increase the viscosity of liquid-like TDP-43 RNP granules in neurons. *Proc Natl Acad*
Sci U S A **114**, E2466-e2475 (2017).

16.　Kim, G.H. & Kwon, I. Distinct roles of hnRNPH1 low-complexity domains in splicing and
transcription. *Proc Natl Acad Sci U S A* **118**(2021).

17.　Wong, L.E., Kim, T.H., Muhandiram, D.R., Forman-Kay, J.D. & Kay, L.E. NMR Experiments for
Studies of Dilute and Condensed Protein Phases: Application to the Phase-Separating Protein
CAPRIN1. *J Am Chem Soc* **142**, 2471-2489 (2020).

18.　Kim, T.H. *et al.* Interaction hot spots for phase separation revealed by NMR studies of a CAPRIN1
condensed phase. *Proc Natl Acad Sci U S A* **118**(2021).

19.　Bierma, J.C. *et al.* Controlling Liquid-Liquid Phase Separation of Cold-Adapted Crystallin
Proteins from the Antarctic Toothfish. *J Mol Biol* **430**, 5151-5168 (2018).

20.　Gui, X. *et al.* Structural basis for reversible amyloids of hnRNPA1 elucidates their role in stress
granule assembly. *Nat Commun* **10**, 2006 (2019).

21.　Zhou, X. *et al.* Mutations linked to neurological disease enhance self-association of low-
complexity protein sequences. *Science* **377**, eabn5582 (2022).

22.　Niaki, A.G. *et al.* Loss of Dynamic RNA Interaction and Aberrant Phase Separation Induced by
Two Distinct Types of ALS/FTD-Linked FUS Mutations. *Mol Cell* **77**, 82-94.e4 (2020).

23.　Hofweber, M. *et al.* Phase Separation of FUS Is Suppressed by Its Nuclear Import Receptor and
Arginine Methylation. *Cell* **173**, 706-719.e13 (2018).

24.　Patel, A. *et al.* A Liquid-to-Solid Phase Transition of the ALS Protein FUS Accelerated by Disease
Mutation. *Cell* **162**, 1066-77 (2015).

25.　Silva, J.L. *et al.* Targeting Biomolecular Condensation and Protein Aggregation against Cancer.
Chem Rev **123**, 9094-9138 (2023).

26.　Xiang, J. *et al.* Development of an α-synuclein positron emission tomography tracer for imaging
synucleinopathies. *Cell* **186**, 3350-3367.e19 (2023).

27.　Fan, Y. *et al.* Generic amyloid fibrillation of TMEM106B in patient with Parkinson's disease
dementia and normal elders. *Cell Res* **32**, 585-588 (2022).

28.　Raimondi, D. *et al.* DEOGEN2: prediction and interactive visualization of single amino acid
variant deleteriousness in human proteins. *Nucleic Acids Res* **45**, W201-w206 (2017).

29.　Fariselli, P., Martelli, P.L., Savojardo, C. & Casadio, R. INPS: predicting the impact of non-
synonymous variations on protein stability from sequence. *Bioinformatics* **31**, 2816-21 (2015).

30.　Rentzsch, P., Witten, D., Cooper, G.M., Shendure, J. & Kircher, M. CADD: predicting the
deleteriousness of variants throughout the human genome. *Nucleic Acids Res* **47**, D886-d894
(2019).

31.　Frazer, J. *et al.* Disease variant prediction with deep generative models of evolutionary data.
Nature **599**, 91-95 (2021).

32.　Cheng, J. *et al.* Accurate proteome-wide missense variant effect prediction with AlphaMissense.
Science **381**, eadg7492 (2023).

33.　Lancaster, A.K., Nutter-Upham, A., Lindquist, S. & King, O.D. PLAAC: a web and command-
line application to identify proteins with prion-like amino acid composition. *Bioinformatics* **30**,
2501-2 (2014).

34. Chong, P.A., Vernon, R.M. & Forman-Kay, J.D. RGG/RG Motif Regions in RNA Binding and
Phase Separation. *J Mol Biol* **430**, 4650-4665 (2018).

35. Vernon, R.M. *et al.* Pi-Pi contacts are an overlooked protein feature relevant to phase separation.
Elife **7**(2018).

36. Martin, E.W. *et al.* Valence and patterning of aromatic residues determine the phase behavior of
prion-like domains. *Science* **367**, 694-699 (2020).

37. Wang, J. *et al.* A Molecular Grammar Governing the Driving Forces for Phase Separation of Prion-
like RNA Binding Proteins. *Cell* **174**, 688-699.e16 (2018).

38. Bolognesi, B. *et al.* A Concentration-Dependent Liquid Phase Separation Can Cause Toxicity
upon Increased Protein Expression. *Cell Rep* **16**, 222-231 (2016).

39. Saar, K.L. *et al.* Learning the molecular grammar of protein condensates from sequence
determinants and embeddings. *Proc Natl Acad Sci U S A* **118**(2021).

40. Dignon, G.L., Best, R.B. & Mittal, J. Biomolecular Phase Separation: From Molecular Driving
Forces to Macroscopic Properties. *Annu Rev Phys Chem* **71**, 53-75 (2020).

41. Brangwynne, Clifford P., Tompa, P. & Pappu, Rohit V. Polymer physics of intracellular phase
transitions. *Nature Physics* **11**, 899-904 (2015).

42. Martin, E.W. & Mittag, T. Relationship of Sequence and Phase Separation in Protein Low-
Complexity Regions. *Biochemistry* **57**, 2478-2487 (2018).

43. Chen, Z. *et al.* Screening membraneless organelle participants with machine-learning models that
integrate multimodal features. *Proc Natl Acad Sci U S A* **119**, e2115369119 (2022).

44. Shen, B. *et al.* Computational Screening of Phase-separating Proteins. *Genomics Proteomics
Bioinformatics* **19**, 13-24 (2021).

45. Monahan, Z. *et al.* Phosphorylation of the FUS low-complexity domain disrupts phase separation,
aggregation, and toxicity. *Embo j* **36**, 2951-2967 (2017).

46. Brandes, N., Goldman, G., Wang, C.H., Ye, C.J. & Ntranos, V. Genome-wide prediction of disease
variant effects with a deep protein language model. *Nat Genet* **55**, 1512-1522 (2023).

47. You, K. *et al.* PhaSepDB: a database of liquid-liquid phase separation related proteins. *Nucleic
Acids Res* **48**, D354-d359 (2020).

48. Li, Q. *et al.* LLPSDB: a database of proteins undergoing liquid-liquid phase separation in vitro.
Nucleic Acids Res **48**, D320-d327 (2020).

49. Wang, X. *et al.* LLPSDB v2.0: an updated database of proteins undergoing liquid-liquid phase
separation in vitro. *Bioinformatics* **38**, 2010-2014 (2022).

50. Yamasaki, A. *et al.* Liquidity Is a Critical Determinant for Selective Autophagy of Protein
Condensates. *Mol Cell* **77**, 1163-1175.e9 (2020).

51. Koehler, L.C. *et al.* TDP-43 Oligomerization and Phase Separation Properties Are Necessary for
Autoregulation. *Front Neurosci* **16**, 818655 (2022).

52. Li, G., Panday, S.K. & Alexov, E. SAAFEC-SEQ: A Sequence-Based Method for Predicting the
Effect of Single Point Mutations on Protein Thermodynamic Stability. *Int J Mol Sci* **22**(2021).

53. Quan, L., Lv, Q. & Zhang, Y. STRUM: structure-based prediction of protein stability changes
upon single-point mutation. *Bioinformatics* **32**, 2936-46 (2016).

54. Geng, C., Vangone, A., Folkers, G.E., Xue, L.C. & Bonvin, A. iSEE: Interface structure, evolution,
and energy-based machine learning predictor of binding affinity changes upon mutations. *Proteins*
87, 110-119 (2019).

55. Iqbal, S. *et al.* Assessing the performance of computational predictors for estimating protein

stability changes upon missense mutations. *Brief Bioinform* **22**(2021).

56. Fisher, R.S. & Elbaum-Garfinkle, S. Tunable multiphase dynamics of arginine and lysine liquid
condensates. *Nat Commun* **11**, 4628 (2020).

57. Ukmar-Godec, T. *et al.* Lysine/RNA-interactions drive and regulate biomolecular condensation.
Nat Commun **10**, 2909 (2019).

58. Qin, Z. *et al.* Deactylation by SIRT1 enables liquid-liquid phase separation of IRF3/IRF7 in innate
antiviral immunity. *Nat Immunol* **23**, 1193-1207 (2022).

59. Erdős, G., Pajkos, M. & Dosztányi, Z. IUPred3: prediction of protein disorder enhanced with
unambiguous experimental annotation and visualization of evolutionary conservation. *Nucleic*
Acids Res **49**, W297-w303 (2021).

60. Adzhubei, I.A. *et al.* A method and server for predicting damaging missense mutations. *Nat*
Methods **7**, 248-9 (2010).

61. van Mierlo, G. *et al.* Predicting protein condensate formation using machine learning. *Cell Rep*
34, 108705 (2021).

62. Hardenberg, M., Horvath, A., Ambrus, V., Fuxreiter, M. & Vendruscolo, M. Widespread
occurrence of the droplet state of proteins in the human proteome. *Proc Natl Acad Sci U S A* **117**,
33254-33262 (2020).

63. Landrum, M.J. & Kattman, B.L. ClinVar at five years: Delivering on the promise. *Hum Mutat* **39**,
1623-1630 (2018).

64. Landrum, M.J. *et al.* ClinVar: improvements to accessing data. *Nucleic Acids Res* **48**, D835-d844
(2020).

65. Manor, U. *et al.* Regulation of stereocilia length by myosin XVa and whirlin depends on the actin-
regulatory protein Eps8. *Curr Biol* **21**, 167-72 (2011).

66. Frittoli, E. *et al.* The signaling adaptor Eps8 is an essential actin capping protein for dendritic cell
migration. *Immunity* **35**, 388-99 (2011).

67. Yap, L.F. *et al.* Upregulation of Eps8 in oral squamous cell carcinoma promotes cell migration
and invasion through integrin-dependent Rac1 activation. *Oncogene* **28**, 2524-34 (2009).

68. Menna, E. *et al.* Eps8 regulates axonal filopodia in hippocampal neurons in response to brain-
derived neurotrophic factor (BDNF). *PLoS Biol* **7**, e1000138 (2009).

69. Hertzog, M. *et al.* Molecular basis for the dual function of Eps8 on actin dynamics: bundling and
capping. *PLoS Biol* **8**, e1000387 (2010).

70. Disanza, A. *et al.* Regulation of cell shape by Cdc42 is mediated by the synergic actin-bundling
activity of the Eps8-IRSp53 complex. *Nat Cell Biol* **8**, 1337-47 (2006).

71. Disanza, A. *et al.* Eps8 controls actin-based motility by capping the barbed ends of actin filaments.
Nat Cell Biol **6**, 1180-8 (2004).

72. Shi, Y., Lin, L., Wang, C. & Zhu, J. Promotion of row 1-specific tip complex condensates by
Gpsm2-Gαi provides insights into row identity of the tallest stereocilia. *Sci Adv* **8**, eabn4556
(2022).

73. Brangwynne, C.P. *et al.* Germline P granules are liquid droplets that localize by controlled
dissolution/condensation. *Science* **324**, 1729-32 (2009).

74. Chen, X., Wu, X., Wu, H. & Zhang, M. Phase separation at the synapse. *Nat Neurosci* **23**, 301-
310 (2020).

75. Shin, Y. *et al.* Spatiotemporal Control of Intracellular Phase Transitions Using Light-Activated
optoDroplets. *Cell* **168**, 159-171.e14 (2017).

76. Bhat, P., Honson, D. & Guttman, M. Nuclear compartmentalization as a mechanism of
quantitative control of gene expression. *Nat Rev Mol Cell Biol* **22**, 653-670 (2021).

77. Ong, J.Y. & Torres, J.Z. Phase Separation in Cell Division. *Mol Cell* **80**, 9-20 (2020).

78. Wu, X. *et al.* Vesicle Tethering on the Surface of Phase-Separated Active Zone Condensates. *Mol*
Cell **81**, 13-24.e7 (2021).

79. Wu, X., Cai, Q., Feng, Z. & Zhang, M. Liquid-Liquid Phase Separation in Neuronal Development
and Synaptic Signaling. *Dev Cell* **55**, 18-29 (2020).

80. Xiao, Q., McAtee, C.K. & Su, X. Phase separation in immune signalling. *Nat Rev Immunol* **22**,
188-199 (2022).

81. Noda, N.N., Wang, Z. & Zhang, H. Liquid-liquid phase separation in autophagy. *J Cell Biol*
219(2020).

82. Su, Q., Mehta, S. & Zhang, J. Liquid-liquid phase separation: Orchestrating cell signaling through
time and space. *Mol Cell* **81**, 4137-4146 (2021).

83. Sanders, D.W. *et al.* Competing Protein-RNA Interaction Networks Control Multiphase
Intracellular Organization. *Cell* **181**, 306-324.e28 (2020).

84. Eddy, S.R. Accelerated Profile HMM Searches. *PLoS Comput Biol* **7**, e1002195 (2011).

85. Mistry, J. *et al.* Pfam: The protein families database in 2021. *Nucleic Acids Res* **49**, D412-d419
(2021).

86. Zbinden, A., Pérez-Berlanga, M., De Rossi, P. & Polymenidou, M. Phase Separation and
Neurodegenerative Diseases: A Disturbance in the Force. *Dev Cell* **55**, 45-68 (2020).

87. Liu, Y., Zhang, T., You, N., Wu, S. & Shen, N. MAGPIE: accurate pathogenic prediction for
multiple variant types using machine learning approach. *Genome Med* **16**, 3 (2024).

88. Rives, A. *et al.* Biological structure and function emerge from scaling unsupervised learning to
250 million protein sequences. *Proc Natl Acad Sci U S A* **118**(2021).

89. Guillen-Boixet, J. *et al.* RNA-Induced Conformational Switching and Clustering of G3BP Drive
Stress Granule Assembly by Condensation. *Cell* **181**, 346-361 e17 (2020).

90. Saito, M. *et al.* Acetylation of intrinsically disordered regions regulates phase separation. *Nat*
Chem Biol **15**, 51-61 (2019).

91. Stenson, P.D. *et al.* The Human Gene Mutation Database (HGMD(®)): optimizing its use in a
clinical diagnostic or research setting. *Hum Genet* **139**, 1197-1207 (2020).

92. Karczewski, K.J., Francioli, L.C., Tiao, G., Cummings, B.B. & Xavier, R.J. The mutational
constraint spectrum quantified from variation in 141,456 humans. *Nature* **581**, 434-443 (2020).

93. Tanford, C. Contribution of Hydrophobic Interactions to the Stability of the Globular
Conformation of Proteins. *J.am.chem.soc* **84**, 4240-4247 (1962).

94. Zimmerman, J.M., Eliezer, N. & Simha, R. The characterization of amino acid sequences in
proteins by statistical methods. *Journal of Theoretical Biology* **21**, 170-201 (1968).

95. Nelson, D.L. & Cox, M.M. *Lehninger Principles of Biochemistry*, (Lehninger principles of
biochemistry., 2005).

96. Guo, Y., Yu, L., Wen, Z. & Li, M. Using support vector machine combined with auto covariance
to predict protein-protein interactions from protein sequences. *Nucleic Acids Res* **36**, 3025-30
(2008).

97. Li, X. *et al.* SDNN-PPI: self-attention with deep neural network effect on protein-protein
interaction prediction. *BMC Genomics* **23**, 474 (2022).

98. Guo, Z. *et al.* 3D genome assisted protein–protein interaction prediction. *Future generations*

computer systems: FGCS (2022).

99.   Zhu-Hong *et al.* Prediction of protein-protein interactions from amino acid sequences with ensemble extreme learning machines and principal component analysis. *BMC Bioinformatics* **14**(2013).

100.   Grantham & R. Amino Acid Difference Formula to Help Explain Protein Evolution. *Science* **185**, 862-864 (1974).

101.   Pontius, J., Richelle, J. & Wodak, S.J. Deviations from standard atomic volumes as a quality measure for protein crystal structures. *Journal of Molecular Biology* **264**, 121-136 (1996).

102.   Lee, B.K. & Richards, F.M. interpretation of protein structures: estimation of static accessibility. *Journal of Molecular Biology* **55**, 379,IN3-400,IN4 (1971).

103.   Zhou, P., Tian, F.F., Li, B., Wu, S.R. & Li, Z.L. Genetic algorithm-based virtual screening of combinative mode for peptide/protein. *Acta Chimica Sinica* **64**, 691-697 (2006).

104.   Piñero, J. *et al.* The DisGeNET knowledge platform for disease genomics: 2019 update. *Nucleic Acids Res* **48**, D845-d855 (2020).

105.   Tunyasuvunakool, K. *et al.* Highly accurate protein structure prediction for the human proteome. *Nature* **596**, 590-596 (2021).

Acknowledgement

This project is supported by the National Key Research and Development Program (2022YFE0125300), Innovation Program of Shanghai Municipal Education Commission (2023ZKZD16), the National Natural Science Foundation of China (82071262, 32300464, 81671326, 19Z103150073), Natural Science Foundation of Shanghai (20ZR1427200, 20511101900, 21ZR1433000), Shanghai Municipal Science and Technology Major Project (2017SHZDZX01, 20JC1418600), the Shanghai Leading Academic Discipline Project (B205), China Postdoctoral Science Foundation (2023M732266) and Shanghai Jiao Tong University STAR Grant (YG2023ZD26, YG2022ZD024, YG2022QN111, YG2021QN135).

Author contributions

QL, GH, YS, and MF conceived the concept; MF, XW, QL, and YS designed experiments; QL led the project with assistance from MF, GH, YS, and XW; XW, MX, and QL collected samples; MF, XW performed experiments with assistance from QL, GH, YS, and LL; MF, XW and QL analyzed data; MF, XW, XZ, QL, and YS wrote the manuscript with input from all co-authors.

Competing Interests Statement

The authors declare no competing interests

Figures

Figure1

 Fig. 1. Overview of the study. The upper green panel illustrates PSMutPred, a machine

 learning approach designed to predict the effect of missense mutations on natural phase

 separation. Each mutation is converted into a feature vector and distinct models were

 employed for two main tasks: Identifying mutations that impact PS (termed 'Impact

 Prediction') and determining whether a mutation strengthens or weakens the PS

 threshold (labeled as 'Strengthen/Weaken Prediction'). Additionally, PS features,

 including the output from PSMutPred, were evaluated for their utility in predicting the

 pathogenicity of missense variants (lower orange panel).

1083 **Figure2**

1084

1085

Fig. 2. Analyses of mutations that impact phase separation (PS).

(A) Comparison of the PS propensity of proteins corresponding to collected mutations

(70 proteins) with the PS propensity of the human proteome. (PScore[35], Left;

PhaSePred[43], Right; ****P < 0.0001, two-sided Mann–Whitney U test).

(B) The proportion of mutations affecting phase separation ('Impact' mutations) (Left)

located in IDRs and Domains, compared with the total proportion of IDRs and Domains

(Right). 'Impact' mutations are more likely to be located in IDRs rather than Domains.

(C) The top 30 high-frequency mutations among collected 'Impact' mutations.

(D) Distribution of amino acid distances from each mutation site to the nearest domain

boundary. Distances of 'Impact' mutations (in red) and 'Background' mutations (in grey)

were compared within Domains (Left) and within IDRs (Right). 'Impact' mutations

exhibit significantly smaller distances in both cases (****P < 0.0001, two-sided Mann–

Whitney U test).

(E) Distribution of eight pi-contact prediction values (PPVs) for mutation sites

predicted by PScore[35]. Values of 'Impact' mutations (in red) and 'Background' mutations

(in grey) were compared. The dot in each violin represents the average of values.

Mutations impacting PS exhibit a higher average frequency of pi-contact, with a

significant difference observed in six out of eight measurements of pi-contact values.

(F) Statistical comparison of the changes of amino acid property index before and after

mutation between collected 'Strengthen' (n = 79, orange) and 'Weaken/Disable' groups

(n = 228, blue) under Kolmogorov–Smirnov D test (WT, wild-type amino acid; MT,

mutant amino acid). The direction of the D statistic was set as positive when the mean

value of the 'Strengthen' group was higher and as negative when that of the 'Weaken'

group was higher. Significant differences were observed in amino acid mass (Left),

hydrophobicity (Middle) as well as polarity (Right).

1111

1112 **Figure3**

1113

1114

Fig. 3. Evaluation of methods' performance on predicting missense mutations that

impact natural phase separation (PS) ('Impact' mutation).

(A) Discriminative power evaluation of representative PS prediction methods for

'Impact' mutations against random 'Background' mutations, comparing absolute score

changes pre- and post-mutation. Each method's score difference values are represented

in paired violin plots (Left for 'Impact' mutations and right for 'Background' mutations.

Each dot within a violin indicates the median value).

(B) Performance evaluation of representative PS prediction methods on discerning

'Impact' mutations against random 'Background' mutations (IP task). AUROC is based

on the absolute score changes.

(C and D) Model performance in identifying 'Impact' mutations evaluated using leave-

one-source-out (LOSO, Left) and an independent test set (Right), measured by AUROC

(C) and AUPR (D).

(E and F) A parallel evaluation similar to (C and D) but the 'Background' mutations

were generated following the same IDRs: Domains ratio as the collected 'Impact'

samples (weighted sampling).

(G and H) A parallel evaluation similar to (C and D) but the 'Background' mutations

were generated by aligning the frequency of different mutations with their frequency in

the impact dataset (AA weighted sampling). We assigned weights to each type of

mutation based on the number of occurrences in the impact dataset, with a minimum

weight of 1 to ensure all mutation types are considered.

1136

1137

1138

 Figure4

1140

Fig. 4. Experimental validation of *Eps8* missense mutations predicted by PSMutPred

to impact PS.

(A) Representative images of overexpressed GFP-*Eps8* and its mutants in HEK293

cells (scale bars: 10 μm; n = 10 randomly picked cells). WT denotes wild type.

(B) Quantification of puncta within the wild type and mutants of *Eps8* in HEK293 cells

(****P < 0.0001 by two-tailed Student's t-test, n = 10 randomly picked cells). Error

bars represent standard deviation (s.d.).

(C) Ribbon diagram representation of mouse EPS8 structure predicted by AlphaFold2,

showing both front (left) and back (right) views.

(D-F) Detailed regions involving missense mutations with their neighboring residues

(D, F), and interaction analysis (E). The mutations are shown with the stick mode in red

while hydrogen bonds are shown as blue dashed lines. Sequence alignments within

critical residues are shown in bold.

1154

1155

1156

 Figure5

1158

Fig. 5. Evaluation of PSMutPred scores across ClinVar[64] variants.

(A-D) Pearson correlations comparison between groups. For each group, the Pearson correlation was computed by comparing the PSMutPred scores of its variants against their respective ClinVar pathogenicity labels (pathogenic or likely pathogenic coded as 1 and benign or likely benign coded as 0). P-values of Pearson correlation (*P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001; NS = no significance).

(A) Comparison between variants from the PS-prone group defined by phase separation proteins (83 known PS proteins, 1,451 variants) and low-PS-prone group variants (8,528 proteins, 84,840 variants).

(B) Comparison between variants from the predicted PS-prone group (1,276 proteins, 30,889 variants) and low-PS-prone group variants (7,335 proteins, 56,853 variants).

(C) Comparison between variants located in IDRs (n = 15,427) and Domains within the predicted PS-prone group (n = 15,462).

(D) Comparison between variants from neurodegenerative disease (ND) related

proteins (19 proteins) and variants from other proteins (non-ND) (within the computationally predicted PS-prone group), variants being predicted 'Impact' by PSMutPred-IP were considered (having a combined IP-LR, IP-SVR, and IP-RF rank score sum above 1.5). Scores of PSMutPred-SP models (n = 252) show a positive correlation with the pathogenicity of ND-related proteins, which differed from the overall pattern.

(E) Performance of PSMutPred-IP models on pathogenicity prediction of missense variants. Variants were selected from the PS-prone group defined by known PS proteins and further narrowed to those prone to locate in IDRs (IUPred3 score > 0.5 and not mapped by PfamScan, 489 variants).

(F) A parallel evaluation of (E) but focuses on variants from the PS-prone group predicted by algorithms (8,188 variants).

(G) Comparison of the proportion of variants predicted as 'Impact' by different PSMutPred-IP models between the PS-prone group and low-PS-prone group. The proportion of 'Uncertain Significance' (VUSs) IDR variants predicted as 'Impact' (PSMutPred-IP rank score > 0.8) for each protein was computed. The distributions of these proportion values are presented in a top-to-bottom sequence: IP-RF (top), IP-LR (middle), and IP-SVR (bottom). We compared the PS-prone group (83 proteins) with the low-PS-prone group (8,528 proteins) defined by known PS proteins (PS proteins, Left) and compared the PS-prone group (1,276 proteins) with the low-PS-prone group (7,335 proteins) predicted by algorithms (Predicted-PS proteins, Right). The differences were calculated based on Kolmogorov's D statistic, the direction of the D statistic was set as positive when the mean proportion of the PS-prone group was higher and as negative when that of the low-PS-prone group was higher. PS-prone proteins have a higher proportion of missense mutations predicted to 'Impact' PS.

1201 **Figure6**

1202

1203

Fig. 6. Analysis of phase separation related feature contributions to pathogenicity

prediction by combining them with other representative unsupervised prediction

metrics.

(A-E) Pathogenicity prediction performance evaluation of the model combining EVE

with PS-related features.

(A) AUROC (Left) and AUPR (Right) evaluations on the independent test set (n =

15,394). Purple line represents the model trained with both EVE and PS features; green

line represents the EVE score alone.

(B) AUROC (Left) and AUPR (Right) evaluations specifically on variants within IDRs

from the data set analyzed in (A) (n = 5,656).

(C, D) The divergence of predicted scores distributions between the standalone EVE

(green) and the combined model (purple), quantified using a two-sided Mann–Whitney

U test on the independent test set.

(C) Score distributions for pathogenic-prone variants (pathogenic and likely pathogenic,

n = 2,044, left graph) and benign-prone variants (benign and likely benign, n = 3,612,

right graph) with a focus on variants located in IDRs.

(D) A parallel evaluation of (C) but focusing on variants located in Domains (6,665

pathogenic or likely pathogenic and 3,073 benign or likely benign).

Except for a performance reduction for benign variants located in Domains, the

pathogenicity power can be significantly improved by incorporating PS features.

(E). AUROC (Left) and AUPR (Right) evaluation of IDRs variants with high

AlphaFold2 pLDDT scores (pLDDT >= 0.7, n = 2,763) and low pLDDT scores

(pLDDT < 0.5, n = 2,407).

(F-I) Pathogenicity prediction performance evaluation of the model combining ESM1b

with PS-related features.

(F) AUROC (Left) and AUPR (Right) evaluation of the model trained with ESM1b and

PS features using 5-fold cross-validation under the ClinVar dataset (n = 140,321;

Compared to EVE, ESM1b is more up-to-date and covers a broader range of proteins

and variants).

(G). AUROC (Left) and AUPR (Right) evaluation of IDRs variants with high

AlphaFold2 pLDDT scores (pLDDT >= 0.7, n = 36,032) and low pLDDT scores

(pLDDT < 0.5, n = 25,755).

(H and I) Combining PS features with ESM1b scores, pathogenicity scores for

1,015,769 ClinVar variants of uncertain significance were predicted.

1238

1239

1240