Mechanical forces pattern endocardial Notch activation via mTORC2-PKC pathway

Yunfei Mu [1,2,3,4#], Shijia Hu [1,2,3,4#], Xiangyang Liu [2,3,4], Xin Tang [2,3,4], Hongjun Shi [2,3,4*]

1. Fudan University, Shanghai, 200433, China.

2. Key Laboratory of Growth Regulation and Translational Research of Zhejiang Province, School of Life Sciences, Westlake University, Hangzhou 310024, China

3. Westlake Laboratory of Life Sciences and Biomedicine, Hangzhou, Zhejiang, China.

4. Institute of Basic Medical Sciences, Westlake Institute for Advanced Study, Hangzhou, Zhejiang, China.

[#]These authors contributed equally to this article.

*Corresponding Author:

Hongjun Shi, Key Laboratory of Growth Regulation and Translational Research of Zhejiang Province, School of Life Sciences, Westlake University. Westlake Laboratory of Life Sciences and Biomedicine. Institute of Basic Medical Sciences, Westlake Institute for Advanced Study. Hangzhou, Zhejiang, China. 310024, Zhejiang, China.

E-mail: shihongjun@westlake.edu.cn

26 **Summary**

27
Notch signaling has been identified as a key regulatory pathway in patterning the

endocardium through activation of endothelial-to-mesenchymal transition (EMT) in the

atrioventricular canal (AVC) and proximal outflow tract (OFT) region. However, the precise

mechanism underlying Notch activation remains elusive. By transiently blocking the

heartbeat of E9.5 mouse embryos, we found that Notch activation in the arterial endothelium

was dependent on its ligand Dll4, whereas the reduced expression of Dll4 in the endocardium

led to a ligand-depleted field, allowing Notch to be specifically activated in AVC and OFT by

regional increased shear stress. The strong shear stress altered the membrane lipid

microdomain structure of endocardial cells, which activated mTORC2 and PKC and

promoted Notch1 cleavage even in the absence of strong ligand stimulation. These findings

highlight the role of mechanical forces as a primary cue for endocardial patterning and

provide insights into the mechanisms underlying congenital heart diseases of endocardial

origin.

Key words:

42
43   Cardiac patterning, Endocardium, EMT, Notch, Shear stress, Mechanosensing

1. INTRODUCTION

The formation of the mammalian heart begins with a simple linear heart tube which undergoes complex morphogenic processes to ultimately reach the adult four-chamber configuration. In mice, the pattern of the adult heart becomes apparent at embryonic day (E) 9.5. At this stage, increased cell proliferation and sarcomeric development at the outer curvature of the primary heart tube lead to the formation of working chamber myocardium, while less proliferative and less differentiated myocardium at the inner curvature characterizes the outflow tract (OFT) and atrioventricular canal (AVC) [1, 2]. Between E9.5 and E10.5, endocardial cells in the OFT and AVC regions undergo extensive endothelial-to-mesenchymal transition (EMT), forming endocardial cushions. Further growth and remodeling of these endocardial cushions contribute to the septation of the OFT and cardiac chambers and give rise to cardiac valves [3]. Malformation of the endocardial cushions underlies the mechanism of the majority of human congenital heart defects (CHD).

Multiple interactive signal transduction pathways are involved in patterning valve versus chamber myocardium and regulating EMT. At E9.5, expression of BMP2 in the AVC and OFT myocardium directly activates TBX2 expression [4, 5], which, in turn, activates TGFβ2 expression [6]. Both BMP2 and TGF-β2 enhance Snail1 expression and stimulate EMT [4, 7]. Notch signaling is also required for EMT. At E9.5, NICD (Notch1 intracellular domain) expression is the highest in endocardial cells of the AVC and OFT region, while in the ventricle, NICD is more restricted to the endocardium at the base of developing trabeculae [8, 9]. Notch, through the transcriptional factor CSL, directly activates Snail2 expression, leading to the repression of VE-cadherin expression [7, 10]. Genetic abrogation of Notch signaling blocks EMT in the AVC endocardium, whereas ectopic endocardial NICD expression leads to partial EMT of ventricular endocardial cells outside AVC [10, 11].

Despite the well-characterized role of Notch in endocardial patterning and EMT, little is

known about the mechanism that specifically activates Notch in the areas of endocardium

that undergo EMT. Among various Notch receptors and ligands, Notch1, Dll4, and Jag1 are

expressed in the endocardium at the onset of EMT [10, 12]. Notch1 transcript appears to be

uniformly expressed throughout the endocardium at E9.5, while Dll4 is most concentrated in

the ventricular endocardium at the base of the trabeculae [9]. Jag1 is highly expressed

throughout the myocardium, with only a few individual AVC endocardial cells positive for

Jag1 [13, 14]. Conditional deletion of Dll4 in all endothelial cells results in the loss of classic

EMT markers and complete absence of AVC cushions at E9.5 [13]. However, it remains unclear

whether this impact on EMT is specifically due to the loss of Dll4-Notch signaling at cushion

endocardium or a consequence of global circulatory failure resulting from vascular

deformation in Dll4-deficient embryos. Therefore, the currently known expression pattern of

Notch receptors and ligands does not fully explain the specific pattern of Notch activation in

the endocardium.

Studies in zebrafish embryos have revealed the essential role of various

mechanosensitive ion channels in modulating Notch1b receptor expression in the AVC

endocardium in response to increased shear stress in this area [15-17]. However, as mentioned

earlier, in the developing mouse endocardium at the onset of EMT, Notch1 is uniformly

expressed throughout the endocardium despite restricted NICD in AVC and OFT

endocardium. Thus, the equivalent mechanosensitive pathways controlling region-specific

pattern of Notch activation, and hence EMT, in the mammalian endocardium are still unclear.

To address this question, we transiently blocked the heartbeat of mouse embryos at E9.5 *in*

vivo using the class III antiarrhythmic drug dofetilide, which selectively blocks the rapid

component of the delayed rectifier K^+ current (I_{Kr}) in cardiomyocytes [18]. We found that Notch

activation in the vascular endothelium is dependent on Dll4, whereas in the endocardium, the overall reduction of Dll4 expression creates a ligand-depleted field that allows establishment of a specific Notch activation pattern in the AVC and OFT regions in response to increased shear stress. Strong shear stress in these valve-forming areas alters the membrane lipid microdomain structure of the endocardial cells, which then activates mTORC2 and PKC, promoting Notch1 cleavage even in the absence of strong ligand stimulation. The results uncover the mechanical force as a primary cue for endocardial patterning in mammals and may provide new ways to explain many congenital heart diseases of endocardial origin.

2. RESULTS

2.1. Notch is strongly activated in the AVC and proximal OFT endocardium in spite of weak ligand expression at the onset of EMT

Although Notch1, Dll4, and Jag1 have been reported as the principal receptor and ligands expressed in the endocardium [12, 13, 19], the relationship between the Notch activity and its ligand expression in various parts of the E8.5-9.5 endocardium has not been examined in sufficient details. To investigate whether regional-specific Notch activation is due to differential ligand expression, we performed whole-mount *in situ* hybridization and immunofluorescence staining of NICD, Dll4, and Jag1. Both *Dll4* transcripts and NICD were uniformly positive throughout all endothelial lining of the dorsal aorta and heart at E8.5 (Figure 1-figure supplement 1A, Video 1, 3). However, at E9.5, when endocardial EMT starts, distinct patterns of NICD and Dll4 expression appeared (Figure 1 and Video 2, 4). Based on the expression pattern of NICD and Dll4, overall, the endocardium at E9.5 can be viewed as three distinct types as summarized in Figure 1. Type I is Dll4-high and NICD-high, including the endocardium lining the distal OFT, the base of the trabeculae, the dorsal wall of the left atrium, and the right atrium. These endocardial cells normally do not undergo EMT. Type II is Dll4-low and NICD-high, including the AVC and proximal OFT endocardium. Endocardial cells in these regions undergo EMT to form endocardial cushion cells. Type III is Dll4-low and NICD-low, including the endocardium flanking the AVC and on the top of ventricular trabeculae. Jag1 was highly expressed in the myocardium but expressed at a very low level in a small subset of the OFT and AVC endocardial cells, consistent with the previous report (Figure 1-figure supplement 1A) [13]. Vascular endothelium continued to express high levels of Dll4 and NICD uniformly, similar to type I endocardial cells. Dll4 protein staining pattern agreed with the *Dll4* transcript and also agreed with VEGF receptor-2 (Flk1/KDR) protein expression

throughout the cardiovascular endothelium at both E8.5 and E9.5 (Figure 1-figure

supplement 1B), consistent with the regulatory role of VEGF signaling in Dll4 expression [20].

Thus, the Dll4 and NICD expression patterns were disconnected in the type II endocardium

at E9.5, suggesting the existence of additional mechanisms for Notch activation.

One previous study showed that endocardial EMT was dependent on Dll4 [13]. To further

dissect the role of Dll4 in regional Notch activation, we conditionally deleted Dll4 in all

endothelial cells using the *Tie2^cre* line. At E9.5, *Dll4* conditional null AVC cushions were

poorly cellularized, consistent with the previous finding. Furthermore, in most areas normally

expressing high Dll4, i.e., vascular endothelium and aortic sac, NICD was completely

abolished, whereas in the AVC, proximal OFT, and the base of the trabeculae, NICD was

reduced but not completely absent (Figure 1-figure supplement 1C). As conditional deletion of

Dll4 resulted in embryos with severely disorganized vasculature (Figure 1-figure supplement

1C), we cannot rule out the possibility that the reduced NICD in the AVC might not be simply

due to Dll4 deletion, but rather affected by circulatory failure. These findings indicate that

vascular endothelium is dependent on Dll4 for Notch activation, whereas in the endocardium,

additional factors are required to pattern Notch activation in the proximal OFT and AVC.

2.2. Notch activation in cushion endocardium is dependent on blood flow

Past work on zebrafish embryos showed that cardiac contraction activates endocardial

Notch signaling[21]. To test if Notch activation is regulated by blood flow in the developing

mouse heart, pregnant mice were gavaged with a single dose of dofetilide at 2mg/kg at E9.5.

The embryos were then either harvested immediately after the treatment or allowed to

develop until E18.5 when cardiac morphology was assessed (Figure 2A). Dofetilide caused

an immediate stop of blood flow in the embryos, as observed by Doppler ultrasound from 1 h

through to 3 h, which completely recovered at 5 h post-treatment (Figure 2B and Video 5, 6).

Three hours cessation of blood flow caused a complete loss of NICD without affecting total

Notch1 receptor protein in the proximal OFT and AVC endocardium. The loss of NICD was completely recovered at 5 h post-treatment, in line with the dynamics of the heart rate changes (Figure 2C). No significant changes in the expressions of Dll4 and Jag1 were noted in AVC endocardium after dofetilide treatment (Figure 2-figure supplement 1A). Pro-EMT markers phospho- Smad1/5, Sox9 and Twist1 were downregulated in the AVC endocardium after cardiac arrest (Figure 2C). Interestingly, NICD in the dorsal aorta was resistant to the cessation of flow.

Consistent with the inhibition of EMT, transient cessation of blood flow resulted in severely hypoplastic OFT and AVC endocardial cushions one day after treatment (Figure 2D), and various heart defects (Figure 2E and Figure 2-figure supplement 1B) in 40% of embryos: ventricular septal defects (VSD), bicuspid semilunar valve, atrioventricular valve defects, and conotruncal defects, consistent with malformation of endocardial cushions.

To rule out the possible direct effect of dofetilide on endocardial cells independent of flow, we conditionally deleted *Tnnt2* in embryonic myocardium, which caused bradycardia in E9.5 embryos and lethality at E10.5 (Video 7). Similar to the effect of dofetilide, reducing blood flow rate by genetic means also caused a significant reduction of NICD in the cushion endocardium at E9.5 (Figure 2-figure supplement 1C). Furthermore, block of heartbeat by *ex vivo* blebbistatin treatment, an inhibitor for non-muscle myosin II ATPase, also prevented Notch cleavage in the cushion endocardium at E9.5 (Figure 2-figure supplement 1D). To rule out the possible effect of hypoxia on Notch activation, we depleted embryonic erythrocytes by crossing Epor[cre] with ROSA-DTA mouse line, which resulted in widespread hypoxia without interfering embryonic heartbeat, yet NICD was normal in the AVC endocardium (Figure 2-figure supplement 1E). We also cultured E9.5 embryos in a medium saturated with 95% O_2 in the presence of dofetilide for 3 h. High levels of O_2 eliminated HIF-1α nuclear staining induced by cardiac arrest,

178   while NICD was still absent in the AVC endocardium (Figure 2-figure supplement 1F).

179   Thus, all subsequent *ex vivo* embryo culture experiments were performed in 95% O_2 to

180   minimize the effect of hypoxia on Notch activation. Collectively, these results indicate

181   that Notch activation and EMT in the cushion endocardium are dependent on the

182   mechanical stimulation from blood flow.

2.3. Flow-responsive mTORC2-PKCε activity is required for Notch activation in the

cushion endocardium

Given the fast response of NICD to flow cessation, we suspect a change in post-

translational modification, such as a phosphorylation event, that might mediate the effect.

PKC, AKT, and ERK have all been reported to have their phosphorylation status altered in

response to fluid shear stress in cultured endothelial cells [22]. In addition, *in vitro* shear stress-

induced Notch activation can be blocked by inhibitors of PKC, AKT, and ERK [23]. Therefore,

we examined the phosphorylation status of these three signaling molecules in the

endocardium before and after dofetilide treatment. Both pPKC ($βII^{Ser660}$) and $pAKT^{Ser473}$ were

restricted to the cushion endocardium and the base of the trabeculae in control embryos and

were almost completely lost as quickly as 1 h after treatment and then both recovered at 5 h

after treatment. Both pPKC and pAKT were not detectable in dorsal aorta endothelium

(Figure 3A and Figure 3-figure supplement 1A). Their response rate in cushion endocardium

was faster than NICD, whose maximum inhibition occurred at 3 h post-treatment (Figure 2C).

pERK was not detectable in the cushion endocardium (Figure 3-figure supplement 1B).

Inhibition of AKT phosphorylation in cultured E9.5 embryos by wortmannin did not inhibit

Notch activation (Figure 3-figure supplement 1C), whereas inhibition of PKC activity by

staurosporine treatment blocked Notch activation (Figure 3-figure supplement 1D),

To further confirm the role of PKC in regulating Notch, we individually knocked out

Prkce and *Prkch* (Figure 3-figure supplement 1E). Among all PKC family members, these

two isozymes are most highly expressed in the endocardium based on RNAseq analysis of

E9.5 embryos' endocardial and vascular endothelial cells (unpublished data). *Prkch*[KO] did not

affect Notch activation or produce any phenotype. *Prkce*[KO] caused a slight but significant

reduction of NICD in the cushion endocardium at E9.5 and caused 25% heart defects at

E18.5.

Prkce/Prkch double knockout resulted in a further loss of NICD at E9.5, leading to pericardial

effusion and complete lethality at E10.5. (Figure 3B and Figure 3-figure supplement 1F). No

pericardial effusion or heartbeating abnormalities were noted at E9.5. Treatment of pregnant

mice at E9.5 with a potent PKC activator, phorbol 12-myristate 13-acetate (PMA), almost

completely rescued dofetilide-induced loss of NICD, pPKC, cushion cellularity, and cardiac

defects in 3 h (Figure 3C-3E). Thus, PKC activity is both necessary and sufficient for Notch

activation in the cushion endocardium.

Both PKC and AKT belong to the AGC kinase family and both have a conserved

hydrophobic motif at the C-terminal tail of the catalytic domain, whose phosphorylation is

dependent on mTORC2 and is required for kinase activity [24]. Therefore, the observed

inhibition of hydrophobic motif phosphorylation of PKC and AKT indicates impairment of

the common upstream activator mTORC2. We conditionally deleted one of the essential

mTORC2 components, Rictor[25], in the endothelial cells (Figure 3-figure supplement 1E). As

expected, ablation of mTORC2 completely blocked hydrophobic motif phosphorylation of

both PKC and AKT and consequently led to a significant decrease of NICD in the cushion

endocardium (Figure 3F). Thus, blood flow activates Notch in the cushion endocardium via

mTORC2-PKC signaling pathway.

2.4. Shear stress-induced alteration of membrane lipid microstructure activates

mTORC2-PKC-Notch signaling pathway

In cultured mammalian endothelial cells, fluid shear stress (FSS) increases the numbers

of cell surface caveolae, a specialized membrane microdomain enriched for cholesterol and

sphingolipids; sequestration of cholesterol inhibits shear-dependent activation of ERK [26].

Thus, we tried to manipulate the membrane lipid microdomain by treating the cultured E9.5

embryos with cholesterol. Staining of Caveolin-1, the major component of caveolae, showed

that Caveolin-1 was normally expressed on the cell surface of the AVC endocardial cells

including the luminal surface and the lateral cell adhesion sites. However, in areas with low

shear stress such as dorsal aorta endothelium, atrial, and ventricular endocardium

downstream of AVC, Caveolin-1 was mainly restricted to the lateral cell adhesion sites and

absent on the luminal surface. *Ex vivo* dofetilide treatment caused retraction of Caveolin-1

from the luminal surface to the lateral cell adhesion sites in the AVC endocardial cells, while

co-treatment with cholesterol rescued the presence of Caveolin-1 to the luminal surface of

AVC endocardial cells in the presence of dofetilide (Figure 4A). The loss of NICD,

$pPKC^{Ser660}$, and $pAKT^{Ser473}$ caused by dofetilide could all be rescued by pretreatment with

cholesterol, indicating reactivation of mTORC2 (Figure 4B). These rescuing effects of

cholesterol disappeared in the Rictor KO embryos, suggesting mTORC2 was an intermediate

signaling molecule linking membrane lipid microstructure and Notch activation.

2.5. Pharmacogenetic interaction in etiology of congenital heart defects

245
Gene-environmental interactions are believed to underlie the complex etiology of many

congenital heart diseases. With the mechanosensitive nature of Notch in mind, we speculated

that reducing the gene dosage of Notch1 may interact with agents that disrupt normal

hemodynamic stimulations to the endocardium and synergistically cause heart defects. To test

this, we crossed Notch1 heterozygous null male (FVB) with wildtype female mice, and

treated the pregnant mice with a lower dose of dofetilide (1.8 mg/kg) at E9.5. In addition, we

tested another drug verapamil, a commonly used FDA-approved L-type calcium channel

blocker for treatment of high blood pressure, heart arrhythmias, and angina[27], in FVB

background. Neither Notch1 heterozygosity alone, nor drug treatment alone at the applied

dosage produced any notable heart defects. However, the combination of Notch1 mutation

and drug exposure of either dofetilide or verapamil resulted in over 50% of fetuses having

various heart defects of endocardial origin (Figure 5A, B).

3. DISCUSSION

In this study, we identified fluid shear stress as the primary activator of Notch signaling in the area of AVC and OFT of the mouse looping heart tube. Strong fluid shear stress in the AVC and OFT promotes formation of caveolae on the luminal surface of the endocardial cells, which enhances PKCε phosphorylation by mTORC2. Activated PKC then augments Notch cleavage under the minimal ligand stimulation, resulting in Notch activation and EMT specifically in areas of valve primordium characteristic of narrow internal diameter and high shear stress. These findings are directly relevant to endocardial patterning. Notch activity is known to be essential for establishing OFT and AVC endocardial cell identity capable of EMT [10, 11]. Activation of Notch in the AVC endocardium is presumed to be mainly driven by the Dll4 ligand because Dll4 has been demonstrated to be expressed in the AVC endocardial cells and endothelial-specific knockout of *Dll4* produced acellularized AV cushions [13]. However, careful examination of the immunostaining pattern revealed that NICD, VEGFR2 and Dll4 were initially expressed with equal levels throughout cardiovascular endothelium at E8.5 prior to endocardial EMT, but at E9.5 with the onset of EMT, endocardial VEGFR2 and Dll4 were markedly reduced and NICD becomes restricted to endocardium overlaying the merging OFT and AVC cushions, implying a transition of the main driver of Notch activation from the pan-endothelial VEGFR2-Dll4 stimulation at E8.5 to the regionally restricted shear force stimulation at E9.5.

We therefore propose the following working model as how mechanical cues guide Notch activation during endocardial patterning. Following gastrulation, high level of VEGF-VEGFR2 signaling in mesodermal tissues initiates Dll4 expression which activates Notch and drives arterial endothelium and endocardium differentiation and proliferation. At E9.5 when endocardial EMT starts, VEGF signaling must dampen due to its suppressive function on

EMT [28, 29]. One way to lower VEGF signaling in the endocardium is through downregulating

VEGFR2 expression. Consequently, Dll4 expression also decreases, rendering Notch less

active in the endocardium compared to the vascular endothelium. In the meantime, cardiac

ballooning on both sides of the AVC expands the myocardial chambers and widens the

endocardial internal diameter, leaving only the endocardium of the OFT and AVC endocardial

cushion region closely attached. Narrow blood passage way creates strong shear stress to

stimulate Notch cleavage only in these narrow areas of future cardiac septation and valve

formation. Thus, through these sophisticated mechanisms, the developing mouse hearts may

achieve three purposes: (1) reducing VEGF signaling to permit endocardial EMT; (2)

reducing Dll4 to prevent widespread endocardial Notch activation and make endocardium

sensitive to flow; (3) maintaining a proper cushion size and shape by limiting the flanking

endocardium to undergo EMT despite physically close to the field of BMP2 derived from of

AVC myocardium (Figure 6).

Studies in zebrafish identified several cation channels including Trpv4, Trpp2, Piezo1,

Piezo2 and P2X that contribute to mechanosensing in the developing endocardium [15, 17, 30].

Blood flow in the looping mammalian heart is predominantly unidirectional whereas early

zebrafish embryonic heart displays oscillatory flow pattern [17]. Mammalian embryonic

endocardium undergoes extensive EMT to form valve primordia while zebrafish valves are

primarily the product of endocardial infolding [15]. It is thus possible that mammalian hearts

may evolve additional mechanisms to guide endocardial patterning and development. Our

data provide evidence to support that membrane lipid microdomain acts as a shear stress

sensor and transduces the mechanical cue to activate intracellular mTORC2-PKC-Notch

signaling pathway in the developing endocardium. Shear stress has been shown to upregulate

the number of caveolae in cultured endocardial cells [26]. Flow induced AKT phosphorylation

is dependent on Caveolin-1 [31]. Both PKCα and PKCε interact with Caveolin-1 and bind with

caveolae membranes [32, 33]. In neuronal cells or neural tissues, accumulation of psychosine in

the plasma membrane disrupted lipid rafts, blocked recruitment of mTORC2 and PKC to the

lipid raft, and inhibited AKT and PKC activation [34, 35]. In line with these previous findings,

we demonstrated that high level of shear stress in the AVC and OFT is required for caveolin

localization to the luminal surface of the endocardial cell membrane, and this membrane lipid

microstructure is both necessary and sufficient for mTORC2-PKC-Notch pathway activation.

The mechanism by which shear stress increases caveolae is unknown. In a previous study,

shear stress exposure for a few minutes rapidly decreases the lipid order and increases the

fluidity of the plasma membranes which appears to contradict our findings [36]. However,

membrane lipid compositions are highly dynamic and regulated. It is possible that acute shear

stress and its associated kinetic energy may initially decrease membrane lipid order, but long-

term shear may evoke cellular feedback mechanisms to increase lipid order and enhance

membrane rigidity.

The positive regulation of Notch activation by PKC has been reported in a number of *in

vitro* systems. Studies in CD4^{+} T cells revealed that Notch is activated within hours of TCR

stimulation independent of Notch ligands and PKC activity is both sufficient and necessary

for the ligand independent Notch activation [37]. The exact mechanism by which PKC regulates

Notch processing is not known, and in need of further investigation in the future.

The vast majority of congenital heart diseases do not have a genetic explanation and are

considered to have a multifactorial origin. Studies conducted on various model organisms

such as chick and zebrafish have demonstrated that alterations in blood flow during early

stages of heart development can lead to heart malformation [38]. Similarly, human studies have

shown that fetal bradycardia or abnormal flow patterns in the Ductus Venosus during the first

trimester of pregnancy are associated with an increased risk of CHD [39, 40]. Our research

findings suggest that abnormal heart contraction or flow patterns, resulting from genetic

mutations or the use of certain drugs during early heart development, may interact with

genetic pathways involved in mechanosensing and endocardial EMT. These interactions

336   could contribute to the complex etiology of congenital heart disease.

4.   MATERIALS AND METHODS

4.1. Reagents

343   A list of the reagents used in this study is provided in Resource Table.

4.2. Animals

All mice used in this study were housed at a constant temperature (23°C) and humidity

(~50%) with a 12-h light/dark cycle and ad libitum access to food and water. All animal

experiments were performed in accordance with the protocol approved by the Westlake

University Institutional Animal Care and Use Committee (approval # 21-007-SHJ). Noon of

the day of vaginal plug detection was defined as embryonic day (E) 0.5. For dofetilide

treatment, mice were gavaged at E9.5 with a single dose of dofetilide at 2 mg/kg dissolved in

saline. For phorbol 12-myristate 13-acetate (PMA, 2 mg/kg) administration, mice were

gavaged at E9.5 with a single dose of PMA dissolved in CMC-Na. Verapamil was given by

intraperitoneal injection at E9.5. At the indicated date post-treatment, the dams were

sacrificed by cervical dislocation, and the embryos were harvested for further analysis. All

sources of our mouse lines are listed in Resource Table. Knockout mice generated in-house

are described and validated in Figure 3-figure supplement 1E.

4.3. Fetal echocardiography

Ultrasound scanning was performed using the Vevo 300 high-frequency ultrasound

machine (FUJI FILM Visual Sonics Inc., Canada). *In utero* echocardiography of fetal mice

was conducted at E9.5. Transducer MX-550D had a central frequency of 40 MHz with axial

and lateral resolutions both about 30 μm. The pregnant mice were sedated using 2%

isoflurane and kept at 1% isoflurane to maintain body temperature at 35.5-37°C and heart rate

at 400 to 500 beats per minute (bpm). Pulse-wave Doppler was used to measure velocity and

time interval parameters to measure the blood flow velocity and time interval parameters of

the fetal hearts. All fetuses were scanned twice in order and their *in utero* positions were

noted. For embryos of *cTnT^{cre}* x *Tnnt2$^{flox/flox}$*, after completing the Doppler ultrasound

imaging, the embryos were dissected out with the recorded positions, and embryonic tissues

were collected for genotype identification to correlate with the heart rate data. During the

measurements, observers were blinded to the assigned groups. Analysis was performed with

Vevo Lab 5.7.1.

4.4. Fetal heart morphology

375
Fetal hearts were imaged using the Zeiss Lightsheet Z.1 microscope. Embryos were

harvested at E18.5, and hearts were dissected in phosphate-buffered saline (PBS), fixed

overnight in a mixed solution of 10% neutral buffered formalin and 2.5% glutaraldehyde,

then rinsed twice in PBS, and dehydrated in 50, 75 and 100% ethanol for 30 mins each at

room temperature (RT). Samples were then transferred into a specially designed glass tube

containing 100 μL of BABB solution (1:2 benzyl alcohol: benzyl benzoate) and incubated for

30 mins to clear the sample. The glass tube was mounted into the sample chamber which was

filled with 87% glycerol (RI~1.45). Hearts were scanned from the apex to the great arteries

for tissue autofluorescence using the 561 nm laser line and the detection optics 5x/0.16. 3D

reconstruction of the image stacks and morphological analyses were performed with Imaris

9.3 software.

387
388 **4.5. *Ex vivo* embryo culture**

389
Mouse embryos at E9.5 were carefully dissected in Hanks Buffer containing calcium

and magnesium to remove decidua without damaging their yolk sacs and placentas. They

were then cultured in 1 mL embryo culture medium (50% rat serum, 50% DMEM), pre-

saturated with a gas mix of 95% O_2 and 5% CO_2 in 5 mL volume glass tubes. Tubes were

sealed and placed in a rotatory shaker maintained at 37 ℃ and 50 rpm. For pharmacological

treatment, stauroporine (100 nM), wortmannin treatment (2 μM), dofetilide (0.2 ug/ml), or

water-soluble cholesterol (1 mg/ml) was diluted in embryo culture medium to the desired

concentrations.

4.6. Immunofluorescence and *in situ* hybridization

399
E9.5 and E10.5 wild embryos were dissected in PBS, fixed in 4% paraformaldehyde

(PFA) overnight at 4 °C, dehydrated in ethanol series, cleared in xylene, and embedded in

paraffin. Tissues were sectioned at 6 μm thickness. Paraffin sections were subjected to a 20

mins heat-induced antigen retrieval using a citric acid solution (pH 6). The sections were

blocked with 5% normal donkey serum (NDS) in Tris-buffered saline with 1% Tween-20

(TBST) for 1 hour, followed by overnight incubation at 4°C with primary antibodies diluted

in the blocking buffer. Bound primary antibodies were detected using fluorescently

conjugated corresponding secondary antibodies. For detection of low-abundance targets such

as NICD and phospho-PKC^{Ser660}, tyramide signal amplification (TSA) was applied. A list of

the antibodies used in this study is provided in Resource Table. Confocal microscopy images

were captured on a Zeiss LSM 800 and image analysis was performed using Zen 2.3.

Fluorescent intensity was calculated using ImageJ software.

To analyze NICD expression in E8.5, E9.5 mouse hearts, whole mouse embryos were

fixed in 4% PFA at 4°C overnight. After blocking in 5% NDS, 0.5% Triton X-100 in TBS at

4°C overnight, samples were incubated at 4°C overnight with the primary NICD antibody

diluted in blocking buffer (1:400). After washes, embryos were incubated with donkey anti

rabbit HRP (1:500) 4°C overnight. Embryos were then washed and reacted with a tyramide-

biotin solution at RT for 30 minutes. After that, samples were washed and incubated with

Cy^{TM}3 streptavidin (1:500). Stained embryos were dehydrated in methanol, cleared in BABB

and imaged by Zeiss LSM 800.

Whole mount in situ hybridization of *Dll4* was performed using the HCR probe set

according to the manufacturer's instruction (Molecular Instruments, Los Angeles). Stained

embryos were dehydrated in methanol, cleared in BABB and imaged by Zeiss LSM 800.

4.7 Statistics

424
425   Difference in expression levels was tested using two-tailed t test. Two-sided Fisher's exact

426   test was used to compare the heart defect rates between two groups. P value <0.05 was

427   considered significant. All statistical analyses were performed using GraphPad Prism 8

428   software.

      5.   FIGURE LEGENDS

   Figure 1: Endocardium and vascular endothelium of dorsal aorta showed different

   patterns of Notch1 activation and Dll4 ligand expression.

   Type I cells are NICD-high and Dll4-high. Type II cells are NICD-high and Dll4-low. Type III

   cells are NICD-low and Dll4- low. Scale bars, 100 μm.

   Figure 1— Figure supplement 1: Notch is strongly activated in the AVC and proximal

   OFT endocardium despite weak ligand expression at the onset of EMT.

   (**A**) Representative images of NICD (green) whole mount immunofluorescence, *Dll4* (red)

   whole mount *in situ* hybridization, and Jag1 (yellow) immunofluorescence of E8.5 and E9.5

   mouse hearts. Arrows indicate arterial endothelium and arrowheads indicate endocardium. (**B**)

   Representative images demonstrating VEGFR2 (Green) and Dll4 (red) protein expression in

   E8.5 and E9.5 mouse arteries (arrow) and endocardium (arrowhead). (**C**) Representative

   images showing Dll4 protein and NICD in wild-type and endothelial-specific *Dll4*-deleted

   (*Tie2^{cre}; Dll4$^{flox/flox}$*) E9.5 mouse hearts. Scale bars, 100 μm.

   Figure 2: Notch activation in cushion endocardium is dependent on blood flow.

   (**A**) Schematic diagram of the experimental design. (**B**) Echocardiography of control and

   dofetilide treated embryos. (0 h: n = 12; 1 h: n = 15; 3 h: n = 12; 5 h: n = 9). (**C**) Expression of

   NICD, total Notch1, p-SMAD1/5 and Sox9, Twist1 in the E9.5 dorsal aorta endothelium

   (arrow) and endocardium (arrowhead). (**D**) Sagittal E10.5 hematoxylin-eosin stained sections

   demonstrated hypocellularity in both superior (arrowhead) and inferior AV cushion (arrow)

   caused by dofetilide treatment. Quantification of mesenchymal cell density in superior (below

   left) and inferior (below right) AV cushion (Control: n = 16 embryos; Dofetilide: n = 14

   embryos). OFT, outflow tract; A, atrium; V, ventricle. (**E**) Representative heart defects induced

   by maternal dofetilide treatment. pmVSD, perimembranous ventricular septal defect; DORV,

double-outlet right ventricle; mVSD, muscular VSD; OA, overriding aorta; BAV, bicuspid

aortic valve; AVSD, atrioventricular septal defect; BPV, bicuspid pulmonary valve; ra, right

atrium; la, left atrium; ao, aorta; rv, right ventricle; lv, left ventricle; s, interventricular septum;

pa, pulmonary artery. Scale bars, 100 μm (**C**, **D**), 500 μm (**E**).

Figure 2—Figure supplement 1: (**A**) Representative figures and statistics of Dll4 and Jag1

expression in AVC endocardium after dofetilide treatment. (**B**) Heart defects induced by maternal

dofetilide treatment. (0 mg/kg: n = 37; 2.0 mg/kg: n =117). pmVSD, perimembranous ventricular

septal defect; DORV, double-outlet right ventricle; mVSD, muscular VSD; OA, overriding aorta;

BAV, bicuspid aortic valve; AVSD, atrioventricular septal defect; BPV, bicuspid pulmonary

valve. (**C**) Representative images of NICD in myocardial-specific cardiac troponin T knockout

mouse heart. (**D**) Representative images of NICD expression in AV canal, proximal OFT and

dorsal aorta after blebbistatin treatment (5 μM) for 3 h. (**E**) HIF-1α and NICD in red blood cell

depleted E9.5 mouse hearts. (**F**) Representative images of HIF-1α and NICD in E9.5 control,

dofetilide-treated, and hyperoxia-treated mouse hearts. Arrow: dorsal aorta endothelium.

Arrowhead: endocardium. Scale bars, 100 μm.

Figure 3: Flow-responsive mTORC2-PKCε activity is required for Notch activation in the

cushion endocardium.

(**A**) Phospho-PKC^{Ser660} and phospho-AKT^{Ser473} levels in the atrioventricular (AV) canal

endocardium (arrowhead) after dofetilide treatment. (**B**) NICD expression in AV canal

endocardium in *Prkce* and *Prkch* double heterozygous, single knockout and double knockout

embryos. Each point in the quantification chart represents one embryo. (**C**) NICD, p-PKC^{Ser660}

and p-AKT^{Ser473} staining in AVC endocardium (arrowhead) after dofetilide treatment and after

rescue by PMA. (**D**) Sagittal E10.5 HE staining sections demonstrate acellularized superior

(arrowhead) and inferior AV cushion (arrow) caused by dofetilide which were rescued by PMA

treatment (2 mg/kg). OFT, outflow tract; A, atrium; V, ventricle. Mesenchymal cell density was

quantitated in superior and inferior AV cushions (Control: $n = 14$ embryos; Dofetilide: $n = 11$

embryos; Dofetilide + PMA: $n = 9$ embryos). (**E**) Heart defect rate caused by maternal dofetilide

and PMA treatment (Control: $n = 37$; Dofetilide: $n = 62$; Dofetilide + PMA: $n = 27$). (**F**) NICD,

phospho-PKC^{Ser660}, and phospho- AKT^{Ser473} in AV canal endocardium (arrowhead) in wild-type

and *Rictor* null mice (*Rictor*$^{+/+}$: $n = 6$; *Rictor*$^{-/-}$: $n = 8$). Scale bars, 100 μm.

Figure 3—Figure supplement 1: (**A**) pPKC^{Ser660} in E9.5 mouse proximal OFT and dorsal

aorta, and pAKT^{Ser473} in E9.5 dorsal aorta endothelium. pPKC^{Ser660} expression in proximal OFT

was quantified. (**B**) pERK1/2 was not detectable in AV canal endocardium. (**C**) NICD, pPKC

and pAKT expression in cultured E9.5 heart in response to *ex vivo* wortmannin treatment (2

μM) for 3 h. (**D**) NICD, pPKC and pAKT expression in cultured E9.5 heart in response to *ex*

vivo staurosporine treatment (100 nM) for 3 h. (**E**) *Prkch*, *Prkce* and *Rictor* knockout mice

strategies and western blot or immunofluorescence confirmation of KO. (**F**) Counts of abnormal

hearts of the descendants of *Prkch*, *Prkce* double heterozygous intercrosses at E18.5. Arrow:

dorsal aorta endothelium. Arrowhead: endocardium. Scale bars, 100 μm.

Figure 4: Shear stress-induced alteration of membrane lipid microstructure activated

mTORC2-PKC-Notch signaling pathway.

(**A**) Caveolin-1 (Cav-1) and VE-cadherin expression in mouse E9.5 AV canal, dorsal aorta,

atrium and ventricle endocardium, demonstrating luminal (arrowhead) and lateral (arrow)

surface localization. *Ex vivo* dofetilide treatment (0.2 μg/ml) of cultured E9.5 embryos for 3 h

caused retraction of Caveolin-1 and VE-cadherin from luminal surface of the AVC

endocardial cells to the lateral cell adhesion sites which could be rescued by co-treatment

with cholesterol (Chol, 1 mg/ml). (**B**) Loss of NICD, phospho-PKC^{Ser660}, and phospho-

AKT^{Ser473} in AVC endocardium (arrowhead) by *ex vivo* dofetilide treatment could be

rescued by cholesterol. The rescue failed in *Rictor* null hearts. Scale bars, 100 μm (**A**, **B**).

Scale bars, 10 μm (**A**, zoom-in).

Figure 5: Pharmacogenetic interaction causing heart defects.

(**A**) Heart defect rate significantly increased in Notch1 heterozygous embryos treated with

dofetilide (1.8 mg/kg) at E9.5. (Saline *Notch1*$^{+/+}$: $n = 48$, Saline *Notch1*$^{+/-}$: $n = 37$, Dofetilide

Notch1$^{+/+}$: $n = 31$, Dofetilide *Notch1*$^{+/-}$: $n = 16$). (**B**) Types of heart defects in the dofetilide and

Notch1$^{+/-}$ combined group. (**C**) Heart defect rate significantly increased in Notch1 heterozygous

embryos treated with Verapamil (15 mg/kg) at E9.5. (Saline *Notch1*$^{+/+}$: $n = 18$, Saline *Notch1*$^{+/-}$:

$n = 9$, Verapamil *Notch1*$^{+/+}$: $n = 8$, Verapamil *Notch1*$^{+/-}$: $n = 12$). (**D**) Types of heart defects in

the verapamil combined with *Notch1*$^{+/-}$ group.

Figure 6. Working model of the establishment of Notch activation pattern by mechanical

cues. The establishment of Notch activation pattern by mechanical cues involves a series of

events in the developing heart tube. At E8.5, the arterial endothelium and non- EMT

endocardium exhibit low shear stress, high VEGF, Dll4, and Notch signaling. One day later, the

endocardium undergoes patterning and becomes capable of epithelial-to- mesenchymal

transition (EMT) only in the AVC and proximal OFT regions. This patterning is achieved

through restricted expression of BMP2 and NICD in these specific areas. EMT requires the

downregulation of VEGF signaling by the endocardium, enabling EMT to occur. Additionally,

Dll4 is downregulated in endocardium to prevent widespread Notch activation. Simultaneously,

the high shear stress present in the AVC and proximal OFT regions leads to increased membrane

lipid order, which activates the mTORC2-PKC-Notch pathway and promotes EMT. On the other

hand, regions flanking these valve-forming areas experience lower shear stress, resulting in

inactive Notch signaling and an inability to undergo EMT.

6. ACKNOWLEDGEMENTS

The authors are grateful to Dr. Xu Li of Westlake University for valuable suggestions regarding Notch experiments. We thank Dr. Zhen Zhang from Shanghai Jiao Tong University, Dr. Bing Zhang, Dr. Jiemin Jia, Dr. Shang Cai from Westlake University and Dr Hui Zhang from Shanghai Tech University for sharing mouse lines. We thank the Microscopy Core Facility of Westlake University for assistance and advice. We thank the Westlake Animal Facility and Youshi Chen for mouse breeding and husbandry. We thank Jianfeng Wang for assistance in *ex vivo* culture. We appreciate help from Jiayi Lin concerning fetal echocardiography. This work was supported by Natural Science Foundation of Zhejiang Province of China (LZ19H040001) and Westlake Education Foundation.

7. AUTHOR CONTRIBUTIONS

H.S. raised hypothesis, designed experiments, interpreted data and drafted the work. Y.M., and S.H. conducted the main body of the experiments, including mouse crosses and treatment, embryo harvest, staining and data analysis. X.L. made contributions by optimizing light sheet microscopy and performed work related to heart phenotyping. S.H. participated in manuscript drafting. X.T. characterized the change of membrane lipid structure induced by flow.

8. DECLARATION OF INTERESTS

The authors declare no competing interests.

9. Supplemental files

Video 1: E8.5 NICD whole mount staining. NICD (green) whole-mount immunofluorescence staining of E8.5 mouse heart.

Video 2: E9.5 NICD whole mount staining. NICD (green) whole-mount immunofluorescence staining of E9.5 mouse heart.

Video 3: E8.5 Dll4 *in situ* hybridization. Whole-mount *Dll4* (red) *in situ* hybridization staining of E8.5 mouse heart.

Video 4: E9.5 Dll4 *in situ* hybridization. Whole-mount *Dll4* (red) *in situ* hybridization staining of E9.5 mouse heart.

Video 5: E9.5 echo of a beating embryonic heart. Echocardiography of a normal E9.5 mouse embryo *in utero* using color Doppler.

Video 6: E9.5 echo of a dofetilide treated embryonic heart. Echocardiography of a 1-hour after dofetilide treated E9.5 mouse embryo *in utero* using color Doppler.

Video 7: E9.5 echo of *cTnT^{cre}* x *Tnnt2$^{flox/flox}$* beating with nonbeating. Echocardiography of two E9.5 embryos from *cTnT^{cre}; Tnnt2$^{flox/+}$* x *Tnnt2$^{flox/flox}$* crossing. One embryo was genotyped to be *Tnnt2$^{flox/flox}$* and the other *cTnT^{cre}; Tnnt2$^{flox/flox}$*.

REFERENCES

1. Moorman AF and Christoffels VM. Cardiac chamber formation: development, genes, and evolution. *Physiological reviews*. 2003;83:1223-67.

2. Miquerol L and Kelly RG. Organogenesis of the vertebrate heart. *Wiley Interdisciplinary Reviews: Developmental Biology*. 2013;2:17-29.

3. Leckband DE, Le Duc Q, Wang N and De Rooij J. Mechanotransduction at cadherin-mediated adhesions. *Current Opinion in Cell Biology*. 2011;23:523-530.

4. Ma L, Lu M-F, Schwartz RJ and Martin JF. Bmp2 is essential for cardiac cushion epithelial-mesenchymal transition and myocardial patterning. *Development*. 2005;132:5601-5611.

5. Rivera-Feliciano J and Tabin CJ. Bmp2 instructs cardiac progenitors to form the heart-valve-inducing field. *Developmental biology*. 2006;295:580-588.

6. Shirai M, Imanaka-Yoshida K, Schneider MD, Schwartz RJ and Morisaki T. T-box 2, a mediator of Bmp-Smad signaling, induced hyaluronan synthase 2 and Tgfbeta2 expression and endocardial cushion formation. *Proceedings of the National Academy of Sciences of the United States of America*. 2009;106:18604-9.

7. Niessen K, Fu Y, Chang L, Hoodless PA, McFadden D and Karsan A. Slug is a direct Notch target required for initiation of cardiac cushion cellularization. *Journal of Cell Biology*. 2008;182:315-325.

8. Del Monte G, Grego-Bessa J, Gonzalez-Rajal A, Bolos V and De La Pompa JL. Monitoring Notch1 activity in development: evidence for a feedback regulatory loop. *Developmental dynamics : an official publication of the American Association of Anatomists*. 2007;236:2594-614.

9. Grego-Bessa J, Luna-Zurita L, Del Monte G, Bolós V, Melgar P, Arandilla A, Garratt AN, Zang H, Mukouyama Y-S, Chen H, Shou W, Ballestar E, Esteller M, Rojas A, Pérez-Pomares JM and De La Pompa JL.

604   Notch Signaling Is Essential for Ventricular Chamber Development. *Developmental cell*. 2007;12:415-429.

605
10.   Timmerman LA, Grego-Bessa J, Raya A, Bertran E, Perez-Pomares JM, Diez J, Aranda S, Palomo S,

McCormick F, Izpisua-Belmonte JC and de la Pompa JL. Notch promotes epithelial-mesenchymal transition

during cardiac development and oncogenic transformation. *Genes & development*. 2004;18:99-115.

11.   Luna-Zurita L, Prados B, Grego-Bessa J, Luxan G, del Monte G, Benguria A, Adams RH, Perez-Pomares

JM and de la Pompa JL. Integration of a Notch-dependent mesenchymal gene program and Bmp2-driven cell

invasiveness regulates murine cardiac valve formation. *The Journal of clinical investigation*. 2010;120:3493-

507.

12.   Krebs LT, Xue Y, Norton CR, Shutter JR, Maguire M, Sundberg JP, Gallahan D, Closson V, Kitajewski J,

Callahan R, Smith GH, Stark KL and Gridley T. Notch signaling is essential for vascular morphogenesis in

mice. *Genes & development*. 2000;14:1343-52.

13.   MacGrogan D, D'Amato G, Travisano S, Martinez-Poveda B, Luxan G, Del Monte-Nieto G, Papoutsi T,

Sbroggio M, Bou V, Gomez-Del Arco P, Gomez MJ, Zhou B, Redondo JM, Jimenez-Borreguero LJ and de la

Pompa JL. Sequential Ligand-Dependent Notch Signaling Activation Regulates Valve Primordium Formation

and Morphogenesis. *Circulation research*. 2016;118:1480-97.

14.   Luxan G, D'Amato G, MacGrogan D and de la Pompa JL. Endocardial Notch Signaling in Cardiac

Development and Disease. *Circulation research*. 2016;118:e1-e18.

15.   Duchemin AL, Vignes H and Vermot J. Mechanically activated piezo channels modulate outflow tract

valve development through the Yap1 and Klf2-Notch signaling axis. *eLife*. 2019;8.

16.   Galvez-Santisteban M, Chen D, Zhang R, Serrano R, Nguyen C, Zhao L, Nerb L, Masutani EM, Vermot J,

Burns CG, Burns CE, Del Alamo JC and Chi NC. Hemodynamic-mediated endocardial signaling controls in

vivo myocardial reprogramming. *eLife*. 2019;8.

627
628   17.   Heckel E, Boselli F, Roth S, Krudewig A, Belting HG, Charvin G and Vermot J. Oscillatory Flow

629   Modulates Mechanosensitive klf2a Expression through trpv4 and trpp2 during Heart Valve Development.

630   *Current biology : CB*. 2015;25:1354-61.

631   18.   Mounsey JP and Dimarco JP. Dofetilide. *Circulation*. 2000;102:2665-2670.

632
633   19.   D'Amato G, Luxan G, del Monte-Nieto G, Martinez-Poveda B, Torroja C, Walter W, Bochter MS,

634   Benedito R, Cole S, Martinez F, Hadjantonakis AK, Uemura A, Jimenez-Borreguero LJ and de la Pompa JL.

635   Sequential Notch activation regulates ventricular chamber development. *Nature cell biology*. 2016;18:7-20.

636   20.   Joshua, Lan, W, Boudreau E, Stanley, He D, Schachterle W, Didier, Oettgen P, Brian, Benoit and Jason.

637
638   ETS Factors Regulate Vegf-Dependent Arterial Specification. *Developmental cell*. 2013;26:45-58.

639
640   21.   Samsa LA, Givens C, Tzima E, Stainier DY, Qian L and Liu J. Cardiac contraction activates endocardial

641   Notch signaling to modulate chamber maturation in zebrafish. *Development*. 2015;142:4080-91.

642   22.   Li YS, Haga JH and Chien S. Molecular basis of the effects of shear stress on vascular endothelial cells.

643
644   *Journal of biomechanics*. 2005;38:1949-71.

645
23.   Masumura T, Yamamoto K, Shimizu N, Obi S and Ando J. Shear stress increases expression of the arterial

endothelial marker ephrinB2 in murine ES cells via the VEGF-Notch signaling pathways. *Arteriosclerosis,*

thrombosis, and vascular biology. 2009;29:2125-31.

24.   Baffi TR, Lorden G, Wozniak JM, Feichtner A, Yeung W, Kornev AP, King CC, Del Rio JC, Limaye AJ,

Bogomolovas J, Gould CM, Chen J, Kennedy EJ, Kannan N, Gonzalez DJ, Stefan E, Taylor SS and Newton AC.

mTORC2 controls the activity of PKC and Akt by phosphorylating a conserved TOR interaction motif. *Sci*

Signal. 2021;14.

25.   Sarbassov DD, Ali SM, Kim DH, Guertin DA, Latek RR, Erdjument-Bromage H, Tempst P and Sabatini

DM. Rictor, a novel binding partner of mTOR, defines a rapamycin-insensitive and raptor-independent pathway

that regulates the cytoskeleton. *Current biology : CB*. 2004;14:1296-302.

26.   Park H, Go Y-M, John PLS, Maland MC, Lisanti MP, Abrahamson DR and Jo H. Plasma Membrane

Cholesterol Is a Key Molecule in Shear Stress-dependent Activation of Extracellular Signal-regulated Kinase.

Journal of Biological Chemistry. 1998;273:32304-32311.

27.   Fahie S and Cassagnol M. Verapamil *StatPearls* Treasure Island (FL) ineligible companies. Disclosure:

Manouchkathe Cassagnol declares no relevant financial relationships with ineligible companies.: StatPearls

PublishingCopyright © 2023, StatPearls Publishing LLC.; 2023.

28.   Dor Y, Camenisch TD, Itin A, Fishman GI, McDonald JA, Carmeliet P and Keshet E. A novel role for

VEGF in endocardial cushion formation and its potential contribution to congenital heart defects. *Development*.

2001;128:1531-1538.

29.   Chang C-P, Neilson JR, Bayle JH, Gestwicki JE, Kuo A, Stankunas K, Graef IA and Crabtree GR. A Field

of Myocardial-Endocardial NFAT Signaling Underlies Heart Valve Morphogenesis. *Cell*. 2004;118:649-663.

30.   Fukui H, Chow RW, Xie J, Foo YY, Yap CH, Minc N, Mochizuki N and Vermot J. Bioelectric signaling

and the control of cardiac cell identity in response to mechanical forces. *Science*. 2021;374:351-354.

31.   Albinsson S, Nordström I, Swärd K and Hellstrand P. Differential dependence of stretch and shear stress

signaling on caveolin-1 in the vascular wall. *American Journal of Physiology-Cell Physiology*. 2008;294:C271-

C279.

32.   Mineo C, Ying Y-S, Chapline C, Jaken S and Anderson RGW. Targeting of Protein Kinase Cα to Caveolae.

673
Journal of Cell Biology. 1998;141:601-610.

33.   Oka N, Yamamoto M, Schwencke C, Kawabe J-I, Ebina T, Ohno S, Couet J, Lisanti MP and Ishikawa Y.

Caveolin Interaction with Protein Kinase C. *Journal of Biological Chemistry*. 1997;272:33416-33421.

34.   Sural-Fehr T, Singh H, Cantuti-Catelvetri L, Zhu H, Marshall MS, Rebiai R, Jastrzebski MJ, Givogri MI,

Rasenick MM and Bongarzone ER. Inhibition of IGF-1-PI3K-Akt-mTORC2 in lipid rafts increases neuronal

vulnerability in a genetic lysosomal glycosphingolipidosis. *Disease models & mechanisms*.

2019;12:dmm036590.

35.   White AB, Givogri MI, Lopez-Rosas A, Cao H, Van Breemen R, Thinakaran G and Bongarzone ER.

Psychosine Accumulates in Membrane Microdomains in the Brain of Krabbe Patients, Disrupting the Raft

Architecture. *The Journal of Neuroscience*. 2009;29:6068-6077.

36.   Yamamoto K and Ando J. Endothelial cell and model membranes respond to shear stress by rapidly

decreasing the order of their lipid phases. *Journal of cell science*. 2013;126:1227-1234.

37.   Steinbuck MP, Arakcheeva K and Winandy S. Novel TCR-Mediated Mechanisms of Notch Activation and

Signaling. *The Journal of Immunology*. 2018;200:997-1007.

38.   Goenezen S, Rennie MY and Rugonyi S. Biomechanics of early cardiac development. *Biomechanics and

modeling in mechanobiology*. 2012;11:1187-204.

39.   Doubilet PM, Benson CB and Chow JS. Long-term prognosis of pregnancies complicated by slow

embryonic heart rates in the early first trimester. *Journal of Ultrasound in Medicine*. 1999;18:537-541.

40.   Braga M, Moleiro ML and Guedes-Martins L. Clinical Significance of Ductus Venosus Waveform as

Generated by Pressure- volume Changes in the Fetal Heart. *Curr Cardiol Rev*. 2019;15:167-176.

 Resource Table

Category	Name	Source	Catalog		
Chemicals	Staruosporine	MCE	HY-15141		
	Phorbol 12-myristate 13-acetate (PMA)	Sigma-Aldrich	P8139		
	Dofetilide	Rhawn	R023594		
	Wortmannin	Selleck	S2758		
	Cholesterol-Water soluble	Sigma-Aldrich	C4951		
	CMC-Na	Shyuanye	S14016		
	Saline	Shyuanye	R27405		
	Isoflurane	RWD	R510-22-10		
	10% neutral buffered formalin	Legene	DF0111		
	Glutaraldehyde	Sigma-Aldrich	G7651		
	Benzyl alcohol	Sinoreagent	30020618		
	Benzyl benzoate	Sinoreagent	30020828		
	Glycerol	Sinoreagent	10010618		
	Hanks buffer	Beyotime	C0218		
	DMEM	Gibco	C11965500BT		
	DMSO	Macklin	D806645		
	Triton X-100	Macklin	I997471		
	Fetal bovine sreum	Solarbio	S9030		
	Verapamil	Selleck	S4202		
Mouse lines	$Dll4^{flox/flox}$	Cyagen	$C57BL/6J-Dll4^{em1cyagen}$		
	$Nfatc1^{cre/cre}$	GemPharmatech	$C57BL/6JGpt-Nfatc1^{em1Cin(P2A-iCre)}/Gpt$		
	$Tie2^{cre}$	GemPharmatech	$C57BL/6JGpt-H11^{em1Cin(Tek-iCre)}/Gpt$		
	$Tnnt2^{flox/flox}$	GemPharmatech	$C57BL/6JGpt-Tnnt2^{em1Cflox}/Gpt$		
	$cTnt^{cre}$	Jax	STOCK Tg(Tnnt2-cre)5Blh/JiaoJ		
	$Prkce^{KO}$	Westlake university			
	$Prkch^{KO}$	Westlake university			
	$Rictor^{KO}$	Westlake university			
	$Notch1^{KO}$	Westlake university			
	FVB	Beijing HFK			
	C57bl/6	Shanghai Jihui Laboratory			
	$Epor^{cre}$	GemPharmatech	$C57BL/6JGpt-Epor^{em1Cin(P2A-iCre-P2A-EGFP)}/Gpt$		
	ROSA-DTA	Jax	$B6.129P2-Gt(ROSA)26Sor^{tm1(DTA)Lky}/J$		
Genotyping PCR primers	Dll4-F: CAACTGACCTAAAATGGGATGGTG				
	Dll4-R: GGTAACTACAAGGCAGAAAGAGGA				
	Tnnt2-F: GATCCTGCCTCCTTAGGTCTCAAGT				
	Tnnt2-R: CAAGTTCCTATGCCACATCTGCATG				
	Nfatc1cre-F: ACAGTCCCAGGAGCAGCAAACC				
	Nfatc1cre-R: CAGGAAGGCCAGGTTCCTGATG				
	Nfatc1WT-F: CCCTGCTATTGACTTGGAGGC				
	Nfatc1WT-R: CCGTCCCTGTTGGAGAGCTACT				
	Tie2cre-F: GCGGTCTGGCAGTAAAAACTATC				
	Tie2cre-R: GTGAAACAGCATTGCTGTCACTT				
	cTnTcre-F: GGACATGTTCAGGGATCGCCAGGCG				
	cTnTcre-R: GCATAACCAGTGAAACAGCATTGCTG				
	Prkce-F: GAGTGTTCAGGGAGCGTATG				
	Prkce-R: CAAGTAGGTGGCCATGAACTTG				
	Prkch-F: GAGACCGCATCTTCAAGC				
	Prkch-R: GTAGGTGGGCTGCCTC				
	Rictor-F: TCCTTCTCTGTTACAGATG				
	Rictor-R: ACCATTCTGTCTCGTTC				
	Rosa-DTA_FWD: GTTATCAGTAAGGGAGCTGCAGTGG				
	Rosa-DTA_wt_RVS: GGCGGATCACAAGCAATAATAACC				
	Rosa-DTA_mt_RVS: AAGACCGCGAAGAGTTTGTCCTC				
	Epor-F: TCCCACTCCACCTCACTTGAAG				
	Epor-R: CTGACTTCATCAGAGGTGGCATC				
	Notch1-F: AAGCTGGGAGGAGAAAAGCAGACC				
	Notch1-R: CACAACCTCCTATAGCCCTTACC				
Immunohistochemistry	Citrate acid solution (50x)	Beyotime			
	Normal donkey serum	Solarbio			
	Paraformaldehyde	Sigma-Aldrich			
	HCR probe set	Molecular instrument			

	Name	Catalog	Species and type	Supplier	Dilution
Antibodies for immunohistochemistry	Primary antibody				
	Cleaved Notch (NICD)[*]	#4147	Rabbit monoclonal	Cell Signaling Technology	1:400
	Phospho-AKT (Ser473) (D9E)[*]	#4060	Rabbit monoclonal	Cell Signaling Technology	1:400
	Phospho-SMAD1/5 (Ser463/465) (41D10)[*]	#9516	Rabbit monoclonal	Cell Signaling Technology	1:400
	Phospho-PKC (pan) (βII Ser660)[*]	#9371	Rabbit monoclonal	Cell Signaling Technology	1:400
	Jagged1 (D4Y1R)[*]	#70109	Rabbit monoclonal	Cell Signaling Technology	1:400
	VEGF receptor 2 (D5B1)	#9698	Rabbit monoclonal	Cell Signaling Technology	1:400
	PKCε (22B10)[*]	#2683	Rabbit monoclonal	Cell Signaling Technology	1:400
	PKC eta (EPR18513)[*]	ab179524	Rabbit monoclonal	Abcam	1:500
	Notch1 (EP1238Y)[*]	ab52627	Rabbit monoclonal	Abcam	1:500
	VE-Cadherin	AF1002	Goat polyclonal	R&D Systems	1:500
	Dll4[*]	AF1389	Goat polyclonal	R&D Systems	1:400
	Troponin T (RV-C2)	AB 2240831	Rabbit monoclonal	DSHB	1:200
	Sox9 (SN74-09)	ET1611-56	Rabbit monoclonal	HuaBio	1:200
	Secondary antibody				
	HRP-Donkey Anti-Rabbit IgG (H+L)	711-035-152	Donkey polyclonal	Jackson ImmunoResearch	1:500
	Peroxidase AffiniPure Donkey Anti-Mouse IgG (H+L)	715-035-150	Donkey polyclonal	Jackson ImmunoResearch	1:500
	Peroxidase AffiniPure Donkey Anti-Goat IgG (H+L)	705-035-003	Donkey polyclonal	Jackson ImmunoResearch	1:500
	Cy™5 AffiniPure Donkey Anti-Goat IgG (H+L)	705-175-147	Donkey polyclonal	Jackson ImmunoResearch	1:500
	Cy™3 AffiniPure Donkey Anti-Goat IgG (H+L)	711-165-152	Donkey polyclonal	Jackson ImmunoResearch	1:500
	Alexa Fluor® 488 AffiniPure Donkey Anti-Mouse IgG (H+L)	715-545-150	Donkey polyclonal	Jackson ImmunoResearch	1:500
	Cy™3 Streptavidin	016-160-084		Jackson ImmunoResearch	1:500
	Alexa Fluor® 488 Streptavidin	016-540-084		Jackson ImmunoResearch	1:500
	DAPI			Beyotime	1:500
	*Antibodies requiring tyramide signal amplification (TSA)				
Software	ImageJ 1.52i				
	Zen 2.3				
	Imaris v9.1.1				
	Adobe Photoshop cc2019				
	Olympus OlyVIA 3.3				
	GraphPad Prism 8				
	Vevo Lab 5.7.1				

Figure 1-figure supplement 1

A

Wildtype pregnant mice

Dofetilide

E9.5   +1 hour   +3 hrs.   +5 hrs.   Histological analysis E10.5   Phenotyping E18.5

B

Control

Dofetilide 1 h

Heart rate (bpm)

0 h   1 h   3 h   5 h

****
****

C

Dofetilide 0 h   Dofetilide 1 h   Dofetilide 3 h   Dofetilide 5 h

AV Canal — NICD

AVC NICD intensity
0 h 1 h 3 h 5 h
***

Dorsal Aorta — NICD

DA NICD intensity
0 h 1 h 3 h 5 h
ns

Proximal OFT — NICD

POFT NICD intensity
0 h 1 h 3 h 5 h
**

AV Canal — Total Notch1

Total Notch1 intensity
0 h 1 h 3 h 5 h
ns

AV Canal — p-SMAD1/5

p-SMAD1/5 intensity
0 h 1 h 3 h 5 h
****

AV Canal — Sox9

Sox9 intensity
0 h 1 h 3 h 5 h
***

AV Canal — Twist1

Twist1 intensity
0h 1h 3h 5h
****

D

Embryonic day 10.5

Control   Dofetilide

OFT   A   V

Superior AV Cushion

Number of cells /mm²
Control   Dofetilide
****

Inferior AV Cushion

Number of cells /mm²
Control   Dofetilide
****

E

OA/pmVSD — ra ao la rv s lv

DORV — ra pa ao rv s lv

mVSD — ra la rv s lv

BAV — ra pa la

AVSD — ra rv s lv

BPV — ra ao la

Figure 2-figure supplement 1

A

Dofetilide 0 h Dofetilide 1 h Dofetilide 3 h Dofetilide 5 h

Dll4

Jag1

B

Heart defect rate (%)

Dofetilide dosage (mg/kg)

☐ Normal ☐ Defect

****

Total No. of abnormal hearts	47
Defect types	pmVSD	30 (63.8%)
	DORV	19 (40.4%)
	mVSD	11 (23.4%)
	OA	10 (20.3%)
	BAV	8 (17.0%)
	AVSD	6 (12.8%)
	BPV	1 (2.1%)

C

Tnnt2^flox/flox *cTnT^Cre;Tnnt2^flox/flox*

AV Canal NICD TnnT2

Proximal OFT NICD TnnT2

Dorsal Aorta NICD VE-cadherin

D

Control Blebbistatin 3 h

AV Canal NICD VE-cad

Proximal OFT NICD VE-cad

Dorsal Aorta NICD VE-cad

E

Wild type *Epor^cre;ROSA-DTA*

E9.5

HIF-1α

NICD

F

Control Dofetilide 3 h Dofetilide 3 h (95% O₂) Dofetilide 5 h

HIF-1α

NICD

Figure 3-figure supplement 1

A

B

	Total No. of abnormal hearts	10
	pmVSD	6 (60.0%)
	BPV	2 (20.0%)
Defect types	DORV	1 (10.0%)
	AVSD	1 (10.0%)
	OA	1 (10.0%)
	BAV	1 (10.0%)

C

D

Total No. of abnormal hearts		6
Defect types	pmVSD	6 (100.0%)

Dorsal aorta

OFT cushions

Myocardium

Atrioventricular canal

Endocardium

Cardiac jelly

AV cushions

Trabecula

BMP2　NICD　Dll4　VEGFR2

E8.5

E9.5

Non-EMT endocardium of early heart tube

High shear stress cushion endocardium capable of EMT

Low shear stress non-cushion endocardium