Mechano-regulation of GLP-1 production by Piezo1 in intestinal L cells

Yanling Huang[1#], Haocong Mo[1#], Jie Yang[2#], Luyang Gao[1#], Tian Tao[1], Qing Shu[1], Wenying Guo[1], Yawen Zhao[1], Jingya Lyu[1], Qimeng Wang[3], Jinghui Guo[4], Hening Zhai[5], Linyan Zhu[6], Hui Chen[3*], Geyang Xu[1,7*]

[1]Department of Physiology, School of Medicine, Jinan University, 601 Huangpu Avenue West, Tianhe District, Guangzhou 510632, Guangdong, China;

[2]Department of Pathology, School of Basic Medicine, Guangzhou Medical University, Guangzhou 511436, Guangdong, China;

[3]Biotherapy Center, Cell-gene Therapy Translational Medicine Research Center, the Third Affiliated Hospital, Sun Yat-Sen University, Guangzhou 510630, Guangdong, China.

[4]School of Medicine, The Chinese University of Hong Kong, Shenzhen 518172, Guangdong, China;

[5]Endoscopy Center, The First Affiliated Hospital of Jinan University, 613 Huangpu Avenue West, Tianhe District, Guangzhou 510630, Guangdong, China;

[6]Department of Pharmacology, School of Medicine, Jinan University, Guangzhou 510632, Guangdong, China.

[7]Key Laboratory of Viral Pathogenesis & Infection Prevention and Control (Jinan University), Ministry of Education, Guangzhou 510632, Guangdong, China.

[#]Yanling Huang, Haocong Mo, Jie Yang and Luyang Gao contributed equally to this work.

* Corresponding Author:

Geyang Xu

Department of Physiology, School of Medicine, Jinan University, 601 Huangpu Avenue West, Tianhe District, Guangzhou, Guangdong, 510632, China.

Tel: 86-20-85220260

Fax: 86-20-85221343

E-mail: xugeyangliang@163.com

Or

Hui Chen

Biotherapy Center; Cell-Gene Therapy Translational Medicine Research Center, the Third Affiliated Hospital of Sun Yat-Sen University, Guangzhou 510630, Guangdong, China.

Tel: 86-20-85252826

E-mail: chenh567@mail.sysu.edu.cn

Abstract

Glucagon-like peptide 1 (GLP-1) is a gut-derived hormone secreted by intestinal L cells and vital for postprandial glycemic control. As open-type enteroendocrine cells, whether L cells can sense mechanical stimuli caused by chyme and thus regulate GLP-1 synthesis and secretion is unexplored. Our study showed expression of Piezo1 in intestinal L cells. Its level varied in different energy status and correlates with blood glucose and GLP-1 levels. Mice with L cell-specific loss of Piezo1 (*IntL-Piezo1*$^{-/-}$) exhibited impaired glucose tolerance, increased body weight, reduced GLP-1 production and decreased CaMKKβ/CaMKIV-mTORC1 signaling pathway under normal chow diet or high fed diet. Activation of the intestinal Piezo1 by its agonist Yoda1 or intestinal bead implantation increased the synthesis and secretion of GLP-1, thus alleviated glucose intolerance in diet-induced-diabetic mice. Overexpression of Piezo1, Yoda1 treatment or stretching stimulated GLP-1 production and CaMKKβ/CaMKIV-mTORC1 signaling pathway, which could be abolished by knockdown or blockage of Piezo1 in primary cultured mouse L cells and STC-1 cells. These findings suggest a previously undiscovered mechano-regulation of GLP-1 production in L cells, which may shed new light on the treatments of diabetes.

Keywords: Piezo1, Glucagon-like peptide 1, Intestinal L cells, CaMKKβ, CaMKIV, Mammalian target of rapamycin

63 **Abbreviations:**

CaMKKβ	Ca^{2+}/calmodulin-dependent protein kinase kinase β
CaMKIV	Ca^{2+}/calmodulin-dependent protein kinases IV
EECs	Enteroendocrine cells
mTOR	mechanistic target of rapamycin
RYGB	Roux-en-Y gastric bypass
S6K	p70 S6 Kinase
S6	S6 Ribosomal Protein
T2DM	Type 2 Diabetes Mellitus
CCK	Cholecystokinin

Introduction

The gastrointestinal (GI) tract represents the largest endocrine organ in the human body. The enteroendocrine cells (EECs) located throughout the GI tract secrete a large number of gastrointestinal hormones to regulate a variety of physiological processes and are key regulators for energy homeostasis (Bany Bakar et al., 2023). GLP-1 is one of the gut-derived peptide hormones essential for postprandial glycemic control (Song et al., 2019). It is produced from Proglucagon (Gcg) by proprotein convertase in the intestinal L cells, a group EECs predominantly situated in the distal gut (Drucker, 2006; Rouillé et al., 1997). The circulating GLP-1 levels rapidly increase after meal and reduce postprandial blood glucose fluctuations by augmenting insulin secretion, suppressing glucagon secretion and slowing gastric emptying (Drucker, 2006; Willms et al., 1996). Nowadays, GLP-1-based therapy is well-recognized and commonly used in treatment of Type 2 Diabetes Mellitus (T2DM) (Saxena et al., 2021; Tan et al., 2022). Although GLP-1 analogs such as exenatide and liraglutide have been approved for clinical use in the treatment of T2DM, these drugs may lead to side effects such as pancreatitis (Elashoff et al., 2011), as well as potential risk of inducing medullary thyroid carcinoma (Bezin et al., 2023). Therefore, elucidation of the mechanism that regulates GLP-1 production is essential for the development of new drug targets for the treatment of diabetes.

EECs can be divided into two categories according to its morphology: open type and closed type. The open type EECs possess microvilli protruding into the gut lumen and have direct contact with the luminal contents. In contrast, the closed type EECs are located basolaterally without direct contact with the lumen (Gribble & Reimann, 2016). Both types of EECs synthesize and store peptides or hormones in secretory granules and release them by exocytosis at the basolateral membrane upon mechanical, chemical or neural stimulation (Atanga et al., 2023). As open-type cells, L cells received both chemical and mechanical signals from the luminal contents, and neural signals from the nerves (Furness et al., 2013). It has been well-documented that nutrients such as glucose, lipids, and amino acids in the intestinal lumen can stimulate

the secretion of GLP-1 from L cells (Diakogiannaki et al., 2012). GLP-1 secretion can

also be stimulated by intrinsic cholinergic nerves (Anini et al., 2002; Drucker, 2006).

However, whether and how L cells coordinate mechanical stimuli from intestinal

lumen to regulate GLP-1 production remain poorly understood.

Piezo channels, including Piezo1 and Piezo2 have recently been identified as a

mechanosensitive ion channel that is involved in the sensation of multiple mechanical

stimuli, such as shear stress, pressure, and stretch (Gudipaty et al., 2017; Li et al.,

2014; Romac et al., 2018). They allow the influx of cations such as Ca^{2+} and Na^+ in

response to mechanical tension and converts mechanical stimuli into various electrical

and chemical signals. Piezo1 plays a crucial role in blood pressure regulation, red

blood cell volume regulation, bone homeostasis, pulmonary and cardiac functions

(Cahalan et al., 2015; Lai et al., 2022; Wang et al., 2023; Wang et al., 2016). Previous

studies have reported that Piezo1 is expressed in the intestinal epithelium, regulating

barrier function, mucus secretion, and inflammation (Jiang et al., 2021; Liu et al.,

2022; Xu et al., 2021). Interestingly, accumulating evidence demonstrates the

regulation of insulin secretion by Piezo1 (Deivasikamani et al., 2019; Ye et al., 2022).

Recent studies have also reported that Piezo2 is expressed in a population of EECs

and convert force into serotonin release (Alcaino et al., 2018; Treichel et al., 2022).

These findings suggest a critical role of Piezo channels in the mechano-regulation of

hormone production. However, whether Piezo channels are expressed L cells and play

a role in GLP-1 production remain unknown.

Here, we present evidence that Piezo1 channels on intestinal L cells mediate the

mechanosensing from intestinal contents and trigger the synthesis and secretion of

GLP-1 through CaMKKβ/CaMKIV-mTORC1 signaling pathway, thus regulating

glucose homeostasis. Our research provides new insights for the treatment of T2DM

and a new theoretical basis for hypoglycemic medicines targeting Piezo1.

Materials and Methods

Collection of Human Intestine Samples

Obese participants with type 2 diabetes (n=6, BMI=45.87±4.889 kg/m^2) and one-year post-RYGB patients (n=6, BMI=25.48±1.085 kg/m^2) were recruited in current study. Written informed consent was obtained from each donor. The study protocol was approved by the Institutional Review Board of Jinan University. Mucosal biopsies were obtained from human intestines by using a colonoscopy (CF-HQ290I; Olympus).

Genetic Mouse Generation

Villin-Flippase (Vil-Flp) mice

Vil-Flp knock-in mouse model was developed by Shanghai Model Organisms Center, Inc. The targeting construct was designed to insert a 2A-Flp-WPRE-pA coexpression cassette into the stop codon of mouse *Vil1* gene via homologous recombination using CRISPR/Cas9 system. 5'-AGCCCCTACCCTGCCTTCAA-3' was chosen as Cas9 targeted guide RNA (sgRNA). The donor vector, sgRNA and Cas9 mRNA was microinjected into C57BL/6J fertilized eggs. F0 generation mice positive for homologous recombination were identified by long PCR. The primers (I-IV) used for detection of the correct homology recombination were I: 5'-ACTTCAGGCCTAACGCTCAC-3' and II: 5'-TGTCCTGCAGGCAGAGAAAG-3' for the correct 5' homology arm recombination, and III: 5'-GTGCCGTCTCTAAGCACAGT-3'and IV: 5'-TGTTGGTGCTTCGGAGTGTT-3'for the correct 3' homology arm recombination. The PCR products were further confirmed by sequencing. F0 mice were crossed with C57BL/6J mice to obtain *Vil-Flp* heterozygous mice.

Flp-dependent Glucagon-Cre (FGC) mice

FGC mouse model was developed by Shanghai Model Organisms Center, Inc. The targeting construct was designed to insert an IRES-F3-Frt-Wpre-pA-Cre-Frt-F3 expression cassette into the 3' UTR of mouse *Gcg* gene of via homologous recombination using CRISPR/Cas9 system. 5'-ATGCAAAGCAATATAGCTTC-3'

was chosen as Cas9 targeted guide RNA (sgRNA). The donor vector, sgRNA and

Cas9 mRNA was microinjected into C57BL/6J fertilized eggs. F0 generation mice

positive for homologous recombination were identified by long PCR. The primers

(I-IV) used for detection of the correct homology recombination were I:

5'-TGCTACACAGGAGGTCTGTC-3' and II: 5'-AGGCATGCTCTGCTATCACG-3'

for the correct 5' homology arm recombination, and III:

5'-CCCTCCTAGTCCCTTCTCAGT-3' and IV:

5'-GCCAAGGACATCTTCAGCGA-3' for the correct 3' homology arm

recombination. The PCR products were further confirmed by sequencing. F0 mice

were crossed with C57BL/6J mice to obtain *FGC* heterozygous mice.

IntL-Cre mice

Vil-Flp mice were crossed with *FGC* mice to obtain Intestinal L cell-specific Cre

(*IntL-Cre*) mice.

IntL-Piezo1$^{-/-}$ mice

Piezo1$^{loxp/loxp}$ mice (B6.Cg-Piezo1$^{tm2.1Apat}$/J) purchased from Jackson laboratory

were crossed with IntL-Cre mice to generate *IntL-Piezo1$^{-/-}$* mice.

PCR is used to identify the genotype of mice during the subsequent mating and

breeding process. The primers required for mouse genotyping are shown in the SI

Appendix, Table S1.

Mouse Validation

mT/mG reporter mice were purchased from Shanghai Model Organisms Center,

Inc. *IntL-Cre* mice were bred with *mT/mG* reporter mice to further validate Cre

recombinase activity and specificity. Every single *IntL-Cre* mouse was confirmed by

mT/mG reporter mice before breeding with *Piezo1$^{loxp/loxp}$* mice to generate Intestinal L

cell-Piezo1$^{-/-}$ (*IntL-Piezo1$^{-/-}$*) mice.

Frozen Tissue Confocal Imaging

IntL-Cre-mT/mG reporter mice were sacrificed. Fresh ileum and pancreas tissues were collected and embedded in O.C.T. compound for histological analysis immediately. Slices of the tissues were cut for confocal imaging, which was protected from light. Fluorescence was detected by laser scanning confocal microscopy (Li et al., 2022).

Animal Housing and Treatment

Male mice were maintained on a 12-hour light/12-hour dark cycle environment. Normal chow diet (NCD) or a high-fat diet (HFD) and water were available ad libitum unless specified otherwise. The animal protocols were approved by the Animal Care and Use Committee of Jinan University.

Male *IntL-Piezo1$^{-/-}$* mice and age-matched control littermates (*Piezo1$^{loxp/loxp}$*, *Vil-Flp*, *FGC*, *IntL-Cre* mice) fed with NCD or HFD were used in the experiments.

Male *Piezo1$^{loxp/loxp}$* and *IntL-Piezo1$^{-/-}$* mice fed with 10 week-high fat diet were intraperitoneally injected with normal saline (NS) or the GLP-1R agonist Exendin-4 (100 ug/kg body weight) for 7 consecutive days.

High fat diet induced diabetic mice were randomly divided into 3 groups. When indicated, animals were injected intraperitoneally with Vehicle, Yoda1 (2 μg per mouse) or GsMTx4 (250 μg/kg) for 7 consecutive days.

High fat diet treated *IntL-Piezo1$^{-/-}$* mice were randomly divided into 2 groups. When indicated, animals were injected intraperitoneally with Vehicle, Yoda1 (2 μg per mouse) for 7 consecutive days.

Diet induce diabetic C57BL/6J mice were divided into sham and intestinal bead implantation groups.

Food and water intake detection

The food and water intake were quantified using metabolic cages (Cat 41853, Ugo Basile, Comerio, Italy). The mice were individually housed in these specialized cages and given a period of 3 days to acclimate before data collection began. They had unrestricted access to food and water, which was continuously monitored

throughout the study. The 41850-010 software/interface package, consisting of EXPEDATA (for data analysis) and METASCREEN (for data collection) software, along with the IM-2 interface module, was employed to record and analyze the data.

Intraperitoneal Glucose Tolerance Test

Mice were fasted for 12 hours before measuring their fasting glucose levels. An intraperitoneal glucose tolerance test (IPGTT) was performed by administering 1.5 g/kg body weight of glucose. Blood glucose concentrations were measured at specified time points using a glucometer by collecting tail vein blood samples.

Insulin Tolerance Test

Mice were subjected to a 4-hour fast before measurement of fasting glucose were taken. Insulin tolerance tests (ITT) were conducted with a dose of 0.75 U/kg body weight of insulin. Blood glucose levels were measured at specified time points.

Intestinal Bead implantation

High-fat diet-induced type 2 diabetic C57BL/6J mice were fasted 6 to 8 h before the operation. Standard aseptic procedures were used throughout the operation. For bead implantation, a 1cm incision was made on the abdominal wall to expose the intestine. A 1 cm incision was made approximately 1cm above the ileocecal region. A 2.5 mm diameter bead was implanted into the ileum of the mouse through an incision. Then the wound was closed with suture. Finally, the abdominal wall was closed with suture. For sham operation, all the procedures were the same as the bead implantation except that the bead was not implanted.

Stretching of isolated ileum

About 2cm ileum was isolated from control and *IntL-Piezo1*$^{-/-}$ mice and kept in the specimen chamber filled with Tyrode's solution (KCl 0.2g/L, NaCl 8g/L, CaCl$_2$ 0.2g/L, MgCl$_2$ 0.1g/L, NaHCO$_3$ 1g/L, NaH$_2$PO$_4$ 0.05g/L, Glucose 1g/L) of 37°C

gassed with oxygen. The specimen was connected to the force transducer of organ bath system (HW200S, Techman, Chengdu, CN). Adjust the transducer to apply traction force of 1.5 grams on the tissue and maintained for two hours.

Measurement of GLP-1 Secretion

The measurement of GLP-1 secretion was carried out according to previously described methods (Zhai et al., 2018). Samples were collected in the presence of aprotinin (2 μg/mL), EDTA (1 mg/mL) and diprotin A (0.1 mmol/L), and stored at -80 °C before use. GLP-1 levels were assayed using enzyme immunoassay kits following the manufacturer's instructions.

Histological Analysis

Tissues were collected, fixed with 4% paraformaldehyde, embedded in paraffin, and cut into 4μm sections. Standard protocols were followed for staining the sections with hematoxylin-eosin. Photomicrographs were captured under an inverted microscope (Leica, Germany).

Immunofluorescence

Paraffin sections were dewaxed and rehydrated. After antigen retrieval in citrate buffer (pH6.0), sections were blocked with normal serum and then incubated with rabbit anti-Piezo1 (1:400) and mouse anti-GLP-1 (1:500) antibodies at 4°C overnight. The sections were then incubated with a mixture of secondary antibodies. Images were taken by laser scanning confocal microscopy (Leica SP8). Fluorescence intensity was quantitated by ImageJ software.

In situ hybridization

Paraffin sections were dewaxed and rehydrated. After antigen retrieval in in citrate buffer (pH6.0), the sections were incubated with Proteinase K (5ug/ml) at 37°C for the 15 minutes. Then the sections were hybridized with the probes overnight in a

temperature-controlled chamber at 40°C. The Piezo1 probe sequences were as follows:

5′-CTGCAGGTGGTTCTGGATATAGCCC-3',5′-AAGAAGCAGATCTCCAGCCCG

AAT-3', 5′-GCCATGGATAGTCAATGCACAGTGC-3'. After washing with SSC

buffers, the sections were hybridized in pre-warmed branch probes at 40°C for 45

minutes. After washing with SSC buffers, the sections were hybridized with signal

probe at 42°C for 3 hours. After washing with SSC buffers, the sections were blocked

with normal serum and then incubated with mouse anti-GLP-1 (1:500) antibody at

4°C overnight followed by secondary antibody. Images were taken laser scanning

confocal microscopy (Leica SP8) and the fluorescence signals were quantified by

ImageJ.

Western Blot Analysis

Tissues and cells were harvested. Ileal mucosa was scraped for protein extraction.

Protein extraction was performed by using RIPA lysis buffer (50mM Tris PH 7.4,

150mM NaCl, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

and protease inhibitor cocktail.), then 40 μg of proteins were loaded onto an

SDS-PAGE gel for separation. After the separation, the proteins were transferred onto

a nitrocellulose membrane. The membrane was then incubated in blocking buffer at

room temperature for 1 hour. For overnight incubation, the membrane and primary

antibody (at the recommended dilution as stated in the product datasheet) were

immersed in primary antibody dilution buffer, with gentle agitation, at 4°C.

Subsequently, the membrane was incubated with a secondary antibody that

specifically recognizes and binds to the primary antibody. Finally, Western blotting

luminol reagent was used to visualize bands. The grey scale values of the bands were

measured using ImageJ software.

RNA extraction, quantitative real-time PCR

RNA was extracted and reverse-transcribed into cDNAs using RT-PCR kit. Real-time PCR was performed as previously described (Zhai et al., 2018). Sequences for the primer pairs used in this study were shown in SI Appendix, Table S2.

Isolation of Mouse Intestinal L Cells

A 5~6cm long ileum segment was collected from the IntL-cre-mT/mG mouse. The tissue was washed with ice-cold PBS twice to remove the chyme in the lumen. The tissue was minced into $0.5mm^3$ pieces in ice-cold PBS and then digested in 100mIU collagenase I and 0.01g/mL trypsin at 37°C for 30min with rotation. After digested tissue was passed through 40μm and 30μm cell strainers sequentially, then centrifuged for 7min at 4°C. The cell pellete was resuspended in red cell lysis buffer and incubated for 10min at room temperature. The unlysed cells were collected by centrifugation and resuspended with 1mL cold PBS. The GFP positive cells was sorted by fluorescence-activated cell sorting (FASC) on Beckman Coulter MoFlo XDP cell sorter system.

Cell Culture and Treatments

STC-1 cells were maintained in DMEM medium supplemented with 2.5% fetal bovine serum and 10% equine serum at 37 °C with 5% CO_2 air. L cells were maintained in DMEM medium supplemented with 10% fetal bovine serum.

For cell transfection, cells were plated at optimal densities and grown for 48 h. Cells were then transfected with GFP, Piezo1-GFP, CaMKKβ, CaMKIV by using lipofectamine reagent according to the manufacturer's instructions.

For stable knockdown of Piezo1 in STC-1 cells, short hairpin RNA (shRNA) sequences for mouse Piezo1 interference were cloned in to pLKO.1 vector. To produce lentivirus, psPAX2, pMD2G and pLKO.1 or pLKO.1-shPiezo1 plasmids (siPiezo1: CCAACCTTATCAGTGACTT) were co-transfected into 293T cells with lipofectamine 2000 reagent. Supernatant containing lentivirus was collected 48 hours

after transfection and filtered through 0.45μm filter. The virus-containing supernatant
was used to infect STC-1 cells. Forty-eight hours after infection, the STC-1 cells were
subjected to 1ug/mL puromycin selection for 2-3 days.

For cell stretching, cells were grown in silicone elastic chambers coated by 0.1 %
gelatin solution. After incubated at 37 °C for 24-48 hours, The chambers were
subjected to mechanical stretch to 120% of their original length.

328 Calcium Imaging

Cells were plated onto confocal dishes at optimal densities and grown for 24 h.
Cells were loaded with the calcium fluorescent probe fluo-4 AM (1 μM) for 1 h at
37 °C, then the cells were treated with Yoda1 (5 μM) or GsMTx4. The intracellular
calcium ions were measured at room temperature using a laser confocal microscope
with an excitation wave length of 494 nm and an emission wave length of 516 nm.
The change of fluorescent signal was presented as$\Delta F/F0$ and plotted against time.

336 Whole-cell Patch-Clamp recording

Borosilicate glass-made patch pipettes (BF150-86-7.5, Sutter Instrument Co,
USA) were pulled with micropipette puller (P-1000, Sutter Instrument Co, USA) to a
resistance of 3–5 MΩ after being filled with pipette solution: 138mM KCl, 10mM
NaCl, 1mM $MgCl_2$, 10mM Glucose and 10mM HEPES (pH 7.4). Cells were bathed
in Margo-Ringer solution: 130mM NaCl, 5mM KCl, 1mM $MgCl_2$, 2.5mM $CaCl_2$,
10mM Glucose, 20mM HEPES (pH7.4). Whole-cell calcium currents of STC-1 cells
were recorded with the EPC10 USB patch-clamp amplifier (HEKA, Germany)
controlled by PatchMaster software.

346 Key resource

347   Key reagents or resources are listed in the SI Appendix, Table S3.

Statistical Analysis

All data were expressed as mean ± S.E.M. Statistical differences were evaluated by one-way ANOVA or Student's t-test. The correlation was determined by Pearson analysis. $P < 0.05$ was considered significant. (*$p < 0.05$, **$p < 0.01$, *** $p < 0.001$, ns=not significance).

Results

Assessment of Piezo1 in human and mouse intestine in different energy status

Piezo1 mRNA was found to be highly expressed in both mouse ileal mucosa and STC-1 cells (Figure supplement 1A). Moreover, Piezo1 was co-localized with GLP-1 in immunofluorescent staining on NCD fed mouse ileal sections, indicating its expression in L cells (Figure supplement 1B). Interestingly, increased body weight and impaired glucose tolerance were observed in high-fat diet-induced diabetic mice, while Piezo1 and Proglucagon expression levels in the ileal mucosa of diabetic mice were significantly lower than that in mice feed with normal chow diet (Figure supplement 1C-F). Moreover, ileal mucosal *Piezo1* mRNA levels were positively correlated with *Proglucagon* mRNA levels (Figure supplement 1G), but negatively correlated with the AUC of glucose tolerance test (Figure supplement 1H). Obese T2MD patients who underwent Roux-en-Y gastric bypass (RYGB) surgery showed decreased BMI (Figure supplement 1I) and increased *Piezo1* and GLP-1 in ileal mucosa (Figure supplement 1J and K) compared to that before surgery. These findings indicated that Piezo1 is expressed in intestinal L cells and its level varies in different energy status.

Generation and characterization of *IntL-Piezo1$^{-/-}$* mice

To investigate the potential role of Piezo1 in GLP-1 production, we tried to knockout *Piezo1* in L cells by Cre-loxP system driven by an L cell-specific promoter. Proglucagon (encoded by *Gcg* gene) is mainly expressed in both L cells and pancreatic α cells (Jin, 2008). Villin-1 (encoded by *Vil1* gene) is expressed in gastrointestinal epithelium, including L cells, but not in pancreatic α cells (Maunoury et al., 1992; Rutlin et al., 2020). Since neither Gcg nor Villin are specific markers for L cells, we tried to generate a new line of mice enabling loss of Piezo1 expression specifically in the intestine L cell by combination of Flp-Frt and Cre-loxP system. We inserted a Flippase (Flp) expression cassette in the 3'UTR of *Vil1* to generate a *Vil1* promoter-driven Flp mice (*Vil-Flp*) (Fig. 1A). Then, we generated Flippase-dependent Gcg promoter driven-Cre (*FGC*) mice by inserting an Frt-flanked Cre expression cassette in reverse orientation within the 3'- UTR of *Gcg* gene (Figure 1A). We further crossed the *Vil-Flp* mice with *FGC* mice to obtain L cell specific Cre mice (*IntL-Cre*), in which *Vil1* promoter-driven Flippase flipped the reverse Cre cassette into a correct orientation in Villin positive cells (including L cells, but not pancreatic α cells), and thus Cre can only be expressed under the *Gcg* promoter in L cells. The genotypes of the *Vil-Flp*, *FGC* and *IntL-Cre* mice were identified by PCR with specific primers (Figure 1B). The flipping of the reverse Cre cassette was validated by PCR, which confirmed that the flipping only occurred the intestine, but not in the pancreas (Figure 1C). To confirm the cell type specificity of Cre activity, we crossed *IntL-Cre* mice to *mT/mG* reporter mice. All tissues and cells of *mT/mG* mice express red fluorescence (membrane-targeted tdTomato; mT) at baseline, and switch to membrane-targeted EGFP in the presence of cell-specific Cre (Figure 1D). EGFP expression was only observed scatteredly in the intestine, but not in the pancreas, indicating the L cell-specific Cre activity in the *IntL-Cre* mice (Figure 1E). Finally, we bred *IntL-Cre* mice with *Piezo1$^{loxp/loxp}$* mice to generate *IntL-Piezo1$^{-/-}$* mice (Figure 1F).

Under normal chow diet, *IntL-Piezo1$^{-/-}$* mice exhibited increased body weight

(Figure 1G) and greater glycemic excursions compared to control groups

(*Piezo1$^{loxp/loxp}$*, *Vil-Flp*, *FGC* and *IntL-cre*) (Figure 1H and I), while the food and

water intake were not changed (Figure 1-figure supplement 2A and B). The

morphology of islet (Figure 1-figure supplement 3A) and ileum (Figure 1-figure

supplement 4A) were not affected. Ileal mucosal Proglucagon expression and plasma

GLP-1 level were significantly lower in *IntL-Piezo1$^{-/-}$* mice than that in all littermate

controls such as *Piezo1$^{loxp/loxp}$*, *Vil-Flp*, *FGC* and *IntL-Cre* mice (Figure 1J and K),

while no significant alteration was observed in the expression of pancreatic Piezo1

and Proglucagon (Figure 1-figure supplement 3B-D), as well as ileal mucosal

cholecystokinin (CCK), another hormone secreted by L cells (Figure 1-figure

supplement 5). According to in situ hybridization of Piezo1 and GLP-1, the

expression of Piezo1 disappeared in GLP-1 positive cells, suggesting successful

knockout of Piezo1 in L cells in *IntL-Piezo1$^{-/-}$* mice (Figure 1L and M).

Piezo1 is a non-selective cationic channel that allows passage of Ca^{2+} and Na^+.

CaMKKβ is the main calcium/calmodulin dependent protein kinase kinase involved

in the regulation of metabolic homeostasis (Marcelo et al., 2016). It is activated by

binding calcium-calmodulin (Ca^{2+}/CaM), resulting in downstream activation of

kinases CaMKIV. The activation of CaMKIV modulate the gene expression of

nutrient- and hormone-related proteins (Ban et al., 2000; Chen et al., 2011;

Takemoto-Kimura et al., 2017). Previous studies have reported that Ca^{2+} and mTOR

signaling regulate the production of GLP-1 (Tolhurst et al., 2011; Xu et al., 2015; Yu

& Jin, 2010). Drawing from these findings, we hypothesized that Piezo1 might

regulate GLP-1 synthesis through the CaMKKβ/CaMKIV-mTOR signaling pathway

(Figure 1N). As shown in Figure 1O, abrogated GLP-1 production was associated

with decreased CaMKKβ/CaMKIV-mTOR signaling in the ileal mucosa of

IntL-Piezo1$^{-/-}$ mice (Figure 1O).

Derangements of glucose metabolism and glp-1 production were induced by HFD in *IntL-Piezo1⁻/⁻* mice, which was mitigated by Exendin-4

We next assessed the effect of L cell-specific *Piezo1* gene deletion on GLP-1 and glucose tolerance in diet-induced diabetic mice. *IntL-Piezo1⁻/⁻* and control mice were exposed to HFD for 10 weeks. Compared to the controls, higher body weight (Figure 2A), greater glucose excursions (Figure 2B) were observed in *IntL-Piezo1⁻/⁻* mice exposed to HFD. Ileal mucosal Proglucagon expression levels were lower in *IntL-Piezo1⁻/⁻* than control mice (Figure 2C-F). Impaired CaMKKβ/CaMKIV-mTORC1 signaling pathway in ileal mucosa as evidenced by a decrease in CaMKKβ, reduced phosphorylation levels of CaMKIV, mTOR, S6K, and S6 was also observed in *IntL-Piezo1⁻/⁻* mice (Figure 2F). No significant alteration in morphology, Piezo1 or Proglucagon levels were observed in the pancreas of *IntL-Piezo1⁻/⁻* mice (Figure 2-figure supplement 3E-H). Together these data demonstrate that *IntL-Piezo1⁻/⁻* mice with prolonged HFD feeding exhibit impaired glucose metabolism phenotype and reduced GLP-1.

Injection of GLP-1 analog Exendin-4 (Ex4) decreased the body weight (Figure 2G) and improved both glucose tolerance (Figure 2H) and insulin resistance (Figure 2I) in control and *IntL-Piezo1⁻/⁻* mice, while endogenous synthesis of GLP-1 was not changed by Ex4 injection in *IntL-Piezo1⁻/⁻* mice (Figure 2J and K). These data suggested that decreased GLP-1 synthesis and secretion contribute to impaired glucose metabolism in *IntL-Piezo1⁻/⁻* mice.

The pharmacological and mechanical activation of ileal Piezo1 stimulates GLP-1 synthesis

We next examined whether activation of Piezo1 could rescued the impaired glucose metabolism in diet-induced diabetic mice. Injection of Piezo1 activator Yoda1 after 10 weeks of high-fat diet, led to reduced body weight and improved the impaired glucose metabolism significantly in diabetic mice, while Piezo1 antagonist GsMTx4 reversed the weight loss and glucose-lowering effect of Yoda1 (Figure 3A and B). Yoda1 remarkably induced an increase in GLP-1 synthesis and secretion (Figure 3C and D), as well as an increment of CaMKKβ/CaMKIV-mTORC1 signaling in ileal mucosa (Figure 3E), while GsMTx4 abolished the effect of Yoda1 (Figure 3C-E). However, weight loss, improved plasma glucose and increased GLP-1 production induced by Yoda1 were not observed in *IntL-Piezo1$^{-/-}$* mice (Figure 3F-J).

The intestine receives mechanical stimulation from the chyme, which may activate Piezo1 in the intestine epithelium, including L cells. To mimic the mechanical pressing and stretching induced by intestinal contents, a small silicon bead was implanted into the high-fat diet-induced diabetic mouse ileum. Intestinal bead implantation improved the impaired glucose metabolism in diabetic mice (Figure 3K and L). Body weight loss, activated ileal mucosal CaMKKβ/CaMKIV-mTOR signaling, increased mRNA and protein levels of ileal mucosal Piezo1 and Proglucagon, as well as the circulating levels of GLP-1 were observed in diabetic mice after operation (Figure 3M-R). The above data suggest that mechanical stimuli induced by intestinal bead implantation activates ileal Piezo1 in diabetic mice, stimulating GLP-1 production via CaMKKβ/CaMKIV-mTOR signaling axis, thus improving glucose homeostasis.

Piezo1 regulates GLP-1 synthesis and secretion in primary cultured mouse L cells and isolated mouse ileum.

To obtain primary L cells, we isolated cell from the ileum of *IntL-Cre-mT/mG* mice, in which tdTomato expression switched to EGFP expression in L cells as shown in Figure 1E. EGFP positive cells (mouse L cells) were then sorted from isolated single cells (Figure 4A). Immunofluorescence showed that the sorted EGFP+ cells were Piezo1 positive (Figure 4B).

Yoda1 at the dose of 5μM triggered an increase in intracellular Ca^{2+} level in primary cultured mouse L cells, which was blocked by pre-incubation of cells with GsMTx4 (0.1 μM) for 15 min (Figure 4C). Yoda1 also stimulated Proglucagon expression and GLP-1 secretion, as well as CaMKKβ/CaMKIV-mTOR signaling pathway in primary cultured mouse L cells (Figure 4D-F). In contrast, knockdown of Piezo1 by shRNA led to significant decrease in Proglucagon expression and GLP-1 secretion, as well as inhibition of CaMKKβ/CaMKIV/mTOR signaling pathway (Figure 4G-J).

Given the ability of Piezo1 in sensing mechanical force, tension of 1.5g was applied to the isolated mouse ileum bathed in Tyrode's solution for two hours. Tension stimulated Proglucagon expression, GLP-1 secretion and activated CaMKKβ/CaMKIV-mTOR signaling pathway in the ileum of control mice, but not in *IntL-Piezo1^{-/-}* mice (Figure 4K-N), suggesting the involvement of Piezo1 of the L cells in mediating the force-induced GLP-1 production and CaMKKβ/CaMKIV-mTOR signaling.

Pharmacological, mechanical and genetic activation of Piezo1 stimulates GLP-1 synthesis and secretion in STC-1 cells

To further validate the role of Piezo1 in regulating GLP-1, we examined the effect of manipulating Piezo1 on GLP-1 production in an intestinal neuroendocrine cell line STC-1. Pharmacological activation of Piezo1 by Yoda1 triggered an inward current in STC-1 cell recorded by whole cell patch-clamp, which could be inhibited by pre-incubation of GsMTx4 (Figure 5A). Yoda1 also triggered an increase in intracellular Ca^{2+} level in STC-1 cells. Pre-incubation of cells with GsMTx4 (0.1 μM) for 15 min inhibited $[Ca^{2+}]_i$ increase (Figure 5B and C). Yoda1 induced a concentration-dependent activation of CaMKKβ/CaMKIV-mTOR pathway and GLP-1 synthesis and secretion (Figure 5D-F). GsMTx4 blocked the effect of Yoda1 on STC-1 cells in both GLP-1 and CaMKKβ/CaMKIV-mTOR activation (Figure 5G-I).

To mimic the activation of Piezo1 by mechanical stretching in vivo, STC-1 cells grown on elastic chambers were subjected to mechanical stretch to 120% of their original length. Mechanical stretch upregulated Piezo1 and Proglucagon expression, promoted GLP-1 secretion (Figure 5J-M), and activated CaMKKβ/CaMKIV- mTOR signaling pathways (Figure 5M).

Consistent to the pharmacological and mechanical activation of Piezo1, over-expression of Piezo1 in STC-1 cells resulted in a significant increase in GLP-1 production, as well as activation of the CaMKKβ/CaMKIV-mTOR signaling pathway (Figure 6A-D). Conversely, knockdown of Piezo1 by shRNA led to a significant decrease in GLP-1 production and inhibition of CaMKKβ/CaMKIV-mTOR signaling pathways (Figure 6E-H).

Piezo1 regulates GLP-1 production through CaMKKβ/CaMKIV and mTOR in STC-1 cells

Next, we examined whether CaMKKβ/CaMKIV and mTOR signaling mediates the effects of Piezo1 on GLP-1 production. Overexpression of CaMKKβ or CaMKIV increased CaMKKβ/CaMKIV and mTOR signaling activity, resulting in increased synthesis and secretion of GLP-1 (Figure 7A-C). In contrast, the CaMKKβ inhibitor STO-609, downregulated CaMKKβ/CaMKIV and mTOR signaling, as well as GLP-1 synthesis and secretion (Figure 7D-F). Inhibition of mTORC1 activity by rapamycin suppressed GLP-1 production induced by Yoda1, which was associated with inhibition of mTOR signalling (Figure 7G-I).

Discussion

It has been known for decades that GLP-1 secretion from the intestinal L cells is stimulated by meal intake and is essential for postprandial glycemic control (Drucker, 2006; Song et al., 2019). However, the mechanism underlying the regulation of GLP-1 production is not completely understood. One of the problems that impeded the investigation of regulation mechanism of GLP-1 is the lack of an L cell-specific genetically engineered animal model. Here, we generated an L cell-specific Cre mouse line for the first time by combination of the Flp-Frt and Cre-LoxP systems, which allows genetic manipulation specifically in the L cells and thus creates a useful tool to investigate molecular mechanisms in L cells.

Previous studies have shown that L cells are able to sense nutrients in the intestinal lumen such as glucose and other carbohydrates, lipids and amino acids, which induce GLP-1 secretion through different mechanisms, including membrane depolarization-associated exocytosis, Ca^{2+}/Calmodulin (Tolhurst et al., 2011), cAMP (Yu & Jin, 2010), mTORC1 (Xu et al., 2015) and AMPK (Jiang et al., 2016) signaling pathways. However, it is innegligible that as open type endocrine cells, L cells not only receive the chemical stimulations from the nutrients, but also mechanical stimulation when the chyme passing through the intestine, including stretching, pressure and shear force (Sensoy, 2021). Here, we showed the expression of the mechano-sensitive ion channel Piezo1 in L cells of human and mouse intestinal sections, mouse primary L cell culture and an intestinal neuroendocrine cell line STC-1, suggesting the mechano-sensing ability of L cells and potential regulation of GLP-1 in response to mechanical stimulation. Indeed, our study showed that the mechano-regulation of GLP-1 secretion did exist as demonstrated by the increased GLP-1 secretion by intestinal bead implantation, intestinal tissue stretching or STC-1 cell stretching. Moreover, mice with selective loss of Piezo1 expression in intestinal L cell (*IntL-Piezo1$^{-/-}$*) exhibited reduced circulating levels of GLP-1, increased body weight and impaired glucose homeostasis, while pharmacological activation of Piezo1 on mice, primary L cells and STC-1 cells did the reverse. More importantly,

IntL-Piezo1^{-/-} mice was unable to response to the tension-induced GLP-1 production.

These further suggested a Piezo1-mediated mechanical sensing mechanism in L cells

that regulates GLP-1 production and glucose metabolism by sensing the stimulation

of intestinal luminal contents.

Interestingly, this intestinal Piezo1-mediated mechanical sensing mechanism

may severely impaired in diabetic patients and rodents. We observed a decreased

Piezo1 expression in the ileal mucosa of diet-induced diabetic mice accompanied by

reduced GLP-1 production. When challenged with high-fat, *IntL-Piezo1^{-/-}* mice

exhibited more severe symptoms of diabetes which was mitigated by Ex-4. These

findings suggest that the impairment of Piezo1-midated mechanical sensing function

in the intestine is an important mechanism for the pathogenesis of T2DM. It is

noteworthy that RYGB, a commonly performed weight-loss and hypoglycemic

surgery (Cummings et al., 2004), significantly increased Piezo1 expression in L cells

of obese diabetic patients. Yoda1 treatment or intestinal bead implantation enhanced

GLP-1 production and improved glucose metabolism in the diet-induced diabetic

mouse model, suggesting that restore the mechano-sensing or enhance the function of

Piezo1 either pharmacologically or mechanically, may be a new strategy to improve

the secretion of GLP-1, thus alleviate T2DM. However, in our study, the

Piezo1-mediated regulation of GLP-1 production is only demonstrated in transgenic

mice, mouse primary L cells and an intestinal neuroendocrine cell line derived from

mouse. Whether Piezo1 plays the same role in human L cells awaits to be investigated.

A number of studies have generated L cells culture from human intestinal organoid

culture or human intestinal stem cell monolayer culture by manipulating the growth

factors in the media (Goldspink et al., 2020; Petersen et al., 2014; Villegas-Novoa et

al., 2022). It is worthy to validate our finding in human L cells in order to prove its

translational potential in T2DM treatment.

The intragastric balloon is a current noninvasive clinical weight loss measure

that involves placing a space-occupying balloon in the stomach to reduce food intake

and generate satiety signals, thus maintaining satiety. Investigations illustrated that

intragastric balloon alter the secretion of hormones such as cholecystokinin and pancreatic polypeptide, delay the emptying of food in the stomach and reduce the appetite (Mathus-Vliegen & de Groot, 2013). Intragastric balloon provides a feasible weight loss intervention for obese people (Kim et al., 2016). In this study, a new intestinal implantation surgery of beads was adopted, which may offer a novel approach for weight loss and glucose control by activating the intestinal Piezo1-GLP-1 signaling pathway in the future.

Mechanistically, our study suggest that Piezo1 regulates GLP-1 production through a CaMKKβ/CaMKIV-mTOR signaling pathway. CaMKKβ/CaMKIV has been reported to mediate the Ca^{2+} signaling in many metabolic processes, including liver gluconeogenesis and de novo lipogenesis, adipogenesis, insulin sensitivity and β cell proliferation (Anderson et al., 2012; Lin et al., 2011; Liu et al., 2012; J. Liu et al., 2022). mTOR plays a central role in nutrient and energy sensing and regulates cellular metabolism and growth in response to different nutrient and energy status (Howell & Manning, 2011). Here we showed that mTOR can also response to mechanical stimuli through a mechano-sensitive Ca^{2+} channel mediated CaMKKβ/CaMKIV activation. Although we did not demonstrate direct phosphorylation of mTOR or S6K by CaMKIV in L cells, previous study reported that CaMKK2 could serve as a scaffold to assemble CaMKIV with key components of the mTOR/S6K pathway and promote liver cancer cell growth (Lin et al., 2015), which lend support to the CaMKKβ/CaMKIV-mTOR signaling in our study. Recently, Knutson et. al. found that ryanodine and IP3-triggered calcium release from intracellular calcium store could amplified the initial Peizo2 - Ca^{2+} signal triggered by mechanical stimulation, and was required for the mechanotransduction in the serotonin release from enterochromaffin cells (Knutson et al., 2023). In our study, we also observed long-lasting intracellular Ca^{2+} increase triggered by Yoda1 in primary L cells and STC-1 cells, which also suggested an involvement of intracellular Ca^{2+} store in the Ca^{2+} relay. Beside Ca^{2+}, cyclic AMP (cAMP) is another signaling molecule that active *Gcg* gene expression and GLP-1 production (Drucker et al., 1994; Jin, 2008; Simpson

et al., 2007). cAMP was found to play a critical role in nutrients-induced GLP-1 secretion, including glucose (Ong et al., 2009), lipids (Hodge et al., 2016), and amino acids (Tolhurst et al., 2011). Previous study reported that Ca^{2+} can activate soluble adenylyl cyclase (sAC) to increase intracellular cAMP (Jaiswal & Conti, 2003). Whether sAC-cAMP can be activated by Piezo1-mediated Ca^{2+} influx and whether it is an alternative signaling pathway that mediates the Piezo1-regulated GLP-1 production remain to be explored.

In summary, our study reveals a previously undiscovered Piezo1-mediated mechano-sensing property of intestinal L cell, which plays an essential role in the regulation of GLP-1 production and glucose metabolism. This finding also suggests a new mechano-regulation in enteroendocrine cells, in addition to chemical and neuronal regulation, which may shed new light on the strategy for metabolic diseases such as diabetes and obesity.

Acknowledgements

Funding

This work was supported by grants from the National Natural Science Foundation

of China (82170818, 81770794), Guangdong Basic and Applied Basic Research

Foundation (2024A1515010686), the Fundamental Research Funds for the Central

Universities (21620423), Guangzhou Science and Technology Plan Project Funding

(202201011353).

Conflicts of interest

On behalf of all authors, the corresponding author states that there is no conflict

of interest.

Author contribution

Geyang Xu and Hui Chen designed research; Yanling Huang, Haocong Mo, Jie

Yang, Luyang Gao, Tian Tao, Qing Shu, Wenying Guo, Yawen Zhao, Jingya Lyu,

Qimeng Wang, Jinghui Guo, Hening Zhai, Linyan Zhu and Hui Chen performed

research; Yanling Huang, Haocong Mo, Jie Yang, Hui Chen and Geyang Xu analyzed

data; Geyang Xu and Hui Chen wrote and edited the paper. All authors contributed to

the discussion and revision of the article and all approved the final versions of the

manuscript. Geyang Xu is responsible for the integrity of the work as a whole.

Ethical Statement

Animals used in this study were handled in accordance with the Guide for the

Care and Use of Laboratory Animals published by the National Institutes of Health

(NIH Publications No. 8023, revised 1978). All animal protocols were approved by

the Animal Care and Use Committee of Jinan University.

Human subject research was approved by the Institutional Review Board of

Jinan University.

Reference

Alcaino, C., Knutson, K. R., Treichel, A. J., Yildiz, G., Strege, P. R., Linden, D. R., . . . Beyder, A. (2018). A population of gut epithelial enterochromaffin cells is mechanosensitive and requires Piezo2 to convert force into serotonin release. *Proc Natl Acad Sci U S A*, *115*(32), E7632-e7641. https://doi.org/10.1073/pnas.1804938115

Anderson, K. A., Lin, F., Ribar, T. J., Stevens, R. D., Muehlbauer, M. J., Newgard, C. B., & Means, A. R. (2012). Deletion of CaMKK2 from the liver lowers blood glucose and improves whole-body glucose tolerance in the mouse. *Mol Endocrinol*, *26*(2), 281-291. https://doi.org/10.1210/me.2011-1299

Anini, Y., Hansotia, T., & Brubaker, P. L. (2002). Muscarinic receptors control postprandial release of glucagon-like peptide-1: in vivo and in vitro studies in rats. *Endocrinology*, *143*(6), 2420-2426. https://doi.org/10.1210/endo.143.6.8840

Atanga, R., Singh, V., & In, J. G. (2023). Intestinal Enteroendocrine Cells: Present and Future Druggable Targets. *Int J Mol Sci*, *24*(10). https://doi.org/10.3390/ijms24108836

Ban, N., Yamada, Y., Someya, Y., Ihara, Y., Adachi, T., Kubota, A., . . . Seino, Y. (2000). Activating transcription factor-2 is a positive regulator in CaM kinase IV-induced human insulin gene expression. *Diabetes*, *49*(7), 1142-1148. https://doi.org/10.2337/diabetes.49.7.1142

Bany Bakar, R., Reimann, F., & Gribble, F. M. (2023). The intestine as an endocrine organ and the role of gut hormones in metabolic regulation. *Nat Rev Gastroenterol Hepatol*. https://doi.org/10.1038/s41575-023-00830-y

Bezin, J., Gouverneur, A., Pénichon, M., Mathieu, C., Garrel, R., Hillaire-Buys, D., . . . Faillie, J. L. (2023). GLP-1 Receptor Agonists and the Risk of Thyroid Cancer. *Diabetes Care*, *46*(2), 384-390. https://doi.org/10.2337/dc22-1148

Cahalan, S. M., Lukacs, V., Ranade, S. S., Chien, S., Bandell, M., & Patapoutian, A. (2015). Piezo1 links mechanical forces to red blood cell volume. *Elife*, *4*. https://doi.org/10.7554/eLife.07370

Chen, K., Yu, X., Murao, K., Imachi, H., Li, J., Muraoka, T., . . . Tokumitsu, H. (2011). Exendin-4 regulates GLUT2 expression via the CaMKK/CaMKIV pathway in a pancreatic β-cell line. *Metabolism*, *60*(4), 579-585. https://doi.org/10.1016/j.metabol.2010.06.002

Cummings, D. E., Overduin, J., & Foster-Schubert, K. E. (2004). Gastric bypass for obesity: mechanisms of weight loss and diabetes resolution. *J Clin Endocrinol Metab*, *89*(6), 2608-2615. https://doi.org/10.1210/jc.2004-0433

Deivasikamani, V., Dhayalan, S., Abudushalamu, Y., Mughal, R., Visnagri, A.,
Cuthbertson, K., . . . Sukumar, P. (2019). Piezo1 channel activation mimics high
glucose as a stimulator of insulin release. *Sci Rep*, *9*(1), 16876.
https://doi.org/10.1038/s41598-019-51518-w

Diakogiannaki, E., Gribble, F. M., & Reimann, F. (2012). Nutrient detection by
incretin hormone secreting cells. *Physiol Behav*, *106*(3), 387-393.
https://doi.org/10.1016/j.physbeh.2011.12.001

Drucker, D. J. (2006). The biology of incretin hormones. *Cell Metab*, *3*(3), 153-165.
https://doi.org/10.1016/j.cmet.2006.01.004

Drucker, D. J., Jin, T., Asa, S. L., Young, T. A., & Brubaker, P. L. (1994). Activation
of proglucagon gene transcription by protein kinase-A in a novel mouse
enteroendocrine cell line. *Mol Endocrinol*, *8*(12), 1646-1655.
https://doi.org/10.1210/mend.8.12.7535893

Elashoff, M., Matveyenko, A. V., Gier, B., Elashoff, R., & Butler, P. C. (2011).
Pancreatitis, pancreatic, and thyroid cancer with glucagon-like peptide-1-based
therapies. *Gastroenterology*, *141*(1), 150-156.
https://doi.org/10.1053/j.gastro.2011.02.018

Furness, J. B., Rivera, L. R., Cho, H. J., Bravo, D. M., & Callaghan, B. (2013). The
gut as a sensory organ. *Nat Rev Gastroenterol Hepatol*, *10*(12), 729-740.
https://doi.org/10.1038/nrgastro.2013.180

Goldspink, D. A., Lu, V. B., Miedzybrodzka, E. L., Smith, C. A., Foreman, R. E.,
Billing, L. J., . . . Gribble, F. M. (2020). Labeling and Characterization of Human
GLP-1-Secreting L-cells in Primary Ileal Organoid Culture. *Cell Rep*, *31*(13), 107833.
https://doi.org/10.1016/j.celrep.2020.107833

Gribble, F. M., & Reimann, F. (2016). Enteroendocrine Cells: Chemosensors in the
Intestinal Epithelium. *Annu Rev Physiol*, *78*, 277-299.
https://doi.org/10.1146/annurev-physiol-021115-105439

Gudipaty, S. A., Lindblom, J., Loftus, P. D., Redd, M. J., Edes, K., Davey, C. F., . . .
Rosenblatt, J. (2017). Mechanical stretch triggers rapid epithelial cell division through
Piezo1. *Nature*, *543*(7643), 118-121. https://doi.org/10.1038/nature21407

Hodge, D., Glass, L. L., Diakogiannaki, E., Pais, R., Lenaghan, C., Smith, D. M., . . .
Reimann, F. (2016). Lipid derivatives activate GPR119 and trigger GLP-1 secretion in
primary murine L-cells. *Peptides*, *77*, 16-20.
https://doi.org/10.1016/j.peptides.2015.06.012

Howell, J. J., & Manning, B. D. (2011). mTOR couples cellular nutrient sensing to organismal metabolic homeostasis. *Trends Endocrinol Metab*, *22*(3), 94-102. https://doi.org/10.1016/j.tem.2010.12.003

Jaiswal, B. S., & Conti, M. (2003). Calcium regulation of the soluble adenylyl cyclase expressed in mammalian spermatozoa. *Proc Natl Acad Sci U S A*, *100*(19), 10676-10681. https://doi.org/10.1073/pnas.1831008100

Jiang, S., Zhai, H., Li, D., Huang, J., Zhang, H., Li, Z., . . . Xu, G. (2016). AMPK-dependent regulation of GLP1 expression in L-like cells. *J Mol Endocrinol*, *57*(3), 151-160. https://doi.org/10.1530/jme-16-0099

Jiang, Y., Song, J., Xu, Y., Liu, C., Qian, W., Bai, T., & Hou, X. (2021). Piezo1 regulates intestinal epithelial function by affecting the tight junction protein claudin-1 via the ROCK pathway. *Life Sci*, *275*, 119254. https://doi.org/10.1016/j.lfs.2021.119254

Jin, T. (2008). Mechanisms underlying proglucagon gene expression. *J Endocrinol*, *198*(1), 17-28. https://doi.org/10.1677/joe-08-0085

Kim, S. H., Chun, H. J., Choi, H. S., Kim, E. S., Keum, B., & Jeen, Y. T. (2016). Current status of intragastric balloon for obesity treatment. *World J Gastroenterol*, *22*(24), 5495-5504. https://doi.org/10.3748/wjg.v22.i24.5495

Knutson, K. R., Whiteman, S. T., Alcaino, C., Mercado-Perez, A., Finholm, I., Serlin, H. K., . . . Beyder, A. (2023). Intestinal enteroendocrine cells rely on ryanodine and IP(3) calcium store receptors for mechanotransduction. *J Physiol*, *601*(2), 287-305. https://doi.org/10.1113/jp283383

Lai, A., Cox, C. D., Chandra Sekar, N., Thurgood, P., Jaworowski, A., Peter, K., & Baratchi, S. (2022). Mechanosensing by Piezo1 and its implications for physiology and various pathologies. *Biol Rev Camb Philos Soc*, *97*(2), 604-614. https://doi.org/10.1111/brv.12814

Li, J., Hou, B., Tumova, S., Muraki, K., Bruns, A., Ludlow, M. J., . . . Beech, D. J. (2014). Piezo1 integration of vascular architecture with physiological force. *Nature*, *515*(7526), 279-282. https://doi.org/10.1038/nature13701

Li, Z., Bowers, E., Zhu, J., Yu, H., Hardij, J., Bagchi, D. P., . . . MacDougald, O. A. (2022). Lipolysis of bone marrow adipocytes is required to fuel bone and the marrow niche during energy deficits. *Elife*, *11*. https://doi.org/10.7554/eLife.78496

Lin, F., Marcelo, K. L., Rajapakshe, K., Coarfa, C., Dean, A., Wilganowski, N., . . . York, B. (2015). The camKK2/camKIV relay is an essential regulator of hepatic cancer. *Hepatology*, *62*(2), 505-520. https://doi.org/10.1002/hep.27832

Lin, F., Ribar, T. J., & Means, A. R. (2011). The Ca2+/calmodulin-dependent protein kinase kinase, CaMKK2, inhibits preadipocyte differentiation. *Endocrinology*, *152*(10), 3668-3679. https://doi.org/10.1210/en.2011-1107

Liu, B., Barbosa-Sampaio, H., Jones, P. M., Persaud, S. J., & Muller, D. S. (2012). The CaMK4/CREB/IRS-2 cascade stimulates proliferation and inhibits apoptosis of β-cells. *PLoS One*, *7*(9), e45711. https://doi.org/10.1371/journal.pone.0045711

Liu, H., Hu, J., Zheng, Q., Feng, X., Zhan, F., Wang, X., . . . Hua, F. (2022). Piezo1 Channels as Force Sensors in Mechanical Force-Related Chronic Inflammation. *Front Immunol*, *13*, 816149. https://doi.org/10.3389/fimmu.2022.816149

Liu, J., Li, Y., Gao, N., Ji, J., & He, Q. (2022). Calcium/calmodulin-dependent protein kinase IV regulates vascular autophagy and insulin signaling through Akt/mTOR/CREB pathway in ob/ob mice. *J Physiol Biochem*, *78*(1), 199-211. https://doi.org/10.1007/s13105-021-00853-6

Marcelo, K. L., Means, A. R., & York, B. (2016). The Ca(2+)/Calmodulin/CaMKK2 Axis: Nature's Metabolic CaMshaft. *Trends Endocrinol Metab*, *27*(10), 706-718. https://doi.org/10.1016/j.tem.2016.06.001

Mathus-Vliegen, E. M., & de Groot, G. H. (2013). Fasting and meal-induced CCK and PP secretion following intragastric balloon treatment for obesity. *Obes Surg*, *23*(5), 622-633. https://doi.org/10.1007/s11695-012-0834-6

Maunoury, R., Robine, S., Pringault, E., Léonard, N., Gaillard, J. A., & Louvard, D. (1992). Developmental regulation of villin gene expression in the epithelial cell lineages of mouse digestive and urogenital tracts. *Development*, *115*(3), 717-728. https://doi.org/10.1242/dev.115.3.717

Ong, W. K., Gribble, F. M., Reimann, F., Lynch, M. J., Houslay, M. D., Baillie, G. S., . . . Pyne, N. J. (2009). The role of the PDE4D cAMP phosphodiesterase in the regulation of glucagon-like peptide-1 release. *Br J Pharmacol*, *157*(4), 633-644. https://doi.org/10.1111/j.1476-5381.2009.00194.x

Petersen, N., Reimann, F., Bartfeld, S., Farin, H. F., Ringnalda, F. C., Vries, R. G., . . . de Koning, E. J. (2014). Generation of L cells in mouse and human small intestine organoids. *Diabetes*, *63*(2), 410-420. https://doi.org/10.2337/db13-0991

Romac, J. M., Shahid, R. A., Swain, S. M., Vigna, S. R., & Liddle, R. A. (2018). Piezo1 is a mechanically activated ion channel and mediates pressure induced pancreatitis. *Nat Commun*, *9*(1), 1715. https://doi.org/10.1038/s41467-018-04194-9

Rouillé, Y., Kantengwa, S., Irminger, J. C., & Halban, P. A. (1997). Role of the prohormone convertase PC3 in the processing of proglucagon to glucagon-like

peptide 1. *J Biol Chem*, *272*(52), 32810-32816.
https://doi.org/10.1074/jbc.272.52.32810

Rutlin, M., Rastelli, D., Kuo, W. T., Estep, J. A., Louis, A., Riccomagno, M. M., . . .
Rao, M. (2020). The Villin1 Gene Promoter Drives Cre Recombinase Expression in
Extraintestinal Tissues. *Cell Mol Gastroenterol Hepatol*, *10*(4), 864-867.e865.
https://doi.org/10.1016/j.jcmgh.2020.05.009

Saxena, A. R., Gorman, D. N., Esquejo, R. M., Bergman, A., Chidsey, K., Buckeridge,
C., . . . Kim, A. M. (2021). Danuglipron (PF-06882961) in type 2 diabetes: a
randomized, placebo-controlled, multiple ascending-dose phase 1 trial. *Nat Med*,
27(6), 1079-1087. https://doi.org/10.1038/s41591-021-01391-w

Sensoy, I. (2021). A review on the food digestion in the digestive tract and the used in
vitro models. *Curr Res Food Sci*, *4*, 308-319.
https://doi.org/10.1016/j.crfs.2021.04.004

Simpson, A. K., Ward, P. S., Wong, K. Y., Collord, G. J., Habib, A. M., Reimann, F.,
& Gribble, F. M. (2007). Cyclic AMP triggers glucagon-like peptide-1 secretion from
the GLUTag enteroendocrine cell line. *Diabetologia*, *50*(10), 2181-2189.
https://doi.org/10.1007/s00125-007-0750-9

Song, Y., Koehler, J. A., Baggio, L. L., Powers, A. C., Sandoval, D. A., & Drucker, D.
J. (2019). Gut-Proglucagon-Derived Peptides Are Essential for Regulating Glucose
Homeostasis in Mice. *Cell Metab*, *30*(5), 976-986.e973.
https://doi.org/10.1016/j.cmet.2019.08.009

Takemoto-Kimura, S., Suzuki, K., Horigane, S. I., Kamijo, S., Inoue, M., Sakamoto,
M., . . . Bito, H. (2017). Calmodulin kinases: essential regulators in health and disease.
J Neurochem, *141*(6), 808-818. https://doi.org/10.1111/jnc.14020

Tan, Q., Akindehin, S. E., Orsso, C. E., Waldner, R. C., DiMarchi, R. D., Müller, T. D.,
& Haqq, A. M. (2022). Recent Advances in Incretin-Based Pharmacotherapies for the
Treatment of Obesity and Diabetes. *Front Endocrinol (Lausanne)*, *13*, 838410.
https://doi.org/10.3389/fendo.2022.838410

Tolhurst, G., Zheng, Y., Parker, H. E., Habib, A. M., Reimann, F., & Gribble, F. M.
(2011). Glutamine triggers and potentiates glucagon-like peptide-1 secretion by
raising cytosolic Ca2+ and cAMP. *Endocrinology*, *152*(2), 405-413.
https://doi.org/10.1210/en.2010-0956

Treichel, A. J., Finholm, I., Knutson, K. R., Alcaino, C., Whiteman, S. T., Brown, M.
R., . . . Beyder, A. (2022). Specialized Mechanosensory Epithelial Cells in Mouse Gut
Intrinsic Tactile Sensitivity. *Gastroenterology*, *162*(2), 535-547.e513.
https://doi.org/10.1053/j.gastro.2021.10.026

Villegas-Novoa, C., Wang, Y., Sims, C. E., & Allbritton, N. L. (2022). Development
of a Primary Human Intestinal Epithelium Enriched in L-Cells for Assay of GLP-1
Secretion. *Anal Chem*, *94*(27), 9648-9655.
https://doi.org/10.1021/acs.analchem.2c00912

Wang, J., Sun, Y. X., & Li, J. (2023). The role of mechanosensor Piezo1 in bone
homeostasis and mechanobiology. *Dev Biol*, *493*, 80-88.
https://doi.org/10.1016/j.ydbio.2022.11.002

Wang, S., Chennupati, R., Kaur, H., Iring, A., Wettschureck, N., & Offermanns, S.
(2016). Endothelial cation channel PIEZO1 controls blood pressure by mediating
flow-induced ATP release. *J Clin Invest*, *126*(12), 4527-4536.
https://doi.org/10.1172/jci87343

Willms, B., Werner, J., Holst, J. J., Orskov, C., Creutzfeldt, W., & Nauck, M. A.
(1996). Gastric emptying, glucose responses, and insulin secretion after a liquid test
meal: effects of exogenous glucagon-like peptide-1 (GLP-1)-(7-36) amide in type 2
(noninsulin-dependent) diabetic patients. *J Clin Endocrinol Metab*, *81*(1), 327-332.
https://doi.org/10.1210/jcem.81.1.8550773

Xu, G., Li, Z., Ding, L., Tang, H., Guo, S., Liang, H., . . . Zhang, W. (2015). Intestinal
mTOR regulates GLP-1 production in mouse L cells. *Diabetologia*, *58*(8), 1887-1897.
https://doi.org/10.1007/s00125-015-3632-6

Xu, Y., Bai, T., Xiong, Y., Liu, C., Liu, Y., Hou, X., & Song, J. (2021). Mechanical
stimulation activates Piezo1 to promote mucin2 expression in goblet cells. *J
Gastroenterol Hepatol*, *36*(11), 3127-3139. https://doi.org/10.1111/jgh.15596

Ye, Y., Barghouth, M., Dou, H., Luan, C., Wang, Y., Karagiannopoulos, A., . . .
Renström, E. (2022). A critical role of the mechanosensor PIEZO1 in glucose-induced
insulin secretion in pancreatic β-cells. *Nat Commun*, *13*(1), 4237.
https://doi.org/10.1038/s41467-022-31103-y

Yu, Z., & Jin, T. (2010). New insights into the role of cAMP in the production and
function of the incretin hormone glucagon-like peptide-1 (GLP-1). *Cell Signal*, *22*(1),
1-8. https://doi.org/10.1016/j.cellsig.2009.09.032

Zhai, H., Li, Z., Peng, M., Huang, Z., Qin, T., Chen, L., . . . Xu, G. (2018). Takeda G
Protein-Coupled Receptor 5-Mechanistic Target of Rapamycin Complex 1 Signaling
Contributes to the Increment of Glucagon-Like Peptide-1 Production after Roux-en-Y
Gastric Bypass. *EBioMedicine*, *32*, 201-214.
https://doi.org/10.1016/j.ebiom.2018.05.026

 Figure legends

Figure 1: Generation, Validation and Characterization of *IntL-Piezo1$^{-/-}$* mice

(**A**) Schematic description for the generation of Villin-Flippase (*Vil-Flp*) and

Flippase-dependent Gcg-Cre (*FGC*) mice. Vil-Flp flip the inverted Cre gene in the

Gcg-Cre cassette in *IntL-Cre* mice to restrict Cre expression in intestinal L cells. As

shown, Locations of genotyping primers are also indicated.

(**B**) Tail DNA genotyping PCR results using genotyping primer for *Vil-Flp*, *FGC* and

Flippase-activated Cre (*IntL-Cre*) mice.

(**C**) Intestine and pancreas DNA genotyping results. The "Original" band represents

the original FGC cassette with inverted Cre, while the "Flipped" band represents

recombined FGC cassette with Cre flipped into the correct direction.

(**D**) Schematic description for the validation of *IntL-Cre* efficacy by crossing with

mT/mG reporter mice.

(**E**) Fluorescence was detected in the ileal and pancreatic tissues from *mT/mG* and

IntL-Cre-mT/mG mice by frozen tissue confocal microscopy. Green fluorescence

represents successful deletion of TdTomato and reactivation of EGFP in the

Cre-expressing cells.

(**F**) Schematic description for the generation of Intestinal L cell-Piezo1$^{-/-}$ mice

(*IntL-Piezo1$^{-/-}$*) by crossing *Piezo1$^{loxp/loxp}$* mice with *IntL-Cre* mice.

(**G**) Body weight of 14- to 16-week-old male mice of the indicated genotypes fed

NCD (n=6/group).

(**H-I**) IPGTT (**H**) and ITT (**I**) and associated area under the curve (AUC) values of

14- to 16-week-old male mice of the indicated genotypes fed NCD (n=6/group).

(**J**) *Proglucagon* mRNA levels in ileum of 14- to 16-week-old male mice of the

indicated genotypes fed NCD. (n=6/group).

(**K**) The plasma GLP-1 levels in 14- to 16-week-old male mice of the indicated

genotypes fed NCD (n=6/group).

(**L**) Representative images for Piezo1 RNA-FISH and GLP-1 immunofluorescent

staining in the ileum of 14-week-old male mice of indicated genotypes fed NCD

(n=6/group).

(**M**) Quantitative analysis of Piezo1-positive GLP-1 cells in the ileal mucosa of

14-week-old male mice of indicated genotypes fed NCD (n=6/group).

(**N**) A schematic diagram depicting the potential mechanisms linking the

CaMKKβ/CaMKIV-mTOR signaling pathway and GLP-1 production.

(**O**) Representative western blots are shown for indicated antibodies in the ileal

mucosa (n = 6/group).

Data are represented as mean ± SEM. Significance was determined by Student's t test

for comparison between two groups, and by one-way ANOVA for comparison among

three groups or more, *p < 0.05, **p < 0.01, ***p < 0.001.

Figure 2: Validation and phenotype of *IntL-Piezo1$^{-/-}$* mice fed with high-fat diet

(**A**) Body weight of 14- to 16-week-old male *Piezo1$^{loxp/loxp}$* and *IntL-Piezo1$^{-/-}$* mice fed with HFD for 10 weeks (n=6/group).

(**B**) IPGTT and associated area under the curve (AUC) values of 14- to 16-week-old male *Piezo1$^{loxp/loxp}$* and *IntL-Piezo1$^{-/-}$* mice fed with HFD (n=6/group).

(**C**) *Proglucagon* mRNA levels in the ileal mucosa of 14- to 16-week-old male *Piezo1$^{loxp/loxp}$* and *IntL-Piezo1$^{-/-}$* mice fed with HFD (n=6/group).

(**D**) The plasma GLP-1 level in 14- to 16-week-old male *Piezo1$^{loxp/loxp}$* and *IntL-Piezo1$^{-/-}$* mice fed with HFD (n=6/group).

(**E**) Double immunofluorescent staining of Piezo1, and GLP-1 in the ilea of 14- to 16-week-old male *Piezo1$^{loxp/loxp}$* and *IntL-Piezo1$^{-/-}$* mice fed HFD (n=6/group).

(**F**) Representative western blots are shown for indicated antibodies in the ileal mucosa (n=6/group).

(**G**) Body weight after 7 consecutive days infusion of saline or Ex4 (100ug/kg body weight) in 14- to 16-week-old male *Piezo1$^{loxp/loxp}$* and *IntL-Piezo1$^{-/-}$* mice fed with HFD (n=6/group).

(**H-I**) IPGTT (**H**) and ITT (**I**) and associated area under the curve (AUC) values after consecutive infusion of saline or Ex4.

(**J**) *Proglucagon* mRNA levels in the ileal mucosa (n=6/group) after consecutive infusion of saline or Ex4.

(**K**) The plasma GLP-1 level after consecutive infusion of saline or Ex4 (n=6/group).

Data are represented as mean ± SEM. Significance was determined by Student's t test for comparison between two groups, and by one-way ANOVA for comparison among three groups or more, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure 3: Chemical and mechanical interventions of Piezo1 regulates GLP-1 synthesis in mice

(**A-E**) 14- to 16-week-old male C57BL/6J mice fed with HFD for 10 weeks were infused with vehicle, Yoda1 (2 μg per mouse) or GsMTx4 (250 μg/kg) by i.p. for 7 consecutive days. (n=6/group).

(**A**) Body weight change after consecutive drug infusion.

(**B**) IPGTT and associated area under the curve (AUC) values.

(**C**) *Proglucagon* mRNA levels in the ileal mucosa.

(**D**) The plasma GLP-1 level.

(**E**) Representative western blots are shown for indicated antibodies in the ileal mucosa.

(**F-J**) 14- to 16-week-old male *IntL-Piezo1$^{-/-}$* mice fed with HFD for 10 weeks were infused with vehicle, Yoda1 (2 μg per mouse) by i.p. for 7 consecutive days. (n=4 or 5/group)

(**F**) Body weight change after 7 consecutive days' drug infusion.

(**G**) Fasting blood glucose levels.

(**H**) Ileal mucosal *Proglucagon* mRNA levels.

(**I**) Plasma GLP-1 levels.

(**J**) Ileal mucosal Proglucagon protein levels.

(**K-R**) 14- to 16-week-old male C57BL/6J mice fed with HFD were subjected to sham operation, or intestinal bead implantation (n=6/group).

(**K**) Fasting blood glucose levels.

(**L**) IPGTT and associated area under the curve (AUC) values.

(**M**) Body weight.

(**N** and **O**) *Piezo1* (**N**) and *Proglucagon* (**O**) mRNA levels in the ileal mucosa.

(**P**) The plasma GLP-1 levels.

(**Q**) Immunofluorescence staining of GLP-1 in ileum and quantification of GLP-1 postive cells.

(**R**) Representative western blots images and densitometry quantification for indicated antibodies in the ileal mucosa.

Data are represented as mean ± SEM. Significance was determined by Student's t test for comparison between two groups, and by one-way ANOVA for comparison among three groups or more, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure 4: Piezo1 regulates GLP-1 synthesis and secretion in primary cultured mouse L cells and isolated mouse ileum.

(**A**) Isolation of mouse L cells (GFP positive) from ileal tissue by FACS. The gating in flowcytometry for sorting of GFP positive cells.

(**B**) Immunofluorescent staining of Piezo1 in sorted GFP positive L cells.

(**C**) Intracellular Ca^{2+} imaging by fluo-4-AM calcium probe. The change of fluorescent intensity ($\Delta F/F0$) was plotted against time.

(**D-F**) L cells were treated with vehicle or Yoda1 (5μM) for 24 hours.

(**D**) *Proglucagon* mRNA expression.

(**E**) GLP-1 concentrations in the culture medium.

(**F**) Western blot images and densitometry quantification for the indicated antibodies.

(**G-J**) Knockdown of Piezo1 in L cells by shRNA for 48 hours.

(**G**) *Piezo1* mRNA expression.

(**H**) *Proglucagon* mRNA expression.

(**I**) GLP-1 levels in the culture medium.

(**J**) Western blot images and densitometry quantification for the indicated antibodies.

(**K-N**) Ileal tissues from *Piezo1^loxp/loxp^* and *IntL-Piezo1^-/-^* mice were subjected to tension force (n=6/group).

(**K**) A representative photograph showing the traction of isolated ileum.

(**L**) *Proglucagon* mRNA levels.

(**M**) GLP-1 concentrations in the medium.

(**N**) Western blot images and densitometry quantification for the indicated antibodies.

Data are represented as mean ± SEM. n=6. Significance was determined by Student's t test for comparison between two groups, and by one-way ANOVA for comparison among three groups or more, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$.

Figure 5: Modulation of GLP-1 synthesis and secretion by pharmacological and
mechanical activation of Piezo1 in STC-1 cells

(**A**) Whole-cell currents induced by Yoda1 (5μM) were recorded from STC-1 cells or
STC-1 cells pretreated with GsMTx4 for 30 min.

(**B** and **C**) Intracellular calcium imaging in STC-1 cells. (**B**) STC-1 cells were loaded
with fluo-4 AM for 1 h. The representative time-lapse image showing the intracellular
Ca^{2+} signals. (**C**) The change of fluorescent intensity ($\Delta F/F0$) was plotted against
time.

(**D-F**) STC-1 cells were treated with various concentrations of Yoda1 for 24 h. (**D**)
Whole-cell extracts underwent western blot with indicated antibodies. (**E**)
Proglucagon mRNA levels. (**F**) GLP-1 concentrations in the culture medium.

(**G-I**) STC-1 cells were treated with Yoda1 (5 μM) in the presence or absence of
GsMTx4 (0.1 μM) for 24 h. (**G**) Whole-cell extracts underwent western blot with
indicated antibodies. (**H**) *Proglucagon* mRNA levels. (**I**) GLP-1 concentrations in the
culture medium.

(**J-M**) STC-1 were subjected to mechanical stretch. (**J**) STC-1 cells were cultured in
elastic chambers and the chambers were subjected to mechanical stretch by 120%
extension of their original length. The medium GLP-1 concentrations were detected at
indicated time. (**K**) *Piezo1* mRNA levels. (**L**) *Proglucagon* mRNA levels (**M**)
Whole-cell extracts underwent western blot with indicated antibodies.

Data are represented as mean ± SEM. n=6. Significance was determined by Student's
t test for comparison between two groups, and by one-way ANOVA for comparison
among three groups or more, *p < 0.05, **p < 0.01, ***p < 0.001.

1022

Figure 6: Genetic interference of Piezo1 regulates GLP-1 production in STC-1 cells

(**A-D**) STC-1 cells were transfected with mouse control or *Piezo1* expression plasmids for 48h. *Piezo1* (**A**) and *Proglucagon* (**B**) mRNA levels in STC-1 cells. (**C**) GLP-1 concentrations in culture medium. (**D**) Whole-cell extracts underwent western blot with indicated antibodies.

(**E-H**) Stable knockdown of Piezo1 in STC-1 cells. *Piezo1* (**E**) and *Proglucagon* (**F**) mRNA levels in STC-1 cells. (**G**) GLP-1 concentrations in culture medium. (**H**) Whole-cell extracts underwent western blot with indicated antibodies.

Data are represented as mean ± SEM. n=6. Significance was determined by Student's t test, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$.

 Figure 7: Modulation of GLP-1 production by CaMKKβ/CaMKIV and mTOR
signaling activity in STC-1 cells

(**A-C**) STC-1 cells were transfected with GFP, CaMKKβ or CaMKIV plasmids for 48h. (**A**) *Proglucagon* mRNA levels in STC-1 cells. (**B**) GLP-1 concentrations in culture medium. (**C**) Whole-cell extracts underwent western blot with indicated antibodies.

(**D-F**) STC-1 cells were treated with CaMKKβ inhibitor STO-609 (10 μmol/L) for 24 h. (**D**) *Proglucagon* mRNA levels in STC-1 cells. (**E**) GLP-1 concentrations in culture medium. (**F**) Whole-cell extracts underwent western blot with indicated antibodies.

(**G-I**) STC-1 cells were pretreated with Rapamycin (50 nmol/L) for 1 h, then treated with Yoda1 (5 μmol/L) for 24 h. (**G**) *Proglucagon* mRNA levels in STC-1 cells. (**H**) GLP-1 concentrations in the culture medium. (**I**) Whole-cell extracts underwent western blot with indicated antibodies.

Data are represented as mean ± SEM. n=6. Significance was determined by Student's t test for comparison between two groups, and by one-way ANOVA for comparison among three groups or more, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7