1
2
Pseudocobalamin production and use in marine *Synechococcus* cultures and communities

Catherine C. Bannon[1], Maria A. Soto[1], Elden Rowland[1], Nan Chen[1], Anna Gleason[1], Emmanuel
Devred[2], Julie LaRoche[1], Erin M. Bertrand[1]*

[1]Department of Biology and Institute for Comparative Genomics, Dalhousie University, Halifax,
Nova Scotia, Canada
[2]Bedford Institute of Oceanography, Fisheries and Oceans Canada, Dartmouth, Nova Scotia,
Canada

* Correspondence:
erin.bertrand@dal.ca

Running head: psB$_{12}$ in *Synechococcus*

Keywords: marine vitamin dynamics, methionine synthase, North Atlantic Ocean,
picocyanobacteria, cobalamin

Abstract

Cobalamin influences marine microbial communities because an exogenous source is required by most eukaryotic phytoplankton, and demand can exceed supply. Pseudocobalamin is a cobalamin analog that is produced and used by most cyanobacteria but is not directly available to eukaryotic phytoplankton. Some microbes can remodel pseudocobalamin into cobalamin, but a scarcity of pseudocobalamin measurements impedes our ability to evaluate its importance for marine cobalamin production. Here, we perform simultaneous measurements of pseudocobalamin and methionine synthase (MetH), the key protein that uses it as a co-factor, in *Synechococcus* cultures and communities. In *Synechococcus* sp. WH8102, pseudocobalamin quota decreases in low temperature (17 °C) and low N:P, while MetH did not. Pseudocobalamin and MetH quotas were influenced by culture methods and growth phase. Despite the variability present in cultures, we found a comparably consistent quota of 300 ± 100 pseudocobalamin molecules per cyanobacterial cell in the Northwest Atlantic Ocean, suggesting that cyanobacterial cell counts may be sufficient to estimate pseudocobalamin inventories in this region. This work offers insights into cellular pseudocobalamin metabolism, and the environmental and physiological conditions that may influence it, and provides environmental measurements to further our understanding of when and how pseudocobalamin can influence marine microbial communities.

Introduction

Cobalamin, B_{12}, is an essential organic micronutrient that impacts microbial community composition and productivity in various parts of the ocean (Koch *et al.*, 2011; Bertrand *et al.*, 2015). Most cyanobacteria produce and use a unique cobalamin analog called pseudocobalamin, psB_{12}, that is poorly bioavailable to most eukaryotic phytoplankton because of its structural differences at the α-ligand (adenine = psB_{12}, DMB = B_{12}) (Helliwell *et al.*, 2016; Heal *et al.*, 2017). Similar to B_{12}, psB_{12} occurs in different forms (Me-, Ado- and OH-psB_{12}), which have unique functions in the cell (Fig 1A). psB_{12} has the potential to impact B_{12} availability through the activities of remodelers, such as select heterotrophic bacteria and eukaryotic algae, which are able to exchange the adenine α-ligand for DMB (Helliwell *et al.*, 2016; Ma *et al.*, 2020). The prevalence of and controls on psB_{12} production and use remains unclear, in part due to the limited measurements of psB_{12} in cultures and in the natural environment.

Figure 1: (A) Structure of pseudocobalamin, psB_{12}, with adenine at the alpha (α) ligand and Me-, OH- or Ado- at the beta (β) ligand. C:N ratios of the molecules are as follows (OH-psB_{12} = 3.6:1, Me-psB_{12} = 3.6:1, Ado-psB_{12} = 3.2:1). (B) Simplified diagram of cobalamin biosynthesis pathway along with the percent of marine *Synechococcus* genomes (n = 15) with the putative gene from Helliwell et al, 2016. (C) Known (MetH) and expected (NrdJ) proteins in *Synechococcus* that use psB_{12} as a co-factor. Genes/proteins that are bolded are investigated in this study as proxy.

Accurately quantifying metabolites can help characterize the composition, activity, and controlling processes in marine microbial communities. However, such measurements remain difficult in part because of the complex and unknown chemical backgrounds they are embedded in (Kido Soule *et al.*, 2015; Moran *et al.*, 2022). Multiple studies have quantified particulate B_{12} (size fraction >0.2 μm), while only one study to date has quantified particulate psB_{12} in the ocean (Heal *et al.*, 2017). Concentrations of particulate psB_{12} ranged from undetectable to 0.15 pM. These values correlated with cyanobacterial carbon and co-occurred with B_{12} (Heal *et al.*, 2017). Quantifying both psB_{12} and B_{12} compounds would provide a more complete picture of the

processes governing cobalamin availability, especially in regions vulnerable to cobalamin co-
limitation and where cyanobacteria are abundant.

Insight into psB_{12} cell biology is restricted because psB_{12} quotas have only been
quantified to date in four cyanobacterial strains (Heal *et al.*, 2017) and the influence specific
biotic or abiotic factors have on psB_{12} production and use have yet to be explored, either in
culture or in the field. The psB_{12} biosynthesis pathway includes up to 30 enzymes (Cob/Cbi
proteins) and is similar to the B_{12} pathway with the exception of the lower ligand synthesis and
activation (DMB for B_{12}, adenine for psB_{12}) (Helliwell et al, 2016; Roth et al, 2003). Corrin ring
biosynthesis is present in the majority of available marine *Synechococcus* genomes (n = 15)
(Helliwell et al, 2016) (Fig 1B) but the specific bottleneck or rate limiting step of this pathway
remains unclear (Rodionov *et al.*, 2003; Lu *et al.*, 2020). One key psB_{12}-dependent enzyme in
cyanobacteria is methionine synthase (MetH or MS), which requires *methyl*-psB_{12} (Me-psB_{12}) as
a cofactor to generate methionine from homocysteine and 5-methyltrahydrofolate (Tanioka *et*
al., 2009, 2010; Heal *et al.*, 2017) (Fig 1C). An alternative non-B_{12} requiring methionine
synthase, MetE, appears to be absent in marine *Synechococcus* genomes (Rodinov et al,
2003). Some cyanobacteria also encode NrdJ, a class II ribonucleotide reductase that is
expected to use Ado-psB_{12} to generate dNTPs from their ribonucleotide metabolites through a
deoxygenation step (Gleason and Olszewski, 2002; Lundin *et al.*, 2010), though the use of
psB_{12} by this enzyme has yet to be specifically examined (Fig 1C). Quantifying proteins involved
in both psB_{12} production and use has the potential to provide insights into psB_{12} cell biology, but
has thus far not been investigated.

Synechococcus is one of the most abundant cyanobacterial genera in the ocean
(Flombaum *et al.*, 2013) and an important member of the microbial community in the Northwest
Atlantic Ocean, particularly in coastal euphotic zone waters during the fall (Li and Wood, 1988;
Zorz et al, 2019; Soto et al, 2023), As such, it is likely to be a major source of psB_{12} in the
region. *Synechococcus* is comprised of many strains that can be grouped into distinct clades
using phylogenetic clustering (Ahlgren and Rocap, 2012). These clades appear to thrive under
distinct levels of light, temperature, macronutrients, and metals (Sohm et al, 2016; Ahlgren and
Rocap, 2012). Environmental trends in this region indicate that the plankton community is
subject to large changes in temperature (from 2 °C in spring up to 25 °C in fall) (Zorz et al,
2019) and nitrogen availability (Loder *et al.*, 2013). Dominant *Synechococcus* ASVs in the
Northwest Atlantic Ocean have been placed in clade III, I and IV (Robicheau *et al.*, 2022).
Investigating the influence that temperature and nutrient availability have on psB_{12} in a
Synechococcus isolate of a regionally important clade could provide insight into how the psB_{12}
inventory may fluctuate in this region.

Here, we leverage targeted metabolomics and proteomics to investigate psB_{12}
production and use in *Synechococcus* in communities from the Northwest Atlantic Ocean
(NWA). To begin, we cultured *Synechococcus* sp. WH8102, a well characterized strain
belonging to a dominant clade in the region (clade III), under a range of conditions (low N:P, low
temperature, diel cycle, growth phase) and distinct culture methods (batch culture,
semicontinuous culture). By measuring both psB_{12} production and use in these conditions, we
obtain quantitative insights into *Synechococcus* psB_{12} cell biology while constraining psB_{12}
quotas and highlighting environmental and physiological conditions that may impact it. Then,
using samples collected from the Northwest Atlantic Ocean, we investigate in situ psB_{12} quotas
and use in environmental *Synechococcus* communities through protein and co-factor
quantification.

Material and methods

Identification of proteins and selection of peptides

We performed a literature search to identify proteins involved in psB$_{12}$ production
(Cob/Cbi proteins) and utilization (MetH and NrdJ). BLASTp was used to search for homologues
of these *Synechococcus* sp. WH8102 proteins in other available marine *Synechococcus*
genomes using the GenBank nr database. Proteins were aligned using COBALT (Papadopoulos
and Agarwala, 2007) and digested in silico with trypsin to reveal conserved tryptic peptides.
Sequences were introduced in the Unipept Tryptic Peptide Analysis search (Singh *et al.*, 2019)
to determine if the peptide is present in a significant number of the *Synechococcus* strains of
interest. Peptides for CobO, cob(I)alamin adenosyl transferase, and MetH, methionine
synthase, were selected for analyses as they met the peptide selection guidelines for SRM
targeted assays described in Hoofnagle et al. (2016) (Hoofnagle *et al.*, 2016) and were largely
conserved across *Synechococcus* clades. Peptides are reported in Table S1. We found no
suitable peptide with adequate coverage across *Synechococcus* clades for NrdJ.

Culture experiments to measure *Synechococcus* peptides and pseudocobalamin quotas

Synechococcus sp. WH8102 (CCMP2370) obtained from the National Center for Marine
Algae and Microbiota (NCMA, Bigelow Laboratory, USA) was grown axenically in plastic, vented
250 mL flasks (10861-576, VWR) with modified Synthetic Ocean Water media (Dupont *et al.*,
2008) void of any added cobalamin, under ~20 μmol photons m^{-2} s^{-1} supplied through white
LED lights (LEDMO-EZ550). Irradiance was measured by a PAR irradiance sensor (QSL2101,
Biospherical Instruments) and set at a light:dark cycle of 12:12 h (Table 1). Throughout
culturing, we monitored growth via relative fluorescence units (RFU) measured in a 10-AU
Fluorometer (Turner Designs, San Jose, CA), and cell counts using a BD Accuri™ C6 flow
cytometer as described below. Cultures were also tested for axenicity once a week using SYBR
Green II dye (Invitrogen) followed by flow cytometry screening. Specific growth rates (d^{-1}) were
calculated using linear regression of ln cell counts during exponential growth.

We maintained semicontinuous cultures of *Synechococcus* sp. WH8102 in exponential
phase by replacing a volume of culture with fresh media to dilute the cell concentration to a fixed
value every other day (Table 1). Semicontinuous cultures were grown in three conditions,
control (24±1 ºC; 26 N:P), low temperature (17±1 ºC; 26 N:P) and low N:P ratio (24±1 ºC;10
N:P). Cultures (n=3) were maintained for ~7 generations in exponential phase under the
conditions described before gentle harvesting using a bottle top vacuum filtration system. We
harvested protein samples (50 mL) on 0.2 μm polycarbonate filters, metabolite samples (50 mL)
on 0.2 μm Nylon filters, and particulate organic carbon (POC) samples (20 mL) on GF/F glass
microfiber filters.

Table 1: Experimental treatments (n=3) performed on *Synechococcus* sp. WH8102 in specific
culture methods.

Treatment	Culture methods	Harvesting time
Control (24±1 ºC; 26 N:P)	Semicontinuous	after ~7 generations (Exponential phase)
Low temperature (17±1 ºC)		
Low N:P ratio (10 N:P)		
Diel cycle (harvest at 9:30, 18:00 and 22:00)	Batch	Exponential phase
		Stationary phase

146
For strain *Synechococcus* sp. WH8102 grown in batch culture, we harvested samples at
both exponential and stationary phase (Fig S1). For early exponential phase, we harvested 35

mL for protein samples, 17 mL for metabolite samples, and 8 mL for POC samples at the three
points in the diel cycle. In stationary phase, we harvested 20 mL for protein samples, 10 mL for
metabolite samples, and 5 mL for POC samples at the three points in the diel cycle. The three
points of the diel cycle evaluated were: 1.5 hours after the lights went on (9:30), 1 hour before
the lights went off (18:00), and 3 hours after the lights went off (22:00).

Synechococcus sp. WH8102 culture cell counts
Culture cell density was measured on an Accuri C6 flow cytometer (BD Biosciences)
equipped with a Red (640 nm) and Blue (488 nm) laser, after labeling with1:1000 SYBR Green
II dye (Invitrogen). Single cell events were recorded on forward side scatter and side scatter
plots and distinguished using optical filters of phycoerythrin (488 nm laser, filter = 572/28 nm)
and chlorophyll *a* (488 nm laser, filter = 675/30 nm). Accuri run limits were set to a maximum of
500,000 events or 50 µL, with a flow rate of 35 µL/min and a core size of 16 µm. The gating
strategy for *Synechococcus* populations was developed based on the Acccuri application note
provided by BD Biosciences.

POC measurements
Samples were acidified for 6 hours in a glass desiccator with an open bottle of
concentrated hydrochloric acid (37% HCl, 250 mL) and later dried overnight at 45 °C. Filters
were packed in tin capsules and analyzed for particulate carbon on an elemental analyzer
(Elementar Vario microcube) coupled to an IRMS (Isoprime 100). The samples were flash
combusted at 1150 °C to convert particulate carbon into CO_2 gas. These gaseous components
were then analyzed by the Isoprime 100. The values were blank-corrected and divided by cell
number to give per cell quota.

Environmental sample collection
We collected environmental samples in Fall 2016 onboard an AZMP (Atlantic Zone
Monitoring Program) cruise aboard CCGS Hudson from September 15th, 2016, to October 6th,
2016 (HUDSON 2016027). Samples were taken at 4 stations on the Halifax line (HL 01, HL 02,
HL 05.5, HL 06). At each of the stations, we collected 1 L of water at each depth for particulate
metabolite analysis. The sample was gently filtered through a 0.2 µm Nylon filter in the dark to
reduce photodegradation. For protein samples, 10 L of water was pre-filtered (330 µm) from
Niskin bottles then filtered gentled through 3 and 0.2 µm polycarbonate filters via peristaltic
pumping. Parallel 1.8 mL seawater samples for analytical flow cytometry were fixed with 1%
paraformaldehyde. Flow cytometry samples and filters were then frozen immediately at -80°C
until analysis.

Environmental cyanobacterial cell counts
Preserved samples were thawed at room temperature, and prefiltered using 35 µm cell
strainers into a 5 mL fluorescence-activated Cell Sorting (FACS) tube. The samples were then
collected and analyzed on a Novocyte 3000 with NovoExpress software (Agilent, USA) for
natural fluorescence cell counts. The Novocyte 3000 is equipped with three lasers [Red (640
nm), Blue (488 nm), and Violet (405 nm)], and the optical filters of interest are chlorophyll-a (488
nm laser, filter = 675/30 nm),divinyl chlorophyll a (405 nm laser, filter = 675/30 nm), phycocyanin
(640 nm, filter =675/30 nm) and phycoerythrin (488 nm laser, filter = 572/28 nm). The samples
were collected with a flow rate of 120 µL/min and a threshold of chlorophyll > 450 nm. The
gating strategy was developed based on the Novocyte application note provided by Agilent and
the *Prochlorococcus* gate was constructed with the aid of pure *Prochlorococcus* culture, an
example of environmental flow cytogram is provided (Fig S2).

Particulate metabolites extraction
Metabolites were extracted from both culture and environmental samples following the
methodology described by Heal et al. (2017) with minor modifications. Briefly, 0.2 mL of each

100 µm and 400 µm silica beads were added to the bead beater tubes containing sample filters.
1 mL of ice-cold solvent mixture (40:20:20 acetonitrile: methanol: MQ water) was added and the
mixture was agitated using a bead beater (MP Biomedicals) in 3 x 40 second pulses at 1800
rotations per minute (RPM) over a 20-minute period. Heavy-CN-B_{12} (Cambridge Isotopes Labs,
CLM-9770-E) was used as an internal standard and was added to a final concentration of 3 nM
in all metabolite samples prior to extraction. Solvent was removed under vacuum (Eppendorf,
Mississauga, ON) at room temperature. Culture samples were re-suspended in buffer A (20 mM
ammonium formate, 0.1% formic acid, 2% acetonitrile) to generate equivalent of extracts from
12,100 cells/µL. Environmental samples were resuspended with 100 µL of buffer A. All samples
were then vortexed and centrifuged at 10,000 x g for 3 min at 4 °C and then diluted 2-fold with
buffer in conical polypropylene HPLC vials (Phenomenex, Torrance, CA) prior to mass
spectrometry analysis.

Metabolite quantification
HPLC-MS was used to quantify cobalamins using a Dionex Ultimate-3000 LC system
coupled to the electrospray ionization source of a TSQ Quantiva triple-stage quadrupole mass
spectrometer in SRM mode. Details of mass spectrometry conditions are reported in
supplement methods and the transition list is reported in Table S2.
For metabolite samples, quality control (QC) samples were prepared by mixing equal
volumes of each sample within its respective group (1 QC for culture samples and 1 QC for
environmental samples). Calibration curves with authentic cobalamin standards were prepared
using the QC sample as a matrix and triplicate injections were performed for 0, 0.1, 5, 10, 50,
and 100 fmol of each authentic cobalamin standard on analytical column with linearity up to
1000 fmol. All cobalamins were normalized by heavy-CN-B_{12} internal standard to account for
variability introduced by matrix effects, during extraction and instrument analysis. Limits of
quantitation and limits of detection were calculated as 10x and 3x the variation in the QC
sample without spike, respectively and recorded in Table S3.
Given that commercial psB_{12} standards are not available, the following assumptions
were made for psB_{12} quantification using associated cobalamin analog standards: (1) B_{12} and
psB_{12} ionize with the same efficiency, (2) B_{12} and psB_{12} fragments are generated in a similar
fashion, (3) B_{12} and psB_{12}'s different elution times do not greatly affect the response.

Protein extraction and digestion
Protein sample filters were submerged in 750 µL of SDS extraction buffer (2% SDS, 0.1
M Tris/HCl pH 7.5, 5% glycerol, 5 mM EDTA) and incubated for 10 min on ice before being
heated for 15 min at 95 °C in a ThermoMixer C (Eppendorf, Mississauga, ON) at 350 RPM.
Samples were then sonicated on ice for 1 min with a Q125 Sonicator (Qsonica Sonicators,
Newton, CT) at 50% amplitude and 125 W (pulse 15 s ON, 15 s OFF) then kept at room
temperature for 30 min and vortexed every 10 min. The filter was then removed, and samples
were centrifuged at 15,000 x g for 30 min at room temperature. 4x volume of ice-cold acetone
was added to the supernatant for overnight precipitation of proteins at -20 °C before washing
with 3 x 400 µL of ice-cold acetone and 1 x 400 µL with ice-cold methanol. Following each
wash, samples were centrifuged at 15,000 x g for 30 min at room temperature and the
supernatant was removed. Pellet was dried down under vacuum (Eppendorf, Mississauga, ON)
for 15 min at room temperature. Extracted protein was resuspended in 20 µL of 8 M urea for 10
min at room temperature then80 µL of freshly made 50 mM ammonium bicarbonate was
gradually added to get a final solution of 1.6 M urea and 40 mM ammonium bicarbonate.
Aliquots of 50 µg of protein were removed from each sample and diluted to 100 µL with
50 mM ammonium bicarbonate for protein digestion. Proteins were reduced with 5mM
dithiothreitol (DTT) at 56 °C for 30 min in a ThermoMixer, then alkylated with 15 mM
iodoacetamide at room temperature in the dark. Residual iodoacetamide was quenched with a

second addition of DTT to give a final concentration of 10 mM DTT. Protein was digested with 1
µg trypsin (Thermo Scientific, Waltham, MA) at 37 ⁰C overnight. 0.5 µL of formic acid was added
to halt the digestion. A modified protein Micro BCA kit (Thermo Scientific - p/n 23235)was
conducted to confirm protein concentration of each sample before injection using a trypsin
digested BSA (Bovine Serum Albumin) standard to make the calibration curve.

Peptide quantification
Targeted mass spectrometry was performed using a Dionex Ultimate 3000 UPLC system
interfaced to a TSQ Quantiva triple-stage quadrupole mass spectrometer (MS) (Thermo
Scientific, Waltham, MA), fitted with a heated, low flow capillary ESI probe (HESI-II). Isotopically
labeled, heavy internal standard versions of each peptide were synthesized by Thermo
Scientific™ at >95% purity. Heavy peptide standard was added to samples to yield 20 fmol on
column for every 1 ug total protein injected. Details of mass spectrometer conditions are
reported in supplement methods. Each sample was analyzed via triplicate injections using the
transition list of peptides that is found in Table S1.

Statistical analysis
Statistical analyses were performed using R (RStudio Team, 2020). A one-way ANOVA
followed by a Post-hoc Tuckey's Test was used to determine the differences in *Synechococcus*
culture experiments. A two-way ANOVA followed by a Post-hoc Tuckey's Test was used to
determine the effects of growth phase and diel cycle on the same variables. Differences were
considered significant when a $p\text{-value} < 0.05$ was observed.
266

Results

Factors that influence psB₁₂ production and use in *Synechococcus* sp. WH8102

Figure 2: Changes in (A) growth rate (per day), (B,C) mol C per cell, (D, E) methyl- , (F, G) hydroxy- pseudocobalamin and (H,I) MetH and (J,K) CobO (nmol per mol carbon) in *Synechococcus* sp. WH8102 under different experimental treatments grown in batch (B, D, F, H, J) and in semi-continuous cultures (C, E, G, I, K). Mean values are represented by solid circles, individual biological replicate (n=3) values by small open circles, and the standard deviation by vertical lines. Different letters over data points indicate statistically significant differences (p-value < 0.05) between pairs of means based on the post-hoc Tukey's Test. Shared letters (i.e. a, b, c) indicate no significant difference, if no letters are displayed then the differences between treatments were insignificant. *Synechococcus* sp. WH8102 culture data provided in Dataset 1.

We performed culture experiments with *Synechococcus* sp. WH8102 (clade III) in triplicate (n=3) to investigate the influence different conditions have on psB₁₂ production and use. Growth rates (per day) were similar to those reported in literature and were consistent across batch grown cultures and semi-continuously grown cultures in control and low nitrogen treatments but significantly lower in low temperature (17°C) (Fig 2A) (Mackey *et al.*, 2013). Mol carbon per *Synechococcus* cell increased during the day and in stationary phase, and significantly increased in low temperatures (Fig 2 B, C). Microgram (µg) protein per *Synechococcus* cell was invariant across the time of day and growth phase, but significantly increased in low temperature compared to other treatments (Fig S3B, C).

Cellular Me-psB₁₂ stoichiometry and quota (intracellular nutrient pool) ranged from 20 to 1500 nmol psB₁₂ per mol C (Fig 2D, E) and 340 to 16,000 Me-psB₁₂ molecules per cell respectively (Dataset 1). OH-psB₁₂ ranged from 7 to <1 nmol psB₁₂ per mol C in all samples (Fig 2F, G), 4 to 67 molecules per cell, and accounted for <1% of total psB₁₂ inventory per cell (Fig 2 D-G). Despite our low limit of detection (LOD) (22 molecules per cell, Table S3), we did not detect Ado-psB₁₂. Me-psB₁₂ quota did not differ across diel cycle but was significantly higher in stationary phase (Fig 2D). Both Me-psB₁₂ and OH-psB₁₂ quotas increased >350% in cultures grown in batch compared to the control cultures grown semi-continuously although there was no difference in growth rate (Fig 2D-G). For semi-continuously grown cultures, Me-psB₁₂ cellular quota significantly decreases at 17 °C compared to 24 °C, and when grown at 10 N:P versus the 26 N:P control (Fig 2E). In contrast, we found that OH-psB₁₂ quota did not increase in

stationary phase and was only significantly lower when grown semi-continuously at 17 °C (Fig
2F, G).

MetH peptide concentration ranged from 5 to 33 nmol per mol C in conditions tested (Fig
2H, I), representing 0.07 to 0.18% of total cellular protein (supplemental methods), and 40 to
370 MetH copies per cell (Dataset 1). MetH peptide concentration increased in the morning
(9:30) at exponential phase when compared samples taken in the afternoon (18:00) and
evening (22:00) (Fig 2H). MetH abundance (fmol per nmol carbon) did not vary in low
temperature and low nitrogen (Fig 2I). CobO concentration ranged from 3 to 24 nmol per mol C
in conditions tested (Fig 2J, KI), representing a 0.006 to 0.02% of total cellular protein
(supplemental methods), and 20 to 240 CobO copies per cell (Dataset 1). CobO concentration
was higher in exponential phase than stationary phase and were significantly lower in cultures
grown in low temperature (Fig 2K). CobO fmol per µg protein was invariant in the conditions
tested (Fig S3J, K).

psB$_{12}$ production and use by cyanobacteria in the Northwest Atlantic Ocean

Figure 3: (A) Location of sampling on the Scotian Shelf and (B) depth profiles of particulate
methyl-psB$_{12}$ (pM, triangle), cyanobacterial abundance (10^6 cells per L, circles), temperature
(°C, yellow) and chlorophyll *a* fluorescence (green) depth profiles in the Fall, 2016. For
pseudocobalamin, error bar represents technical replicates of mass spectrometry
measurements. Circles are shaded based on the calculated % population of *Synechococcus*
cells amongst all cyanobacteria (*Synechococcus* plus *Prochlorococcus*) determined through
flow cytometry. Environmental data provided in Dataset 2.

To investigate psB$_{12}$ production and use in the Northwest Atlantic Ocean, we measured
cyanobacterial cells and particulate psB$_{12}$ concentrations at four stations on the Scotian Shelf
(HL 01, 02) and Slope (HL 05.5, 06) (Fig 3A). Peak chlorophyll *a* fluorescence (green) was
between 10-30 m for HL 01 and HL 02 stations but were below 40 m for HL 05.5 and HL 06.
Temperature decreased with depth at all stations but increased in surface waters from coastal to

shelf break stations. Additional environmental data is provided in Dataset 2.

Me-psB$_{12}$ was the only quantifiable particulate psB$_{12}$ form, despite our method's low
limits of detection for both OH- and Ado-psB$_{12}$ (Table S3). Me-psB$_{12}$ measurements were up to
0.08 pM and were undetectable below 50 m depth (Fig 3B). We found a higher particulate psB$_{12}$
concentration at shallow depths and coastal waters, which generally mirrored cyanobacterial
distribution. We also measured particulate B$_{12}$ which ranged from undetectable to 0.4 pM total,
with Me-B$_{12}$, Ado-B$_{12}$, and OH-B$_{12}$ dominant in that order (Dataset 2).

Synechococcus cells per L were highest in the surface at HL 01 and HL 02.
Synechococcus dominated cyanobacterial communities in most samples except for HL 06 at 20
m and 50 m (32% and 65% respectively) where *Prochlorococcus* also contributed (calculated
from cyanobacteria cell counts, Dataset 2). *Synechococcus* comprised >65% of the
cyanobacterial community in 1 m samples from all stations and these samples were therefore
chosen to be screened for the *Synechococcus*-specific MetH peptide.

Figure 4: Assessing MetH peptide coverage in Scotian Shelf *Synechococcus* communities. (A)
Recruitment of cobalamin-related reads per nucleic acid by two dominant *Synechococcus*
metagenome assembled bins, shown as a proportion of all cobalamin-associated reads from the
entire 0.2 um prokaryotic community at HL 02, HL 06 in Northwest Atlantic Ocean in Fall, 2016
(see Soto et al, 2023). (B) Alignment of MetH protein sequences from the two dominant
assembled *Synechococcus* genome bins from Halifax Line in Fall 2016 (black) and
representative *Synechococcus* genomes from Clade I (orange), Clade III (blue), Clade IV
(green). Peptide used to quantify *Synechococcus* MetH in this study is highlighted in grey.

We examined the presence of the MetH peptide in environmental *Synechococcus*
communities, leveraging a previous metagenomics study which identified potential prokaryotic
cobalamin producers on the Scotian Shelf (Soto et al., 2023). Two *Synechococcus* genome bins
(BB16_Bin_49-*Synechoccus* and AZOF_MAG_68-*Synechoccus*) are present at surface depths
at HL 02 and HL 06 in Fall 2016 (Soto *et al.*, 2023) (Fig 4A). The peptide we used to quantify
MetH in *Synechococcus* in this study, highlighted in gray, is present in these bins (Fig 4A) and
within available marine *Synechococcus* genomes from clade III, IV and I which are the dominant
Synechococcus clades present in this region (Robicheau et al, 2022) (Fig 4B). The high
contribution of cobalamin-associated reads from these two dominant *Synechococcus* bins at HL
02 and HL 06 in Fall 2016, and the presence of the MetH peptide in these bins, suggests that
the MetH peptide we used in this study is likely to capture the majority of *Synechococcus* MetH
present in these locations.

Figure 5: Pseudocobalamin production and use in the Northwest Atlantic Ocean. Relationship
between (A) particulate psB$_{12}$ (pM) and cyanobacterial cell counts (10^6 L^{-1}) in environmental
samples. (B) Correlation between *Synechococcus* cell (10^6 L^{-1}) and *Synechococcus* MetH
abundance (fmol L^{-1}) in the surface ocean, (C) Protein (MetH) to cofactor (Me-psB$_{12}$) ratio per
Synechococcus cell in semicontinuous culture studies (grey circle, triangle, diamond) and
environmental samples (blue squares). Horizontal and vertical error bars represent technical
replicates measured using LC-MS/MS.

Assuming that all Me-psB$_{12}$ originates from cyanobacteria, cyanobacteria in our samples had
a Me-psB$_{12}$ quota of 300 ± 100 molecules Me-psB$_{12}$ per cell, with the exception of HL 05.5 at 1
m where there was ~1150 Me-psB$_{12}$ molecules per cyanobacterial cell (Fig 5A). Cyanobacterial
cells count was used in place of *Synechococcus* cell counts as *Prochlorococcus* contributed to
the cyanobacterial community in some of these samples (i.e. HL 06, Fig 3B). All *in situ* quotas
fall into a range set by culture studies (Fig 2D, E). We also found an average of 460 ± 75
molecules of MetH per *Synechococcus* cell in surface samples measured (Fig 5B).

We then investigated the MetH:Me-psB$_{12}$ stoichiometry in our culture and environmental
community samples. Stoichiometry allows us to determine the proportion of Me-psB$_{12}$ quota that
is theoretically being used by MetH and provides insights into the *in situ* psB$_{12}$ related status of
the community and how it compared to cultures. Crystal structures of MetH have revealed that
there is a 1:1 MetH to Me-B$_{12}$ stoichiometry, and that MetH exclusively uses one Me-B$_{12}$
molecule as a co-factor (Matthews *et al.*, 2008; Koutmos *et al.*, 2009). Although the structure
has not been explicitly determined for cyanobacterial MetH, a study has demonstrated that
cobalamin binding pocket of cyanobacterial psB$_{12}$ accommodates the adenine of psB$_{12}$ in place
of the DMB (Heal *et al.*, 2017). From this, we assume that MetH and psMe-B$_{12}$ have a 1:1
protein to cofactor.

Notably, the sample from HL 02 was the only one to have a ratio lower than 1:1 (HL 02 = 0.4
Me-psB$_{12}$: 1 MetH). This suggests that there was insufficient Me-psB$_{12}$ to occupy the MetH
enzymes present. In contrast, HL 05.5 had 2.8 Me-psB$_{12}$:1 MetH. Both *Synechococcus* sp.
WH8102 samples grown semi-continuously at 17 °C (low temp treatment) and environmental
samples from HL 06 have close to the theoretical ratio of 1:1, indicating tight coupling of Me-
psB$_{12}$ production and use in these samples. Cultures grown in batch had significantly more Me-
psB$_{12}$ per cell than environmental samples and where therefore omitted from this graph but are
displayed in Fig S4.

Discussion

Is protein abundance a suitable proxy for psB$_{12}$ quotas?

There is increasing interest in leveraging transcriptomic and proteomic data to infer rates and processes in microbial communities (Hawco *et al.*, 2020; Saito *et al.*, 2020; Roberts *et al.*, 2024). In theory, measuring coenzyme use (MetH) or production (CobO) could help approximate cellular concentrations or production rates of Me-psB$_{12}$. In practice, this requires a thorough understanding of psB$_{12}$ production, use, and changes in cofactor to protein ratios.

Here we show that CobO peptide per mol carbon did not always mirror psB$_{12}$ quotas. For example, in stationary phase CobO was the lowest while Me-psB$_{12}$ was the highest (Fig 2). Interestingly, CobO abundance and psB$_{12}$ decreased in unison under low temperatures. However, CobO is just one step of approximately 30 that are in the pathway to produce corrinoids *de novo* (Warren *et al.*, 2002, Fig 1B), and is responsible for the production of Ado-psB$_{12}$ which we were unable to measure in both culture and environmental samples. Contrary to what can be observed in many heterotrophic bacteria, Cob/Cbi genes in *Synechococcus* (including the strain used in this study) are not clustered in operons and therefore may not be tightly co-regulated in *Synechococcus* (Rodionov *et al.*, 2003). The same environmental or physiological condition may thus have a different effect on different Cob/Cbi genes. Recent research determined that CobSV, CobQ and CobW are key genes that enhanced cobalamin synthesis in *Ensifer adhaerens* HY-1 when more highly expressed (Xu *et al.*, 2022). Future investigations into these genes, and others, may provide insight into the rate limiting step and regulation of psB$_{12}$ and B$_{12}$ biosynthesis.

MetH peptide abundance, perhaps a proxy for Me-psB$_{12}$ use, rarely followed the general variations we observed in Me-psB$_{12}$ in batch grown *Synechococcus* sp. WH8102 cultures. For example, in stationary phase MetH abundance was lowest while Me-psB$_{12}$ was highest (Fig 2D, H). In semi-continuously grown cultures, MetH was invariant while Me-psB$_{12}$ was significantly decreased in low N:P and temperature treatments (Fig 2E, I). However, the protein:co-factor stoichiometry for cultures grown at 17 °C were surprisingly close to 1:1, suggesting that under certain conditions inferring Me-psB$_{12}$ quota from MetH abundance may be reasonable (Fig 5C). Although neither MetH nor CobO abundance sufficiently estimated or corresponded to Me-psB$_{12}$ quota in cultured *Synechococcus* sp. WH8102, the observed protein and co-factor variability still provides insight into psB$_{12}$-related cellular biology.

Variability in psB$_{12}$ quotas and protein inventories in Synechococcus sp. WH8102 cultures

Culture experiments here were performed on *Synechococcus* sp. WH8102 (clade III), a representative strain of a dominant clade in the region (Robicheau et al, 2022) to determine which variables may affect psB$_{12}$ production and use. While not representative of all *Synechococcus* in this region, this culture work provides the first insights into the plasticity of psB$_{12}$ quota and use in *Synechococcus* which can help contextualize environmental measurements. However, there is significant within and between clade variability in even cell size and macronutrient quotas (Harcourt et al, 2024), and so within and between-clade variability in psB$_{12}$ is also expected. In general, further investigation into the relationship between different factors and psB$_{12}$ related activity in different clades is an area of interest that can be pursued leveraging the peptides described here.

The relationship between temperature and psB$_{12}$ quota is particularly important when conceptualizing the spatio-temporal patterns psB$_{12}$ inventory and dynamics. Here, we

demonstrate that *Synechococcus* sp. WH8102 grew significantly slower at 17 °C compared to
the control 24 °C (Fig 2A), a result that is consistent with previous reports on this strain (Mackey
et al., 2013; Varkey *et al.*, 2016). *Synechococcus* sp. WH8102 cultures grown at low
temperatures have a significant increase of mol carbon per cell (Fig 2B), possibly due to an
increased cell size in lower temperatures (Moran et al, 2010) or decreased growth rate (Finkel
et al, 2009), or both. Previous research has shown that C, N, and P quotas in various
Synechococcus strains increase with decreased growth rate (Harcourt et al, 2024). In contrast,
we show that mol Me- and OH-psB$_{12}$ per mol carbon (Fig 2C-K) and molecules of psB$_{12}$ per cell
(Fig 5C, Dataset 2) significantly *decreased* in cultures grown at 17 °C compared to the control
24 °C. The decrease in psB$_{12}$ per cell at lower temperatures suggests that there is 1) reduced
requirements for psB$_{12}$ or 2) a decrease in psB$_{12}$ production, or both. In the low temperature
culture, we did not observe a change in MetH but we did observe a significant decrease in
CobO at low temperature (Fig 2H, J). This suggests that MetH abundance, and possibly psB$_{12}$
demand, remains constant, and that instead perhaps enzymes involved in psB$_{12}$ production may
be more impacted by low temperature than most enzymes, creating a metabolic bottleneck in
psB$_{12}$ production. We speculate that this could be a result of high sensitivity to low temperature
in the psB$_{12}$ production pathway compared to other pathways. Or, potentially, during low
temperature growth, resources could be allocated away from psB$_{12}$ production. Given that the
thermal optima for growth is variable across *Synechococcus* strains, further investigation into
the relationship between growth rate, temperature, size, and psB$_{12}$ quota in different clades is
an area of interest.

We also demonstrate that psB$_{12}$ quota in *Synechococcus* sp. WH8102 significantly
decreased in low N:P, despite no change in growth rate or significant change in demand (MetH
abundance) (Fig 2A, C). This result is of potential importance for the Scotian Shelf region, as a
decrease in dissolved nitrogen relative to phosphorus has been observed since 2010 and is
expected to continue in the future (Lehmann *et al.*, 2023). On average, the N:P ratio in
environmental samples was (3.5 ± 4.5 N:P) (Table S2) however no correlation between N:P and
psB$_{12}$ (pM) was observed. Further investigations of the relationships between N:P ratios, psB$_{12}$
production, and remodeling activity are warranted, especially given that B_{12} and nitrogen have
been shown to co-limit productivity and influence community composition in various oceanic
regions, including the coast of the Gulf of Alaska (Koch *et al.*, 2011), Long Island Embayment
(Gobler *et al.*, 2007) and Eastern boundary of the South Atlantic gyre (Browning *et al.*, 2017)..

Me-psB$_{12}$ (nmol per mol carbon) was significantly higher in *Synechococcus* sp. WH8102
harvested at stationary phase compared to exponential phase (Fig 2B). In stationary phase,
there were 20 Me-psB$_{12}$ for every MetH, suggesting that there are significant pools of psB$_{12}$ in
cells that are not associated with MetH (Dataset 1, Fig S4). This striking result points to either
intracellular storage or alternative uses of Me-psB$_{12}$ in *Synechococcus*, neither of which have
been investigated in psB$_{12}$ producers. Although roles for psB$_{12}$ remain unknown, some
alternative roles of B_{12} in the cell are beginning to be uncovered. For example, B$_{12}$ is a sensitive
light receptor and transcription regulator, albeit the benefit of these functions in stationary phase
is not obvious (Ortiz-Guerrero *et al.*, 2011; Cheng *et al.*, 2016). Although increased psB$_{12}$ quota
in stationary phase was unexpected, this trend is similar to that of some harmful algal biotoxin
producers where biotoxins accumulate in stationary phase for which reasons remain unclear
(Tong *et al.*, 2011; Song *et al.*, 2023). Investigating the reservoirs of storage, accumulation, and
alternative roles of psB$_{12}$ could provide additional insight into the cellular processes that impact
psB$_{12}$ concentrations in cells and psB$_{12}$ inventory in the ocean.

We observed a surprising >350% increase of psB$_{12}$ quota in cultures grown in batches
compared to cultures grown semicontinuously (Fig 2B, C; Fig 6A). Although psB$_{12}$ quota at the
beginning of the culture experiments was not measured, it could be expected that batch grown

cells may have begun the experiment with a higher psB_{12} quota after accumulation in previous
stationary phases (as seen in Fig 2D). Semi-continuously grown cultures were harvested after
at least 7 generations (Fig S1) that did not reach stationary phase, and as such may have
depleted their psB_{12} reserve. Unlike semi-continuous cultures, no new media was added to
batch cultures during growth, therefore the higher psB_{12} concentrations may be due to the
accumulation of either passively or actively released psB_{12} in the media or, potentially, the
buildup of substances that may promote psB_{12} production. Unfortunately, we were unable to test
this hypothesis due to the difficulties of measuring psB_{12} in the dissolved (<0.2 µm) phase.

Interestingly, we observed the same influence of culture method when investigating B_{12}
quota a recently isolated flavobacterium, *Pibocella* sp.. *Pibocella sp.* is a facultative B_{12}
consumer, it does not make nor absolutely require B_{12}, but accumulates significantly more B_{12}
per cell when grown in batch compared to semi-continuously – even at the same dissolved B_{12}
concentration (supplemental methods, Fig S5). Moreover, an increase in pg B_{12} per µg carbon
was also observed when a haptophyte, *T. lueta,* was grown in batch cultures compared to in
chemostats when grown at the same B_{12} concentration (Nef et al, 2019). The striking influence
culture method has on both B_{12} and psB_{12} quota across different species merits further research
and begs the question - *Which culture method best represents different environmental*
conditions?

Semi-continuous cultivation: most comparable to *in-situ* quotas

Comparing micronutrients quota between lab-grown cultures and environmental
communities is rare but interest is growing. To determine which culture method of
Synechococcus sp. WH8102 best reflected psB_{12} dynamics in an environmental context, we first
compared psB_{12} quotas per *Synechococcus* cell. We found that psB_{12} per cell of batch grown
cultures here fell in the same range of *Synechococcus* sp. WH8102 (clade III) and
Synechococcus sp. WH7803 (clade V) reported in literature (Heal et al, 2017) (Fig 6A).
However, batch cultures had much greater psB_{12} per cell concentrations than those measured
semicontinuously grown cultures and in the environment (Fig 6B). We also compared the MetH
to Me-psB_{12} stoichiometry from both culture and environmental studies. We found once again
that samples from semi-continuously grown *Synechococcus* sp. WH8102 most closely
resembled the *in-situ* stoichiometry of *Synechococcus* communities in this region (Fig 5C).
Semi-continuously grown *Synechococcus* sp. WH8102 ranged from 1.2 to 8.2 Me-psB_{12} per
MetH while environmental ranged from 0.4 to 2.8 Me-psB_{12} per MetH (Fig 5C). These results
combined suggest that semi-continuously grown cultures appeared to be the most appropriate
experimental approach for obtaining environmentally relevant data regarding psB_{12} in
Synechococcus communities in our region (Fig 5C). Our results highlight that careful
consideration is required when trying to infer metabolite dynamics in the field through lab
cultures.

Figure 6: psB$_{12}$ per cell in (A) cyanobacterial cultures and (B) the environment. (A) PsB$_{12}$ per cell reported by Hawco et al, 2020* (dark gray), Heal et al, 2017 (light grey) and this study (black) in *Prochlorococcus* (square) and *Synechococcus* (circle) strains grown in batch or semicontinuously. Exp = exponential phase, stat = stationary phase. (B) psB$_{12}$ quotas from Heal et al, 2017 from the North Pacific Ocean (NPO) (light grey) and from the Northwest Atlantic Ocean (NWA) (black) included in this study.
*psB$_{12}$ was inferred from targeted proteomic measurements of MetH

Using cyanobacterial cell counts as a proxy for psB$_{12}$ inventory

In this study, we provide the first set of environmental psB$_{12}$ measurements in the Atlantic Ocean (Fig 3, 5). Here, psB$_{12}$ accounted for 9 to 23% of total particulate cobalamin-related compounds (Fig7A, Table S2) - a significant part of total cobalamin that is missed if only B$_{12}$ is measured. Missing psB$_{12}$ measurements, especially in regions where cyanobacteria are abundant, restricts our ability to evaluate the importance psB$_{12}$ and remodeling have on cobalamin production. However, measuring psB$_{12}$, like many marine metabolites, is inherently challenging due to low concentration in a complex matrix, and lack of commercially available standard (Moran *et al.*, 2022). Therefore, a proxy to estimate psB$_{12}$ inventory could be a useful tool for researchers and further our understanding of when and how psB$_{12}$ can influence marine microbial communities.

Figure 7: (A) psB$_{12}$ (dark gray) and combined (light gray) total particulate B$_{12}$ (pM) down the water column at HL 02 and HL 05.5. Environmental data presented in Table S2. (B) Estimated particulate psB$_{12}$ inventory (pM) from 1997-2013 at HL 02 (1m) calculated from S*ynechococcus* cell counts from Li, 2014.

Here we found that 91% of our samples (10/11) contained 300 ± 100 Me-psB$_{12}$ molecules per cyanobacteria cell (Fig 5A). The range of psB$_{12}$ per cyanobacteria cell quantified in study was similar to the range of environmental measurements made from samples in the North Pacific Ocean (Heal *et al.*, 2017) (Fig 6B). The consistency of psB$_{12}$ measured inspired us to consider whether cyanobacterial cell counts may be sufficient to predict psB$_{12}$ inventory. For this case study, we estimated total psB$_{12}$ inventory over years at station HL 02 based *Synechococcus* cell counts from (Li, 2014), prepared as described in (Li and Dickie, 2001) (Fig 7B). While we used a narrower quota for this case study, we suggest that a general order of 10^3

psB$_{12}$ molecules per cyanobacterial cell could be useful for estimating psB$_{12}$ inventory from
cyanobacterial cell counts.

The 10^3 psB$_{12}$ molecules per cyanobacterial cell estimate suggested here is *a* quota, not
the quota. Outliers from this range will undoubtedly appear with additional environmental
measurements. However, just like the outliers from HL 05.5, we believe that these outliers
provide valuable information that can guide future research. The value of psB$_{12}$ per
cyanobacterial cell at 1 m from HL 05.5 still fall into a reasonable range reported by this study
and literature (Fig 6A, B) (Hawco et al, 2020; Heal et al, 2017). HL 05.5 is located on the
Scotian Shelf break, an oceanographically complex region where the microbial community
present is more diverse than other stations (Zorz *et al.*, 2019). Indeed, when examining the
previously analyzed cyanobacterial (16S) mapped ASVs of the samples used in this study (Fig
S6), we see that samples from HL 05.5 had highest Shannon diversity and chao values
compared to the others sampled (Fig S7) (Robicheau *et al.*, 2022). This indicates that
community diversity may contribute to the increased Me-psB$_{12}$ cell quota seen in these samples
– emphasizing the importance of psB$_{12}$ measurements from a range of cyanobacteria and
Synechococcus clades if more precise quota predictions are of utility.

Conclusion

Although psB$_{12}$ is a potentially significant source of B$_{12}$ in the ocean, there are limited
measurements of it, and the factors that influence its production and use are unknown. Here, we
reveal that in culture conditions, psB$_{12}$ quota decreases in low temperature and low N:P and that
both growth phase and culturing methods can impact psB$_{12}$ quota in *Synechococcus* sp.
WH8102. In samples from the Northwest Atlantic Ocean, we found that psB$_{12}$ quota per
cyanobacteria cell was surprisingly consistent. This provides researchers with a proxy to
systematically incorporate cyanobacteria and remodellers into conceptual and numerical models
of environmental B$_{12}$ dynamics by estimating psB$_{12}$ inventory. These quantitative values are a
step towards understanding of the role psB$_{12}$ has in marine microbial communities.

Acknowledgements

Thanks to Dr. Jenni Tolman for the advice on processing flow cytometry samples and to Dr.
William Li for the use of his long-term record of *Synechococcus* cell counts and his seminal
work on picophytoplankton in this region. We are grateful to the Bedford Institute of
Oceanography, Fisheries and Oceans Canada Atlantic Zone Monitoring Program (AZMP) for
access to the sea for the collection of samples analyzed in this study. This study was funded by
NSERC Discovery Grant RGPIN2015-05009 to EMB, Simons Foundation Grant 504183 to
EMB, Simons Foundation CBIOMES Award ID 1001702 to EMB, Canada Research Chair
Support to EMB, Ocean Frontier Institute support to EMB and JLR, and NSERC CGS-D to CB.

References

Bertrand, E.M., McCrow, J.P., Moustafa, A., Zheng, H., McQuaid, J.B., Delmont, T.O., et al. (2015)
Phytoplankton–bacterial interactions mediate micronutrient colimitation at the coastal Antarctic
sea ice edge. *Proc Natl Acad Sci USA* **112**: 9938–9943.
Browning, T.J., Achterberg, E.P., Rapp, I., Engel, A., Bertrand, E.M., Tagliabue, A., and Moore, C.M. (2017)
Nutrient co-limitation at the boundary of an oceanic gyre. *Nature* **551**: 242–246.
Cheng, Z., Yamamoto, H., and Bauer, C.E. (2016) Cobalamin's (Vitamin B12) Surprising Function as a
Photoreceptor. *Trends Biochem Sci* **41**: 647–650.
Dupont, C.L., Barbeau, K., and Palenik, B. (2008) Ni Uptake and Limitation in Marine *Synechococcus*
Strains. *Appl Environ Microbiol* **74**: 23–31.

Flombaum, P., Gallegos, J.L., Gordillo, R.A., Rincon, J., Zabala, L.L., Jiao, N., et al. (2013) Present and future global distributions of the marine Cyanobacteria Prochlorococcus and Synechococcus. *Proceedings of the National Academy of Sciences* **110**: 9824–9829.

Gleason, F.K. and Olszewski, N.E. (2002) Isolation of the Gene for the B12-Dependent Ribonucleotide Reductase from Anabaena sp. Strain PCC 7120 and Expression in Escherichia coli. *Journal of Bacteriology* **184**: 6544–6550.

Gobler, C., Norman, C., Panzeca, C., Taylor, G., and Sañudo-Wilhelmy, S. (2007) Effect of B-vitamins (B1, B12) and inorganic nutrients on algal bloom dynamics in a coastal ecosystem. *Aquat Microb Ecol* **49**: 181–194.

Gurdeep Singh, R., Tanca, A., Palomba, A., Van der Jeugt, F., Verschaffelt, P., Uzzau, S., et al. (2019) Unipept 4.0: Functional Analysis of Metaproteome Data. *J Proteome Res* **18**: 606–615.

Hawco, N.J., McIlvin, M.M., Bundy, R.M., Tagliabue, A., Goepfert, T.J., Moran, D.M., et al. (2020) Minimal cobalt metabolism in the marine cyanobacterium *Prochlorococcus*. *Proc Natl Acad Sci USA* **117**: 15740–15747.

Heal, K.R., Qin, W., Ribalet, F., Bertagnolli, A.D., Coyote-Maestas, W., Hmelo, L.R., et al. (2017) Two distinct pools of B $_{12}$ analogs reveal community interdependencies in the ocean. *Proc Natl Acad Sci USA* **114**: 364–369.

Helliwell, K.E., Lawrence, A.D., Holzer, A., Kudahl, U.J., Sasso, S., Kräutler, B., et al. (2016) Cyanobacteria and Eukaryotic Algae Use Different Chemical Variants of Vitamin B12. *Current Biology* **26**: 999–1008.

Hoofnagle, A.N., Whiteaker, J.R., Carr, S.A., Kuhn, E., Liu, T., Massoni, S.A., et al. (2016) Recommendations for the Generation, Quantification, Storage, and Handling of Peptides Used for Mass Spectrometry–Based Assays. *Clinical Chemistry* **62**: 48–69.

Kido Soule, M.C., Longnecker, K., Johnson, W.M., and Kujawinski, E.B. (2015) Environmental metabolomics: Analytical strategies. *Marine Chemistry* **177**: 374–387.

Koch, F., Marcoval, M.A., Panzeca, C., Bruland, K.W., Sañudo-Wilhelmy, S.A., and Gobler, C.J. (2011) The effect of vitamin B12 on phytoplankton growth and community structure in the Gulf of Alaska. *Limnology and Oceanography* **56**: 1023–1034.

Koutmos, M., Datta, S., Pattridge, K.A., Smith, J.L., and Matthews, R.G. (2009) Insights into the reactivation of cobalamin-dependent methionine synthase. *Proceedings of the National Academy of Sciences* **106**: 18527–18532.

Lehmann, N., Reed, D.C., Buchwald, C., Lavoie, D., Yeats, P.A., Mei, Z.-P., et al. (2023) Decadal Variability in Subsurface Nutrient Availability on the Scotian Shelf Reflects Changes in the Northwest Atlantic Ocean. *Journal of Geophysical Research: Oceans* **128**: e2023JC019928.

Li, W.K.W. (2014) The state of phytoplankton and bacterioplankton on the Scotian Shelf and Slope: Atlantic Zone Off-Shelf Monitoring Program 1994-2013, Dartmouth, Nova Scotia: Can. Tech. Rep. Hydrogr. Ocean. Sci.

Li, W.K.W. and Dickie, P.M. (2001) Monitoring phytoplankton, bacterioplankton, and virioplankton in a coastal inlet (Bedford Basin) by flow cytometry. *Cytometry* **44**: 236–246.

Li, W.K.W. and Wood, A.M. (1988) Vertical distribution of North Atlantic ultraphytoplankton: analysis by flow cytometry and epifluorescence microscopy. *Deep Sea Research Part A Oceanographic Research Papers* **35**: 1615–1638.

Loder, J., Han, G., Galbraith, P., Chasse, J., and van der Baaren, A. eds. (2013) Aspects of climate change in the Northwest Atlantic off Canada. *Canadian Technical Report of Fisheries and Aquatic Sciences 3045* 202.

Lu, X., Heal, K.R., Ingalls, A.E., Doxey, A.C., and Neufeld, J.D. (2020) Metagenomic and chemical characterization of soil cobalamin production. *ISME J* **14**: 53–66.

Lundin, D., Gribaldo, S., Torrents, E., Sjöberg, B.-M., and Poole, A.M. (2010) Ribonucleotide reduction -
horizontal transfer of a required function spans all three domains. *BMC Evol Biol* **10**: 383.

Ma, A.T., Tyrell, B., and Beld, J. (2020) Specificity of cobamide remodeling, uptake and utilization in Vibrio
cholerae. *Molecular Microbiology* **113**: 89–102.

Mackey, K.R.M., Paytan, A., Caldeira, K., Grossman, A.R., Moran, D., McIlvin, M., and Saito, M.A. (2013)
Effect of Temperature on Photosynthesis and Growth in Marine Synechococcus spp. *Plant*
Physiology **163**: 815–829.

Mackey, K.R.M., Post, A.F., McIlvin, M.R., Cutter, G.A., John, S.G., and Saito, M.A. (2015) Divergent
responses of Atlantic coastal and oceanic Synechococcus to iron limitation. *Proc Natl Acad Sci U*
S A **112**: 9944–9949.

Matthews, R.G., Koutmos, M., and Datta, S. (2008) Cobalamin-dependent and cobamide-dependent
methyltransferases. *Current Opinion in Structural Biology* **18**: 658–666.

Moran, M.A., Kujawinski, E.B., Schroer, W.F., Amin, S.A., Bates, N.R., Bertrand, E.M., et al. (2022)
Microbial metabolites in the marine carbon cycle. *Nature Microbiology* **7**: 508–523.

Ortiz-Guerrero, J.M., Polanco, M.C., Murillo, F.J., Padmanabhan, S., and Elías-Arnanz, M. (2011) Light-
dependent gene regulation by a coenzyme B$_{12}$-based photoreceptor. *Proc Natl Acad Sci USA*
108: 7565–7570.

Papadopoulos, J.S. and Agarwala, R. (2007) COBALT: constraint-based alignment tool for multiple protein
sequences. *Bioinformatics* **23**: 1073–1079.

Roberts, M.E., Bhatia, M.P., Rowland, E., White, P.L., Waterman, S., Cavaco, M.A., et al. (2024) Rubisco in
high Arctic tidewater glacier-marine systems: A new window into phytoplankton dynamics.
Limnology and Oceanography **n/a**:

Robicheau, B.M., Tolman, J., Bertrand, E.M., and LaRoche, J. (2022) Highly-resolved interannual
phytoplankton community dynamics of the coastal Northwest Atlantic. *ISME COMMUN* **2**: 1–12.

Rodionov, D.A., Vitreschak, A.G., Mironov, A.A., and Gelfand, M.S. (2003) Comparative Genomics of the
Vitamin B$_{12}$ Metabolism and Regulation in Prokaryotes. *J Biol Chem* **278**: 41148–41159.

Saito, M.A., McIlvin, M.R., Moran, D.M., Santoro, A.E., Dupont, C.L., Rafter, P.A., et al. (2020) Abundant
nitrite-oxidizing metalloenzymes in the mesopelagic zone of the tropical Pacific Ocean. *Nat*
Geosci **13**: 355–362.

Song, W., Song, X., Chi, L., Zhu, J., Cao, X., and Yu, Z. (2023) Novel insights into toxin changes associated
with the growth of *Alexandrium pacificum*: Revealing active toxin-secretion ability and toxin cell
quota variation. *Harmful Algae* **129**: 102516.

Soto, M.A., Desai, D., Bannon, C., LaRoche, J., and Bertrand, E.M. (2023) Cobalamin producers and
prokaryotic consumers in the Northwest Atlantic. *Environmental Microbiology* **n/a**:

Suffridge, C.P., Gómez-Consarnau, L., Monteverde, D.R., Cutter, L., Arístegui, J., Alvarez-Salgado, X.A., et
al. (2018) B Vitamins and Their Congeners as Potential Drivers of Microbial Community
Composition in an Oligotrophic Marine Ecosystem. *Journal of Geophysical Research:*
Biogeosciences **123**: 2890–2907.

Tanioka, Y., Miyamoto, E., Yabuta, Y., Ohnishi, K., Fujita, T., Yamaji, R., et al. (2010)
Methyladeninylcobamide functions as the cofactor of methionine synthase in a Cyanobacterium,
Spirulina platensis NIES-39. *FEBS Letters* **584**: 3223–3226.

Tanioka, Y., Yabuta, Y., Yamaji, R., Shigeoka, S., Nakano, Y., Watanabe, F., and Inui, H. (2009) Occurrence of
Pseudovitamin B12 and Its Possible Function as the Cofactor of Cobalamin-Dependent
Methionine Synthase in a Cyanobacterium Synechocystis sp. PCC6803. *J Nutr Sci Vitaminol* **55**:
518–521.

Tong, M., Kulis, D.M., Fux, E., Smith, J.L., Hess, P., Zhou, Q., and Anderson, D.M. (2011) The effects of
growth phase and light intensity on toxin production by Dinophysis acuminata from the
northeastern United States. *Harmful Algae* **10**: 254–264.

Varkey, D., Mazard, S., Ostrowski, M., Tetu, S.G., Haynes, P., and Paulsen, I.T. (2016) Effects of low
703   temperature on tropical and temperate isolates of marine Synechococcus. *ISME J* **10**: 1252–
704   1263.
705   Warren, M.J., Raux, E., Schubert, H.L., and Escalante-Semerena, J.C. (2002) The biosynthesis of
706   adenosylcobalamin (vitamin B12). *Nat Prod Rep* **19**: 390–412.
707   Xu, S., Xiao, Z., Yu, S., Zeng, W., Zhu, Y., and Zhou, J. (2022) Enhanced cobalamin biosynthesis in Ensifer
708   adhaerens by regulation of key genes with gradient promoters. *Synthetic and Systems*
709   *Biotechnology* **7**: 941–948.
710   Zorz, J., Willis, C., Comeau, A.M., Langille, M.G.I., Johnson, C.L., Li, W.K.W., and LaRoche, J. (2019) Drivers
711   of Regional Bacterial Community Structure and Diversity in the Northwest Atlantic Ocean. *Front*
712   *Microbiol* **10**: 281.
713