Intrapartum antibiotic prophylaxis selects for mutators in group B streptococci among persistently colonized patients

Macy E. Pell[1], Heather M. Blankenship[2], Jennifer A. Gaddy[3,4,5],

H. Dele Davies[6], Shannon D. Manning[1]*

[1]Michigan State University, Department of Microbiology, Genetics, and Immunology (MGI), E. Lansing, MI;

[2]Michigan Department of Health and Human Services, Bureau of Laboratories, Division of Infectious Diseases, Lansing, MI;

[3]Department of Pathology, Microbiology and Immunology, Vanderbilt University Medical Center, Nashville, TN;

[4]Department of Medicine, Vanderbilt University School of Medicine, Nashville, TN;

[5]Tennessee Valley Healthcare Systems, Department of Veterans Affairs, Nashville, TN; and

[6]University of Nebraska Medical Center, Omaha, NE.

*Materials and correspondence to mannin71@msu.edu

ABSTRACT

Through vaginal colonization, GBS causes severe pregnancy outcomes including neonatal

sepsis and meningitis. Although intrapartum antibiotic prophylaxis (IAP) has reduced early-onset

disease rates, persistent GBS colonization has been observed in patients following prophylaxis.

To determine whether IAP selects for genomic signatures that enhance GBS survival and

persistence in the vaginal tract, whole-genome sequencing was performed on 97 isolates from

58 patients before (prenatal) and after (postpartum) IAP/childbirth. Core-gene mutation analysis

identified 7,025 mutations between the paired isolates. Three postpartum isolates accounted for

98% of mutations and were classified as "mutators" because of point mutations within DNA

repair systems. *In vitro* assays revealed stronger biofilms in two mutators. These findings

suggest that antibiotics select for mutations that promote survival *in vivo*, which increases the

likelihood of transmission to neonates. They also demonstrate how mutators can provide a

reservoir of beneficial mutations that enhance fitness and genetic diversity in the GBS

population.

INTRODUCTION

As an opportunistic bacterial pathogen, Group B *Streptococcus* (GBS) colonizes ~35% of

pregnant patients[1] and colonization is the main risk factor for invasive neonatal infection[2,3].

During pregnancy, rectovaginal colonization can also lead to sepsis of the pregnant person and

other adverse outcomes including preterm birth and stillbirth[4,5]. GBS disease in neonates occurs

as early-onset disease (EOD), which presents as sepsis and/or pneumonia within the first week

of life, or late-onset disease (LOD) that presents as bacteremia and/or meningitis between 1

week and 3 months after birth[6].

In the U.S., screening for vaginal-rectal colonization is recommended at 35-37 weeks'

gestation[7]. To prevent neonatal infections, GBS-positive patients are given intrapartum antibiotic

prophylaxis (IAP) during labor, for which penicillin is the most effective[3]. While IAP has

successfully reduced the incidence of EOD, it has not impacted LOD rates and is not helpful for

preventing preterm births or stillbirths[8]. It can also negatively impact the microbiome[9,10].

Furthermore, GBS colonization is intermittent during pregnancy and colonization frequencies

vary across locations[11,12]. One study of EOD cases, for example, showed that 60% were born to

individuals who were GBS-negative at the prenatal screening[13], thereby highlighting the need

for improved treatment and prevention methods.

Classifying GBS based on the polysaccharide capsule (cps), which dictates the serotype, and

the multilocus sequence type (ST), has identified associations between molecular traits and

disease severity[14,15]. Capsule type III ST-17 strains predominate among neonates with invasive

disease[14–16] and possess unique features that help withstand oxidative stress, survive inside

macrophages, associate with host cells, and elicit proinflammatory responses relative to other

STs[17–21]. Less definitive epidemiological associations have been observed for other lineages

(e.g., STs 19 and 23) despite their high prevalence[15]. The effectiveness of IAP was also found

to vary across STs in our prior study[22], with STs 17 and 19 more commonly colonizing patients

before IAP and persisting up to 6 weeks postpartum than other STs. ST-12 strains, however,

were more frequently lost after IAP, which was also independently associated with persistent

colonization. Despite these associations, the mechanisms that promote GBS persistence in the

vaginal tract and evasion of antibiotic-mediated killing remain elusive. This is a serious concern

as persistent colonization enhances the risk of invasive disease, especially for subsequent

pregnancies and babies after IAP cessation[10,22], and may partly explain why rates of GBS LOD

have not decreased in the U.S.[8,10].

Altogether, these findings led us to hypothesize that IAP imposes a selective pressure on GBS,

such that genomic features that promote colonization are selected for to enhance fitness. We

used whole-genome sequencing (WGS) to compare GBS isolates recovered from pregnant

patients sampled before (prenatal) and after (postpartum) IAP[23]. This study has enhanced

understanding of how IAP impacts the colonizing GBS population, concurrently providing insight

for treatment and prevention improvements.

RESULTS

GBS colonization dynamics and antibiotic exposures differ across patients

Isolates (n=97) were collected from 58 pregnant patients in a prior study[23]. Forty-one (70.7%)

patients had GBS at the prenatal and postpartum samplings indicating persistent colonization

despite IAP. Two postpartum strains from two persistently colonized patients (IDs 11 and 14)

were not available for characterization. The remaining 17 patients were colonized only at the

prenatal sampling indicating loss of GBS colonization. Most (n=53; 91.4%) patients received

IAP with ampicillin (n=26), penicillin (n=17), cefazolin (n=2), or clindamycin (n=6). One patient

received both ampicillin and clindamycin, and another had penicillin and metronidazole (Flagyl).

Five (8.6%) patients did not receive IAP, including three with persistent colonization and two

with prenatal colonization only; four of these patients had neonates with signs of infection.

Moreover, eight (13.8%) patients received postpartum antibiotics, four of whom had persistent

colonization.

As determined previously[22], the 97 GBS strains represent 20 STs comprising clonal complex

(CC)-1 (n=25), CC-12 (n=15), CC-17 (n=16), CC-19 (n=17), CC-22 (n=2), CC-23 (n=20), and

CC-26 (n=1). One strain represented a singleton (ST-67). Eight distinct cps types were

represented, with cps III (n=27) predominating followed by Ia (n=21) and V (n=17); four isolates

were non-typeable (NT).

Pangenome analysis reveals a large accessory genome and clustering by CCs

Ninety-two high-quality genomes were included in this analysis; three prenatal and two

postpartum genomes from five patients were excluded due to poor assembly quality. Among the

92 genomes, 68 represented paired strains from 34 patients with persistent colonization

recovered from both the prenatal and postpartum samplings in the same patients. Additionally,

seven unpaired strains from seven persistently colonized patients collected at the prenatal (n=4)

or postpartum (n=3) sampling were included plus the 17 strains from patients who lost GBS

following IAP and were only colonized at the prenatal sampling.

The GBS pangenome consists of 5,054 unique genes comprising a small core-genome with

1,368 core and 213 soft-core genes. It has a large accessory genome, which contained 945

shell and 2,528 cloud genes accounting for 69% of the total genes. A maximum likelihood (ML)

tree based on the 1,368 core genes resolved five sequence clusters grouping together with

>95% bootstrap support (**Figure 1**). Each cluster was associated with a CC. The 17 CC-19

strains and 16 CC-17 strains grouped into two separate clusters with 100% bootstrap support,

while the CC-1 (n=23) and CC-12 (n=23) strains were part of two separate clusters with 99%

and 95% bootstrap support, respectively. Although the 19 CC-23 strains clustered together, they

comprised a mixture of outgroups and smaller clades with 2-5 genomes each. A core-gene

neighbor-net tree confirmed these groupings (**Figure S1**) and detected evidence for

recombination (pairwise homoplasy index (PHI) p-value = 0.00) across the 17,825 parsimony-

informative (PI) sites.

Among the 34 pairs of strains from patients with persistent colonization, 29 (85.3%) clustered

together with >91% bootstrap support and five did not. Both strains from two of these five

patients (IDs 3 and 8) had different STs and were on different branches of the tree, suggesting

acquisition of a new strain type following childbirth and IAP. For the remaining three patients

(IDs 22, 43, and 56), the prenatal and postpartum strains shared a ST but did not cluster

together in the ML tree. This finding suggests that the phylogenetic distance between these

paired strains is due to nucleotide-level variation within the core gene sequences.

112
113
Figure 1: Maximum likelihood tree of 92 GBS genomes based on 1,368 core genes.
Patient strain IDs are noted at the end of each branch such that persistent strains contain a ".1"
(prenatal) or ".2" (postpartum) to indicate the sampling timepoint with respect to IAP and
childbirth. The clonal complex (CC) is represented by the colored strips at the outside of the tree
and bootstrap support values >70% are noted on the tree branches. The maroon circles on the
outside of the phylogeny show the antibiotic(s) used for IAP; the paired strains from the same
patient have the same antibiotic usage profiles. IDs in red bold face font highlight the paired
prenatal and postpartum strains from five patients (IDs 3, 8, 22, 43, and 56) that do not cluster
together in the tree.

Acquisition of antibiotic resistance genes is not important for persistent colonization

To determine whether GBS residing in the vaginal tract acquired antibiotic resistance genes

(ARGs) to promote colonization, an assembly-based analysis was performed to extract these

genes from the 92 genomes. Nine distinct ARGs were identified conferring resistance to

tetracyclines (*tetL, tetM, tetO, tetW, lmrP*)*,* macrolides-lincosamides-streptogramins (MLS)

(*ermA, lmrP, mreA*), fluoroquinolones (*norB*), and cationic peptides (*mprF*). Both *lmrP* and *mprF*

were found in all strains, while *norB* and *mreA* were found in all but one (**Figure 2**).

Among the variably present ARGs, *tetM* predominated and was found in 63 (68.5%) genomes

including 47 of the 68 (69.1%) paired strains from the 34 patients. A similar proportion (n=11;

64.7%) was detected in the 17 prenatal strains that were lost following IAP. Although the

prenatal and postpartum strains from patient 22 had the same ST, *tetM* was only detected in the

prenatal strain (22.1), indicating gene loss. This result is not surprising since *tetM* is often

associated with plasmids and the two strains did not cluster together in the core-gene ML tree

(**Figure 1**). The same was true for patient 3, though this finding is less surprising in that the two

paired strains had different STs, indicating acquisition of a new strain by the postpartum visit.

Specific virulence gene profiles are not linked to persistent colonization

An assembly-based analysis was also performed to identify virulence genes in the 92 genomes;

50 distinct genes were detected and classified based on function. Genes important for

adherence and invasion of host cells (n=18, 36%), immune modulation (n=17, 34%),

metabolism (n=1, 2%), toxin production (n=13, 26%), and dissemination (n=1, 2%) were found.

Figure 2: Antibiotic resistance genes (ARGs) detected in the 92 genomes. ARGs were extracted from the 92 genome assemblies using five different ARG databases. The x-axis shows the patient strains with a ".1" (prenatal) and ".2" (postpartum) after the "Patient ID" indicating the sampling timepoint. The gene profiles for the paired isolates from the same 32 patients with isolates of the same sequence type (ST) are shown next to each other on the left side of the vertical purple line. The two sets of paired strains (IDs 3 and 8) with different STs along with seven unpaired strains are shown between the purple and red vertical lines. The ARG profiles for the 17 "lost" strains, which were recovered only at the prenatal sampling and did not persist, are to the right of the red vertical line. Gene names are on the y-axis. The color gradient shows the average percent identity of a gene across the five databases, with black representing 100% identity and mint green representing <10% identity. ARGs identified in this analysis confer resistance to tetracyclines (*tetL, tetM, tetO, tetW, lmrP*), macrolides-lincosamides-streptogramins (*ermA, lmrP, mreA*), fluoroquinolones (*norB*), and cationic peptides (*mprF*).

The greatest variation in virulence genes was observed for those encoding capsule (*cps*)

production and pili (**Figure 3**). Among the 32 persistently colonized patients with strains of the

same ST, several discrepancies in gene presence were observed. Most of these differences,

however, involved genes that fell below the identity and coverage thresholds, indicating they are

present but may be distinct. Patients 3 and 8 with prenatal and postpartum strains belonging to

different STs, also had distinct virulence gene profiles, indicating acquisition of a new strain.

To further confirm that the prenatal and postpartum strain populations had distinct virulence

gene profiles, a principal component analysis was performed. No distinct clustering patterns

were observed for the prenatal versus postpartum strains (**Figure S2A**) or after stratifying by

antibiotic, cps type, and colonization phenotype (persistent or lost) (**Figure S2B-E**). Additionally,

no difference was observed between colonization phenotype and the number of adherence

(n=1,031, Chi-square p = 0.10) or immune modulatory (n=1,231, Chi-square p=0.52) genes, as

determined by coverage and nucleotide identity thresholds. These data suggest that possession

of specific combinations of virulence genes is not linked to a strain's ability to persistently

colonize the vaginal tract after IAP.

Point mutations in persistent strains highlight microevolution after IAP

To detect nucleotide-level variation, we conducted a reads-based analysis of the 1,368 core

genes in the 64 paired prenatal and postpartum strains that shared STs and were recovered

from the 32 patients with persistent colonization. In this analysis, each prenatal genome was

used as the reference genome for its paired postpartum genome and the number and type of

mutations in the postpartum genomes was determined. Notably, 7,025 mutations were identified

in 24 (75.0%) of the 32 postpartum genomes. No mutations were detected in the remaining

eight postpartum genomes, which had identical core gene sequences as the paired prenatal

genomes from the same patients.

Figure 3: Virulence genes detected in 92 GBS genomes. The x-axis shows the patient ID with the ".1" (prenatal) and ".2" (post-partum) indicating the sampling timepoint. Gene names are displayed on the y-axis. The gene profiles for the paired isolates from 32 patients with isolates of the same sequence type (ST) are next to each other, left of the purple line. The seven unpaired strains plus the two paired strains with different STs from patients 3 and 8 are shown between the purple and red vertical lines. Profiles for the 17 "lost" strains are to the right of the red line. The color gradient displays the average percent identity of a gene, with black representing 100% identity and mint green representing <10% identity.

Among the 24 genomes with ≥1 mutation, most mutations (n=6,065; 86.3%) were in coding regions, though an additional 958 (13.6%) were in uncharacterized regions and two were in rRNAs. The type of mutations varied across genomes (**Figure S3A**). Most (n=5,982; 85.2%) were single nucleotide polymorphisms (SNPs) including synonymous (n=3,469; 58.0%), missense (n=1,771; 29.6%), or nonsense (n=26; 0.4%) mutations (**Figure 4**). Complex

mutations (n=712), defined as a combination of multiple nucleotide polymorphisms (MNPs) and

an insertion or deletion (indels), were also detected and classified as synonymous (n=311;

43.7%) and missense (n=299; 41.9%) mutations. Among the indels (172 insertions and 124

deletions), frameshift mutations (n=107; 36.1%) were most common. Similar frequencies of

conservative outcomes (n=6), defined as an indel of one or more entire codons, and disruptive

outcomes (n=7), defined as a codon change plus a codon indel, were among the deletions.

Another deletion resulted in the loss of a stop codon, while two others resulted in the loss of a

start codon. For the insertions, more were conservative (n=7) than disruptive (n=4) mutations

with two resulting in nonsense mutations and four in the loss of a stop codon. Among the 35

MNPs detected, most were missense mutations (n=25; 71.4%).

Figure 4: Core gene mutations detected in the postpartum genomes of GBS strains from
patients following IAP. Five types of mutations (y-axis) were detected in 24 of the 32
postpartum genomes when each genome was compared to its respective prenatal genome from
the same patient. No mutations were identified in eight of the 32 postpartum genomes. Types of
mutations identified include multiple nucleotide polymorphisms (MNP), complex mutations,
deletions, insertions, and single nucleotide polymorphisms (SNP). The raw mutation count is
noted in parentheses for each type of mutation, while the mutation outcomes are distinguished
by color and are shown as a proportion of each mutation type (x-axis). The categories are not
mutually exclusive as some mutations were classified as multiple types (i.e., an insertion
classified as disruptive and as a frameshift will be included in both mutation categories).

A skewed distribution of mutations reveals mutator strains in GBS

Comparing mutation frequencies across the postpartum genomes revealed a skewed

distribution with three outliers (**Figure S3B**). These outlier genomes contained 204, 262, and

6,454 mutations relative to an average of five mutations and range of 1-16 mutations for the 21

postpartum genomes with at least one mutation (**Figure S3C**). Based on the total assembly

length, we calculated the mutation rate, or the number of mutations per number of base pairs

(bp) for each postpartum genome relative to its respective prenatal genome. The rates for the

three outlier genomes are 9.37×10^{-5} (patient genome ID=36.2), 1.32×10^{-4} (ID=43.2) and 0.003

(ID=22.2). These rates were 53-, 75-, and 1,747-fold higher, respectively, than the average

mutation rate (1.77×10^{-6} mutations/bp) for the remaining 29 genomes, including the eight

genomes lacking any detectable mutations.

Because mutators arise at a low frequency in the population due to mutations within genes

encoding DNA replication and proofreading machinery, mismatch repair (MMR) pathways, and

the 8-oxo-dG (GO) system[24,25], we examined these genes in the three outlier postpartum

genomes. Genome ID 22.2, which contains the most mutations, also has the greatest number of

nonsynonymous mutations in genes encoding DNA repair systems including the MMR (n=14)

and GO (n=8) pathways, and in genes involved in DNA replication (n=18), DNA repair (n=7),

and DNA recombination (n=9). Similarly, genome ID 36.2 has a deletion in the gene encoding

DNA polymerase IV, a missense SNP in *mutY* of the GO system, and a 33 bp insertion within

an MMR locus, while genome ID 43.2 has a missense SNP in the DNA topoisomerase gene

and a 1 bp deletion downstream of the MMR loci. The increased mutation rates along with

detectable mutations in these key regions meet the criteria[24,26] for classifying these three strains

as "mutators" that arose after IAP.

Distinct classes of MMR loci exist but none are associated with hypermutability

Since mutations in MMR loci were identified in all three mutators, we examined the six MMR

pathway genes (7,467 bp region) in the 64 paired genomes from the 32 patients. These genes

encode DNA mismatch repair proteins, MutS (*mutS*) and MutL (GBS_230), a cold shock protein,

a MFS transporter/LmrP homolog (GBS_436), the Holliday junction branch migration protein

(RuvA), and DNA-3-methyladenine glycosylase I (GBS_235). A multiple sequence alignment

using MUSCLE[27] detected only 104 variable nucleotides.

Construction of a neighbor joining (NJ) phylogeny based on the MMR gene sequences (7,467

bp) revealed four clusters, which we arbitrarily named Class 1-4 (**Figure 5**); three of these

classes had subclasses grouping with >60% bootstrap support. Class 4 predominated (n=28,

43.7%) and contains three subclasses (4A-4C), followed by Class 2 (n=21, 32.8%) with

subclasses 2A-2E. Class 1 (n=12, 18.7%) has two subclasses, while Class 3 has none.

Although a neighbor-net tree based on 103 PI sites within the 7,467 bp MMR region provided no

evidence of recombination (PHI p-value=0.0899), the four distinct MMR classes are still evident

(**Figure S4**).

The MMR loci classes and subclasses are associated with specific CCs. All CC-17 strains, for

instance, possess Class 1 MMR loci, with subclasses 1A and 1B clustering separately in the

core gene phylogeny (**Figure S5**). By contrast, the Class 2 and Class 4 subclasses are split

between CCs, with subclass 4A being found exclusively within CC-23 strains and 4B and 4C

clustering separately within the CC-19 strains. The subclasses of the Class 2 MMR loci did not

group together within the phylogeny, as 2B-2E were found exclusively in the CC-1 or CC-12

strains and subclass 2A strains were unrelated and split between the two CCs.

Figure 5: Neighbor joining phylogeny based on the MMR loci (7,467 bp) extracted from 32 pairs of strains from pregnant patients collected before and after IAP/childbirth. The evolutionary distances are shown as the number of base substitutions per site. Four sequence clusters (Classes 1-4) were detected using the bootstrap test with 1,000 replicates. Subclasses were identified with three classes, which grouped with >60% bootstrap support; the number in parentheses indicates the number of the genomes with the same subclass (100% identity) among the 64 genomes. Bootstrap values are shown on the tree branches and the three mutator strains are in red, bolded text.

A comparison of the strain pairs (prenatal versus postpartum) from the same patients revealed

that the MMR classes and subclasses were identical in all but two pairs. The two postpartum

mutator strains (IDs 22.2 and 36.2) differed from the prenatal genomes (IDs 22.1 and 36.1) by

56 nucleotides (21 amino acids) and 1 nucleotide (1 amino acid), respectively. Although two of

the three mutators belong to MMR Class 4, both had distinct MMR subclasses representing 4A

(genome ID 43.2) and 4B (genome ID 36.2). The third mutator (genome ID 22.2) belonged to

Class 3 (**Figure 5**), suggesting that a single MMR allele is not associated with hypermutability.

To determine how common each MMR class is among GBS genomes and identify novel

classes, a Basic Local Alignment Search Tool (BLAST)[28] analysis was performed on GBS

genomes available in the National Center for Biotechnology Information (NCBI; access date

3/21/2023). Among ~100 GBS genomes with sequence matches, MMR subclass 2A

predominated as 43 genomes were identified with 100% nucleotide identity. Five MMR loci

subclasses (1B, 2B, 2C, 2D, and 4C), however, were unique to our strains, as no published

GBS genomes had matching sequences. The Class 3 loci, which was found in mutator strain

22.2, returned three GBS genomes with 100% identity including strains B111 (CP021772.1),

SG-M8 (CP021868.1), and GBS-M002 (CP013908.1), all of which represent invasive, human-

derived strains from Asia. Similar to strain 22.2, each was classified as

serotype VI and belong to ST-1 using PubMLST[29].

IAP elicits a selective pressure that differentially impacts lineages

To better understand which point mutations could alter protein functionality, we further classified

the subset of 5,291 characterized SNPs out of the 5,982 SNPs identified in the 1,368 core

genes among the 32 postpartum strains from patients with persistent colonization. While

specific mutations were not associated with individual lineages, CCs 1 (n=4,953), 19 (n=121),

and 23 (n=199) had the greatest number of SNPs. These numbers, however, are skewed by

320 inclusion of the three mutator strains. Excluding the mutators revealed that the CC-23 strains

321 had the most SNPs (n=21) followed by CCs 12 (n=15) and 1 (n=12), averaging 3, 5, and 1

322 SNP(s) per strain, respectively, and ranging from 0-10, 0-9, and 0-7 SNPs per strain. By

323 comparison, the postpartum non-mutator strains of CCs 17 (n=3) and 19 (n=7) acquired fewer

324 SNPs overall, averaging between 0-1 SNPs per strain with a range of 0-3. Among the few SNPs

325 identified in CC-17 strains, only one was nonsynonymous, indicating a pattern of negative or

326 purifying selection for strains in this lineage. Comparatively, the non-mutator strains of other

327 lineages had more nonsynonymous (n=38) than synonymous (n=17) SNPs, suggesting a

328 pattern of positive/diversifying selection and a trend of adaptive evolution for these lineages.

330 Based on prior medical record reviews[23], the three mutator strains had been exposed to either

331 penicillin (patient IDs 36 and 43) or ampicillin (ID 22) for IAP. Both antibiotics contributed to an

332 average of 292 and 31 SNPs in the postpartum strains from patients given ampicillin (n=17) or

333 penicillin (n=10), respectively. After excluding the mutators, however, the average decreased to

334 two SNPs/strain for both drugs. While comparisons could not be made for the other antibiotics

335 due to low numbers, the three strains from patients who did not receive IAP, had an average of

336 only one SNP/strain. Collectively, these data demonstrate that the emergence of mutators can

337 impact mutation frequencies, and further, they suggest that certain strain types are more

338 susceptible to IAP-associated evolutionary pressures.

340 **Mutators have a high frequency of missense SNPs and genes under positive selection**

341 Since mutators can serve as a reservoir of beneficial mutations for the surrounding bacterial

342 population, we sought to further examine those SNPs identified within coding sequences in the

343 three mutator strains. Among the subset of 5,291 characterized SNPs, 5,233 (98.9%) were in

344 the three mutators and mapped to 1,167 genes, yielding an average of two SNPs/gene (**Figure**

345 **6A**). Comparatively, 58 (1.1%) SNPs were detected within 56 genes in the non-mutator strains,

resulting in an average of 1 SNP/gene (**Figure 6B**). Within the mutator genomes, most genes

(n=683) acquired ≤4 SNPs relative to the respective prenatal genomes with 390, 142, 86, and

65 genes having 1, 2, 3, and 4 SNPs, respectively. Moreover, >4 SNPs were detected within

366 genes; 26 of these genes had >20 SNPs and were important for DNA replication/repair,

antimicrobial resistance, phage proteins, and virulence. Among all 366 genes, most (77.6%,

n=284) had evidence of negative selection based on the ratio of nonsynonymous (dN) to

synonymous (dS) mutations. Another 61 genes (16.7%) were under positive selection, while the

remaining 21 (5.74%) genes had evidence for neutral selection.

Restricting our analysis to genes with >4 nonsynonymous SNPs, we identified 662 SNPs in 91

genes, with most (98.3%, n=651) representing missense mutations (**Figure 6C**). Eight genes

had strong signals of positive selection (dN/dS >4) and are important for oxidative stress

(*yaaA_2,* GBS_639), competition (GBS_534), translation (*asnS*), virulence (GBS_675), and

aminoglycoside resistance (GBS_318) Two hypothetical genes (GBS_724, GBS_776) were also

of interest (Table S9) along with *aspS* (tRNA ligase), *recX* (recombination regulator), and *esaA*

(type VII secretion protein). Because multiple virulence and stress response genes were

identified, it is likely that such processes play a role in persistent colonization following IAP.

Of the 651 missense SNPs identified in the mutator strains, several were located in the same

set of genes, ranging from 4-26 missense SNPs/gene (**Figure 6C**). Virulence genes like *esaA,*

hylB (hyaluronate lyase)*,* and *pilA* (PI-2a pilus adhesin), which are important for colonization,

acquired >10 missense SNPs/gene. Furthermore, 10 nonsense mutations were acquired in

eight genes including *asp3* (accessory Sec-system protein), *fbsA* (fibrinogen-binding adhesin),

and GBS_550 (pullulanase), which also had a mutation contributing to the loss of a start codon.

Figure 6: Number and distribution of SNPs detected within genes from the 32 postpartum
genomes relative to the respective prenatal genomes. The distribution of SNP frequencies
(x-axis) by the number of genes (y-axis) in the A) mutator and B) non-mutator genomes. C) The
number of nonsynonymous SNP types (missense=blue, start lost=green, nonsense=black) is
shown by gene (x-axis) for the subset of genes with >4 nonsynonymous SNPs in the three
mutator genomes.

Postpartum strains have enhanced biofilm production relative to prenatal strains

Because biofilm formation is a common colonization strategy and nonsynonymous mutations

were found in biofilm-related genes (e.g., *pilA*, *gtfB*, GBS_067), we examined this phenotype in

the paired strains. Indeed, an increase in biofilm production was observed in over half (54.8%,

n=17) of the 31 postpartum strains relative to each paired prenatal strain (**Figure 7**); five of

these increases were significant. Among the mutators, two had increased biofilm production,

one of which (pair 22) had the largest increase observed across all pairs evaluated (fold-change

= 8.14) (**Figure S6**). Two postpartum isolates from pairs 26 and 27 exhibited a significant

decrease in biofilm formation; these isolates had acquired 11 and 2 mutations, respectively,

some of which occurred within genes encoding binding and surface proteins such as penicillin-

binding protein Pbp2B, capsule protein CpsC, and a LPXTG anchor domain surface protein.

393
Figure 7: A comparison of biofilm formation between prenatal and postpartum strains
from 31 patients with persistent colonization. Biofilm formation was assessed across the 31
paired strains by measuring the absorbance (OD_{595}) values (y-axis) and averaging the biological
replicates (n=3). Prenatal (mint green) and postpartum (dark blue) isolates are shown by patient
ID (x-axis). One-tailed, paired t-tests were calculated to identify significant differences in biofilm
formation between the paired strains (*=p <0.05, **=p<0.01). Each measurement was
normalized to a negative media control. Error bars display standard deviation.
401

DISCUSSION

Through this analysis, we have identified genomic traits and alterations that influence persistent colonization by GBS, which we had observed in 59.4% of colonized pregnant people following IAP[22], and increases the risk of consequent invasive disease. The pangenome analysis of 1,368 core genes uncovered five distinct sequence clusters, with most paired strains of the same ST clustering together. This finding along with a comprehensive evaluation of point mutations indicates that in most cases, the strains that persisted in each patient are highly similar to those recovered prior to IAP. Nonetheless, we identified evidence of microevolution in GBS as point mutations were found in 72% (n=21) of the 29 postpartum strains that clustered together with each respective prenatal strain. It is therefore likely that the accumulation of key mutations may lead to enhanced fitness in some strains, thereby promoting persistent colonization.

Indeed, we observed increased biofilm formation in over half of the postpartum strains, meaning that the strains exposed to antibiotics formed better biofilms than the respective prenatal strains. Consistent with this finding is the identification of point mutations in pili-associated genes (*pilA-C, srtC1*-4) and Sec-system related genes (GBS_067, *gtfB*, *asp3*), which were linked to biofilms in *Streptococcus pneumoniae*[30] as well as *cps* loci, which we found to be important for GBS biofilms and ascending infection[31]. Although only five of the biofilm increases were significant, many prenatal strains exhibited robust biofilm phenotypes prior to antibiotic exposure (OD_{595} >1); hence, it is possible that these strains had been persisting in each host prior to our first prenatal sampling. The variation in biofilm production observed is consistent with our study showing differences across lineages and pilus types[32]. These data suggest that biofilms may enhance persistence in some strain backgrounds, however, confirmation is required in more clinically relevant conditions.

Antibiotic resistance is another mechanism for survival in antibiotics, yet we failed to detect

newly acquired ARGs in the postpartum strains relative to the prenatal strains from each patient.

Therefore, resistance to the antibiotics used for IAP was ruled out as a mechanism of persistent

colonization. Among those postpartum genomes from individuals (88.6%) who received β-

lactam antibiotics for IAP, none had β-lactam resistance genes, which is not surprising since

GBS is universally susceptible to the β-lactams[10]. Although resistance to the MLS antibiotics is

more common in GBS[10], the 11.3% of patients given clindamycin had postpartum strains that

were neither phenotypically resistant[23] or harbored macrolide resistance genes. Acquisition of

specific virulence genes was also not associated with persistence since distinct genes were not

found in the postpartum versus prenatal strains. Nonetheless, this assembly-based analysis has

some limitations in that genes labeled as absent may have fallen below our detection thresholds

due to poor coverage or high nucleotide variation.

Most noteworthy was our observation that three pairs (IDs 22, 43, 56) of strains from patients

with persistent colonization did not cluster together in the core gene phylogeny despite having

the same ST. The reads-based nucleotide analysis of the 1,368 core genes identified 6454,

262, and 4 mutations in the three postpartum genomes relative to each prenatal genome and

the two strains from pairs 22 and 43 were classified as mutators. This classification is based on

the increased mutation rates and identification of mutations in genes encoding DNA replication

and proofreading machinery[24,26]. The same was true for the postpartum genome from patient

36, which acquired 204 mutations despite having the same ST and clustering with the prenatal

genome in the core gene phylogeny. While mutators have been described in other streptococcal

species including *Streptococcus pyogenes*[33], *S. pneumoniae*[34], *S. mutans*[25], and *S. iniae*[35], this

study is the first to describe mutators in GBS.

Mutators (or hypermutators) represent bacterial cells within a population that have a higher-

than-average mutation rate[24] and typically arise in bacterial populations undergoing adaptation

to stressful conditions[36,37]. They have also been linked to persistent bacterial infections, such as

Pseudomonas aeruginosa infections in the lungs of cystic fibrosis patients[38]. In most cases,

mutators arise at low frequencies due to DNA polymerase errors, dysfunctional proofreading

mechanisms, and/or failure of mismatch/error correcting machinery, including the MMR

pathway[24,26]. Dysfunction of this system increases the likelihood of horizontal gene transfer and

spread of antibiotic resistance[26]. Notably, all three mutators had mutations within the MMR-

encoding region, which comprised four distinct classes. Each mutator strain had a different

subclass, demonstrating that hypermutability is not associated with a specific MMR genotype,

but is more likely due to the mutations that disrupt the functionality of this system.

Each mutator also belongs to a different CC. More SNPs were detected among the non-mutator

persistent pairs belonging to CC-23 compared to those within CCs 17 and 19. The average

mutation rate across non-mutator strains of CCs 1, 12, and 23 (n=11, 3.66×10^{-6}) was also

significantly higher than the average mutation rate of the CC17 and CC19 non-mutator strains

(n=6, 1.37×10^{-6}; T-test p=0.02). Together, these observations support the classification of CCs

1, 12, and 23 as generalists, which are more vulnerable to genomic alterations with an

enhanced ability to colonize a wide range of hosts[14,39]. In contrast, CCs 17 and 19, which cause

higher frequencies of neonatal disease[14–16], are likely to be more specialized. Nonetheless, this

core-gene mutation analysis has some limitations including the possibility of infrequent

sequencing errors. To ensure that the variation observed between paired genomes was not due

to sequencing artifacts, however, we manually confirmed each mutation while determining that

each was in an area of >10x read coverage.

Studies in *S. pneumoniae* have identified increased mutation rates after antibiotic exposure[37,40],

suggesting that antibiotic stress may drive genetic diversity and contribute to more robust and fit

populations. Although three postpartum strains still acquired mutations without IAP exposure,

the average mutation rate (1.50×10^{-6}) was considerably lower than for the strains with IAP

exposure (1.16×10^{-4}). It is notable that one of these patients (ID=56) received cefazolin

following childbirth instead and the postpartum strain did not cluster with the prenatal strain in

the core-gene phylogeny, highlighting genomic variation. Consistent with these findings, >16%

of genes with mutations had evidence of positive selection, suggesting that IAP exerts a

selective pressure on GBS, fostering selection and expansion of more resilient strains. As

nonsynonymous SNPs are more likely to influence protein functionality and we observed

multiple genes with similar mutations, it is also likely that these genes and/or gene combinations

have evolved in parallel and enhance the colonization potential of strains in the vaginal tract.

Among the three mutators, one strain (genome ID 22.2) underwent a capsule switch from cps V

to VI after IAP, which was observed previously[41] and was suggested to occur via extensive

recombination events that introduce genomic variation beyond the *cps* loci[41,42]. Intriguingly, this

mutator strain also possessed MMR loci belonging to Class 3, which was only detected in two

cps VI genomes in our dataset and three others among ~100 published GBS genomes

evaluated. Although cps VI is less common in North America, it has been shown to predominate

in some regions[15,43], which could be attributable to differential selective pressures across

locations. Identification of capsule switching *in vivo* following IAP is concerning in that cps V

strains along with types Ia, Ib, II, III, and IV are responsible for 98% of invasive infections in

babies worldwide and are targeted in the GBS6 conjugate vaccine[44,45].

Indeed, a cps switch from type V to type VI in a mutator strain background could threaten the

effectiveness of this vaccine in some populations due to evasion of vaccine-induced immunity

along with the effectiveness of IAP. The same is true for mutations outside of the *cps* loci, as

mutators can increase the potential for recombination events through which rare alleles and

genotypes can arise. Because current GBS vaccines are based on only a subset of circulating

cps types[44], there are concerns that vaccination could introduce a selective pressure from which

rare and more resilient strains can prevail as has been observed for *S. pneumoniae*[46].

Nonetheless, using vaccines and antibiotics to protect newborns from invasive GBS infections

far outweighs the risks imposed by evolutionary events that could enhance immune evasion, but

also highlights the need for continuously monitoring strains circulating across populations.

Future studies should therefore focus on better characterizing these strains to fully understand

the role they play in colonization and disease.

METHODS

Bacterial isolate selection and characterization

A subset of 97 GBS isolates recovered from vaginal-rectal swabs from 58 pregnant patients[23]

were included in this analysis. Isolates were recovered from the same patient during a prenatal

(35-37 weeks gestation) and postpartum (6-weeks after childbirth) visit[23], and were previously

characterized by MLST, serotyping, and cps typing[22]. Patients with GBS at both samplings were

considered to have persistent colonization resulting in 80 strains from 41 patients. The

remaining 17 patients had GBS only at the prenatal sampling and were considered to have lost

the pathogen by the postpartum sampling.

Whole-genome sequencing (WGS)

A set of 41 of the 97 isolates was sequenced previously and the raw reads were downloaded

from the NCBI Sequence Read Archive (SRA) (**Table S2**). The remaining 56 isolates were

sequenced for this study following DNA extraction from overnight cultures of GBS grown in

Todd-Hewitt broth (THB) at 37°C + 5% CO_2 using the E.Z.N.A. (Omega Bio-tek Inc., Norcross,

GA, USA) or Wizard HMW (Promega, Madison, WI, USA) kits. Construction of DNA libraries

was performed by the Michigan Department of Health and Human Services using the Nextera

XT library prep kit (Illumina, San Diego, CA, USA) followed by sequencing on the MiSeq

(Illumina) with 2x250 bp paired end reads.

Raw reads for the 97 genomes were trimmed with Trimmomatic v0.39[47] using "gentle trimming"

parameters to trim adapters and remove sequences with average quality scores <15 or less

than 36 nucleotides in length. The paired-end reads, or single-end reads for SRR517012 that

was sequenced using an unpaired method, were assessed for quality using FastQC v0.11.9

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/); high-quality genomes were

defined as those that failed <10% of quality modules. High-quality trimmed reads were

assembled *de novo* using SPAdes v3.13.1 (kmers 21, 33, 55, 77, 99, 127) with mismatch error

correction[48] and quality was assessed using QUAST v5.0.2[49] and MultiQC v1.15[50] with default

parameters. Assemblies were considered high-quality if N50 and L50 scores were >15,000 and

<50, respectively. Assembled genomes were annotated using Prokka v1.14.6 with the –proteins

option that allows for the use of a custom database[51]. This database was curated from 25

closed *S. agalactiae* genomes downloaded from NCBI to ensure that gene naming was specific

for GBS (**Table S10**). STs were confirmed using PubMLST[29], though previously determined

STs[22] were used for three strains with poor assembly quality for one of the seven MLST loci.

Pangenome and phylogenetic analysis

Roary v3.11.2 was used to create a multiFASTA alignment of core genes from the 92 high-

quality assemblies using the -i (blastp of 95%) and -e (PRANK aligner 170427) parameters[52].

ML phylogenies were generated with RAxML v8.2.12 (parameters -m GTRGAMMA, 500

bootstrap replicates)[53], and phylogeny visualizations and annotations were performed using the

Interactive tree of life (iToL, https://itol.embl.de/). Analysis of the six MMR loci was performed by

manually extracting each gene from the assemblies using Geneious v2022.2.2

(https://www.geneious.com/) followed by a MUSCLE[27] nucleotide alignment and construction of

a neighbor-joining phylogeny in MEGA X[54]. Neighbor-net trees were generated in SplitsTree

v4.19.0[55] and the PHI test was used to detect recombination with a window size of 100 and a k

of 1; a p-value of <0.05 was considered significant.

Gene and nucleotide-level analyses

ARGs and virulence genes were extracted from the 92 assemblies with ABRicate

(https://github.com/tseemann/abricate) using five ARG databases (CARD[56], ARG-annot[57],

MEGARes[58], Resfinder[59], NCBI AMRFinderPlus[60]) and the virulence-finder-database (VFDB)[61],

respectively. A gene was considered present if it had ≥10% identity and ≥80% coverage and for

the ARGs, if it was detected in two or more databases. Antibiotic resistance phenotypes were

determined previously by disk diffusion[23] and were compared to the ARG data.

Core gene mutations were extracted from the assemblies of each pair of strains recovered from

patients with persistent colonization (n=64, 32 pairs). For each pair, the assembled prenatal

genome was used as the reference, while the trimmed reads from the postpartum isolate were

mapped against the reference. Each core genome was interrogated for mutations including

SNPs, insertions, deletions, MNPs, and complex mutations using Snippy v4.6.0

(https://github.com/tseemann/snippy). For discrepancies in gene presence/absence within a pair

of strains, Geneious v2022.2.2 was used to confirm ≥15X read coverage and the up and

downstream regions were mapped with respect to a given gene to confirm location and

completeness of the gene within the contig. Core gene mutations were validated using a ≥10X

read coverage cutoff.

Biofilm Assays

Biofilm assays were performed in triplicate on all but one pair of strains from persistently

colonized patients (n=62 isolates, 31 pairs). Values were normalized to blank media controls as

described[32] with the modification of stagnant growth for 24hrs. The magnitude (fold-change) of

the difference in biofilm formation was determined for the postpartum strain relative to the paired

prenatal strain.

Data analysis and statistics

Raw output from WGS analyses were managed in Microsoft Excel. R v4.1.2 (https://www.r-

project.org/) was used in RStudio v2022.07.1+554 (https://www.rstudio.com/) for importing data

(readr v2.1.4), data wrangling (devtools v2.4.5, dplyr v1.1.3, forcats v1.0.0, plyr v1.8.8, tidyr

v1.3.0, tidyverse v2.0.0), and visualization (ggplot2 v3.4.3, viridis v0.6.4) (https://cran.r-

project.org/). R package factoextra v1.0.7 was used to perform and visualize PCA with 95%

confidence ellipses for virulence genes using the prcomp() and fviz_pca() functions

(https://cran.r-project.org/). Chi-square tests were performed in Microsoft Excel or Epi Info™ v7

(Centers for Disease Control and Prevention, Atlanta, GA, USA); statistically significant

associations were classified at $p≤0.05$. For unnamed genes that acquired mutations, arbitrary

"gene ids" using the GBS_00X notation were assigned for easier reference. For all SNP-only

analyses, SNPs were filtered to exclude those that occurred in rRNA and/or uncharacterized,

non-coding regions.

As characterized by Snippy, nonsynonymous SNPs included those classified as "initiator codon

variant", "missense", "start lost", "stop lost", and "stop gained" (nonsense), while synonymous

SNPs included those referred to as "stop retained" and "synonymous variant". Measures of

evolutionary selection were calculated based on the nonsynonymous to synonymous SNP ratio

(πN/πS) across the genomes, as well as across each gene of interest to discern areas under

positive (πN/πS >1), neutral (πN/πS = 1), or negative (πN/πS <1) selection. Calculations for

evolutionary selection across genes were limited to genes that acquired a total of 5 or more

SNPs. Student t-tests were used to analyze differences in biofilm formation between isolate

pairs and mutation rates across CCs; a p-value <0.05 was deemed significant.

Data availability

The datasets are available in the supplementary tables and the raw sequencing reads for all 97

genomes have been deposited in GenBank.

Acknowledgements

We thank the Research Technology Support Facility and Michigan Department of Agriculture

and Rural Development for DNA sequencing, as well as the National Center for *Streptococcus*

(Edmonton, Alberta), particularly Marguerite Lovgren, for strain collection and serotyping.

Financial support was provided by the Michigan State University (MSU) Research Foundation

(to SDM) and in part, by the Michigan Sequencing and Academic Partnerships for Public Health

Innovation and Response (MI-SAPPHIRE) initiative at the MDHHS via CDC through the

Epidemiology and Laboratory Capacity for Prevention and Control of Emerging Infectious

Diseases Enhancing Detection Expansion program (6NU50CK000510-02-07; to HMB and

SDM). Additional support was provided by the National Institutes of Health (R01HD090061 to

JAG and SDM, R01AI134036 to JAG) and Department of Veterans Affairs Merit Award

I01BX005352 (Office of Research) to JAG. Student support was provided to MEP by the MSU

Department of MGI via the Philipp and Vera Gerhardt Travel and Ralph Evans Awards and the

MSU College of Natural Science via two Outstanding Scholar Fellowships.

Author contributions

Conceptualization: MEP, SDM; methodology: HMB, MEP; validation and formal analysis: MEP,

SDM; resources: SDM, HMB, HDD; supervision and insight: SDM, HMB, JAG, HDD; data

curation: MEP, SDM; writing—original draft preparation, MEP; writing—review and editing and

data visualization: MEP, SDM; funding acquisition: SDM, HMB, JAG. All authors have read and

agreed to the final version of the manuscript.

Competing interests

The authors declare no competing interests or other interests that could influence the

interpretation of the article.

REFERENCES

1. Russell, N. J. *et al.* Maternal colonization with group B *Streptococcus* and serotype distribution worldwide: Systematic review and meta-analyses. *Clin Infect Dis* **65** (suppl_2), S100-S111 (2017).

2. Boyer, K. M. & Gotoff, S. P. Prevention of early-onset neonatal group B streptococcal disease with selective intrapartum chemoprophylaxis. *N Engl J Med* **314**, 1665–9 (1986).

3. Verani, J. R., McGee, L., Schrag, S. J. Prevention of perinatal group B streptococcal disease--revised guidelines from CDC, 2010. *MMWR Recomm Rep* **59**, 1–36 (2010).

4. Phares, C. R. *et al.* Epidemiology of invasive group B streptococcal disease in the United States, 1999-2005. *JAMA* **299**, 2056-65 (2008).

5. Seale, A. C. *et al.* Stillbirth with group B *Streptococcus* disease worldwide: Systematic review and meta-analyses. *Clin Infect Dis* **65,** S125–S132 (2017).

6. Schuchat, A. Epidemiology of group B streptococcal disease in the United States: Shifting paradigms. *Clin Microbiol Rev* 11, 497-513 (1998).

7. Mei, J. Y. & Silverman, N. S. Group B *Streptococcus* in pregnancy. *Obstet Gynecol Clin North Am* **50**, 375–387 (2023).

8. Schrag, S. J. *et al.* Group B streptococcal disease in the era of intrapartum antibiotic prophylaxis. *N Engl J Med* **342**, 15–20 (2000).

9. Azad, M. B. *et al.* Impact of maternal intrapartum antibiotics, method of birth and breastfeeding on gut microbiota during the first year of life: A prospective cohort study. *BJOG* **123**, 983–993 (2016).

10. Schrag, S. J. & Verani, J. R. Intrapartum antibiotic prophylaxis for the prevention of perinatal group B streptococcal disease: Experience in the United States and implications for a potential group B streptococcal vaccine. *Vaccine* **31**, D20–D26 (2013).

11. Anthony, B. F., Okada, D. M. & Hobel, C. J. Epidemiology of the group B *Streptococcus*: Maternal and nosocomial sources for infant acquisitions. *J Pediatr* **95**, (1979).

12. Kwatra, G. *et al.* Prevalence of maternal colonisation with group B *Streptococcus*: a systematic review and meta-analysis. *Lancet Infect Dis* **16**, 1076–1084 (2016).

13. Van Dyke, M. K. *et al.* Evaluation of Universal Antenatal Screening for Group B *Streptococcus*. *Obstet Gynecol Surv* **64**, 703–704 (2009).

14. Jones, N. *et al.* Multilocus sequence typing system for group B *Streptococcus*. *J Clin Microbiol* **41**, 2530–2536 (2003).

15.   Bianchi-Jassir, F. *et al.* Systematic review of group B streptococcal capsular types,
sequence types and surface proteins as potential vaccine candidates. *Vaccine* **38**,
(2020).

16.   Manning, S. D. *et al.* Multilocus sequence types associated with neonatal Group B
streptococcal sepsis and meningitis in Canada. *J Clin Microbiol* **47**, 1143–1148 (2009).

17.   Springman, A. C. *et al.* Selection, recombination, and virulence gene diversity among
group B streptococcal genotypes. *J Bacteriol* **191**(17):5419-27 (2009).

18.   Korir, M. L. *et al.* Association and virulence gene expression vary among serotype III
group B *Streptococcus* isolates following exposure to decidual and lung epithelial cells.
Infect Immun **82**, (2014).

19.   Korir, M. L. *et al.* Differing mechanisms of surviving phagosomal stress among group B
Streptococcus strains of varying genotypes. *Virulence* **8**, 924–937 (2017).

20.   Korir, M. L. *et al. Streptococcus agalactiae cadD* alleviates metal stress and promotes
intracellular survival in macrophages and ascending infection during pregnancy. *Nat*
Commun **13**, (2022).

21.   Flaherty, R. A., Aronoff, D. M., Gaddy, J. A., Petroff, M. G. & Manning, S. D. Distinct
Group B *Streptococcus* sequence and capsule types differentially impact macrophage
stress and inflammatory signaling responses. *Infect Immun* **89**, e00647-20 (2021).

22.   Manning, S. D. *et al.* Genotypic diversity and serotype distribution of group B
Streptococcus isolated from women before and after delivery. *Clin Infect Dis* **46**, 1829–
1837 (2008).

23.   Spaetgens, R. *et al.* Perinatal antibiotic usage and changes in colonization and
resistance rates of group B *Streptococcus* and other pathogens. *Obstet Gynecol* **100**,
525–533 (2002).

24.   Miller, J. H. Spontaneous mutators in bacteria: Insights into pathways of mutagenesis and
repair. *Annu Rev Microbiol* **50**, 625–643 (1996).

25.   Gonzalez, K., Faustoferri, R. C. & Quivey, R. G. Role of DNA base excision repair in the
mutability and virulence of Streptococcus mutans. *Mol Microbiol* **85**, 361–377 (2012).

26.   Chopra, I., O'Neill, A. J. & Miller, K. The role of mutators in the emergence of antibiotic-
resistant bacteria. *Drug Resist Updat* **6**, (2003).

27.   Edgar, R. C. MUSCLE: Multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res* **32**, 1792–1797 (2004).

28.   Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment
search tool. *J Mol Biol* **215**, (1990).

29. Jolley, K. A., Bray, J. E. & Maiden, M. C. J. Open-access bacterial population genomics:
BIGSdb software, the PubMLST.org website and their applications. *Wellcome Open Res*
3, 1–20 (2018).

30. Bandara, M. *et al.* The accessory Sec system (SecY2A2) in *Streptococcus pneumoniae*
is involved in export of pneumolysin toxin, adhesion and biofilm formation. *Microbes*
Infect **19**, 402–412 (2017).

31. Noble, K. *et al.* Group B *Streptococcus cpsE* is required for serotype V capsule
production and aids in biofilm formation and ascending infection of the reproductive tract
during pregnancy. *ACS Infect Dis* **7**, (2021).

32. Parker, R. E. *et al.* Association between genotypic diversity and biofilm production in
group B *Streptococcus*. *BMC Microbiol* **16**, 1–12 (2016).

33. Scott, J., Thompson-Mayberry, P., Lahmamsi, S., King, C. J. & McShan, W. M. Phage-
associated mutator phenotype in group A *Streptococcus*. *J Bacteriol* **190**, 6290–6301
(2008).

34. Samrakandi, M. M. & Pasta, F. Hyperrecombination in *Streptococcus pneumoniae*
depends on an atypical *mutY* homologue. *J Bacteriol* **182**, (2000).

35. Rudenko, O., Engelstädter, J. & Barnes, A. C. Evolutionary epidemiology of
Streptococcus iniae: Linking mutation rate dynamics with adaptation to novel
immunological landscapes. *Infect Gen Evol* **85**, (2020).

36. Sniegowski, P. D., Gerrish, P. J. & Lenski, R. E. Evolution of high mutation rates in
experimental populations of *E. coli*. *Nature* **387**, (1997).

37. Mehta, H. H. *et al.* The essential role of hypermutation in rapid adaptation to antibiotic
stress. *Antimicrob Agents Chemother* **63**, (2019).

38. Oliver, A., Cantón, R., Campo, P., Baquero, F. & Blázquez, J. High frequency of
hypermutable *Pseudomonas aeruginosa* in cystic fibrosis lung infection. *Science (1979)*
288, (2000).

39. Richards, V. P. *et al.* Population gene introgression and high genome plasticity for the
zoonotic pathogen *Streptococcus agalactiae*. *Mol Biol Evol* **36**, (2019).

40. Henderson-Begg, S. K., Livermore, D. M. & Hall, L. M. C. Effect of subinhibitory
concentrations of antibiotics on mutation frequency in *Streptococcus pneumoniae*. *J*
Antimicrob Chemother **57**, 849–854 (2006).

41. Neemuchwala, A., Teatero, S., Athey, T. B. T., Mcgeer, A. & Fittipaldi, N. Capsular
switching and other large-scale recombination events in invasive sequence type 1 group
B *Streptococcus*. *Emerg Infect Dis* **22**, 1941–1944 (2016).

42.   Brochet, M. *et al.* Shaping a bacterial genome by large chromosomal replacements, the
evolutionary history of *Streptococcus agalactiae*. *Proc Natl Acad Sci U S A* **105**, 15961–
15966 (2008).

43.   Lachenauer, C. S. *et al.* Serotypes VI and VIII predominate among group B streptococci
isolated from pregnant Japanese women. *J Infect Dis* **179**, (1999).

44.   Buurman, E. T. *et al.* A novel hexavalent capsular polysaccharide conjugate vaccine
(GBS6) for the prevention of neonatal group B streptococcal infections by maternal
immunization. *J Infect Dis* **220**, (2019).

45.   Madhi, S. A. *et al.* Potential for Maternally Administered Vaccine for Infant Group B
Streptococcus. *N Engl J Med* **389**, (2023).

46.   Brueggemann, A. B., Pai, R., Crook, D. W. & Beall, B. Vaccine escape recombinants
emerge after pneumococcal vaccination in the United States. *PLoS Pathog* **3**, 1628–1636
(2007).

47.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: A flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

48.   Bankevich, A. *et al.* SPAdes: A new genome assembly algorithm and its applications to
single-cell sequencing. *J Comput Biol* **19**, 455–477 (2012).

49.   Gurevich, A., Saveliev, V., Vyahhi, N. & Tesler, G. QUAST: Quality assessment tool for
genome assemblies. *Bioinformatics* **29**, 1072–1075 (2013).

50.   Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: Summarize analysis results
for multiple tools and samples in a single report. *Bioinformatics* **32**, 3047–3048 (2016).

51.   Seemann, T. Prokka: Rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068–
2069 (2014).

52.   Page, A. J. *et al.* Roary: Rapid large-scale prokaryote pan genome analysis.
Bioinformatics **31**, 3691–3693 (2015).

53.   Stamatakis, A., Ludwig, T. & Meier, H. RAxML-III: A fast program for maximum likelihood-
based inference of large phylogenetic trees. *Bioinformatics* **21**, 456–463 (2005).

54.   Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. MEGA X: Molecular evolutionary
genetics analysis across computing platforms. *Mol Biol Evol* **35**, 1547–1549 (2018).

55.   Huson, D. H. SplitsTree: Analyzing and visualizing evolutionary data. *Bioinformatics* **14**,
68–73 (1998).

56.   Jia, B. *et al.* CARD 2017: Expansion and model-centric curation of the comprehensive
antibiotic resistance database. *Nucleic Acids Res* **45**, D566–D573 (2017).

57.   Gupta, S. K. *et al.* ARG-annot, a new bioinformatic tool to discover antibiotic resistance

genes in bacterial genomes. *Antimicrob Agents Chemother* **58**, 212–220 (2014).

58.   Doster, E. *et al.* MEGARes 2.0: A database for classification of antimicrobial drug, biocide

and metal resistance determinants in metagenomic sequence data. *Nucleic Acids Res*

48, D561–D569 (2020).

59.   Zankari, E. *et al.* Identification of acquired antimicrobial resistance genes. *J Antimicrob*

Chemother **67**, 2640–2644 (2012).

60.   Feldgarden, M. *et al.* Validating the AMRFINder tool and resistance gene database by

using antimicrobial resistance genotype-phenotype correlations in a collection of isolates.

Antimicrob Agents Chemother **63**, 1–19 (2019).

61.   Chen, L., Zheng, D., Liu, B., Yang, J. & Jin, Q. VFDB 2016: Hierarchical and refined

dataset for big data analysis - 10 years on. *Nucleic Acids Res* **44**, D694–D697 (2016).