Ontology-based modeling, integration, and analysis of heterogeneous clinical, pathological, and molecular kidney data for precision medicine

Yongqun Oliver He, PhD[1]*; Laura Barisoni, MD[2]; Avi Z Rosenberg, MD, PhD[3]; Peter Robinson, MD[4]; Alexander D. Diehl, PhD[5]; Yichao Chen, BA[6]; Jimmy P. Phuong, PhD[7]; Jens Hansen, MD[8]; Bruce W. Herr II, BS[9]; Katy Börner, PhD[9]; Jennifer Schaub, MD[1]; Nikki Bonevich[1]; Ghida Arnous[1]; Saketh Boddapati[1]; Jie Zheng, PhD[1]; Fadhl Alakwaa, PhD[1]; Pinaki Sarder, PhD[10]; William D. Duncan, PhD[10]; Chen Liang, PhD[11]; M. Todd Valerius, PhD[12]; Sanjay Jain, MD, PhD[13]; Ravi Iyengar, PhD[8]*; Jonathan Himmelfarb, MD[7]; Matthias Kretzler, MD[1]; and Kidney Precision Medicine Project
(* Co-corresponding authors)

[1]University of Michigan, Ann Arbor, MI, USA; [2]Duke University, Durham, NC, USA; [3]Johns Hopkins University School of Medicine, MD, USA; [4]The Jackson Laboratory for Genomic Medicine, Farmington, CT, USA; [5]University at Buffalo, Buffalo, NY, USA; [6]Pennsylvania State University, State College, PA, USA; [7]University of Washington, Seattle, WA, USA; [8]Icahn School of Medicine at Mount Sinai, New York, NY, USA; [9]Indiana University, Bloomington, IN, USA; [10]University of Florida, Gainesville, FL, USA; [11]University of South Carolina, Columbia, SC, USA; [12]Brigham and Women's Hospital, Boston, Massachusetts, USA; [13]Washington University, St. Louis, MO, USA.

Abstract
Many data resources generate, process, store, or provide kidney related molecular, pathological, and clinical data. Reference ontologies offer an opportunity to support knowledge and data integration. The Kidney Precision Medicine Project (KPMP) team contributed to the representation and addition of 329 kidney phenotype terms to the Human Phenotype Ontology (HPO), and identified many subcategories of acute kidney injury (AKI) or chronic kidney disease (CKD). The Kidney Tissue Atlas Ontology (KTAO) imports and integrates kidney-related terms from existing ontologies (e.g., HPO, CL, and Uberon) and represents 259 kidney-related biomarkers. We also developed a precision medicine metadata ontology (PMMO) to integrate 50 variables from KPMP and CZ CellxGene data resources and applied PMMO for integrative kidney data analysis. The gene expression profiles of kidney gene biomarkers were specifically analyzed under healthy control or AKI/CKD disease statuses. This work demonstrates how ontology-based approaches support multi-domain data and knowledge integration in precision medicine.

Introduction

The Kidney Precision Medicine Project (KPMP) (https://www.kpmp.org/) is an NIH/NIDDK-funded consortium that aims to characterize with precision the complexity of chronic kidney disease (CKD) and acute kidney injury (AKI) at the patient level to increase our ability to improve personalized treatment (1). While AKI is a sudden and often temporary loss of kidney function, CKD causes reduced kidney function over a long period of time and can lead to end-stage kidney disease. However, transition from AKI to CKD can also be observed. Both AKI and CKD are associated with complex pathogenesis involving the interactions among genetic, pathological, molecular, social, and environmental factors. Despite significant effort from the nephrology community, the mechanisms underlying the development and progression of kidney diseases are not yet fully understood, in part because of challenges to integrate data from multiple knowledge domains. Hence, integration of different types of data related to kidney diseases deserves to be the subject of further in-depth investigations.

Significant effort has been made recently to generate kidney related data and make them publicly available to the community of investigators. The Human BioMolecular Atlas Program (HuBMAP) aims to develop an open and global platform to map healthy cells in the human body (2). The Human Cell Atlas project (HCA) is a global consortium focused on mapping every cell type in the human body and developing a 3-dimensional atlas of human cells to

transform our understanding of biology and disease (3). The CellxGene resource, which is closely associated with the HCA, is a suite of computational tools that help scientists to deposit, download, query, and visually explore curated and standardized single-cell biology datasets (4).

Data integration remains a big challenge due to heterogeneity of data within and between domains. Interoperable ontologies support data standardization, sharing, and integration. Many reference biomedical ontologies such as Cell Ontology (CL) (5), Uber-anatomy ontology (Uberon) (6), and Human Phenotype Ontology (HPO) (7) have been widely used across different data resource projects, which significantly promote standard data annotation. We have developed the Kidney Tissue Atlas Ontology (KTAO) (8) to support kidney tissue knowledge representation. The HuBMAP Common Coordinate Framework Atlas ontology (CCFO) is an application ontology built to support the development of the Human Reference Atlas (HRA) by the HuBMAP project (9). KTAO and CCFO use the same set of reference ontologies such as CL and Uberon, which supports the shared knowledge query and analysis. However, still missing are ontologies to integrate metadata that can support data standardization and integration across domains.

This study describes our ontology-based strategy to address the data integration challenge to support kidney precision medicine research. We have developed an integrated precision medicine metadata ontology (PMMO) to collect, map, standardize, and integrate various metadata variables from data resources. PMMO, KTAO, and CCFO are used together to support our kidney precision medicine use case studies. This work demonstrates how ontology-based approaches support data and knowledge integration in precision medicine.

Methods

Figure 1 illustrates the general project design for integration of data across domains. First, we developed PMMO to map, standardize, and integrate metadata variables from multiple resources including KPMP, HuBMAP, CellxGene, and HCA. HuBMAP and CellxGene only contain single cell transcriptomic data; it is notable that the metadata types related to the cell types and the tissues/species/phenotypes of these cell types may differ between HuBMAP and CellxGene. PMMO, however, well represents all types of cell types and the tissues/species/phenotypes. Second, we focus on the ontology-based knowledgebase, which represents knowledge with ontologies such as CL, Uberon, HPO, GO, and OPMI. As a related use case, we have identified various healthy and kidney disease gene/protein biomarkers from CCFO and KTAO, respectively. These biomarker genes' expression profiles from various datasets stored in CellxGene will be queried and analyzed. Since CellxGene does not contain all the metadata of KPMP kidney disease studies (e.g., diabetes phenotype), the metadata ontology will be used to cross-link metadata details in an ontology-standardized manner. The metadata ontology will include sufficient granularity to ensure the cross integration among different data resources.

Figure 1. General project design.

Kidney phenotype annotation and submission to HPO. Starting from 2017, KPMP has organized multiple workshops in which clinical nephrologists, pathologists, and ontologists worked together to discuss and apply a high level framework for annotating, collecting, and proposing new terms or modifications to HPO (7) and underlying ontologies such as Uberon (6). After internal collection of terminologies and discussion within KPMP, Excel spreadsheets of new terms, modifications, definitions, and proposed hierarchies were prepared and sent to HPO developers. Individual HPO GitHub issue trackers were then generated for individually proposed terms and tagged with "KPMP". After further discussion in GitHub and with KPMP investigators, the newly proposed kidney related HPO terms were added to the HPO and subject to public checking. The released terms were then further imported back to the KTAO using the tool Ontofox (10).

Precision Medicine Metadata Ontology (PMMO) – Ontological representation and integration of metadata types from data resources in KPMP and CellxGene. KPMP includes various metadata types associated with clinical, pathological, and molecular aspects (https://www.kpmp.org/available-data). Clinical data were obtained from electronic health records and stored using RedCap (11). Kidney tissue-related data were obtained and stored at the KPMP Tissue Interrogation Sites (TIS) data and pathology data resource. The CellxGene has 10 common variables (e.g., age, disease) (4). The PMMO ontologically models and represents the KPMP-derived data, the 10 common variables (e.g., age, disease) derived from CellxGene, and their semantic relations. Additional new terms along with these variable terms were also added to enable us to build up the semantic relations. Pre-existing ontology terms are reused. The ontology of precision medicine and investigation (OPMI) (8) was used as the default ontology for the metadata ontology development.

Kidney disease modeling and analysis with heterogeneous data from multiple resources. Different types of kidney diseases (including AKI, CKD, and others) and their associated pathological features were first modeled using an ontological approach. Then the data from different resources using KPMP and CellxGene were used for identifying, integrating, and analyzing the data for the patients related to the modeled kidney diseases.

Kidney biomarkers collection and representation in KTAO. Gene/protein biomarkers related to kidney cell types and diseases were manually collected from three resources: (i) KPMP consortium, (ii) our manual literature mining, and (iii) HuBMAP (for reference kidney cell biomarkers). The annotated biomarkers were then added to the KTAO ontology using a standard format. SPARQL (a recursive acronym for SPARQL Protocol and RDF Query Language) queries were generated to extract the biomarker records from HuBMAP CCFO and KPMP KTAO ontologies. The Ontobee SPARQL endpoint (12) was used for the query generation.

Extraction of kidney biomarker gene expression profiles using CellxGene and KPMP Atlas explorer. The CellxGene API was used to extract gene expression profiles from specific datasets. KPMP atlas explorer was also used for data browsing and analysis. Expression levels for the SPP1 gene were mapped onto relevant cell types in CL using the Graphviz dot language and the ontology hierarchy was visualized with Graphviz (13).

Results

Ontological modeling of kidney diseases and associated pathological phenotypes:
We collected hundreds of kidney specific phenotypes and found many of them were not yet in the HPO. Since 2017, our KPMP kidney community has been working closely with HPO developers on the ontological modeling of kidney phenotypes. As of March 16, 2024, HPO had received 329 kidney specific phenotype terms contributed by KPMP and added most of these terms to HPO. These and other kidney related HPO phenotype terms were later imported back to the Kidney Tissue Atlas Ontology (KTAO), which aims to collect and interlink kidney phenotypes and other types of entities (e.g., cells, anatomical entities, biomarkers) all together. By doing so, the KPMP team contributes to the community effort of kidney phenotype collection and modeling, and meanwhile uses the newly developed phenotype ontology models to support KPMP research goals.

A more recent effort is to further model and analyze novel and more specific kidney phenotypes that cross and overlap different existing phenotypes. Our KPMP team has been focusing on AKI and CKD. Interestingly, AKI and CKD are not always distinct entities and overlap or transition of clinical and pathologic characteristics between AKI and CKD can be observed. This definition results in a spectrum of phenotypes (Figure 2). Both AKI and CKD patients may have diabetes and/or hypertension (Figure 2A). For example, patients without any other pathologic conditions may experience AKI alone (i.e., following use of nephrotoxic antibiotics), but the same AKI phenotype may be

superimposed on an underlying diabetic nephropathy phenotype under similar circumstances (diabetic patient with recent use of nephrotoxic antibiotics developing acute renal failure). Similarly, patients with CKD may have characteristics typical of diabetic nephropathy, which is generally a chronic damage phenotype; however, some patients present with a mixed phenotype of chronic active damage, where, in the absence of an obvious trigger, a smoldering acute process occurs hand to hand with the development of chronicity (Figure 2B).

Figure 2. Pathological kidney disease categorization. (A) Venn diagram illustrating overlap between CKD and AKI, and how clinical diabetes and hypertension may overlap with both kidney diseases. (B) Venn diagram illustrating partial overlap in kidney disease manifestation of AKI and CKD. Each of kidney disease states may be associated with specific features indicated by the different terms, some of which are AKI or CKD specific (A,B and I,J, respectively), and others reflects overlap (C, D; G, H; K, L; M, N; O, P; S, T; U, V).

Our research found that shared phenotypes may come with specific features newly identified in different domains . As a result, the overlapping phenotypes may deserve new ontological term representation (Figure 2B). Specifically, we were able to identify two specific disease categories associated with AKI (Figure 2B): Acute Interstitial Nephritis (AIN) and Acute Interstitial Nephritis superimposed on Diabetic Nephropathy (AINs-DN); and two specific disease categories associated with CKD: Diabetic Nephropathy with Chronic Active Interstitial Nephritis (DN-CAIN) and Diabetic Nephropathy (DN).

Many of these detailed kidney disease categories are not yet in HPO, and our modeling of these diseases (Figure 2B) is also unique in that it provides a strategy to identify various disease characteristics that are indicative of these specific disease categories, thus, enabling a more precise definition of disease.

The identification of these distinct disease states and their characteristics is a focus of our current KPMP research. One step towards this goal is the work of the KPMP adjudication committee, which routinely conducts a pathology focused adjudication process that defines disease categories for individual human subject patients enrolled in the KPMP study. A descriptor scoring approach is also used to score different categories of morphologic parameters. After the adjudication based categorization, these kidney patients are being closely monitored to identify unique parameters indicative of their disease categories.

Integrative KTAO development:
As shown in Figure 3, KTAO is an application ontology that serves as the knowledgebase of kidney tissue atlas. As of March 16, 2024, KTAO has 11,882 terms and 175,926 axioms. The KTAO terms include 11,210 classes, 399 object properties, 26 data type properties, 207 annotation properties, and 40 instance terms. In ontology, an axiom represents a relation among terms or entities. Here is an example of an axiom representing the anatomical location of the podocyte (CL_0000653): *'part of' some 'visceral layer of glomerular capsule'* (UBERON_0005751). With so many axioms represented, KTAO serves as its own knowledgebase of the kidney tissue atlas.

Figure 3. KTAO development workflow and statistics. As of February 26, 2024, our KPMP team had contributed to HP 329 terms and 327 of them had been added to HPO and then imported back to KTAO. We also collected 252 kidney related biomarkers and represented them in KTAO. KTAO has also imported more than 10,000 terms from over 40 existing ontologies. Furthermore, KTAO generates KTAO-specific terms not available in other ontologies.

KTAO reuses existing ontologies such as CL, HPO, and Uberon. KTAO imports terms from 32 ontologies with at least 10 terms included (https://ontobee.org/ontostat/KTAO). Table 1 lists the statistics of terms in KTAO.

Table 1. KTAO statistics of terms including those imported from other ontologies and KTAO-specific terms.

Ontology used in KTAO (selected)	No. of terms in KTAO
Cell Ontology (CL)	386
Human Phenotype Ontology (HPO)	960
Uber-anatomy ontology (Uberon)	1,171
Gene Ontology (GO)	1,225
Protein Ontology (PR)	316
Ontology of Genes and Genomes (OGG)	483
Ontology of Precision Medicine and Investigation (OPMI)	864
KTAO-specific (with KTAO namespace)	53
Total in KTAO (version 1.0.121, 2024-03-16)	11,882

PMMO development and application for precision medicine metadata integration:
To efficiently analyze heterogeneous data from different data resources, it is critical to integrate the data types at the metadata level. However, such a task is often challenging. As illustrated in Figure 4, we focused on the task of integrating three specific metadata types: (i) 18 KPMP clinical metadata types focused on clinical data, (ii) 29 KPMP biopsy TIS metadata, and (iii) 10 CellxGene metadata types. Our PMMO systematically models and represents these metadata types in an integrative semantic framework.

We realized that even with small numbers of metadata elements, significant issues arise in our integration and modeling of the three metadata sets (Figure 4). For example, the core data type 'patient ID' is defined using different labels: 'patient ID' in the KPMP Biopsy TIS data, 'RedCap study ID' in KPMP Clinical data, and 'donor ID' in

CellxGene data. While these terms are used in different scenarios, we found that they refer to the same IDs and can be used as the most important metadata elements to interlink different data resources. Another example is the representation of 'disease' in different data resources: 'disease' in CellxGene data, 'tissue type' in KPMP clinical data, and 'condition' in KPMP Biopsy TIS data (Figure 4). In addition to 'tissue type', the KPMP clinical data file has another tissue related metadata element called "Tissue Source". Here, "tissue type" refers to specific disease information (e.g., AKI, CKD, and healthy), and "Tissue Source" refers to where to obtain the tissue (e.g., KPMP TIS). Mapped to the "Tissue" and "Species" in CellxGene, the two KPMP datasets have two hidden values: kidney for "Tissue" and human for "Species". Therefore, the three metadata sets share four data types: patient ID, disease, tissue, and species, which have been properly identified and modeled in PMMO.

In addition to the core terms shared by all the three metadata sets, we also found many metadata elements unique to each of these sets or shared by only between two of the three sets (Figure 4).

The PMMO representation reuses terms from reference ontologies. We first mapped the different metadata elements to terms from reference ontologies, and then integrated them together in one ontology format. The OPMI is used as the default ontology for the PPMO ontology development.

Figure 4. Venn Diagram of metadata variables from three specific data resources.

An important use case of the PPMO is in integrating information from different sources. Guided by the PPMO modeling, we were able to trace the kidney precision medicine data across different data resources using KPMP/CellxGene web browsers (Figure 5) or API. For example, using the KPMP repository report program, we are able to identify the records of an anonymized human subject with the patient ID of "27-10039" (Figure 5A). This human subject is a CKD patient (female, white, 60-69 years old) with a diabetes history of 30-40 years and a hypertension history of 30-40 years (Figure 5B). The expression profiles of the gene *SLC5A2* (solute carrier family 5 member 2, commonly known as SGLT2) are shown in KPMP spatial transcriptomics format (Figure 5C) or KPMP gene expression format (Figure 5D). Furthermore, CellxGene also provides specific profiles for this human donor with the donor ID as the same as the KPMP patient ID (Figure 5E). The PMMO to be built will include the linkage information to interlink datasets between different datasets.

Figure 5. Integrative exploration of the records of an anonymized KPMP human subject (patient ID: 27-10039) using KPMP or CellxGene web programs. (A) Basic KPMP information of records associated with this patient. (B) Clinical data. (C) KPMP spatial transcriptomics related to the example gene SLC5A2. (D) KPMP gene expression profile for SLC5A2. (E) CellxGene record whose donor ID equals to KPMP patient ID.

Using the same approach as laid out above, we can essentially search for all the datasets across the different sources including KPMP clinical data, KPMP biopsy TIS data, and CellxGene. KPMP systematically collects and analyzes the records and kidney biopsy samples from well-studied human subjects with AKI, CKD or reference healthy controls. The related cell gene expression data from the KPMP studies have also been submitted to CellxGene. It is important to note that different data sources use the same IDs although these IDs have different labels (patient ID, RedCap study ID, or donor ID) (Figure 4). Different databases also have different variables. For example, the CellxGene source has 10 variables (e.g., sex, age, disease), some of which are unique or shared with other databases (Figure 4). These variables can be analyzed using the web browsers or via the CellxGene API.

KPMP kidney gene/biomarker collection, annotation, and analysis:
Our research identified 529 kidney gene biomarkers (Figure 6A). These biomarkers include 192, 26, and 19 gene biomarkers unique for healthy reference kidney or kidney cell types, CKD, or AKI, respectively. In addition, these biomarkers include three sets of biomarkers shared by two groups: 3 shared by healthy and CKD groups, 5 shared by healthy and AKI groups, and 14 shared by AKI and CKD groups (Figure 6A). These biomarkers were ontologically represented in KTAO.

Kidney biomarkers were also analyzed using different resources. Specifically, we focused on the 38 AKI biomarkers as shown in Figure 6A using the KPMP datasets described in a *Nature* paper (14). The dataset is also stored in the CellxGene with a specific Dataset_id. Note that only 26 out of these 38 AKI biomarker genes had data in the CellxGene dataset. These 26 genes include 5 genes shared by healthy and AKI groups (red color), 8 genes shared by CKD and AKI groups (blue color), and 13 gene markers unique to AKI (green color) (Figure 6B-D). Our results show that some biomarkers can be confirmed to be good markers, but some not. These biomarkers have different profiles.

Figure 6. Kidney gene biomarker analysis. (A) Kidney biomarkers collected in KTAO. (B-D) The gene expression profiles of 26 AKI biomarker genes using the KPMP study labeled "Single-nucleus RNA-seq of the Adult Human Kidney (Version 1.0)" (Dataset_id in CellxGene: 07854d9c-5375-4a9b-ac34-fa919d3c3686). (B) AKI state. (C) CKD state. (D) healthy state. CellxGene API was used to generate the (B-D) results. The colors in (A) match the same colored boxes in (B-D).

While Figure 6 provides a detailed presentation of 26 gene biomarkers' gene expression profiles among 16 kidney cell types, these 16 cell types have additional relationships that are not observable in the figure. To address relationships between cell types, we mapped these cell types to the Cell Ontology (CL) and used the CL hierarchy to identify the relations among these cell types (Figure 7). By doing so, we were able to obtain more insights. For example, with the help of the CL hierarchical structure, we were able to find that SPP1 was highly expressed in the high level class term "nephron tubule epithelial cell" (CL_1000494) in the healthy controls and AKI patients (but not in CKD patients), esp. in its subclass of "kidney loop of Henle ascending limb epithelial cell" (CL_1000849), which includes two specific cell types: "kidney loop of Henle thin ascending limb epithelial cell" (CL_1001107) and "kidney loop of Henle thick ascending limb epithelial cell" (CL_1001106). Furthermore, the SPP1 expression profile in the "kidney loop of Henle thick ascending limb epithelial cell" appears to be distinct among three disease statuses (healthy, AKI, and CKD) (Figure 7). Since SSP1 protein functions as a cytokine, these differential expression patterns can help us identify changes in signaling functions across cell types and the different regions of the nephron. If these changes in signaling leads to changes in metabolism and/or transport, it is reasonable to hypothesize these changes could be reflected in new metabolites in the urine that could serve as biomarkers.

Figure 7. Ontology-based analysis of SPP1 gene expression profiles in 16 different cell types. Cell types arranged in the Cell Ontology (CL) hierarchy for kidney cell types were colorized based on their measured expression of the SPP1 gene, with darker shades indicating higher expression levels, based on the $\log_{10}$ of the expression value.

Discussion and Conclusions

This study describes several substantive advances we have made in providing a semantic framework for the kidney atlas. First, we are developing an approach for kidney-specific histological and pathological phenotypes modeling, leading to a collaborative effort of adding more than 300 kidney-specific phenotype terms to the HPO ontology. We also report our continued identification and ontological modeling of many precise kidney disease subcategories (Figure 2). Second, KTAO was applied to systematically integrate various types of kidney related knowledge at the ontological level. For example, KTAO imports terms and axioms related to kidney cells, anatomic entities, and phenotypes from CL, Uberon, and HPO, respectively. It also collects and annotates various kidney related biomarkers from KPMP and HuBMAP collections and our own literature mining. Third, we have developed the PPMO and applied it to seamlessly integrate data from different data resources (e.g., KPMP clinical data resource, KPMP tissue interrogation and Omics data resource, and CellxGene resource). Fourth, we have applied our ontological modeling and analysis strategy for kidney disease use case studies, with a focus on the AKI and CKD disease characteristics and gene biomarker profile analysis using different resources.

We find continuous challenges in systematically modeling and analyzing many specific kidney disease sub-categories. For example, our KPMP disease focused analyses enumerated many kidney disease subcategories by overlapping higher level disease categories such as including AKI, CKD, and diabetes (Figure 2). These disease subcategories appear to have their unique features, and our precision medicine approaches provide a feasible strategy to find these unique features by profiling and integrating clinical, pathological, cellular, and molecular characteristics related to an increasing pool of human subjects with these specific diseases. To further categorize and identify unique features of these disease subcategories in multiple domains, a collaborative effort involving kidney clinicians, pathologists, ontologists and bioinformaticians is required.

To address the challenge of integrating multi-domain features we propose to further develop KTAO or separately develop a new integrative application ontology tentatively called "Pathomics Ontology" (or Histology Descriptors and Pathomics Ontology - HDPO). The KPMP analysis of tissue includes a variety of methodologies, including i) disease adjudication, which takes in account the clinical data and the overall histology, ii) the descriptor scoring, which is a detailed visual assessment and scoring of all kidney tissue compartments, and iii) the pathomic features, which uses computational image analysis to quantify morphometric and textural tissue characteristics extracted from automatically segmented functional tissue units. The HDPO will combine the pathology phenotyping and descriptor scoring together with the pathomic features units and omics data. The goal is to identify unique features associated with specific pathological phenotypes at the cell, structure, and biopsy level. The newly proposed ontology will reuse terms from existing ontologies (e.g., CL, Uberon, and HPO), and it will also incorporate many other concepts such as histological descriptors, pathomic features, and Omics-related terms from spatial transcriptomics, with the goal to better support systematic and integrative modeling and analysis of newly identified disease or phenotype subcategories and unique features associated with these subcategories. The KPMP descriptor scoring and computational image analysis processes provide a solid base and model for such ontology development and application.

Data from different resources can be integrated in different ways. At the level of knowledge (a special type of annotated data), an efficient way is to incorporate the learned knowledge into ontologies and knowledge graphs, and then use programming languages such as SPARQL to perform queries and analysis. For example, we successfully developed SPARQL scripts to query and analyze biomarker knowledge from the KTAO and HuBMAP CCFO. At the instance data level, we posit that metadata integration is critical for integration of data from different data sources. Our research has identified the critical need for a metadata ontology for integrating different precision medicine related data sources, which led further to our initiation of the PPMO metadata ontology development. We demonstrate applications of PPMO in our kidney precision medicine research. This type of metadata ontology and its associated methods can also be applied to study other organs and diseases including the heart, liver, and vasculature.

Our ontology-based integrative approaches using knowledge from multiple domains can drive discovery of new biomarkers. While hundreds of kidney gene biomarkers have been found, our research focused on data integration has found that not all these biomarkers are equally expressed well in kidney cells. Using our ontology based methodology, it should be possible to identify the key cell types involved in the different kidney disease states and find a small number of gene biomarkers that work well at individual cell levels.

Acknowledgements
The KPMP is funded by the following grants from the NIDDK: U01DK133081, U01DK133091, U01DK133092, U01DK133093, U01DK133095, U01DK133097, U01DK114866, U01DK114908, U01DK133090, U01DK133113, U01DK133766, U01DK133768, U01DK114907, U01DK114920, U01DK114923, U01DK114933, U24DK114886, UH3DK114926, UH3DK114861, UH3DK114915, UH3DK114937. Human Reference Atlas (HRA) research and development is funded by the NIH Common Fund through the Office of Strategic Coordination/Office of the NIH Director under awards OT2OD033756 and OT2OD026671, by the Cellular Senescence Network (SenNet) Consortium through the Consortium Organization and Data Coordinating Center (CODCC) under award number U24CA268108, and by the NIDDK under awards U24DK135157 and U01DK133090. The work has also been supported by the Kidney Precision Medicine Project grant U2CDK114886, and the NIH National Institute of Allergy and Infectious Diseases (NIAID), Department of Health and Human Services under BCBB Support Services Contract HHSN316201300006W/HHSN27200002.

References

1. de Boer IH, Alpers CE, Azeloglu EU, Balis UGJ, Barasch JM, Barisoni L, et al. Rationale and design of the Kidney Precision Medicine Project. Kidney Int. 2021;99(3):498-510.
2. Hu BC. The human body at cellular resolution: the NIH Human Biomolecular Atlas Program. Nature. 2019;574(7777):187-92.
3. Rozenblatt-Rosen O, Stubbington MJT, Regev A, Teichmann SA. The Human Cell Atlas: from vision to reality. Nature. 2017;550(7677):451-3.
4. Biology CS-C, Abdulla S, Aevermann B, Assis P, Badajoz S, Bell SM, et al. CZ CELLxGENE Discover: A single-cell data platform for scalable exploration, analysis and modeling of aggregated data. bioRxiv. 2023:2023.10. 30.563174.
5. Diehl AD, Meehan TF, Bradford YM, Brush MH, Dahdul WM, Dougall DS, et al. The Cell Ontology 2016: enhanced content, modularization, and ontology interoperability. J Biomed Semantics. 2016;7(1):44.
6. Mungall CJ, Torniai C, Gkoutos GV, Lewis SE, Haendel MA. Uberon, an integrative multi-species anatomy ontology. Genome Biol. 2012;13(1):R5.
7. Kohler S, Gargano M, Matentzoglu N, Carmody LC, Lewis-Smith D, Vasilevsky NA, et al. The Human Phenotype Ontology in 2021. Nucleic Acids Res. 2021;49(D1):D1207-D17.
8. Ong E, Wang LL, Schaub J, O'Toole JF, Steck B, Rosenberg AZ, et al. Modelling kidney disease using ontology: insights from the Kidney Precision Medicine Project. Nat Rev Nephrol. 2020;16(11):686-96.
9. Herr BW, 2nd, Hardi J, Quardokus EM, Bueckle A, Chen L, Wang F, et al. Specimen, biological structure, and spatial ontologies in support of a Human Reference Atlas. Sci Data. 2023;10(1):171.
10. Xiang Z, Courtot M, Brinkman RR, Ruttenberg A, He Y. OntoFox: web-based support for ontology reuse. BMC Res Notes. 2010;3:175.
11. Harris PA, Taylor R, Minor BL, Elliott V, Fernandez M, O'Neal L, et al. The REDCap consortium: Building an international community of software platform partners. J Biomed Inform. 2019;95:103208.
12. Ong E, Xiang Z, Zhao B, Liu Y, Lin Y, Zheng J, et al. Ontobee: A linked ontology data server to support ontology term dereferencing, linkage, query and integration. Nucleic Acids Res. 2017;45(D1):D347-D52.
13. Gansner ER, North SC. An open graph visualization system and its applications to software engineering. Software: practice and experience. 2000;30(11):1203-33.
14. Lake BB, Menon R, Winfree S, Hu Q, Melo Ferreira R, Kalhor K, et al. An atlas of healthy and injured cell states and niches in the human kidney. Nature. 2023;619(7970):585-94.