Lipopeptide antibiotics disrupt interactions of undecaprenyl phosphate with UptA

Abraham O. Oluwole[1,2*], Neha Kalmankar[1,2], Michela Guida[2,3], Jack L. Bennett[1,2], Giovanna Poce[3], Jani R. Bolla[2,4], Carol V. Robinson[1,2*]

[1]Department of Chemistry, University of Oxford, South Parks Road, Oxford, OX1 3QZ, UK. [2]The Kavli Institute for Nanoscience Discovery, Oxford, South Parks Road, Oxford, OX1 3QU, UK. [3]Department of Chemistry and Technologies of Drug, Sapienza University of Rome, 00185 Rome, Italy. [4]Department of Biology, University of Oxford, South Parks Road, Oxford, OX1 3RB, UK.

Email: abraham.oluwole@chem.ox.ac.uk; carol.robinson@chem.ox.ac.uk

Keywords: Peptidoglycan, undecaprenyl phosphate transporter A, UptA, native mass spectrometry

Abstract: The peptidoglycan pathway represents one of the most successful antibacterial targets with the last critical step being the flipping of carrier lipid, undecaprenyl phosphate (C_{55}-P), across the membrane to re-enter the pathway. This translocation of C_{55}-P is facilitated by DedA and DUF368 domain-containing family membrane proteins via unknown mechanisms. Here we employ native mass spectrometry to investigate the interactions of UptA, a member of the DedA family of membrane protein from *Bacillus subtilis*, with C_{55}-P, membrane phospholipids and cell wall-targeting antibiotics. Our results show that UptA, expressed and purified in *E. coli*, forms monomer-dimer equilibria, and binds to C_{55}-P in a pH-dependent fashion. Specifically, we show that UptA interacts more favourably with C_{55}-P over shorter-chain analogues and membrane phospholipids. Moreover, we demonstrate that lipopeptide antibiotics, amphomycin and aspartocin D, can directly inhibit UptA function by out-competing the substrate for the protein binding, in addition to their propensity to form complex with free C_{55}-P. Overall, this study shows that UptA-mediated translocation of C_{55}-P is potentially mediated by pH and anionic phospholipids and provides insights for future development of antibiotics targeting carrier lipid recycling.

Introduction

The global threat of drug-resistant bacterial pathogens calls for renewed efforts in the search for new antibiotics, along with the identification of new biochemical pathways and interactions that can be effectively inhibited. A key antibiotic target is peptidoglycan, the cell wall polymer that provides bacteria with resistance to osmotic stress and environmental assaults (1, 2). Accordingly, topline antibiotics such as amoxicillin and vancomycin (3, 4) inhibit the transpeptidation step of the peptidoglycan synthesis (5-7). Moreover, promising antibiotic candidates, such as ramoplanin and moenomycin, target the transglycosylation step (8, 9). Peptidoglycan biosynthesis begins with the formation of uridine diphosphate-*N*-acetylmuramyl-pentapeptide (UM5) in the cytosol (10), followed by its coupling

to the lipid carrier undecaprenyl monophosphate (C_{55}-P) on the cytosolic side of the cytoplasmic membrane (11, 12). The resulting lipid I is further decorated with a GlcNAc residue to form lipid II (13) before being flipped across the cytoplasmic membrane (14-16). In the periplasm, the headgroup of lipid II is polymerized and cross-linked into the existing meshwork (17-19). The lipid carrier is then released as undecaprenyl diphosphate (C_{55}-PP). To participate in the next round of precursor transfer, C_{55}-PP must be dephosphorylated to form C_{55}-P (20-22) and then flipped across the cytoplasmic membrane such that its phosphate headgroup returns to the cytoplasmic side. The mechanism by which C_{55}-P is translocated across the cytoplasmic membrane is the least understood of the membrane-associated steps of the peptidoglycan pathway (Fig. 1A).

Recently, proteins belonging to the DedA- and the DUF368 domain-containing families were shown to facilitate trans-bilayer transport of C_{55}-P in the Gram-positive bacteria *Bacillus subtilis* and *Staphylococcus aureus* (23), and in the Gram-negative bacteria *Vibrio cholerae* (24). DedA proteins are highly conserved, having 8 homologues in the model bacteria *E. coli* and 6 homologues in *B. subtilis* (25-27). Members of the DedA transporters in *B. subtilis* include UptA and PetA, which are proposed to facilitate trans-bilayer flipping of C_{55}-P and phosphatidylethanolamine, respectively (23, 28). Compared to the DedA family, the DUF368-domain containing proteins, exemplified by PopT, are less conserved, being found in *S. aureus* and *V. cholerae,* but are absent in *B. subtilis* and *E. coli* (29). *V. cholerae* PopT becomes important for cell survival only under alkaline conditions (24). Several Gram-positive and Gram-negative bacteria encode one or multiple UptA-type transporters suggesting that the C_{55}-P recycling process, mediated by these flippases, is broadly conserved across different bacterial species. For example, the single DedA homologue in *Borrelia burgdorferi* is an essential protein required for proper cell division (30). Therefore, the recycling step of peptidoglycan biosynthesis mediated by UptA-type flippases make promising antibiotics target (31). This necessitates a molecular-level understanding of how UptA interacts with its cognate substrate, C_{55}-P, and potentially with membrane phospholipids, to elucidate the mechanism of C_{55}-P translocation.

Herein we employ native mass spectrometry (native MS) to elucidate how *B. subtilis* UptA interacts with multiple lipidic substrates and antibiotics. We show that purified UptA exists as an equilibrium of monomers and dimers. We further show that UptA interacts more favorably with C_{55}-P than C_{55}-PP and membrane phospholipids, and then provide molecular-level evidence on the amino acid residues involved in C_{55}-P binding. We find that UptA binds to its cognate ligand C_{55}-P in a pH-dependent manner and show that lipopeptide antibiotics, such as amphomycin and aspartocin D, bind to UptA with higher affinity than other cell-wall targeting antimicrobial peptides such as bacitracin and vancomycin. Interestingly, amphomycin and its analogues out-compete the flippase UptA for C_{55}-P binding, raising the possibility of a new mode of action and avenue for the development of novel antibiotics to inhibit peptidoglycan biosynthesis.

Results

Monomer-dimer equilibrium of UptA. We expressed *B. subtilis* UptA in *E. coli* and introduced the purified protein into the mass spectrometer from a buffer containing 0.05% LDAO and 200 mM ammonium acetate (pH 8.0). The mass spectrum of UptA displays two charge state distributions assigned to monomers (22963.49±0.70 Da) and dimers (45925.21±0.65 Da) (Fig. 1B). To confirm the non-covalent association of UptA monomers, we isolated the 9+ charge state assigned to the dimer and subjected these ions to high-energy collisional dissociation (HCD, Fig. 1C). The resulting MS^2 spectrum displays new peaks that correspond by mass to UptA protomers, and bear charges complementary to the isolated parent ions (+5 and +4) (Fig. 1D). Of note, the experimental mass for the UptA monomer is higher than the theoretical sequence mass (22936.36 Da) by 28 Da, suggesting an endogenous modification. To identify this unknown modification, we activated the monomer (6+ charge state) using HCD, yielding a plethora of *b*- and *y*-type fragment ions (Fig. S1). The observed fragments are consistent with the formylation of the N-terminal methionine of UptA (Fig. S1). Bacterial inner membrane proteins are normally co-translationally formylated on the N-terminus, however, this modification is rarely observed *in vitro* due to subsequent removal by the cytosolic deformylase (32). Therefore, retainment of N-terminal formylation on UptA suggests periplasmic localisation of the N-terminus of UptA, thus supporting the predicted N-out topology (23). Additionally, we observe UptA in monomeric and dimeric forms in other detergents (Fig. S2), confirming the oligomeric composition of UptA, which might apply to other bacterial DedA homologues (33).

UptA binds C$_{55}$-P with high affinity. We explored the binding affinity of UptA towards C$_{55}$-P by recording mass spectra for solutions containing 2.5 µM UptA and 0 to 40 µM C$_{55}$-P in 200 mM ammonium acetate, 0.05% LDAO at pH 8.0. The mass spectrum for the equimolar mixture of UptA and C$_{55}$-P yielded peaks corresponding to UptA in its apo form and in complex with C$_{55}$-P (Fig. 2A), an indication of high-affinity binding interactions. Further increase in the concentration of C$_{55}$-P enhanced the intensity of the UptA:C$_{55}$-P complex, together with the emergence of additional binding events (Fig. 2A). We deconvoluted the spectra (34) to extract mean relative intensities of bound and unbound forms of UptA in the spectral series. The data reflected an increase in the relative abundance of UptA:(C$_{55}$-P)$_n$ complexes as a function of the concentration of C$_{55}$-P (Fig. 2B). For the first binding event, the relative intensity plateaued at a protein/ligand molar ratio of ~4 (Fig. 2B). By fitting the experimentally observed intensity ratios to the Hill equation, we obtained an apparent K_d = 5.7 µM (95% CI: 5.0 to 6.4 µM) for the binding of one C$_{55}$-P molecule to UptA. This K_d value for UptA/C$_{55}$-P binding is of the same order of magnitude as the binding interactions for MurG/lipid I (K_d = 1.89±0.6 µM) (35) and MurJ/lipid II (K_d = 2.9±0.6 µM) (14), indicating that UptA exhibits a similarly high affinity binding with respect to C$_{55}$-P.

To investigate the binding selectivity of UptA for monophosphate (C_{55}-P) versus diphosphate (C_{55}-PP) forms of the lipid carrier, we performed comparative binding assays. First, we recorded mass spectra for a mixture of UptA with increasing concentrations of C_{55}-PP (Fig. S3). We observed peaks assigned to UptA in ligand-free and C_{55}-PP-bound forms, and the relative intensity of the UptA:C_{55}-PP complex increased with increasing concentration of C_{55}-PP (Fig. 2C). We then fit the data to Hill equation to obtain an apparent K_d for this system. The relative intensities of UptA:C_{55}-PP yielded a significantly higher apparent K_d = 15.3 µM (95% CI: 9.5 to 32.8 µM) than the K_d obtained for UptA:C_{55}-P (K_d = 5.7±0.7 µM). Being within an order of magnitude, these dissociation constants suggest that UptA forms extensive contact with both forms of the lipid carriers, but that the protein interacts more favourably with C_{55}-P than C_{55}-PP. We therefore conclude that UptA is more selective towards C_{55}-P than C_{55}-PP, thus confirming its preferred endogenous substrate.

C_{55}-P binding to UptA is sensitive to pH. UptA is a member of the DedA family of transporters whose cellular functions are potentially driven by proton-motive force and provide conditional fitness to bacteria under alkaline conditions (25, 26, 36). Membrane flippases such as MurJ, which is driven by proton-motive force (37), exhibit pH-dependency in ligand binding (14). We therefore considered the possibility that substrate interactions of UptA might exhibit pH dependency. To this end, we equilibrated C_{55}-P with UptA at a protein/ligand molar ratio of 1:4 in a buffer of the same composition but at pH 8.0 and 5.0. The resulting spectrum at pH 8.0 exhibits peaks consistent with monomeric UptA in an apo form, and in complex with 1 and 2 molecules of C_{55}-P (Fig. S4). We also observed UptA dimer in complex with up to two C_{55}-P molecules, and the relative intensities of C_{55}-P bound to the dimer is higher than to the monomer (Fig. S4). This suggested that dimer interacts more favourably with C_{55}-P than the monomer UptA. However, the peaks assigned to ligand-bound UptA are more intense at pH 8.0 compared to pH 5.0 by a factor of 3. These results imply that more favourable protein-ligand interactions occur under physiologically relevant pH conditions. As a negative control, we tested for pH sensitivity of C_{55}-P binding to MraY, the latter being the downstream enzyme in the peptidoglycan pathway that couples UM5 to C_{55}-P. To this end, we equilibrated MraY and C_{55}-P at a molar ratio of 1:4 in a buffer of pH 8.0 and pH 5.0. In this case, the spectra show no significant difference in the intensity of C_{55}-P bound to MraY at pH 8.0 and at pH 5.0 (Fig. S4). We therefore conclude that environmental pH modulates the interaction of C_{55}-P with UptA, and potentially its transport mechanism.

Amino acid residues mediating C_{55}-P interactions with UptA.
Conserved arginine residues in the putative membrane re-entrant loops of *E. coli* DedA proteins YqjA (R130) and YghB (R136) are important for the cellular role of these proteins (36). The corresponding arginine residues (R112 and R118) in *B. subtilis* UptA have been suggested to impair the cellular function of UptA by making the cells more susceptible to inhibition by amphomycin (23). How these

residues impact substrate interactions, however, is not known. Based on the model structure of UptA predicted by AlphaFold, the R112 residue should engage Q64 while R118 is able to form hydrogen bonds with E32, H119 and W146 to potentially stabilise the membrane re-entrant loops (Fig. 3A). To test for the impact of these residues on C_{55}-P binding, we generated the E32A, Q64A, R112A, R118A, H119A and W146A UptA single mutants. After purification of the proteins, using the same protocol as the wild type (WT) (See Methods and Fig. S5), we equilibrated aliquots of each protein variant (5 µM) with C_{55}-P (10 µM) and recorded spectra under the same instrument settings. The resulting spectra exhibited peaks consistent with C_{55}-P binding in all cases; but only a modest reduction in C_{55}-P binding intensity was observed in the case of the Q64A and H119A mutants compared to the wild type (Fig. S5, Fig. 3B). Compared to the wild type UptA, we observed more significant reduction (27-30%) in C_{55}-P binding with the mutants E32A, W146A, R112A and R118A that participate directly in the predicted hydrogen bonding network (Figure 3A, B), suggesting that these residues are important for C_{55}-P binding. We probe this observation further by investigating the double mutants R112A/R118A and R118A/R119A where the C_{55}-P binding reduced more significantly (43% and 46%, respectively) than in the single mutants. We additionally generated a double mutant R112E/R118E in which the polarity of both native arginine residues in the wild type are reversed. We find that the R112E/R118E UptA exhibited 63% reduction in the intensity of C_{55}-P binding compared to the wild type (Fig. 3B). These observations suggest that impaired hydrogen bonding between R118 and Q64 on the one hand, and R118 and E32 could have also disrupted the C_{55}-P binding sites, highlighting the roles of R112 and R118 in the binding of C_{55}-P to UptA.

Since wild-type UptA displayed monomer-dimer equilibria and the dimer bind to C_{55}-P more intensely than the monomer (cf. Fig. S4), we probed this observation further using the UptA single amino acid mutant E32A since the latter consistently retained a higher dimer population than the wild type (Fig. S6). We prepared and analysed solutions containing 2.5 µM E32A UptA and 10 µM C_{55}-P. The resulting spectra exhibited peaks assigned to UptA monomers and dimers in complex with C_{55}-P (Fig. S6). We find that the relative intensity of ligand-bound dimers is significantly higher than the ligand-bound monomers, and the ligand binding has only a modest impact on the monomer-dimer ratios. Together these results suggest that the ligand C_{55}-P binds to dimeric UptA with a higher affinity than to the monomeric form.

UptA interacts more favourably with C_{55}-P than its shorter-chain analogues. Next, we investigated the chain length selectivity of UptA by testing its interaction with geranylgeranyl phosphate (C_{20}-P), hexaprenyl phosphate (C_{30}-P), and C_{55}-P; the latter representing the native form of the lipid carrier in the model organisms *B. subtilis* and *E. coli* (38). For this experiment, we incubated 4 µM UptA with a 10-fold molar excess of C_{20}-P, C_{30}-P and C_{55}-P at pH 8.0 and recorded spectra under the same conditions. For the UptA/C_{20}-P sample, the spectrum exhibits low-intensity peaks corresponding by mass to UptA:C_{20}-P complexes, leaving most of the protein in the apo form (Fig.

S7). In the case of UptA/C_{30}-P, we observed more intense protein/ligand complexes, indicating a higher binding affinity of UptA to C_{30}-P compared to C_{20}-P (Fig. S7). The spectra recorded for the UptA/C_{55}-P mixture present peaks with the highest intensity of protein/ligand complexes (Fig. S7), indicating a more favourable interaction for C_{55}-P compared to the lipidic substrate with shorter aliphatic chains. The length and stereochemistry of lipid tails can influence their interactions with membrane proteins (39). Our data shows that the lipid carrier with longer hydrophobic tails binds more favourably to UptA than the shorter ones, supporting our hypothesis that C_{55}-P can form extensive contact with UptA to favour its binding interactions. Accordingly, the relative abundance of UptA in complex with C_{55}-P is higher than in the case of UptA-bound C_{30}-P and C_{20}-P (Fig. S7). Together, these data are consistent with C_{55}-P being the preferred endogenous substrate for UptA.

UptA binds more favourably to phosphatidylglycerols than phosphatidylethanolamines. Proteins belonging to the DedA family are a widespread family of transporters and are proposed to flip diverse lipids, including carrier lipids and membrane phospholipids. We therefore probed the interaction of UptA with membrane phospholipids to understand if the latter could modulate its function. For this purpose, we selected anionic phosphatidylglycerols (PG) and zwitterionic phosphatidylethanolamines (PE), representing the major classes of phospholipid headgroups found in *B. subtilis* (40, 41). We recorded mass spectra of solutions containing 4 µM UptA incubated with 10 µM a15:0-i15:0 PE and 10 µM a15:0-i15:0 PG (a15:0, anteiso-pentadecanoic acid; i15:0, iso-pentadecanoic acid). The fatty acid composition of these lipids closely mimics those of native *B. subtilis* lipids (see below). The resulting spectra display peaks corresponding to UptA in apo and lipid-bound forms in both cases (Fig. 4A,B). We observed the formation of 1:1 and 1:2 protein-lipid complexes of UptA with a15:0-i15:0 PG, but only a 1:1 complex in the case of a15:0-i15:0 PE. Accordingly, the mean relative intensity of UptA in complex with a15:0-i15:0 PG is higher than for a15:0-i15:0 PE (Fig. 4C). We therefore hypothesise that UptA interacts more favourably with PG than with PE. We tested additional phospholipids PG and PE with comparable acyl chains. We find that 16:0-18:1 PG and 18:1-18:1 PG bind to UptA more intensely than 16:0-18:1 PE and 18:1-18:1 PE, respectively (Fig. 4C), supporting the hypothesis. Importantly, UptA binds to C_{55}-P more intensely than any of these diacylglycerol phospholipids (Fig. 4C), in line with C_{55}-P being its native substrate.

To verify the lipid binding preference of UptA with more native-like lipids, we prepared lipid extracts from *B. subtilis* membranes (see Methods). We then incubate an aliquot of the resulting crude lipids (Fig. S8) with UptA. The spectra yielded peaks assigned to UptA in complex with cardiolipins (~1326 Da) and a range of phospholipids, predominantly 694 Da, 721 Da, 748 Da and 776 Da species (Fig. 4D, 4E). To identify these lipids, we performed tandem MS/MS, isolating the lipid species and subjecting it to collisional activation. The resulting spectra exhibited fragments at 152.99 Da (Fig. S9), a diagnostic feature of the phosphatidylglycerol headgroup (42). The fatty acid fragments are mixed populations, predominantly 15:0-15:0, 15:0-17:0, 16:0-16:0, and 16:0-18:1 (Fig. S9) which are the

typical fatty acids from *B. subtilis* membrane phospholipids (41). We therefore conclude that UptA interacts more favourably with the anionic cardiolipins and phosphatidylglycerols than with the zwitterionic phosphatidylethanolamines in the membrane lipid extracts.

C_{55}-P outcompetes phospholipids for UptA binding. To investigate whether or not C_{55}-P can outcompete phospholipids in binding to UptA, we incubated solutions containing UptA-DOPG with increasing concentrations of C_{55}-P. Spectra were recorded after aliquots of the solution were equilibrated on ice for 15 mins. The spectrum of UptA/DOPG mixture in the absence of C_{55}-P reflected the binding of 1-2 molecules of DOPG per UptA (Fig. S10). The presence of C_{55}-P in the sample yielded a new peak corresponding to UptA:C_{55}-P complex, and simultaneously caused attenuation of the intensity DOPG-bound UptA in a concentration-dependent manner. At the highest C_{55}-P concentration tested (20 µM), C_{55}-P has displaced most of DOPG from UptA (Fig. S10). We therefore hypothesised that phospholipids should be unable to outcompete C_{55}-P bound to UptA. To test this hypothesis, we incubated UptA with C_{55}-P and then equilibrated aliquots with an increasing concentration of DOPG. Indeed, C_{55}-P remains bound to UptA even in the presence of excess DOPG (20 µM) (Fig. S10), indicating that phospholipids could not significantly displace C_{55}-P from UptA. Analogous experiments with DOPE show that C_{55}-P effectively outcompete DOPE for UptA binding but the latter could not efficiently displace C_{55}-P from UptA (Fig. S10). Together this study shows that UptA interacts more favourably with C_{55}-P than with membrane phospholipids, thereby establishing C_{55}-P as the preferred endogenous substrate.

Lipopeptide antibiotics disrupt UptA:C_{55}-P complex. We next investigate how UptA and its complex with C_{55}-P interacts with a range of cell-wall targeting antibiotics including the lipopeptides daptomycin and amphomycin. Daptomycin is known to kill bacteria by depolarising the membrane (43) and by complexing with lipid II (44). We investigated if daptomycin affects C_{55}-P binding to UptA by recording spectra for solutions containing 2.5 µM UptA, 10 µM C_{55}-P and different concentrations of daptomycin. In the absence of antibiotics, the spectrum exhibited peaks assigned to UptA in apo form, and in complex with 1 and 2 C_{55}-P molecules (Fig. S11). In the presence of daptomycin, we observed additional peaks including those corresponding to the binary complex UptA:daptomycin and the ternary complex UptA:C_{55}-P:daptomycin (Fig. S11). Further increases in daptomycin concentration resulted in higher intensity of these complexes, indicating that daptomycin interacts with UptA but it does not directly compete against C_{55}-P binding to the protein.

Amphomycin and derivatives are known to kill bacteria by forming complexes with free C_{55}-P (45), however, whether or not this involves a direct interaction with a protein cellular target is not known. We therefore investigated how amphomycin affects the UptA:C_{55}-P complex. We recorded spectra for solutions containing 2.5 µM UptA, 10 µM C_{55}-P and then equilibrated aliquots with increasing concentrations of amphomycin. The spectra exhibited peaks consistent with amphomycin binding to

UptA, in addition to the C_{55}-P binding (Fig. 5A). We find that the peak intensities of the UptA:C_{55}-P complex reduce with respect to an increase in the concentration of amphomycin (Fig. 5B). Compared to the case of daptomycin, little-to-no UptA:C_{55}-P complexes remained in the presence of 10-fold molar excess of amphomycin (Fig. 5B), suggesting that amphomycin destabilise the ternary complex to sequester C_{55}-P. We then performed similar experiments with aspartocin D, an analogue of amphomycin. The resulting spectrum shows that aspartocin D similarly binds to UptA and also induces dissociation of UptA:C_{55}-P complex (Fig. 5C).

The amphiphilic nature of daptomycin and amphomycin raises the possibility that these lipopeptide antibiotics may promiscuously bind to UptA in a nonspecific manner. We probe this possibility by recording spectra for UptA:C_{55}-P solutions in the presence of 100 μM polymyxin B and 100 μM polymyxin E. These polymyxins are membrane-disrupting lipopeptides that target the outer membrane lipopolysaccharides in Gram-negative bacteria (46) but without a known affinity for C_{55}-P or UptA. We find that, unlike daptomycin and amphomycins, the polymyxins do not bind to UptA (Fig. S11), indicating that not every lipopeptide can bind to UptA. The ability of daptomycin and amphomycin analogues to directly interact with UptA therefore suggests that these lipopeptides may render a fraction of the cellular pool of UptA inaccessible to C_{55}-P. Notably, polymyxin B and polymyxin E similarly caused attenuation of C_{55}-P binding to UptA (Fig. S10), but this effect is less significant compared to the impact of amphomycin analogues (Fig. 5C). The reduction of C_{55}-P binding to UptA in the presence of polymyxins can be attributed to potential electrostatic interactions between the phosphate group of C_{55}-P molecules and the polymyxins which are cationic at physiological pH (47).

To verify these results, we studied the impact of bacitracin and vancomycin on UptA:C_{55}-P interactions. Bacitracin and vancomycin are peptide antibiotics whose mode of action involves sequestration of C_{55}-PP and lipid II, respectively (48, 49). To this end, we prepared UptA:C_{55}-P mixtures and then equilibrated aliquots with bacitracin and vancomycin. We find that bacitracin and vancomycin bind to UptA but with very low intensities (Fig. S11), indicating weak affinity interactions. In line with the specificity of bacitracin for C_{55}-PP rather than C_{55}-P, this antibiotic modestly competes against C_{55}-P binding to UptA (Fig. S11, Fig. 5C). By contrast, vancomycin has no observable impact on C_{55}-P binding (Fig. S11, Fig. 5C). Thus, the non-lipopeptide antibiotics exhibit weak affinity for UptA and had only a minor impact on UptA:C_{55}-P interactions. Taken together, our results highlight that the lipopeptides such as amphomycin and aspartocin D induce dissociation of UptA:C_{55}-P complexes, and may therefore interfere with peptidoglycan pathway by binding directly to UptA in addition to their known propensity to form complexes with free C_{55}-P molecules (45).

Discussions

In this study, using native MS, we investigated the interplay of multiple lipidic substrates and antibiotics with UptA (summarised in Fig. 6). Specifically, we report that purified UptA exists as an equilibrium of monomers and dimers, suggesting that DedA proteins might form dimers in the native membrane when lateral pressures in the lipid bilayers (50) might prevent their dissociation. In line with the dependency of DedA family proteins on proton motive force (51), we observed less C_{55}-P binding to UptA in the presence of excess H^+ and that the monomer-dimer ratio of UptA is sensitive to pH rather than to C_{55}-P binding. This feature contrasts with those of the upstream peptidoglycan enzymes MraY (35, 52) and UppP (53, 54) whose dimerization are promoted by lipid binding.

Our studies also show that UptA selectively binds to C_{55}-P over C_{55}-PP and that the protein interacts more favourably with C_{55}-P than the shorter chain analogues C_{30}-P and C_{20}-P. This implies that the hydrophobic tail of C_{55}-P makes extensive contact with UptA, promoting the binding interactions and potentially facilitating the transport of carrier lipids. The higher affinity of UptA towards the monophosphate form of the carrier lipid, rather than to the diphosphate form, favours a mechanism in which UptA mediates the transport of C_{55}-P rather than both C_{55}-P and C_{55}-PP. We further show that UptA interacts more favourably with the anionic lipids than the zwitterionic lipids and that these lipids are readily displaced by C_{55}-P. This binding preference agrees with UptA being responsible for C_{55}-P rather than the phospholipid translocation (23). Competition between anionic lipids, which are more abundant in the native lipid bilayer than C_{55}-P, suggests that both lipids bind near the same sites on UptA and that the local lipid environment of UptA can potentially modulate its cellular function.

We provided molecular-level evidence that the conserved arginine (R112 and R118) residues in the putative membrane re-entrant helices of UptA are important for its substrate binding interactions, confirming earlier reports that *B. subtilis* strains bearing these mutations on UptA are more susceptible to inhibition by MX2401, a derivative of amphomycin (23). Despite the combined cellular pool of C_{55}-PP and C_{55}-P being limited to ~1.5×10^5 molecules per cell (55), they mediate the transport of glycan-bearing components of the cell wall peptidoglycan, teichoic acid and capsular polysaccharide (56). Therefore, antibiotics that block C_{55}-P and its flippases are potential drug targets that can potentially inhibit multiple pathways and thus mitigate drug resistance. Accordingly, bacitracin and teixobactin form complex with C_{55}-PP (57, 58) while amphomycin analogues inhibit bacterial cell wall formation by forming complexes with C_{55}-P (23, 45, 59, 60). Herein we show that amphomycin and aspartocin D can additionally bind directly to UptA and induce dissociation of the UptA:C_{55}-P complex, a finding that may guide the development of new antibiotic analogues to inhibit the flippase function of UptA and its homologues.

MATERIALS AND METHODS

Detailed experimental procedures are provided in the SI Appendix. Briefly, UptA (wild type and mutants) was expressed in *E. coli* C43(DE3) cells (Lucigen) with GFP fusion on the C-terminus and solubilised from the membrane fraction by *n*-dodecyl-*β-D*-maltopyranoside (DDM). After purification by affinity chromatography, the GFP was cleaved by TEV protease, removed, and UptA was further purified by size-exclusion chromatography in a buffer containing 20 mM Tris-HCl (pH 8.0), 200 mM NaCl, 0.02% DDM. Before measurements, proteins were buffer-exchanged into 0.05% LDAO, 200 mM ammonium acetate at the desired pH using a centrifugal buffer exchange device (Micro Bio-Spin 6, Bio-Rad). Stock solutions of lipids were prepared from chloroform/methanol solution by evaporating aliquots of known volume in a SpeedVac. After drying, the lipid films were weighed and resuspended to a final concentration of 0.5 mM in 200 mM ammonium acetate and 0.05% LDAO by vertexing. Stock solutions of 0.5 mM amphomycin, aspartocin D, daptomycin, polymyxin B, polymyxin E, and bacitracin were made in the same buffer. All experiments were repeated three times from newly prepared stock solutions. Native mass spectrometry was performed on a Q-Exactive hybrid quadrupole-Orbitrap mass spectrometer (Thermo Fisher Scientific, Bremen, Germany) and lipid identification was performed on an Orbitrap Eclipse Tribrid mass spectrometer (Thermo).

ACKNOWLEDGMENTS

Research in the C.V.R. laboratory is supported by a Wellcome Trust Award (221795/Z/20/Z); and a Medical Research Council (MRC) programme grant (MR/V028839/1) on which A.O.O. is a researcher co-investigator. J.R.B. acknowledges the Royal Society University Research Fellowship grant (URF\R1\211567). A.O.O. is a Junior Research Fellow at St Anne's College, University of Oxford, United Kingdom.

REFERENCES

1.   T. Schneider, H.-G. Sahl, An oldie but a goodie – cell wall biosynthesis as antibiotic target pathway. *Int. J. Med. Microbiol.* **300**, 161-169 (2010).
2.   J. M. C. Kwan, Y. Qiao, Mechanistic Insights into the Activities of Major Families of Enzymes in Bacterial Peptidoglycan Assembly and Breakdown. *ChemBioChem* **24**, e202200693 (2023).
3.   A. Coleman, Y. Vohra, K. Rascati, S. Kubes, B. Moffett, Antibiotic Utilization and Efficacy Associated With Treating Pediatric Urinary Tract Infections in Texas Medicaid Patients in the First Year of Life. *Pediatr. Infect. Dis. J.* **40**, 993-996 (2021).
4.   E. Sauvage, M. Terrak, Glycosyltransferases and Transpeptidases/Penicillin-Binding Proteins: Valuable Targets for New Antibacterials. *Antibiotics* **5**, 12 (2016).
5.   A. J. Meeske *et al.*, MurJ and a novel lipid II flippase are required for cell wall biogenesis in Bacillus subtilis. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 6437-6442 (2015).
6.   A. C. Y. Kuk, A. Hao, S. Y. Lee, Structure and Mechanism of the Lipid Flippase MurJ. *Annu. Rev. Biochem.* **91**, 705-729 (2022).
7.   S. Kumar, A. Mollo, D. Kahne, N. Ruiz, The Bacterial Cell Wall: From Lipid II Flipping to Polymerization. *Chem. Rev.* **122**, 8884-8910 (2022).
8.   M.-C. Lo *et al.*, A New Mechanism of Action Proposed for Ramoplanin. *J. Am. Chem. Soc.* **122**, 3540-3541 (2000).

9.   X. Fang *et al.*, The mechanism of action of ramoplanin and enduracidin. *Mol. BioSyst.* **2**, 69-76 (2006).

10.   H. Barreteau *et al.*, Cytoplasmic steps of peptidoglycan biosynthesis. *FEMS Microbiol. Rev.* **32**, 168-207 (2008).

11.   B. C. Chung *et al.*, Crystal structure of MraY, an essential membrane enzyme for bacterial cell wall synthesis. *Science* **341**, 1012-1016 (2013).

12.   A. Bouhss, A. E. Trunkfield, T. D. Bugg, D. Mengin-Lecreulx, The biosynthesis of peptidoglycan lipid-linked intermediates. *FEMS Microbiol. Rev.* **32**, 208-233 (2007).

13.   S. Ha, D. Walker, Y. Shi, S. Walker, The 1.9 A crystal structure of Escherichia coli MurG, a membrane-associated glycosyltransferase involved in peptidoglycan biosynthesis. *Protein Sci.* **9**, 1045-1052 (2000).

14.   J. R. Bolla *et al.*, Direct observation of the influence of cardiolipin and antibiotics on lipid II binding to MurJ. *Nat. Chem.* **10**, 363-371 (2018).

15.   S. Kumar, A. Mollo, D. Kahne, N. Ruiz, The bacterial cell wall: from lipid II flipping to polymerization. *Chemical Reviews* **122**, 8884-8910 (2022).

16.   K. Bupp, J. van Heijenoort, The final step of peptidoglycan subunit assembly in Escherichia coli occurs in the cytoplasm. *J. Bacteriol.* **175**, 1841-1843 (1993).

17.   A. J. F. Egan, J. Errington, W. Vollmer, Regulation of peptidoglycan synthesis and remodelling. *Nat. Rev. Microbiol.* **18**, 446-460 (2020).

18.   V. M. Hernández-Rocamora *et al.*, Real-time monitoring of peptidoglycan synthesis by membrane-reconstituted penicillin-binding proteins. *eLife* **10**, e61525 (2021).

19.   Y. Park, A. Taguchi, V. Baidin, D. Kahne, S. Walker, A Time-Resolved FRET Assay Identifies a Small Molecule that Inhibits the Essential Bacterial Cell Wall Polymerase FtsW. *Angew. Chem. Int. Ed.* **62**, e202301522 (2023).

20.   M. El Ghachi, A. Bouhss, D. Blanot, D. Mengin-Lecreulx, The bacA gene of Escherichia coli encodes an undecaprenyl pyrophosphate phosphatase activity. *J. Biol. Chem.* **279**, 30106-30113 (2004).

21.   M. El Ghachi, A. Derbise, A. Bouhss, D. Mengin-Lecreulx, Identification of multiple genes encoding membrane proteins with undecaprenyl pyrophosphate phosphatase (UppP) activity in Escherichia coli. *J. Biol. Chem.* **280**, 18689-18695 (2005).

22.   R. Bernard, M. El Ghachi, D. Mengin-Lecreulx, M. Chippaux, F. Denizot, BcrC from Bacillus subtilis acts as an undecaprenyl pyrophosphate phosphatase in bacitracin resistance. *J. Biol. Chem.* **280**, 28852-28857 (2005).

23.   I. J. Roney, D. Z. Rudner, Two broadly conserved families of polyprenyl-phosphate transporters. *Nature* **613**, 729-734 (2023).

24.   B. Sit *et al.*, Undecaprenyl phosphate translocases confer conditional microbial fitness. *Nature* **613**, 721-728 (2023).

25.   W. T. Doerrler, R. Sikdar, S. Kumar, L. A. Boughner, New Functions for the Ancient DedA Membrane Protein Family. *J. Bacteriol.* **195**, 3-11 (2013).

26.   P. R. Panta, W. T. Doerrler, A Burkholderia thailandensis DedA Family Membrane Protein Is Required for Proton Motive Force Dependent Lipid A Modification. *Front. Microbiol.* **11** (2021).

27.   S. Kumar, W. T. Doerrler, Members of the conserved DedA family are likely membrane transporters and are required for drug resistance in Escherichia coli. *Antimicrob. Agents Chemother.* **58**, 923-930 (2014).

28. I. J. Roney, D. Z. Rudner, The DedA superfamily member PetA is required for the transbilayer distribution of phosphatidylethanolamine in bacterial membranes. *Proc. Nat. Acad. Sci. U. S. A.* **120**, e2301979120 (2023).

29. A. Galinier, C. Delan-Forino, E. Foulquier, H. Lakhal, F. Pompeo, Recent Advances in Peptidoglycan Synthesis and Regulation in Bacteria. *Biomolecules* **13**, 720 (2023).

30. F. T. Liang *et al.*, BB0250 of Borrelia burgdorferi is a conserved and essential inner membrane protein required for cell division. *J. Bacteriol.* **192**, 6105-6115 (2010).

31. M. Arai, L. Liu, T. Fujimoto, A. Setiawan, M. Kobayashi, DedA protein relates to action-mechanism of halicyclamine A, a marine spongean macrocyclic alkaloid, as an anti-dormant mycobacterial substance. *Mar Drugs* **9**, 984-993 (2011).

32. A. Ranjan, E. Mercier, A. Bhatt, W. Wintermeyer, Signal recognition particle prevents N-terminal processing of bacterial membrane proteins. *Nat. Commun.* **8**, 15562 (2017).

33. H. L. Scarsbrook, R. Urban, B. R. Streather, A. Moores, C. Mulligan, Topological analysis of a bacterial DedA protein associated with alkaline tolerance and antimicrobial resistance. *Microbiology (Reading)* **167** (2021).

34. M. T. Marty *et al.*, Bayesian deconvolution of mass and ion mobility spectra: from binary interactions to polydisperse ensembles. *Anal. Chem.* **87**, 4370-4376 (2015).

35. A. O. Oluwole *et al.*, Peptidoglycan biosynthesis is driven by lipid transfer along enzyme-substrate affinity gradients. *Nat. Commun.* **13**, 2278 (2022).

36. S. Kumar, C. L. Bradley, P. Mukashyaka, W. T. Doerrler, Identification of essential arginine residues of Escherichia coli DedA/Tvp38 family membrane proteins YqjA and YghB. *FEMS Microbiology Letters* **363** (2016).

37. F. A. Rubino, S. Kumar, N. Ruiz, S. Walker, D. E. Kahne, Membrane Potential Is Required for MurJ Function. *J. Am. Chem. Soc.* **140**, 4481-4484 (2018).

38. G. Manat *et al.*, Deciphering the Metabolism of Undecaprenyl-Phosphate: The Bacterial Cell-Wall Unit Carrier at the Membrane Frontier. *Microbial drug resistance (Larchmont, N.Y.)* **20** (2014).

39. X. Cong, Y. Liu, W. Liu, X. Liang, A. Laganowsky, Allosteric modulation of protein-protein interactions by individual lipid binding events. *Nat Commun* **8**, 2203 (2017).

40. J. A. den Kamp, I. Redai, L. L. van Deenen, Phospholipid composition of Bacillus subtilis. *J. Bacteriol.* **99**, 298-303 (1969).

41. J. D. Nickels *et al.*, Bacillus subtilis Lipid Extract, A Branched-Chain Fatty Acid Model Membrane. *The Journal of Physical Chemistry Letters* **8**, 4214-4217 (2017).

42. J. Pi, X. Wu, Y. Feng, Fragmentation patterns of five types of phospholipids by ultra-high-performance liquid chromatography electrospray ionization quadrupole time-of-flight tandem mass spectrometry. *Analytical methods* **8**, 1319-1332 (2016).

43. W. R. Miller, A. S. Bayer, C. A. Arias, Mechanism of action and resistance to daptomycin in Staphylococcus aureus and enterococci. *Cold Spring Harb. Perspect. Med.* **6** (2016).

44. F. Grein *et al.*, Ca2+-Daptomycin targets cell wall biosynthesis by forming a tripartite complex with undecaprenyl-coupled intermediates and membrane lipids. *Nat. Commun.* **11**, 1455 (2020).

45. T. M. Wood *et al.*, Mechanistic insights into the C 55-P targeting lipopeptide antibiotics revealed by structure–activity studies and high-resolution crystal structures. *Chem. Sci.* **13**, 2985-2991 (2022).

46. S. C. Nang, M. A. Azad, T. Velkov, Q. T. Zhou, J. Li, Rescuing the last-line polymyxins: achievements and challenges. *Pharmacological reviews* **73**, 679-728 (2021).

47. C. J. Slingerland, I. Kotsogianni, C. M. J. Wesseling, N. I. Martin, Polymyxin Stereochemistry and Its Role in Antibacterial Activity and Outer Membrane Disruption. *ACS Infect Dis* **8**, 2396-2404 (2022).

48. J. Smith, E. Weinberg, Mechanisms of antibacterial action of bacitracin. *Microbiol.* **28**, 559-569 (1962).

49. W. P. Hammes, F. C. Neuhaus, On the mechanism of action of vancomycin: inhibition of peptidoglycan synthesis in Gaffkya homari. *Antimicrob. Agents Chemother.* **6**, 722-728 (1974).

50. R. S. Cantor, Lateral pressures in cell membranes: a mechanism for modulation of protein function. *J. Phys. Chem. B* **101**, 1723-1725 (1997).

51. P. R. Panta, W. T. Doerrler, A Burkholderia thailandensis DedA family membrane protein is required for proton motive force dependent lipid a modification. *Frontiers in Microbiology* **11**, 618389 (2021).

52. E. Henrich *et al.*, Lipid Requirements for the Enzymatic Activity of MraY Translocases and in Vitro Reconstitution of the Lipid II Synthesis Pathway. *J. Biol. Chem.* **291**, 2535-2546 (2016).

53. S. D. Workman, L. J. Worrall, N. C. J. Strynadka, Crystal structure of an intramembranal phosphatase central to bacterial cell-wall peptidoglycan biosynthesis and lipid recycling. *Nat. Commun.* **9**, 1159 (2018).

54. A. O. Oluwole *et al.*, Real-time biosynthetic reaction monitoring informs the mechanism of action of antibiotics. *J. Am. Chem. Soc.* **146**, 7007-7017 (2024).

55. H. Barreteau *et al.*, Quantitative high-performance liquid chromatography analysis of the pool levels of undecaprenyl phosphate and its derivatives in bacterial membranes. *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.* **877**, 213-220 (2009).

56. M. Rausch *et al.*, Coordination of capsule assembly and cell wall biosynthesis in Staphylococcus aureus. *Nat. Commun.* **10**, 1404 (2019).

57. L. L. Ling *et al.*, A new antibiotic kills pathogens without detectable resistance. *Nature* **517**, 455-459 (2015).

58. R. Shukla *et al.*, Teixobactin kills bacteria by a two-pronged attack on the cell envelope. *Nature* **608**, 390-396 (2022).

59. T. Schneider *et al.*, The lipopeptide antibiotic Friulimicin B inhibits cell wall biosynthesis through complex formation with bactoprenol phosphate. *Antimicrob. Agents Chemother.* **53**, 1610-1618 (2009).

60. Z. Wang, B. Koirala, Y. Hernandez, M. Zimmerman, S. F. Brady, Bioinformatic prospecting and synthesis of a bifunctional lipopeptide antibiotic that evades resistance. *Science* **376**, 991-996 (2022).

61. J. Jumper *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-589 (2021).

62. N. Ferruz, S. Schmidt, B. Höcker, ProteinTools: a toolkit to analyze protein structures. *Nucleic Acids Res.* **49**, W559-w566 (2021).

Fig. 1: UptA is primarily monomeric with a low population of dimer. A) Schematic depiction of the lipid II cycle of peptidoglycan biosynthesis. C_{55}-P, Undecaprenyl monophosphate; C_{55}-PP, Undecaprenyl diphosphate; UM5, UDP-*N*-acetylmuramyl-pentapeptide; PBPs, penicillin-binding proteins. **B**) Native mass spectrum of UptA (2.2 µM) liberated from a buffer containing 200 mM ammonium acetate (pH 8.0), 0.05% LDAO. Peaks are assigned to UptA in monomeric and dimeric forms. Insert, the model structure of UptA (AFO31823-F1) predicted by AlphaFold (61). Predicted N- and C- termini are indicated. **C**) Quadrupole isolation of the UptA dimer (9+ charge state). **D**) MS/MS of the isolated UptA dimer upon collisional activation (190 V). The dimer dissociates into individual UptA protomers confirming their non-covalent association.

Fig 2. UptA binds C$_{55}$-P with a high affinity. A) Spectra for UptA incubated with different concentrations of C$_{55}$-P in a buffer containing 200 mM ammonium acetate (pH 8), 0.05% LDAO. **B-C**) The relative abundance of UptA bound to one (1x) or two (2x) molecules of C$_{55}$-P (panel B) and C$_{55}$-PP (panel C). Inserts, chemical structures of C$_{55}$-P and C$_{55}$-PP. Each data point represents an average of 3 replicate measurements, and the curves represent best fits of the data to Hill's equation (see methods), error bars are standard deviations.

Fig. 3. Probing the amino acid residues involved in UptA-C$_{55}$-P binding. A. Model UptA structure predicted by AlphaFold (AFO31823-F1) (61) and a zoomed view showing hydrogen bonding networks surrounding R118 and R112. R112--Q64: 3.1 Å; R118--E32: 2.7 Å; and E32--W146: 3.5 Å. Hydrogen bonding distances were computed using ProteinTools(62) and the image was processed in PyMOL. **B.** Relative intensity of C$_{55}$-P bound to UptA wild type and mutants. Bars represent the mean of three biological replicates with each data point shown in black, error bars are standard deviations. The double mutant at residues R112 and R118 caused the most significant reduction in C$_{55}$-P binding. $*p$ = 0.02 to 0.04, $**p$ = 0.004 to 0.0012, $****p$ = 0.0002, $****p < 0.0001$ from unpaired two-tailed t-test by comparing mutants with wild type, n = 3.

Fig 4. UptA binds **phosphatidylglycerols**. A) Native mass spectrum for UptA equilibrated with a15:0-i15:0 PE and a15:0-i15:0 PG. UptA interacts more favourably with PG than PE. **B)** Chemical structure of a15:0-i15:0 PE and a15:0-i15:0 PG. **C)** Relative intensity of UptA bound to phospholipids and C_{55}-P. UptA interacts more favourably with C_{55}-P than with the phospholipids. Bar represents the mean of three independent replicates shown as data points and the error bars are standard deviations. *p*-values are from two-tailed t-tests with n=3, ***p=0.0003 to 0.0005, ****p<0.0001. D. Native mass spectrum of 4 μM equilibrated with ~0.1 mg/mL total lipid extracts of *B. subtilis.* Peaks labelled # is a 43.3 kDa contaminant. **E**. Zoomed view of a change state (6+), showing UptA adducts with phosphatidylglycerols.

Fig. 5 UptA function can be inhibited by lipopeptide antibiotics. A) Mass spectra (6+ charge state) for solutions containing 2.5 µM UptA and 10 µM C_{55}-P incubated with 0-100 µM amphomycin. **B)** Relative intensity of C_{55}-P bound to UptA in the presence of amphomycin and daptomycin. Bar represents the mean of three independent replicates shown as data points and error bars are standard deviations. Amphomycin destabilises the UptA:C_{55}-P complex whereas daptomycin has a modest impact. **C)** Fraction of C_{55}-P bound to UptA in the presence of cell-wall targeting antibiotics. Each sample contains 2.5 µM UptA, 10 µM C_{55}-P and 100 µM of indicated antibiotic, and spectra were acquired using the same instrument settings and collisional activation of 75 V. Bar represents the mean of three independent replicates shown as data points, and error bar represents standard deviations. P-values from two-tailed t-tests with n=3; **p=0.001 to 0.006; ****p<0.0001, ns, not significant.

Fig. 6 Insights from native MS into the interaction of UptA with lipidic substrates and antibiotics. (i) UptA exhibits monomer-dimer equilibria, with more dimer at pH 8.0 than pH 5.0. (ii) UptA interacts with C_{55}-P more favourably than C_{55}-PP, C_{30}-P, and C_{20}-P. (iii) Amphomycin destabilises the UptA:C_{55}-P complex and sequesters C_{55}-P, thus effectively out-competing the flippase for its carrier lipid substrate. (iv) UptA binds phospholipids, with a higher preference for the anionic than zwitterionic phospholipids. (v) UptA interacts with UptA more favourably than phospholipids, highlighting its preferred lipid substrate. (iv) Based on the proton-dependency of other DedA proteins (52) and the pH-dependency of UptA – C_{55}-P interactions, we propose that the binding of H^+ to UptA enables translocation of C_{55}-P and its subsequent release for the next round of PG precursor synthesis.

Supporting Information

Methods

Chemicals

C_{55}-PP, C_{55}-P, C_{30}-P and C_{20}-P were purchased from Larodan. Phospholipids – a15:0-i15:0 PE, a15:0-i15:0 PG, 16:0-18:1 PE, 16:0-18:1 PG, 18:1-18:1 PE, 18:1-18:1 PG – were from Avanti Polar Lipids. Amphomycin and aspartocin D were purchased from Cayman; while Daptomycin and bacitracin were purchased from Merck. *n*-dodecyl-*β*-D-maltoside (DDM), *n*-decyl-*β*-D-maltoside (DDM), n-octylglucoside (OG) and *n*-dodecyl-*N,N*-dimethylamine-*N*-oxide (LDAO) were from Anatrace.

Plasmids

A codon-optimised version of *Bacillus subtilis* UptA (yngC) (Integrated DNA Technologies) was inserted into a modified pET15b vector between NdeI and NheI endonuclease restriction sites by In-Fusion HD cloning kit (Takara Bio). The resulting plasmid encodes UptA fused to GFP-His$_6$ on the C-terminus via a TEV-protease cleavable site. Plasmids for UptA mutants were generated by site-directed mutagenesis with CloneAmp HiFi PCR Premix (Takara Bio) using pairs of oligonucleotides listed in Table S1. The plasmid pET26-MBP-MraYaa bearing HRV3C protease cleavage site between the MBP and the wild-type MraY was a kind gift from S-Y Lee (Addgene plasmid #100166) (1).

Table S1: Oligonucleotides used to generate plasmids for the expression of UptA mutants.

Primers	Sequence
E32AF	TCCAAGCGCAATTGTATTGGCCTACGGTGG
E32AR	ACAATTGCGCTTGGAATAATTTCAATCATCAGT
Q64AF	GATCGCCGCAATTTTCATTTACTGGATTGGCCGT
Q64AR	AAAATTGCGGCGATCGTTCCCCCGATA
R112AF	TTCTGCTGCCTTCATCCCTGTAGTCCGTCATGC
R112AR	ATGAAGGCAGCAGAAAACACAACCCCCG
R118AF	TGTAGTCGCACATGCTATTTCGATCCCAGCCG
W146AF	CATTCCTGCCTCCATCCTGTTCGTTTACCTGG
W146AR	ATGGAGGCAGGAATGATGGCAAGTACTGTC
R112A/R118A_Fwd	TTCTGCTGCCTTCATCCCTGTAGTCGCACATGCTATTTCGATCCCAGCCG
R112A/R118A_Rev	GCATGTGCGACTACAGGGATGAAGGCAGCAGAAAACACAACCCCCG
R112E/R118E_Fwd	ATTCATCCCTGTAGTCGAACATGCTATTTCGATCCCAGCCG
R112E/R118E_Rev	ACTACAGGGATGAATTCAGCAGAAAACACAACCCCCG

Expression and purification of UptA

Plasmids were used to transform chemically competent *E. coli* C43(DE3) cells (Lucigen) and selected on LB/Agar containing 100 µg/ml ampicillin at 37 °C. Single colonies were used to inoculate 60 mL LB containing 100 µg/ml ampicillin and were grown at 37 °C overnight for 16 h. For large-scale protein expression, 10 mL of the overnight culture was aseptically transferred into 1 L of LB/ampicillin and cells were allowed to grow at 37°C until the $OD_{600\,nm}$ reached 0.6-0.8 after which the temperature was lowered to 18 °C. Expression of UptA-GFP was induced by adding isopropyl-β-D-thiogalactopyranoside (IPTG) at a final concentration of 0.5 mM UptA-GFP and shaking for 20 h at 18 °C. Cells were harvested by centrifugation at 5000 × *g* and lysed immediately or stored at -80 ° C until required.

Cells were thawed on ice and resuspended in a buffer containing 50 mM Tris-HCl (pH 7.4), 200 mM NaCl, 10 µg/mL deoxyribonuclease I (DNAse I), 5 mM $MgCl_2$, 5 mM β-mercaptoethanol and EDTA-free protease inhibitor cocktail tablets. The cells were then disrupted with a microfluidiser (Microfluidics) operated at 20,000 psi. Non-lysed cells and debris were pelleted by centrifugation at 20,000 × *g* for 20 min and the clarified lysate was retained. The membranes fraction was then pelleted by ultracentrifugation at 100,000 x g for 1 h. They were resuspended in a buffer containing 20 mM Tris-HCl (pH 8.0), 300 mM NaCl, 20% glycerol. Membrane aliquots were either solubilised immediately or flash-frozen in liquid nitrogen and stored at -80°C. The membrane was solubilised by incubation with DDM at a final concentration of 1% (w/v) for 60 min. Non-solubilised aggregates were removed by centrifugation at 20,000×g for 20 min and the supernatant was loaded onto a 5-ml HisTrap HP column preequilibrated in buffer A (20 mM Tris-HCl (pH 8.0), 200 mM NaCl, 10% glycerol, 0.03% DDM and 20 mM imidazole). After binding of the His-tagged protein, the column was washed with 100 mL of buffer A, and then with 50 mL of buffer B (20 mM Tris-HCl (pH 8.0), 200 mM NaCl, 10% glycerol, 0.03% DDM and 50 mM imidazole). UptA-GFP was eluted from the column with buffer C (20 mM Tris-HCl (pH 8.0), 200 mM NaCl, 10% glycerol, 0.02% DDM and 300 mM imidazole). To remove imidazole and to cleave the GFP fusion, eluted protein was mixed with TEV protease and dialysed against buffer D (20 mM Tris-HCl (pH 8.0), 150 mM NaCl, 10% glycerol, 0.02% DDM) at 4°C for 16 h.

The digestion mixture was subsequently incubated with 2-mL bed volume of nickel nitriloacetic acid (NTA) agarose resin to remove His-tagged GFP and TEV protease. Tag-free UptA was collected as the flowthrough, concentrated using an Amicon centrifugal concentrator with nominal molecular weight cut-off 50 kDa. The protein was finally purified by size-exclusion chromatography (SEC) using a 24-mL Superdex 200 Increase column. The SEC buffer was 20 mM Tris-HCl (pH 8.0), 200 mM NaCl, 0.02% DDM. UptA was concentrated to ~200 uM, aliquoted, flash-frozen flash in liquid nitrogen, and stored at -80 °C until use. Concentration was measured using a Biomate UV spectrophotometer at 280 nm using extinction coefficients calculated from their predicted amino acid sequences. UptA mutants were purified in 3 biological replicates, using the same procedures as for the wild type.

Expression and purification of MraY.

MraY from *Aquifex aeolicus* was purified as previously described (3). Briefly, the plasmid pET26-HisMBP-MraYaa was used to transform chemically competent *E. coli* C41(DE3) cells (Lucigen) and selected on LB/Agar containing 50 µg/ml kanamycin at 37 °C. Cells were grown in Terrific Broth media supplemented with 50 µg/ml kanamycin at 37°C until the $OD_{600\,nm}$ reached 0.8. Protein expression was induced by adding IPTG at a final concentration of 1 mM for 4 h at 37 °C. Cells were harvested by centrifugation at 5000 × *g* and lysed and solubilized by incubation with 1% DDM for 1 h. His-tagged MBP-MraY was purified by affinity chromatography and digested by HRV3C protease. MraY was isolated by SEC into a buffer containing 20 mM Tris-HCl (pH 8.0), 200 mM NaCl, 0.02% DDM.

Denaturing SDS-PAGE

Denaturing SDS-PAGE was performed using a precast NuPAGE™ Novex 12% Bis-Tris Mini protein gel (Invitrogen) according to the manufacturer's specifications. 30 µL of 0.3 µg/µL protein were mixed with 10 µL of 4X LDS sample loading buffer (Invitrogen) and heated at 80 °C for 5 min. 20 µL of the resulting samples were then loaded into in each well. 5 µL PageRuler™ prestained protein ladder (Thermo Scientific) was loaded into a well as molecular weight marker. Electrophoresis was performed at room temperature for 36 min using a constant voltage (200 V) in 1X solution of NuPAGE™ MES SDS running buffer (Invitrogen). The gel was stained with Quick Coomassie stain solution (Neo Biotech) for 60 min and rinsed with water for 24 h before imaging on iBright™ FL1500 imaging system (Thermo).

Lipid extracts from *B. Subtilis*

B. subtilis strain 168 (ATCC 23857) was cultivated in 200 mM LB media at 37 °C until $OD_{600\,nm}$ 1.2 under aurobic conditions. Cells were harvested by centrifugation at 4,000 *g* for 20 minutes and resuspended in 20 mM Tris-HCl (pH 8.0), 150 mM NaCl. Cells were lysed by 5-6 passes through a microfluidizer at 20,000 psi. The lysates were clarified by centrifugation at 4,000 *g* for 10 minutes, and the totoal membrane was pelleted at 120,000 *g*

for 1 h. The membrane pellet was ressupended in 20% sucrose, 50 mM Tris-HCl (pH 8.0), and loaded as the topmost layer of a 4-layer sucrose gradient: 20% (4 mL), 36% (3 mL), 51% (8 mL) and 55% (2 mL) (4). After centrifugation at 120,000 g for 20 hrs, the cytosolic membrane was collected at the interface of 36% and 45% sucrose layers, and washed 3 times with 1 M ammonium acetate by dilution-centrigugation at 120,000 g for 30 min. The membrane pellet was homogenised in 2 mL of 1 M ammonicum acetate and stored in -80 °C until use. Lipids were extracted by homogenising 0.8 mL of membrane with 1 mL of 1:1 vol/vol) chloroform-methanol mixture in a glass vial followed by vortexing. To separate the aqueous and the organic pahses, adding 1 mL chloroform was added, and the mixture was centrifuged at 3,000 g for 5 min. The chloroform layer containing the total lipids was recovered into a clean glass vial, and the organic solvent was evaporated in a SpeedVac for 16 h. Lipid film was resuspended 200 mM ammonium acetate, 0.05% LDAO (pH 8.0) to ~0.5 mg/mL for the binding experiments.

Sample preparation for native MS
Before measurements, the protein was buffer-exchanged into 0.05% LDAO, 200 mM ammonium acetate ("MS buffer") at the desired pH using a centrifugal buffer exchange device (Micro Bio-Spin 6, Bio-Rad). Stock solutions of lipids were prepared from chloroform/methanol solution by evaporating aliquots of known volume in a SpeedVac. After drying, the lipid films were weighed and resuspended to a final concentration of 0.5 mM in 200 mM ammonium acetate (pH8) and 0.05% LDAO by vertexing. Amphomycin and aspartocin D were purchased from Cayman while Daptomycin and bacitracin were purchased from Merck. Stock solutions of 0.5 mM amphomycin, aspartocin D, daptomycin and bacitracin, were made in the same buffer. All experiments were repeated three times from newly prepared stock solutions.

Native mass spectrometry
About 3 μL protein aliquot was transferred into a gold-coated borosilicate capillary (Harvard Apparatus) and was mounted on the nano ESI source of a Q-Exactive hybrid quadrupole-Orbitrap mass spectrometer (Thermo Fisher Scientific, Bremen, Germany). The instrument settings were 1.2 kV capillary voltage, S-lens RF 200%, argon UHV pressure 3.3×10^{-10} mbar, the capillary temperature was set to 200 °C, and resolution of the instrument was set to 17,500 at a transient time of 64 ms. Voltages of the ion transfer optics –injection flatapole, inter-flatapole lens, bent flatapole, and transfer multipole were set to 5, 3, 2, and 30 V respectively. The noise level was set at 3. Unless otherwise stated, proteins were activated by applying 75 V in the high-energy collisional dissociation cell without in-source trapping. For the MS^2 spectra, parent ions were isolated using an in-source trapping voltage of -200 V, and then fragmented by applying 150-200 V in the high-energy collision-induced dissociation (HCD) cell. Lipids bound to UptA from the total lipid extracts were identified on an Orbitrap Eclipse Tribrid mass spectrometer (Thermo). Peaks corresponding to the parent ions in the negative ESI spectra were isolated within ±1.0 Th and then activated with HCD voltage 10-30%. To identify endogenous modifications using native top-down mass spectrometry, the 6+ charge state of monomeric UptA was isolated (3828 ± 5 Th) and activated using HCD (200 V) with reduced trapping pressures. The resulting fragments were detected in the Orbitrap (R = 280,000 @ m/z 200), averaging ~500 transients to maximise the signal-to-noise ratio. Theoretical fragments from N-terminally formylated UptA were matched to the MS^2 spectrum (3 ppm) using TDValidator (Proteinaceous) and manual validation. Data were visualised and exported for processing using the Qual browser of Xcalibur 4.1.31.9 (Thermo Scientific).
Spectral deconvolution was performed using UniDec 6.0.2 (2). Relative binding affinities were obtained from deconvoluted spectra by dividing the intensity of ligand-bound protein peaks by the sum of the intensities of ligand-bound and ligand-free protein peaks. The mean and standard deviation of these fractional binding intensities from three independent experiments were plotted against the concentrations of the ligand C_{55}-P or C_{55}-PP. To obtain the apparent dissociation constant K_d, the mean relative intensities at different concentrations were fitted globally using GraphPad Prism 10.2.0 to Eq. (1):

$$y = \frac{B_{max} x^h}{x^h + K_d{}^h} \tag{1}$$

Where y is the mean relative intensity and x is the corresponding ligand concentration. B_{max} is the maximum specific binding, K_d is the apparent dissociation constant, and h is the Hill coefficient.

Supplementary Figures

Fig. S1. UptA is formylated at the N-terminus. A. HCD MS/MS spectrum of UptA (6+). The eight most abundant assigned ions are labelled. **B**. Sequence map of the UptA construct used in this study. Terminal *b*- and *y*-type fragment ions observed in the MS/MS spectrum are indicated. **C.** Expanded view of the peak corresponding to b$_{80}$$^{2+}$ ion**.** The theoretical isotopic envelope is overlaid.

Fig. S2. Oligomeric state of UptA in different detergents. Shown are the spectra recorded for UptA in buffer containing 200 mM Ammonium acetate (pH 8.0) supplemented with indicated detergents. The instrument was tuned differently in each case to maximise the transmission of dimeric species.

Fig. S3. UptA:C$_{55}$-PP binding interactions. Spectra for UptA incubated with different concentrations of C$_{55}$-PP in a buffer containing 200 mM ammonium acetate (pH 8), 0.05% LDAO. UptA bind C$_{55}$-PP but with lesser intensities compared to C$_{55}$-P (cf. Fig. 2A).

Fig. S4. pH dependency of C_{55}-P binding interactions with UptA and MraY. A,B. Spectra for UptA/C_{55}-P and MraY/C_{55}-P mixtures at pH 8.0 and pH 5.0**.** Peaks corresponding to the dimeric form of UptA is observed mostly in complex with C_{55}-P. **C.** Relative intensity of UptA:C_{55}-P at pH 8.0 and pH 5.0 as a function of concentrations of C_{55}-P in a buffer containing 200 mM ammonium acetate,0.05% LDAO. Data point is an average of 3 independent replicates, and the error bars are standard deviations. **D.** Relative intensity of C_{55}-P (10 μM) bound to UptA (2.5 μM) and MraY (2.5 μM) at different pH. Unlike UptA, there is no pH-dependency in C_{55}-P binding by MraY. Bar represents the mean of 3 replicate measurements shown as data points, and the error bars are standard deviations.

Fig. S5. Native MS analysis of C$_{55}$-P binding to UptA mutants. **A.** SDS-PAGE image of UptA wild-type (WT) and mutants. M, PageRuler^{TM} prestained protein ladder. UptA is ~22 kDa but migrate on SDS-PAGE as ~17 kDa protein. **B.** Representative spectra for UptA wild type and mutants (5 µM) incubated with C$_{55}$-P (10 µM). Proteins were liberated from a buffer containing 0.05% LDAO and 200 mM ammonium acetate (pH 8.02) using a collisional activation of 100 V.

Fig. S6. Dimer fraction of UptA binds C$_{55}$-P more than the monomeric form. **A.** Spectrum of UptA E32A mutant display peaks corresponding to monomers and dimers. **B.** Relative intensity of dimer populations in spectra for the wild-type UptA and the E32A mutant**.** In both cases, the dimer fraction was higher at pH 8.0 than at pH 5.0. **C.** Spectrum of UptA E32A mutant equilibrated with 10 μM C$_{55}$-P. **D.** Corresponding intensities of apo and C$_{55}$-P bound UptA. Dimeric form of UptA bind to the ligand C$_{55}$-P more significantly than the monomeric form. **E.** The relative proportion of UptA monomers and dimers in a mixture of 2.5 μM E32A UptA and 10 μM C$_{55}$-P at pH 8.0. The ligand C$_{55}$-P caused a modest increase in the dimer fraction.

Fig. S7. UptA binds carrier lipids of longer chain lengths more intensely than those of shorter chain, and phospholipids. A. Spectra for 4 μM UptA equilibrated with 40 μM undecaprenyl phosphate, C_{55}-P hexaprenyl phosphate, C_{30}-P and geranylgeranyl phosphate, C_{20}-P. Samples were prepared in a buffer containing 200 MM ammonium acetate (pH8.0) and 0.05% LDAO. **B.** Relative intensity of ligand-free and ligand-bound UptA. UptA binds to C_{55}-P more intensely than the shorter chain analogues.

Fig. S8. Mass spectrum of total lipid extract from B. subtilis membrane. Lipid extracts from B. subtilis membrane was dissolved in 0.2 M ammonium acetate, 0.05% LDAO (pH8.0) and analysed in the positive ESI mode using source activation of 25 V. Main phospholipids (PE, blue; PG, green) are highlighted in the expanded view.

Fig. S9. MS/MS spectra (negative ESI mode) for lipid species interacting with UptA from the total lipid extracts of *B. subtilis*. Peaks corresponding by mass to each of the parent ions were isolated with a source activation of 150 V, and then fragmented with 15-25 V in the HCD cell. All species carry a single negative charged.

Fig. S10. C55-P outcompetes phospholipids for UptA binding. A,B) Deconvoluted native mass spectra for 4 µM UptA and10 µM 18:1-18:1 PG (DOPG, panel A) or 18:1-18:1 PE (DOPE, panel B) in the presence of increasing concentrations of C_{55}-P. The intensities of peaks assigned to lipid bound to UptA attenuates upon titration with C_{55}-P. **C,D**) Spectra for 4 µM UptA and 10 µM C_{55}-P, then incubated with increasing

concentrations of DOPG (Panel B) or DOPE (panel (D). The peaks assigned to UptA-bound C_{55}-P remained in the presence of lipids, indicating that C_{55}-P binds to UptA more favourable than the phospholipids.

Fig. S11. Impact of cell-wall antibiotics on UptA:C$_{55}$-P interactions. **A.** Spectra (6+ charge state) of 2.5 µM UptA and 10 µM C$_{55}$-P in the presence of different daptomycin concentrations. Unannotated peaks correspond to multiple daptomycin binding events in the 5+ charge state. **B**. Spectra (6+ charge state) of UptA

and C_{55}-P in the presence of different antibiotics and acquired using the same instrument settings and collisional activation of 75 V. Relative intensity of UptA-bound C_{55}-P in the spectra shown in Figure 5. **C.** Chemical structures of lipopeptide antibiotics used in this study.

SI References

1. B. C. Chung *et al.*, Crystal structure of MraY, an essential membrane enzyme for bacterial cell wall synthesis. *Science* **341**, 1012-1016 (2013).
2. A. O. Oluwole *et al.*, Peptidoglycan biosynthesis is driven by lipid transfer along enzyme-substrate affinity gradients. *Nat. Commun.* **13**, 2278 (2022).
3. M. T. Marty *et al.*, Bayesian deconvolution of mass and ion mobility spectra: from binary interactions to polydisperse ensembles. *Anal. Chem.* **87**, 4370-4376 (2015).
4. L. A. Baker *et al.*, Magic-angle-spinning solid-state NMR of membrane proteins. *Methods Enzymol.* **557**, 307-328 (2015).