Lipopeptide antibiotics disrupt interactions of undecaprenyl phosphate with UptA

Abraham O. Oluwole[1,2*], Neha Kalmankar[1,2], Michela Guida[2,3], Jack L. Bennett[1,2], Giovanna Poce[3], Jani R. Bolla[2,4] Carol V. Robinson[1,2*]

[1]Department of Chemistry, University of Oxford, South Parks Road, Oxford, OX1 3QZ, UK. [2]The Kavli Institute for Nanoscience Discovery, Oxford, South Parks Road, Oxford, OX1 3QU, UK. [3]Department of Chemistry and Technologies of Drug, Sapienza University of Rome, 00185 Rome, Italy. [4]Department of Biology, University of Oxford, South Parks Road, Oxford, OX1 3RB, UK.

Email: abraham.oluwole@chem.ox.ac.uk; carol.robinson@chem.ox.ac.uk

Abstract: The peptidoglycan pathway represents one of the most successful drug targets with the last but critical step being the recycling of undecaprenyl phosphate. This translocation of undecaprenyl phosphate is facilitated by DedA and DUF386 domain-containing family membrane proteins via unknown mechanisms. Here we employ native mass spectrometry to investigate the interactions of *Bacillus subtilis* UptA, a putative C_{55}-P flippase, with C_{55}-P, membrane phospholipids and cell wall targeting antibiotics. Our results show that UptA, expressed and purified in *E. coli*, forms monomer-dimer equilibria, and binds to ligands in a pH-dependent fashion. Specifically, we show that UptA interacts more favourably with C_{55}-P over shorter-chain analogues and membrane phospholipids. Moreover, we demonstrate that lipopeptide antibiotics, amphomycin and aspartocin D, can directly intercept UptA function by outcompeting the substrate for the protein binding site. Overall, this study suggests that UptA-mediated translocation of C_{55}-P is potentially regulated by anionic phospholipids and provides insights for future development of antibiotics targeting carrier lipid recycling.

Introduction

The global threat of drug-resistant bacterial pathogens calls for renewed efforts in the search for new antibiotics, along with the identification of new biochemical pathways and interactions that can be effectively targeted. A key antibiotic target is peptidoglycan, the cell wall polymer that provides bacteria with resistance to osmotic stress and environmental assaults [1, 2]. Accordingly, topline antibiotics such as amoxicillin and vancomycin [3, 4] intercept the transpeptidation step of the peptidoglycan synthesis [5-7]. Moreover, promising antibiotic candidates, such as ramoplanin and moenomycin, target the transglycosylation step [8, 9]. Peptidoglycan biosynthesis begins with the formation of UDP-N-acetylmuramyl-pentapeptide in the cytosol [10], followed by its coupling to the lipid carrier undecaprenyl monophosphate (C_{55}-P) on the cytosolic side of the inner membrane (Fig.1A) [11, 12]. The resulting lipid I is further decorated with a GlcNAc residue to form lipid II [13] before being flopped across the inner membrane [14-16]. In the periplasm, the glycopeptide moieties of lipid II are cross-linked into a nascent peptidoglycan polymer and the lipid carrier is released as undecaprenyl diphosphate (C_{55}-PP) [17-19]. To participate in the next round of precursor transfer, C_{55}-PP must be dephosphorylated to form C_{55}-P [20-22] and then flipped across the inner membrane such that its phosphate headgroup returns to the cytoplasmic side. The mechanism by which C_{55}-P is translocated across the cytoplasmic membrane is the least understood of the membrane-associated steps of the peptidoglycan pathway (Fig. 1A).

Recently, proteins belonging to the DedA- and the DUF368 domain-containing families were shown to facilitate trans-bilayer transport of C_{55}-P in the Gram-positive bacteria *Bacillus subtilis* and *Staphylococcus aureus* [23], and in the Gram-negative bacteria *Vibrio cholerae* [24]. DedA proteins are highly conserved, having 8 homologues in the model bacteria *E. coli* and 6 homologues in *B. subtilis*, respectively [25-27]. *B. subtilis* DedA paralogues include UptA and PetA, which are proposed to facilitate trans-bilayer flipping of C_{55}-P and anionic phospholipids, respectively [23, 28]. The DUF368-domain containing proteins, exemplified by PopT, are less conserved, being found in *S. aureus* and *V. cholerae,* but are absent in *B. subtilis* and *E. coli* [29]. *V. cholerae* PopT becomes essential for cell survival only in alkaline conditions [24]. Several Gram-positive and Gram-negative bacteria encode one or multiple UptA-type transporters suggesting that C_{55}-P recycling, mediated by these flippases, is broadly conserved across different bacterial species. For example, the single DedA homologue in *Borrelia burgdorferi* is an essential protein required for proper cell division in this organism [30]. Accordingly, the recycling step of peptidoglycan biosynthesis mediated by UptA-type flippases can be targeted by antibiotics. This necessitates a molecular-level understanding of how UptA interacts with its cognate substrate, C_{55}-P, and potentially with membrane phospholipids, to elucidate the mechanism of C_{55}-P translocation for antibiotic targeting.

Herein we employ native mass spectrometry to elucidate how *B. subtilis* UptA interacts with multiple lipidic substrates and antibiotics. We show that purified UptA exists as an equilibrium of monomers and dimers. We further show that UptA interacts more favourably with C_{55}-P than C_{55}-PP and membrane phospholipids and

provide molecular-level evidence on the amino acid residues involved in C_{55}-P binding. We find that UptA

binds to its cognate ligand C_{55}-P in a pH-dependent manner and show that lipopeptide antibiotics, such as

amphomycin and aspartocin D, bind to UptA with higher affinity than other cell-wall targeting antimicrobial

peptides. Interestingly, these lipopeptides can intercept peptidoglycan biosynthesis by out-competing the

flippase for C_{55}-P, highlighting avenues for the development of novel antibiotics.

Fig. 1: UptA is primarily monomeric with a low population of dimer. A) A schematic depiction of

the lipid II cycle of peptidoglycan biosynthesis. **B**) Native mass spectrum of UptA (2.2 µM) liberated

from a buffer containing 200 mM ammonium acetate (pH 8.0), 0.05% LDAO. Peaks are assigned to

UptA in monomeric and dimeric forms. Insert, the model structure of UptA (AFO31823-F1) predicted

by AlphaFold (31). Predicted N- and C- termini are indicated. **C**) Quadrupole isolation of the UptA

dimer (9+ charge state).   **D**) MS/MS of the isolated UptA dimer charge state upon collisional

activation (190 V). The dimer dissociates into individual UptA protomers confirming their non-

covalent association.

Results

Monomer-dimer equilibrium of UptA. We expressed *B. subtilis* UptA in *E. coli* and introduced the

purified protein into the mass spectrometer from a buffer containing 0.05% LDAO and 200 mM

ammonium acetate (pH 8.0). The mass spectrum of UptA displays two charge state distributions

assigned to monomers (22963.49±0.70 Da) and dimers (45925.21±0.65 Da) (Fig. 1B). To confirm
the non-covalent association of UptA monomers, we isolated the 9+ charge state assigned to the
dimer and subjected these ions to high-energy collisional dissociation (HCD, Fig. 1C). The resulting
MS^2 spectrum displays new peaks that correspond by mass to UptA protomers, and bear charges
complementary to the isolated parent ions (+5 and +4) (Fig. 1D). Of note, the experimental mass for
the UptA monomer is higher than the theoretical sequence mass (22936.36 Da) by 28 Da,
suggesting an endogenous modification. To identify this unknown modification, we activated the
monomer (6+ charge state) using HCD, yielding a plethora of *b*- and *y*-type fragment ions. The
observed fragments are consistent with the formylation of the N-terminal methionine of UptA (Fig.
S1). Bacterial inner membrane proteins are normally co-translationally formylated on the N-
terminus, however, this modification is rarely observed *in vitro* due to subsequent removal by the
cytosolic deformylase (32). Therefore, retainment of N-terminal formylation on UptA suggests
periplasmic localisation of the N-terminus of UptA, thus supporting the predicted N-out topology
(23). Additionally, we observe UptA in monomeric and dimeric forms in other detergents (Fig. S2),
confirming the oligomeric composition of UptA, which might apply to other bacterial DedA
homologues.

UptA binds C_{55}-P with high affinity. We explored the binding affinity of UptA towards C_{55}-P by
recording mass spectra for solutions containing 2.5 µM UptA and 0 to 40 µM C_{55}-P in 200 mM
ammonium acetate, 0.05% LDAO at pH 8.0. The mass spectrum for the equimolar mixture of UptA
and C_{55}-P yielded peaks corresponding to UptA in its apo form and in complex with C_{55}-P (Fig. 2A),
an indication of high-affinity binding interactions. Further increase in the concentration of C_{55}-P
enhanced the intensity of the UptA:C_{55}-P complex, together with the emergence of additional
binding events (Fig. 2A). We deconvoluted the spectra (33) to extract mean relative intensities of
bound and unbound forms of UptA in the spectral series. The data reflected an increase in the
relative abundance of UptA:$(C_{55}$-P$)_n$ complexes as a function of the concentration of C_{55}-P (Fig. 2B).
For the first binding event, the relative intensity plateaued at a protein/ligand molar ratio of ~4 (Fig.
2B). By fitting the experimentally observed intensity ratios to the Hill equation, we obtained an
apparent K_d = 5.7 µM (95% CI: 5.0 to 6.4 µM) for the binding of one C_{55}-P molecule to UptA. This K_d
value for UptA/C_{55}-P binding is of the same order of magnitude as the binding interactions for
MurG/lipid I (K_d = 1.89±0.6 µM) (34) and MurJ/lipid II (K_d = 2.9±0.6 µM) (14), indicating that UptA
exhibits a similarly high affinity binding with respect to C_{55}-P.
To investigate the binding selectivity of UptA for monophosphate (C_{55}-P) versus diphosphate (C_{55}-
PP) forms of the lipid carrier, we performed competitive binding assays. First, we recorded mass
spectra for a mixture of UptA with increasing concentrations of C_{55}-PP. We observed peaks
assigned to UptA in ligand-free and C_{55}-PP-bound forms, and the relative intensity of the
UptA:C_{55}-PP complex increased with increasing concentration of C_{55}-PP (Fig. 2C). However, the

relative intensities of UptA:C_{55}-PP yielded a significantly higher apparent dissociation constant K_d =

15.3 µM (95% CI: 9.5 to 32.8 µM) than that obtained for C_{55}-P (K_d = 5.7±0.7 µM). Being within an

order of magnitude, these dissociation constants suggest that UptA forms extensive contact with

both forms of the lipid carriers, but that the protein interacts more favourably with C_{55}-P than C_{55}-PP.

We therefore conclude that UptA is more selective towards C_{55}-P than C_{55}-PP, thus confirming its

preferred endogenous substrate.

Fig 2. UptA binds C_{55}-P with a high affinity. A) Spectra for UptA incubated with different

concentrations of C_{55}-P in a buffer containing 200 mM ammonium acetate (pH 8), 0.05% LDAO. **B-**

C) The relative abundance of UptA bound to one (1x) or two (2x) molecules of C_{55}-P (panel B) and

C_{55}-PP (panel C). Inserts, chemical structures of C_{55}-P and C_{55}-PP. Each data point represents an

average of 3 replicate measurements, and the curves represent the best fits of the data to Hill's

equation (see methods), error bars are standard deviations. **D**) Mean relative intensity of ligand-free

and ligand-bound UptA in spectra recorded for 4 µM UptA incubated with 40 µM C_{20}-P, 40 µM C_{30}-P,

or 40 µM C_{55}-P. UptA binds the 'true' carrier lipid C_{55}-P in preference to the shorter chain analogues.

Bar is the mean of three (3) independent replicates, error bars are standard deviations.

UptA interacts more favourably with C_{55}-P than its shorter-chain analogues. Next, we

investigated the chain length selectivity of UptA by testing its interaction with C_{20}-P, C_{30}-P and C_{55}-P.

For this experiment, we incubated 4 µM UptA with a 10-fold molar excess of C_{20}-P, C_{30}-P and C_{55}-P

at pH 8.0 and recorded spectra under the same conditions. For the UptA/C_{20}-P sample, the

spectrum exhibits low-intensity peaks corresponding by mass to UptA:C_{20}-P complexes, leaving

most of the protein in the apo form (Fig. 2D, Fig. S3). We observed more intense protein/ligand

complexes in the case of UptA/C_{30}-P, indicating a higher binding affinity of C_{30}-P compared to C_{20}-P.

The spectra recorded for the UptA/C_{55}-P mixture under the same condition resolved up to four

binding events per UptA molecule (Fig. S3), indicating a more favourable interaction compared to

the lipidic substrate with shorter chains. Accordingly, the relative intensity of UptA in complex with

C_{55}-P is higher than in the case of UptA-bound C_{30}-P and C_{20}-P (Fig. 2D, Fig. S3). Thus, the

interaction of UptA with the lipid carrier is enhanced with the hydrophobic tail of C_{55}-P, supporting

our hypothesis that the long polyprenyl tail of the lipid carrier forms extensive contact with UptA.

Together, these data are consistent with C_{55}-P being the preferred endogenous substrate for UptA.

Amino acid residues mediating C_{55}-P interactions with UptA.

Two conserved arginine residues (R112 and R118) in the membrane re-entrant helices of UptA have

been suggested to impair the cellular function of UptA by making the cells more susceptible to

inhibition by amphomycin (23). Based on the model structure of UptA predicted by Alphafold, the

R112 residue should engage Q64 while R118 is able to form hydrogen bonds with E32 and W146 to

stabilise the intramembrane helices (Fig. 3A). To test for the impact of these residues on C_{55}-P

binding, we generated the E32A, Q64A, R112A, and R118A UptA point mutants. After purification of

the proteins, using the same protocol as the wild type, we equilibrated aliquots of each protein

variant (5 µM) with C_{55}-P (10 µM) and recorded spectra under the same instrument settings (Fig.

S4). The resulting spectra exhibited peaks consistent with C_{55}-P binding in all cases; only a modest

reduction in C_{55}-P binding intensity was observed in the case of the R112A mutant compared to the

wild type (Fig. 3B). We surmised that single point mutation was insufficient to attenuate binding and

generated a double mutant (R112A/R118A). Comparing the C_{55}-P binding affinity of the double

mutant with the wild type we found a more significant reduction in C_{55}-P binding, suggesting that

both residues are important for C_{55}-P binding. We additionally generated a double mutant
R112E/R118E in which the polarity of both native arginine residues in the WT are reversed. This
mutant exhibited the weakest C_{55}-P binding of all variants tested (Fig. 3B). This observation
suggests that Coulombic repulsion between the glutamates and C_{55}-P destabilises binding
interactions. Alternatively, impaired hydrogen bonding between R118 and EQ64 on the one hand,
and R118 and E32 could also disrupt C_{55}-P binding sites. Either or both mechanisms could be at
play, highlighting the critical roles of R112 and R118 in the binding of C_{55}-P to UptA.
169

170
171
Fig. 3. Probing the amino acid residues involved in UptA-C_{55}-P binding. A. Zoomed views of
UptA model structure showing hydrogen bonding networks surrounding R118 and R112. **B.** Relative
intensity of C_{55}-P bound to UptA wild type and mutants. Bars represent the mean of three biological
replicates with each data point shown in black, error bars are standard deviations. The double
mutant at residues R112 and R118 caused the most significant reduction in C_{55}-P binding.

C_{55}-P binding to UptA is sensitive to pH. Membrane flippases, such as MurJ, are driven by
proton-motive force (35) and exhibit pH-dependency in ligand binding (14). We considered the
possibility that substrate interactions of UptA might exhibit a pH dependency. To this end, we
equilibrated C_{55}-P with UptA at a protein/ligand molar ratio of 1:4 in a buffer of the same composition
but at a more acidic pH. The resulting spectrum exhibits peaks consistent with monomeric UptA in
an apo form, and in complex with 1 and 2 molecules of C_{55}-P (Fig. S5). We also observed UptA
dimer in complex with up to two C_{55}-P molecules (Fig. S5). However, the intensity of peaks assigned
to ligand-bound UptA is more intense at pH 8.0 compared to pH 5.0 (Fig. 4A, Fig. S5). These results
imply that more favourable interactions occur under physiologically relevant pH conditions. As a
negative control, we tested for pH sensitivity of C_{55}-P binding to MraY, the latter being the
downstream enzyme in the peptidoglycan pathway that couples UDP-MurNAc pentapeptide to

C_{55}-P. To this end, we equilibrated MraY and C_{55}-P at a molar ratio of 1:4 in a buffer of pH 8.0 and

pH 5.0 In this case, the spectra show no significant difference in the intensity of C_{55}-P bound to

MraY at pH 8.0 and at pH 5.0 (Fig. 4B, Fig. S5). We therefore conclude that environmental pH

modulates the interaction of C_{55}-P with UptA, and potentially its transport mechanism.

Fig 4. UptA preferentially binds C_{55}-P over membrane phospholipids. A-B) Mean relative

intensity of apo and lipid-bound UptA and MraY at pH 8.0 and pH 5.0. UptA binds C_{55}-P more

favourably at pH 8 than pH 5 but MraY exhibit no pH-dependency. **C**) Mean relative intensity of lipid-
bound UptA in the spectra for 4 μM UptA and 10 μM of each lipid. DOPG and POPG interact more
favourably with UptA than DOPE and POPE. C_{55}-P binds to UptA more intensely than the
phospholipids. Bar represents the mean of three independent replicates, data points are shown as
open circles, and error bars are standard deviations. **D**) Deconvoluted mass spectra of mixtures
containing 4 μM UptA,10 μM DOPG and increasing concentrations of C_{55}-P. C_{55}-P out-competes
phospholipids for UptA binding. **E**) Spectra for 4 μM UptA and 10 μM C_{55}-P, then incubated with
increasing concentrations of DOPG. The latter could not displace UptA-bound C_{55}-P.

C_{55}-P binds preferentially to dimeric forms of UptA. Since purified UptA displayed monomer-
dimer equilibria, we considered the possibility that C_{55}-P might preferentially interact with one
oligomeric species over the other. To probe this hypothesis, we used the UptA point mutant E32A
since the latter consistently retained a higher dimer population than the wild type (Fig. S6). We
prepared and analysed solutions containing 2.5 μM E32A UptA and 0-20 μM C_{55}-P at pH 8.0. The
resulting spectra exhibited peaks assigned to UptA monomers and dimers in apo and C_{55}-P bound
forms (Fig. S6). The mean relative intensity of ligand-bound dimers is however more intense than
the corresponding peaks for the ligand-bound monomers (Fig. S6). Together these data indicate that
the ligand C_{55}-P binds to the dimer fraction with a higher affinity than to monomers. Notably, the total
monomer-dimer fraction does not change significantly in the presence of C_{55}-P, and the preferential
ligand binding to the dimer fractions was observed at both pH 8.0 and pH 5.0 (Fig. S6). Together
these observations suggested that dimerisation promotes ligand interactions with UptA, and
potentially its cellular function.

C_{55}-P outcompetes phospholipids for UptA binding. DedA family proteins are proposed to either
translocate carrier lipids, membrane phospholipids, or both. We therefore probed the interaction of
UptA with membrane phospholipids to understand if the latter could modulate its function. For this
purpose, we selected the anionic lipids phosphatidylglycerols (PG) and phosphatidylethanolamines
(PE), being the major classes of phospholipid headgroups found in *B. subtilis* (36). We then
recorded mass spectra of solutions containing 4 μM UptA incubated with 10 μM POPE, DOPE,
POPG, DOPG and C_{55}-P. The resulting data shows that UptA binds to POPG and DOPG more
intensely than POPE and DOPE, respectively (Fig. 4C). Together these observations suggest that
UptA interacts more favourably with anionic than cationic lipids. Importantly, C_{55}-P binds to UptA
with higher relative affinity than any of these diacylglycerol phospholipids (Fig. 4C), in line with C_{55}-P
being its native substrate.

To investigate whether or not C_{55}-P can outcompete phospholipids in binding to UptA we incubated
solutions containing UptA-DOPG with increasing concentrations of C_{55}-P. Spectra were recorded

after aliquots of the solution were equilibrated on ice for 15 mins. Notably, spectra in the absence of

C_{55}-P reflected the binding of up to two molecules of DOPG per UptA (Fig. 4D). The presence of

C_{55}-P caused attenuation of the intensity DOPG-bound UptA in a concentration-dependent manner.

At the highest C_{55}-P concentration tested (20 µM), C_{55}-P has displaced most of DOPG from UptA

(Fig. 4D). We therefore hypothesised that phospholipids should be unable to outcompete C_{55}-P

bound to UptA. To test this hypothesis, we incubated UptA with C_{55}-P and then titrated in an

increasing concentration of DOPG. Indeed, C_{55}-P remains bound to UptA even in the presence of

excess DOPG (20 µM) (Fig. 4E), indicating that phospholipids could not displace C_{55}-P from UptA.

Analogous experiments with DOPE show that C_{55}-P effectively outcompete DOPE for UptA binding

but the latter could not displace C_{55}-P from UptA (Fig. S7). Together this study shows that UptA

interacts more favourably with C_{55}-P than with membrane phospholipids, thereby establishing C_{55}-P

as the preferred endogenous substrate.

Lipopeptide antibiotics disrupt UptA:C_{55}-P complex. We next sought to identify an antibiotic that

can intercept flippase function by disrupting its interaction with the carrier lipid substrate. We tested

a range of cell-wall targeting antibiotics before selecting daptomycin, amphomycin, and aspartocin,

as these lipopeptides bind to UptA with a high affinity (see below). Daptomycin is known to kill

bacteria by depolarising the membrane (37) and by complexing with lipid II (38). We investigated if

daptomycin affects C_{55}-P binding to UptA by recording spectra for solutions containing 2.5 µM UptA,

10 µM C_{55}-P and different concentrations of daptomycin. In the absence of antibiotic, the spectrum

exhibited peaks assigned to UptA in apo form, and in complex with 1 and 2 C_{55}-P molecules (Fig.

5A). In the presence of daptomycin, we observed additional peaks corresponding to the binary

complex UptA:daptomycin and the ternary complex UptA:C_{55}-P:daptomycin. Further increases in

daptomycin concentration resulted in higher intensity of these complexes, indicating that the

antibiotic does not compete against C_{55}-P binding to UptA. The peak intensity of

UptA:C_{55}-P:daptomycin ternary complex suggested that daptomycin does not compete for the same

binding sites as C_{55}-P on UptA, and does not sequester C_{55}-P significantly from UptA. Accordingly,

the reduction in the relative intensity of UptA:C_{55}-P complex is compensated for by the abundance

of UptA: C_{55}-P:daptomycin complex. (Fig. 5A).

Amphomycin and derivatives are known to kill bacteria by forming complexes with free C_{55}-P (39),

however, whether or not this involves a direct interaction with a protein cellular target is not known.

We therefore investigated how amphomycin affects the UptA:C_{55}-P complex. We recorded spectra

for solutions containing 2.5 µM UptA, 10 µM C_{55}-P and then equilibrated aliquots with increasing

concentrations of amphomycin. The spectra exhibited peaks consistent with amphomycin binding to

UptA, in addition to C_{55}-P binding. We find that the peak intensities of the UptA:C_{55}-P complex

reduce with respect to an increase in the concentration of amphomycin (Fig. 5B). Compared to the

case of daptomycin, however, little-to-no UptA:C_{55}-P complexes remained in the presence of 10-fold

molar excess of amphomycin (Fig. 5B). To verify these results, we performed similar experiments

with aspartocin D, an analogue of amphomycin. We find that aspartocin D similarly binds to UptA

and induces dissociation of UptA:C_{55}-P complex (Fig. S8). Bacitracin is a peptide antibiotic whose

mode of action is to sequester C_{55}-PP thereby halting carrier lipid homeostasis (40). To test the

specificity of amphomycin interaction with UptA and C_{55}-P, we prepared UptA/C_{55}-P mixtures and

then equilibrated aliquots with increasing concentrations of bacitracin. In line with the specificity of

bacitracin for C_{55}-PP rather than C_{55}-P, bacitracin binds to UptA only with a low intensity and does

not compete against C_{55}-P, even at elevated concentrations of 200 µM (Fig. S8). Taken together,

these data indicate that amphomycin and aspartocin D induce dissociation of UptA:C_{55}-P

complexes, by binding directly to UptA in addition to their known propensity to form complexes with

free C_{55}-P molecules (39).

Fig. 5 UptA function can be intercepted by lipopeptide antibiotics. A) Mass spectra (6+ charge

state) for solutions containing 2.5 µM UptA and 10 µM C_{55}-P incubated with 0-100 µM daptomycin.

Complexes of UptA with C_{55}-P and daptomycin are labelled. **B**) Spectra (6+ charge state) for
solutions containing 2.5 μM UptA and 10 μM C_{55}-P incubated with 0-100 μM amphomycin.
Amphomycin destabilises the UptA:C_{55}-P complex whereas daptomycin has a modest impact. Data
are displayed as the mean of three independent replicates, and error bars are standard deviations.
291

Discussions
In this study, using native MS, we investigated the interplay of multiple lipidic substrates and
antibiotics for UptA (summarised in Fig. 6). UptA is a representative member of the DedA family of
transporters that mediate carrier lipid recycling in peptidoglycan biosynthesis. Specifically, we report
that purified UptA exists as an equilibrium of monomers and dimers, suggesting that DedA proteins
might form dimers in the native membrane when lateral pressures in the lipid bilayers (41) might
prevent their dissociation. In line with the dependency of DedA family proteins on proton motive
force (42), we found that the monomer-dimer ratio of UptA is sensitive to pH. This feature contrasts
with those of the upstream peptidoglycan enzymes MraY (34, 43) and UppP (44, 45) whose
dimerization are promoted by lipid binding.
Our studies also show that UptA selectively binds to C_{55}-P over C_{55}-PP and that the protein interacts
more favourably with C_{55}-P than the shorter chain analogues C_{30}-P and C_{20}-P. This implies that the
hydrophobic tail of C_{55}-P makes extensive contact with UptA, promoting the binding interactions and
potentially facilitating the transport of carrier lipids. The higher affinity of UptA towards the
monophosphate form of the carrier lipid, rather than the diphosphate form, favours a mechanism in
which UptA mediates unidirectional flipping of C_{55}-P rather than acting as a scramblase for both C_{55}-
P and C_{55}-PP. We further show that UptA interacts more favourably with the anionic lipids DOPG
and POPG than the cationic lipids DOPE and POPE and that these lipids are readily displaced by
C_{55}-P. This binding preference agrees with UptA being responsible for C_{55}-P rather than the
phospholipid translocation (23). Competition between anionic lipids, which are more abundant in the
native lipid bilayer than C_{55}-P, suggests that both lipids bind near the same sites on UptA and that
the local lipid environment of UptA can potentially modulate its cellular function. We observed less
C_{55}-P binding to UptA in the presence of excess H^+. This suggests that the UptA undergoes a
proton-driven conformational change accompanying C_{55}-P flipping and release, similar to the
proton-coupled amino acid (46) and antimicrobial peptide symporters (47).
We provided molecular-level evidence that the conserved arginine (R112 and R118) residues in the
putative membrane re-entrant helices of UptA are important for its substrate binding interactions,
confirming earlier reports that *B. subtilis* strains bearing these mutations on UptA are more
susceptible to inhibition by MX2401, a derivative of amphomycin (23). We further showed that
amphomycin and daptomycin bind with high affinity to UptA. Accordingly, the chemical scaffolds of
these lipopeptides can therefore inspire the development of analogues that can intercept the cellular
function of UptA and thus compromise the cell wall formation.

Despite the combined cellular pool of C_{55}-PP and C_{55}-P being limited to ~1.5×10^5 molecules per cell

(48), they mediate the transport of glycan-bearing components of the cell wall peptidoglycan,

teichoic acid and capsular polysaccharide (49). Therefore, antibiotics that intercept C_{55}-P and its

flippases are potential drug targets that can potentially intercept multiple pathways and thus mitigate

drug resistance. Accordingly, C_{55}-PP is the target of bacitracin and teixobactin (50, 51). Friulimicin

and amphomycin analogues intercept bacterial cell wall formation by forming complexes with C_{55}-P

(23, 52-54). Herein we show that amphomycin and aspartocin D can additionally induce dissociation

of the UptA: C_{55}-P complex, and bind directly to UptA, a finding that can therefore guide the

development of new antibiotic analogues to intercept the flippase function of UptA and its

homologues.

Fig. 6 Insights from native MS into the interaction of UptA with lipidic substrates and antibiotics.

(i) UptA exhibits monomer-dimer equilibria, with more dimer at pH 8.0 than pH 5.0. (ii) UptA interacts

with C_{55}-P more favourably than C_{55}-PP, C_{30}-P, and C_{20}-P; dimerization promotes C_{55}-P binding. (iii)

Amphomycin destabilises the UptA:C_{55}-P complex, effectively out-competing the flippase for its

carrier lipid substrate. (iv) UptA binds phospholipids, with a higher preference for the anionic than

cationic lipids. (v) C_{55}-P is able to displace UptA-bound phospholipids, suggesting that both ligands

interact with UptA near the same sites. (iv) Binding of H^+ to UptA enables translocation of C_{55}-P and

its subsequent release for the next round of PG precursor synthesis.

344   REFERENCES
345
1.   T. Schneider, H.-G. Sahl, An oldie but a goodie – cell wall biosynthesis as antibiotic target
pathway. *International Journal of Medical Microbiology* **300**, 161-169 (2010).
2.   J. M. C. Kwan, Y. Qiao, Mechanistic Insights into the Activities of Major Families of Enzymes
in Bacterial Peptidoglycan Assembly and Breakdown. *ChemBioChem* **24**, e202200693 (2023).
3.   A. Coleman, Y. Vohra, K. Rascati, S. Kubes, B. Moffett, Antibiotic Utilization and Efficacy
Associated With Treating Pediatric Urinary Tract Infections in Texas Medicaid Patients in the
First Year of Life. *Pediatr Infect Dis J* **40**, 993-996 (2021).

4. E. Sauvage, M. Terrak, Glycosyltransferases and Transpeptidases/Penicillin-Binding Proteins:
Valuable Targets for New Antibacterials. *Antibiotics* **5**, 12 (2016).

5. A. J. Meeske *et al.*, MurJ and a novel lipid II flippase are required for cell wall biogenesis in
Bacillus subtilis. *Proc Natl Acad Sci U S A* **112**, 6437-6442 (2015).

6. A. C. Y. Kuk, A. Hao, S. Y. Lee, Structure and Mechanism of the Lipid Flippase MurJ. *Annu Rev*
Biochem **91**, 705-729 (2022).

7. S. Kumar, A. Mollo, D. Kahne, N. Ruiz, The Bacterial Cell Wall: From Lipid II Flipping to
Polymerization. *Chem Rev* **122**, 8884-8910 (2022).

8. M.-C. Lo *et al.*, A New Mechanism of Action Proposed for Ramoplanin. *Journal of the*
American Chemical Society **122**, 3540-3541 (2000).

9. X. Fang *et al.*, The mechanism of action of ramoplanin and enduracidin. *Molecular*
BioSystems **2**, 69-76 (2006).

10. H. Barreteau *et al.*, Cytoplasmic steps of peptidoglycan biosynthesis. *FEMS Microbiology*
Reviews **32**, 168-207 (2008).

11. B. C. Chung *et al.*, Crystal structure of MraY, an essential membrane enzyme for bacterial cell
wall synthesis. *Science* **341**, 1012-1016 (2013).

12. A. Bouhss, A. E. Trunkfield, T. D. Bugg, D. Mengin-Lecreulx, The biosynthesis of peptidoglycan
lipid-linked intermediates. *FEMS microbiology reviews* **32**, 208-233 (2007).

13. S. Ha, D. Walker, Y. Shi, S. Walker, The 1.9 A crystal structure of Escherichia coli MurG, a
membrane-associated glycosyltransferase involved in peptidoglycan biosynthesis. *Protein Sci*
9, 1045-1052 (2000).

14. J. R. Bolla *et al.*, Direct observation of the influence of cardiolipin and antibiotics on lipid II
binding to MurJ. *Nat Chem* **10**, 363-371 (2018).

15. S. Kumar, A. Mollo, D. Kahne, N. Ruiz, The bacterial cell wall: from lipid II flipping to
polymerization. *Chemical Reviews* **122**, 8884-8910 (2022).

16. K. Bupp, J. van Heijenoort, The final step of peptidoglycan subunit assembly in Escherichia
coli occurs in the cytoplasm. *J Bacteriol* **175**, 1841-1843 (1993).

17. A. J. F. Egan, J. Errington, W. Vollmer, Regulation of peptidoglycan synthesis and remodelling.
Nat Rev Microbiol **18**, 446-460 (2020).

18. V. M. Hernández-Rocamora *et al.*, Real-time monitoring of peptidoglycan synthesis by
membrane-reconstituted penicillin-binding proteins. *eLife* **10**, e61525 (2021).

19. Y. Park, A. Taguchi, V. Baidin, D. Kahne, S. Walker, A Time-Resolved FRET Assay Identifies a
Small Molecule that Inhibits the Essential Bacterial Cell Wall Polymerase FtsW. *Angewandte*
Chemie International Edition **62**, e202301522 (2023).

20. M. El Ghachi, A. Bouhss, D. Blanot, D. Mengin-Lecreulx, The bacA gene of Escherichia coli
encodes an undecaprenyl pyrophosphate phosphatase activity. *J Biol Chem* **279**, 30106-
30113 (2004).

21. M. El Ghachi, A. Derbise, A. Bouhss, D. Mengin-Lecreulx, Identification of multiple genes
encoding membrane proteins with undecaprenyl pyrophosphate phosphatase (UppP)
activity in Escherichia coli. *J Biol Chem* **280**, 18689-18695 (2005).

22. R. Bernard, M. El Ghachi, D. Mengin-Lecreulx, M. Chippaux, F. Denizot, BcrC from Bacillus
subtilis acts as an undecaprenyl pyrophosphate phosphatase in bacitracin resistance. *J Biol*
Chem **280**, 28852-28857 (2005).

23. I. J. Roney, D. Z. Rudner, Two broadly conserved families of polyprenyl-phosphate
transporters. *Nature* **613**, 729-734 (2023).

24. B. Sit *et al.*, Undecaprenyl phosphate translocases confer conditional microbial fitness.
Nature **613**, 721-728 (2023).

25.   W. T. Doerrler, R. Sikdar, S. Kumar, L. A. Boughner, New Functions for the Ancient DedA
Membrane Protein Family. *Journal of Bacteriology* **195**, 3-11 (2013).
26.   P. R. Panta, W. T. Doerrler, A Burkholderia thailandensis DedA Family Membrane Protein Is
Required for Proton Motive Force Dependent Lipid A Modification. *Frontiers in Microbiology*
11 (2021).
27.   S. Kumar, W. T. Doerrler, Members of the conserved DedA family are likely membrane
transporters and are required for drug resistance in Escherichia coli. *Antimicrob Agents*
Chemother **58**, 923-930 (2014).
28.   I. J. Roney, D. Z. Rudner, The DedA superfamily member PetA is required for the transbilayer
distribution of phosphatidylethanolamine in bacterial membranes. *Proceedings of the*
National Academy of Sciences **120**, e2301979120 (2023).
29.   A. Galinier, C. Delan-Forino, E. Foulquier, H. Lakhal, F. Pompeo, Recent Advances in
Peptidoglycan Synthesis and Regulation in Bacteria. *Biomolecules* **13**, 720 (2023).
30.   F. T. Liang *et al.*, BB0250 of Borrelia burgdorferi is a conserved and essential inner membrane
protein required for cell division. *J Bacteriol* **192**, 6105-6115 (2010).
31.   J. Jumper *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
583-589 (2021).
32.   A. Ranjan, E. Mercier, A. Bhatt, W. Wintermeyer, Signal recognition particle prevents N-
terminal processing of bacterial membrane proteins. *Nature Communications* **8**, 15562
(2017).
33.   M. T. Marty *et al.*, Bayesian deconvolution of mass and ion mobility spectra: from binary
interactions to polydisperse ensembles. *Anal Chem* **87**, 4370-4376 (2015).
34.   A. O. Oluwole *et al.*, Peptidoglycan biosynthesis is driven by lipid transfer along enzyme-
substrate affinity gradients. *Nature Communications* **13**, 2278 (2022).
35.   F. A. Rubino, S. Kumar, N. Ruiz, S. Walker, D. E. Kahne, Membrane Potential Is Required for
MurJ Function. *Journal of the American Chemical Society* **140**, 4481-4484 (2018).
36.   J. A. den Kamp, I. Redai, L. L. van Deenen, Phospholipid composition of Bacillus subtilis. *J*
Bacteriol **99**, 298-303 (1969).
37.   W. R. Miller, A. S. Bayer, C. A. Arias, Mechanism of action and resistance to daptomycin in
Staphylococcus aureus and enterococci. *Cold Spring Harbor perspectives in medicine* **6**
(2016).
38.   F. Grein *et al.*, Ca2+-Daptomycin targets cell wall biosynthesis by forming a tripartite complex
with undecaprenyl-coupled intermediates and membrane lipids. *Nature Communications* **11**,
1455 (2020).
39.   T. M. Wood *et al.*, Mechanistic insights into the C 55-P targeting lipopeptide antibiotics
revealed by structure–activity studies and high-resolution crystal structures. *Chemical*
Science **13**, 2985-2991 (2022).
40.   J. Smith, E. Weinberg, Mechanisms of antibacterial action of bacitracin. *Microbiology* **28**,
559-569 (1962).
41.   R. S. Cantor, Lateral pressures in cell membranes: a mechanism for modulation of protein
function. *The journal of physical chemistry B* **101**, 1723-1725 (1997).
42.   P. R. Panta, W. T. Doerrler, A Burkholderia thailandensis DedA family membrane protein is
required for proton motive force dependent lipid a modification. *Frontiers in Microbiology*
11, 618389 (2021).
43.   E. Henrich *et al.*, Lipid Requirements for the Enzymatic Activity of MraY Translocases and in
Vitro Reconstitution of the Lipid II Synthesis Pathway. *J Biol Chem* **291**, 2535-2546 (2016).

44. S. D. Workman, L. J. Worrall, N. C. J. Strynadka, Crystal structure of an intramembranal
phosphatase central to bacterial cell-wall peptidoglycan biosynthesis and lipid recycling.
Nature Communications **9**, 1159 (2018).
45. A. O. Oluwole *et al.*, Real-time biosynthetic reaction monitoring informs the mechanism of
action of antibiotics. *J. Am. Chem. Soc.* **146**, 7007-7017 (2024).
46. M. Löbel *et al.*, Structural basis for proton coupled cystine transport by cystinosin. *Nature*
Communications **13**, 4845 (2022).
47. D. Ghilarov *et al.*, Molecular mechanism of SbmA, a promiscuous transporter exploited by
antimicrobial peptides. *Science Advances* **7**, eabj5363 (2021).
48. H. Barreteau *et al.*, Quantitative high-performance liquid chromatography analysis of the
pool levels of undecaprenyl phosphate and its derivatives in bacterial membranes. *J*
Chromatogr B Analyt Technol Biomed Life Sci **877**, 213-220 (2009).
49. M. Rausch *et al.*, Coordination of capsule assembly and cell wall biosynthesis in
Staphylococcus aureus. *Nat Commun* **10**, 1404 (2019).
50. L. L. Ling *et al.*, A new antibiotic kills pathogens without detectable resistance. *Nature* **517**,
455-459 (2015).
51. R. Shukla *et al.*, Teixobactin kills bacteria by a two-pronged attack on the cell envelope.
Nature **608**, 390-396 (2022).
52. T. M. Wood *et al.*, Mechanistic insights into the C55-P targeting lipopeptide antibiotics
revealed by structure–activity studies and high-resolution crystal structures. *Chemical*
Science **13**, 2985-2991 (2022).
53. T. Schneider *et al.*, The lipopeptide antibiotic Friulimicin B inhibits cell wall biosynthesis
through complex formation with bactoprenol phosphate. *Antimicrob Agents Chemother* **53**,
1610-1618 (2009).
54. Z. Wang, B. Koirala, Y. Hernandez, M. Zimmerman, S. F. Brady, Bioinformatic prospecting and
synthesis of a bifunctional lipopeptide antibiotic that evades resistance. *Science* **376**, 991-
996 (2022).
473
474

Supporting information

Lipopeptide antibiotics disrupt interactions of undecaprenyl phosphate with UptA

Abraham O. Oluwole[1,2*], Neha Kalmankar[1,2], Michela Guida[2,3], Jack L. Bennett[1,2], Giovanna Poce[3],

Jani R. Bolla[2,4] Carol V. Robinson[1,2*]

[1]Department of Chemistry, University of Oxford, South Parks Road, Oxford, OX1 3QZ, UK. [2]The Kavli Institute

for Nanoscience Discovery, Oxford, South Parks Road, Oxford, OX1 3QU, UK. [3]Department of Chemistry and

Technologies of Drug, Sapienza University of Rome, 00185 Rome, Italy. [4]Department of Biology, University of

Oxford, South Parks Road, Oxford, OX1 3RB, UK.

Email: abraham.oluwole@chem.ox.ac.uk; carol.robinson@chem.ox.ac.uk

Table of Contents

                                          Methods

Chemicals

C_{55}-PP, C_{55}-P, C_{30}-P and C_{20}-P were purchased from Larodan. Phospholipids – POPE, DOPE, POPG,

DOPG – were purchased from Avanti Polar Lipids. Amphomycin and aspartocin D were purchased

from Cayman; while Daptomycin and bacitracin were purchased from Merck. *n*-dodecyl-*β*-D-

maltoside (DDM), *n*-decyl-*β*-D-maltoside (DDM), n-octylglucoside (OG) and *n*-dodecyl-*N,N*-

dimethylamine-*N*-oxide (LDAO) were from Anatrace.

Plasmids

A codon-optimised version of *Bacillus subtilis* UptA (yngC) (Integrated DNA Technologies) was

inserted into a modified pET15b vector between NdeI and NheI endonuclease restriction sites by In-

Fusion HD cloning kit (Takara Bio). The resulting plasmid encodes UptA fused to GFP-His$_6$ on the C-

terminus via a TEV-protease cleavable site. Plasmids for UptA mutants were generated by site

directed mutagenesis with CloneAmp HiFi PCR Premix (Takara Bio) using pairs of oligonucleotides

listed in Table S1. The plasmid pET26-MBP-MraYaa bearing HRV3C protease cleavage site between

the MBP and the wild-type MraY was a kind gift from S-Y Lee (Addgene plasmid #100166) (1). MraY

was expressed and purified as previously described (2).

Table S1: Oligonucleotides using to generate plasmids for the expression of UptA mutants.

Primers	Sequence
E32AF	TCCAAGCGCAATTGTATTGGCCTACGGTGG
E32AR	ACAATTGCGCTTGGAATAATTTCAATCATCAGT
Q64AF	GATCGCCGCAATTTTCATTTACTGGATTGGCCGT
Q64AR	AAAATTGCGGCGATCGTTCCCCCGATA
R112AF	TTCTGCTGCCTTCATCCCTGTAGTCCGTCATGC
R112AR	ATGAAGGCAGCAGAAAACACAACCCCCG
R118AF	TGTAGTCGCACATGCTATTTCGATCCCAGCCG
W146AF	CATTCCTGCCTCCATCCTGTTCGTTTACCTGG
W146AR	ATGGAGGCAGGAATGATGGCAAGTACTGTC
R112A/R118A_Fwd	TTCTGCTGCCTTCATCCCTGTAGTCGCACATGCTATTTCGATCCCAGCCG
R112A/R118A_Rev	GCATGTGCGACTACAGGGATGAAGGCAGCAGAAAACACAACCCCCG
R112E/R118E_Fwd	ATTCATCCCTGTAGTCGAACATGCTATTTCGATCCCAGCCG
R112E/R118E_Rev	ACTACAGGGATGAATTCAGCAGAAAACACAACCCCCG

Expression and purification of UptA

Plasmids were used to transform chemically competent *E. coli* C43(DE3) cells (Lucigen) and selected

on LB/Agar containing 100 µg/ml ampicillin at 37 °C. Single colonies were used to inoculate 60 mL

LB containing 100 µg/ml ampicillin and were grown at 37 °C overnight for 16 h. For large-scale protein

expression, 10 mL of the overnight culture was aseptically transferred into 1 L of LB/ampicillin and cells were allowed to grow at 37°C until the $OD_{600\ nm}$ reached 0.6-0.8 after which the temperature was lowered to 18 °C. Expression of UptA-GFP was induced by adding isopropyl-β-D-thiogalactopyranoside (IPTG) at a final concentration of 0.5 mM UptA-GFP and shaking for 20 h at 18 °C. Cells were harvested by centrifugation at 5000 × *g* and lysed immediately or stored at -80 ° C until required.

Cells were thawed on ice and resuspended in a buffer containing 50 mM Tris-HCl (pH 7.4), 200 mM NaCl, 10 µg/mL deoxyribonuclease I (DNAse I), 5 mM $MgCl_2$, 5 mM β-mercaptoethanol and EDTA-free protease inhibitor cocktail tablets. The cells were then disrupted with a microfluidiser (Microfluidics) operated at 20,000 psi. Non-lysed cells and debris were pelleted by centrifugation at 20,000 × *g* for 20 min and the clarified lysate was retained. The membranes fraction was then pelleted by ultracentrifugation at 100,000 x g for 1 h. They were resuspended in a buffer containing 20 mM Tris-HCl (pH 8.0), 300 mM NaCl, 20% glycerol. Membrane aliquots were either solubilised immediately or flash-frozen in liquid nitrogen and stored at -80°C. The membrane was solubilised by incubation with DDM at a final concentration of 1% (w/v) for 60 min. Non-solubilised aggregates were removed by centrifugation at 20,000×g for 20 min and the supernatant was loaded onto a 5-ml HisTrap HP column preequilibrated in buffer A (20 mM Tris-HCl (pH 8.0), 200 mM NaCl, 10% glycerol, 0.03% DDM and 20 mM imidazole). After binding of the His-tagged protein, the column was washed with 100 mL of buffer A, and then with 50 mL of buffer B (20 mM Tris-HCl (pH 8.0), 200 mM NaCl, 10% glycerol, 0.03% DDM and 100 mM imidazole). UptA-GFP was eluted from the column with buffer C (20 mM Tris-HCl (pH 8.0), 200 mM NaCl, 10% glycerol, 0.02% DDM and 300 mM imidazole). To remove imidazole and to cleave the GFP fusion, eluted protein was mixed with TEV protease and dialysed against buffer D (20 mM Tris-HCl (pH 8.0), 150 mM NaCl, 10% glycerol, 0.02% DDM) at 4°C for 16 h.

The digestion mixture was subsequently incubated with 2-mL bed volume of nickel nitriloacetic acid (NTA) agarose resin to remove His-tagged GFP and TEV protease. Tag-free UptA was collected as the flowthrough, concentrated using an Amicon centrifugal concentrator with nominal molecular weight cut-off 50 kDa. The protein was finally purified by size-exclusion chromatography (SEC) using a 24-mL Superdex 200 Increase column. The SEC buffer was 20 mM Tris-HCl (pH 8.0), 200 mM NaCl, 0.02% DDM. UptA was concentrated to ~200 uM, aliquoted, flash-frozen flash in liquid nitrogen, and stored at -80 °C until use. Concentration was measured using a Biomate UV spectrophotometer at 280 nm using extinction coefficients calculated from their predicted amino acid sequences. UptA mutants were purified in 3 biological replicates, using the same procedures as for the wild-type.

Sample preparation for native MS
Prior to measurement, the protein was buffer-exchanged into 0.05% LDAO, 200 mM ammonium acetate ("MS buffer") at desired pH in using a centrifugal buffer exchange device (Micro Bio-Spin 6, Bio-Rad). Stock solutions of lipids were prepared from chloroform/methanol solution by evaporating

aliquots of known volume in a SpeedVac. After drying, the lipid films were weighed and resuspended

to a final concentration of 0.5 mM in 200 mM ammonium acetate (pH8) and 0.05% LDAO by vertexing.

Amphomycin and aspartocin D were purchased from Cayman while Daptomycin and bacitracin were

purchased from Merck. Stock solutions of 0.5 mM amphomycin, aspartocin D, daptomycin and

bacitracin, were made in the same buffer. All experiments were repeated three times from newly

prepared stock solutions.

Native mass spectrometry

101
About 3 µL protein aliquot was transferred into a gold-coated borosilicate capillary (Harvard

Apparatus) and was mounted on the nano ESI source of a Q-Exactive hybrid quadrupole-Orbitrap

mass spectrometer (Thermo Fisher Scientific, Bremen, Germany). The instrument settings were

1.2 kV capillary voltage, S-lens RF 200%, argon UHV pressure 3.3×10^{-10} mbar, the capillary

temperature was set to 200 °C, and resolution of the instrument was set to 17,500 at a transient time

of 64 ms. Voltages of the ion transfer optics –injection flatapole, inter-flatapole lens, bent flatapole,

and transfer multipole were set to 5, 3, 2, and 30 V respectively. The noise level was set at 3. Unless

otherwise stated, proteins were activated by applying 75 V in the high-energy collisional dissociation

cell without in-source trapping. For the MS^2 spectra, parent ions were isolated using an in-source

trapping voltage of -200 V, and then fragmented by applying 150-200 V in the high-energy collision-

induced dissociation cell. To identify endogenous modifications using native top-down mass

spectrometry, the 6+ charge state of monomeric UptA was isolated (3828 ± 5 Th) and activated using

HCD (200 V) with reduced trapping pressures. The resulting fragments were detected in the Orbitrap

(R = 280,000 @ m/z 200), averaging ~500 transients to maximise the signal-to-noise ratio. Theoretical

fragments from N-terminally formylated UptA were matched to the MS^2 spectrum (3 ppm) using

TDValidator (Proteinaceous) and manual validation. Data were visualised and exported for processing

using the Qual browser of Xcalibur 4.1.31.9 (Thermo Scientific). Spectral deconvolution was

performed using UniDec 6.0.2 (3). Relative binding affinities were obtained from deconvoluted spectra

by dividing the intensity of ligand-bound protein peaks by the sum of the intensities of ligand-bound

and ligand-free protein peaks. The mean and standard deviation of these fractional binding intensities

from three independent experiments were plotted against the concentrations of the ligand C_{55}-P or

C_{55}-PP. To obtain the apparent dissociation constant K_d, the mean relative intensities at different

concentrations were fitted globally using GraphPad Prism 10.2.0 to Eq. (1):

$$y = \frac{B_{max}x^h}{x^h + K_d{}^h} \qquad\qquad (1)$$

Where y is the mean relative intensity and x is the corresponding ligand concentration. B_{max} is the

maximum specific binding, K_d is the apparent dissociation constant, and h is the Hill coefficient.

Supplementary Figures

Fig. S1. UptA is formylated at the N-terminus. A. HCD MS/MS spectrum of UptA (6+). The eight most abundant assigned ions are labelled. **B**. Sequence map of the UptA construct used in this study. Terminal *b*- and *y*-type fragment ions observed in the MS/MS spectrum are indicated. **C.** Expanded view of the peak corresponding to b_{80}^{2+} ion**.** The theoretical isotopic envelope is overlaid.

Fig. S2. Oligomeric state of UptA in different detergents. Shown are the spectra recorded for UptA in buffer containing 200 mM Ammonium acetate (pH 8.0) supplemented with indicated detergents. The instrument was tuned differently in each case to maximise the transmission of dimeric species.

Fig. S3. UptA binds carrier lipids of longer chain lengths more intensely than those of shorter chain. Shown are the spectra recorded for 4 µM UptA equilibrated with 10-fold molar excess of C_{20}-P, C_{30}-P and C_{55}-P.

Fig. S4. Native MS analysis of C$_{55}$-P binding to UptA mutants. Shown are the representative

spectra for UptA wild type and mutants (5 uM) incubated with C$_{55}$-P (10 uM). Proteins were liberated

from a buffer containing 0.05% LDAO and 200 mM ammonium acetate (pH8. 02) using a collisional

activation of 100 V.

Fig. S5. UptA binds C₅₅-P more favourably at pH 8.0 than at pH 5.0. **A**. Spectra for UptA/C_{55}-P

mixtures at indicated pH. Inserts, the relative abundance of apo UptA and C_{55}-P bound forms. **B.**

Spectrum for MraY/C_{55}-P mixture at pH 8.0 and pH 5.0. There is no pH-dependency in C_{55}-P binding

by MraY.

Fig. S6. Dimer fraction of UptA binds C$_{55}$-P more than the monomeric form. **A.** Spectra from UptA

E32A mutant display peaks corresponding to monomers and dimers. Insert, the relative intensity of

monomer-dimer populations in the E32A mutant and the wild-type UptA**.** More dimers are observed

at pH 8.0 than at pH 5.0. **B-C.** Spectrum of UptA E32A mutant equilibrated with 10 µM and 20 µM C$_{55}$-

P. Insert, the relative intensity of apo and C$_{55}$-P bound proteins in each spectrum. Dimer fractions bind

the ligand more than the monomers. **D.** The total proportion of monomers and dimers in a mixture of

E32A UptA and 0-20 µM C$_{55}$-P. The ligand C$_{55}$-P does not cause a significant difference in the

monomer-dimer ratios. **E.** Mean relative intensity of apo and C$_{55}$-P bound E32A UptA at pH5 and pH8.

C$_{55}$-P bind to UptA more intensely at pH 8.0 than at pH 5.0.

Fig. S7. Competition for UptA between DOPE and C₅₅-P. A. Deconvoluted mass spectra for

solutions containing 4 µM UptA and 10 µM DOPE but different concentrations of C_{55}-P. The DOPE

bound to UptA was displaced by the added C_{55}-P. **B.** Spectra for solutions containing 4 µM UptA and

10 µM C_{55}-P but different concentrations of DOPE. C_{55}-P remains bound to UptA in the presence of

added phospholipids.

Fig. S8. Aspartocin binds to UptA and disrupts UptA:C_{55}-P interactions. **A.** Spectra (6+ charge state) of 2.5 µM UptA and 10 µM C_{55}-P in the presence of 0-80 µM aspartocin D. Peaks corresponding to UptA: C55-P (highlighted red) decreased in intensity as a function of aspartocin D concentration. **B**. Spectrum recorded for 2.5 µM UptA and 10 µM C_{55}-P but in the presence of 0, 100 and 200 µM bacitracin. Bacitracin binds to UptA only weakly and does not affect C_{55}-P binding.

REFERENCES

1. B. C. Chung *et al.*, Crystal structure of MraY, an essential membrane enzyme for bacterial cell wall synthesis. *Science* **341**, 1012-1016 (2013).
2. A. O. Oluwole *et al.*, Peptidoglycan biosynthesis is driven by lipid transfer along enzyme-substrate affinity gradients. *Nature Communications* **13**, 2278 (2022).
3. M. T. Marty *et al.*, Bayesian Deconvolution of Mass and Ion Mobility Spectra: From Binary Interactions to Polydisperse Ensembles. *Analytical Chemistry* **87**, 4370-4376 (2015).