Dynamic Tracking of Native Precursors in Adult Mice

Suying Liu[1,2], Sarah E. Adams[1,2], Haotian Zheng[3], Juliana Ehnot[1,2], Seul K. Jung[1,2], Greer Jeffrey[1,2],

Theresa Menna[1,2], Louise E. Purton[4,5], Hongzhe Lee[3], and Peter Kurre[1,2*]

[1] Comprehensive Bone Marrow Failure Center, Children's Hospital of Philadelphia, Philadelphia, PA

19104, USA

[2]Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA

[3]Department of Biostatistics, Epidemiology and informatics, University of Pennsylvania, Philadelphia, PA

19104, USA

[4]Stem Cell Regulation Unit, St. Vincent's Institute of Medical Research, Fitzroy, VIC 3065, Australia

[5]Department of Medicine, The University of Melbourne, Parkville, VIC 3052, Australia

*Correspondence: kurrep@chop.edu

Abstract
Hematopoietic dysfunction has been associated with a reduction in the number of active precursors.
However, precursor quantification at homeostasis and under diseased conditions is constrained by
the scarcity of available methods. To address this issue, we optimized a method for quantifying a
wide range of hematopoietic precursors. Assuming the random induction of a stable label in
precursors following a binomial distribution, estimates depend on the inverse correlation between
precursor numbers and the variance of precursor labeling among independent samples.
Experimentally validated to cover the full dynamic range of hematopoietic precursors in mice (1 to
10^5), we utilized this approach to demonstrate that thousands of precursors, which emerge after
modest expansion during fetal-to-adult transition, contribute to native and perturbed hematopoiesis.
We further estimated the number of precursors in a mouse model of Fanconi Anemia, showcasing
how repopulation deficits can be classified as autologous (cell proliferation) and non-autologous
(lack of precursor). Our results support an accessible and reliable approach for precursor
quantification, emphasizing the contemporary perspective that native hematopoiesis is highly
polyclonal.

Introduction

Continuous self-renewal and differentiation of hematopoietic stem and progenitor cells (HSPCs) is

fundamental to blood production. Even though rare cases exist where a single HSPC clone supports

hematopoiesis, the HSPC population contributing to homeostatic hematopoiesis is usually highly

polyclonal [1,2]. For example, the number of HSPCs actively participating in white blood cell

production was estimated to range from 20,000 to 200,000 in humans [3,4]. Conversely, a decline in

the number of active hematopoietic precursors has been linked to hematopoietic dysfunction. For

example, in humans older than 70 years of age, hematopoietic dysfunction coincides with an abrupt

reduction in the HSPCs population actively contributing to blood production [4].

These observations underscore the association between a low number of active hematopoietic

precursors and hematological function. Yet, few methods are suitable for quantifying active

hematopoietic precursors in a native environment [3–8]. In mice, methods utilizing *in situ* barcodes

often suffer from barcode homoplasy, preventing total precursor number estimation [9,10]. In

humans, estimations based on somatic mutations and computational modeling carry a high degree

of uncertainty [3,4]. The absence of precise precursor quantification methods in their natural

environment hampers the study of precursor numbers across different conditions and their potential

use as predictive functional markers.

In mice, the quantification of developmental hematopoietic precursors has been achieved using

animals engineered with a Confetti cassette [11]. This cassette can randomly recombine and

express one of four fluorescence proteins (FPs, being RFP, CFP, YFP, or GFP) upon Cre induction,

resulting in variability in Confetti expression pattern in mice that inversely correlates with precursor

numbers (Figure 1A) [11]. While suitable for developmental hematopoiesis, this method has limited

linear range (50-2500), impeding its potential application to adult hematopoiesis or clonally restricted

hematopoiesis. To measure hematopoietic precursors in various conditions, expansion of the

detection range is desired.

Interestingly, the random induction of Confetti colors in HSPCs bears similarity with X-chromosome

inactivation (XCI). In XCI, one of the X chromosomes is randomly inactivated in precursor cells, a

process that adheres to a binomial distribution. Formula derivation from binomial distribution implies

that a higher XCI variance among individuals correlates with a smaller number of precursors at the

time of inactivation (Figure 1B). As XCI is faithfully maintained in the progeny, this correlation has

been used to estimate precursor numbers from mature blood cells in human and mouse, leading to

the discovery of the first mutation (Tet2) that contributes to clonal hematopoiesis [7,12–14].

Nonetheless, XCI occurs exclusively in females, limiting its applicability in males. Additionally, XCI

takes place during early development when few precursors are present, resulting in low sensitivity in
adulthood [15].

Building on insights from Confetti mice and XCI studies, we investigated whether random induction
of Confetti FPs in precursor cells can be similarly modeled by a binomial distribution, whereby the
variance of FPs inversely correlates with precursor numbers. Based on this relationship, we asked if
we could broaden the measurable range of precursor number, which would allow us to expand the
existing correlation range.

Examining the premises of binomial distribution and setting some assumptions, we report that the
random induction of FPs among a group of mice or cells can be modeled by a binomial distribution.
Experimental validation establishes a broadened linear range, covering the full spectrum of
precursor numbers ($1\sim10^5$) and overcoming the prior range limit. We leverage this correlation to
probe the number of hematopoietic precursors at homeostasis, post-myeloablation, during
developmental expansion, and in a mouse model of inherited bone marrow failure.

<u>Results</u>
<u>Binomial distribution underlies the inverse linear correlation between FP% variance and the</u>
<u>precursor numbers</u>
To broaden the limited correlation range, we aimed to elucidate the mathematical relationship
between variance and precursor numbers [11]. Inspired by XCI, we asked how the stochastic
induction of Confetti FPs in precursors may satisfy the premises of binomial distribution [16,17]. We
found that when we assume the number of precursors remain the same for the condition tested, then
induction of FPs meets the required premises: (1) the number of precursors, denoted as *n*, remains
constant within a group of mice or cells; (2) each observation (e.g. each precursor) is independent;
(3) each observation yields one of two outcomes (e.g., it signifies the presence or absence of RFP in
a precursor); (4) the probability of success, represented as *p* (e.g., the likelihood of a precursor
expressing RFP upon Confetti induction), remains consistent across all observations.

Should the induction of a FP adhere to a binomial distribution, precursor numbers can be estimated
using the following equation:

$$\lg(\hat{n}) = \lg\left[\frac{(1-\widehat{FP_{mean}\%})}{\widehat{FP_{mean}\%}}\right] - 2 \times \lg\left(\widehat{CV_{FP}}\right) \qquad \text{(equation 1, see Methods)}$$

where $\hat{n}$ signifies the estimated number of precursors, $\widehat{FP_{mean}}\%$ represents the estimated probability

of a precursor being one given FP (e.g., $\widehat{RFP_{mean}}\%$), and $\widehat{CV_{FP}}$ denotes the estimated coefficient of

variance (CV) of one given FP (e.g., $\widehat{CV_{RFP}}$). CV is the standard deviation divided by the mean,

utilized by the empirical correlation formula.

The equation proposes a direct linear relationship between the precursor number and the CVs of

individual FPs, supporting the empirical correlation formula [11]. Indeed, by analyzing a published

dataset providing precursor numbers and the corresponding distribution of FPs, we found that the

correlation between n and individual CVs of FP exhibits superior fit (>0.93), higher than the empirical

correlation equation based on all three CVs (0.75) (Figure S1A) [11]. Moreover, according to

equation 1, the y-intercept of correlation between precursor number and CV was influenced by

$FP_{mean}\%$, while the slope was the same (-2) regardless of $FP_{mean}\%$. Given the unequal value of

$RFP_{mean}\%$, $CFP_{mean}\%$, and $YFP_{mean}\%$ in the published dataset, the correlation equations show

significantly distinct y-intercepts (p<0.0001) (Figure S1A and B). In contrast, the slopes of these

correlation equations were similar (p=0.49, with all 95% confidence intervals encompassing -2, the

theoretical value) (Figure S1A).

In conclusion, the FP induction in precursor cells can be modeled by a binomial distribution with the

assumption that precursor numbers are constant among a group of mice. This sets the mathematical

basis for the inverse linear correlation between variance of FP% and the number of precursors.

<u>A broad range of precursor numbers correlates with variance of FP%</u>

While the correlation formula derived from binomial distribution does not impose any range limitation,

experimental errors may confound the measurement of variance. We next aimed to experimentally

confirm this correlation. To streamline the calculation, we used standard deviation instead of CV to

compute variance. The correlation between standard deviation and precursor numbers is expressed

through the following equation:

$\lg(\hat{n}) = \lg\left[\widehat{FP_{mean}}\% \times (1 - \widehat{FP_{mean}}\%)\right] - 2 \times \lg(\widehat{\sigma_{FP}})$ (equation 2, see Methods)

where $\widehat{\sigma_{FP}}$ denotes the estimated standard deviation of a given FP% (e.g. $\widehat{\sigma_{RFP}}$).

Given this equation, $\lg(\widehat{\sigma_{FP}})$ exhibits a linear and inverse correlation with $\lg(\hat{n})$. Indeed, $\lg(\widehat{\sigma_{RFP}})$,

$\lg(\widehat{\sigma_{CFP}})$ and $\lg(\widehat{\sigma_{YFP}})$ exhibited a linear and inverse correlation with $\lg(\hat{n})$ in the published data

(Figure S1C) [11].

To simplify the validation, we used a two-color cell model HL-60 bearing one of the two FPs (BFP

and GFP, wherein GFP represents non-BFP) (Figure 1C). Although the Confetti cassette

recombination can generate one of four colors, our estimation is based on a single given FP (e.g. a

precursor expresses RFP or not), making this simplification justifiable.

We first proved that neither BFP nor GFP HL-60 cells had a competitive growth advantage over the

other, ensuring that HL-60 progeny mirrored the seeding population (Figure S1D). We next sorted

one to 10^7 cells into individual wells and allowed them to proliferate for at least three generations

before assessing BFP% at the end of the culture (Figure 1C). We focus on cell range up to 10^5 cells

as 1.4×10^5 non-HSC LSK and 5.2×10^3 active HSCs were previously estimated in an adult female

mouse, which corresponds to the highest possible precursor numbers in mice [18]. We considered

the survival rate of cells in wells receiving a single HL-60 cell (53% to 60%), to ensure the accuracy

of the number of cells seeded.

Consistent with the proposed correlation, as seeded numbers increase, BFP% standard deviation

decreases (Figure 1D and E). The variance of BFP% reveals an inverse correlation with the seeding

numbers ($R^2 = 0.996$) (Figure 1F). After normalizing for cell survival rates, the calculated numbers

closely align with the expected numbers from one to 10^6 cells (Figure 1G). We had anticipated that

measurement of small variance in high precursor numbers may be confounded by experimental

errors. Indeed, at cell numbers larger than 10^6, the standard deviation of BFP% and the expected

number reached a plateau (Figure 1G, grey circles). To ensure accurate measurement without

confounding errors, we suggest 10^5 to be the upper limit for experimental measurement.

<u>Experimental practices for accurate precursor number measurement</u>

It is crucial to ensure the accuracy of variance measured because variance of FP% is solely used for

estimation. Based on the data from the two-color HL-60 model, we conclude that there are at least

three experimental practices required for accurate cell estimates: (1) exclusion of outliers; (2)

sufficient flow cytometry recorded events; (3) sufficient sample size per group. Both outliers and

insufficient recorded events inaccurately inflate sample variance, leading to underestimation of

precursor numbers (Figure S1E and F). We found that the minimum recorded events increase in

tandem with the seeded number, in contrast to the reported 500-event threshold (Figure S1F) [11].

While having a small sample size per biological replicate is possible, the variability of estimates

among replicates would be high (Figure S1G). Practically, achieving a large sample size (>20) is

cumbersome in mice. To determine a feasible sample size for relatively accurate estimates, we

resampled datapoints to calculate the variability reduction as sample size increases (Figure S1H).

Our analyses reveal that five samples per group are sufficient to substantially minimize error,

whereas an additional increase in sample size leads to marginal error reduction. We therefore use at

least five mice per biological replicates for our precursor estimations.

Inducible HSC-Scl-CreER^T-targeted Confetti labeling in the active blood precursors at the adult

stage

To label hematopoietic precursors and quantify their number *in vivo*, we considered the choice of

Cre mouse line to be crossed with Confetti. Historically, hematopoietic precursors are thought to be

hematopoietic stem cells (HSCs) capable of long-term repopulation. However, recent studies

indicate that multipotent progenitors (MPPs) also contribute alongside HSCs in maintaining blood

production [19–22]. Hence, both HSCs and MPPs should be labeled.

Among Cre mouse lines capable of labeling HSCs and MPPs simultaneously at the adult stage

(such as Rosa26^{CreERT2}, Mx1-Cre, and HSC-Scl-CreER^T), we chose HSC-Scl-CreER^T because of its

preference for labeling HSPCs (immunophenotypically defined as Lin$^-$Sca-1$^+$cKit$^+$ (LSK)) [23]. To

validate the specificity of HSC-Scl-CreER^T, we generated mice possessing a single Confetti allele

and homozygous HSC-Scl-CreER^T alleles. We then examine Confetti expression one day after a

two-day tamoxifen administration. Consistent with prior reports of HSC-Scl-CreER^T activity, we

observed Confetti expression in the LSK population and T cells (Figure 2A, Figure S2A-E).

Unexpectedly, Confetti expression was additionally detected in NK cells, CD41$^+$ cells, non-

inflammatory monocytes (SSC-A^{low}Ly6C$^-$Ly6G$^-$CD11b$^+$), common myeloid progenitors (CMP, Lin$^-$

Sca-1$^-$cKit$^+$CD16/32$^{-/low}$CD34$^+$),   and   granulocyte-monocyte   progenitors   (GMP,   Lin$^-$Sca-1$^-$

cKit$^+$CD16/32$^+$CD34$^+$) [21] (Figure S2D-G). Considering the relatively shorter lifespan of these cells

compared to HSC/MPP, the labeling in these cells should minimally impact precursor calculations,

especially with a chase period post-labeling. While the Confetti induction in the relatively long-lived T

cells precludes estimation based on mature T cells, we conclude that HSC-Scl-CreER^T remains a

practical choice for labeling active hematopoietic precursors.

The stability of Confetti labeling after induction is required for binomial distribution. Therefore, the

absence of background Cre activity after Confetti induction is critical. We determine the background

Cre activity in HSC-Scl-CreER^T/Confetti animals by two approaches: (1) detecting Confetti

expression in non-induced animals; (2) identifying cells co-expressing two Confetti FPs months after

tamoxifen induction due to cassette "flipping" from background Cre activity. We observe no Confetti-

expressing cells without tamoxifen treatment in mice up to 50 weeks old and minimal co-expression

of two Confetti FPs in induced HSC-Scl-CreER^T/Confetti animals (Figure S3A-B). Conversely, Vav-

Cre/Confetti animals constitutively express Cre, resulting in ~15% cells co-expressing two Confetti
colors in peripheral blood (PB) T cells (Figure S3C-D). Both lines of evidence support minimal Cre
background activity in HSC-Scl-CreER^T and its use as a Cre driver for precursor number calculation.
218

<u>The variance of FP% inversely correlates with precursor numbers *in vivo*</u>
To ascertain the inverse correlation between variance of FP% and the number of hematopoietic
precursors *in vivo*, we first investigate the variance of FP% in mice hosting various numbers of
hematopoietic precursors. We generate these mice by non-competitively transplanting $0.25x10^6$ or
$4x10^6$ induced CD45.2$^+$ HSC-Scl-CreER^T/Confetti mouse bone marrow (BM) cells into CD45.1$^+$
recipient mice (Figure 2B). Given the constant frequency of precursors in the donor BM ($1/10^5$) and
the varying doses of donor BM cells, these mice received approximately 2.5 ("low precursor number")
or 40 ("high precursor number") precursors [16,24–26]. While similar experiments transplanting
defined numbers of precursors have been reported, the estimation for precursor numbers lower than
50 has not been explored.
229

Due to the scarcity of hematopoietic precursors in the BM, the precursor number seeded in
recipients followed Poisson distribution instead of being constant, violating the premise (1) of
binomial distribution. Nonetheless, our simulations show that the seeding number variation among
recipient mice is relatively small, therefore the variance resulting from precursor number differences
still dominates and inversely correlates with precursor numbers. A minor problem is that the
estimated numbers are not at one-to-one ratio to expected numbers at low precursor number range
(<10, see Methods) (Figure S4A).
237

238 As cells are not 100% labeled with Confetti, for clarity, we use "FP$_{PB}$%" or "FP$_{BM}$%" to represent the
239 frequency of a given FP in the PB or BM, and use "FP$_{Conf}$%" to represent the frequency of a given
240 FP in the Confetti$^+$ population. For total precursor number calculations, we normalize the number
241 calculated from the Confetti-labeled population to Confetti labeling efficiency.
242

As expected, we observe higher RFP$_{Conf}$% variance across all PB cell types in mice belonging to the
"low precursor number" groups (Figure 2C and Figure S4B-C). Since variance of any FP following a
binomial distribution inversely correlates with precursor numbers, we also find higher CFP$_{Conf}$% and
YFP$_{Conf}$% variance in the "low precursor number" groups (Figure S4D-E). Following the same
principle, we observe higher RFP$_{PB}$%, CFP$_{PB}$%, YFP$_{PB}$% and Confetti% variance in "low precursor
number" mice (Figure S4F).
249

Applying equation 2, we find that the precursor numbers of B cells and myeloid cells are noticeably
higher in the first two months than at four months post-transplantation, suggesting transient
progenitor contributions at early time points (Figure 2D). At four months post-transplantation, the
estimated precursor numbers align with the expected values (in myeloid cells, 94 for "high precursor
number" groups, three for "low precursor number" groups, Figure 2D). Again, for any FP following
binomial distribution, variance of FP% inversely correlates with precursor numbers and thus can be
leveraged to calculate precursor numbers (Methods). Since different FPs measure the same
population of precursors, we expect the estimations from different FPs to be similar. Indeed,
precursor numbers derived from RFP_{Conf}% highly correlate with those estimated from YFP_{Conf}% or
CFP_{Conf}% (R^2 = 0.885 for CFP% in Confetti$^+$, R^2 = 0.769 for YFP% in Confetti$^+$), as well as those
from RFP_{PB}%, YFP_{PB}% and CFP_{PB}% (Figure 2E, Figure S4G).
261
Consistent with PB data, higher variance of RFP_{Conf}% is also observed in the BM of "low precursor
number" mice (Figure S5A). The precursor numbers estimated from various BM HSPC
subpopulations align well with each other and are consistent within the same group (Figure 2F).
However, in "high precursor number" groups, estimates from BM HSPCs are lower than those
derived from PB myeloid cells (27 for BM HSPC, 94 for PB myeloid cells, Figure 2F). This
discrepancy may reflect uneven seeding of precursors to the BM throughout the body after
transplantation, and the fact that we only sample a part of the BM (femur, tibia, and pelvis)[27]. In
summary, we validate the inverse correlation between variance of FP% and precursor numbers *in
vivo*. Moreover, the quantification of precursor numbers is feasible even for precursor numbers
outside the empirical correlation range.
272
<u>Cell frequency measurements fail to reflect the differences in precursor numbers</u>
Given that transplantation studies are performed non-competitively and the recipients are lethally
irradiated, we expect to see minimal differences in donor chimerism between two groups, despite
drastic differences in donor precursor numbers. Indeed, although we observe substantially lower
donor chimerism in mice belonging to "low precursor number" groups during the first two months
post-transplantation, the chimerism differences are very small by four months post-transplantation
(5.5±2.8%, Figure 2G). The initial differences in donor precursor numbers do not affect PB cell
frequencies, except for the first month post-transplantation, when higher PB myeloid frequencies are
observed in the "low precursor number" groups (Figure 2H). This suggests myeloid cell production is
enhanced when very few precursors are available after irradiation-mediated injury (Figure 2C).
283
In the BM, nucleated cell counts, donor chimerism and HSPC frequencies are mostly similar
regardless of donor precursor numbers, suggesting that even extremely low numbers of

hematopoietic precursors can still effectively repopulate a non-competitive environment (Figure S5B-
D). Therefore, in cases where hematopoietic precursors expand or decline without competition, stem
cell frequencies are less informative to study precursor activity. Measuring precursor numbers is
more meaningful as low number of active precursors may be constrained by compensatory
proliferation.

<u>Thousands of hematopoietic precursors contribute to native hematopoiesis</u>
While the empirical formula, which measures 50-2500 precursors, supports quantification of
precursors labeled at fetal stages, it may not be applicable to native hematopoiesis, when the total
precursor number may exceed 2500. Having validated the feasibility for measuring precursor
numbers *in vivo* by Confetti animals, we next sought to investigate the number of active precursors
contributing to native hematopoiesis. In HSC-Scl-CreER^T/Confetti animals, we opt to label a smaller
portion of HSPCs with a two-day treatment, despite the potential to label up to 60% HSPCs with
Confetti (CFP+YFP+RFP) by 14-day tamoxifen treatment (Figure 3A, Figure S6A). This
precautionary measure aims to mitigate potential toxicity arising from prolonged tamoxifen treatment.

Unexpectedly, we observed a successive decrease of CFP$_{Conf}$% and CFP$_{PB}$% in some animals,
which distort the distribution of CFP$_{Conf}$% (Figure S6B and C). A similar decline is not observed in
RFP$_{Conf}$% or YFP$_{Conf}$%, nor in RFP$_{RFP+YFP}$% (calculated by RFP$_{PB}$%/(RFP$_{PB}$%+YFP$_{PB}$%)) (Figure
S6D and E). A similar loss of CFP$_{Conf}$% is neither observed in animals with Confetti induction during
fetal development, suggesting a potential immune response to CFP in adult-induced animals (Figure
S6F). To circumvent this caveat, we decided to calculate precursor numbers solely based on the
variance of RFP$_{RFP+YFP}$%. Since only ~2% of myeloid cells are labeled with CFP, we reason that the
decline of CFP in some animals is unlikely to affect the calculation of total precursor numbers based
on other FPs, such as RFP$_{RFP+YFP}$%. Moreover, in transplantation studies, precursor numbers
calculated with variance of RFP$_{RFP+YFP}$% linearly correlate with those calculated using variance of
other FPs (Figure S4G).

To further ensure accuracy, we focus on PB myeloid cells, as (1) Confetti labeling of T cells result
from direct induction by HSC-Scl-CreER^T, but not differentiation from HSPCs (Figure S2D); (2)
Confetti labeling (RFP and YFP) of B cells does not plateau at seven months post-induction (Figure
3B); (3) myeloid cells are of shorter lifespan. In myeloid cells, we focus on precursor numbers after
four months post-induction, when their labeling reaches a plateau (Figure 3B). The stability of
average RFP$_{RFP+YFP}$% suggests an equal contribution of RFP$^+$ and YFP$^+$ cells to the PB, a
prerequisite for estimating precursor numbers from FP% distributions in the progenies (the myeloid

cells) (Figure 3C). Like those in PB, the average $RFP_{RFP+YFP}$% in BM HSPC subpopulations is also

stable, supporting non-biased differentiation of RFP^+ and YFP^+ precursor cells (Figure 3C).

Fitting data to equation 2, we estimate an average of 2667 precursors contributing to native

myelopoiesis (Figure 3D, average of numbers at five to seven months post-induction). This number

closely aligns with the clone number estimated by transposon-based barcodes statistic in

granulocytes (831/30% = 2770) [1].

The average precursor number calculated from PB myeloid cells at the time of BM analysis (1958)

matches those calculated from BM myeloid progenitors (MP, $Lin^-Sca-1^-cKit^+$) and HSPCs (1773 and

1917), but it is five-fold higher than that of LT-HSC (Figure 3E). Although LT-HSCs are thought to

sustain steady-state hematopoiesis, recent studies suggest that ST-HSC and MPPs persist long-

term as primary contributors to adult hematopoiesis [1,19,21,28]. The observed discrepancy

between the number of precursors contributing to LT-HSC and those contributing to MPPs indicates

that at least some MPPs are not replaced by progenitors directly differentiated from LT-HSC at

seven months post-induction, affirming the persistence of MPPs (Figure 3E). Furthermore, the fact

that PB myeloid and BM MP precursor numbers are closer to those of MPPs than LT-HSC confirms

active and long-term (at least seven months) contributions from MPPs to steady-state myelopoiesis.

Of note, the quantity of precursor does not necessarily correlate with the frequency of cell type in the

BM. For example, the cell frequency of MPP2 ($Lineage^-cKit^+Sca-1^+Flk2^-CD48^+CD150^+$) in BM is very

low, but MPP2 comprises a modest number of precursors, making its frequency-to-precursor-

number ratio the lowest among the HSPC subtypes (Figure 3F and S6G). The frequency-to-

precursor-number ratio reflects how well precursors of a particular type proliferate to expand their

absolute cell count. The low frequency-to-precursor-number ratio of MPP2 suggests that it expands

more poorly than other HSPC subtypes.

In summary, we detect thousands of hematopoietic precursors contributing to adult hematopoiesis.

At the time of BM analysis, the number of PB myeloid precursors is comparable to those observed in

BM MP and HSPCs.

<u>Precursor numbers determined by FP% variance confirm reduced clonality of progenitors after</u>

<u>myeloablation</u>

Myeloablation through 5-fluorouracil (5-FU) treatment depletes most actively cycling cells, forcing

quiescent stem cells to proliferate [29]. Previous studies have indicated a significant reduction in the

number of clones detected within the BM c-Kit$^+$ population (consisting of MP and HSPC) following a

single dose of 5-FU treatment [9]. However, questions arise regarding whether this is an artifact

stemming from potential under-sampling of the highly expanded c-Kit$^+$ population after 5-FU

treatment by single-cell sequencing. Given the quantitative estimation of a wide range of precursor

numbers through equation 2, we aimed to investigate whether precursor numbers within progenitor

populations indeed reduce ten days post-5-FU treatment (Figure 4A).

To determine the precursor changes post-5-FU treatment, we use the animals described in Figure

3A as untreated (UT) benchmark cohort, which were collected at the same age. At ten days post

injection, the efficacy of 5-FU treatment is validated by lower PB myeloid cell frequency post-

treatment and higher frequencies of BM progenitors compared to UT animals (Figure S7A and B).

The high progenitor frequencies in the BM results from the depletion of cycling cells in the BM and

the enhanced proliferation of HSPCs following 5-FU treatment.

Compared to UT animals, the precursor numbers of MP (including CMP, GMP, MEP) and HSPC

significantly decrease, confirming a reduction in clonality in the c-Kit$^+$ population (Figure 4B) [9]. By

contrast, the precursor numbers of LT-HSC do not show a decreasing trend. While transplantation

studies support unchanged clonality of primitive stem cells after a single dose of 5-FU, a similar

investigation in a native environment has not been conducted [30]. Our findings support the notion

that native LT-HSC clonality remains unaltered following one-dose 5-FU treatment. Together,

myeloablation treatment reinforces how the dynamics of precursor numbers can be tracked through

Confetti pattern variations.

<u>Modest developmental expansion of active lifelong precursors</u>

HSCs are thought to undergo substantial expansion in the fetal liver during fetal development [31–

33]. However, a recent study employing the empirical formula challenges this notion by quantifying

endogenous lifelong hematopoietic precursors labeled at various developmental stages [34]. It

revealed limited expansion of hematopoietic precursors during the fetal liver stage (from E10 to

E15,1.8- to 2.7-fold) as well as a gradual and moderate increase from the fetal liver to the post-natal

stage (2.4- to 10-fold) [34]. Although intriguing, most post-natal measurements in this study fell

outside the empirical linear range, leaving the genuine degree of post-natal precursor expansion

uncertain. As our formula quantitatively assesses a wide range of precursor numbers, we set out to

directly compare the numbers of precursor labeled at various development stages (E11.5 and E14.5,

by one dose of tamoxifen; adult-stage, benchmark shown in in Figure 3A, all analyzed at the same

age) (Figure 4C).

For accuracy, we first examine the dynamics of Confetti labeling. In E14.5-induced animals, the

Confetti labeling of T cells almost double from one month to four months of age (4.7% at one month,

9.3% at four months), suggesting Confetti$^+$-labeled precursors contribute more to post-natal T cells

than non-labeled precursors (Figure 4D). Therefore, for T cells in E14.5-induced animals, we focus

on precursor numbers generated after three months of age. The Confetti labeling for E11.5-induced

animals remain stable for all timepoints examined (Figure S7C).

Similar to adult-induction, the distribution of average FP% is stable for fetal-induction, supporting

equal proliferation of RFP$^+$ and YFP$^+$ cells (Figure S7D). Unlike adult-induction, fetal-labeled

precursor numbers can be calculated with various combination of FP%, since the expression of CFP

introduced at the fetal stages is stable (Figure S6F). Nonetheless, for consistency, all precursor

numbers are calculated with variance of RFP$_{RFP+YFP}$%, the one employed in adult-induction animals.

The resulting precursor number estimates for E11.5 and E14.5-labeled cohorts are similar in all PB

cell types except at two and three months of age, echoing the previous study reporting limited

lifelong precursor expansion in the fetal liver (Figure 4E) [34]. Precursor numbers calculated from

BM subpopulations are also comparable between the two timepoints (Figure 4F). Although E9.5-

labeled clones are reported to seed non-uniformly across bones, the precursor numbers calculated

from PB myeloid cells do not significantly differ from those calculated from BM MP and HSPC in

E14.5-labeled animals (Figure 4F) [9].

For comparison between fetal- and adult-induced animals, we focus on PB myeloid cells, as adult-

induced animals have non-saturated labeling in B cells and non-HSPC-rooted Confetti labeling in T

cells (Figure 3B). Here, we observe a relatively small increase of precursor numbers in adult-induced

animals in PB and BM compared to those labeled at the fetal stage (Figure 4E and F). The increase

between E11.5/E14.5- and adult-labeled precursors is less than ten-fold, similar to a previous report

(Figure 4G) [34].

Together, we confirm minimal to no expansion of lifelong precursors in the fetal liver stage and a

minor expansion of expansion of lifelong precursors from fetal liver to adult stage.

Fancc$^{-/-}$ mice have normal numbers of hematopoietic precursors at steady-state

Our approach provides an opportunity to investigate the number of active HSPC precursors in the

native environment. To showcase precursor quantification in genetic mouse models, we focus on

Fanconi Anemia (FA), the most common inherited bone marrow failure syndrome [35]. Current FA

mouse models exhibit mostly normal adult hematopoiesis at steady state but demonstrate reduced

repopulation ability upon BM transplantation [36–41]. It remains unclear if their steady-state blood

production is sustained by a reduced number of precursors, predisposing them to repopulation

defects after transplantation.

To quantify precursor number in a mouse model of FA, we generate $Fancc^{+/+}$, $Fancc^{+/-}$ and $Fancc^{-/-}$

mice with a single Confetti allele and homozygous HSC-Scl-CreER^T alleles (Fancc$^{+/+}$Confetti$^{fl/+}$HSC-

Scl-CreER$^{T/T}$, hereafter $Fancc^{+/+}$; Fancc$^{+/-}$Confetti$^{fl/+}$HSC-Scl-CreER$^{T/T}$, hereafter $Fancc^{+/-}$; Fancc$^{-/-}$

Confetti$^{fl/+}$HSC-Scl-CreER$^{T/T}$, hereafter $Fancc^{-/-}$) [42]. Consistent with previous literature, we observe

similar PB cell frequencies and blood counts between $Fancc^{+/+}$ and $Fancc^{-/-}$ mice (Figure S8A-C) [41].

BM nucleated cell counts, as well as HSPC and MP frequencies, are mostly identical, except for ST-

HSC frequencies, which are significantly reduced in $Fancc^{-/-}$ mice compared to $Fancc^{+/-}$ mice (Figure

S8D-E).

To label hematopoietic precursors in FA animals, Confetti expression was induced at two months of

age, and FP% was monitored over seven months (Figure 5A). The absence of $Fancc$ did not affect

Confetti labeling efficiency, Confetti labeling dynamics or average FP% (RFP$_{RFP+YFP}$%), suggesting

precursor numbers in $Fancc^{-/-}$ can be similarly calculated (Figure 5B and Figure S8F). As in previous

adult-induction animals, RFP$_{RFP+YFP}$% was used to estimate precursor numbers. The resulting

myeloid and BM precursor numbers were comparable regardless of $Fancc$ genotype, albeit $Fancc^{-/-}$

mice had a slight reduction in HSPC precursor numbers (Figure 5C and D). These observations

collectively suggest that a normal number of precursors sustains blood production in $Fancc^{-/-}$ mice.

The number of $Fancc^{-/-}$ precursors remains unchanged in mice transplanted with young donor cells

While $Fancc^{-/-}$ mice have a similar number of precursors as their wildtype counterparts at

homeostasis, it is unknown whether the reduced repopulation ability post-transplantation stems from

diminished precursor numbers, reduced cell expansion, or a combination of both [37]. To understand

the underlying mechanism, we performed competitive transplantation using BM cells from three-

month-old Confetti-induced $Fancc^{+/+}$, $Fancc^{+/-}$ or $Fancc^{-/-}$ mice along with CD45.1$^+$ competitor cells

(Figure 5E, Figure S9A). Consistent with previous studies, recipient mice of $Fancc^{-/-}$ cells show

significantly lower donor chimerism in the PB and BM compared to those of $Fancc^{+/+}$ or $Fancc^{+/-}$ cells

(Figure 5F, Figure S9B). Despite lower donor chimerism, PB and BM precursor numbers were

unaffected in $Fancc^{-/-}$ recipient mice (Figure 5G, Figure S9C-E). This suggests that reduced $Fancc^{-/-}$

cell proliferation instead of fewer active precursors is likely the cause for the reduced repopulation

capacity post-transplantation.

Ageing $Fancc^{-/-}$ mice have reduced lymphoid hematopoietic precursors upon transplantation

Ageing *Fancc*[-/-] mice have been reported to develop hematologic neoplasms, resulting in decreased

survival [43]. To determine if aging *Fancc*[-/-] cells also maintain similar precursor numbers post-

transplantation, we competitively transplanted BM cells from nine-month-old Confetti-induced

Fancc[+/+] or *Fancc*[-/-] mice with CD45.1[+] competitor cells, a stage when aging *Fancc*[-/-] mice start to

show decreased survival [43] (Figure S9F).

The diminished repopulation ability of *Fancc*[-/-] cells is reaffirmed by lower PB and BM chimerism

(Figure 5H, Figure S9I). While no differences in precursor numbers are noted during the initial post-

transplantation period, a slight yet consistent reduction in lymphoid precursors is observed in *Fancc*[-/-]

recipient mice at three to five months post-transplantation (Figure 5I, Figure S9G and H). Changes in

precursor numbers in the BM of KO recipients are less clear, as we observed high variance in

several HSPC subtypes (Figure S9J). Nonetheless, precursor numbers of MEP, LSK and ST-HSC

showed a consistent reduction. In conclusion, aging *Fancc*[-/-] mice showed a modest but consistent

loss of active PB lymphoid precursors post-transplantation, implying decreased lymphoid precursors

additionally compromise repopulation capacity as *Fancc*[-/-] mice age.

Discussion

The polyclonal nature of endogenous hematopoiesis imposes methodological problems on a robust

dynamic measurement. Inspired by the XCI studies, we successfully employed the correlation

formula informed by binominal distribution to quantify precursor number in native hematopoiesis.

Using HSC-Scl-CreER[T]/Confetti animals, we estimate thousands of precursors contributing to adult

native hematopoiesis, a number comparable to the previous report [1]. This number results from a

moderate increase during fetal-to-adult transition and respond dynamically to myeloablation by 5-FU.

Applied to a mouse model of inherited bone marrow failure (Fancc[-/-] mice), we detected normal

precursor numbers at steady state and decreased lymphoid precursors upon transplantation of aging

donors.

Although the linear relationship between variance of FP% and precursor numbers has been

described, its measurable range has been limited [11]. Estimates outside this linear range may

erroneously fall within it, making it challenging to distinguish accurate measurements within the

range from inaccurate ones beyond it [11]. Based on binomial distribution, we expanded the

measurable range to encompass the full spectrum of hematopoietic precursors in mice, minimizing

the likelihood of inaccuracies. Furthermore, the linear correlation based on binomial distribution can

accommodate any labeling system that follows the underlying premises, without the need for the

multispectral Confetti cassette.

Having established a quantitative measurement, we were initially surprised by the moderate
differences between fetal and adult-precursors [34]. The maximum increase observed (seven-fold
observed in CMP) was substantially lower than the increase of competitive repopulation units (CRU)
from E12 to E16 determined by limited dilution assays (~38-fold) (Figure 4G), as well as the
escalation from immunophenotype-defined HSPC counts (28- to 47-fold, from 3-5x10^3 at E14.5 to
1.4x10^5 at two months old) (Figure 4G) [18,32,44,45]. It is possible that we overestimated E11.5
precursors, as the numbers of lifelong E11.5 precursors were substantially higher (on average 753
PB myeloid precursor, 512 HSPC precursors) than the one to two repopulation units estimated with
transplantation [46]. Nonetheless, our results were comparable to the ~870 hematopoietic cluster
cells observed at E11.5 [47]. We may have underestimated precursor numbers in adult-labeled
animals, as most adult HSCs are quiescent [29]. However, HSC precursor numbers did not increase
after induced proliferation by one dose 5-FU treatment (Figure 4B). Therefore, we confirm the
moderate precursor numbers difference between fetal- and adult-stage, emphasizing the analysis of
hematopoiesis in a native environment. Future studies using different methods to investigate
precursor activity locally should validate this result.

Novel methods tracing clones *in situ* offer new opportunities to study native hematopoiesis, yet most
are challenging to apply in genetic models [1,9,10,27,48]. Since only two mouse lines are required
(Confetti and Cre, sometimes Cre is already included for conditional deletion of alleles), our
approach is particularly convenient to study hematopoiesis in mouse models of genetic disorders.
Applying precursor measurements to *Fancc$^{-/-}$* mice, we observe a minor decrease in precursor
numbers after transplantation of aging donors. Although donor chimerism differences had been
linked to differences in precursor numbers, reduced proliferation capacity may also contribute to
reduced competitive repopulation capacity [26]. In this case, definitive precursor number analysis is
necessary to differentiate these possibilities.

Currently, concerns regarding clonal restriction in the context of FA gene therapy arises, as the
engraftment of gene-corrected stem cells has yielded marginal clinical benefits [49]. For the first time,
we demonstrate that FA cells maintain a normal precursor quantity post-transplantation, thereby
disproving clonal attrition, including putative homing deficits post-transplantation as a cause in this
model [50]. However, it is imperative to acknowledge that the majority of murine models of FA,
including the *Fancc$^{-/-}$* model utilized herein, do not recapitulate the pathophysiology observed in FA
patients. Future investigations in other FA mice and studies leveraging FA patient-derived materials
will be pivotal in corroborating and validating the findings presented here.

While we implemented a careful data processing procedure, one pitfall of our analyses was that we

inferred precursor numbers from their progenies, assuming uniform and linear expansion from

precursor to progenies. A recent study showed non-uniform precursor clone sizes, although the level

of non-uniformity is low [27]. In cases where the non-uniformity is high, according to mathematical

modeling, we measured the major contributors to hematopoiesis[51]. Another potential caveat is that

the relative contribution of Confetti-labeled precursors to blood production compared to non-labeled

precursors remains unknown. Future studies using different Cre drivers should validate the

precursor numbers for steady-state hematopoiesis.

In summary, we substantially broadened the applicable range of the correlation between variance of

FP% and the precursor based on binomial distribution. We discovered thousands of precursors

contributing to steady-state adult murine hematopoiesis and validated that fetal-to-adult precursor

expansion is indeed limited. This analysis highlights active precursor numbers as an important

metric in both normal and genetic mouse models.

<u>Methods</u>

<u>Variance modeling with two-color HL-60</u>

HL-60 cells were cultured with IMDM containing 20% FBS (Gemini) and 5% Penicillin-Streptomycin

(Gibco) and maintained at 1 x 10^5 and 1 x 10^6 cells/ml. Mycoplasma tests (Lonza) were performed

routinely to rule out mycoplasma contamination. BFP-HL-60 and GFP-HL-60 were generated with

lentivirus transduction of pGK-BFP (Genscript) and LeGO-V2 (a gift from Dr. Stefano Rivella lab).

For seeding of one to 10,000 cells, sorting of HL-60 mixtures was performed using a BD FACS Aria

III. For seeding of 100,000 cells, cell counts and dilution was used. After expansion, HL-60 cells

were fixed with BD fixation buffer before Aurora (Cytek) or Cytoflex (Beckman Coulter) analysis.

<u>Mice</u>

HSC-Scl-CreER^T mice [23] were crossed with R26R-Confetti mice (B6.129P2-

Gt(ROSA)26Sor$^{tm1(CAG-Brainbow2.1)Cle}$/J) to generate HSC-Scl-CreER$^{T/T}$Confetti$^{fl/+}$ (HSC-Scl-

CreER^T/Confetti) animals. HSC-Scl-CreER^T/Confetti animals were crossed with *Fancc$^{+/-}$* mice [42] to

generate HSC-Scl-CreER$^{T/T}$Confetti$^{fl/+}$*Fancc$^{+/+}$* (*Fancc$^{+/+}$*), HSC-Scl-CreER$^{T/T}$Confetti$^{fl/+}$*Fancc$^{+/-}$*

(*Fancc$^{+/-}$*), and HSC-Scl-CreER$^{T/T}$Confetti$^{fl/+}$Fancc$^{-/-}$ (*Fancc$^{-/-}$*) mice. Vav-Cre was generously offered

by Dr. Wei Tong (Children's Hospital of Philadelphia). Vav-Cre/Confetti animals were generated by

crossing Vav-Cre with R26R-Confetti mice. All animals are in the B6 (CD45.2) strain background,

unless otherwise stated. Six- to twelve-week-old females were used for timed pregnancies. Eight-

week-old mice were used for adult Confetti induction. Both female and male mice were used [11].

Six- to eight-week-old old female B6 CD45.1 mice (B6.SJL-*Ptprc^a Pepc^b*/BoyJ, Jackson laboratories)

were used as competitors and recipients for transplantation studies. All mice were maintained in the

conventional small animal facility at the Children's Hospital of Philadelphia (CHOP). All procedures

involving animals were approved by the Institutional Animal Care and Use Committee at the

Children's Hospital of Philadelphia.

<u>Animal identification</u>

Tail snip DNA was extracted using KAPA Express Extract Kit (Roche). Genotyping PCR was

performed with HotStarTaq Master Mix (Qiagen) according to manufacturer's instruction. Genotyping

primers used are summarized in Table S3. To determine the zygosity of HSC-Scl-CreER^T, qPCR

was additionally performed with purified tail snip DNA using SYBR™ Green Universal Master Mix

(Applied Biosystems).

<u>Animal procedures</u>

For fetal induction, timed matings of HSC-Scl-CreER$^{T/T}$Confetti$^{fl/fl}$ mice and HSC-Scl-CreER$^{T/T}$ mice

were set up. The mice were separated the next morning and noon of the day of separation was

considered E0.5. Tamoxifen was delivered at 100mg/kg to the dam orally at E11.5 or E14.5. Pups

were C-sectioned and cross-fostered at E18.5 due to reported delivery difficulties caused by

tamoxifen treatment [52]. For mice used for defined ("low" versus "high") number of transplantation

(Figure 2), tamoxifen was delivered at 70mg/kg orally once per day for 14 days at eight weeks old.

For adult induction (Figure 3), tamoxifen was delivered at 70mg/kg orally once per day for two days

at eight-week-old. For one dose 5-fluorouracil (5-FU, Sigma) treatment, 5-FU was intraperitoneally

injected once ten days before BM harvest (37-week-old) at 150 mg/kg. To obtain peripheral blood for

Confetti analysis, mice were anesthetized using isoflurane and retro-orbitally bled or

submandibularly bled for *Fancc$^{-/-}$* mice with occasional congenital eye defects[53] for 1 capillary of

blood (50ul). For peripheral blood counts, blood was collected in EDTA tubes using similar bleeding

methods and was analyzed by the Translational Core Lab (Children's Hospital of Philadelphia).

<u>Bone marrow sample processing</u>

To collect BM cells, mice were euthanized by CO$_2$ inhalation. Tibia, femur and pelvis were dissected,

and the bone marrow cells were flushed with 26-gauge needles. The single-cell suspension

generated with 18-gauge needles was then filtered through 70μm strainers. Erythrocytes in the bone

marrow cells were hemolyzed by RBC lysis buffer before antibody staining or transplantation. For

stem cell enrichment, bone marrow cells were further lineage depleted using EasySep™ Mouse

Hematopoietic Progenitor Cell Isolation Kit (STEMCELL Technologies).

<u>Flow cytometry analysis</u>

Peripheral blood and bone marrow cells were analyzed using Aurora (Cytek) or BD FACS Aria III

and the flow cytometry data were analyzed using FlowJo (Tree Star). The combinations of the

following cell surface markers were used to define the peripheral blood populations: myeloid cells:

CD11b$^+$ or Gr-1$^+$; T-cell: CD3ε$^+$; B-cell: B220$^+$. The following combinations of cell surface markers

were used to define the bone marrow stem and progenitor cells (Lineage/Lin: CD11b, Gr-1, B220,

CD3ε,   Ter119):   LTHSC: Lin$^-$c-Kit$^+$Sca1$^+$Flk2$^-$CD150$^+$CD48$^-$;   MPP2:   Lin$^-$c-Kit$^+$Sca1$^+$Flk2$^-$

CD150$^+$CD48$^{+;}$ MPP3: Lin$^-$c-Kit$^+$Sca1$^+$Flk2$^-$CD150$^-$CD48$^+$; MPP4: Lin$^-$c-Kit$^+$Sca1$^+$Flk2$^+$CD150$^-$CD48$^+$;

STHSC: Lin$^-$c-Kit$^+$Sca1$^+$Flk2$^-$CD150$^-$CD48$^-$; MEP: Lin$^-$c-Kit$^+$Sca1$^-$CD34$^-$CD16/32$^-$; CMP: Lin$^-$c-

Kit$^+$Sca1$^-$CD34^{mid}CD16/32^{mid}; GMP: Lin$^-$c-Kit$^+$Sca1$^-$CD34$^+$CD16/32$^+$. For bone marrow stem and

progenitor cell analysis, DAPI (Biolegend) was used to distinguish dead cells. Representative

examples of flow cytometry gating can be found in Figure S2C. The antibodies were used at

optimized dilutions listed in Table S1.

<u>Bone marrow transplantation</u>

The day before transplantation, female CD45.1 recipient mice were lethally irradiated (5.2Gy x 2, 3h

apart) using an X-ray irradiator (Precision). On the day of transplantation, bone marrow cells from

donor mice (CD45.2$^+$) were collected under sterile conditions as described, RBC-lysed and counted

for cell number. For Figure 2 (transplantation of defined precursor numbers), donor mice were

induced to express Confetti as described; donor bone marrow cells were (non-competitively) injected

into irradiated recipient mice; each matched high- and low- precursor number group received donor

bone marrow cells pooled from three to five mice. For transplantation of young *Fancc* mice, donor

mice were induced to express Confetti at E14.5; 1.5 x 10^6 donor bone marrow cells were mixed with

2.5 x 10^5 CD45.1 supporting bone marrow cells and injected into irradiated recipient mice via tail

vein. For transplantation of aging Fancc mice, donor mice were induced to express Confetti at two-

months of age; 2 x 10^6 donor bone marrow cells were mixed with 5 x 10^5 CD45.1 supporting bone

marrow cells.

<u>Derivation of the correlation between variance of confetti FP% and precursor numbers based on</u>

<u>binomial distribution</u>

When a random variable adheres to binomial distribution, studies have established that the following

equation holds true:

$$n = \frac{p(1-p)}{\sigma^2}$$

638   where *n* signifies the number of precursors, *p* represents the probability of an individual being one of

639   the FPs (e.g., RFP), and σ^2 denotes the variance of specific FP% (e.g. σ$_{RFP\%}$^{2})[7]. In experiments, *p*

640   will be estimated with the average FP% in the Confetti$^+$ cells (e.g. RFP$_{mean}$%), and σ^2 will be

641 estimated with the variance of FP% among a group of individual or mice (e.g. $\widehat{\sigma_{RFP}}^2$). Consequently,

642 the estimation of precursors number n is calculated using the following equation:

$$\hat{n} = \frac{\widehat{FP_{mean}}\% \times (1 - \widehat{FP_{mean}}\%)}{\widehat{\sigma_{FP}}^2}$$

643 A logarithmic transformation can then be performed, allowing us to establish a linear relationship

644 between the variance of FP% and the number of precursors:

645 $$\lg(\hat{n}) = \lg\left[\widehat{FP_{mean}}\% \times (1 - \widehat{FP_{mean}}\%)\right] - 2 \times \lg(\widehat{\sigma_{FP}}) \quad \text{(equation 2)}$$

647 In a previous study[11],

$$\sigma_{FP} = CV_{FP} \times FP_{mean}\%$$

Therefore,

$$\lg(\hat{n}) = \lg\left[\frac{(1 - \widehat{FP_{mean}}\%)}{\widehat{FP_{mean}}\%}\right] - 2 \times \lg(\widehat{CV_{FP}}) \text{ (equation 1)}$$

<u>Resampling to determine sample size per replicate</u>

FP% data generated from HL-60 was used for re-sampling. For each seeded number, FP% was

resampled for different sample sizes from all the FP% with replacement. Variance of estimated n

was then calculated by the standard deviation of the estimated precursor numbers generated from

the same sample size. Relative error was determined by dividing variance of estimated n with the

average of the estimated n. Refer to "varying well numbers.rmd" for detailed R code.

<u>Simulation to determine the effect of varying FP$_{mean}$% on correlation between variance of FP% and</u>

<u>precursor numbers</u>

Simulation of binomial distribution of varying FP$_{mean}$% (probability of being a FP) was performed in R,

generating corresponding FP% values used for variance calculation. The correlation between

variance and precursor number n was then determined by linear correlation, and the slopes and

intercepts were compared to each other. Refer to "Simulation of varying FPmean_percent.rmd" for

detailed R code.

<u>Simulation to determine the effect of precursor numbers following Poisson post-transplantation</u>

Precursor numbers in individual samples were simulated to follow a Poisson distribution, where the

mean of precursor numbers is the expected precursor numbers (expected n). The induction of a FP

in each precursor was then simulated by random assignment, where probability of being a FP was

set to be FP$_{mean}$%. For each expected precursor numbers, variance of FP% among samples was

then calculated, and equation 2 was used to estimate precursor numbers (estimated n). The

correlation between expected n and estimated n was then plotted. Refer to "Double layer binomial

simulation.rmd" for detailed R code.

<u>Data processing and normalization</u>

Peripheral blood cell subset frequencies were normalized to the total % of myeloid cells, T-cell and

B-cell, to avoid an underestimate due to incomplete RBC lysis. For transplant animals, peripheral

blood $CD45.1^+$% and $CD45.2^+$% are normalized to total $CD45.1^+$% and $CD45.2^+$% to avoid an

underestimate due to incomplete RBC lysis. After normalization of cell frequencies, the sum of

Confetti% and FP_{Conf}% (diving FP_{PB}% or FP_{BM}% by sum of Confetti%) were then calculated for each

cohort. At each step, potential outliers were removed based on Tukey method. The variance of

FP_{Conf}% and average FP_{Conf}% are then fitted into equation 2 to calculate precursor numbers if

sample size is at least five. The precursor number is then used to compared with the minimum flow

cytometry recorded event of samples. If the estimated number is higher than the minimum flow count

of samples, the sample with the minimum flow count will be excluded, and the calculation is

performed again on the rest of samples. After exclusion, the resulting precursor number will be

compared again with minimum flow count of samples, until it is lower than the minimum flow count or

sample size is smaller than five.

<u>Statistical analysis</u>

<u>Statistical significance for precursor number estimates</u>

As the distribution of precursor numbers is not predetermined, to compare mean precursor number

differences between two conditions from a limited number of biological replicates, we employed

permutation test. For unpaired permutation test, if there are three to five biological replicates per

condition, the smallest p values possible are 0.05 ~ 0.004 ($1/\binom{6}{3}$ ~ $1/\binom{5}{10}$). For paired permutation

test, if there are three to five biological replicates per condition, the smallest p value possible are

0.125 ~ 0.03125 ($1/2^3$ ~ $1/2^5$). Thus, for three to five biological replicates per condition, even though

some of the p values may not reach the commonly used alpha level (0.05), it still represents

substantial number differences. For those p values that were lowest possible but did not reach the

alpha level, we specifically labeled with a "#" in the figures and legends.

<u>Statistical analyses of other data</u>

All other two-sample statistical analyses were performed using Student's t test, if the sample was

normally distributed, or Welch's t test, when the sample was not normally distributed (F test). For

multiple comparison, one-way ANOVA or two-way ANOVA was used.

<u>Data and code availability</u>

The original Confetti% measured in the peripheral blood and the bone marrow and R code to
analyze and reproduce all the results, numeric and figures can be found at
https://doi.org/10.5281/zenodo.8222789. The re-analyzed data from a previous study can be found
in the online version of papers[11].

References

1. Sun J, Ramos A, Chapman B, Johnnidis JB, Le L, Ho Y-J, et al. Clonal dynamics of native
haematopoiesis. Nature. 2014;514: 322–327. doi:10.1038/nature13824

2. Yamamoto R, Morita Y, Ooehara J, Hamanaka S, Onodera M, Rudolph KL, et al. Clonal Analysis
Unveils Self-Renewing Lineage-Restricted Progenitors Generated Directly from Hematopoietic Stem
Cells. Cell. 2013;154: 1112–1126. doi:10.1016/j.cell.2013.08.007

3. Lee-Six H, Øbro NF, Shepherd MS, Grossmann S, Dawson K, Belmonte M, et al. Population
dynamics of normal human blood inferred from somatic mutations. Nature. 2018;561: 473–478.
doi:10.1038/s41586-018-0497-0

4. Mitchell E, Chapman MS, Williams N, Dawson KJ, Mende N, Calderbank EF, et al. Clonal
dynamics of haematopoiesis across the human lifespan. Nature. 2022;606: 343–350.
doi:10.1038/s41586-022-04786-y

5. Breivik H. Haematopoietic stem cell content of murine bone marrow, spleen, and blood. Limiting
dilution analysis of diffusion chamber cultures. J Cell Physiol. 1971;78: 73–78.
doi:10.1002/jcp.1040780111

6. Campbell P, Chapman MS, Wilk M, Boettcher S, Mitchell E, Dawson K, et al. Clonal dynamics
after allogeneic haematopoietic cell transplantation using genome-wide somatic mutations. 2023.
doi:10.21203/rs.3.rs-2868644/v1

7. WALLIS VJ, LEUCHARS E, CHWALINSKI S, DAVIES AJS. ON THE SPARSE SEEDING OF
BONE MARROW AND THYMUS IN RADIATION CHIMAERAS. Transplantation. 1975;19: 2–11.
doi:10.1097/00007890-197501000-00002

8. Szilvassy SJ, Humphries RK, Lansdorp PM, Eaves AC, Eaves CJ. Quantitative assay for
totipotent reconstituting hematopoietic stem cells by a competitive repopulation strategy. Proc
National Acad Sci. 1990;87: 8736–8740. doi:10.1073/pnas.87.22.8736

9. Bowling S, Sritharan D, Osorio FG, Nguyen M, Cheung P, Rodriguez-Fraticelli A, et al. An
Engineered CRISPR-Cas9 Mouse Line for Simultaneous Readout of Lineage Histories and Gene
Expression Profiles in Single Cells. Cell. 2020;181: 1410-1422.e27. doi:10.1016/j.cell.2020.04.048

10. Pei W, Feyerabend TB, Rössler J, Wang X, Postrach D, Busch K, et al. Polylox barcoding
reveals haematopoietic stem cell fates realized in vivo. Nature. 2017;548: 456–460.
doi:10.1038/nature23653

11. Ganuza M, Hall T, Finkelstein D, Chabot A, Kang G, McKinney-Freeman S. Lifelong
haematopoiesis is established by hundreds of precursors throughout mammalian ontogeny. Nat Cell
Biol. 2017;19: 1153–1163. doi:10.1038/ncb3607

12. Buescher ES, Alling DW, Gallin JI. Use of an X-linked human neutrophil marker to estimate
timing of lyonization and size of the dividing stem cell pool. J Clin Investig. 1985;76: 1581–1584.
doi:10.1172/jci112140

13. Berger CN, Sturm KS. Estimation of the Number of Hematopoietic Precursor Cells During Fetal
Mouse Development by Covariance Analysis. Blood. 1996;88: 2502–2509.
doi:10.1182/blood.v88.7.2502.bloodjournal8872502

14. Busque L, Patel JP, Figueroa M, Vasanthakumar A, Provost S, Hamilou Z, et al. Recurrent
Somatic TET2 Mutations in Normal Elderly Individuals With Clonal Hematopoiesis. Nat Genet.
2012;44: 1179–1181. doi:10.1038/ng.2413

15. Gandini E, Gartler SM, Angioni G, Argiolas N, Dell'Acqua G. Developmental implications of
multiple tissue studies in glucose-6-phosphate dehydrogenase-deficient heterozygotes. Proc Natl
Acad Sci United States Am. 1968;61: 945–8. doi:10.1073/pnas.61.3.945

16. Micklem HS, Lennon JE, Ansell JD, Gray RA. Numbers and dispersion of repopulating
hematopoietic cell clones in radiation chimeras as functions of injected cell dose. Exp Hematol.
1987;15: 251–7.

17. Harrison DE, Lerner C, Hoppe PC, Carlson GA, Alling D. Large Numbers of Primitive Stem Cells
Are Active Simultaneously in Aggregated Embryo Chimeric Mice. Blood. 1987;69: 773–777.
doi:10.1182/blood.v69.3.773.773

18. Cosgrove J, Hustin LSP, Boer RJ de, Perié L. Hematopoiesis in numbers. Trends Immunol.
2021;42: 1100–1112. doi:10.1016/j.it.2021.10.006

19. Fanti A-K, Busch K, Greco A, Wang X, Cirovic B, Shang F, et al. Flt3- and Tie2-Cre tracing
identifies regeneration in sepsis from multipotent progenitors but not hematopoietic stem cells. Cell
Stem Cell. 2023;30: 207-218.e7. doi:10.1016/j.stem.2022.12.014

20. Patel SH, Christodoulou C, Weinreb C, Yu Q, Rocha EL da, Pepe-Mooney BJ, et al. Lifelong
multilineage contribution by embryonic-born blood progenitors. Nature. 2022; 1–7.
doi:10.1038/s41586-022-04804-z

21. Schoedel KB, Morcos MNF, Zerjatke T, Roeder I, Grinenko T, Voehringer D, et al. The bulk of
the hematopoietic stem cell population is dispensable for murine steady-state and stress
hematopoiesis. Blood. 2016;128: 2285–2296. doi:10.1182/blood-2016-03-706010

22. Solomon M, Song B, Govindarajah V, Good S, Arasu A, Hinton EB, et al. Slow cycling and
durable Flt3+ progenitors contribute to hematopoiesis under native conditions. J Exp Med. 2023;221:
e20231035. doi:10.1084/jem.20231035

23. Göthert JR, Gustin SE, Hall MA, Green AR, Göttgens B, Izon DJ, et al. In vivo fate-tracing
studies using the Scl stem cell enhancer: embryonic hematopoietic stem cells significantly contribute
to adult hematopoiesis. Blood. 2005;105: 2724–2732. doi:10.1182/blood-2004-08-3037

24. Harrison DE, Astle CM, Lerner C. Number and continuous proliferative pattern of transplanted
primitive immunohematopoietic stem cells. Proc Natl Acad Sci. 1988;85: 822–826.
doi:10.1073/pnas.85.3.822

25. Harrison DE, Stone M, Astle CM. Effects of transplantation on the primitive
immunohematopoietic stem cell. J Exp Med. 1990;172: 431–437. doi:10.1084/jem.172.2.431

26. Harrison DE, Jordan CT, Zhong RK, Astle CM. Primitive hemopoietic stem cells: direct assay of
most productive populations by competitive repopulation with simple binomial, correlation and
covariance calculations. Exp Hematol. 1993;21: 206–19.

27. Li L, Bowling S, McGeary SE, Yu Q, Lemke B, Alcedo K, et al. A mouse model with high clonal
barcode diversity for joint lineage, transcriptomic, and epigenomic profiling in single cells. Cell.
2023;186: 5183-5199.e22. doi:10.1016/j.cell.2023.09.019

28. Busch K, Klapproth K, Barile M, Flossdorf M, Holland-Letz T, Schlenner SM, et al. Fundamental
properties of unperturbed haematopoiesis from stem cells in vivo. Nature. 2015;518: 542–546.
doi:10.1038/nature14242

29. Wilson A, Laurenti E, Oser G, Wath RC van der, Blanco-Bose W, Jaworski M, et al.
Hematopoietic Stem Cells Reversibly Switch from Dormancy to Self-Renewal during Homeostasis
and Repair. Cell. 2008;135: 1118–1129. doi:10.1016/j.cell.2008.10.048

30. Lerner C, Harrison DE. 5-Fluorouracil spares hemopoietic stem cells responsible for long-term
repopulation. Exp Hematol. 1990;18: 114–8.

31. Morrison SJ, Hemmati HD, Wandycz AM, Weissman IL. The purification and characterization of
fetal liver hematopoietic stem cells. Proc National Acad Sci. 1995;92: 10302–10306.
doi:10.1073/pnas.92.22.10302

32. Ema H, Nakauchi H. Expansion of hematopoietic stem cells in the developing liver of a mouse
embryo. Blood. 2000;95: 2284–2288. doi:10.1182/blood.v95.7.2284

33. Rybtsov S, Ivanovs A, Zhao S, Medvinsky A. Concealed expansion of immature precursors
underpins acute burst of adult HSC activity in foetal liver. Dev (Camb, Engl). 2016;143: 1284–1289.
doi:10.1242/dev.131193

34. Ganuza M, Hall T, Myers J, Nevitt C, Sánchez-Lanzas R, Chabot A, et al. Murine foetal liver
supports limited detectable expansion of life-long haematopoietic progenitors. Nat Cell Biol. 2022;24:
1475–1486. doi:10.1038/s41556-022-00999-5

35. Ceccaldi R, Parmar K, Mouly E, Delord M, Kim JM, Regairaz M, et al. Bone Marrow Failure in
Fanconi Anemia Is Triggered by an Exacerbated p53/p21 DNA Damage Response that Impairs
Hematopoietic Stem and Progenitor Cells. Cell Stem Cell. 2012;11: 36–49.
doi:10.1016/j.stem.2012.05.013

36. Carreau M, Gan OI, Liu L, Doedens M, Dick JE, Buchwald M. Hematopoietic compartment of
Fanconi anemia group C null mice contains fewer lineage-negative CD34+ primitive hematopoietic
cells and shows reduced reconstitution ability. Exp Hematol. 1999;27: 1667–1674.
doi:10.1016/s0301-472x(99)00102-2

37. Haneline LS, Gobbett TA, Ramani R, Carreau M, Buchwald M, Yoder MC, et al. Loss of FancC
Function Results in Decreased Hematopoietic Stem Cell Repopulating Ability. Blood. 1999;94: 1–8.
doi:10.1182/blood.v94.1.1.413k03_1_8

38. Zhang Q-S, Marquez-Loza L, Eaton L, Duncan AW, Goldman DC, Anur P, et al. Fancd2–/– mice
have hematopoietic defects that can be partially corrected by resveratrol. Blood. 2010;116: 5140–
5148. doi:10.1182/blood-2010-04-278226

39. Du W, Amarachintha S, Erden O, Wilson A, Meetei AR, Andreassen PR, et al. Fancb deficiency
impairs hematopoietic stem cell function. Sci Rep-uk. 2015;5: 18127. doi:10.1038/srep18127

40. Dubois EL, Guitton-Sert L, Béliveau M, Parmar K, Chagraoui J, Vignard J, et al. A Fanci
knockout mouse model reveals common and distinct functions for FANCI and FANCD2. Nucleic
Acids Res. 2019;47: 7532–7547. doi:10.1093/nar/gkz514

41. Zha J, Kunselman L, Fan J-M, Olson TS. Bone marrow niches of germline FANCC/FANCG
deficient mice enable efficient and durable engraftment of hematopoietic stem cells after
transplantation. Haematologica. 2019;104: e284–e287. doi:10.3324/haematol.2018.202143

42. Chen M, Tomkins DJ, Auerbach W, McKerlie C, Youssoufian H, Liu L, et al. Inactivation of Fac in
mice produces inducible chromosomal instability and reduced fertility reminiscent of Fanconi
anaemia. Nat Genet. 1996;12: 448–451. doi:10.1038/ng0496-448

43. Cerabona D, Sun Z, Nalepa G. Leukemia and chromosomal instability in aged Fancc–/– mice.
Exp Hematol. 2016;44: 352–357. doi:10.1016/j.exphem.2016.01.009

44. Young I-C, Wu B, Andricovich J, Chuang S-T, Li R, Tzatsos A, et al. Differentiation of fetal
hematopoietic stem cells requires ARID4B to restrict autocrine KITLG/KIT-Src signaling. Cell Rep.
2021;37: 110036–110036. doi:10.1016/j.celrep.2021.110036

45. Mochizuki-Kashio M, Yoon Y me, Menna T, Grompe M, Kurre P. FANCD2 Alleviates Physiologic
Replication Stress in Fetal Liver HSC. Biorxiv. 2020; 2020.09.30.320796.
doi:10.1101/2020.09.30.320796

46. Kumaravelu P, Hook L, Morrison AM, Ure J, Zhao S, Zuyev S, et al. Quantitative developmental
anatomy of definitive haematopoietic stem cells/long-term repopulating units (HSC/RUs): role of the
aorta-gonad-mesonephros (AGM) region and the yolk sac in colonisation of the mouse embryonic
liver. Development. 2002;129: 4891–4899. doi:10.1242/dev.129.21.4891

47. Yokomizo T, Ideue T, Morino-Koga S, Tham CY, Sato T, Takeda N, et al. Independent origins of
fetal liver haematopoietic stem and progenitor cells. Nature. 2022;609: 779–784.
doi:10.1038/s41586-022-05203-0

48. Rodriguez-Fraticelli AE, Wolock SL, Weinreb CS, Panero R, Patel SH, Jankovic M, et al. Clonal
analysis of lineage fate in native haematopoiesis. Nature. 2018;553: 212–216.
doi:10.1038/nature25168

49. Río P, Navarro S, Wang W, Sánchez-Domínguez R, Pujol RM, Segovia JC, et al. Successful
engraftment of gene-corrected hematopoietic stem cells in non-conditioned patients with Fanconi
anemia. Nat Med. 2019;25: 1396–1401. doi:10.1038/s41591-019-0550-z

50. Zhang X, Shang X, Guo F, Murphy K, Kirby M, Kelly P, et al. Defective homing is associated with
altered Cdc42 activity in cells from patients with Fanconi anemia group A. Blood. 2008;112: 1683–
1686. doi:10.1182/blood-2008-03-147090

51. Stone M. A general statistical model for clone--tissue studies, using X-chromosome inactivation
data. Biometrics. 1983;39: 395–409.

52. Lizen B, Claus M, Jeannotte L, Rijli FM, Gofflot F. Perinatal induction of Cre recombination with
tamoxifen. Transgenic Res. 2015;24: 1065–1077. doi:10.1007/s11248-015-9905-5

53. Joly S, Rodriguez L, Mdzomba JB, Carreau M, Pernet V. The ocular manifestations of Fanconi
anemia in a genetic mouse model. Investigative Ophthalmology & Visual Science. 60: 4738–4738.

Acknowledgments

Work was supported by R01-HL150882. We thank Florin Tuluc (CHOP Flow Cytometry Core) and

Jessica Gucwa (Cytek bioscience) for assistance in flow cytometry; Kaosheng Lv (CHOP) for

preparation of reagents; Zilu Zhou (Penn) for help in statistical analysis; Nancy Speck (Penn), Wei

Tong (Penn), Julia Warren (Penn), Ding-wen (Roger) Chen (CHOP), Stephanie N Hurwitz (Penn),

Hua Qing (Genentech), Hui Chen (Penn) for in-depth discussion. Some figure panels were

generated using Biorender.

Author information

Authors and Affiliations

Comprehensive Bone Marrow Failure Center, Children's Hospital of Philadelphia, Philadelphia, PA,

USA

Suying Liu, Sarah Adams, Juliana Ehnot,Seul K. Jung, Greer Jeffery, Theresa Menna, Peter Kurre

Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA

Suying Liu, Sarah Adams, Juliana Ehnot,Seul K. Jung, Greer Jeffery, Theresa Menna, Peter Kurre

Department of Biostatistics, Epidemiology and informatics, University of Pennsylvania, Philadelphia,

PA, USA

Haotian Zheng, Hongzhe Lee

Stem Cell Regulation Unit, St. Vincent's Institute of Medical Research, Fitzroy, VIC 3065, Australia

Louise E. Purton

Department of Medicine, The University of Melbourne, Parkville, VIC 3052, Australia

Louise E. Purton

900
901 Correspondence authors
902 Correspondence to Peter Kurre kurrep@chop.edu
903
Figure Legends
Figure 1. The Principle of Inverse Correlation between Variance of FP% and precursor numbers.
(A) Schematic of the Confetti cassette at the mouse *Rosa26* locus. Sequences of four fluorescence
protein are interspaced by loxP sites. Upon Cre-expression, the Confetti cassette will recombine
and express one of the four fluorescence proteins.
(B) The inverse relationship between variance of markers and precursor numbers. The distribution
of FP% is determined in the progeny to estimate the number of precursors.
(C) Workflow to validate the correlation formula between variance of FP% and precursor numbers
by a two-color cell system.
(D) BFP frequencies in wells seeded with 1 to 100,000 HL-60. Each dot represents a well. Two
replicates were shown. Each replicate consists of at least 20 wells per seeded number. Error
bars represent mean ± SD.
(E) The standard deviation of BFP% in wells seeded with various numbers of HL-60. Error bars
represent mean ± SD. N = 3 replicates.
(F) The correlation between seeded numbers and standard deviation of BFP%. N = 3 replicates.
(G) The correlation between seeded numbers and the numbers estimated with standard deviation of
BFP% and equation 2. N = 3 replicates except for seeded numbers 10^6 and 10^7 (grey circles).
For 10^6 and 10^7, N= 1 replicates.
For (A-C), images were created with Biorender.
923
Figure 2. Transplantation of Defined Number of Precursors
(A) Representative flow plots showing Confetti induction by HSC-Scl-CreER^T in the LSK population.
(B) Schematic for transplantation of defined number of precursors. Briefly, 4×10^6 Confetti-induced
WBM for "high precursor number" groups, or 0.25×10^6 WBM for "low precursor number" groups
were transplanted into lethally irradiated recipient mice. Image was created with Biorender.

(C) RFP_{Conf}% distribution in PB myeloid cells. Each dot represents one animal. Recipient sample
size = 7-14 mice per replicate, N = 3 replicates per group. Error bars represent mean ± SD.

(D) The precursor numbers of B cells, T cells, and myeloid cells in recipient mice. Dash lines mark
the level of historically estimated transplantable clone numbers. Each dot represents a precursor
number calculated from multiple mice. Error bars represent mean ± SD.

(E) The correlation of precursor numbers calculated from RFP_{Conf}%, YFP_{Conf}% or CFP_{Conf}%. Each
dot represents a precursor number calculated from multiple mice.

(F) The precursor number of BM MP and HSPC in recipient mice. Dash lines represent historically
estimated transplantable clone numbers. Each dot represents a precursor number calculated
from multiple mice. Error bars represent mean ± SD.

(G) The donor chimerism of recipient mice, and the $CD45.2^{+}$ chimerism differences between high-
and low- precursor number groups. Each solid dot represents one animal. Each outlined dot
represents the average value of a replicate containing multiple animals. Recipient sample size =
7-14 mice per replicate, N = 3 replicates for each precursor number group. Two-way ANOVA for
paired samples was performed.

(H) The frequency of myeloid cells in the peripheral blood of recipients. Each dot represents one
animal. Recipient sample size = 7-14 mice per replicate, N = 3 replicates for each precursor
number group. Two-way ANOVA for paired samples was performed.

ns, non-significant, * $p < 0.05$, ** $p < 0.001$.

Figure 3 Quantification of active hematopoietic precursor at steady-state

(A) Experiment schematic for Confetti induction in adult-induced animals. Image was created with
Biorender.

(B) Confetti labeling in B cells, myeloid cells and BM HSPCs. Statistics for comparisons between
labeling of month seven and other months were shown. One-way ANOVA for paired samples
was performed.

(C) RFP_{Conf}% in PB myeloid cells and BM HSPCs. Statistics for comparisons between labeling of
month seven and other months were shown. One-way ANOVA for paired samples was
performed.

(D) The number of myeloid precursors and the average precursor number from month five to seven.

(E) The number of BM precursors. Paired permutation was performed.

(F) The frequency-to-clone ratio of bone marrow HSPCs. Paired permutation test was performed.

Each solid dot represents one animal. Each outlined dot represents the average value of a replicate
containing multiple animals. Error bars represent mean ± SD. PB sample size = 14-22 mice per
replicate; BM sample size = 8-10 mice per replicate; N = 4 replicates. ns, non-significant, * $p < 0.05$,
** $p < 0.001$.

Figure 4. Number of precursor post-5-FU treatment and along developmental ontogeny

(A) Experiment schematic for one-dose 5-FU treatment. Image was created with Biorender.

(B) The number of BM precursors post-5FU treatment. Permutation test was performed. For UT,

sample size = 8-10 mice per replicate; N = 4 replicates; for 5-FU, sample size = 6-8 mice per

replicate; N = 5 replicates.

(C) Experiment schematic for Confetti induction at fetal stages. Image was created with Biorender.

(D) The Confetti labeling of B cells, T cells and myeloid cells. Each dot represents one animal. Each

outlined dot represents the average value of a replicate containing multiple animals. Sample size

= 4-11 mice per replicate; N = 5 replicates.

(E) The number of precursors in PB myeloid cells, B cells and T cells. Permutation test was

performed. For E14.5 and adult-induction, N = 4 replicates; for E11.5, n = 3 replicates.

(F) The number of BM precursors. Each dots represent one precursor number. Permutation test was

performed. For E14.5 and adult-induction, N = 4 replicates; for E11.5, N = 3 replicates.

(G) The relative fold change increase of precursor numbers from fetal- to adult-stage, as well as fold

change of CRU and cell counts.

Error bars represent mean ± SD. ns, non-significant, * $p < 0.05$, ** $p < 0.001$.

Figure 5. Precursor numbers in a mouse model of FA

(A) Experimental workflow to generate $Fancc^{+/+}$ and $Fancc^{-/-}$ mice, induce Confetti expression and

track precursor number dynamics. Image was created with Biorender.

(B) Confetti labeling in PB B cells and myeloid cells. Each dot represents one animal. Each outlined

dot represents the average value of a replicate containing multiple animals. Dashed lines

connect the values for the same replicate.

(C) The number of PB myeloid precursor.

(D) The number of BM precursor. Permutation test was performed.

(E) Experiment schematic for competitive transplantation. Image was created with Biorender.

(F) PB donor chimerism in recipient mice of young FA donors. Each dot represents one animal.

Dashed lines connect the average donor chimerism of three replicates. For $Fancc^{+/+}$ or $Fancc^{+/-}$,

n = 10 per replicate, N = 3 replicates; for $Fancc^{-/-}$, N = 7-9 per replicate, N = 3 replicates. Two-

way ANOVA was performed.

(G) The number of PB myeloid precursor post-transplantation of young FA donors. Dashed lines

connect the average precursor numbers. Permutation test was performed. N = 3 for each group.

(H) PB donor chimerism recipient mice of aging FA donor cells. Each dot represents one animal.

Dashed lines connect the average donor chimerism of three replicates. For both genotype, N =

10 per replicate, N = 3 replicates. Two-way ANOVA was performed.

(I) The number of PB B precursor post-transplantation of aging FA donors. Dashed lines connect

the average precursor numbers. Permutation test was performed. N = 3 for each genotype.

For (A-C), $Fancc^{+/+}$, sample size = 11-17 mice per replicate, N =5 replicates; $Fancc^{+/-}$, sample size =

10-22 mice per replicate, N =7 replicates; $Fancc^{-/-}$, sample size = 9-17 mice per replicate, N =4

replicates. For (D), $Fancc^{+/+}$, sample size = 5-9 mice per replicate, N =5 replicates; $Fancc^{+/-}$, sample

size = 6-12 mice per replicate, N =5 replicates; $Fancc^{-/-}$, sample size = 6-11 mice per replicate, N =4

replicates. Error bars represent mean ± SD. nd, not determined; ns, non-significant; * p < 0.05, ** p <

0.001. # represents lowest p value possible for permutation test.

1009

<u>Supplementary Figure Legend</u>

Figure S1: Establishing the correlation between variance of FP and precursor numbers.

(A) The correlation between precursor numbers and CVs of individual FP or all CVs (original

correlation formula), replotted from Ganuza et al., 2017.

(B) FP_{mean}% in a previously published dataset, replotted from Ganuza et al., 2017.

(C) The correlation between precursor numbers and standard deviation (σ) of individual FP,

replotted from Ganuza et al., 2017.

(D) The BFP% in HL-60 mix during two weeks of growth.

(E) The effect of outliers on precursor number estimation. Left, BFP% in samples with/without

outliers. Quartiles by Tukey method was shown. The outlier is highlighted by circle. Middle,

standard deviation of BFP% with/without outliers. Right, estimated precursor numbers

with/without outliers.

(F) The changes of standard deviation when recorded events varied. "&" denotes the potential

overestimated standard deviation due to low flow recorded events.

(G) The variance of estimated precursor numbers when varying sample sizes were used.

(H) The relative error of estimation as sample sizes increased. Each dot represents data from one

seeded number. Error bars represent mean ± SD.

1027

Figure S2: The induction specificity of HSC-Scl-CreER^T

(A) Experiment schematic to examine the induction specificity of HSC-Scl-CreER^T. N =10 for

induced animals, N =3 for non-induced animals. Image was created with Biorender.

(B) Flow gating strategy for BM HSPC and MPs.

(C) Confetti labeling in BM HSPCs. Error bars represent mean ± SD.

(D) Flow gating strategy for BM lineage cells.

(E) Confetti labeling in BM lineage cells. Error bars represent mean ± SD.

(F) Flow gating strategy to examine spleen cells.

(G) Confetti labeling in spleen cells. Error bars represent mean ± SD.

See Methods and Table S1 for the detailed antibody, fluorophore and marker combination used for

each cell type.

1039

Figure S3: Analysis of HSC-Scl-CreER^T background Cre activity

(A) Representative flow plot showing Confetti expression in PB T cells in 48-50-week-old HSC-Scl-

CreER^T/Confetti animals compared to an animal induced with one-day tamoxifen treatment.

(B) Quantification of RFP and YFP% in PB T cells. N =8 for non-induced 48-50-week-old animals.

(C) Representative flow plot for Confetti labeling in PB T cells post HSC-Scl-CreER^T-mediated

induction.

(D) The level of CFP$^+$ and RFP$^+$ double positive cells in PB T cells of animals induced by HSC-Scl-

CreER^T compared to Vav-Cre/Confetti animals analyzed at the same age. N =3 for Vav-Cre

animals, N= 8 for HSC-Scl-CreER^T animals.

1049

Figure S4: Variance of FP% inversely correlates with precursor numbers *in vivo*

(A) The correlation between expected numbers and the estimated numbers calculated from FP% of

precursors following Poisson distribution. See Methods for simulation details.

(B) Flow gating strategy for PB B cell, T cell and myeloid cell. See Methods and Table S1 for the

detailed antibody, fluorophore and marker combination used for each cell type.

(C) RFP$_{Conf}$% distribution in B cell and T cell.

(D) YFP$_{Conf}$% distribution in myeloid cell.

(E) CFP$_{Conf}$% distribution in myeloid cell.

(F) RFP$_{PB}$%, CFP$_{PB}$%, YFP$_{PB}$% and Confetti% distribution in myeloid cells.

(G) Linear correlation of PB precursor numbers calculated from RFP$_{Conf}$%, RFP$_{PB}$%, CFP$_{PB}$%,

YFP$_{PB}$%, Confetti% or RFP$_{RFP+YFP}$%, Each dot represents a precursor number calculated from

multiple mice.

For (C-F), each dot represents one animal. Recipient sample size = 7-14 mice per replicate, N =3

replicates per group. Error bars represent mean ± SD.

1064

Figure S5 BM analysis for high- and low-precursor number groups

(A) RFP$_{Conf}$% distribution in BM HSPCs. Each dot represents one animal

(B) BM donor chimerism.

(C) BM nucleated cell numbers. N =15 mice for high precursor number group, n= 14 for low

precursor number group. Unpaired t-test was performed.

(D) BM HSPC and MP frequency in donor CD45.2$^+$ population. Two-way ANOVA for paired samples

was performed.

For (B) and (D), each solid dot represents one animal; each outlined dot represents the average

value of a replicate containing multiple animals. Except for (C), recipient sample size = 7-14 mice per

replicate, N =3 replicates per group. For all graphs, Error bars represent mean ± SD. ns, non-

significant, * $p < 0.05$.

1076

Figure S6 Analysis of FP induction in adult-induced animals

(A) Confetti labeling in LSK after 14days of tamoxifen treatment. N =5 animals. Error bars represent

mean ± SD.

(B) CFP_{Conf}% in PB myeloid cells.

(C) CFP_{PB}% in PB myeloid cells.

(D) $RFP_{RFP+YFP}$% in PB myeloid cells.

(E) RFP_{PB}% and YFP_{PB}% in PB myeloid cells.

(F) CFP_{Conf}% in PB myeloid cells of fetal-induced animals

(G) BM HSPC frequencies. Error bars represent mean ± SD.

For (B-E) sample size = 14-22 mice per replicate; for (F), sample size = 4-11 mice per replicate; for

(G) sample size = 8-10 mice per replicate. For (B-E) and (G), N =4 replicates. For (F), N =5

replicates.

1089

Figure S7: Analysis related to 5-FU treatment and Confetti fetal induction

(A) PB myeloid cells frequencies before 5-FU treatment and ten days post-5-FU treatment. Dashed

lines connect the values for the same replicate. Sample size = 6-8 mice per replicate, N =5

replicates for 5-FU. Paired t test was performed.

(B) The BM HSPC frequencies of UT and 5-FU treated groups ten days post-5-FU treatment. For UT,

sample size = 8-10 mice per replicate, N =4 replicates; for 5-FU, sample size = 6-8 mice per

replicate, N =5 replicates.

(C) Confetti% in PB in E11.5-labeled animals. Dashed lines connect values for the same replicate.

(D) The distribution of FPs in Confetti[+] population of B cells. Data showing average from all mice. N

=24-34 for E14.5, N =7-24 for E11.5. Error bars represent mean ± SD.

For (A-C), each solid dot represents one animal; each outlined dot represents the average value of a

replicate containing multiple animals. ns, non-significant; * $p < 0.05$, ** $p < 0.001$.

1102

Figure S8 Analysis of FA mice at steady-state

(A) PB myeloid cell frequencies. Each dot represents one animal. Each outlined dot represents the

average value of a replicate containing multiple animals.

(B) PB B cell frequencies. Each dot represents one animal. Each outlined dot represents the

average value of a replicate containing multiple animals.

(C) Complete blood count analysis. Each dot represents one animal. N =13 for *Fancc*$^{+/+}$, N =16 for

Fancc$^{-/-}$ mice. Unpaired t test was performed.

(D) BM HSPC frequencies. Each dot represents one animal. Each outlined dot represents the

average value of a replicate containing multiple animals. One-way ANOVA was performed.

(E) BM nucleated cells counts. Each dot represents one animal. Unpaired t test was performed. N

=12 for *Fancc*$^{+/+}$ and *Fancc*$^{-/-}$ mice.

(F) RFP$_{RFP+YFP}$% in PB myeloid cells and B cells. Each dot represents one animal. Each outlined dot

represents the average value of a replicate containing multiple animals.

For (A-B) and (F), Fancc$^{+/+}$, sample size = 11-17 mice per replicate, N =5 replicates; Fancc$^{+/-}$,

sample size = 10-22 mice per replicate, N =7 replicates; Fancc$^{-/-}$, sample size = 9-17 mice per

replicate, N =4 replicates. For (D), Fancc$^{+/+}$, sample size = 5-9 mice per replicate, N =5 replicates;

Fancc$^{+/-}$, sample size = 6-12 mice per replicate, N =5 replicates; Fancc$^{-/-}$, sample size = 6-11 mice

per replicate, N =4 replicates. Error bars represent mean ± SD. ns, non-significant; * p < 0.05.

1121

Figure S9 Analysis of FA recipient mice

(A) Young *Fancc* donor BM HSPC frequency analysis. N = 5 for Fancc$^{+/+}$ or Fancc$^{+/-}$, N = 2 for

Fancc$^{-/-}$.

(B) Donor chimerism in recipient BM of young *Fancc* donor. Each dot represents one animal. Each

outlined dot represents the average value of a replicate containing multiple animals. Two-way

ANOVA was performed. Recipient sample size = 7-10 mice per replicate, N =3 replicates per

group.

(C) Number of PB B cell precursor in recipients of young FA donor.

(D) Number of PB T cell precursor in recipients of young FA donor.

(E) Number of BM HSPC precursor in recipients of young FA donor.

(F) Ageing *Fancc* donor BM HSPC frequency analysis. N = 3 for Fancc$^{+/+}$, N = 3 for Fancc$^{-/-}$.

(G) Number of PB myeloid cell precursor in recipients of aging FA donor.

(H) Number of PB T cell precursor in recipients of aging FA donor.

(I) Donor chimerism in recipient BM of ageing *Fancc* donor. Each dot represents one animal. Each

outlined dot represents the average value of a replicate containing multiple animals. Two-way

ANOVA was performed. Recipient sample size = 7-10 mice per replicate, N =3 replicates per

group.

(J) Number of BM HSPC precursor in recipients of aging FA donor.

For (C-E), (G-H) and (J), N =3 replicates per group; permutation test was performed; dashed lines

connect the average precursor numbers. Error bars represent mean ± SD. nd, not determined; ns,

non-significant; * p < 0.05, ** p < 0.001. # represents lowest p value possible for permutation test.

1143

Figure 1

Figure 2

Figure 4

Figure 5