MGPfact[XMBD]: A Model-Based Factorization Method for scRNA Data Unveils Bifurcating Transcriptional Modules Underlying Cell Fate Determination

Jun Ren[1,2,3], Ying Zhou[2,3], Yudi Hu[2], Jing Yang[2], Hongkun Fang[2], Xuejing Lyu[2], Jintao Guo[2], Xiaodong Shi[1], Qiyuan Li[2,3*]

1 School of Informatics, Xiamen University, Xiamen, 361105, China.

2 National Institute for Data Science in Health and Medicine, School of Medicine, Xiamen University, Xiamen, 361102, China.

3 Department of Hematology, The First Affiliated Hospital of Xiamen University and Institute of Hematology, School of Medicine, Xiamen University, Xiamen, 361102, China.

*Corresponding authors

Qiyuan Li, National Institute for Data Science in Health and Medicine, School of Medicine, Xiamen University, Xiamen, 361102, China

Fax: 0592-2185175; Tel: 0592-2185175; Email: qiyuan.li@xmu.edu.cn

Abstract: Manifold-learning is particularly useful to resolve the complex cellular state space from single-cell RNA sequences. While current manifold-learning methods provide insights into cell fate by inferring graph-based trajectory at cell level, challenges remain to retrieve interpretable biology underlying the diverse cellular states. Here, we described MGPfact[XMBD], a model-based manifold-learning framework and capable to factorize complex development trajectories into independent bifurcation processes of gene sets, and thus enables trajectory inference based on relevant features. MGPfact[XMBD] offers more nuanced understanding of the biological processes underlying cellular trajectories with potential determinants. When bench-tested across 239 datasets, MGPfact[XMBD] showed advantages in major quantity-control metrics, such as branch division accuracy and trajectory topology, outperforming most established methods. In real datasets, MGPfact[XMBD] recovered the critical pathways and cell types in microglia development with experimentally valid regulons and markers. Furthermore, MGPfact[XMBD] discovered evolutionary trajectories of tumor-associated CD8[+] T cells and yielded new subtypes of CD8[+] T cells with gene expression signatures significantly predictive of the responses to immune checkpoint inhibitor in independent cohorts. In summary, MGPfact[XMBD] offers a manifold-learning framework in scRNA-seq data which enables feature selection for specific biological processes and contributing to advance our understanding of

biological determination of cell fate.

Key Words: scRNA-seq, factorization, manifold-learning, bifurcation process, mixtures of Gaussian processes

Introduction

Data-mining of single-cell RNA sequencing (scRNA-seq) is often transformed into learning of lower-dimensional embedding (Becht et al., 2019; Haghverdi et al., 2015; Van der Maaten and Hinton, 2008) of the expression vectors, which represents the variation in the cellular space and helps explain the biological background. Previous single-cell studies used various embedding methods to characterize and visualize clustering of cells with unique biological functions(Saelens et al., 2019). Among the existing methods, graph-based embedding can better capture nonlinear biological signals among cells hence yielded more insights of the diversity of cells. More recent studies also use graph-based embedding (Costa et al., 2018) to reveal the dependency among cells and thereby reconstruct the evolutionary trajectory in the cellular space, which helps in understanding the determination of cell fate in development, differentiation and cancer.

To date, more and more manifold-learning methods are developed to infer lower-dimensional graphic embedding (manifolds) of scRNA-seq data, and yielded a number of trajectories corresponding to important cellular processes, such as TSCAN (Ji and Ji, 2016), DPT (Haghverdi et al., 2016), and scShaper (Smolander et al., 2022) belong to linear topological classes and reveal major linear pathways based on embedding spaces or cell clustering. TSCAN employs the construction of minimum spanning trees to discover pathways, while DPT reconstructs cellular trajectories using random walks, and scShaper integrates multiple pseudo-temporal solutions to deduce the shortest trajectory within a linear context. Additionally, there have been many approaches capable of inferring complex tree topological structures, such as the widely used Monocle series of algorithms includes Monocle 2 and 3 (Cao et al., 2019; Qiu et al., 2017b, 2017a). They leverage sophisticated graphing techniques to map intricate cell hierarchies; Monocle 2 creates DDRtree based on reversed graph embedding techniques, while Monocle 3 utilizes UMAP(Becht et al., 2019) for embedding. TinGa (Todorov et al., 2020) and scFates (Faure et al., 2023) represent more recent innovations. TinGa utilizes the Growing Neural Gas (GNG) algorithm (Fritzke, 1994) to construct an adaptive graph structure that effectively captures the density structure of the dataset. scFates, streamlines pseudotime analysis with flexible tree learning options,

advanced feature extraction tasks, and specific functionalities for fork analysis.

Moreover, recent studies based on RNA velocity has provided insights into cell state transitions.

These methods measure RNA synthesis and degradation rates based on the abundance of spliced

and unspliced mRNA, such as CellRank (Lange et al., 2022). Nevertheless, current RNA velocity

analyses are still unable to resolve cell-fates with complex branching trajectory. Deep learning

methods such as scTour (Li, 2023) and TIGON (Sha, 2024) circumvent some of these limitations,

offering continuous state assumptions or requiring prior cell sampling information.

Despite these advances, trajectory prediction remains a major challenge in single cell analysis.

Firstly, graph-based trajectories represent synergic effects multiple biological processes, making it

difficult to disentangle the effects of specific process, hence limited model interpretability and the

power to gain novel biological insights. Secondly, the inference of trajectory is highly dependent on

the biases in the gene-selection, whereas conventional statistical feature-selection methods are less

efficient for the learning of complex topologies, and adds to the difficulty of suggesting candidate

genes for downstream functional study. Additionally, many approaches require additional prior

information, which further limits the applicability.

Here, we describe MGPfact^XMBD (Factorization based on Mixtures of Gaussian Processes), a

model-based, unsupervised manifold-learning method which factorize complex cellular trajectories

into interpretable bifurcation Gaussian processes of transcription, and thereby enable discovery of

specific biological determinants of cell fate. In the validation datasets, MGPfact recapitulated

developmental trajectory of microglia and recovered key regulatory factors which have been proved

experimentally. Moreover, MGPfact discovered highly specific subtypes of tumor-associated CD8$^+$

T cells which associated with benefit to cancer immunotherapy.

Bring together, MGPfact is a knowledge discovery tool which conducts manifold-learning and

factorization simultaneously. MGPfact offers two advantages in future scRNA-seq analyses: first, it

provides highly interpretable, factorizable cellular trajectories with matched gene expression

modules; then, it provides efficient feature-selection for graph-based embedding, thus enhancing

our understanding of the determination of cell fate.

Results

Design of MGPfact

Fig 1. Overview of MGPfact workflow. The complete workflow comprises two major stages: MURP downsampling with preprocessed data and trajectory reconstruction. In the stage of trajectory reconstruction, the scRNA-seq data were first factorized into independent bifurcation processes based on mixtures of Gaussian processes, which were then merged into a consensus trajectory.

The analytical pipeline of MGPfact consists two major stages (Fig. 1). An algorithmic description is given in Algorithm 1.

First, we performed downsampling of the preprocessed scRNA-seq data Y to yield a M-by-N expression matrix Y' based on the "minimum unbiased representative points" (MURP) as described previously (Ren et al., 2022), where M representative points were considered as landmarks of the cellular trajectory and N is the number of genes. Then, we computed L Principal Components (PCs) of the downsampled expression matrix to obtain the matrix $Y^* = \{y_1^*, y_2^*, y_3^*, \dots y_L^*\}$ (M-by-L),

$$y_l^* = Y' \cdot \mathbf{v}_l \tag{1}$$

where $\mathbf{v}_l$ is projection vector, y_l^* serve as the l-th initial state of embedding.

Next, we used typical Gaussian Process Regression of y_l^* on pseudotime T:

$$y_l^* = f(T) + \varepsilon \tag{2}$$

where $f(T)$ is a Gaussian Process (GP) with covariance matrix S.

$$f(\mathbf{T}) = \mathcal{GP}(0, \mathbf{S} + \sigma_S^2 \cdot \mathbf{I}) \tag{3}$$

And for all features:

$$p(\mathbf{Y}^*, f(\mathbf{T})) = p(\mathbf{Y}^*|f(\mathbf{T})) \cdot p(f(\mathbf{T})) \tag{4}$$

where $p(\mathbf{Y}^*|f(\mathbf{T}))$ is defined as follow:

$$p(\mathbf{Y}^*|f(\mathbf{T})) = \prod_{l=1}^{L} p(\mathbf{y}_l^*|f(\mathbf{T})) = \prod_{l=1}^{L} \mathcal{N}(\mathbf{y}_l^*|0, \mathbf{S} + \sigma_S^2 \cdot \mathbf{I}) \tag{5}$$

Specifically, we consider $\mathbf{S}$ as a mixture of L independent bifurcation Gaussian processes(Schulz et al., 2018),

$$\mathbf{S} = \sum_{l=1}^{L} \mathbf{s}_l \tag{6}$$

To cope with the bifurcation processes in cell fate, each of the Gaussian processes is defined with a bifurcation point at b_l, branching labels $\mathbf{c}_l$, and the necessary hyperparameters. The branching labels $\mathbf{c}_l \in \{0,1,2\}$, correspond to different phases and states of cell fate, where $\mathbf{c}_l = \mathbf{0}$ corresponds the phase before branching, and $\mathbf{c}_l \in \{1,2\}$. corresponds to the two cellular states of the bifurcation process, respectively. For any landmark (MURP) x,

$$\begin{cases} c_{l,x} = 0, & if \ t_x < b_l \\ c_{l,x} \in \{1,2\}, & if \ t_x \geq b_l \end{cases} \tag{7}$$

The covariance matrix $\mathbf{s}_l$ for trajectory l can be expressed as follows,

$$[\mathbf{s}_l]_{x,y} = \mathcal{K}(t_x, t_y) \tag{8}$$

where K is a kernel function. We employ radial basis function (rbf) and polynomial kernel function (pl). We chose these two kernel functions for the effectiveness in handling nonlinear and polynomial relationships, achieving a balance between model performance and computational efficiency.

$$k_{rbf}(t_x, t_y) = \lambda_{rbf} \cdot e^{\left(-\alpha_{rbf}\|t_x - t_y\|^2\right)} \tag{9}$$

$$k_{pl}(t_x, t_y) = \left(\lambda_{pl}t_x^T t_y + c_{pl}\right)^{d_{pl}} \tag{10}$$

And the $\mathcal{K}(t_x, t_y)$ is calculated as follows:

135
$$\mathcal{K}(t_x, t_y) = \begin{cases} k_{rbf}(t_x, t_y) + k_{pl}(t_x, t_y) & t_x, t_y < b_l \\ k_{rbf}(t_x, t_y) + k_{pl}(t_x - b_l, t_y - b_l) & t_x, t_y > b_l, c_{l,x} = c_{l,y} \\ \dfrac{k_{rbf}(t_x, b_l) \cdot k_{rbf}(b_l, t_y)}{k_{rbf}(b_l, b_l)} & \\ & c_{l,x} \neq c_{l,y} \end{cases} \tag{11}$$

136 Therefore, $p\big(\mathbf{Y}^* \big| f(\mathbf{T})\big)$ is updated as follows:

137
$$p\big(\boldsymbol{Y}^* \big| f(\boldsymbol{T})\big) = \prod_{l=1}^{L} \mathcal{N}\big(\boldsymbol{y}_l^* \big| 0, \textstyle\sum_{l=1}^{L} \boldsymbol{s}_l^* + \sigma_S^2\big) \tag{12}$$

138 We infer all parameters by maximizing the posterior likelihood using Markov Chain Monte Carlo

139 (MCMC) methods available in Mamba(B J, 2014). The posterior distribution of pseudotime $\boldsymbol{T}$ can

140 be represented as:

141
$$p(\boldsymbol{T} \mid \boldsymbol{Y}^*) \propto p(\boldsymbol{Y}^* \mid f(\boldsymbol{T})) \cdot p(f(\boldsymbol{T})) \tag{13}$$

where $p(\boldsymbol{Y}^* \mid f(\boldsymbol{T}))$ is the likelihood function of the observed data $\boldsymbol{Y}^*$, and $p(f(\boldsymbol{T}))$ is the prior

distribution of the Gaussian process. This posterior distribution integrates the observed data with

model priors, enabling inference of pseudotime and trajectory simultaneously. Due to the high

autocorrelation of T in the posterior distribution, we use Adaptive Metropolis within Gibbs (AMWG)

sampling (Roberts and Rosenthal, 2009; Tierney, 1994). Other parameters are estimated using the

more efficient SLICE sampling technique (Neal, 2003).

Algorithm 1 MGPfact: infer cell fate trajectory

INPUT: expression matrix $\boldsymbol{Y}$, independent trajectories number L
OUTPUT: $\boldsymbol{\theta} = \{\boldsymbol{T}, \boldsymbol{B}, \boldsymbol{C}, \text{other hyperparameters}\}$; trajectory toplogy
1: initialize all parameters in $\boldsymbol{\theta}$
2: covariance matrix $S = 0$
3: object function $\mathcal{L} = 0$
4: $\boldsymbol{Y}' \leftarrow$ MURP downsampling $Y_{p,n}$;
5: $\boldsymbol{Y}^* \leftarrow$ PCA analysis
6: $\boldsymbol{\theta} \leftarrow$ Opmized ObjectF
7: $Graph \leftarrow$ Greate independent trajectory using $\boldsymbol{\theta}$
8: **function** OBJECTF($\boldsymbol{Y}^*, \boldsymbol{T}, b_l, \boldsymbol{c}_l$)
9:      $Q \leftarrow dim(\boldsymbol{Y}^*)[2]$
10:     **for** $q = 1; q < Q; q++$ **do**   ▷ object function
11:        $\boldsymbol{v} \leftarrow [\boldsymbol{Y}^*]_q$
12:        **for** $l = 1; l < L; l++$ **do**
13:           $s_l \leftarrow$ COV($\boldsymbol{T}, b_l, \boldsymbol{c}_l$))
14:           $\boldsymbol{S} \leftarrow \boldsymbol{S} + \boldsymbol{s}_l$
15:        **end for**
16:        $\mathcal{L} \leftarrow \mathcal{L} +$ MultivariateNormalPDF($\boldsymbol{v}, 0, \boldsymbol{S}$)
17:     **end for**
18:     **return** $\mathcal{L}$
19: **end function**
20: **function** COV($\boldsymbol{T}, b_l, \boldsymbol{c}_l$)   ▷ construct covariance matrix
21:     $P \leftarrow length(\boldsymbol{T})$
22:     **for** $i = 1; i < P; i++$ **do**
23:        **for** $j = 1; j < P; j++$ **do**
24:           $s_l[i,j] \leftarrow$ KERNEL($t_i, t_j, b_l, \boldsymbol{c}_l$)
25:        **end for**
26:     **end for**
27:     **return** s_l
28: **end function**
29: **function** KERNEL($t_i, t_j, b_l, c_{l,i}, c_{l,j}$)   ▷ kernel function
30:     **if** $t_i, t_j < b_l$ **then**
31:        **return** $k_{rbf}(t_i, t_j) + k_{pl}(t_i, t_j)$
32:     **else if** $t_i, t_j > b_l \wedge c_{l,i} = c_{l,j}$ **then**
33:        **return** $k_{rbf}(t_i, t_j) + k_{pl}(t_i - b_l, t_j - b_l)$
34:     **else if** $c_{l,i} \neq c_{l,j}$ **then**
35:        **return** $\frac{k_{rbf}(t_i, b_l) \cdot k_{rbf}(b_l, t_j)}{k_{rbf}(b_l, b_l)}$
36:     **end if**
37: **end function**

Then, MGPfact can identify genes that have significant impacts on the branching events in the trajectories. we introduce a rotation matrix $R = \{r_1, r_2, \ldots, r_L\}$ to obtain factor score w_l for each trajectory l by rotating $\mathbf{Y}^*$.

$$\mathbf{w}_l = \mathbf{Y}^* \cdot \mathbf{r}_l + e_l^2 \tag{14}$$

For all trajectories,

$$p(\boldsymbol{W}|\boldsymbol{Y}^*) = \prod_{l=1}^{L} [\mathcal{N}(\boldsymbol{Y}^* \cdot \boldsymbol{r}_l + e_l|0, \boldsymbol{s}_l) \cdot \mathcal{N}(e_l|0, \sigma_{error}^2)] \tag{15}$$

where e_l is the error term for the l-th trajectory.

Specifically, the factor scores w_l for each gene onto the l-th trajectory can be expressed using equations (1) and (14) as follows,

$$\mathbf{w}_l = [\boldsymbol{Y}' \cdot \mathbf{v}_l] \cdot \mathbf{r}_l + e_l^2 = \boldsymbol{Y}' \cdot \mathbf{u}_l + e_l^2 \tag{16}$$

Here, $\mathbf{u}_l$ is used to represent the contribution (gene weight) of each gene to the l-th trajectory, thus enabling gene-selection based on the inferred trajectories.

162   Additionally, we can combine independent bifurcation processes to form a consensus diffusion

163   tree to represent the trajectory of cell fate (Supplementary Methods, Supplementary Table 1).

Performance evaluation of MGPfact

Robustness analysis of MGPfact

Before the performance evaluation, we performed a grid search on the number of independent

trajectories in 100 training datasets and selected $L = 3$ for downstream testing (Supplementary Fig.

1, Supplementary Fig. 2, Methods).

To further validate the efficacy of MURP downsampling step in MGPfact, we employed an

alternative downsampling using randomly selected cells for trajectory inference (Methods). This

comparison revealed that the prediction accuracy substantially diminished without MURP, evidenced

by a notable reduction in branch assignment ($F1_{branches}$, 20.5%) and cell ordering (cor_{dist}, 64.9%)

(Supplementary Fig. 3). In contrast, trajectory predictions utilizing MURP-based downsampling

realized an overall score increase of 5.31% to 185%, underscoring the indispensable role of MURP

in the trajectory inference capabilities of MGPfact.

Furthermore, we performed a robustness analysis on the topological consistency of the

predicted consensus trajectory by comparing prediction results from randomly sampled subsets of

the original data. As a result, the consensus trajectory predicted from random subsets by MGPfact

retained a high degree of congruence with those from the original datasets (Supplementary Fig. 4,

$HIM_{mean} = 0.686$). This outcome attests to MGPfact's robustness and generalizability under varying

data conditions, hence the capability of retrieving conserved bifurcative trajectories in the data.

MGPfact predicts cellular trajectories

Then, we assessed the performance of MGPfact for prediction of cellular trajectories in 239 test

datasets, including 171 synthetic and 68 real datasets, alongside with another 7 existing algorithms

(Supplementary Fig. 1). For overall performance score, MGPfact ($Overall_{mean} = 0.534$) ranked

second only to TinGa ($Overall_{mean} = 0.563$) and outperformed the rest of 6 algorithms (Fig. 2a).

Particularly, MGPfact demonstrated the highest accuracy in predicting cell fate in branching

trajectory (Fig. 2b, $F1_{branches\,mean} = 0.482$). As for other three individual metrics, MGPfact ranked

the 4[th] in HIM ($HIM_{mean} = 0.606$), the 6[th] in cor_{dist} ($cor_{dist\,mean} = 0.507$), and the 4[th] in

$wcor_{features}$ $(wcor_{features_{mean}} = 0.712)$ (Fig.2c-e).

Next, we compared the performance of MGPfact with the other algorithms in 9 different trajectory types, respectively, for predicting differentiated cell fate ($F1_{branches}$). As a result, MGPfact significantly outperformed more than half of the algorithms tested in the following trajectory types (T-test P<0.1, Table 1): disconnected graph (N=5), linear (N=5), bifurcation (N=4), multifurcation (N=4) and tree (N=4). As for the other three metrics, MGPfact also showed advantages in HIM in linear (N=5), bifurcation (N=3), convergence (N=3); and in $wcor_{features}$ in multifurcation (N=6), bifurcation (N=5). Nevertheless, MGPfact showed limited predictive performance for cor_{dist} (Supplementary Table 2).

Fig. 2. Trajectory inference (TI) performance of state-of-the-art methods in 239 test datasets. a. Overall scores；b. $F1_{branches}$； c. HIM； d. cor_{dist}； e. $wcor_{features}$. All results are color-coded based on the trajectory types, with the black line representing the mean value. The "Overall" assessment is calculated as the geometric mean of all four metrics.

Table 1 MGPfact outperformed state-of-the-art methods in $F1_{branches}$. P-values based one-sided paired t-tests suggest that the $F1_{branches}$ scores of MGPfact were significantly higher than those of the other

208 **methods for different trajectory types in the test set.**

Trajectory Type	DPT	Monocle DDRTree	Monocle3	scFates Tree	scShaper	TinGa	TSCAN
Acyclic Graph	0.169	0.735	0.341	0.167	**0.006**	0.485	**0.002**
Bifurcation	0.559	**0.001**	**0.015**	0.298	**0.010**	0.772	**0.002**
Convergence	0.275	**0.059**	**0.000**	**0.027**	0.120	0.871	**0.008**
Cycle	**0.000**	**0.000**	**0.002**	0.114	0.991	0.659	0.982
Disconnected Graph	**0.020**	**0.001**	0.108	**0.007**	**0.001**	0.475	**0.000**
Connected Graph	**0.053**	0.214	0.114	0.285	**0.005**	**0.057**	**0.001**
Linear	**0.000**	**0.000**	**0.000**	**0.000**	1.000	**0.000**	0.980
Multifurcation	**0.033**	**0.001**	**0.041**	0.717	0.552	0.758	**0.051**
Tree	**0.021**	0.809	0.918	**0.086**	**0.000**	0.993	**0.000**

It is also worth noting that in 68 test datasets of real cell populations, MGPfact ranked the top of

all 7 algorithms for "overall score" (Fig. 3a). As for the individual metric, MGPfact ranked to top for

predicting trajectory topology ($HIM_{mean} = 0.721$, Fig. 3c); and the 2nd in trajectory branching

($F1_{branches_{mean}} = 0.600$, Fig. 3b) with no significant difference with that of the top predictor

(scShaper, T-test, P = 0.829). These data show that MGPfact can effectively reconstruct the

trajectory of cell fate and retrieve relevant biological processes. As for the other metrics in real test

datasets, MGPfact was the 5th in the similarity of cell locations ($cor_{dist_{mean}} = 0.46$), and the 3rd in

the similarity of gene significance ($wcor_{features_{mean}} = 0.735$). The differences between the metrics

of MGPfact and those of the top performers were subtle (Fig. 3d-e, $\Delta cor_{dist_{mean}} =$

$0.07, \Delta wcor_{features_{mean}} = 0.05$).

We also analyzed three real-world datasets (Saelens et al., 2019) , each representing a unique

topology of trajectory: linear, single bifurcation, and multiple bifurcations. MGPfact excelled in

capturing key developmental trajectory with branch points (Supplementary Fig. 5). In the linear

trajectory, MGPfact accurately predicted the absence of bifurcations, aligning well with the ground

truth ($overall = 0.871$). For the bifurcation trajectory, MGPfact successfully identified the main

bifurcation ($overall = 0.636$). As for the multifurcation trajectory, MGPfact's prediction is also close

to the ground truth, as reflected by the overall score ($overall = 0.566$).

In summary, our data suggest that MGPfact is highly efficient in predicting cell fate in branching

trajectory ($F1_{branches}$) and topological structure (HIM). These capabilities align with the primary objectives of the algorithm, namely, effective identification of the branching events in the development processes of cells. In addition, MGPfact performed better in real datasets, suggesting its robustness to noise from real experimental conditions. Nevertheless, trajectory inference by MGPfact is based on factorization of the covariance matrix, hence less performance in $wcor_{features}$ and cor_{dist} than methods based on full covariance matrix.

Fig 3. Trajectory inference (TI) performance of state-of-the-art methods in 68 test datasets of real cell population. a. Overall scores；b. $F1_{branches}$ ； c. HIM；d. cor_{dist}；e. $wcor_{features}$ ．All results are color-coded based on the trajectory types, with the black line representing the mean value for ranking all methods. The "Overall" assessment is calculated as the geometric mean of all four metrics.

Comparative of Time Efficiency and Memory Consumption

We also compared the runtime and memory usage of different algorithms across 239 test datasets (Supplementary Table 3). MGPfact's average maximum memory consumption

243 ($memory_{mean}^{(max)} = 0.75GB$) is comparable to those of the other algorithms ($memory_{mean}^{(max)} \in$

244 $[0.55, 0.91]GB$). As a trade-off for its advantages in feature-selection and factorization, MGPfact

245 requires moderately longer execution time than the other algorithms ($time_{mean} = 3.42\ min$).

MGPfact recovers the trajectory of early postnatal microglia development

247 The main advantage of MGPfact lies in the capability to factorize a complex cellular trajectory

248 into bifurcation processes of selected co-expressed genes. To illustrate how MGPfact elucidates the

249 biological process underlying cell fate determination, we applied MGPfact to a scRNA-seq data of

250 microglia development and validated the results with experimental evidences (Li et al., 2019).

MGPfact recovers the determinants of the microglia development

Utilizing MGPfact, we reconstructed the developmental trajectories of microglia from immature

microglia (IM at pseudotime 0) to homeostatic microglia (HM) and proliferative-region-associated

microglia (PAM) (Fig. 4a-c, left panel). MGPfact identified three bifurcation processes

(Supplementary Table 4), each corresponding to 74-90 highly weighted genes (HWG, absolute gene

weight > 0.05) (Fig. 4a-c, right panel).

The first bifurcation determines the differentiated cell fates of PAM and HM, which involves a set

of notable marker genes of both cell types, such as Apoe, Selplg (HM), and Gpnmb (PAM). The

second bifurcation determines the proliferative status, which is crucial for the development and

function of PAM and HM (Guzmán, n.d.; Li et al., 2019). The genes affected by the second bifurcation

are associated with cell cycle and proliferation, such as Mki67, Tubb5, Top2a. The third bifurcation

influences the development and maturity of microglia, of which the highly weighted genes, such as

Tmem119, P2ry12, and Sepp1 are all previously annotated markers for establishment of the fates

of microglia(Anderson et al., 2022; Li et al., 2019) (Supplementary Table 4).

Moreover, we retrieved highly active regulons from the HWG by MGPfact, of which the

significance is quantified by the overall weights of the member genes. These data unveiled highly

active transcription regulations in each bifurcation processes, which further traced back to the

influential transcription factors as determinants of microglia development, such as Hif1a, E2f5, and

Nfkb1 (Fig. 4d-e, Methods). Specifically, Hif1a is crucial for microglial activation and directly linked

to neurodegenerative disease progression (Wang et al., 2022). Our data showed an upregulation of

Hif1a in the PAM-branch (phase 1) of the first bifurcation, reaffirming the role of Hif1a in PAM

differentiation. The other two transcription factors, E2f5 and Nfkb1, were active in phase 2 of the 2nd

bifurcation and the 3rd bifurcation, respectively. Both are known for the roles in microglial

development (Dresselhaus and Meffert, 2019; Nawal, n.d.).

Fig. 4. **MGPfact reconstructed the developmental trajectory of microglia, recovering known determinants of microglia fate. a-c.** The inferred independent bifurcation processes with respect to the unique cell types (color-coded) of microglia development, where phase 0 corresponds to the state before bifurcation; and phases 1 and 2 correspond to the states post-bifurcation. Each colored dot represents a metacell of unique cell type defined by MURP. The most highly weighted regulons in each trajectory were labeled by the corresponding transcription factors (left panels). The HWG of each bifurcation process include a set of highly weighted genes, of which the expression levels differ significantly among phases 1, 2, and 3 (right panels). **d.** The most highly weighted regulons influencing the three developmental trajectories of microglia. **e.** The expression levels of the transcription factors of highly weighted regulons in each trajectory significantly differ among different phases. **f.** The consensus developmental trajectory by merging the three bifurcation processes. Point 0 denotes the initial of differentiation, whereas the notion of "*n-m*" denotes the *m*-th branch from the branching point *n*. Each colored circle represents a landmark (MURP) of the trajectory, showing the fraction of cell types. The transcription factors of highly weighted regulons in each bifurcation process were used to label each branching point. Particularly, PAM-T1 and PAM-T2 are the two newly defined subtypes of PAM. **g.** Selected differently expressed genes between PAM-T1 and PAM-T2 ($|logfc| > 0.25$, adjusted P-value < 0.1) are shown by colored-dots corresponding to the mean expression levels in either cell type. The IDs validated marker genes for PAM are labeled in green. In all box plots, the horizontal line represents the median value, and the whisker extends to the furthest data point within 1.5 times the interquartile range. Significance is denoted as: ns, not significant, * P < 0.05, ** P < 0.01, *** P < 0.001, **** P < 0.0001, Wilcoxon rank-sum test.

Using consensus trajectories to delineate the development of microglial cells

We generated a consensus trajectory of microglial development from three independent bifurcation processes (Methods). Of note, the consensus trajectory revealed two distinct subtypes of proliferative-region-associated microglia (PAM), PAM-T1 (Hif1a+/E2f5-/Nfkb1+) and PAM-T2 (Hif1a+/E2f5-/Nfkb1-) (Fig. 4f). Particularly, the highly expressed genes in PAM-T2, including Spp1, Gpnmb, Lgals1, and Cd63, are previously identified in disease-associated microglia (DAM)(Li et al., 2019). Thus, our finding reaffirmed the connection between the two cell types, DAM and PAM, and suggested Nfkb1 as a potential determinant of differentiation of PAM (Fig. 4g, Supplementary Table 5).

In conclusion, MGPfact reconstructed the cellular trajectory of microglial development, identified distinct cell types with marker genes and key regulators which are highly consistent to the experimental evidences (Dresselhaus and Meffert, 2019; Li et al., 2019; Nawal, n.d.; Wang et al., 2022) .

Using MGPfact to decipher the evolution of tumor-associated CD8$^+$ T cells

Next, we applied MGPfact to two populations of tumor-associated CD8$^+$ T cells from non-small cell lung cancer (NSCLC)(Guo et al., 2018) and colorectal cancer (CRC)(Zhang et al., 2018), respectively. Using the same analytical pipeline as above, we identified a set of CD8$^+$ T cell gene expression signatures (GES) from MGPfact-inferred trajectories, which are significantly predictive of clinical outcomes and immune treatment responses. Additionally, our data unveiled novel subtypes of tumor-associated CD8$^+$ T cells with strong clinical implications.

MGPfact better explains the fate of tumor-associated CD8$^+$ T cells

We assessed the goodness-of-fit (adjusted R-square) of the consensus trajectory derived by MGPfact and three methods (Monocle 2, Monocle 3 and scFates Tree) for the CD8$^+$ T cell subtypes described in the original studies (Guo et al., 2018; Zhang et al., 2018). The data showed that MGPfact significantly improved the explanatory power for most CD8+ T cell subtypes over Monocle 2, which was used in the original studies (P < 0.05, see Table 2 and Supplementary Table 6), except for the CD8-GZMK cells in the CRC dataset. Additionally, MGPfact demonstrated better explanatory power in specific cell types when compared to Monocle 3 and scFates Tree. For instance, in the NSCLC dataset, MGPfact exhibited higher explanatory power for CD8-LEF1 cells (Table 2, R-squared = 0.935), while Monocle 3 and scFates Tree perform better in other cell types.

Table 2 Comparison of the explanatory power for CD8$^+$ T cell fate for MGPfact and three other different methods. Adjusted R-squared values and P-values based on F-tests demonstrate the relative performance of MGPfact, Monocle 2, Monocle 3, and scFates Tree in fitting the experimentally characterized and annotated CD8+ T cell subtypes.

		MGPfact		Monocle 2		Monocle 3		scFates Tree	
		Adjust R^2	P	Adjust R^2	P	Adjust R^2	P	Adjust R^2	P
	CD8-LEF1	**0.935**	0.000	0.176	0.000	0.089	0.08	0.902	0.000
	CD8-CD28	**0.195**	0.002	0.170	0.000	0.108	0.06	0.006	0.145
NSCLC	CD8-CX3CR1	0.634	0.000	0.259	0.000	0.629	0.000	**0.882**	0.000
(GSE99254)	CD8-GZMK	0.259	0.000	0.189	0.000	**0.855**	0.000	0.547	0.000
	CD8-ZNF683	0.232	0.001	0.051	0.043	**0.625**	0.003	0.039	0.003
	CD8-LAYN	0.435	0.000	0.031	0.027	0.503	0.018	**0.523**	0.000
CRC	CD8-LEF1	0.311	0.000	0.027	0.036	0.461	0.007	**0.99**	0.000
(GSE108989)	CD8-GPR183	0.380	0.000	0.032	0.025	**0.474**	0.006	0.139	0.000

CD8-CX3CR1	0.648	0.000	0.047	0.008	0.454	0.007	**0.817**	0.000
CD8-GZMK	0.130	0.013	0.550	0.000	**0.855**	0.000	0.236	0.000
CD8-CD6	0.277	0.000	0.109	0.007	**0.45**	0.008	0.054	0.000
CD8-CD160	0.124	0.016	0.080	0.025	**0.856**	0.000	0.707	0.000
CD8-LAYN	**0.741**	0.000	0.172	0.000	0.373	0.021	0.505	0.000

MGPfact identifies T-cell gene expression signatures with clinical implications

MGPfact discerned the different cellular fates of tumor-associated CD8$^+$ T cells by distinct bifurcation processes. To reveal the clinical implications of these bifurcation processes, we retrieved the mean expression vectors corresponding to either phase (branch) as Gene Expression Signatures (GES), and stratified cancer cohorts by quantifying the propensities to certain destiny of CD8$^+$ T cells (Methods). Our data demonstrated pronounced disparities in the clinical outcomes associated with different fates of CD8$^+$ T cells among patients (Fig. 5a-b, Supplementary Fig. 8e). In NSCLC, trajectory 1 is associated with cytotoxic T cells (96%, phase 1) and higher overall survival in TCGA-lung adenocarcinoma (LUAD) patients. Trajectory 2 is associated with exhausted T cells (91%, phase 2), and lower overall survival in the same cohort (Fig. 5a, Supplementary Fig. 6a-b). Similarly, in the CRC, trajectory 1 is associated with exhausted T cells (95%, phase 1) and poor overall survival in TCGA-COAD patients (Fig. 5b, Supplementary Fig. 8a-b).

In addition, for each trajectory identified in NSCLC and CRC, we selected a set of HWG (absolute gene weight > 0.05) to characterize the underlying biological processes and key transcription factors determining the bifurcation (Supplementary Table 7-8, Supplementary Fig. 6c-d, Supplementary Fig. 8c-d). In NSCLC, the HWG of trajectory 1 are primarily implicated in immune responses associated with antigen processing and presentation (Supplementary Table 9), while trajectory 2-HWG involve processes of immune cell migration (Supplementary Table 9). For CRC, trajectory 1 is enriched for genes of T cell activation and regulation (Supplementary Table 10).

Notably, our data showed that the weighted mean expression of the HWG of CD8$^+$ T cell trajectories (Methods) are associated with responses to immune checkpoint inhibitors (ICIs) in multiple independent cohorts (Fig. 5c-d, Supplementary Fig. 8f). For instance, the weighted means of HWG of trajectories 1 and 2 in NSCLC which are associated with high activities of cytotoxic T cells, predicted better responses to anti-PD-130 and anti-CTLA-4 (Cho et al., 2020; Liu et al., 2018), as well as their combination therapies (Auslander et al., 2018) ($AUC \in \{0.813, 0.964\}$, P < 0.1, Supplementary Fig. 7). Similarly, HWG pertaining to trajectory 1 in CRC is associated with high proportion of $EMRA$(87%, phase 1), hence better responses to immunotherapies in 4 cohorts

(Auslander et al., 2018; Cho et al., 2020; Jung et al., 2019; Liu et al., 2018) ($AUC \in \{0.796, 0.964\}$, $P < 0.01$, Supplementary Fig. 9).

Taken together, the factorization of scRNA-seq data by MGPfact provides highly relevant gene expression signatures of the fate of tumor associated CD8$^+$ T cells, which advances the understanding of the evolution of tumor immune microenvironment (TIME) and predicts clinical outcomes.

Fig. 5. Highly weighted genes (HWG) of the bifurcation processes of CD8$^+$ T cells serve as reliable indicators for clinical outcome and ICI treatment response. a. Gene expression signatures (GES) corresponding to HWG in CD8$^+$ T cells trajectory 1 and 2 in NSCLC predict overall survival of the TCGA-LUAD cohort. **b.** Gene expression signatures (GES) corresponding to HWG in CD8$^+$ T cells trajectory 1 in CRC predict overall survival of the TCGA-COAD cohort. **c.** ROC curve showing the weighted mean of HWG in Trajectories 1 and 2 in NSCLC significantly associated with ICI response across 3 independent studies. **d.** ROC curve showing the weighted mean of HWG in trajectories 1 and 2 in CRC significantly associated with ICI response across 4 independent studies.

MGPfact identified new subtypes of CD8⁺ T cells with clinical implications

Furthermore, the consensus trajectories of tumor-associated CD8⁺ T cells inferred by MGPfact from NSCLC and CRC revealed new subtypes of lymphocytes. In NSCLC, we characterized CD8-ZNF683-T1 (LEF+/TBX21-) and CD8-ZNF683-T2 (LEF+/TBX21+) from CD8-ZNF683 (Fig. 6a, Supplementary Fig. 6c-d). The CD8-ZNF683-T2 cells highly expressed genes associated with "pre-exhausted" state, such as ITGAL, SAMD3, and SLAMF7 (Pritchard et al., 2023), many of which are target genes of TBX21. In contrast, CD8-ZNF683-T1 showed lower expression of these genes, hence repellency to the "pre-exhausted" state (Fig. 6b, Supplementary Table 11). In CRC, we identified two subtypes of effector memory T cells (CD8-GZMK), CD8-GZMK-T1 (EOMES-/BHLHE40+) and CD8-GZMK-T2 (EOMES-/BHLHE40-) (Fig. 6d, Supplementary Fig. 8c-d). CD8-GZMK-T2 strongly resembles CD8-GZMK, and potentially differentiating into CD8-CD6 (resident memory T cells, T_{RM}) and CD8-CD160 (intraepithelial lymphocytes, IEL); whereas CD8-GZMK-T1 cells demonstrated higher expression of GZMB, indicating an active cytotoxic cell-killing capability (Trapani, 2001). Simultaneously, these cells also marked by high expression levels of immune related genes, including OASL, RBPJ, and CTLA4, which are known targets of BHLHE40 (Lutter et al., 2022; Salmon et al., 2022), implying that BHLHE40 is a modulator of the higher effector activity in CD8-GZMK-T1 (Fig. 6e, Supplementary Table 12).

We further derived scores based on the differentially expressed genes of CD8-ZNF683-T1 and CD8-GZMK-T1 (Methods), as measures of the fraction of each subtype in cancer cohorts. In the LUAD cohort of TCGA, increased fraction of CD8-ZNF683-T1 in TIME was associated with favorable outcomes (Fig. 6c). And increased fractions of CD8-GZMK-T1 in TIME were associated with better responses to ICI therapy across 3 independent cohorts[31–33] (Fig. 6f, Supplementary Fig. 10, $AUC \in \{0.782, 0.917\}$, P < 0.1), which were treated with anti-PD-1, anti-CTLA-4 and their combination therapies.

In conclusion, our data showed the cellular trajectory inferred by MGPfact can be used to elucidate the complex evolutionary processes of tumor-associated CD8⁺ T cells, and further inform the characterization of new subtypes of T cells with significant clinical implications.

Fig. 6. MGPfact serves as an effective approach for characterization of new cellular subtypes.
a. The consensus trajectory of tumor-associated CD8+ T cells in NSCLC identified CD8-ZNF683-T1
and CD8-ZNF683-T2 as two subtypes of CD8-ZNF683, which are influenced by TBX21. **b.** Selected
differently expressed genes between CD8-ZNF683-T1 and CD8-ZNF683-T2 (|logfc| > 0.25,

adjusted P-value < 0.1). **c.** High expression of CD8-ZNF683-T1 signatures predicts good overall survival in the TCGA LUAD cohort (Methods). P-values were calculated through multivariate Cox regression analysis, and HR represents hazard ratio. **d.** The consensus trajectory of tumor-associated CD8$^+$ T cells in CRC identified CD8-GZMK-T1 and CD8-GZMK-T2 as two subtypes of CD8-GZMK. **e.** Selected differently expressed genes between CD8-GZMK-T1 and CD8-GZMK-T2 (|logfc| > 0.25, adjusted P-value < 0.1). **f.** ROC curve showing high expression of CD8-GZMK-T1 signature associated with ICI treatment response in three independent studies. The consensus trajectory is formed by merging three bifurcation processes. Each colored circle represents a landmark (MURP), indicating the of cell type.

Discussion

Single-cell RNA sequencing (scRNA-seq) provides a direct, quantitative snapshot of a population of cells in certain biological conditions, thereby revealing the actual cell states and functions. Although existing clustering and embedding algorithms can effectively reveal discrete biological states of cells, these methods become less efficient when depicting continuous evolving of cells over the temporal domain. The introduction of manifold learning offers a new dimension for discovery of relevant biological knowledge in cell fate determination, allowing for a better representation of continuous changes in cells, especially in time-dependent processes such as development, differentiation, and clonal evolution. However, current manifold learning methods face major limitations, such as the need for prior information on pseudotime and cell clustering, and lack of explainability, which restricts their applicability. Additionally, many existing trajectory inference methods do not support gene selection, making it difficult to annotate the results to known biological entities, thereby hindering the interpretation of results and subsequent functional studies.

We developed MGPfact to overcome the limitations of the existing methods. Inspired by recent studies, MGPfact model the cell fate as mixture of Gaussian processes, which accommodate both continuous evolution pathway and biphasic destiny of cell fate. Thus, MGPfact is capable to distinguish discrete and continuous events in the same trajectory. In addition, by factorizing the mixture Gaussian processes, MGPfact offers the advantage to select genes corresponding to each bifurcation process and thereby enable full biological annotation and interpretation of the trajectory. As a validation, we showed that gene-selection by MGPfact consistently recapitulated the development of microglia and tumor-associated CD8$^+$ T cells; and recovered key regulators of distinct cell fate. So far, MGPfact is the only model-based manifold-learning framework which

factorizes complex development trajectories into independent bifurcation processes of gene sets.

We conducted a comprehensive comparison of MGPfact with existing TI methods from various perspectives. This comparison included the correlation of cell sorting, accuracy of branch allocation, similarity of topological structures and differentially expressed features. It ought to be mentioned that number of principal components used should be determined by the intrinsic biological characteristics of the cell fate-determination. Our experiment based on a limited number of datasets may not represent more complex scenarios in other cell types. For the overall TI-performance, MGPfact demonstrated leading performance across 239 datasets, second only to TinGa. For the performance in branch allocation, which directly reflect the fitness to the outcomes of cell-fate, MGPfact outperformed its counterparts, especially in the topology groups of linear and bifurcation. As for $wcor_{features}$ and cor_{dist}, MGPfact performed less well mainly for two reasons. First, MGPfact is designed for bifurcation topology in the cellular trajectory, hence less efficient in inferring complex topologies. Then, MGPfact inference is based on selected gene sets instead of the whole transcriptome, the resulted trajectory correspond only to the bifurcation processes of interest hence does not necessarily reflect the whole topology of cellular trajectory. Furthermore, MGPfact performed significantly better in trajectory prediction in real cell population compared to the synthetic ones, suggesting the algorithm fit better to the true biological variation and noises.

To reconstruct the trajectory of cell fate, we merged all the bifurcation processes into a consensus trajectory. In the validation by microglia and tumor-associated CD8[+] T cells, the consensus trajectory revealed highly consistent findings recovering known biology and the marker genes of specific cell states which further inform the transcription factor (TF) determining the fate of cell. In addition, the consensus trajectory revealed new subtypes of cells demonstrating highly relevant transcriptional characteristics. Particularly, we reported new subtypes of tumor associated CD8[+] T cells characterized by different TBX21 and BHLHE40 activity, both of which are known regulators of CD8[+] T cell functionality (Lutter et al., 2022; Pritchard et al., 2023; Salmon et al., 2022; Trapani, 2001). These data suggest that MGPfact is capable to discover gene modules with strong and consistent transcriptional background hence better interpretability. Moreover, the results of MGPfact demonstrated strong clinical relevance. Using MGPfact we retrieved gene expression signatures (GES) which quantitatively measure the propensity and fraction of different fates of CD8[+] T cell. These signatures correspond to important biological processes of T cell activity; and predict clinical outcome and ICI treatment responses from transcriptome data of bulk tumor biopsies, independent of any endogenous feature of the tumor cells.

Nevertheless, MGPfact also has some limitations, which shall be addressed in future study.

Firstly, the complex definition of the bifurcation kernel introduces unfavorable singularity to the
Gaussian kernel when considering highly complex trajectories. Additionally, the current trajectory
inference by MGPfact is solely based on the temporal domain, neglecting bifurcation processes
occurring in space. To overcome these limitations, future models should incorporate spatial
dynamics of transcription and RNA velocity data to provide more comprehensive insights of cell fate.
Moreover, the reconstruction of cellular trajectories by MGPfact implies independence of each
bifurcation processes, which may not reflect real cellular behavior. Therefore, predictions from
MGPfact should be interpreted with caution and validated experimentally.

Methods

Benchmarking MGPfact to state-of-the-art methods

We adopted a comprehensive evaluation framework from previous scRNA-seq study to assess
the TI performance of MGPfact (Saelens et al., 2019; Smolander et al., 2022; Todorov et al., 2020).
The validation dataset comprises 110 real data and 229 synthetic data, encompassing 9 different
cellular trajectory topologies. The ground truth of cellular trajectories of each dataset were inferred
and validated by the original study (Saelens et al., 2019). The synthetic datasets were generated
using four simulators: dyngen (Saelens et al., 2019), dyntoy (Saelens et al., 2019), PROSSTT
(Papadopoulos et al., 2019), and Splatter (Zappia et al., 2017), each modeling different trajectory
topologies such as linear, branching, and cyclic. Splatter simulates branching events by setting
expression states and transition probabilities, dyntoy generates random expression gradients to
reflect dynamic changes, and dyngen focuses on complex branching structures within gene
regulatory networks.

The evaluation of TI performance was based on five metrics.

1) The Hamming-Ipsen-Mikhailov (HIM) distance is a metric for assessing similarity between
two topological structures. It integrates the normalized Hamming distance, which highlights
differences in edge lengths, with the Ipsen-Mikhailov distance, which focuses on similarities
in degree distributions. By linearly combining these two measures, the HIM distance offers
a comprehensive evaluation of both local and global structural differences.

2) The $F1_{branches}$ is a metric used to evaluate a model's accuracy in branch allocation. It
represents the harmonic mean of precision and recall, effectively capturing the performance
of branch identification. In trajectory inference, $F1_{branches}$ are calculated by assessing the

similarity between predicted and actual trajectory branches, emphasizing the Jaccard similarity of branch pairs.

3) This cor_{dist} metric measures similarity in intercellular distances between predicted and actual trajectories. It evaluates model accuracy in cell ordering by comparing relative positions of paired cells, highlighting changes in cell positions across states and reflecting cell differentiation dynamics.

4) The $wcor_{features}$ metric evaluates the similarity of key features, such as differentially expressed genes, between predicted and actual trajectories. Using weighted Pearson correlation, it highlights consistent features, reflecting the model's ability to capture biological variation. This helps identify crucial genes in trajectories and understand the molecular mechanisms of cell state transitions.

5) The $overall$ score is calculated by taking the geometric mean of the four aforementioned metrics, providing an assessment of overall performance.

The dataset is divided into two groups: a training set and a testing set (Supplementary Fig. 1). We use 100 training datasets to perform the following tasks:

1) Determine the optimal number of trajectories: With 3 set as the default for the number of factorized trajectories, we tested other values (1, 2, 4, and 5) and used paired T-tests to assess whether there are significant changes in MGPfact's prediction results under different parameter settings.

2) Verify the critical role of MURP: Randomly select 20, 40, 60, 80, and 100 cells for trajectory inference, map the inference results back to the original data using the KNN graph structure, and compare the prediction results with those obtained through MURP downsampling.

3) Robustness analysis of the consensus trajectory topology: Perform 60%, 70%, 80%, and 90% sampling on the original data, and then calculate the HIM similarity between the consensus trajectory predictions of MGPfact with and without sampling. A higher score indicates better robustness of the method.

Subsequently, in 239 test datasets, we compare MGPfact with 7 state-of-the-art TI methods using the aforementioned metrics, including Monocle DDRTree (Qiu et al., 2017b, 2017a), TSCAN (Ji and Ji, 2016), and DPT (Haghverdi et al., 2016), as well as four new methods from recent studies: Monocle 3 (Cao et al., 2019), scShaper (Smolander et al., 2022), scFates Tree (Faure et al., 2023), and TinGa (Todorov et al., 2020).

The experimental comparisons were conducted on a CentOS system equipped with 48 CPU cores running at 2.2GHz and 250GB of memory. To ensure a uniform comparison, all experiments

were performed using a single CPU core. For MGPfact, we tested each resulted trajectory and selected the one with the best "overall" score for comparison. For the other 7 methods, default settings are used unless otherwise specified.

Application of MGPfact in a Microglia Single-cell RNA-seq Dataset

We utilized the MGPfact reconstructed the developmental trajectory of microglia from a scRNA-seq dataset, including immature microglia (IM), proliferation-associated microglia (PAM), and homeostatic microglia (HM) (Li et al., 2019). In this analysis, we provided a detailed explanation of the analytical steps of MGPfact and the key results. Firstly, we identified three independent developmental pathways and pinpointed HWG associated with each bifurcation process. Then, we retrieved highly active regulons within each bifurcation process, tracing back to the potential influential determinants (transcription factors) in the development of microglia. Finally, we combined all the bifurcation processes into a consensus trajectory (Supplementary methods), which recovered the known biology of disease-related microglia (PAM), as represented by distinct cellular state and marker genes.

Predicting the evolutionary trajectory of tumor-associated CD8$^+$ T cells

We utilized MGPfact to conducted an exploratory analysis of the evolution of tumor-associated CD8$^+$ T cells of non-small cell lung cancer (NSCLC) and colorectal cancer (CRC). We evaluated the goodness-of-fitness of the consensus trajectories from MGPfact to the CD8$^+$ T cell subtypes identified in the original studies. For comparison, we used Monocle 2 (Qiu et al., 2017b, 2017a) as a baseline model.

For the survival analysis, we extracted gene expression signatures (GES) from each independent bifurcation process to develop classifiers for evolutionary propensity of CD8$^+$ T cells towards specific fates, based on which we stratified TCGA cancer cohorts and verified their association with clinical outcome.

To evaluated the association to ICI responses, we used HWG to retrieve key transcription factors related to each bifurcation process and characterized the underlying biological processes. We then assessed their connection to immunotherapy response using weighted means of the HWG.

Finally, we identified new subtypes based on distinct endpoints of the consensus trajectory and validated their association with clinical outcome and immunotherapy response using mean of the differently expressed gene (DEG, Supplementary Methods).

Single-cell sequencing data processing

We obtained the original mouse developmental microglia single-cell sequencing data from the GEO accession number GSE123025 (Li et al., 2019). Using Seurat (Butler et al., 2018; Stuart et al., 2019), we replicated the processing steps described in the original study: 1) Normalization by dividing gene expression values by total RNA count, followed by log2 transformation; 2) Selection of highly variable genes (HVGs) using Seurat's mean.var.plot function, with controlled average expression $[0.0125,3]$ and variance $[0.5, \infty]$; 3) Scaling and centering of the normalized matrix for HVGs, with regression of cell cycle effects. After preprocessing, we grouped cells into IM (P7-C0), PAM (P7-C1 and P7-GPNMB$^+$CLEC7A$^+$), and HM (P60), resulting in a 4,889-gene expression matrix across 1,009 cells.

For analyzing tumor-associated CD8$^+$ T cells, we utilized scRNA-seq data from lung cancer (GSE99254) (Guo et al., 2018) and colorectal cancer (GSE108989) (Zhang et al., 2018) in the GEO database. We extracted preprocessed and centralized gene expression matrices of CD8$^+$ T cells and analyzed them using the same genes and the same method (Monocle 2[9,10]) as in the original papers or MGPfact for   trajectory construction for a direct comparison. The NSCLC data yielded an 888-gene expression matrix across 3,700 cells, while the CRC data resulted in a 700-gene expression matrix across 3,177 cells.

Functional enrichment of highly weighted genes

For the highly weighted genes (HWG, absolute gene weight > 0.05) obtained from independent bifurcation processes, we utilized the R package clusterprofiler (Yu et al., 2012) to perform functional annotation using GO terms (Consortium, 2004), including biological process (BP), cellular component (CC), and molecular function (MF). The results with a Benjamini–Hochberg-adjusted P value less than 0.05 were retained.

Transcription factor program analysis

To comprehensively assess key regulatory factors within each independent trajectory, we performed SCENIC (Aibar et al., 2017) transcription factor regulatory program estimation for each analysis case. GENIE3 (Huynh-Thu et al., 2010) was used to identify co-expressed modules from the results of MURP (Ren et al., 2022) downsampling. RCisTarget (Aerts et al., 2010; Aibar et al., 2017) was then used to identify regulons before AUCell (Aibar et al., 2017) was used to estimate the

activity of each regulon. Each regulon comprises a specific transcription factor and its target genes. Finally, we utilize gene weights obtained from MGPfact analysis to evaluate the distinct impact of top regulons on each trajectory.

Generating the consensus trajectory

Following MGPfact decomposition, we obtained multiple independent bifurcative trajectories, each corresponds to a binary tree within the temporal domain. These trajectories were then merged to construct a coherent diffusion tree, representing the consensus trajectory of cells' fate. The merging process involves initially sorting all trajectories by their bifurcation time. The first (earliest) bifurcative trajectory is chosen as the initial framework, and subsequent trajectories are integrated to the initial framework iteratively by adding the corresponding branches at the bifurcation timepoints. As a result, the trajectories are ultimately merged into a comprehensive binary tree, serving as the consensus trajectory.

Assessing consistency of MGPfact-derived CD8$^+$ T Cell subtype trajectories

In the case study of CD8$^+$ T cells, by combining independent trajectories, we derive a consensus trajectory representing the complex developmental pathway. To assess the goodness-of-fitness to the CD8$^+$ T cell subtypes from the original study, we classified the trajectories into several states based on bifurcation points, each corresponding to a distinct stage of the evolutionary process. Then, we evaluated the interactive effects between the states and pseudotime on the fraction of the cell types using F-test (ANOVA). The resulted R-squared (R2), P-values, and F-statistics were used to evaluate the goodness-of-fitness of the models tested hence the explanatory power. For comparison, we used the Monocle 2 as the baseline model for trajectory inference. The differentiation trajectories of Monocle 2 were replicated following the workflow in the original study (Guo et al., 2018; Zhang et al., 2018).

Survival analysis

We assessed the association of bifurcation processes and specific cell types with the clinical outcomes two cohorts of lung adenocarcinoma (LUAD, N=563) and colon adenocarcinoma (COAD, N=320) data from The Cancer Genome Atlas (TCGA). The gene expression and clinical data were

downloaded from UCSC Xena platform (http://xena.ucsc.edu/).

We assessed the survival impacts of NSCLC and CRC bifurcation processes in the TCGA LUAD and COAD cohorts, respectively. For each independent bifurcation process, we defined Gene Expression Signatures (GES) by the mean expression vectors of all trajectories in phase 1 and 2, respectively. Subsequently, we calculated the Peason's correlation coefficients for each individual expression profile in TCGA LUAD or COAD cohorts to the phase 1 and 2 GES, where higher correlation correspond to stronger propensity to specific cell fate. This allowed us to classify patients into two groups: those exhibiting propensity to phase 1 and those exhibiting propensity to phase 2.

To assess the survival impacts of specific cell states defined by the consensus trajectory, we developed a CD8-ZNF683-T1 score based on the signed average expression level of DEG associated with CD8-ZNF683-T1 (Supplementary methods). Subsequently, we classified the TCGA LUAD cohorts into two groups using the median of CD8-ZNF683-T1 scores, identifying those demonstrating a propensity towards CD8-ZNF683-T1 and those demonstrating a propensity towards CD8-ZNF683-T2.

To adjust for possible confounding effects, the relevant clinical features including age, sex and tumor stage were used as covariates. The Cox regression model was implemented using R-4.2 package "survival". And we generated Kaplan-Meier survival curves based on different classifiers to illustrate differences in survival time and report the statistical significance based on Log-rank test.

Immune-checkpoint inhibitor treatment response analysis

For the prediction of Immune-checkpoint inhibitor treatment response, we collected four datasets containing ICI treatment responses. These datasets consist of two non-small cell lung cancer related datasets (GSE135222 (Jung et al., 2019), n=27; GSE126044 (Cho et al., 2020), n=16) and two melanoma related datasets (GSE115821 (Auslander et al., 2018), n=14; TCGA-MENO (Liu et al., 2018), n=10). All data were processed with DESeq2 (Love et al., 2014) to fit gene dispersion to a negative binomial distribution, normalize raw counts, and stabilize variance, achieving standardization.

To validate the bifurcation processes identified by MGPfact predicting patients' response to immune-checkpoint inhibitor (ICI) treatments, we selected highly weighted genes (HWG) with absolute weights greater than 0.05 from each independent bifurcation process. Then, we calculated the weighted mean expression of HWG in each ICI dataset to generate ROC curves for patient drug response.

To validate the specific cell states defined by the consensus trajectory predicting patients' response to ICI treatments, we used CD8-GZMK-T1 score based on the average expression level differences of upregulated and downregulated genes associated with CD8- GZMK-T1. Then, we calculated the CD8-GZMK-T1 score in each ICI dataset to generate ROC curves for patient drug response.

Code availability

We have developed a comprehensive workflow for MGPfact. Firstly, a Docker container enables one-click program execution (details at: https://github.com/renjun0324/ti_mgpfact). Additionally, to fully harness MGPfact's capabilities, we have created the R package MGPfactR, accessible at: https://github.com/renjun0324/MGPfactR. Within this workflow, MCMC sampling for model parameter estimation is carried out using the Mamba library in the Julia environment. The related Julia package can be found here: https://github.com/renjun0324/MGPfact.jl. Additionally, other analysis scripts can be found on GitHub at https://github.com/renjun0324/mgpfact_paper. And the scFates Tree used in this paper is available as a performance comparison Docker container, constructed using the dendritic trajectory process in scFates (Faure et al., 2023), accessible at: https://github.com/renjun0324/ti_scfates_tree.

Data availability

The datasets used for performance comparison are archived on Zenodo by Saelens *et al*. (Saelens et al., 2019), with processed real and synthetic datasets available at https://doi.org/10.5281/zenodo.1443566. Data for specific cell instances of microglia and CD8$^+$ T cells can be obtained from the GEO database with the following accession numbers: GSE123025 (Li et al., 2019), GSE99254 (Guo et al., 2018), and GSE108989 (Zhang et al., 2018). Expression matrices related to immune-checkpoint inhibitors (ICI) and clinical response information are downloadable from the CTR-DB (https://ctrdb.cloudna.cn).

Acknowledgments

We would like to thank Qingyun Li for providing the cell labels for the microglial dataset(Li et al., 2019). We would also like to express our gratitude to Nengming Xiao, and Guo Fu for their valuable

Sources of Funding

This work was supported by the Fundamental Research Funds for the National Natural Science Foundation of China [82272944 to QL]; the National Natural Science Foundation of China [82203420 to JG].

Author Contributions

The project was conceived by Qiyuan Li. The model construction, data collection and analytical validation was done by Jun Ren, with the assistance of Yudi Hu and Jintao Guo. The manuscript was written by Jun Ren and Xuejing Lyu, with the assistance of Ying Zhou and Xiaodong Shi. All authors read and approved the final manuscript.

Reference

Aerts S, Quan X-J, Claeys A, Naval Sanchez M, Tate P, Yan J, Hassan BA. 2010. Robust Target Gene Discovery through Transcriptome Perturbations and Genome-Wide Enhancer Predictions in Drosophila Uncovers a Regulatory Basis for Sensory Specification. *PLoS Biol* **8**:e1000435. doi:10.1371/journal.pbio.1000435

Aibar S, González-Blas CB, Moerman T, Huynh-Thu VA, Imrichova H, Hulselmans G, Rambow F, Marine J-C, Geurts P, Aerts J, van den Oord J, Atak ZK, Wouters J, Aerts S. 2017. SCENIC: single-cell regulatory network inference and clustering. *Nat Methods* **14**:1083–1086. doi:10.1038/nmeth.4463

Anderson SR, Roberts JM, Ghena N, Irvin EA, Schwakopf J, Cooperstein IB, Bosco A, Vetter ML. 2022. Neuronal apoptosis drives remodeling states of microglia and shifts in survival pathway dependence. *Elife* **11**:e76564.

Auslander N, Zhang G, Lee JS, Frederick DT, Miao B, Moll T, Tian T, Wei Z, Madan S, Sullivan RJ, Boland G, Flaherty K, Herlyn M, Ruppin E. 2018. Robust prediction of response to immune checkpoint blockade therapy in metastatic melanoma. *Nat Med* **24**:1545–1549. doi:10.1038/s41591-018-0157-9

B J S. 2014. Mamba: Markov Chain Monte Carlo for Bayesian Analysis in julia.

Becht E, McInnes L, Healy J, Dutertre C-A, Kwok IWH, Ng LG, Ginhoux F, Newell EW. 2019. Dimensionality reduction for visualizing single-cell data using UMAP. *Nat Biotechnol* **37**:38–44. doi:10.1038/nbt.4314

Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. 2018. Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol* **36**:411–420. doi:10.1038/nbt.4096

Cao J, Spielmann M, Qiu X, Huang X, Ibrahim DM, Hill AJ, Zhang F, Mundlos S, Christiansen L, Steemers FJ, Trapnell C, Shendure J. 2019. The single-cell transcriptional landscape of mammalian organogenesis. *Nature* **566**:496–502. doi:10.1038/s41586-019-0969-x

Cho J-W, Hong MH, Ha S-J, Kim Y-J, Cho BC, Lee I, Kim HR. 2020. Genome-wide identification of differentially methylated promoters and enhancers associated with response to anti-PD-1 therapy in non-small cell lung cancer. *Exp Mol Med* **52**:1550–1563. doi:10.1038/s12276-020-00493-8

Consortium GO. 2004. The Gene Ontology (GO) database and informatics resource. *Nucleic acids research* **32**:D258–D261.

Costa F, Grün D, Backofen R. 2018. GraphDDP: a graph-embedding approach to detect differentiation pathways

in single-cell-data using prior class knowledge. *Nat Commun* **9**:3685. doi:10.1038/s41467-018-05988-7

Dresselhaus EC, Meffert MK. 2019. Cellular Specificity of NF-κB Function in the Nervous System. *Front Immunol* **10**:1043. doi:10.3389/fimmu.2019.01043

Faure L, Soldatov R, Kharchenko PV, Adameyko I. 2023. scFates: a scalable python package for advanced pseudotime and bifurcation analysis from single-cell data. *Bioinformatics* **39**:btac746. doi:10.1093/bioinformatics/btac746

Fritzke B. 1994. A growing neural gas network learns topologies. *Advances in neural information processing systems* **7**.

Guo X, Zhang Y, Zheng L, Zheng C, Song J, Zhang Q, Kang B, Liu Z, Jin L, Xing R, Gao R, Zhang L, Dong M, Hu X, Ren X, Kirchhoff D, Roider HG, Yan T, Zhang Z. 2018. Global characterization of T cells in non-small-cell lung cancer by single-cell sequencing. *Nat Med* **24**:978–985. doi:10.1038/s41591-018-0045-3

Guzmán AU. n.d. Single-cell RNA sequencing of spinal cord microglia in a mouse model of neuropathic pain.

Haghverdi L, Buettner F, Theis FJ. 2015. Diffusion maps for high-dimensional single-cell analysis of differentiation data. *Bioinformatics* **31**:2989–2998. doi:10.1093/bioinformatics/btv325

Haghverdi L, Büttner M, Wolf FA, Buettner F, Theis FJ. 2016. Diffusion pseudotime robustly reconstructs lineage branching. *Nat Methods* **13**:845–848. doi:10.1038/nmeth.3971

Hugo W, Zaretsky JM, Sun L, Song C, Moreno BH, Hu-Lieskovan S, Berent-Maoz B, Pang J, Chmielowski B, Cherry G, Seja E, Lomeli S, Kong X, Kelley MC, Sosman JA, Johnson DB, Ribas A, Lo RS. 2016. Genomic and Transcriptomic Features of Response to Anti-PD-1 Therapy in Metastatic Melanoma. *Cell* **165**:35–44. doi:10.1016/j.cell.2016.02.065

Huynh-Thu VA, Irrthum A, Wehenkel L, Geurts P. 2010. Inferring Regulatory Networks from Expression Data Using Tree-Based Methods. *PLoS ONE* **5**:e12776. doi:10.1371/journal.pone.0012776

Ji Z, Ji H. 2016. TSCAN: Pseudo-time reconstruction and evaluation in single-cell RNA-seq analysis. *Nucleic Acids Res* **44**:e117–e117. doi:10.1093/nar/gkw430

Jung H, Kim HS, Kim JY, Sun J-M, Ahn JS, Ahn M-J, Park K, Esteller M, Lee S-H, Choi JK. 2019. DNA methylation loss promotes immune evasion of tumours with high mutation and copy number load. *Nat Commun* **10**:4278. doi:10.1038/s41467-019-12159-9

Lange M, Bergen V, Klein M, Setty M, Reuter B, Bakhti M, Lickert H, Ansari M, Schniering J, Schiller HB, Pe'er D, Theis FJ. 2022. CellRank for directed single-cell fate mapping. *Nat Methods* **19**:159–170. doi:10.1038/s41592-021-01346-6

Li Q. 2023. scTour: a deep learning architecture for robust inference and accurate prediction of cellular dynamics. *Genome Biology*.

Li Q, Cheng Z, Zhou L, Darmanis S, Neff NF, Okamoto J, Gulati G, Bennett ML, Sun LO, Clarke LE, Marschallinger J, Yu G, Quake SR, Wyss-Coray T, Barres BA. 2019. Developmental Heterogeneity of Microglia and Brain Myeloid Cells Revealed by Deep Single-Cell RNA Sequencing. *Neuron* **101**:207-223.e10. doi:10.1016/j.neuron.2018.12.006

Liu Jianfang, Lichtenberg T, Hoadley KA, Poisson LM, Lazar AJ, Cherniack AD, Kovatich AJ, Benz CC, Levine DA, Lee AV, Omberg L, Wolf DM, Shriver CD, Thorsson V, Hu H, Caesar-Johnson SJ, Demchok JA, Felau I, Kasapi M, Ferguson ML, Hutter CM, Sofia HJ, Tarnuzzer R, Wang Z, Yang L, Zenklusen JC, Zhang J (Julia), Chudamani S, Liu Jia, Lolla L, Naresh R, Pihl T, Sun Q, Wan Y, Wu Y, Cho J, DeFreitas T, Frazer S, Gehlenborg N, Getz G, Heiman DI, Kim J, Lawrence MS, Lin P, Meier S, Noble MS, Saksena G, Voet D, Zhang Hailei, Bernard B, Chambwe N, Dhankani V, Knijnenburg T, Kramer R, Leinonen K, Liu Y, Miller M, Reynolds S, Shmulevich I, Thorsson V, Zhang W, Akbani R, Broom BM, Hegde AM, Ju Z, Kanchi RS, Korkut A, Li J, Liang H, Ling S, Liu W, Lu Y, Mills GB, Ng K-S, Rao A, Ryan M, Wang Jing, Weinstein JN, Zhang J, Abeshouse A, Armenia J, Chakravarty D, Chatila WK, De Bruijn I, Gao J, Gross BE, Heins ZJ, Kundra R, La K, Ladanyi M, Luna A, Nissan MG, Ochoa A, Phillips SM, Reznik E, Sanchez-Vega F, Sander C, Schultz N, Sheridan R, Sumer SO, Sun Y, Taylor BS, Wang Jioajiao, Zhang Hongxin, Anur P, Peto M, Spellman P, Benz C, Stuart JM, Wong CK, Yau C, Hayes DN, Parker JS, Wilkerson MD, Ally A, Balasundaram M, Bowlby R, Brooks D, Carlsen R, Chuah E, Dhalla N, Holt R, Jones SJM, Kasaian K, Lee D, Ma Y, Marra MA, Mayo M, Moore RA, Mungall AJ, Mungall K, Robertson AG, Sadeghi S, Schein

JE, Sipahimalani P, Tam A, Thiessen N, Tse K, Wong T, Berger AC, Beroukhim R, Cherniack AD, Cibulskis C, Gabriel SB, Gao GF, Ha G, Meyerson M, Schumacher SE, Shih J, Kucherlapati MH, Kucherlapati RS, Baylin S, Cope L, Danilova L, Bootwalla MS, Lai PH, Maglinte DT, Van Den Berg DJ, Weisenberger DJ, Auman JT, Balu S, Bodenheimer T, Fan C, Hoadley KA, Hoyle AP, Jefferys SR, Jones CD, Meng S, Mieczkowski PA, Mose LE, Perou AH, Perou CM, Roach J, Shi Y, Simons JV, Skelly T, Soloway MG, Tan D, Veluvolu U, Fan H, Hinoue T, Laird PW, Shen H, Zhou W, Bellair M, Chang K, Covington K, Creighton CJ, Dinh H, Doddapaneni H, Donehower LA, Drummond J, Gibbs RA, Glenn R, Hale W, Han Y, Hu J, Korchina V, Lee S, Lewis L, Li W, Liu X, Morgan M, Morton D, Muzny D, Santibanez J, Sheth M, Shinbro E, Wang L, Wang M, Wheeler DA, Xi L, Zhao F, Hess J, Appelbaum EL, Bailey M, Cordes MG, Ding L, Fronick CC, Fulton LA, Fulton RS, Kandoth C, Mardis ER, McLellan MD, Miller CA, Schmidt HK, Wilson RK, Crain D, Curley E, Gardner J, Lau K, Mallery D, Morris S, Paulauskis J, Penny R, Shelton C, Shelton T, Sherman M, Thompson E, Yena P, Bowen J, Gastier-Foster JM, Gerken M, Leraas KM, Lichtenberg TM, Ramirez NC, Wise L, Zmuda E, Corcoran N, Costello T, Hovens C, Carvalho AL, De Carvalho AC, Fregnani JH, Longatto-Filho A, Reis RM, Scapulatempo-Neto C, Silveira HCS, Vidal DO, Burnette A, Eschbacher J, Hermes B, Noss A, Singh R, Anderson ML, Castro PD, Ittmann M, Huntsman D, Kohl B, Le X, Thorp R, Andry C, Duffy ER, Lyadov V, Paklina O, Setdikova G, Shabunin A, Tavobilov M, McPherson C, Warnick R, Berkowitz R, Cramer D, Feltmate C, Horowitz N, Kibel A, Muto M, Raut CP, Malykh A, Barnholtz-Sloan JS, Barrett W, Devine K, Fulop J, Ostrom QT, Shimmel K, Wolinsky Y, Sloan AE, De Rose A, Giuliante F, Goodman M, Karlan BY, Hagedorn CH, Eckman J, Harr J, Myers J, Tucker K, Zach LA, Deyarmin B, Hu H, Kvecher L, Larson C, Mural RJ, Somiari S, Vicha A, Zelinka T, Bennett J, Iacocca M, Rabeno B, Swanson P, Latour M, Lacombe L, Têtu B, Bergeron A, McGraw M, Staugaitis SM, Chabot J, Hibshoosh H, Sepulveda A, Su T, Wang T, Potapova O, Voronina O, Desjardins L, Mariani O, Roman-Roman S, Sastre X, Stern M-H, Cheng F, Signoretti S, Berchuck A, Bigner D, Lipp E, Marks J, McCall S, McLendon R, Secord A, Sharp A, Behera M, Brat DJ, Chen A, Delman K, Force S, Khuri F, Magliocca K, Maithel S, Olson JJ, Owonikoko T, Pickens A, Ramalingam S, Shin DM, Sica G, Van Meir EG, Zhang Hongzheng, Eijckenboom W, Gillis A, Korpershoek E, Looijenga L, Oosterhuis W, Stoop H, Van Kessel KE, Zwarthoff EC, Calatozzolo C, Cuppini L, Cuzzubbo S, DiMeco F, Finocchiaro G, Mattei L, Perin A, Pollo B, Chen C, Houck J, Lohavanichbutr P, Hartmann A, Stoehr C, Stoehr R, Taubert H, Wach S, Wullich B, Kycler W, Murawa D, Wiznerowicz M, Chung K, Edenfield WJ, Martin J, Baudin E, Bubley G, Bueno R, De Rienzo A, Richards WG, Kalkanis S, Mikkelsen T, Noushmehr H, Scarpace L, Girard N, Aymerich M, Campo E, Giné E, Guillermo AL, Van Bang N, Hanh PT, Phu BD, Tang Y, Colman H, Evason K, Dottino PR, Martignetti JA, Gabra H, Juhl H, Akeredolu T, Stepa S, Hoon D, Ahn K, Kang KJ, Beuschlein F, Breggia A, Birrer M, Bell D, Borad M, Bryce AH, Castle E, Chandan V, Cheville J, Copland JA, Farnell M, Flotte T, Giama N, Ho T, Kendrick M, Kocher J-P, Kopp K, Moser C, Nagorney D, O'Brien D, O'Neill BP, Patel T, Petersen G, Que F, Rivera M, Roberts L, Smallridge R, Smyrk T, Stanton M, Thompson RH, Torbenson M, Yang JD, Zhang L, Brimo F, Ajani JA, Angulo Gonzalez AM, Behrens C, Bondaruk J, Broaddus R, Czerniak B, Esmaeli B, Fujimoto J, Gershenwald J, Guo C, Lazar AJ, Logothetis C, Meric-Bernstam F, Moran C, Ramondetta L, Rice D, Sood A, Tamboli P, Thompson T, Troncoso P, Tsao A, Wistuba I, Carter C, Haydu L, Hersey P, Jakrot V, Kakavand H, Kefford R, Lee K, Long G, Mann G, Quinn M, Saw R, Scolyer R, Shannon K, Spillane A, Stretch J, Synott M, Thompson J, Wilmott J, Al-Ahmadie H, Chan TA, Ghossein R, Gopalan A, Levine DA, Reuter V, Singer S, Singh B, Tien NV, Broudy T, Mirsaidi C, Nair P, Drwiega P, Miller J, Smith J, Zaren H, Park J-W, Hung NP, Kebebew E, Linehan WM, Metwalli AR, Pacak K, Pinto PA, Schiffman M, Schmidt LS, Vocke CD, Wentzensen N, Worrell R, Yang H, Moncrieff M, Goparaju C, Melamed J, Pass H, Botnariuc N, Caraman I, Cernat M, Chemencedji I, Clipca A, Doruc S, Gorincioi G, Mura S, Pirtac M, Stancul I, Tcaciuc D, Albert M, Alexopoulou I, Arnaout A, Bartlett J, Engel J, Gilbert S, Parfitt J, Sekhon H, Thomas G, Rassl DM, Rintoul RC, Bifulco C, Tamakawa R, Urba W, Hayward N, Timmers H, Antenucci A, Facciolo F, Grazi G, Marino M, Merola R, De Krijger R, Gimenez-Roqueplo A-P, Piché A, Chevalier S, McKercher G, Birsoy K, Barnett G, Brewer C, Farver C, Naska T, Pennell NA, Raymond D, Schilero C, Smolenski K, Williams F, Morrison C, Borgia JA, Liptay MJ, Pool M, Seder CW, Junker K, Omberg L, Dinkin M, Manikhas G, Alvaro D, Bragazzi MC, Cardinale V, Carpino G, Gaudio E, Chesla D, Cottingham S, Dubina M, Moiseenko F, Dhanasekaran R, Becker K-F, Janssen K-P, Slotta-Huspenina J, Abdel-Rahman MH, Aziz D, Bell S, Cebulla CM, Davis A, Duell R, Elder JB, Hilty J, Kumar B, Lang J, Lehman NL, Mandt R, Nguyen P, Pilarski R, Rai K, Schoenfield L, Senecal K, Wakely P,

Hansen P, Lechan R, Powers J, Tischler A, Grizzle WE, Sexton KC, Kastl A, Henderson J, Porten S, Waldmann J, Fassnacht M, Asa SL, Schadendorf D, Couce M, Graefen M, Huland H, Sauter G, Schlomm T, Simon R, Tennstedt P, Olabode O, Nelson M, Bathe O, Carroll PR, Chan JM, Disaia P, Glenn P, Kelley RK, Landen CN, Phillips J, Prados M, Simko J, Smith-McCune K, VandenBerg S, Roggin K, Fehrenbach A, Kendler A, Sifri S, Steele R, Jimeno A, Carey F, Forgie I, Mannelli M, Carney M, Hernandez B, Campos B, Herold-Mende C, Jungk C, Unterberg A, Von Deimling A, Bossler A, Galbraith J, Jacobus L, Knudson M, Knutson T, Ma D, Milhem M, Sigmund R, Godwin AK, Madan R, Rosenthal HG, Adebamowo C, Adebamowo SN, Boussioutas A, Beer D, Giordano T, Mes-Masson A-M, Saad F, Bocklage T, Landrum L, Mannel R, Moore K, Moxley K, Postier R, Walker J, Zuna R, Feldman M, Valdivieso F, Dhir R, Luketich J, Mora Pinero EM, Quintero-Aguilo M, Carlotti, Jr CG, Dos Santos JS, Kemp R, Sankarankuty A, Tirapelli D, Catto J, Agnew K, Swisher E, Creaney J, Robinson B, Shelley CS, Godwin EM, Kendall S, Shipman C, Bradford C, Carey T, Haddad A, Moyer J, Peterson L, Prince M, Rozek L, Wolf G, Bowman R, Fong KM, Yang I, Korst R, Rathmell WK, Fantacone-Campbell JL, Hooke JA, Kovatich AJ, Shriver CD, DiPersio J, Drake B, Govindan R, Heath S, Ley T, Van Tine B, Westervelt P, Rubin MA, Lee JI, Aredes ND, Mariamidze A. 2018. An Integrated TCGA Pan-Cancer Clinical Data Resource to Drive High-Quality Survival Outcome Analytics. *Cell* **173**:400-416.e11. doi:10.1016/j.cell.2018.02.052

Love M, Anders S, Huber W. 2014. Differential analysis of count data–the DESeq2 package. *Genome Biol* **15**:10–1186.

Lutter L, Van Der Wal MM, Brand EC, Maschmeyer P, Vastert S, Mashreghi M, Van Loosdregt J, Van Wijk F. 2022. Human regulatory T cells locally differentiate and are functionally heterogeneous within the inflamed arthritic joint. *Clin & Trans Imm* **11**:e1420. doi:10.1002/cti2.1420

Nawal HS. n.d. A Systems Biology Perspective of Stem Cell Differentiation into Microglia.

Neal RM. 2003. Slice sampling. *The annals of statistics* **31**:705–767.

Papadopoulos N, Gonzalo PR, Söding J. 2019. PROSSTT: probabilistic simulation of single-cell RNA-seq data for complex differentiation processes. *Bioinformatics* **35**:3517–3519. doi:10.1093/bioinformatics/btz078

Pritchard GH, Phan AT, Christian DA, Blain TJ, Fang Q, Johnson J, Roy NH, Shallberg L, Kedl RM, Hunter CA. 2023. Early T-bet promotes LFA1 upregulation required for CD8+ effector and memory T cell development. *Journal of Experimental Medicine* **220**:e20191287. doi:10.1084/jem.20191287

Qiu X, Hill A, Packer J, Lin D, Ma Y-A, Trapnell C. 2017a. Single-cell mRNA quantification and differential analysis with Census. *Nat Methods* **14**:309–315. doi:10.1038/nmeth.4150

Qiu X, Mao Q, Tang Y, Wang L, Chawla R, Pliner HA, Trapnell C. 2017b. Reversed graph embedding resolves complex single-cell trajectories. *Nat Methods* **14**:979–982. doi:10.1038/nmeth.4402

Ren J, Zhang Q, Zhou Y, Hu Y, Lyu X, Fang H, Yang J, Yu R, Shi X, Li Q. 2022. A downsampling method enables robust clustering and integration of single-cell transcriptome data. *Journal of Biomedical Informatics* **130**:104093. doi:10.1016/j.jbi.2022.104093

Roberts GO, Rosenthal JS. 2009. Examples of adaptive MCMC. *Journal of computational and graphical statistics* **18**:349–367.

Saelens W, Cannoodt R, Todorov H, Saeys Y. 2019. A comparison of single-cell trajectory inference methods. *Nat Biotechnol* **37**:547–554. doi:10.1038/s41587-019-0071-9

Salmon AJ, Shavkunov AS, Miao Q, Jarjour NN, Keshari S, Esaulova E, Williams CD, Ward JP, Highsmith AM, Pineda JE, Taneja R, Chen K, Edelson BT, Gubin MM. 2022. BHLHE40 Regulates the T-Cell Effector Function Required for Tumor Microenvironment Remodeling and Immune Checkpoint Therapy Efficacy. *Cancer Immunology Research* **10**:597–611. doi:10.1158/2326-6066.CIR-21-0129

Schulz E, Speekenbrink M, Krause A. 2018. A tutorial on Gaussian process regression: Modelling, exploring, and exploiting functions. *Journal of Mathematical Psychology* **85**:1–16. doi:10.1016/j.jmp.2018.03.001

Sha Y. 2024. Reconstructing growth and dynamic trajectories from single-cell transcriptomics data **6**.

Smolander J, Junttila S, Venäläinen MS, Elo LL. 2022. scShaper: an ensemble method for fast and accurate linear trajectory inference from single-cell RNA-seq data. *Bioinformatics* **38**:1328–1335. doi:10.1093/bioinformatics/btab831

Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, Hao Y, Stoeckius M, Smibert P, Satija R.

2019. Comprehensive Integration of Single-Cell Data. *Cell* **177**:1888-1902.e21. doi:10.1016/j.cell.2019.05.031

Tierney L. 1994. Markov chains for exploring posterior distributions. *the Annals of Statistics* 1701–1728.

Todorov H, Cannoodt R, Saelens W, Saeys Y. 2020. TinGa: fast and flexible trajectory inference with Growing
Neural Gas. *Bioinformatics* **36**:i66–i74. doi:10.1093/bioinformatics/btaa463

Trapani JA. 2001. Granzymes: a family of lymphocyte granule serine proteases. *Genome Biol* **2**:reviews3014.1.
doi:10.1186/gb-2001-2-12-reviews3014

Van der Maaten L, Hinton G. 2008. Visualizing data using t-SNE. *Journal of machine learning research* **9**.

Wang Q, Lu M, Zhu X, Gu X, Zhang T, Xia C, Yang L, Xu Y, Zhou M. 2022. The role of microglia immunometabolism
in neurodegeneration: Focus on molecular determinants and metabolic intermediates of metabolic reprogramming.
Biomedicine & Pharmacotherapy **153**:113412. doi:10.1016/j.biopha.2022.113412

Yu G, Wang L-G, Han Y, He Q-Y. 2012. clusterProfiler: an R package for comparing biological themes among gene
clusters. *Omics: a journal of integrative biology* **16**:284–287.

Zappia L, Phipson B, Oshlack A. 2017. Splatter: simulation of single-cell RNA sequencing data. *Genome Biol*
18:174. doi:10.1186/s13059-017-1305-0

Zhang L, Yu X, Zheng L, Zhang Y, Li Y, Fang Q, Gao R, Kang B, Zhang Q, Huang JY, Konno H, Guo X, Ye Y, Gao
S, Wang S, Hu X, Ren X, Shen Z, Ouyang W, Zhang Z. 2018. Lineage tracking reveals dynamic relationships of
T cells in colorectal cancer. *Nature* **564**:268–272. doi:10.1038/s41586-018-0694-x

881