Negative regulation of APC/C activation by MAPK-mediated attenuation of Cdc20^{Slp1} under stress

Li Sun†, Xuejin Chen†, Wenjing Shi†, Chunlin Song†, Libo Liu†, Shuang Bai†, Xi Wang, Jiali Chen, Chengyu Jiang, Shuang-min Wang, Zhou-qing Luo, Yamei Wang*, and Quan-wen Jin*

State Key Laboratory of Cellular Stress Biology, School of Life Sciences, Faculty of Medicine and Life Sciences, Xiamen University, Xiamen 361102, Fujian, China

†These authors contributed equally to this work.

*Address correspondence to: Yamei Wang (wangyamei@xmu.edu.cn), or Quan-wen Jin (jinquanwen@xmu.edu.cn)

One-sentence summary: Inhibitory effect on the activation of anaphase promoting complex/cyclosome (APC/C) by MAPK Pmk1

Keywords: anaphase promoting complex/cyclosome (APC/C); spindle assembly checkpoint (SAC); MAPK; Slp1^{Cdc20}; phosphorylation; ubiquitylation

Abstract

Mitotic anaphase onset is a key cellular process tightly regulated by multiple kinases. The involvement of mitogen-activated protein kinases (MAPKs) in this process has been established in *Xenopus* egg extracts. However, the detailed regulatory cascade remains elusive, and also it is unknown whether the MAPKs-dependent mitotic regulation is evolutionarily conserved in the single cell eukaryotic organism such as fission yeast (*Schizosaccharomyces pombe*). Here we show that two MAPKs in *S. pombe* indeed act in concert to restrain anaphase-promoting complex/cyclosome (APC/C) activity upon activation of the spindle assembly checkpoint (SAC). One MAPK, Pmk1, binds and phosphorylates Slp1^{Cdc20}, the co-activator of APC/C. Phosphorylation of Slp1^{Cdc20} by Pmk1, but not by Cdk1, promotes its subsequent ubiquitylation and degradation. Intriguingly, Pmk1-mediated phosphorylation event is also required to sustain SAC under environmental stress. Thus, our study establishes a new underlying molecular mechanism of negative regulation of APC/C by MAPK upon stress stimuli, and provides an unappreciated framework for regulation of anaphase entry in eukaryotic cells.

INTRODUCTION

The evolutionarily conserved mitogen-activated protein kinase (MAPK) signaling pathways regulate multiple cellular functions in eukaryotic organisms in response to a wide variety of environmental cues (Plotnikov et al., 2011). However, different MAPK pathways have been evolved in an organism to integrate diverse signals and fulfill different regulations on various effectors (Cansado et al., 2021; Ronkina and Gaestel, 2022). This makes specifying the MAPKs and substrates involved in a specific physiological process rather complex and challenging.

In late mitosis, the timely polyubiquitylation and subsequent degradation of securin and cyclin B by anaphase-promoting complex/cyclosome (APC/C) play a critical role for anaphase onset and chromosome segregation (Peters, 2006; Sullivan and Morgan, 2007; Yamano, 2019). Given the essential role of APC/C in triggering chromosome segregation, it is not surprising that APC/C is a key molecular target of the spindle assembly checkpoint (SAC), which is an intricate surveillance mechanism that prolongs mitosis until all chromosomes achieve correct bipolar attachments to spindle microtubules (McAinsh and Kops, 2023; Murray, 2011; Musacchio, 2015). Previous studies have revealed that the active MAPK is required for spindle checkpoint activation and APC/C inhibition in *Xenopus* egg extracts or tadpole cells (Chen, 2004; Chung and Chen, 2003; Minshull et al., 1994; Takenaka et al., 1997; Zhao and Chen, 2006). These studies suggested that phosphorylation of several components of SAC, including Cdc20, Mps1 and Bub1, by MAPK serves as regulatory signals to block APC/C activation upon SAC activation (Chen, 2004; Chung and Chen, 2003; Zhao and Chen, 2006). However, all these studies used either the anti-MAPK antibodies or small molecule inhibitor UO126 to inhibit MAPK or MAPK-specific phosphatase MKP-1 to antagonize MAPK activity, these reagents could not distinguish the contributions of different MAPKs with high specificity. Thus, those seemingly defined mechanisms might suffer the ambiguity due to potential specificity issues.

The fission yeast *Schizosaccharomyces pombe* has three MAPK-signaling cascades:

the pheromone signaling pathway (PSP), the stress-activated pathway (SAP) and the

cell integrity pathway (CIP), with Spk1, Sty1 or Pmk1 as MAPK respectively (Figure

1A) (Cansado et al., 2021; Perez and Cansado, 2010). In *S. pombe*, whether MAPKs

play key roles in regulation of the APC/C and spindle checkpoint remains largely

unexplored. Intriguingly, all genes encoding the major components of three

MAPK-signaling cascades in fission yeast are non-essential (Kim et al., 2010), this

provides the possibility for investigations on their potential functions in a genetic

background with complete deletion of relevant genes.

In this work, we set out to directly examine the requirement of the fission yeast

MAPKs in mitotic APC/C activation and anaphase entry. Our results uncovered a

previously unappreciated mechanism in which Pmk1, the MAPK of CIP,

phosphorylates the APC/C activator Slp1 (the Cdc20 homologue in *S. pombe*) upon

SAC activation. Phosphorylation of Slp1^{Cdc20} results in its ubiquitylation and

degradation and thus subsequently impedes APC/C activation and anaphase entry.

This mechanism also operates in response to spindle defects under environmental

stress. Therefore, our finding extends cell-cycle control stages linked to MAPKs from

previously recognized G_2/M transition and cytokinesis to anaphase entry and mitotic

exit in fission yeast. It will also be interesting to address whether this

MAPK-mediated Cdc20 regulation exists as a general mechanism for balancing

spindle checkpoint activity in higher eukaryotes.

RESULTS

SAP and CIP signaling components are required for spindle checkpoint activity

As the first step to examine the possible requirement of the fission yeast MAPKs in

mitotic progression, we tested the sensitivity of deletion mutants of three fission yeast

MAPKs to microtubule-destabilizing drug thiabendazole (TBZ), which has been

routinely used as an indicator of defective kinetochore or spindle checkpoint (Akera et

al., 2015; Saitoh et al., 1997). We noticed that *pmk1Δ* and *sty1Δ* but not *spk1Δ* cells

were extremely sensitive to TBZ, and constitutively activated CIP signaling by

overexpressing Pek1[DD] (Pek1-S234D;T238D) (Sugiura et al., 1999) or activated SAP

signaling by overexpressing Wis1[DD] (Wis1-S469D;T473D) (Shiozaki et al., 1998)

also caused yeast cells to be sensitive to TBZ (Figure 1B; Supplementary Figure 1A).

These observations indicated that Pmk1 and Sty1 pathways might be involved in

mitosis-related processes.

To extend our analysis and precisely quantify whether the SAC is properly activated

and maintained in MAPK mutants, we adopted one well-established assay using the

cold-sensitive β-tubulin mutant *nda3-KM311*, which compromises the

kinetochore-spindle microtubule attachment upon cold treatment (Hiraoka et al.,

1984), to analyze the ability of cells to arrest in mitosis in the absence of spindle

microtubules and enter anaphase when spindle microtubules are reassembled (Bai et

al., 2022; May et al., 2017; Vanoosthuyse and Hardwick, 2009). In this assay, the

SAC was first robustly activated by the *nda3-KM311* mutant at 18 °C and then

inactivated simply by shift mitotically arrested cells back to permissive temperature

(30 °C) (Figures 1C). Because Cdc13 (cyclin B in *S. pombe*) localizes to the spindle

pole bodies (SPBs) in early mitosis and should be degraded by APC/C to promote

metaphase-anaphase transition, accumulation of Cdc13-GFP on SPBs after cold

treatment serves as the read-out of the SAC activation, and the disappearance rate of

Cdc13-GFP spot under permissive temperature reflects the SAC inactivation efficacy

(Figures 1C). It is noteworthy that the ATP analogue-sensitive mutant *sty1-T97A* (i.e.

sty1-as2) (Zuin et al., 2010), instead of *sty1Δ* deletion mutant, was used in this assay.

That was due to the G_2 delay that has been demonstrated in the *sty1Δ* deletion mutant

(Shiozaki and Russell, 1995a), which could interfere mitotic timing for further

analysis in this assay. We found that inactivation of CIP or SAP signaling by *pmk1Δ*

or *sty1-T97A* compromised full SAC activation, as these two mutants had only less

than 60% of cells with Cdc13-GFP on SPBs after 6 hours of cold treatment, whereas

the percentage in wild-type and *spk1Δ* cells after the same treatment was roughly 90%

(Figure 1D; Supplementary Figure 1C). Consistently, the disappearance of

Cdc13-GFP from SPBs in *pek1^{DD}* and *wis1^{DD}* mutant cells released from

metaphase-arrest was severely delayed (Figure 1D; Supplementary Figure 1B),

suggesting delayed SAC inactivation and anaphase onset after release from spindle

checkpoint arrest in these mutants.

All above data suggested that MAPK pathways are evolutionarily conserved in the

fission yeast to participate in spindle checkpoint activation and maintenance, of which

the CIP and SAP signaling pathways, but not the Spk1 pathway, are actively involved.

CIP and SAP signaling attenuate protein levels of Slp1^{Cdc20} and facilitate

association of MCC with APC/C respectively

The mitotic checkpoint complex (MCC, consisting of Mad3-Mad2-Slp1^{Cdc20} in *S.*

pombe) is the most potent inhibitor of the APC/C prior to anaphase (Kapanidou et al.,

2017; Primorac and Musacchio, 2013; Sudakin et al., 2001). Previous study has

revealed that *Xenopus* MAPK phosphorylates Cdc20 and facilitates its binding by

Mad2, BubR1 (Mad3 in *S. pombe*) and Bub3 to form a tight MCC to prevent Cdc20

from activating the APC/C (Chung and Chen, 2003). In fission yeast, the key SAC

components Mad2 and Mad3 and one molecule of Slp1^{Cdc20} form MCC which binds

to APC/C through another molecule of Slp1^{Cdc20} upon checkpoint arrest, and the

recovery from mitotic arrest accompanies the loss of MCC-APC/C binding (May et

al., 2017; Sczaniecka et al., 2008; Sewart and Hauf, 2017; Vanoosthuyse and

Hardwick, 2009). Since we found that constitutive activation of CIP or SAP signaling

in fission yeast by *pek1^{DD}* or *wis1^{DD}* mutations caused prolonged SAC activation, we

were suspicious that these two MAPK pathways might also execute their functions

cooperatively through a similar mechanism to that in *Xenopus*.

To test this possibility, we analyzed the MCC occupancy on APC/C by

immunoprecipitations of the APC/C subunit Lid1 (Yoon et al., 2002) in *pek1^{DD}* and

wis1^{DD} cells arrested by activated checkpoint. Compared to wild-type, more MCC

components (Mad2 plus Mad3 and Slp1^{Cdc20}) were co-immunoprecipitated in *wis1^{DD}*

cells, while less of them were co-immunoprecipitated in *pek1^{DD}* cells (Figure 2A),

suggesting that activation of the SAP but not the CIP pathway enhanced the

association of MCC with APC/C.

The decrease of co-immunoprecipitated Mad2 and Mad3 in *pek1^{DD}* cells was largely

out of expectation. Further analysis of the inputs of these immunoprecipitations

showed that Slp1^{Cdc20} levels were significantly reduced in *pek1^{DD}* but not *wis1^{DD}* cells

(Figure 2A). Our immunoblotting analyses using the total lysate of *pek1^{DD}* or *wis1^{DD}*

cells arrested at metaphase also confirmed these results (Figure 2B). Consistently, the

Slp1^{Cdc20} level was significantly increased in *pmk1Δ* and *pmk1Δ sty1-T79A* cells but

not in *sty1-T79A* cells (Figure 2C). The elevated Slp1^{Cdc20} level in *pmk1Δ* cells is

similar to that we observed in cells harboring two copies of *slp1^{+}*, which leads to

artificially increased Slp1^{Cdc20} level and also causes a SAC strength defect (Bai et al.,

2022). It is noteworthy that attenuated levels of Slp1^{Cdc20} in *pek1^{DD}* cells was not due

to decreased transcription of *slp1^{+}* gene (Figure 2D). These results collectively

revealed a negative effect of CIP but not SAP signaling on Slp1^{Cdc20} levels, which

potentially results in the weakened MCC-APC/C interaction detected in CIP mutants

(Figure 2A).

Together, our above data suggested a new mechanism involves division of labor

between the two fission yeast MAPK pathways CIP (Pek1-Pmk1) and SAP

(Wis1-Sty1) upon SAC activation, which is required in concert to lower Slp1^{Cdc20}

levels and enhance MCC affinity for APC/C to strongly inhibit APC/C activity

(Figure 2E). In this study, we only focused on characterizing the role of the CIP/Pmk1

in regulating SAC inactivation and anaphase onset hereafter.

207   Pmk1 directly binds Slp1^{Cdc20} to attenuate its levels upon SAC activation

To gain further insight into how Slp1^{Cdc20} levels are modulated by Pmk1, we

examined the potential interaction of Pmk1 with Slp1^{Cdc20}. Through incubating

MBP-tagged Slp1^{Cdc20} with metaphase-arrested yeast cell lysates, we found that Pmk1

expressed from yeast cells could be pull-downed by Slp1^{Cdc20} (Figure 3A).

In previous reports, MAPK-docking sites, which are important for interaction with

MAPKs, have been identified at the N termini of different MAPKKs from yeast to

humans with a notable feature of a cluster of at least two basic residues (mainly Lys/K

and Arg/R) separated by a spacer of 2-6 residues from a hydrophobic-X-hydrophobic

sequence (Bardwell et al., 2001; Bardwell and Thorner, 1996). Thus, we attempted to

identify the potential key residues in Slp1^{Cdc20} that mediate the Pmk1-Slp1^{Cdc20}

interaction. By visual scanning of Slp1^{Cdc20} sequence, we noticed five basic-residue

patches which are also present in several fungal species and loosely resemble the

consensus sequence of the known MAPK-docking sites, and four of them are within

N-terminal portion of the protein (Figure 3B and Supplementary Figure 2). However,

only mutating either of the two most N-terminal basic-residue patches (19-KKR-21

and 47-KR-48) to glutamates (Glu, E) or alanines (Ala, A) in Slp1^{Cdc20} resulted in

decreased amount of Slp1^{Cdc20} pulled down by Pmk1-GST *in vitro* using the

bacterially expressed recombinant proteins, and mutating both patches or mutating

only two arginine residues (R21 and R48) in those patches was sufficient to

completely disrupt their interaction (Figure 3C and Supplementary Figure 3).

Importantly, mutations of these Pmk1-docking sequences in Slp1 (PDSS for short) to

alanines mimicked *pmk1Δ* mutant in both elevated Slp1^{Cdc20} protein levels (Figure 3D)

and defective activation and maintenance of the SAC upon *nda3*-mediated checkpoint

arrest (Figure 3E and Supplementary Figure 4). Also, similar to removal of Pek1

downstream MAPK Pmk1, the lowered Slp1^{Cdc20} protein levels and delayed SAC

inactivation in *pek1^{DD}* mutant could be relieved by *slp1(DS-5A)* mutant, which is

deficient in Pmk1-docking (Figure 3D, 3E and Supplementary Figure 4).

Previously, one conserved non-canonical Pmk1-docking sequence which bears

positively charged residues and IYT motif has been identified in MAPK phosphatase

Pmp1 (Sacristan-Reviriego et al., 2014). Very strikingly, insertion of a 15 amino acid

sequence containing the Pmk1-docking sequence in Pmp1 (or PDSP for short) right in

front of the first basic-residue patch in Slp1(DS-5A) mutant protein restored its

interaction with Pmk1 (Figure 3B and 3C). And this artificially forced Pmk1-Slp1^{Cdc20}

interaction via PDSP was able to lower Slp1^{Cdc20} protein levels and restore higher

percentage of metaphase arrest upon SAC activation in *slp1(DS-5A)* mutant, which is

Pmk1-docking-deficient, although the rescuing effect was less efficient than the

presence of genuine PDSS (Figure 3D, 3E and Supplementary Figure 4). These data

firmly supported the idea that Pmk1 binds Slp1^{Cdc20} to downregulate its levels upon

SAC activation.

Collectively, these results demonstrated that Pmk1 directly binds to Slp1^{Cdc20} through

two short N-terminal basic patches serving as the major docking sites for Pmk1, and

the physical interaction of Pmk1-Slp1^{Cdc20} is involved in attenuation of Slp1^{Cdc20}

abundance upon SAC activation (Figure 3F).

Pmk1 synergizes with Cdk1 to phosphorylate Slp1^{Cdc20}

The association of Pmk1 with Slp1^{Cdc20} and the negative effect of constitutively

activated CIP signaling on SAC inactivation relies on Pmk1 kinase suggested that

Slp1^{Cdc20} may be a direct substrate of Pmk1. To test this possibility, we performed *in*

vitro phosphorylation reactions and found that bacterially expressed recombinant

GST-Slp1^{Cdc20} could be efficiently phosphorylated in the presence of Pmk1-HA-His

purified from yeast cells (Figure 4A).

To identify potential phosphorylation sites *in vivo*, we purified Apc15-GFP from wild

type, *pmk1Δ* and *pek1^{DD}* cells arrested by activated checkpoint with intention that the

Slp1^{Cdc20}-containing APC/C complexes could be captured (Supplementary Figure 5A

and 5B). Purifications were performed in the presence of phosphatase inhibitors to

enrich for phosphopeptides. Mass spectrometry analysis revealed that the

phosphorylation of Thr480 residue at the most C-terminus of Slp1^{Cdc20} was dependent

on the presence of Pmk1, and this modification was present in *pek1^{DD}* but not in wild

type cells (Figure 4B, Supplementary Figure 5C-E and Supplementary Figure 6),

indicating that phosphorylation of Thr480 is very likely triggered by CIP signaling.

We raised phospho-specific antibodies against pT480, it could recognize Slp1^{Cdc20}

fragment phosphorylated by Pmk1 *in vitro* and mutation of Thr480 to alanine (T480A)

abolished site-specific phosphorylation (Figure 4C). We also confirmed that

endogenous Slp1^{Cdc20} was indeed phosphorylated at Thr480 in metaphase-arrested

mts3-1 cells, which was enhanced in *pek1^{DD}* mutant and diminished in *pmk1Δ* mutant

(Figure 4D). Thus, our results established Thr480 as a major Pmk1 phosphorylation

site *in vivo* upon SAC activation.

In addition to pT480, our mass spectrometry analysis also identified Ser28, Thr31 and

Ser59 as Pmk1-independent and Ser76 as Pmk1-dependent phosphorylation sites

(Figure 4B, Supplementary Figure 5C-E and Supplementary Figure 6). All these four

residues are located within the N-terminal unstructured region, fit with the Cdk1 core

motif (SP or TP), and more importantly, they are conserved not only in four

Schizosaccharomyces species but also in several other fungal species (Figure 4B and

Supplementary Figure 2B). Interestingly, they were also proposed as Cdk1

phosphorylation sites based on quantitative phosphoproteomic analysis in a recent

study (Swaffer et al., 2018) (Figure 4B). By using the phospho-specific antibodies

against both pS28 and pT31, we could detect the phosphorylation of Ser28 and Thr31

in vitro by Cdc13 (cyclin B)-containing Cdk1 complexes purified from

metaphase-arrested yeast cells, which was diminished when the kinase activity of ATP

analogue-sensitive Cdk1 [cdc2-as1(F84G)] was inhibited by 1-NM-PP1 or when

Slp1$^{(S28A;T31A)}$ was added as the substrate in *in vitro* kinase reactions (Figure 4E). Next,

we examined whether Ser28 and Thr31 were phosphorylated *in vivo* using

immunoblotting with phospho-specific antibodies to directly detect Slp1^{Cdc20}, which

was co-immunoprecipitated with Apc15 from *nda3*-arrested yeast cells. We observed

phosphorylation of Ser28/Thr31 in wild type cells, but it disappeared following

phosphatase treatment of wild type samples after immunoprecipitation, 1-NM-PP1

treatment of *cdc2-as1* cells before being collected for immunoprecipitation, or when

cells expressing Slp1$^{(S28A;T31A)}$ were used (Figure 4F). Thus, Slp1^{Cdc20} is also phosphorylated by Cdk1 at least at Ser28 and Thr31 in fission yeast mitotic cells.

We wondered whether Slp1^{Cdc20} phosphorylation by Pmk1 affects its stability and APC/C activation. Indeed, phospho-deficient Slp1^{T480A} mutant showed elevated Slp1^{Cdc20} levels in both metaphase-arrested and asynchronously grown cells (Figure 4G and Supplementary Figure 7) and compromised SAC activation compared to wild type cells, likely due to more efficient APC/C activation (Figure 4H and Supplementary Figure 8). Furthermore, Slp1^{T480A} mutant also reversed the inhibitory effect of *pek1^{DD}* on Slp1^{Cdc20} stability and APC/C activation after release from SAC-mediated arrest (Figure 4G and 4H).

Ser76 was identified as a Pmk1-dependent phosphorylation residue in our mass spectrometry analysis, but it was considered as a Cdk1 targeting site in an earlier study (Swaffer et al., 2018). It is fairly possible that phosphorylation of Ser76 is promoted by both Cdk1 and Pmk1, which is conceivable as CDKs and MAPKs recognize the same core motif (SP/TP) (Pinna and Ruzzene, 1996), and crosstalk between CDK and MAPK has recently been reported for pheromone signaling in budding yeast (Repetto et al., 2018). We tested how the double phosphorylation sites (S76A;T480A) mutant affects Slp1^{Cdc20} levels or APC/C activity, and observed that the protein levels of Slp1$^{S76A;T480A}$ were further elevated and SAC activation efficiency in this mutant was further lowered compared to Slp1^{T480A} mutant (Figure 4G and 4H). Of note, Slp1^{T480A} and Slp1$^{S76A;T480A}$ mutant proteins became easily detected in interphase cells (Supplementary Figure 7), this is very different from wild type Slp1 which normally accumulates exclusively in mitosis (Yamada et al., 2000).

We further tested whether replacement of other identified phospho-serine/threonine residues with alanines affects Slp1^{Cdc20} protein levels and APC/C activity. We constructed strains carrying combinations of mutations of S28A, T31A and S59A with S76A, and also with confirmed Pmk1 phosphorylation-deficient T480A, and found

that mutations of S28A, T31A and S59A did not affect Slp1^{Cdc20} protein levels, though

they could also profoundly compromise SAC activation efficiency similarly to

S76A;T480A mutant (Figure 4G, 4H and Supplementary Figure 7). We noticed that

the presence of mutations of S28A, T31A and S59A also abolished the elevated

Slp1^{Cdc20} protein levels conferred by S76A;T480A mutations (Figure 4G and

Supplementary Figure 8). Thus, phosphorylation of S28, T31 and S59 targeted by

Cdk1 most likely influences APC/C activation through distinct mechanism from

Pmk1-mediated S76/T480 phosphorylation.

Taken together, our data suggested that activated Pmk1 phosphorylates Slp1^{Cdc20} *in*

vivo, which leads to lowered Slp1^{Cdc20} levels, delayed SAC inactivation and APC/C

activation as we observed in *pek1^{DD}* cells. Also, Pmk1 can join Cdk1 to enhance the

multi-site phosphorylation of their shared substrate Slp1^{Cdc20} and collaboratively

dampen APC/C activity.

Pmk1- but not Cdk1-mediated Slp1^{Cdc20} phosphorylation promotes its

ubiquitylation

Our above data support the notion that the direct binding of Slp1^{Cdc20} by Pmk1

mediates its phosphorylation, which possibly results its subsequent rapid turnover. It

has been shown in both human and fission yeast cells that Apc15, one of the

conserved components of APC/C, promotes Cdc20/Slp1^{Cdc20} autoubiquitylation and

its turnover by APC/C (Mansfeld et al., 2011; May et al., 2017; Sewart and Hauf,

2017; Uzunova et al., 2012). The increased Slp1^{Cdc20} levels in *pmk1Δ* cells (Figure 2C)

is reminiscent of the previous observations in *apc15Δ* mutant, which also shows

elevated levels of Slp1^{Cdc20} (May et al., 2017; Sewart and Hauf, 2017). We reasoned

that the absence of Apc15 may reverse the negative effect of *pek1^{DD}* on Slp1^{Cdc20}

levels and spindle checkpoint inactivation and APC/C activation. As expected, we

indeed observed that deletion of *apc15* recovered Slp1^{Cdc20} levels in *pek1^{DD}* cells

(Figure 5A), and also lowered the percentage of cells with Cdc13-GFP retained on

SPBs in *nda3-KM311 pek1^{DD}* cells after cold treatment (Figure 5B and Supplementary

Figure 9A). These observations are consistent with previous reports showing the relative abundance between checkpoint proteins and the checkpoint target Slp1^{Cdc20} is an important determinant of checkpoint robustness (Heinrich et al., 2013) and *apc15Δ* mutant is spindle checkpoint-defective (May et al., 2017).

During the course of our mass spectrometry analysis on Slp1^{Cdc20} phosphorylation, we also detected Lys 479 (K479) as an ubiquitin-modifying site (Figure 4B and Supplementary Figure 5 and 6). We wondered whether K479 is involved in Slp1^{Cdc20} autoubiquitylation in fission yeast. In agreement with the mass spectrometry analysis, mutation of K479 to arginine indeed conferred stability of Slp1^{Cdc20} (Figure 5C), even in asynchronously growing cells (Supplementary Figure 10), which is reminiscent of the stabilized Slp1^{Cdc20} in *apc15Δ* mutant cells at interphase (Sewart and Hauf, 2017). These results suggested that K479 is likely Slp1^{Cdc20} ubiquitylation-relevant. Furthermore, *nda3-KM311 slp1^{K479R}* cells also demonstrated lowered metaphase arrest rate after cold treatment (Figure 5D and Supplementary Figure 9B), indicating defective SAC activation. Previous studies have identified two neighbouring lysine residues close to the carboxy terminus of human Cdc20, Lys485 and Lys490, as ubiquitylation sites in prometaphase cells, and mutating these two lysines to arginine completely prevented Cdc20 ubiquitylation (Danielsen et al., 2011; Mansfeld et al., 2011). Very interestingly, a second lysine residue, K472, is also present in fission yeast Slp1^{Cdc20} adjacent to K479 (Figure 4B and Supplementary Figure 11). Although K472 was not identified as an ubiquitin-modifying residue in our mass spectrometry analysis, replacement of K472 with arginine also efficiently stabilized Slp1^{Cdc20}, similar to K479R mutant (Figure 5C and Supplementary Figure 10). More strikingly, simultaneously mutating both K472 and K479 caused additive effects in elevating Slp1^{Cdc20} levels and lowering SAC activation rate in both wild type and *pek1^{DD}* strain backgrounds (Figure 5C, 5D and Supplementary Figure 10), suggesting that likely both these two lysine residues at the C-terminal tail of Slp1^{Cdc20} contribute to its turnover via ubiquitylation, which is very similar to the mechanism in human Cdc20. Consistently, using TUBEs (tandem ubiquitin-binding entities) (Hjerpe et al., 2009) to

enrich for ubiquitinated proteins in metaphase-arrested *mts3-1* cells expressing

sfGFP-Slp1, we were able to demonstrate that mutating both K472 and K479 largely

removed higher molecular weight bands detected by immunoblotting, which were

corresponding to ubiquitylated species of Slp1^{Cdc20} (Figure 5E, F). We also noticed

that Slp1$^{K472R; K479R}$ rendered more severely compromised viability than either single

mutants (Supplementary Figure 12), indicating blocking Slp1^{Cdc20} ubiquitylation

could bring about detrimental consequence to the cells.

Our observation that Slp1^{Cdc20} levels were lowered in *pek1^{DD}* mutant (Figure 2B)

suggested that attenuation of Slp1^{Cdc20} levels upon constitutive activation of CIP

signaling is very likely through facilitating its autoubiquitylation. To test this

possibility, we examined and compared the ubiquitylation levels of sfGFP-Slp1 in

wild type, *pmk1Δ* and *pek1^{DD}* cells by affinity pull-down assays using TUBEs (Figure

5E). Indeed, we observed a much weaker or stronger discrete laddering pattern of

sfGFP-Slp1 in *pmk1Δ* and *pek1^{DD}* cells than that of wild type cells, indicating abated

or enhanced ubiquitylation respectively (Figure 5F). Furthermore, mutations of

T480A, S76A/T480A and Pmk1-docing site, but not S28A/T31A- or

S28A/T31A/S59A-containing mutants, also reduced polyubiquitylation of Slp1^{Cdc20}

(Figure 5F). All these results confirmed that Pmk1- but not Cdk1-mediated Slp1^{Cdc20}

phosphorylation promotes its ubiquitylation.

Together, these findings lent further support to the idea that the CIP signaling pathway

restrain APC/C activity through a cascade mechanism, in which Pmk1 directly binds

and phosphorylates Slp1^{Cdc20} to facilitate its autoubiquitylation, and eventually lowers

its levels (Figure 5G).

Osmotic stress and cell wall damage trigger rapid Slp1^{Cdc20} downregulation and

mitotic exit delay

It has been well established that two MAPKs Sty1 and Pmk1 can be activated by

environmental perturbations including osmotic stress, cell wall damage, thermal stress,

oxidative stress, and glucose limitation, among others (Cansado et al., 2021; Madrid

et al., 2006; Shiozaki and Russell, 1995b). We wondered whether the phosphorylation

of Slp1^{Cdc20} by Pmk1 could tune the strength of SAC activation and APC/C activity in

response to extracellular stress. To investigate this possibility, we performed

arrest-and-release experiments with *nda3-KM311* mutants and added 0.6 M KCl or 2

μg/mL of caspofungin 1 hr before release from metaphase arrest to elicit strong

osmotic saline stress (KCl) or cell wall damage stress (caspofungin) response (Madrid

et al., 2006; Madrid et al., 2016), respectively (Figure 6A). As previously reported,

treatment with KCl or caspofungin resulted in full activation of both Pmk1 and Sty1

(for KCl treatment) or only Pmk1 (for caspofungin treatment) as determined using

anti-phospho p42/44 and anti-phospho p38 antibodies, which serve as indicative of

respective Pmk1 and Sty1 activation (Figure 6B). Intriguingly, treatment with KCl or

caspofungin also correlated with a marked and rapid decrease in Slp1^{Cdc20} levels

(Figure 6B).

Surprisingly, we did not see any alteration of APC/C-MCC interaction accompanying

Pmk1 activation when cells were treated by caspofungin (Figure 6C), which was

different from what we observed in P_{adh11}-*pek1^{DD}* cells (Figure 2A). Also, we failed to

detect enhanced APC/C-MCC association like we observed in P_{adh11}-*wis1^{DD}* cells

when cells were treated by KCl, where Sty1 was activated (Figure 6C). The latter

observation was most likely due to the disruptive effect of high concentration of KCl

on protein-protein interactions, because we observed similar effect when we

immunoprecipitated APC/C in the presence of 0.6 M KCl in the lysis buffer (Figure

6C, right panel). With this data, we could not exclude the possibility that activated

Sty1 transiently enhanced APC/C-MCC association in yeast cells, which was soon

counteracted by the disruptive effect of saline. Nevertheless, the high SAC activation

efficiency in mitotis-arrested cells exposed to osmotic stress (KCl) or cell wall

damage stress (caspofungin) was largely lowered by *pmk1* deletion (Figure 6D),

indicating that at least activated Pmk1 exerts regulation of the spindle checkpoint and

APC/C activation under environmental stimuli. Further, Pmk1-docking site mutant

slp1(DS-5A) and Pmk1 phosphorylation-deficient mutants *slp1^{T480A}* and *slp1$^{S76A;T480A}$*

could partially suppress the sustained SAC activation responding to osmotic and cell

wall damage stress (Figure 6E and Supplementary Figure 13).

DISCUSSION

Involvement of MAPKs in cell cycle regulation

So far, the only known cell-cycle control stages linked to MAPKs in fission yeast are

at G$_2$/M transition and during cytokinesis (Gomez-Gil et al., 2020; Petersen and

Hagan, 2005; Petersen and Nurse, 2007). The p38 MAPK family member Sty1 either

arrests or promotes mitotic commitment in unperturbed, stressed or resumed cell

cycles after stress depending on the downstream kinases it associates with. Sty1 can

phosphorylate kinase Srk1 at threonine 463 (T463), which further phosphorylates and

inhibits Cdc25 phosphatase to negatively regulate the onset of mitosis (Lopez-Aviles

et al., 2005; Lopez-Aviles et al., 2008; Smith et al., 2002). Sty1 also mediates

phosphorylation of Polo kinase Plo1 at serine 402 (S402), which then promotes its

recruitment to the spindle pole bodies (SPBs), where it positively modulates Cdc25

control of Cdc2 (fission yeast Cdk1) activity for mitotic entry (Petersen and Hagan,

2005; Petersen and Nurse, 2007). During cytokinesis, two fission yeast formins For3

and Cdc12 cooperate to nucleate actin filaments for assembly of contractile

actomyosin ring (CAR) (Coffman et al., 2013). It has been shown that Sty1 activity

negatively regulates protein levels of For3 in response to actin cytoskeleton damage to

downregulate CAR assembly (Gomez-Gil et al., 2020). Very recently, Pmk1 has also

been proposed to act as a negative regulator of cytokinesis, which delays CAR

constriction upon cell wall stress (e.g. treatment with blankophor, caspofungin and

caffeine) through a novel checkpoint mechanism (Edreira et al., 2020). However, the

detailed mechanisms how Sty1 and Pmk1 pose negative effect on cytokinesis have not

been elucidated.

In the current study, we have expanded MAPK signaling-controlled cell-cycle stage to

metaphase-anaphase transition in fission yeast. In response to unattached or

tensionless kinetochores, the SAC generates the MCC, which inhibits the APC/C and

block anaphase onset by suppressing APC/C-catalyzed ubiquitylation and subsequent

degradation of securin and cyclin B. As a critical APC/C inhibitor, MCC fulfills its

function through the physical binding of BubR1/Mad3 and Mad2 to two molecules of

Cdc20/Slp1^{Cdc20} (*i.e.* Cdc20^{MCC} and Cdc20$^{APC/C}$), and this mechanism has been

confirmed at least in both humans and fission yeast (Alfieri et al., 2016; Chao et al.,

2012; Izawa and Pines, 2015; Sewart and Hauf, 2017; Yamaguchi et al., 2016). In this

study, we have uncovered a dual mechanism, in which two fission yeast MAPKs

Pmk1 and Sty1 restrain APC/C activity through distinct manners, either by

phosphorylating Slp1^{Cdc20} to lower its levels or by phosphorylating unknown

substrate(s) to promote MCC-APC/C association and thus SAC signaling strength,

respectively (Figure 7). Therefore, our study has established another critical

APC/C-inhibitory mechanism in the spindle checkpoint, though it will be important

for future studies to identify the phosphorylation target(s) posed by SAP signaling

pathway. Nevertheless, our current study together with earlier report (Edreira et al.,

2020) have implicated fission yeast Pmk1 in both the mitotic and cytokinesis

surveillance mechanisms in response to unfavorable environmental stimuli.

The regulation of mitotic entry by MAPK in fission yeast is triggered by many

environmental insults, including high osmolarity, oxidative stress, heat shock,

centrifugation, nutrient starvation and rapamycin treatment (Hartmuth and Petersen,

2009; Petersen and Hagan, 2005; Petersen and Nurse, 2007; Shiozaki and Russell,

1995a; Shiozaki et al., 1998). In this study, we found that treatment of fission yeast

cells with KCl or caspofungin, which provokes a strong osmotic stress or cell wall

damage response and quick Pmk1 activation, correlated with an attenuated Slp1^{Cdc20}

protein levels and delayed APC/C activation. Deletion of *pmk1$^+$* largely overrides the

negative effects of KCl or caspofungin treatment on APC/C activation (Figure 6D),

confirming that activated Pmk1 is responsible for prolonged metaphase arrest under

environmental stress. Thus, our finding implicates MAPK Pmk1 as an important

regulator for anaphase entry in response to adverse extracellular conditions. Intriguingly, one previous study showed that treatment of human cells with the carcinogen cadmium activates MAPK p38 (fission yeast Sty1 homologue) and triggers accelerated ubiquitination and proteolysis of Cdc20, which is essential for prometaphase arrest under SAC (Yen and Yang, 2010). Therefore, although different MAPKs might be employed to fulfil the task in yeast and higher eukaryotes, it is still possible that phosphorylation and regulation of Cdc20 levels by MAPKs is an evolutionarily conserved mechanism to adjust mitotic exit to extracellular or environmental stimuli.

Negative phospho-regulation of Cdc20 by MAPK as a spindle assembly checkpoint-dependent APC/C-inhibitory mechanism

APC/C activity can be regulated at multiple levels, Cdk1-dependent APC/C phosphorylation is one of the major prerequisites for APC/C activation and anaphase onset, in which Apc3 and Apc1 are key subunits for positive phospho-regulation (Fujimitsu et al., 2016; Kraft et al., 2003; Qiao et al., 2016; Steen et al., 2008; Zhang et al., 2016). However, it has been shown that Cdk1-dependent phosphorylation of APC/C co-activator Cdc20 lessens its interaction with APC/C, thus serves as a negative phospho-regulation (Labit et al., 2012). In addition to its phosphorylation by Cdk1, phosphorylation of Cdc20 by multiple other kinases, such as Bub1, PKA, MAPK and Plk1, has also been demonstrated or suggested in *Xenopus*, humans or budding yeast to be inhibitory to APC/C activation and improper anaphase onset upon SAC activation ((Jia et al., 2016) and reviewed in (Yu, 2007)). Therefore, Cdc20 generally serves as an integrator of multiple intracellular signaling cascades that regulate progression through mitosis. In this work, we demonstrated that phosphorylation of Slp1^{Cdc20} by Pmk1 but not Cdk1 promotes its autoubiquitylation, this mechanism is distinct from that reported in *Xenopus* egg extracts and tadpole cells, in which phosphorylation of Cdc20 by MAPK is instead responsible for facilitated Cdc20 binding by MCC to prevent it from activating the APC/C (Chung and Chen, 2003).

Although multiple kinases have been proposed to be directly involved in phosphorylation of Cdc20 proteins (Yu, 2007), precisely identifying the *in vivo* phosphorylation sites in Cdc20 has turned out to be challenging, most likely due to its protein scarcity and instability. Based on known Cdk1 core consensus (SP/TP), 9 residues in human Cdc20 were originally suggested as putative Cdk1 sites, but among them only Ser41 has been proven to be phosphorylated *in vivo* by mass spectrometry analysis (Kramer et al., 2000; Yudkovsky et al., 2000). However, subsequent mass spectrometry analyses identified Ser41 as one of the Bub1 phosphorylation sites both *in vivo* and *in vitro* and another putative Cdk1 site Thr106 as one of the Plk1 sites (Jia et al., 2016; Tang et al., 2004). It is very likely that there are not as many genuine Cdk1 sites in Cdc20 as originally assumed. One supporting evidence is that mutation of only one (Thr32) of the three putative Cdk sites to alanine in N terminus of *C. elegans* CDC-20 is sufficient to phenocopy 3A mutant for accelerated APC/C activation (Kim et al., 2017). Interestingly, Thr32 of *C. elegans* CDC-20 is conserved as Thr 59 and Thr68 in human and *Xenopus* Cdc20, respectively (see Supplementary Figure 14). Based on two-dimensional tryptic phosphopeptide analysis, Thr68 in *Xenopus* Cdc20 has also been regarded as a MAPK target site (Chung and Chen, 2003). These studies in diverse organisms raised an issue that whether multiple kinases share the same phosphorylation sites in Cdc20? It would not be so surprising if it turns out to be the case eventually, at least for Cdk1 and MAPKs, because CDKs and MAPKs recognize the same core motif (SP/TP) (Pinna and Ruzzene, 1996).

By purifying Apc15-aasociated fission yeast Slp1^{Cdc20} proteins from checkpoint-arrested *pek1^{DD}* and *pmk1Δ* cells, we were able to identify Thr480 and Ser76 as *in vivo* Pmk1-relevant phosphorylation sites and 3 Pmk1-independent phosphorylation sites (S28, T31, S59) based on mass spectrometry analyses. All our identified sites except Thr480 are located in the N terminal disordered portion of Slp1^{Cdc20} sequence, this is similar to the phosphorylation residues identified by biochemical and functional analyses in other organisms, in which the major

phosphorylation sites all fall within the N-terminus of Cdc20 homologues. We further provided evidence that S28/T31 are phosphorylated by Cdk1 both *in vitro* and *in vivo*, and T480 phosphorylated by Pmk1 both *in vitro* and *in vivo*. We noticed that there is a tendency for reported phosphorylation residues to be concentrated in the N terminus of Cdc20 proteins (Jia et al., 2016; Tang et al., 2004; Yu, 2007). However, due to the lack of clear sequence homology in their N-terminal and C-terminal tails between Slp1[Cdc20] and its Cdc20 homologues in higher eukaryotes (see Supplementary Figure 14), it is impossible to certainly assign any identified sites from different organisms as homologous sites. But it is possible that multiple kinases modify shared sites in Cdc20, this collaborative effect may be especially helpful under adverse environmental stress. Nevertheless, all reported Cdc20 phosphorylation events constitute an APC/C-inhibitory mechanism, no matter what kinases are responsible for the modifications (Chung and Chen, 2003; D'Angiolella et al., 2003; Jia et al., 2016; Kim et al., 2017; Kramer et al., 2000; Labit et al., 2012; Tang et al., 2004; Yu, 2007; Yudkovsky et al., 2000).

The current model of inhibitory signaling from Cdc20 posits that its phosphorylation either promotes the formation of MCC or inhibits APC/C[Cdc20] catalytically, both are required for proper spindle checkpoint signaling (Jia et al., 2016; Labit et al., 2012; Yamano, 2019). Interestingly, our current study suggests Slp1[Cdc20] phosphorylation by fission yeast MAPK Pmk1 may represent another critical APC/C-inhibitory mechanism in the spindle checkpoint through reducing the protein levels of Slp1[Cdc20], which is achieved via autoubiquitylation-mediated degradation (Figure 7). Actually, previous studies in fission and budding yeast have underlined the importance of accurate relative abundance between checkpoint proteins and Cdc20/Slp1[Cdc20], which sets an important determinant of checkpoint robustness (Heinrich et al., 2013; Pan and Chen, 2004). A benefit of allowing the association of MCC-bound Cdc20 with APC/C is to provide an opportunity for Cdc20 autoubiquitylation by APC/C, a process aided by APC/C subunit Apc15 (Mansfeld et al., 2011; May et al., 2017; Sewart and Hauf, 2017; Uzunova et al., 2012). Notably, the depletion of *apc15+* restores Slp1[Cdc20] levels

in *pek1^{DD}* cells (Figure 5A), consistent with the idea that Pmk1-mediated Slp1^{Cdc20} phosphorylation very likely facilitates its autoubiquitylation. Indeed, our biochemical and functional analyses further establish K479 and K472 in Slp1^{Cdc20} as two key residues directly involved in its ubiquitylation, which likely follows its phosphorylation, particularly at adjacent Thr480. Although K485 and K490 are similarly located at C terminus of human Cdc20 and can be ubiquitylated and responsible for Cdc20 degradation (Danielsen et al., 2011; Mansfeld et al., 2011), there has been no reports on phosphorylation-modified residues in this region. However, although the sequence homology within N- and C-terminal regions between Slp1^{Cdc20} and Cdc20 homologues in higher eukaryotes is extremely low, we do notice that there are several lysine-flanking serine or threonine residues within Cdc20 C-terminal tails of higher eukaryotes (see Supplementary Figure 11), this raises a possibility that this region may be involved in phosphorylation-coupled ubiquitylation. Currently, we do not know how T480 phosphorylation accelerate ubiquitylation of K479 and K472 in Slp1^{Cdc20}. It is possible that phosphorylation of Slp1^{Cdc20} might alter the mode of Cdc20 binding to APC/C which becomes conducive to catalyze Slp1^{Cdc20} ubiquitylation.

In summary, our studies revealed a distinct dual mechanism of MAPK-dependent anaphase onset delay imposed directly on APC/C co-activator Slp1^{Cdc20} and an unidentified substrate, which may emerge as an important regulatory layer to fine tune APC/C and MCC activity upon SAC activation. Importantly, although this mechanism of APC/C regulation by two MAPKs has not been examined in other organisms, it is very likely widespread across eukaryotes including humans, given the involvement of MAPKs in general response to wide range of adverse environmental situations and similar regulations of APC/C and SAC in eukaryotes characterized so far. It is also plausible to assume that the MAPK-mediated APC/C inhibition pathway may also contribute to the pathogenesis of multiple cancers. Indeed, Cdc20 upregulation has been demonstrated in several types of cancers, including breast and colon cancer, and it contributes to aggressive tumor progression and poor prognosis in gastric cancer

and primary non-small cell lung cancer (Chi et al., 2019). It would be interesting to
address whether the altered levels of Cdc20 in these cancers are MAPK relevant.
Nevertheless, inhibition of APC/C^{Cdc20} activity or induction of Cdc20 degradation
have already been listed as alternative therapeutic strategies to control cancer (Greil et
al., 2022; Jeong et al., 2022). Thus, although the detailed molecular mechanisms may
vary, further studies on MAPK-mediated APC/C^{Cdc20} inhibition mechanism in human
would lead to better understanding of its oncogenic role in tumor progression.

Materials and Methods

Fission yeast strains, media and genetic methods

Fission yeast cells were grown in either YE (yeast extract) rich medium or EMM
(Edinburgh minimal medium) containing the necessary supplements. G418 disulfate
(Sigma-Aldrich; A1720), hygromycin B (Sangon Biotech; A600230) or
nourseothiricin (clonNAT; Werner BioAgents; CAS#96736-11-7) was used at a final
concentration of 100 μg/ml and thiabendazole (TBZ) (Sigma-Aldrich; T8904) at 5-15
μg/ml in YE media where appropriate. For serial dilution spot assays, 10-fold
dilutions of a mid-log-phase culture were plated on the indicated media and grown for
3 to 5 days at indicated temperatures.

To create strain with an ectopic copy of *slp1$^+$* at *lys1$^+$* locus on chromosome 1, the
open reading frame of *slp1$^+$* (1467 bp) and its upstream 1504 bp sequence was first
cloned into the vector pUC119-*P$_{adh21}$*-MCS-*hphMX6-lys1*** with *adh21* promoter
(*P$_{adh21}$*) removed using the 'T-type' enzyme-free cloning method (Chen et al., 2017).
The sequence of *hphMX6* in above plasmid was replaced by fragment corresponding
to nourseothiricin resistance gene sequence by *Bgl*II-*EcoR*I cut and re-ligation,
generating pUC119-*P$_{slp1}$-slp1$^+$-T$_{adh1}$::natMX6-lys1***. The resultant plasmids were
linearized by *Apa*I and integrated into the *lys1$^+$* locus, generating the strains
lys1Δ::P$_{slp1}$-slp1$^+$-T$_{adh1}$::hphMX6 and *lys1Δ::P$_{slp1}$-slp1$^+$-T$_{adh1}$::nat^R*.

To generate the mutant strains harboring lysine/arginine (K/R) or serine/threonine to

glutamic acid (E) or alanine (A) mutations, or lysine (K) to arginine (R) mutations

within Slp1, the desired mutations were introduced into

pUC119-P_{slp1}-$slp1^+$-T_{adh1}::$hphMX6$ using standard methods of site-directed

mutagenesis.

To generate the strains carrying superfolder GFP (sfGFP)-tagged wild type or mutant

Slp1, a fragment corresponding to the sequence of *sfGFP* was inserted in front of *slp1*

in pUC119-P_{slp1}-$slp1$-T_{adh1}::$hphMX6$-$lys1^*$ using the 'T-type' enzyme-free cloning

method. The resulting plasmids were linearized and integrated into the $lys1^+$ locus of

chromosome 1 using the hyg^r marker. Then endogenous $slp1^+$ was deleted and

replaced with $ura4^+$.

To generate the strains expressing *ade6*::P_{adh1}-*GST-slp1(456-488aa)*::*natR*,

C-terminal sequence of $slp1^+$ corresponding to 456-488aa was first cloned into the

vector pUC119-P_{adh1}-MCS-$natMX6$-$lys1^*$ and the sequence of $lys1^*$ was replaced

with $ade6^+$ from pAde6-NotI-hphMX (a kind gift of Li-lin Du), and then the sequence

of GST from pGEX-4T-1 was inserted in front of *slp1*. The resulting plasmid

pUC119-P_{adh1}-*GST-slp1(456-488aa)*-$natMX6$-$ade6$ was linearized and integrated into

the $ade6^+$ locus of chromosome III using the nat^r marker.

To generate the constitutively active MAPKK strains $pek1^{DD}$ or $wis1^{DD}$, the Ser234

and Thr238 of $pek1^+$ in pUC119-P_{adh21}-*6HA-pek1$^+$-T$_{adh1}$-hphMX6-lys1**,

pUC119-P_{adh11}-*6HA-pek1$^+$-T$_{adh1}$-hphMX6-lys1**   and

pUC119-P_{adh11}-*6HA-pek1$^+$-T$_{adh1}$-kanMX6-ura4$^+$* and the Ser469 and Thr473 of $wis1^+$

in   pUC119-P_{adh21}-*6HA-wis1$^+$-T$_{adh1}$-hphMX6-lys1**,

pUC119-P_{adh11}-*6HA-wis1$^+$-T$_{adh1}$-hphMX6-lys1**   and

pUC119-P_{adh11}-*6HA-wis1$^+$-T$_{adh1}$-kanMX6-ura4$^+$* were changed to aspartic acids (D)

by standard methods of site-directed mutagenesis. The resulting plasmids were

linearized and integrated into the $lys1^+$ locus of chromosome I or the $ura4^+$ locus of

chromosome III using the *kan^r* marker. A list of the yeast strains used is in

Supplementary Table S1.

Fission yeast cell synchronization methods

For *nda3-KM311* strains, cells were grown at the permissive temperature for

nda3-KM311 (30 °C) to mid-log phase, synchronized at S phase by adding HU

(Sangon Biotech; A600528) to a final concentration of 12 mM for 2 hours followed

by a second dose of HU (6 mM final concentration) for 3.5 hours. HU was then

washed out and cells were released at specific temperatures as required by subsequent

experiments.

Spindle checkpoint activation and silencing assays in fission yeast

For checkpoint silencing assay in the absence of microtubules, mid-log *cdc13-GFP*

nda3-KM311 cells were first synchronized with HU at 30 °C and then arrested in early

mitosis by shifting to 18 °C for 6 hours, followed by incubation at 30 °C to allow

spindle reformation and therefore spindle checkpoint inactivation. For osmotic stress

or cell wall damage stress experiments, 0.6 M KCl or 2 μg/mL caspofungin was added

into cultures during 18 °C incubation 1 hr before being shifted to 30 °C. Cells were

withdrawn at certain time intervals and fixed with cold methanol and stained with

DAPI. 200-300 cells were analyzed for each time point. Each experiment was

repeated at least three times.

Immunoblotting and immunoprecipitation

For routine Western blot and immunoprecipitation experiments, yeast cells were

collected from *nda3-KM311* arrested and released cultures, followed by lysing with

glass bead disruption using Bioprep-24 homogenizer (ALLSHENG Instruments,

Hangzhou, China) in NP40 lysis buffer (6 mM Na_2HPO_4, 4 mM NaH_2PO4, 1% NP-40,

150 mM NaCl, 2 mM EDTA, 50 mM NaF, 0.1 mM Na_3VO_4) plus protease inhibitors

as previously described (Wang et al., 2012). Proteins were immunoprecipitated by

IgG Sepharose beads (GE Healthcare; 17-0969-01) (for Lid1-TAP) or anti-GFP

 nanobody agarose beads (AlpalifeBio, Shenzhen, China; KTSM1301) (for

721 Mad3-GFP). Immunoblot analysis of cell lysates and immunoprecipitates was

722 performed using appropriate antibodies at 1:500 - 1:5000 dilutions and was read out

723 using chemiluminescence.

725 Detection of *in vivo* phosphorylation of Slp1^{Cdc20} at Thr480 and Ser28/Thr31

For detection of phosphorylation of Slp1^{Cdc20} at Thr480 *in vivo*, *mts3-1* mutant strains

expressing *ade6::P$_{adh1}$-GST-slp1(456-488aa)::natR* in wild type, *pmk1Δ* and

lys1Δ::P$_{adh11}$-6xHA-pek1^{DD} background were arrested at 36°C for 3.5 hours, followed

by pull-down of GST-Slp1(456-488aa) by glutathione Sepharose 4B (GE Healthcare).

Purified GST-Slp1(456-488aa) was detected with either goat anti-GST

HRP-conjugated antibodies (GE Healthcare) or rabbit polyclonal anti-pThr480

antibodies.

For detection of phosphorylation of Slp1^{Cdc20} at Ser28/Thr31 *in vivo*, *nda3-KM311*

apc15-13myc::kanR strains expressing *slp1$^+$* or *slp1(S28A;T31A)* in wild type or

cdc2-asM17 background were first synchronized with HU at 30 °C and then arrested

in early mitosis by shifting to 18 °C for 6 hours. For *cdc2-asM17* strains, 1 μM

1-NM-PP1 (Toronto Research Chemicals; A603003) was added into one of the

cultures to inactivate Cdc2 (Cdk1) 20 min before being collected. Apc15-13myc was

immunoprecipitated by anti-c-Myc magnetic beads (MedChemExpress (MCE);

HY-K0206). To remove Slp1^{Cdc20} phosphorylation, one of the immunoprecipitated

samples was treated with 400 units of Lambda protein phosphatase (New England

Biolabs, P0753S) at 30 °C for 30 min. Co-immunoprecipitated Slp1^{Cdc20} was detected

with rabbit polyclonal anti-Slp1 antibodies, and the phosphorylated form of Slp1^{Cdc20}

was detected with rabbit polyclonal anti-pSer28/pThr31 antibodies respectively.

747 Purification of APC/C and analyses by mass spectrometry

To prepare APC/C for mass spectrometry analyses, Apc15-GFP was purified from

6-liter cultures of *nda3-KM311* arrested cells with desired genetic background. Cells

were disrupted and cell lysates were prepared as described above for routine
immunoprecipitation experiments.

For mass spectrometry analyses, purified samples were first run on PAGE gels, after
staining of gels with Coomassie blue, excised gel segments were subjected to in-gel
trypsin (Promega, V5111) digestion and dried. Samples were then analyzed on a
nanoElute (plug-in V1.1.0.27; Bruker, Bremen, Germany) coupled to a timsTOF Pro
(Bruker, Bremen, Germany) equipped with a CaptiveSpray source. Peptides were
separated on a 15cm X 75μm analytical column, 1.6 μm C18 beads with a packed
emitter tip (IonOpticks, Australia). The column temperature was maintained at 55 °C
using an integrated column oven (Sonation GmbH, Germany). The column was
equilibrated using 4 column volumes before loading sample in 100% buffer A (99.9%
MilliQ water, 0.1% FA) (Both steps performed at 980 bar). Samples were separated at
400 nL/min using a linear gradient from 2% to 25% buffer B (99.9% ACN, 0.1% FA)
over 90 min before ramping to 37% buffer B (10min), ramp to 80% buffer B (10min)
and sustained for 10 min (total separation method time 120 min). The timsTOF Pro
(Bruker, Bremen, Germany) was operated in PASEF mode using Compass Hystar 6.0.
Mass Range 100 to 1700m/z, 1/K0 Start 0.6 V·s/cm^2 End 1.6 V·s/cm^2, Ramp time
110.1ms, Lock Duty Cycle to 100%, Capillary Voltage 1600V, Dry Gas 3 L/min, Dry
Temp 180°C, PASEF settings: 10 MS/MS scans (total cycle time 1.27sec), charge
range 0-5, active exclusion for 0.4 min, Scheduling Target intensity 10000, Intensity
threshold 2500, CID collision energy 42eV. All raw files were analyzed by PEAKS
Studio Xpro software (Bioinformatics Solutions Inc., Waterloo, ON, Canada). Data
was searched against the *S. pombe* proteome sequence database (Uniprot database
with 5117 entries of protein sequences at
https://www.uniprot.org/proteomes/UP000002485). *De novo* sequencing of peptides,
database search and characterizing specific PTMs were used to analyze the raw data;
false discovery rate (FDR) was set to ≤1%, and [-10*log(p)] was calculated
accordingly where p is the probability that an observed match is a random event. The
PEAKS used the following parameters: (i) precursor ion mass tolerance, 20 ppm; (ii)

fragment ion mass tolerance, 0.05 Da (the error tolerance); (iii) tryptic enzyme

specificity with two missed cleavages allowed; (iv) monoisotopic precursor mass and

fragment ion mass; (v) a fixed modification of cysteine carbamidomethylation; and

(vi) variable modifications including N-acetylation of proteins and oxidation of Met.

Ubiquitin affinity pull-down assays

For detection of *in vivo* ubiquitylation of Slp1^{Cdc20}, strains all carried *mts3-1* mutation

were used. sfGFP-tagged wild type Slp1 or Slp1 mutants were expressed in the

genomically integrated form of *P$_{slp1}$-sfGFP-slp1(WT or mutants)::hyg^R* in wild type,

pmk1Δ or *P$_{adh11}$-6xHA-pek1^{DD}* background. Cultures were first grown at 25 °C to

mid-log phase and then shifted to 36°C for 3.5 hours to block cells in mitosis prior to

harvesting and cell lysis. Ubiquitinated proteins were pulled down from yeast lysates

using Tandem Ubiquitin Binding Entities (TUBEs) (LifeSensors, UM402) according

to the manufacturer's instructions. The bound proteins were eluted from washed

Agarose-TUBE2 beads and analyzed by western blotting using a mouse monoclonal

anti-GFP antibody (Beijing Ray Antibody Biotech; RM1008). Cell extracts were also

immunoblotted for GFP and Cdc2 as loading controls (inputs).

Expression and purification of recombinant proteins

GST or MBP fusion constructs of fission yeast full-length or truncated Slp1^{Cdc20},

Pmk1, Pek1 and Atf1 were generated by PCR of the corresponding gene fragments

from yeast genomic DNA or cDNA and cloned in frame into expression vectors

pGEX-4T-1 (GE Healthcare) or pMAL-c2x (New England Biolabs), respectively.

Site-directed mutagenesis was done by standard methods to generate vectors carrying

Pek1-DD(S234D, T238D) or Slp1^{Cdc20} harboring K(Lys)/R(Arg) to E(Glu) or A(Ala)

mutations at individual or combined sites including K19, K20, R21, K47, R48, K140,

K141, K161, R162, R163, K472, R473. Integrity of cloned DNA was verified by

sequencing analysis. All recombinant GST- or MBP-fusion proteins were expressed in

Escherichia coli BL21 (DE3) cells. The recombinant proteins were purified by

glutathione Sepharose 4B (GE Healthcare) or Amylose Resin High Flow (New

England Biolabs) respectively according to the manufacturers' instructions.

In vitro **pulldown assay**

The recombinant MBP-Slp1[Cdc20] and MBP, or GST-Pmk1 and GST proteins were

produced in bacteria and purified as described above. To detect the interaction

between bacterially expressed Slp1[Cdc20] and Pmk1-HA expressed in yeast, about 1 μg

of MBP-Slp1[Cdc20] or MBP immobilized on amylose resin was incubated with cleared

cell lysates prepared from yeast strains *nda3-KM311 pmk1-6His-HA*, which were HU

synchronized and arrested at prometaphase by being grown at 18 °C for 6 hours. To

detect the direct interaction between bacterially expressed Slp1 and Pmk1, about 1 μg

of GST-Pmk1 or GST immobilized on glutathione Sepharose 4B resin was incubated

with cleared bacterial lysates expressing MBP-Slp1[Cdc20]. Samples were incubated for

1-3 hours at 4 °C. Resins were washed 3 times with lysis buffer (6 mM Na_2HPO_4, 4

mM NaH_2PO_4, 1% NP-40, 150 mM NaCl, 2 mM EDTA, 50 mM NaF, 0.1 mM

Na_3VO_4, plus protease inhibitors), suspended in SDS sample buffer, and then subject

to SDS-PAGE electrophoresis and Coomassie brilliant blue (CBB) staining.

Non-radioactive *in vitro* kinase assay

GST-Slp1[Cdc20], GST-Atf1, GST-Pmk1, and GST-Pek1[DD] fusions were expressed and

purified from *Escherichia coli* with glutathione Sepharose 4B as described above.

Pmk1-HA-His and Cdc13-GFP were purified from yeast lysates by immuprecipitation

with anti-HA or anti-GFP antibodies respectively.

For detection of phosphorylation of Slp1[Cdc20] and Atf1 by yeast derived Pmk1,

glutathione Sepharose 4B beads bound with GST-Slp1[Cdc20] or GST-Atf1 and agarose

beads bound with Pmk1-HA-His were mixed and washed 3 times with kinase buffer

(50 mM Tris-HCl, pH 7.5, 10 mM $MgCl_2$, 1 mM EGTA), then the reactions were

incubated in kinase buffer with 20 μM ATPγS (Sigma-Aldrich; A1388) at 30°C for

45 min. The kinase reactions were stopped by adding 20 mM EDTA, and the reaction

mixture was alkylated after incubation at room temperature with 2.5 mM

p-nitrobenzyl mesylate (PNBM) (Abcam; ab138910) for 1 hr. The alkylation reaction

was stopped by boiling in SDS-PAGE loading buffer. Phosphorylated proteins were

detected with rabbit monoclonal anti-Thiophosphate ester antibody (Abcam,

ab239919).

For detection of phosphorylation of Slp1^{Cdc20} at Thr480 by bacterially expressed

Pmk1, beads bound with GST-Pmk1, GST-Pek1(DD) and fused substrates

GST-Slp1$^{(456-488aa)}$ or GST-Slp1$^{(456-488aa, T480A)}$ were washed and incubated in kinase

buffer (50 mM Tris-HCl, pH 7.5, 10 mM MgCl$_2$, 1 mM EGTA) with 10 mM ATP

(Sangon Biotech; A600020-0005) at 30°C for 45 min. The reaction was stopped by

boiling in SDS-PAGE loading buffer. Phosphorylated proteins were detected with

rabbit polyclonal anti-pThr480 antibodies.

For detection of phosphorylation of Slp1^{Cdc20} at Ser28/Thr31 by Cdk1, beads bound

with bacterially expressed MBP-Slp1$^{(1-190aa)}$ or MBP-Slp1$^{(1-190aa,S28A;T31A)}$ and agarose

beads bound with Cdc13-GFP/Cdc2 from yeast lysate were mixed and washed 3 times

with kinase buffer (10 mM Tris-HCl, pH 7.4, 10 mM MgCl$_2$, 1 mM DTT), then the

reactions were incubated in kinase buffer with 10 mM ATP (Sangon Biotech;

A600020-0005) at 30°C for 45 min. To inactivate Cdc2-as1, 5 μM 1-NM-PP1

(Toronto Research Chemicals; A603003) was added into the reaction. The reaction

was stopped by boiling in SDS-PAGE loading buffer. Phosphorylated proteins were

detected with rabbit polyclonal anti-pSer28/pThr31 antibodies.

Antibodies for immunoblotting

The phospho-site specific antibodies were made in an in-house facility (for

anti-Slp1-pS28/pT31) or by ABclonal Technology Co., Ltd. (Wuhan, China) (for

anti-Slp1-pT480) by immunizing rabbits with pS28/pT31- or pT480-containing

peptides (LVFPN(S-p)PI(T-p)PLHQQ or PITK(T-p)PSSS, respectively) coupled to

haemocyanin (Sigma). The purified antibodies were concentrated to 0.2 mg/mL and

used at 1:100 dilution. The following antibodies used for immunoblot analyses were

purchased from the indicated commercial sources and were used at the indicated dilution: peroxidase-anti-peroxidase (PAP) soluble complex (Sigma-Aldrich; P1291; 1:10,000); rabbit polyclonal anti-Myc (GeneScript; A00172-40; 1:2,000); mouse monoclonal anti-GFP (Beijing Ray Antibody Biotech; RM1008; 1:2,000); rat monoclonal anti-HA (Roche, Cat. No. 11 867 423 001; 1:2,000); rabbit polyclonal anti-Slp1 (Kim et al., 1998) (1:500); rabbit polyclonal anti-phospho-p44/42 (detecting activated Pmk1) (Cell Signaling Technology; #9101; 1:1,000); rabbit monoclonal anti-phospho-p38 (Cell Signaling Technology; #4511; 1:1,000); rabbit polyclonal anti-PSTAIRE (detecting Cdc2) (Santa Cruz Biotechnology; sc-53; 1:1,000); rabbit monoclonal anti-Thiophosphate ester antibody (Abcam; ab239919; 1:5,000); goat anti-GST HRP-conjugated antibody (RRID:AB_771429; GE Healthcare; 1:10,000). Secondary antibodies used were goat anti-mouse or goat anti-rabbit polyclonal IgG (H+L) HRP conjugates (Thermo Fisher Scientific; #31430 or #32460; 1:10,000).

RT-qPCR

For RT-qPCR of $slp1^+$ mRNA, *nda3-KM311* strains were grown, synchronized by HU and arrested at metaphase with SAC being activated as described above. Cells were collected and total cellular RNA was isolated by using TRIZOL method. cDNA was prepared using oligo(dT) and PrimeScript reverse transcriptase (Takara, D2680A). Real-time PCR was performed in the presence of SYBR Green using an Applied Biosystems 7900HT light cycler. Relative RNA levels were calculated using the *ΔCT* method and normalized to $act1^+$ levels.

Fluorescence microscopy

Cdc13-GFP proteins were observed in cells after fixation with cold methanol. For DAPI staining of nuclei, cells were fixed with cold methanol, washed in PBS and resuspended in PBS plus 1 μg/ml DAPI. Photomicrographs of cells were obtained using a Nikon 80i fluorescence microscope coupled to a cooled CCD camera (Hamamatsu, ORCA-ER) or a Perkin Elmer spinning-disk confocal microscope (UltraVIEW® VoX) with a 100x NA 1.49 TIRF oil immersion objective (Nikon)

900 coupled to a cooled CCD camera (9100-50 EMCCD; Hamamatsu Photonics) and

901 spinning disk head (CSU-X1, Yokogawa). Image processing and analysis were carried

902 out using Element software (Nikon), ImageJ software (National Institutes of Health)

903 and Adobe Photoshop.

Statistical analyses and reproducibility

No statistical methods were used to predetermine sample size. Sample size was as

large as practicable for all experiments. The experiments were randomized and the

investigators were blinded to group allocation during experiments but not blinded to

outcome assessment and data analysis.

All experiments were independently repeated two to more than three times with

similar results obtained. For quantitative analyses of each time course experiment,

300 cells were counted for each time point or sample. The same sample was not

measured or counted repeatedly. No data were excluded from our studies. Data

collection and statistical analyses were performed using Microsoft Office Excel or

GraphPad Prism 6 softwares. Data are expressed as mean values with error bars

(± standard deviation/s.d.) and were compared using two-tailed Student's *t*-tests unless

indicated otherwise.

Acknowledgments:

We thank Kathy Gould, Li-lin Du, Silke Hauf, Jonathan Millar, Takashi Toda,

Yoshinori Watanabe, Elena Hidalgo and National BioResource Project (NBRP),

Japan (http://yeast.nig.ac.jp/yeast/) for fission yeast strains or plasmids; Rafael Daga

for communication on unpublished results; Jose Cansado for advice on active MAPK

detection. We also thank Yaying Wu, Zheni Xu and Chang-chuan Xie for mass

spectrometry experiments and data analyses; Yu-ting He for contributing a few time

course experimental data; and Shiyu Huang, Jun-ming Zhuang and Cancan Zhang for

help with construction of some plasmids or yeast strains.

Funding:

This work was supported by grants from the National Natural Science Foundation of China (No. 32170731, No. 31671411) to Q.W. Jin.

Author Contributions Statement:

W.S., J.C., and C.J. performed all microscopy experiments with the help from Z.L.. L.S., X.C., W.S., C.S., L.L., S.B. and S.W. performed all immunoblotting, co-immunoprecipitation experiments, *in vitro* binding assays and *in vitro* kinase aassays. S.W. and X.W. performed large-scale protein purifications. X.C. and S.W. carried out genetic analyses. Q.J. and Y.W. conceived the study. Q.J. acquired funding. Y.W. and Q.J. designed the experiments and supervised the research. Y.W. and Q.J. wrote the paper with the inputs from Z.L..

Competing Interests Statement:

The authors declare no competing or financial interests.

Data availability statement

The authors confirm that all data supporting the findings of this study are available within the manuscript main figures, supplementary figures, supplementary tables and source data files. The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD048800.

References

Akera, T., Goto, Y., Sato, M., Yamamoto, M., and Watanabe, Y. (2015). Mad1 promotes chromosome congression by anchoring a kinesin motor to the kinetochore. Nat Cell Biol *17*, 1124-1133.

Alfieri, C., Chang, L., Zhang, Z., Yang, J., Maslen, S., Skehel, M., and Barford, D. (2016). Molecular basis of APC/C regulation by the spindle assembly checkpoint. Nature *536*, 431-436.

Bai, S., Sun, L., Wang, X., Wang, S.M., Luo, Z.Q., Wang, Y.M., and Jin, Q.W. (2022). Recovery from spindle checkpoint-mediated arrest requires a novel Dnt1-dependent APC/C activation mechanism. Plos Genetics *18*, e1010397.

Bardwell, A.J., Flatauer, L.J., Matsukuma, K., Thorner, J., and Bardwell, L. (2001). A conserved

docking site in MEKs mediates high-affinity binding to MAP kinases and cooperates with a scaffold

protein to enhance signal transmission. J Biol Chem *276*, 10374-10386.

Bardwell, L., and Thorner, J. (1996). A conserved motif at the amino termini of MEKs might mediate

high-affinity interaction with the cognate MAPKs. Trends in biochemical sciences *21*, 373-374.

Cansado, J., Soto, T., Franco, A., Vicente-Soler, J., and Madrid, M. (2021). The Fission Yeast Cell

Integrity Pathway: A Functional Hub for Cell Survival upon Stress and Beyond. J Fungi (Basel) *8*,

1-32.

Chao, W.C., Kulkarni, K., Zhang, Z., Kong, E.H., and Barford, D. (2012). Structure of the mitotic

checkpoint complex. Nature *484*, 208-213.

Chen, R.H. (2004). Phosphorylation and activation of Bub1 on unattached chromosomes facilitate the

spindle checkpoint. EMBO J *23*, 3113-3121.

Chen, Y.H., Wang, G.Y., Hao, H.C., Chao, C.J., Wang, Y., and Jin, Q.W. (2017). Facile manipulation of

protein localization in fission yeast through binding of GFP-binding protein to GFP. Journal of Cell

Science *130*, 1003-1015.

Chi, J.J., Li, H., Zhou, Z., Izquierdo-Ferrer, J., Xue, Y., Wavelet, C.M., Schiltz, G.E., Zhang, B.,

Cristofanilli, M., Lu, X.*, et al.* (2019). A novel strategy to block mitotic progression for targeted

therapy. EBioMedicine *49*, 40-54.

Chung, E., and Chen, R.H. (2003). Phosphorylation of Cdc20 is required for its inhibition by the

spindle checkpoint. Nat Cell Biol *5*, 748-753.

Coffman, V.C., Sees, J.A., Kovar, D.R., and Wu, J.Q. (2013). The formins Cdc12 and For3 cooperate

during contractile ring assembly in cytokinesis. J Cell Biol *203*, 101-114.

D'Angiolella, V., Mari, C., Nocera, D., Rametti, L., and Grieco, D. (2003). The spindle checkpoint

requires cyclin-dependent kinase activity. Genes Dev *17*, 2520-2525.

Danielsen, J.M.R., Sylvestersen, K.B., Bekker-Jensen, S., Szklarczyk, D., Poulsen, J.W., Horn, H.,

Jensen, L.J., Mailand, N., and Nielsen, M.L. (2011). Mass Spectrometric Analysis of Lysine

Ubiquitylation Reveals Promiscuity at Site Level. Molecular & Cellular Proteomics *10*.

Edreira, T., Celador, R., Manjón, E., and Sánchez, Y. (2020). A novel checkpoint pathway controls

actomyosin ring constriction trigger in fission yeast. eLife *9*.

Fujimitsu, K., Grimaldi, M., and Yamano, H. (2016). Cyclin-dependent kinase 1-dependent activation

of APC/C ubiquitin ligase. Science *352*, 1121-1124.

Gomez-Gil, E., Martin-Garcia, R., Vicente-Soler, J., Franco, A., Vazquez-Marin, B., Prieto-Ruiz, F.,

Soto, T., Perez, P., Madrid, M., and Cansado, J. (2020). Stress-activated MAPK signaling controls

fission yeast actomyosin ring integrity by modulating formin For3 levels. eLife *9*, e57951.

Greil, C., Engelhardt, M., and Wasch, R. (2022). The Role of the APC/C and Its Coactivators Cdh1 and

Cdc20 in Cancer Development and Therapy. Front Genet *13*, 941565.

Hartmuth, S., and Petersen, J. (2009). Fission yeast Tor1 functions as part of TORC1 to control mitotic

entry through the stress MAPK pathway following nutrient stress. J Cell Sci *122*, 1737-1746.

Heinrich, S., Geissen, E.M., Kamenz, J., Trautmann, S., Widmer, C., Drewe, P., Knop, M., Radde, N.,

Hasenauer, J., and Hauf, S. (2013). Determinants of robustness in spindle assembly checkpoint

signalling. Nat Cell Biol *15*, 1328-1339.

Hiraoka, Y., Toda, T., and Yanagida, M. (1984). The NDA3 gene of fission yeast encodes beta-tubulin:

a cold-sensitive nda3 mutation reversibly blocks spindle formation and chromosome movement in

mitosis. Cell *39*, 349-358.

Hjerpe, R., Aillet, F., Lopitz-Otsoa, F., Lang, V., England, P., and Rodriguez, M.S. (2009). Efficient

protection and isolation of ubiquitylated proteins using tandem ubiquitin-binding entities. EMBO Rep

10, 1250–1258.

Izawa, D., and Pines, J. (2015). The mitotic checkpoint complex binds a second CDC20 to inhibit

active APC/C. Nature *517*, 631-634.

Jeong, S.M., Bui, Q.T., Kwak, M., Lee, J.Y., and Lee, P.C. (2022). Targeting Cdc20 for cancer therapy.

Biochim Biophys Acta Rev Cancer *1877*, 188824.

Jia, L., Li, B., and Yu, H. (2016). The Bub1-Plk1 kinase complex promotes spindle checkpoint

signalling through Cdc20 phosphorylation. Nature communications *7*, 10818.

Kapanidou, M., Curtis, N.L., and Bolanos-Garcia, V.M. (2017). Cdc20: At the Crossroads between

Chromosome Segregation and Mitotic Exit. Trends in biochemical sciences *42*, 193-205.

Kim, D.U., Hayles, J., Kim, D., Wood, V., Park, H.O., Won, M., Yoo, H.S., Duhig, T., Nam, M.,

Palmer, G*., et al.* (2010). Analysis of a genome-wide set of gene deletions in the fission yeast

Schizosaccharomyces pombe. Nat Biotechnol *28*, 617-623.

Kim, S.H., Lin, D.P., Matsumoto, S., Kitazono, A., and Matsumoto, T. (1998). Fission yeast Slp1: an

effector of the Mad2-dependent spindle checkpoint. Science *279*, 1045-1047.

Kim, T., Lara-Gonzalez, P., Prevo, B., Meitinger, F., Cheerambathur, D.K., Oegema, K., and Desai, A.

(2017). Kinetochores accelerate or delay APC/C activation by directing Cdc20 to opposing fates. Genes

Dev *31*, 1089-1094.

Kraft, C., Herzog, F., Gieffers, C., Mechtler, K., Hagting, A., Pines, J., and Peters, J.M. (2003). Mitotic

regulation of the human anaphase-promoting complex by phosphorylation. EMBO J *22*, 6598-6609.

Kramer, E.R., Scheuringer, N., Podtelejnikov, A.V., Mann, M., and Peters, J.M. (2000). Mitotic

regulation of the APC activator proteins CDC20 and CDH1. Mol Biol Cell *11*, 1555-1569.

Labit, H., Fujimitsu, K., Bayin, N.S., Takaki, T., Gannon, J., and Yamano, H. (2012).

Dephosphorylation of Cdc20 is required for its C-box-dependent activation of the APC/C. EMBO J *31*,

3351-3362.

Lopez-Aviles, S., Grande, M., Gonzalez, M., Helgesen, A.L., Alemany, V., Sanchez-Piris, M., Bachs,

O., Millar, J.B., and Aligue, R. (2005). Inactivation of the Cdc25 phosphatase by the stress-activated

Srk1 kinase in fission yeast. Mol Cell *17*, 49-59.

Lopez-Aviles, S., Lambea, E., Moldon, A., Grande, M., Fajardo, A., Rodriguez-Gabriel, M.A., Hidalgo,

E., and Aligue, R. (2008). Activation of Srk1 by the mitogen-activated protein kinase Sty1/Spc1

precedes its dissociation from the kinase and signals its degradation. Mol Biol Cell *19*, 1670-1679.

Madrid, M., Soto, T., Khong, H.K., Franco, A., Vicente, J., Pérez, P., Gacto, M., and Cansado, J.

(2006). Stress-induced response, localization, and regulation of the Pmk1 cell integrity pathway in.

Journal of Biological Chemistry *281*, 2033-2043.

Madrid, M., Vázquez-Marín, B., Franco, A., Soto, T., Vicente-Soler, J., Gacto, M., and Cansado, J.

(2016). Multiple crosstalk between TOR and the cell integrity MAPK signaling pathway in fission

yeast. Sci Rep-Uk *6*.

Mansfeld, J., Collin, P., Collins, M.O., Choudhary, J.S., and Pines, J. (2011). APC15 drives the turnover

of MCC-CDC20 to make the spindle assembly checkpoint responsive to kinetochore attachment. Nat

Cell Biol *13*, 1234-1243.

May, K.M., Paldi, F., and Hardwick, K.G. (2017). Fission Yeast Apc15 Stabilizes MCC-Cdc20-APC/C

Complexes, Ensuring Efficient Cdc20 Ubiquitination and Checkpoint Arrest. Curr Biol *27*, 1221-1228.

McAinsh, A.D., and Kops, G.J.P.L. (2023). Principles and dynamics of spindle assembly checkpoint

signalling. Nat Rev Mol Cell Bio *24*, 543-559.

Minshull, J., Sun, H., Tonks, N.K., and Murray, A.W. (1994). A MAP kinase-dependent spindle
assembly checkpoint in Xenopus egg extracts. Cell *79*, 475-486.

Murray, A.W. (2011). A brief history of error. Nat Cell Biol *13*, 1178-1182.

Musacchio, A. (2015). The Molecular Biology of Spindle Assembly Checkpoint Signaling Dynamics.
Curr Biol *25*, R1002-1018.

Pan, J., and Chen, R.H. (2004). Spindle checkpoint regulates Cdc20p stability in Saccharomyces
cerevisiae. Genes Dev *18*, 1439-1451.

Perez, P., and Cansado, J. (2010). Cell integrity signaling and response to stress in fission yeast. Curr
Protein Pept Sci *11*, 680-692.

Peters, J.M. (2006). The anaphase promoting complex/cyclosome: a machine designed to destroy. Nat
Rev Mol Cell Biol *7*, 644-656.

Petersen, J., and Hagan, I.M. (2005). Polo kinase links the stress pathway to cell cycle control and tip
growth in fission yeast. Nature *435*, 507-512.

Petersen, J., and Nurse, P. (2007). TOR signalling regulates mitotic commitment through the stress
MAP kinase pathway and the Polo and Cdc2 kinases. Nat Cell Biol *9*, 1263-1272.

Pinna, L.A., and Ruzzene, M. (1996). How do protein kinases recognize their substrates? Bba-Mol Cell
Res *1314*, 191-225.

Plotnikov, A., Zehorai, E., Procaccia, S., and Seger, R. (2011). The MAPK cascades: signaling
components, nuclear roles and mechanisms of nuclear translocation. Biochim Biophys Acta *1813*,
1619-1633.

Primorac, I., and Musacchio, A. (2013). Panta rhei: the APC/C at steady state. J Cell Biol *201*, 177-189.

Qiao, R., Weissmann, F., Yamaguchi, M., Brown, N.G., VanderLinden, R., Imre, R., Jarvis, M.A.,
Brunner, M.R., Davidson, I.F., Litos, G., *et al.* (2016). Mechanism of APC/CCDC20 activation by
mitotic phosphorylation. Proc Natl Acad Sci U S A *113*, E2570-2578.

Repetto, M.V., Winters, M.J., Bush, A., Reiter, W., Hollenstein, D.M., Ammerer, G., Pryciak, P.M., and
Colman-Lerner, A. (2018). CDK and MAPK Synergistically Regulate Signaling Dynamics via a Shared
Multi-site Phosphorylation Region on the Scaffold Protein Ste5. Molecular Cell *69*, 938-+.

Ronkina, N., and Gaestel, M. (2022). MAPK-Activated Protein Kinases: Servant or Partner? Annu Rev
Biochem, 12.11–12.36.

Sacristan-Reviriego, A., Madrid, M., Cansado, J., Martin, H., and Molina, M. (2014). A conserved
non-canonical docking mechanism regulates the binding of dual specificity phosphatases to cell
integrity mitogen-activated protein kinases (MAPKs) in budding and fission yeasts. PLoS One *9*,
e85390.

Saitoh, S., Takahashi, K., and Yanagida, M. (1997). Mis6, a fission yeast inner centromere protein, acts
during G1/S and forms specialized chromatin required for equal segregation. Cell *90*, 131-143.

Sczaniecka, M., Feoktistova, A., May, K.M., Chen, J.S., Blyth, J., Gould, K.L., and Hardwick, K.G.
(2008). The spindle checkpoint functions of Mad3 and Mad2 depend on a Mad3 KEN box-mediated
interaction with Cdc20-anaphase-promoting complex (APC/C). J Biol Chem *283*, 23039-23047.

Sewart, K., and Hauf, S. (2017). Different Functionality of Cdc20 Binding Sites within the Mitotic
Checkpoint Complex. Curr Biol *27*, 1213-1220.

Shiozaki, K., and Russell, P. (1995a). Cell-cycle control linked to extracellular environment by MAP
kinase pathway in fission yeast. Nature *378*, 739-743.

Shiozaki, K., and Russell, P. (1995b). Counteractive roles of protein phosphatase 2C (PP2C) and a
MAP kinase kinase homolog in the osmoregulation of fission yeast. Embo J *14*, 492-502.

Shiozaki, K., Shiozaki, M., and Russell, P. (1998). Heat stress activates fission yeast Spc1/StyI MAPK
by a MEKK-independent mechanism. Mol Biol Cell *9*, 1339-1349.

Smith, D.A., Toone, W.M., Chen, D., Bahler, J., Jones, N., Morgan, B.A., and Quinn, J. (2002). The
Srk1 protein kinase is a target for the Sty1 stress-activated MAPK in fission yeast. J Biol Chem *277*,
33411-33421.

Steen, J.A., Steen, H., Georgi, A., Parker, K., Springer, M., Kirchner, M., Hamprecht, F., and Kirschner,
M.W. (2008). Different phosphorylation states of the anaphase promoting complex in response to
antimitotic drugs: a quantitative proteomic analysis. Proc Natl Acad Sci U S A *105*, 6069-6074.

Sudakin, V., Chan, G.K., and Yen, T.J. (2001). Checkpoint inhibition of the APC/C in HeLa cells is
mediated by a complex of BUBR1, BUB3, CDC20, and MAD2. J Cell Biol *154*, 925-936.

Sugiura, R., Toda, T., Dhut, S., Shuntoh, H., and Kuno, T. (1999). The MAPK kinase Pek1 acts as a
phosphorylation-dependent molecular switch. Nature *399*, 479-483.

Sullivan, M., and Morgan, D.O. (2007). Finishing mitosis, one step at a time. Nat Rev Mol Cell Biol *8*,
894-903.

Swaffer, M.P., Jones, A.W., Flynn, H.R., Snijders, A.P., and Nurse, P. (2018). Quantitative
Phosphoproteomics Reveals the Signaling Dynamics of Cell-Cycle Kinases in the Fission Yeast. Cell
Reports *24*, 503-514.

Takenaka, K., Gotoh, Y., and Nishida, E. (1997). MAP kinase is required for the spindle assembly
checkpoint but is dispensable for the normal M phase entry and exit in Xenopus egg cell cycle extracts.
J Cell Biol *136*, 1091-1097.

Tang, Z., Shu, H., Oncel, D., Chen, S., and Yu, H. (2004). Phosphorylation of Cdc20 by Bub1 provides
a catalytic mechanism for APC/C inhibition by the spindle checkpoint. Mol Cell *16*, 387-397.

Uzunova, K., Dye, B.T., Schutz, H., Ladurner, R., Petzold, G., Toyoda, Y., Jarvis, M.A., Brown, N.G.,
Poser, I., Novatchkova, M.*, et al.* (2012). APC15 mediates CDC20 autoubiquitylation by APC/C(MCC)
and disassembly of the mitotic checkpoint complex. Nat Struct Mol Biol *19*, 1116-1123.

Vanoosthuyse, V., and Hardwick, K.G. (2009). A novel protein phosphatase 1-dependent spindle
checkpoint silencing mechanism. Curr Biol *19*, 1176-1181.

Wang, Y., Li, W.Z., Johnson, A.E., Luo, Z.Q., Sun, X.L., Feoktistova, A., McDonald, W.H., McLeod, I.,
Yates, J.R., 3rd, Gould, K.L.*, et al.* (2012). Dnt1 acts as a mitotic inhibitor of the spindle checkpoint
protein dma1 in fission yeast. Mol Biol Cell *23*, 3348-3356.

Yamada, H.Y., Matsumoto, S., and Matsumoto, T. (2000). High dosage expression of a zinc finger
protein, Grt1, suppresses a mutant of fission yeast slp1(+), a homolog of CDC20/p55CDC/Fizzy. J Cell
Sci *113 (Pt 22)*, 3989-3999.

Yamaguchi, M., VanderLinden, R., Weissmann, F., Qiao, R., Dube, P., Brown, N.G., Haselbach, D.,
Zhang, W., Sidhu, S.S., Peters, J.M.*, et al.* (2016). Cryo-EM of Mitotic Checkpoint Complex-Bound
APC/C Reveals Reciprocal and Conformational Regulation of Ubiquitin Ligation. Mol Cell *63*,
593-607.

Yamano, H. (2019). APC/C: current understanding and future perspectives. F1000Res *8*, 725.

Yen, A.H., and Yang, J.L. (2010). Cdc20 Proteolysis Requires p38 MAPK Signaling and
Cdh1-Independent APC/C Ubiquitination During Spindle Assembly Checkpoint Activation by
Cadmium. Journal of Cellular Physiology *223*, 327-334.

Yoon, H.J., Feoktistova, A., Wolfe, B.A., Jennings, J.L., Link, A.J., and Gould, K.L. (2002). Proteomics
analysis identifies new components of the fission and budding yeast anaphase-promoting complexes.
Curr Biol *12*, 2048-2054.

Yu, H. (2007). Cdc20: a WD40 activator for a cell cycle degradation machine. Mol Cell *27*, 3-16.

Yudkovsky, Y., Shteinberg, M., Listovsky, T., Brandeis, M., and Hershko, A. (2000). Phosphorylation

of Cdc20/fizzy negatively regulates the mammalian cyclosome/APC in the mitotic checkpoint.

Biochem Biophys Res Commun *271*, 299-304.

Zhang, S., Chang, L., Alfieri, C., Zhang, Z., Yang, J., Maslen, S., Skehel, M., and Barford, D. (2016).

Molecular mechanism of APC/C activation by mitotic phosphorylation. Nature *533*, 260-264.

Zhao, Y., and Chen, R.H. (2006). Mps1 phosphorylation by MAP kinase is required for kinetochore

localization of spindle-checkpoint proteins. Curr Biol *16*, 1764-1769.

Zuin, A., Carmona, M., Morales-Ivorra, I., Gabrielli, N., Vivancos, A.P., Ayte, J., and Hidalgo, E.

(2010). Lifespan extension by calorie restriction relies on the Sty1 MAP kinase stress pathway. EMBO

J *29*, 981-991.

1150

A

Activated signaling

		MAPKKK	MAPKK	MAPK
CIP	*WT*	Mkh1 →	Pek1 →	Pmk1
SAP	*WT*	Wak1/Win1 →	Wis1 →	Sty1
PSP	*WT*	Byr2 →	Byr1 →	Spk1

Disrupted signaling

	MAPKKK	MAPKK	MAPK
pmk1Δ	Mkh1 →	Pek1 →	P̶m̶k1
sty1Δ / sty1^T97A	Wak1/Win1 →	Wis1 →	S̶t̶y1
spk1Δ	Byr2 →	Byr1 →	S̶p̶k1

Constitutively activated signaling

	MAPKKK	MAPKK	MAPK
pek1^DD	Mkh1 →	Pek1 ⟹	Pmk1
wis1^DD	Wak1/Win1 →	Wis1 ⟹	Sty1

B

YE + TBZ, 30 °C

	0 µg/ml	15 µg/ml
wild-type		
pmk1Δ		
sty1Δ		
spk1Δ		
wild-type		
mad2Δ		
P_{adh21}-pek1^{DD}		
P_{adh11}-pek1^{DD}		
P_{adh21}-wis1^{DD}		
P_{adh11}-wis1^{DD}		

C

nda3-KM311 background strains:

12 mM HU, 2 hr;
6 mM HU, 3.5 hr

30 °C
18 °C
w/o HU, 6 hr

0 min, 10 min, 20 min, 30 min, 40 min, 50 min

Cdc13-GFP | DNA (DAPI) | merge

→ Cdc13 at SPB enriched   * Cdc13 at SPB degraded

D

nda3-KM311 background

- ● *wild-type*
- ○ *pmk1Δ*
- ○ *sty1-T97A* (BrB-PP1 treated)
- ○ *spk1Δ*

Cells with Cdc13-GFP on SPBs (%)
Time after release at 30 °C (min)

- ● *wild-type*
- ● *P_{adh11}-pek1^{DD}*
- ● *P_{adh11}-wis1^{DD}*

Cells with Cdc13-GFP on SPBs (%)
Time after release at 30 °C (min)

Figure 1. *pmk1Δ* and *sty1-T97A* **mutants display spindle checkpoint activation defects, while** *pek1^{DD}* **and** *wis1^{DD}* **mutants are defective in checkpoint silencing.**

(**A**) Schematic of core modules of three *Schizosaccharomyces pombe* mitogen-activated protein kinase (MAPK) signaling pathways. Each cascade consists of three core kinases (MAP kinase kinase kinase (MAPKKK), MAP kinase kinase (MAPKK), and MAPK). CIP, the cell integrity pathway; SAP, the stress-activated pathway; PSP, pheromone signaling pathway. These pathways can be disrupted by *pmk1Δ*, *sty1Δ* or ATP analogue-sensitive mutation *sty1-T97A* or *spk1Δ*, and constitutively activated by mutations in MAPKKs Pek1 (*pek1^{DD}*, *pek1-S234D;T238D*) or Wis1 (*wis1^{DD}*, *wis1-S469D;T473D*) respectively.

(**B**) Serial dilution assay on TBZ sensitivity of all MAPK deletion mutants and *pek1^{DD}*- or *wis1^{DD}*-overexpressing mutants. *mad2Δ* is a positive control. Note P_{adh11} is a stronger version of P_{adh21} promoter.

(**C**) Schematic depiction of the experiment design for time-course analyses on SAC or APC/C activation. *nda3-KM311* cells carrying Cdc13-GFP were grown, synchronized with HU and treated at 18 ℃ for 6 hours to activate SAC, and finally shifted back to the permissive temperature 30 ℃. Samples were collected at 10 min intervals and subjected to microscopy analyses. Example pictures of cells with Cdc13-GFP signals enriched or disappeared at spindle pole bodies (SPBs) are shown. Scale bar, 5 μm.

(**D**) Time-course analyses of SAC activation and inactivation in *nda3-KM311 cdc13-GFP* strains with indicated genotypes. *sty1-T97A* was inactivated by 5μM 3-BrB-PP1.

Figure 2. Upon spindle checkpoint activation, Slp1^{Cdc20} levels are reduced in *pek1^{DD}* and the MCC-APC/C association is enhanced in *wis1^{DD}* cells compared to wild-type cells.

(**A**) Co-immunoprecipitation analysis on MCC-APC/C association. Cells with indicated genotypes were grown at 30 ℃ to mid-log phase and arrested at 18 ℃ for 6 hours. Lid1-TAP was immunoprecipitated and associated Mad2, Mad3 and Slp1^{Cdc20} were detected by immunoblotting. The amount of co-immunoprecipitated Mad2, Mad3 and Slp1^{Cdc20} was quantified by being normalized to those of total immunoprecipitated Lid1 in each sample, with the relative ratio between Mad2-GFP plus Mad3-GFP or Slp1^{Cdc20} and Lid1-TAP in wild-type sample set as 1.0. Blots are representative of three independent experiments. *p* values were calculated against wild-type cells. *, *p*<0.05; **, *p*<0.01; ***, *p*<0.001; ****, *p*<0.0001; n.s., not significant.

(**B, C**) Immunoblot analysis of Slp1^{Cdc20} abundance in *nda3-KM311* cells treated at 18 ℃ for 6 hr. Slp1^{Cdc20} levels were quantified with the relative ratio between Slp1^{Cdc20} and Cdc2 in wild-type strain set as 1.0. Phosphorylated Pmk1 (Pmk1-P) or phosphorylated Sty1 (Sty1-P) in (A) were detected using anti-phospho p42/44 and anti-phospho p38 antibodies and represents activated CIP or SAP signaling, respectively. *sty1-T97A* was inactivated by 5μM 3-BrB-PP1. Blots shown are the representative of three independent experiments. *p* values were calculated against wild-type cells. ***, *p*<0.001; n.s., not significant.

(**D**) RT-qPCR analysis of mRNA levels of *slp1^{+}*. Cells with indicated genotypes were grown and treated as in (A-C) before RNA extraction. The relative fold-change (*slp1^{+}*/*act1^{+}*) in mRNA expression was calculated with that in wild-type cells being normalized to 1.0. Note mRNA level of *slp1^{+}* in *pek1^{DD}* mutant is not decreased. Error bars indicate mean ± standard deviation of three independent experiments. Two-tailed unpaired *t*-test was used to derive *p* values. n.s, not significant.

(**E**) Schematic summary of the negative effect of activated CIP and SAP signaling on APC/C activation based on primary phenotype characterization of *pmk1Δ*, *sty1-T97A*, *pek1^{DD}* and *wis1^{DD}* mutants.

Figure 3. Pmk1 directly binds Slp1^{Cdc20} to attenuate its levels upon spindle checkpoint activation.

(**A**) *In vitro* binding assay using bacterially expressed MBP-Slp1^{Cdc20} and yeast lysates prepared from *nda3-KM311 pmk1-HA-6His* cells arrested at 18 °C for 6 hr. Note that weak band detected in MBP sample was due to unspecific background binding to amylose beads. Coomassie blue staining shows inputs for MBP and MBP-Slp1^{Cdc20}.

(**B**) Schematic depiction of the *S. pombe* Slp1^{Cdc20} protein structure with the positions of two confirmed basic-residue patches mediating Slp1^{Cdc20}-Pmk1 association indicated by pink bars. Alignment highlights the conservation of basic-residue patches within 4 *Schizosaccharomyces* species. The deduced Pmk1-docking motifs and different versions of motif mutations are shown. MIM, Mad2-interaction motif; IR, isoleucine-arginine tail.

(**C**) *In vitro* GST pull-down assays with bacterially expressed recombinant GST-Pmk1 and MBP-fusions of wild-type Slp1^{Cdc20} or Slp1^{Cdc20} mutants harboring Pmk1-docking motif mutations. An aliquot of the same amount of MBP-Slp1^{Cdc20} as that added in each GST pull-down reactions was immobilized by amylose resin as the input control. Asterisks indicate unspecific or degraded protein bands.

(**D**) Immunoblot analysis of Slp1^{Cdc20} abundance in *nda3-KM311* cells with indicated genotypes treated at 18 °C for 6 hr. Slp1^{Cdc20} levels were quantified as in Figure 2B and 2C. The experiment was repeated 3 times and the mean value for each sample was calculated. *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$; ****, $p<0.0001$; n.s., not significant.

(**E**) Time-course analyses of SAC activation and inactivation in *nda3-KM311 cdc13-GFP* strains with indicated genotypes. The experiment was repeated 3 times and the mean value for each sample was calculated as in Figure 1D.

(**F**) Schematic summarizing the negative effect of Pmk1-Slp1^{Cdc20} association on Slp1^{Cdc20} abundance and APC/C activation.

Figure 4. Pmk1 synergizes with Cdk1 to phosphorylate and reduce Slp1^{Cdc20} abundance.

(**A**) Non-radioactive *in vitro* phosphorylation assays with bacterially expressed recombinant GST-Slp1^{Cdc20} and Pmk1-HA-His purified from yeast cells. The incorporation of the thiophosphate group was determined using anti-thiophosphate ester antibodies (anti-thioP) as indicative of phosphorylation. Note that the presence or absence of GST-Pek1^{DD} does not affect Slp1^{Cdc20} phosphorylation efficiency. A known Pmk1 substrate Atf1 was used as a positive control (Atf1-P). Asterisks indicate bands corresponding to unspecific or likely degraded proteins.

(**B**) Summary of mass spectrometry data on Slp1^{Cdc20} phosphorylation and ubiquitylation *in vivo* in *nda3-KM311*-arrested cells. Red arrows and filled blue circles denote all detected phosphorylated or ubiquitylated sites respectively, while gray arrows and unfilled circles indicate the absence of phosphorylation or ubiquitylation in some of these sites respectively. Alignment highlights the conservation of most of the detected phosphorylation and ubiquitylation sites within 4 *Schizosaccharomyces* species.

(**C**) *In vitro* phosphorylation assays with bacterially expressed recombinant GST-fusion of Slp1^{Cdc20} fragment (456-488aa), GST-Pmk1 and GST-Pek1^{DD}. The reactions were blotted with pT480 antibodies. Asterisks indicate bands corresponding to unspecific or likely degraded proteins.

(**D**) Immunoblot detection of Slp1^{Cdc20} phosphorylation at T480 *in vivo*. GST-slp1(456-488aa) was purified from *mts3-1* cells with indicated genotypes arrested at 36°C for 3.5 hours, and detected with anti-GST and anti-pThr480 antibodies. Note that pThr480 is absent in *pmk1Δ* cells, and enhanced in *pek1^{DD}* cells relative to that in wild type cells.

(**E**) *In vitro* phosphorylation assays with bacterially expressed recombinant MBP-fusion of Slp1^{Cdc20} fragment (1-190aa) and Cdc13 (cyclin B)-containing Cdk1 complexes purified from metaphase-arrested *nda3-KM311* yeast cells. 1-NM-PP1 was added as inhibitor for analogue-sensitive Cdc2-as. The reactions were blotted with pS28/pT31 antibodies. Asterisks indicate bands corresponding to unspecific or likely degraded proteins.

(**F**) Immunoblot detection of Slp1^{Cdc20} phosphorylation at S28/T31 *in vivo*. Apc15-13myc was immunoprecipitated from *nda3-KM311* cells treated at 18 ℃ for 6 hr and the samples were blotted with pS28/pT31 antibodies. One IP sample from wild

type background was treated with λ-phosphatase. For *cdc2-asM17* cells, 1-NM-PP1

was added to inactivate Cdc2 during culturing.

(**G**) Immunoblot analysis of Slp1^{Cdc20} abundance in *nda3-KM311* cells treated at

18 ℃ for 6 hr. Slp1^{Cdc20} levels were quantified as in Figure 2B and 2C. The

experiment was repeated 3 times and the mean value for each sample was calculated.

*, $p<0.05$; **, $p<0.01$; ***, $p<0.001$; ****, $p<0.0001$; n.s., not significant.

(**H**) Time-course analyses of SAC activation and inactivation in *nda3-KM311*

cdc13-GFP strains with indicated genotypes. The experiment was repeated 3 times

and the mean value for each sample was calculated as in Figure 1D.

(**I**) Schematic summarizing the negative effect of Slp1^{Cdc20} phosphorylation by Pmk1

on its abundance and APC/C activation.

Figure 5. Pmk1- but not Cdk1-mediated Slp1^{Cdc20} phosphorylation promotes its ubiquitylation.

(**A**) Immunoblot analysis of Slp1^{Cdc20} abundance in *apc15$^+$* or *apc15Δ* background cells with indicated genotypes after being treated at 18 ℃ for 6 hr. Slp1^{Cdc20} levels were quantified as in Figure 2B. The experiment was repeated 3 times and the mean value for each sample was calculated. ***, $p<0.001$.

(**B**) Time-course analyses of SAC activation and inactivation in *nda3-KM311 cdc13-GFP* strains with indicated genotypes. The experiments were performed as in Figure 1D.

(**C**) Immunoblot analysis of Slp1^{Cdc20} abundance in K472R, K479R or K472R/K479R mutants. The experiment was repeated 3 times and the mean value for each sample was calculated. *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$; ****, $p<0.0001$.

(**D**) Time-course analyses of SAC activation and inactivation in *nda3-KM311 cdc13-GFP* strains with K472R, K479R or K472R/K479R mutations. The experiments were performed as in (B).

(**E**) Schematic depiction of affinity pull-down assays using TUBE (Tandem Ubiquitin Binding Entity) agarose beads to detect Slp1^{Cdc20} ubiquitylation.

(**F**) TUBE pull-down assays in *mts3-1* strains carrying sfGFP-tagged wild type or mutants of Slp1^{Cdc20} or *pmk1Δ* or *pek1^{DD}* mutations. The TUBE beads-bound samples were blotted with anti-GFP antibodies. Asterisk indicate unspecific bands recognized by anti-GFP antibodies.

(**G**) Schematic summarizing the negative effect of the coupling of Pmk1 phosphorylation and K472/K479-mediated ubiquitylation on Slp1^{Cdc20} abundance and APC/C activation.

Figure 6. Osmotic stress and cell wall damage trigger rapid Pmk1 phosphorylation-dependent Slp1^{Cdc20} downregulation and mitotic exit delay.

(**A**) Schematic depiction of the experimental design for treatment with KCl or caspofungin during *nda3*-mediated spindle checkpoint activation to activate MAPKs. Samples were collected at indicated time points for subsequent analyses including immunoblotting, co-IP and time-course analysis on SAC or APC/C activation.

(**B**) Immunoblot analysis of activation of MAPKs and Slp1^{Cdc20} protein levels. Samples with or without indicated treatments were blotted with anti-phospho p42/44 and anti-phospho p38 antibodies as indicative of phosphorylated Pmk1 (Pmk1-P) or phosphorylated Sty1 (Sty1-P), respectively. Slp1^{Cdc20} levels were detected with anti-Slp1 antibodies and anti-Cdc2 was used as loading control.

(**C**) Co-immunoprecipitation analysis of APC/C-MCC association upon environmental stress. Lid1-TAP was immunoprecipitated from *nda3-KM311*-arrested cells and associated Mad2-GFP, Mad3-GFP and Slp1^{Cdc20} were detected as in Figure 2A. Note that APC/C-MCC association was disrupted when 0.6 M KCl was present during cell culturing or during immunoprecipitation procedures.

(**D**) Time-course analyses of SAC activation and inactivation in *nda3-KM311 cdc13-GFP* strains with indicated genotypes after arrest at 18 ℃ and KCl or caspofungin treatments.

(**E, F**) Immunoblot analyses of Slp1^{Cdc20} abundance and time-course analyses of SAC activation and inactivation efficiency in phosphorylation- and ubiquitylation-deficient mutants upon SAC activation and environmental stress.

Figure 7. Summary of the mechanisms that how MAPKs negatively regulate APC/C activity in fission yeast.

(**A**) Schematic depiction of a possible dual mechanism that how activated MAPK signaling pathways are involved in delaying APC/C activation and SAC inactivation. Upon SAC and MAPK signaling activation, division of labor between the CIP and the SAP pathways enables the phosphorylation of Slp1^{Cdc20} and unidentified substrate(s) by Pmk1 and Sty1 respectively, which leads to lowered Slp1^{Cdc20} level and enhanced MCC affinity for APC/C.

(**B**) Summary of the alteration of Slp1^{Cdc20} protein levels and MCC-APC/C association strength under various conditions examined in this study. White wavy line indicates the absence of Apc15.