Downregulation of Semaphorin 4A in keratinocytes reflects the features of non-lesional psoriasis

Authors: Miki Kume [1], Hanako Koguchi-Yoshioka [1,2], Shuichi Nakai [1,3], Yutaka Matsumura [1], Atsushi Tanemura [1], Kazunori Yokoi [1], Shoichi Matsuda [1,3], Yuumi Nakamura [1.4], Naoya Otani [5], Mifue Taminato [5], Koichi Tomita [5,6], Tateki Kubo [5], Mari Wataya-Kaneda [1,2], Atsushi Kumanogoh [7], Manabu Fujimoto [1], Rei Watanabe [1,8]

Author Affiliations:

[1] Department of Dermatology, Course of Integrated Medicine, Graduate School of Medicine, Osaka University, Osaka, Japan

[2] Department of Neurocutaneous Medicine, Division of Health Sciences, Graduate School of Medicine, Osaka University, Osaka, Japan

[3] Research Department, Maruho Co, Ltd., Kyoto, Japan

[4] Cutaneous Allergy and Host Defense, Immunology Frontier Research Center (iFReC), Osaka University, Osaka, Japan

[5] Department of Plastic Surgery, Course of Organ Regulation Medicine, Graduate School of Medicine, Osaka University, Osaka, Japan

[6] Department of Plastic and Reconstructive Surgery, Kindai University, Osaka, Japan

[7] Department of Respiratory Medicine and Clinical Immunology, Course of Internal

Medicine, Graduate School of Medicine, Osaka University, Osaka, Japan

[8] Department of Medicine for Cutaneous Immunological Diseases, Course of Integrated

Medicine, Graduate School of Medicine, Osaka University, Osaka, Japan

Corresponding author:

Rei Watanabe, MD, PhD.

Department of Medicine for Cutaneous Immunological Diseases, Course of Integrated

Medicine, Graduate School of Medicine, Osaka University, Osaka, Japan

2-2 Yamadaoka, Suita, Osaka 565-0871, Japan

Phone number: +81-6-6879-3031

Fax number: +81-6-6879-3038

E-mail address: rwatanabe@derma.med.osaka-u.ac.jp

Competing interests

SN and SM are affiliated with Maruho Co. as employees, but have declared no

conflicts of interest related to this research. The remaining authors also declare no

conflicts of interest.

Funding statement: This study was supported by a Grant-in-Aid for JSPS Fellows

22KJ2071 (to MK), a Grant-in-Aid for Scientific Research (KAKENHI) 16K19705 (to

RW), and a collaborative research grant from Maruho Co (Osaka, Japan).

Abstract

Psoriasis is a multifactorial disorder mediated by IL-17-producing T cells,

involving immune cells and skin-constituting cells. Semaphorin 4A (Sema4A), an

immune semaphorin, is known to take part in T helper type 1/17 differentiation and

activation. However, Sema4A is also crucial for maintaining peripheral tissue

homeostasis and its involvement in skin remains unknown. Here, we revealed that while

Sema4A expression was pronounced in psoriatic blood lymphocytes and monocytes, it

was downregulated in the keratinocytes of both psoriatic lesions and non-lesions

compared to controls. Imiquimod application induced more severe dermatitis in

Sema4A knockout (KO) mice compared to wild-type (WT) mice. The naïve skin of

Sema4AKO mice showed increased T cell infiltration and IL-17A expression along with

thicker epidermis and distinct cytokeratin expression compared to WT mice, which are

hallmarks of psoriatic non-lesions. Analysis of bone marrow chimeric mice suggested

that Sema4A expression in keratinocytes plays a regulatory role in imiquimod-induced

dermatitis. The epidermis of psoriatic non-lesion and Sema4AKO mice demonstrated

mTOR complex 1 upregulation, and the application of mTOR inhibitors reversed the

skewed expression of cytokeratins in Sema4AKO mice. Conclusively, Sema4A-

mediated signaling cascades can be triggers for psoriasis and targets in the treatment

and prevention of psoriasis.

INTRODUCTION

While the infiltration of immune cells into skin plays a critical role in the development of psoriasis, as evidenced by interleukin (IL) -23/IL-17 axis (Fitch, Harper, Skorcheva, Kurtz, & Blauvelt, 2007; Hawkes, Yan, Chan, & Krueger, 2018; Kim & Krueger, 2017), recent studies have revealed that cells constructing skin structure, such as keratinocytes, fibroblasts, and endothelial cells, also play pivotal roles in the development (Heidenreich, Röcken, & Ghoreschi, 2009; Lowes, Russell, Martin, Towne, & Krueger, 2013; Zhang et al., 2023) and maintenance (Arasa et al., 2019; Francis et al., 2024; Q. Li et al., 2023; Ma et al., 2023; Tan et al., 2015; Zhu et al., 2020) of psoriasis. Among these cells, keratinocytes function as a barrier and produce cytokines, chemokines, and antimicrobial peptides against foreign stimuli, resulting in the activation of immune cells (Ni & Lai, 2020; Zhou, Chen, Cui, Shi, & Guo, 2022).

Semaphorins were initially identified as guidance cues in neural development (Kolodkin, Matthes, & Goodman, 1993; Pasterkamp & Kolodkin, 2003) but are now regarded to play crucial roles in other physiological processes including angiogenesis (Iragavarapu-Charyulu, Wojcikiewicz, & Urdaneta, 2020; Serini, Maione, Giraudo, & Bussolino, 2009), tumor microenvironment (Hui, Tam, Jiao, & Ong, 2019; Jiang et al., 2022; Nakayama, Kusumoto, Nakahara, Fujiwara, & Higashiyama, 2018; Rajabinejad

et al., 2020), and immune systems (Garcia, 2019; Kanth, Gairhe, & Torabi-Parizi, 2021; M. Naito & Kumanogoh, 2023). Semaphorin 4A (Sema4A), one of the immune semaphorins, plays both pathogenic and therapeutic roles in autoimmune diseases (Cavalcanti et al., 2020; He et al., 2023), allergic diseases (Maeda et al., 2019), and cancer (Iyer & Chapoval, 2018; Liu et al., 2018; Y. Naito et al., 2023; Pan, Wang, & Ye, 2016). While Sema4A expression on T cells is essential for T helper type 1 differentiation in the murine *Propionibacterium acnes*-induced inflammation model and delayed-type hypersensitivity model (Kumanogoh et al., 2005), Sema4A amplifies only T helper type 17 (Th17)-mediated inflammation in the effector phase of murine experimental autoimmune encephalomyelitis (Koda et al., 2020). Accordingly, in multiple sclerosis, high serum Sema4A levels correlate with the elevated serum IL-17A, earlier disease onset, and increased disease severity (Koda et al., 2020; Nakatsuji et al., 2012). In anti-tumor immunity, research involving human samples and murine models suggests that Sema4A expressed in cancer cells and regulatory T cells promotes tumor progression (Delgoffe et al., 2013; Liu et al., 2018; Pan et al., 2016), while other reports reveal that Sema4A in cancer cells and dendritic cells bolsters anti-tumor immunity by enhancing CD8 T cell activity (Y. Naito et al., 2023; Suga et al., 2021). In addition to these roles in immune reactions, mice with a point mutation in Sema4A develop retinal

degeneration, suggesting that Sema4A is also crucial for peripheral tissue homeostasis

(Nojima et al., 2013). These reports suggest that the role of Sema4A can differ based on

the disease, phase, and involved cells.

Herein, we investigated the roles of Sema4A in the pathogenesis of psoriasis by

analyzing skin and blood specimens from psoriatic subjects and using a murine model.

RESULTS

Epidermal Sema4A expression is downregulated in psoriasis

The analysis of previously-published single-cell RNA-sequencing data from

control (Ctl) and psoriatic lesional (L) skin specimens (Kim et al., 2023) revealed

detectable expression of *SEMA4A* in keratinocytes, dendritic cells, and macrophages in

both Ctl and L. *SEMA4A* expression was low in neural crest-like cells, fibroblasts, CD4

T cells, CD8 T cells, NK cells, and plasma cells, making the comparison of expression

levels impractical (Figure 1A and B; Figure 1-figure supplement 1A-C). Dendritic cells

and macrophages showed comparable *SEMA4A* expression levels between Ctl and L

(Figure 1C). The adjusted *p*-value (*padj*) for *SEMA4A* in keratinocytes between Ctl and

L was 2.83×10^{-39}, indicating a statistically significant difference despite not being

visually prominent in the volcano plot, which shows comprehensive differential gene

expression in keratinocytes (Figure 1C; Figure 1-figure supplement 1D).

Immunohistochemistry of Ctl and psoriasis demonstrated Sema4A expression

in keratinocytes (Figure 1D). The staining intensity of Sema4A in epidermis was

significantly lower in both non-lesions (NL) and L than in Ctl (Figure 1E). Relative

mRNA expression of *SEMA4A* was also decreased in the epidermis of NL and L

compared to Ctl, while it remained comparable in the dermis (Figure 1F). In contrast,

the proportions of Sema4A-positive cells were significantly higher in blood CD4 and

CD8 T cells, and monocytes in psoriasis compared to Ctl (Figure 1G; Figure 1-figure

supplement 2). Serum Sema4A levels, measured by enzyme-linked immunosorbent

assay (ELISA), were comparable between Ctl and psoriasis (Figure 1H). These findings

demonstrate that the expression profile of Sema4A in psoriasis varies across cell types.

Psoriasis-like dermatitis is augmented in Sema4AKO mice

When psoriasis-like dermatitis was induced in wild-type (WT) mice and

Sema4A knockout (KO) mice by imiquimod application on ears (Figure 2A), ear

swelling on day 4 was more pronounced in Sema4AKO mice (Figure 2B) with

upregulated *Il17a* gene expression (Figure 2C). Flow cytometry analysis of cells

isolated from the ears revealed increased proportions of $V\gamma2^+$ T cells, $V\gamma2^-V\gamma3^-$ double-

negative (DN) $\gamma\delta$ T cells, and IL-17A-producing cells of those fractions in Sema4AKO

epidermis (Figure 2D and E; Figure 2-figure supplement 1). In Sema4AKO dermis,

there was also an increase in the proportions of $V\gamma2^+$ T cells and IL-17A-producing

$V\gamma2^+$ T cells (Figure 2D and E). These results suggest that Sema4A deficiency in mice

accelerates psoriatic profile. IL-17A-producing T cells in skin-draining lymph nodes

(dLN) remained comparable between WT mice and Sema4AKO mice (Figure 2F).

Though the imiquimod model is well-established and valuable murine psoriatic model

(van der Fits et al., 2009), the vehicle of imiquimod cream can activate skin

inflammation that is independent of toll-like receptor 7, such as inflammasome

activation, keratinocyte death and interleukin-1 production (Walter et al., 2013). This

suggests that the imiquimod model involves complex pathway. Therefore, we

subsequently induced IL-23-mediated psoriasis-like dermatitis (Figure2-figure

supplement 2A), a much simpler murine psoriatic model, because IL-23 is thought to

play a central role in psoriasis pathogenesis (Krueger et al., 2007; Lee et al., 2004).

Although ear swelling on day 4 was comparable between WT mice and Sema4AKO

mice (Figure2-figure supplement 2B), the epidermis, but not the dermis, was

significantly thicker in Sema4AKO mice compared to WT mice (Figure2-figure

supplement 2C). We found that the proportion of CD4 T cells among T cells was

significantly higher in Sema4A KO mice compared to WT mice, while the proportion of

Vγ2 and DNγδ T cells among T cells was comparable between them (Figure 2-figure

supplement 2D). On the other hand, focusing on IL-17A-producing cells, the proportion

of IL-17A-producing Vγ2 and DNγδ T cells in CD3 fraction in the epidermis was

significantly higher in Sema4A KO mice, consistent with the results from imiquimod-

induced psoriasis-like dermatitis. (Figure 2-figure supplement 2E).

Sema4A in keratinocytes may play a role in preventing murine psoriasis-like dermatitis

To investigate the cells responsible for the augmented ear swelling in Sema4AKO mice, bone marrow chimeric mice were next analyzed (Figure 3A). Since it has already been reported that bone marrow cells contain keratinocyte stem cells (Harris et al., 2004; Wu, Zhao, & Tredget, 2010), we confirmed that epidermis of mice deficient in non-hematopoietic Sema4A (WT→KO) showed no obvious detection of *Sema4a*, thereby ruling out the impact of donor-derived keratinocyte stem cells infiltrating the host epidermis (Figure 3-figure supplement 1A). WT→KO mice displayed more pronounced ear swelling than mice with intact Sema4A expression (WT→WT) following imiquimod application (Figure 3B). Similarly, mice with a systemic deficiency of Sema4A (KO→KO) showed severe ear swelling compared to mice deficient in hematopoietic Sema4A (KO→WT) (Figure 3B). Ear swelling was comparable between WT→WT mice and KO→WT mice (Figure 3B). Flow cytometry analysis revealed increased infiltration of IL-17A-producing DNγδ T cells in the epidermis, as well as Vγ2$^+$ T cells and IL-17A-producing Vγ2$^+$ T cells in the dermis, in WT→KO mice compared to WT→WT mice (Figure 3C; Figure 3-figure supplement

1B). These findings suggest that non-hematopoietic cells, possibly keratinocytes, are

primarily responsible for the increased imiquimod-induced Sema4AKO mice ear

swelling.

Sema4AKO epidermis is thicker than WT epidermis with increased γδ T17

infiltration

Even without imiquimod application, Sema4AKO ears turned out to be slightly

but significantly thicker than WT ears on week 8 while their appearance remained

normal (Figure 4A). While epidermal thickness of back skin was comparable at birth

(Figure 4B), on week 8, epidermis of Sema4AKO back and ear skin was notably thicker

than that of WT mice (Figure 4B), suggesting that acanthosis in Sema4AKO mice is

accentuated post-birth. Dermal thickness remained comparable between WT mice and

Sema4AKO mice at both times (Figure 4B). The epidermis of WT ear at week 8 showed

significantly higher *Sema4a* mRNA expression compared to the dermis (Figure 4C).

Based on these observations, Sema4A appears to play a more pronounced role in

epidermis than in dermis.

Sema4AKO epidermis exhibited increased expression of *Ccl20, Tnfα*, and

Il17a and a trend of upregulation of *S100a8* compared to WT epidermis (Figure 4-figure

supplement 1A). These differences were not observed in dermis (Figure 4-figure

supplement 1A). Flow cytometry analysis revealed increased infiltration of $\gamma\delta$ T cells in

Sema4AKO ear (Figure 4-figure supplement 1B). These cells predominantly expressed

resident memory T cell (T_{RM})-characteristic molecules, CD69 and CD103 (Figure 4-

figure supplement 1B). Sema4AKO skin also had a higher number of T_{RM} in both CD4

and CD8 T cells (Figure 4-figure supplement 1B). The percentages of V$\gamma2^+$ T cells,

DN$\gamma\delta$ T cells, and CD8 T cells in epidermis were higher in Sema4AKO mice than in

WT mice, which was not the case in dermis (Figure 4-figure supplement 2A). The

proportion of V$\gamma3^+$ dendritic epidermal T cells was comparable between WT mice and

Sema4AKO mice (Figure 4-figure supplement 2A). Epidermal V$\gamma2^+$ T cells and DN$\gamma\delta$ T

cells in Sema4AKO mice showed higher IL-17A-producing capability (Figure 4D),

while IFNγ and IL-4 production was comparable between WT mice and Sema4AKO

mice (Figure 4-figure supplement 2B and C). Conversely, the frequency of IL-17A-

producing T cells from dLN was comparable (Figure 4E). The production of IL-17A

and IFNγ from splenic T cells under T17-polarizing conditions remained consistent

between WT mice and Sema4AKO mice (Figure 4-figure supplement 3).

Taken together, it is suggested that T17 cells are specifically upregulated in

epidermis, indicating that the epidermal microenvironment plays a pivotal role in

facilitating the increased T cell infiltration observed in naïve Sema4AKO mice.

Sema4AKO skin shares features with human psoriatic NL

Previous literatures have identified certain features common to psoriatic L and

NL, such as thickened epidermis (Figure 5-figure supplement 1) (Gallais Sérézal et al.,

2019), CCL20 upregulation (Gallais Sérézal et al., 2019), and accumulation of IL-17A-

producing T cells (Vo et al., 2019), which were detected in Sema4AKO mice.

Gene expression analysis using public RNA-sequencing data (Tsoi et al., 2019)

with RaNA-seq (Prieto & Barrios, 2019) showed upregulation of keratinization and

antimicrobial peptide genes in NL compared to Ctl (Figure 5A). Gene Ontology analysis

highlighted an upregulation in biological processes, predominantly in peptide cross-

linking involved in epidermis formation and keratinocyte differentiation, with a

secondary increase in the defense response to viruses in NL (Figure 5B). While the

expression of *Keratin (KRT) 10* was comparable between NL and Ctl, upregulation in

KRT5, *KRT14*, and *KRT16* was observed in NL (Figure 5C).

In the murine model, relative expression levels of *Krt10*, *Krt14*, *Krt16*, and

Filaggrin were elevated in Sema4AKO epidermis. (Figure 5D). Immunofluorescence

analysis showed that Sema4AKO epidermis had a higher density of keratinocytes

positive for Krt5, Krt10, Krt14, and Krt16 compared to WT epidermis (Figure 5E). This

upregulation was not observed in back skin at birth (Figure 5-figure supplement 2).

Comparable transepidermal water loss between WT mice and Sema4AKO mice

indicated preserved skin barrier function in Sema4AKO mice (Figure 5F).

Based on these results, it is implied that the epidermis of human psoriatic NL

and Sema4AKO mice exhibit shared pathways, potentially leading to an acanthotic state.

Combined with the observed acanthosis and increased T17 infiltration in Sema4AKO

mice, Sema4AKO skin is regarded to demonstrate the features characteristic of human

psoriatic NL.

mTOR signaling is upregulated in the epidermis of psoriatic NL and Sema4AKO mice

Previous reports have shown that mTOR pathway plays a critical role in

maintaining epidermal homeostasis, as evidenced by mice with keratinocyte-specific

deficiencies in Mtor, Raptor, or Rictor, which exhibit a hypoplastic epidermis with

impaired differentiation and barrier formation (Asrani et al., 2017; Ding et al., 2016;

Ding et al., 2020). We thus investigated mTOR pathway in both human and murine

epidermis.

Immunohistochemical analyses highlighted the increase in phospho-S6 (p-S6),

indicating the upregulation of mTOR complex (C) 1 signaling, in the epidermal upper

layers of both L and NL compared to Ctl (Figure 6A). The activation of mTORC2 in the

epidermis, inferred from phospho-Akt (p-Akt), was scarcely detectable in L, NL, and

Ctl (Figure 6A). In the epidermis of WT mice and Sema4AKO mice, the upregulation of

both mTORC1 and mTORC2 signaling was observed in Sema4AKO epidermis by

immunohistochemistry (Figure 6B). Western blot analysis showed upregulated

mTORC1 signaling in Sema4AKO epidermis compared to WT epidermis (Figure 6C).

The enhancement of mTORC1 and mTORC2 signaling became obvious in the

Sema4AKO epidermis after developing psoriatic dermatitis by imiquimod application

(Figure 6D).

Inhibition of mTOR signaling modulates cytokeratin expression in Sema4AKO mice

To investigate the contribution of mTORC1 and mTORC2 signaling in the

development of psoriatic features in the Sema4AKO epidermis, mTORC1 inhibitor

rapamycin and mTORC2 inhibitor JR-AB2-011 were intraperitoneally applied to

Sema4AKO mice for 14 days. Although epidermal thickness remained unchanged by

the inhibitors (Figure 7A and B), relative gene expression of *Krt5* was significantly

upregulated and that of *Krt16* was significantly downregulated after rapamycin

application (Figure 7C). While the upregulation of *Il17a* in Sema4AKO epidermis was

not clearly modified by rapamycin (Figure 7C), immunofluorescence revealed the

decrease in the number of CD3 T cells in Sema4AKO epidermis by rapamycin (Figure

7D). We additionally conducted topical application of rapamycin gel and vehicle gel on

the left and right ears of Sema4AKO mice, respectively. Although there were no

detectable changes in epidermal thickness and epidermal T cell counts, the upregulation

of *Krt5* and downregulation of *Krt16* was observed again (Figure 7-figure supplement

1). Conversely, the application of JR-AB2-011 resulted in decreased expression of *Krt5*,

Krt10, and *Krt14* with a trend towards increased *Krt16* expression (Figure 7E). JR-

AB2-011 did not influence the number of infiltrating T cells in the epidermis (Figure

7F). Next, we investigated whether intraperitoneal rapamycin treatment effectively

downregulates inflammation in the IMQ-induced murine model of psoriasis in

Sema4AKO mice (Figure 7-figure supplement 2A). Rapamycin significantly reduced

epidermal thickness compared to vehicle treatment (Figure 7-figure supplement 2B).

Additionally, rapamycin treatment downregulated the expression of *Krt10*, *Krt14*, and

Krt16 (Figure 7-figure supplement 2C). While the upregulation of *Il17a* in the

Sema4AKO epidermis in IMQ model was not clearly modified by rapamycin (Figure 7-

figure supplement 2C), immunofluorescence revealed a decrease in the number of CD3

T cells in Sema4AKO epidermis by rapamycin (Figure 7-figure supplement 2D). In the

naive states, mTORC1 primarily regulates keratinocyte proliferation, whereas mTORC2

mainly involved in the keratinocyte differentiation through Sema4A-related signaling

pathways. Conversely, in the psoriatic dermatitis state, rapamycin downregulated both

keratinocyte differentiation and proliferation markers. The observed similarities in *Il17a*

expression following treatment with rapamycin and JR-AB2-011, regardless of

additional IMQ treatment, suggest that *Il17a* production is not significantly dependent

on Sema4A-related mTOR signaling.

DISCUSSION

Recent studies have highlighted the significance of IL-17A-producing T_{RM} in psoriatic NL (Cheuk et al., 2014; Gallais Sérézal et al., 2018; Vo et al., 2019; Vu, Koguchi-Yoshioka, & Watanabe, 2021). This condition is also characterized by epidermal thickening with elevated CCL20 expression (Gallais Sérézal et al., 2019) and a distinct cytokeratin expression pattern, marked by increased levels of Krt5, Krt14, and Krt16 (Tsoi et al., 2019) (Figure 5C). Here, we identified that murine Sema4A deficiency could induce these features of psoriatic NL.

Sema4A expression in epidermis, but not in dermis, was diminished in psoriatic L and NL compared to Ctl. While Sema4A expression on blood immune cells was upregulated in psoriasis, its serum levels were comparable between Ctl and psoriasis. Despite the reported involvement of increased Sema4A expression on immune cells in the pathogenesis of multiple sclerosis (Koda et al., 2020; Nakatsuji et al., 2012), our results suggest that the diminished expression of Sema4A in skin-constructing cells plays a more prominent role in the pathogenesis of psoriasis than its increased expression on immune cells.

In our murine model, regardless of imiquimod application, IL-17A-producing T cells were increased in Sema4AKO skin although their frequency was comparable in

dLN of WT mice and Sema4AKO mice. Additionally, the potential for *in vitro* T17

differentiation did not differ between T cells from WT mice and Sema4AKO mice.

These findings suggest that the absence of Sema4A in the skin microenvironment plays

a crucial role in the localized upregulation of IL-17A-producing T cells in Sema4AKO

mice.

It is well-documented that upregulated mTORC1 signaling promotes the

pathogenesis of psoriasis (Buerger, 2018; Karagianni, Pavlidis, Malakou, Piperi, &

Papadavid, 2022; Ruf, Andreoli, Itin, Pluschke, & Schmid, 2014), and that rapamycin

can partially ameliorate the disease activity in psoriatic subjects and murine models

(Bürger et al., 2017; Gao & Si, 2018; Reitamo et al., 2001). Our results using

Sema4AKO mice revealed that inhibition of mTORC1 leads to the downregulation of

Krt16, and inhibition of mTORC2 leads to the downregulation of undifferentiated

keratinocytes, while the Sema4AKO epidermal thickening and the upregulated IL-17A

signaling are not reversed by mTOR blockade in the naïve state. It is plausible that the

downregulation of Sema4A can lead to the upregulation of mTORC1 and mTORC2

signaling in keratinocytes, and the augmented signaling leads to the psoriatic profile of

proliferation and differentiation of keratinocytes, which is part of the psoriatic NL

disposition.

This study has limitations. Sema4A expression in skin cells other than

keratinocytes was not thoroughly investigated. However, since single-cell RNA-

sequencing has shown that Sema4A is predominantly expressed in keratinocytes,

dendritic cells, and macrophages, with a notable reduction in keratinocytes in psoriasis,

we infer that keratinocytes are the primary cells responsible for the psoriatic features

resulting from Sema4A downregulation. We were not able to determine whether

Sema4A functions as a ligand or a receptor in epidermis (Ito & Kumanogoh, 2016;

Kumanogoh et al., 2002; Lu et al., 2018; Sun et al., 2017) in this study, either. We were

not able to reveal how Sema4A expression is regulated. Although we showed that

downregulation of Sema4A is related to the abnormal cytokeratin expression observed

in psoriasis, we could not determine the relationships between Sema4A expression and

the essential molecules upregulated in psoriatic keratinocytes. While both mTORC1 and

mTORC2 signals are upregulated in Sema4AKO epidermis, we were not able to

confirm mTORC2 signaling from human skin due to technical limitation and sample

limitation, while the results from augmented mTORC2 signal in Sema4AKO mice and

the normalization of *Krt5* and *Krt14* by mTORC2 inhibitor imply the involvement of

mTORC2 signal in psoriatic epidermis. The role of Sema4A other than mTOR signaling,

which may be involved in the regulation of T17 induction in skin, is not discussed,

either. These limitations should be overcome in the near future.

In summary, epidermal Sema4A downregulation can reflect the psoriatic non-

lesional features. Thus, targeting the downregulated Sema4A and upregulated mTOR

signaling in psoriatic epidermis can be a promising strategy for psoriasis therapy and

modification of psoriatic diathesis in NL for the prevention of disease development.

MATERIALS AND METHODS

Human sample collection

Psoriatic L and NL skin specimens were acquired from 17 psoriasis patients. Ctl specimens were obtained from 19 subjects who underwent tumor resection or reconstructive surgery. For epidermal-dermal separation, specimens were incubated overnight at 4℃ with 2.5 mg/mL dispase Ⅱ (Wako, Osaka, Japan) in IMDM (Wako). Blood samples were collected from 73 psoriasis and 33 Ctl. In addition to ELISA (Nakatsuji et al., 2012), peripheral blood mononuclear cells were isolated using Ficoll-Paque PLUS density gradient media (Cytiva, Tokyo, Japan). Patient details are provided in Supplemental Table 1.

Mice

C57BL/6J WT mice were procured from CLEA Japan (Tokyo, Japan). Sema4AKO mice with C57BL/6J background were generated as previously described (Kumanogoh et al., 2005). We examined female mice in order to reduce the result variation. Neonatal mice and female mice aged 8 to 12 weeks, maintained under specific pathogen-free conditions, were used.

To develop psoriasis-like dermatitis, 10 mg of 5％ imiquimod cream (Mochida,

Tokyo, Japan) was applied to both ears for 4 consecutive days. To induce IL-23-

mediated psoriasis-like dermatitis, 20 μl of phosphate-buffered saline containing 500 ng

of recombinant mouse IL-23 (BioLegend, San Diego, CA) was injected intradermally

into both ears of anesthetized mice using a 29-gauge needle for 4 consecutive days. Ear

thickness was measured using a thickness gauge (Peacock, Tokyo, Japan).

For bone marrow reconstitution, 2×10^5 bone marrow cells obtained from the

indicated strains were intravenously transferred to recipient mice that had received a

single 10 Gy irradiation. The reconstituted mice were subjected to the experiments in 8

weeks.

In the specified experiments, skin specimens were separated into epidermis and

dermis by 5 mg/mL dispase (Wako) in IMDM for 30 minutes at 37℃.

Transepidermal water loss measurements were performed on the back skin of

mice at week 8 using a Tewameter (Courage and Khazaka Electronic GmbH, Cologne,

Germany), according to the manufacturer's instructions.

Immunofluorescence and immunohistochemistry

Specimens were fixed in 4% paraformaldehyde phosphate buffer solution

(Wako), embedded in paraffin, and sliced into 3-μm thickness on glass slides. After

deparaffinization and rehydration, antigen retrieval was performed using citrate buffer

(pH 6.0, Nacalai tesque, Kyoto, Japan) or TE buffer (pH 9.0, Nacalai tesque).

For immunohistochemistry, samples were incubated with 3% H_2O_2 (Wako) for

5 minutes. After blocking (Agilent, Santa Clara, CA), the specimens were incubated

with the indicated primary antibodies under specified conditions (Supplemental Table 2).

The specimens were applied with the Dako REAL EnVision Detection System,

Peroxidase/DAB, Rabbit/Mouse, HRP kit (Agilent) and counterstained with

hematoxylin (Wako). For immunofluorescence, the blocked specimens were incubated

with the indicated primary antibodies followed by the secondary antibodies (Table E4).

Mounting medium with DAPI (Vector Laboratories, Burlingame, CA) was used. Slides

were observed using a fluorescence microscope (BZ-X700, Keyence, Osaka, Japan).

The staining intensity was measured using Fiji software (ImageJ, National Institutes of

Health, Bethesda, MD) over lengths of 200 μm in human samples and 100 μm in murine

samples. This measurement was taken from five areas of each specimen, and the

average score is presented. The average thickness of epidermis and dermis from 10

spots is presented. The number of epidermal cells positive for the indicated cytokeratins

was counted per 100 μm width, with the average number from 5 areas being presented.

Additionally, the numbers of CD3-positive cells in murine ears were counted across 10

fields of 400 μm width, with the total sum being presented.

Quantitative reverse transcription polymerase chain reaction (qRT-PCR)

Total RNA was extracted from homogenized skin tissue using Direct-zol RNA

MiniPrep Kit (ZYMO Research, Irvine, CA). cDNA was synthesized by High-Capacity

RNA-to-cDNA[TM] Kit (Thermo Fisher Scientific, Waltham, MA). qPCR was performed

using TB Green Premix Ex Taq[TM] II (Takara-bio, Shiga, Japna) on a ViiA 7 Real-Time

PCR System (Thermo Fisher Scientific). The primers are listed in Supplemental Table 3.

All samples were run in triplicate, and the median CT value was calculated. Relative

gene expression levels were normalized to the housekeeping gene *GAPDH* using the

ΔΔCT technique.

Flow cytometry analysis

Single-cell suspensions from murine skin-dLN and spleen were prepared by

grinding, filtering, and lysing red blood cells (BioLegend). Skin specimens were minced

and digested with 3 mg/mL collagenase type III (Worthington Biochemical Corporation,

Lakewood, NJ) in RPMI 1640 medium (Wako) at 37℃ for 10 minutes for epidermis,

and 30 minutes for dermis or whole skin. Cells were surface-stained with directly

conjugated monoclonal antibodies (Supplemental Table 4). Dead cells were identified

using LIVE/DEAD^{TM} Fixable Dead Cell Stain Kit (Thermo Fisher Scientific). To

evaluate cytokine production, cells were stimulated with Phorbol 12-Myristate 13-

Acetate (PMA; 50 ng/mL, Wako) and ionomycin (1000 ng/mL, Wako), plus BD

Golgiplug^{TM} (BD Biosciences, San Jose, CA) for 4 hours before surface staining.

Fixation, permeabilization, and intracellular cytokine staining were performed using BD

Cytofix/Cytoperm^{TM} Fixation/Permeabilization Kit (BD Biosciences) according to the

manufacturer's protocol. In specified experiments, the numbers of each cell subset per

ear were estimated using CountBright^{TM} Absolute Counting Beads (Thermo Fisher

Scientific). Sample analysis was conducted using BD FACSCanto II (BD Biosciences),

and data were analyzed with Kaluza software (Beckman Coulter, Brea, CA).

***In vitro* Th17 differentiation**

Murine splenic T cells were isolated using Pan T Cell Isolation Kit II (Miltenyi

Biotec, Bergisch Gladbach, Germany). Two hundred thousand T cells per well were

cultured in 96-well plates in the presence of T Cell Activation/Expansion Kit (Miltenyi

Biotec) for 2 weeks. The medium was supplemented twice per week with the following

recombinant cytokines: mouse recombinant IL-6 (20 ng/mL), IL-1β (20 ng/mL), and

IL-23 (40 ng/mL) for the IL-23 dependent Th17 cell condition; IL-6 (20 ng/mL) and

TGFβ1 (3 ng/mL) for the IL-23 independent Th17 cell condition; or IL-23

(40 ng/mL) for the IL-23 only Th17 cell condition. The cytokines listed were

purchased from BioLegend (Supplemental Table 5). Afterward, the cultured cells were

processed for flow cytometry analysis.

Western blotting

Murine epidermis was lysed using RIPA buffer (Wako) containing

phosphatase-and protease-inhibitor cocktail (Nacalai tesque). Protein lysates were

separated by 10% SuperSepTM Ace (Wako), transferred onto polyvinylidene difluoride

membranes (0.45 μm, Merck, Darmstadt, Germany) by Trans-Blot Turbo Transfer

System (Bio-Rad, Hercules, CA). The membrane was blocked with 5% bovine serum

albumin and subjected to immunoblotting targeting the indicated proteins overnight 4℃,

followed by the application of HRP-conjugated secondary antibody for one hour at

room temperature (Supplemental Table 2). WB stripping solution (Nacalai tesque) was

used to remove the antibodies for further evaluation.

Inhibition of mTOR

Rapamycin (4 mg/kg, Sanxin Chempharma, Hebei, China), JR-AB2-011 (400

µg/kg, MedChemExpress, Monmouth Junction, NJ) and vehicle were applied

intraperitoneally to mice once daily for 14 consecutive days. Rapamycin and JR-AB2-

011 were dissolved in 100% DMSO and diluted with 40% Polyethylene Glycol 300

(Wako), 5% Tween-80 (Sigma-Aldrich, St. Louis, MO) and 45% saline in sequence. For

topical application, 0.2% rapamycin gel and vehicle gel were prepared by the

pharmaceutical department as previously described (Wataya-Kaneda et al., 2017).

Rapamycin gel was applied to the left ear, and vehicle gel to the right ear (10mg/ear) of

Sema4AKO, once daily for 14 consecutive days. To analyze the preventive

effectiveness of rapamycin in an IMQ-induced murine model of psoriatic dermatitis,

Sema4AKO mice were administered either vehicle or rapamycin intraperitoneally from

Day 0 to Day 17, and IMQ was topically applied to both ears for 4 days starting on Day

14. Then, on Day 18, ears were collected for further analysis.

Data processing of single-cell RNA-sequencing and bulk RNA-sequencing

The raw count matrix data from the previously reported single-cell RNA-

sequencing data from GSE220116 (Kim et al., 2023) were imported into Scanpy (1.9.6)

using Python for further analyses. For each sample, cells and genes meeting the

following criteria were excluded: cells expressing over 200 genes (sc.pp.filter_cells),

cells with a high proportion of mitochondrial genes (> 5%), and genes expressed in

fewer than 3 cells (sc.pp.filter_genes). Counts were normalized using

sc.pp.normalize_per_cell, logarithmized (sc.pp.log1p), and scaled (sc.pp.scale). Highly

variable genes were selected using sc.pp.filter_genes_dispersion with the options

min_mean = 0.0125, max_mean = 2.5, and min_disp = 0.7.

Principal Component analysis was conducted with sc.pp.pca, selecting the 1st

to 50th Principal Components for embedding and clustering. Neighbors were calculated

with batch effect correction by BBKNN (Polański et al., 2020). Embedding was

performed by sc.tl.umap, and cells were clustered using sc.tl.leiden. Some cells were

further subclassified in a similar manner. The data was integrated into an h5ad file,

which can be visualized in Cellxgene VIP (K. Li et al., 2022). We then performed

differential analysis between two groups of cells to identify differential expressed genes

using Welch's t-test. Multiple comparisons were controlled using the Benjamini-

Hochberg procedure, with the false discovery rate set at 0.05 and significance defined as

$padj < 0.05$.

Bulk RNA-sequencing data from GSE121212 (Tsoi et al., 2019) were re-

analyzed using RaNAseq (Prieto & Barrios, 2019) for differential expression in Ctl

versus psoriatic NL. Our analysis included normalization, differential gene expression

(defining significance as $padj < 0.05$), and focused on Gene Ontology biological

process analysis. The gene expression was calculated with the transcripts per million

values.

Statistical analysis

GraphPad Prism 10 software (GraphPad Software, La Jolla, CA) was used for

all statistical analyses except for RNA-sequencing. Mann-Whitney test was used for

two-group comparisons, while Kruskal-Wallis test followed by Dunn's multiple

comparisons test was applied for comparison among three or more groups. Statistical

significance was defined as $p < 0.05$ (*), $p < 0.01$ (**), and $p < 0.001$ (***). In the

analysis of RNA-sequencing data, statistical significance was defined as $padj < 0.01$

(**), and $padj < 0.001$ (***).

Study approval

All experiments involving human specimens were in accordance with the

Declaration of Helsinki and were approved by the Institutional Review Board in Osaka

University Hospital (20158-6). Written informed consent was obtained from all

participants. All murine experiments were approved by Osaka University Animal

Experiment Committee (J007591-013) and all procedures were conducted in

compliance with the Guidelines for Animal Experimentation established by Japanese

Association for Laboratory Animal Science.

Data availability

The single-cell RNA-sequencing datasets generated by Kim et al. (Kim et al.,

2023) and Tsoi et al. (Tsoi et al., 2019) used in this study are available in the NCBI

Gene Expression Omnibus under accession codes GSE220116 and GSE121212,

respectively. Values for all data points in graphs are reported in the Source data.

Acknowledgments

We express our gratitude to all patients for their participation in this study.

REFERENCES

Arasa, J., Terencio, M. C., Andrés, R. M., Marín-Castejón, A., Valcuende-Cavero, F.,

Payá, M., & Montesinos, M. C. (2019). Defective Induction of COX-2 Expression

by Psoriatic Fibroblasts Promotes Pro-inflammatory Activation of Macrophages.

Front Immunol, 10, 536. doi:10.3389/fimmu.2019.00536

Asrani, K., Sood, A., Torres, A., Georgess, D., Phatak, P., Kaur, H., . . . Lotan, T.

L. (2017). mTORC1 loss impairs epidermal adhesion via TGF-beta/Rho kinase

activation. *J Clin Invest, 127*(11), 4001-4017. doi:10.1172/JCI92893

Bürger, C., Shirsath, N., Lang, V., Diehl, S., Kaufmann, R., Weigert, A., . . . Wolf,

P. (2017). Blocking mTOR Signalling with Rapamycin Ameliorates Imiquimod-

induced Psoriasis in Mice. *Acta Derm Venereol, 97*(9), 1087-1094.

doi:10.2340/00015555-2724

Buerger, C. (2018). Epidermal mTORC1 Signaling Contributes to the Pathogenesis of

Psoriasis and Could Serve as a Therapeutic Target. *Front Immunol, 9*, 2786.

doi:10.3389/fimmu.2018.02786

Carvalheiro, T., Affandi, A. J., Malvar-Fernández, B., Dullemond, I., Cossu, M.,

Ottria, A., . . . García, S. (2019). Induction of Inflammation and Fibrosis

by Semaphorin 4A in Systemic Sclerosis. *Arthritis Rheumatol, 71*(10), 1711-

1722. doi:10.1002/art.40915

Cavalcanti, C. A. J., Germoglio, V., de Azevêdo Silva, J., Glesse, N., Vianna, P.,

Cechim, G., . . . Sandrin-Garcia, P. (2020). T-cell specific upregulation of

Sema4A as risk factor for autoimmunity in systemic lupus erythematosus and

rheumatoid arthritis. *Autoimmunity, 53*(2), 65-70.

doi:10.1080/08916934.2019.1704273

Cheuk, S., Wikén, M., Blomqvist, L., Nylén, S., Talme, T., Ståhle, M., & Eidsmo, L.

(2014). Epidermal Th22 and Tc17 cells form a localized disease memory in

clinically healed psoriasis. *J Immunol, 192*(7), 3111-3120.

doi:10.4049/jimmunol.1302313

Delgoffe, G. M., Woo, S. R., Turnis, M. E., Gravano, D. M., Guy, C., Overacre, A.

E., . . . Vignali, D. A. (2013). Stability and function of regulatory T cells

is maintained by a neuropilin-1-semaphorin-4a axis. *Nature, 501*(7466), 252-

256. doi:10.1038/nature12428

Ding, X., Bloch, W., Iden, S., Rüegg, M. A., Hall, M. N., Leptin, M., . . . Eming, S.

A. (2016). mTORC1 and mTORC2 regulate skin morphogenesis and epidermal

barrier formation. *Nat Commun, 7*, 13226. doi:10.1038/ncomms13226

Ding, X., Willenborg, S., Bloch, W., Wickström, S. A., Wagle, P., Brodesser, S., . . .
Eming, S. A. (2020). Epidermal mammalian target of rapamycin complex 2
controls lipid synthesis and filaggrin processing in epidermal barrier
formation. *J Allergy Clin Immunol, 145*(1), 283-300.e288.
doi:10.1016/j.jaci.2019.07.033
Fitch, E., Harper, E., Skorcheva, I., Kurtz, S. E., & Blauvelt, A. (2007).
Pathophysiology of psoriasis: recent advances on IL-23 and Th17 cytokines.
Curr Rheumatol Rep, 9(6), 461-467. doi:10.1007/s11926-007-0075-1
Francis, L., McCluskey, D., Ganier, C., Jiang, T., Du-Harpur, X., Gabriel, J., . . .
Mahil, S. K. (2024). Single-cell analysis of psoriasis resolution
demonstrates an inflammatory fibroblast state targeted by IL-23 blockade. *Nat*
Commun, 15(1), 913. doi:10.1038/s41467-024-44994-w
Gallais Sérézal, I., Classon, C., Cheuk, S., Barrientos-Somarribas, M., Wadman, E.,
Martini, E., . . . Eidsmo, L. (2018). Resident T Cells in Resolved Psoriasis
Steer Tissue Responses that Stratify Clinical Outcome. *J Invest Dermatol,*
138(8), 1754-1763. doi:10.1016/j.jid.2018.02.030
Gallais Sérézal, I., Hoffer, E., Ignatov, B., Martini, E., Zitti, B., Ehrström, M., &
Eidsmo, L. (2019). A skewed pool of resident T cells triggers psoriasis-
associated tissue responses in never-lesional skin from patients with
psoriasis. *J Allergy Clin Immunol, 143*(4), 1444-1454.
doi:10.1016/j.jaci.2018.08.048
Gao, M., & Si, X. (2018). Rapamycin ameliorates psoriasis by regulating the
expression and methylation levels of tropomyosin via ERK1/2 and mTOR pathways
in vitro and in vivo. *Exp Dermatol, 27*(10), 1112-1119. doi:10.1111/exd.13745
Garcia, S. (2019). Role of Semaphorins in Immunopathologies and Rheumatic Diseases.
Int J Mol Sci, 20(2). doi:10.3390/ijms20020374
Harris, R. G., Herzog, E. L., Bruscia, E. M., Grove, J. E., Van Arnam, J. S., &
Krause, D. S. (2004). Lack of a fusion requirement for development of bone
marrow-derived epithelia. *Science, 305*(5680), 90-93.
doi:10.1126/science.1098925
Hawkes, J. E., Yan, B. Y., Chan, T. C., & Krueger, J. G. (2018). Discovery of the IL-
23/IL-17 Signaling Pathway and the Treatment of Psoriasis. *J Immunol, 201*(6),
1605-1613. doi:10.4049/jimmunol.1800013
He, R., Tan, X., Xiang, J., Zhu, J., Jiang, Y., Liu, W., . . . Xing, Y. (2023).
Semaphorin 4A as a Potential Biomarker for Diagnosis of Systemic Lupus
Erythematosus. *Immunol Invest, 52*(1), 104-116.

600   doi:10.1080/08820139.2022.2134024

601 Heidenreich, R., Röcken, M., & Ghoreschi, K. (2009). Angiogenesis drives psoriasis
602   pathogenesis. *Int J Exp Pathol, 90*(3), 232-248. doi:10.1111/j.1365-
603   2613.2009.00669.x

604 Hui, D. H. F., Tam, K. J., Jiao, I. Z. F., & Ong, C. J. (2019). Semaphorin 3C as a
605   Therapeutic Target in Prostate and Other Cancers. *Int J Mol Sci, 20*(3).
606   doi:10.3390/ijms20030774

607 Iragavarapu-Charyulu, V., Wojcikiewicz, E., & Urdaneta, A. (2020). Semaphorins in
608   Angiogenesis and Autoimmune Diseases: Therapeutic Targets? *Front Immunol, 11*,
609   346. doi:10.3389/fimmu.2020.00346

610 Ito, D., & Kumanogoh, A. (2016). The role of Sema4A in angiogenesis, immune responses,
611   carcinogenesis, and retinal systems. *Cell Adh Migr, 10*(6), 692-699.
612   doi:10.1080/19336918.2016.1215785

613 Iyer, A. S., & Chapoval, S. P. (2018). Neuroimmune Semaphorin 4A in Cancer
614   Angiogenesis and Inflammation: A Promoter or a Suppressor? *Int J Mol Sci,*
615   *20*(1). doi:10.3390/ijms20010124

616 Jiang, J., Zhang, F., Wan, Y., Fang, K., Yan, Z. D., Ren, X. L., & Zhang, R. (2022).
617   Semaphorins as Potential Immune Therapeutic Targets for Cancer. *Front Oncol,*
618   *12*, 793805. doi:10.3389/fonc.2022.793805

619 Kanth, S. M., Gairhe, S., & Torabi-Parizi, P. (2021). The Role of Semaphorins and
620   Their Receptors in Innate Immune Responses and Clinical Diseases of Acute
621   Inflammation. *Front Immunol, 12*, 672441. doi:10.3389/fimmu.2021.672441

622 Karagianni, F., Pavlidis, A., Malakou, L. S., Piperi, C., & Papadavid, E. (2022).
623   Predominant Role of mTOR Signaling in Skin Diseases with Therapeutic
624   Potential. *Int J Mol Sci, 23*(3). doi:10.3390/ijms23031693

625 Kim, J., & Krueger, J. G. (2017). Highly Effective New Treatments for Psoriasis
626   Target the IL-23/Type 17 T Cell Autoimmune Axis. *Annu Rev Med, 68*, 255-269.
627   doi:10.1146/annurev-med-042915-103905

628 Kim, J., Lee, J., Li, X., Lee, H. S., Kim, K., Chaparala, V., . . . Krueger, J. G.
629   (2023). Single-cell transcriptomics suggest distinct upstream drivers of IL-
630   17A/F in hidradenitis versus psoriasis. *J Allergy Clin Immunol, 152*(3), 656-
631   666. doi:10.1016/j.jaci.2023.05.012

632 Koda, T., Namba, A., Kinoshita, M., Nakatsuji, Y., Sugimoto, T., Sakakibara, K., . . .
633   Okuno, T. (2020). Sema4A is implicated in the acceleration of Th17 cell-
634   mediated neuroinflammation in the effector phase. *J Neuroinflammation, 17*(1),
635   82. doi:10.1186/s12974-020-01757-w

Kolodkin, A. L., Matthes, D. J., & Goodman, C. S. (1993). The semaphorin genes encode
a family of transmembrane and secreted growth cone guidance molecules. *Cell,*
75(7), 1389-1399. doi:10.1016/0092-8674(93)90625-z

Krueger, G. G., Langley, R. G., Leonardi, C., Yeilding, N., Guzzo, C., Wang, Y., . . .
Lebwohl, M. (2007). A human interleukin-12/23 monoclonal antibody for the
treatment of psoriasis. *N Engl J Med, 356*(6), 580-592.
doi:10.1056/NEJMoa062382

Kumanogoh, A., Marukawa, S., Suzuki, K., Takegahara, N., Watanabe, C., Ch'ng,
E., . . . Kikutani, H. (2002). Class IV semaphorin Sema4A enhances T-cell
activation and interacts with Tim-2. *Nature, 419*(6907), 629-633.
doi:10.1038/nature01037

Kumanogoh, A., Shikina, T., Suzuki, K., Uematsu, S., Yukawa, K., Kashiwamura,
S., . . . Kikutani, H. (2005). Nonredundant roles of Sema4A in the immune
system: defective T cell priming and Th1/Th2 regulation in Sema4A-deficient
mice. *Immunity, 22*(3), 305-316. doi:10.1016/j.immuni.2005.01.014

Lee, E., Trepicchio, W. L., Oestreicher, J. L., Pittman, D., Wang, F., Chamian,
F., . . . Krueger, J. G. (2004). Increased expression of interleukin 23 p19
and p40 in lesional skin of patients with psoriasis vulgaris. *J Exp Med,*
199(1), 125-130. doi:10.1084/jem.20030451

Li, K., Ouyang, Z., Chen, Y., Gagnon, J., Lin, D., Mingueneau, M., . . . Zhang, B.
(2022). Cellxgene VIP unleashes full power of interactive visualization and
integrative analysis of scRNA-seq, spatial transcriptomics, and multiome data.
bioRxiv, 2020.2008.2028.270652. doi:10.1101/2020.08.28.270652

Li, Q., Shao, S., Zhu, Z., Chen, J., Hao, J., Bai, Y., . . . Wang, G. (2023). An
IGFBP7hi endothelial cell subset drives T cell extravasation in psoriasis via
endothelial glycocalyx degradation. *J Clin Invest, 133*(9).
doi:10.1172/jci160451

Liu, X., Sun, Y., Tian, W., Wang, F., Lv, X., Wang, M., . . . Han, M. (2018). Sema4A
Responds to Hypoxia and Is Involved in Breast Cancer Progression. *Biol Pharm*
Bull, 41(12), 1791-1796. doi:10.1248/bpb.b18-00423

Lowes, M. A., Russell, C. B., Martin, D. A., Towne, J. E., & Krueger, J. G. (2013).
The IL-23/T17 pathogenic axis in psoriasis is amplified by keratinocyte
responses. *Trends Immunol, 34*(4), 174-181. doi:10.1016/j.it.2012.11.005

Lu, N., Li, Y., Zhang, Z., Xing, J., Sun, Y., Yao, S., & Chen, L. (2018). Human
Semaphorin-4A drives Th2 responses by binding to receptor ILT-4. *Nat Commun,*
9(1), 742. doi:10.1038/s41467-018-03128-9

Ma, F., Plazyo, O., Billi, A. C., Tsoi, L. C., Xing, X., Wasikowski, R., . . .

Gudjonsson, J. E. (2023). Single cell and spatial sequencing define processes

by which keratinocytes and fibroblasts amplify inflammatory responses in

psoriasis. *Nat Commun, 14*(1), 3455. doi:10.1038/s41467-023-39020-4

Maeda, Y., Tsuda, T., Takeda, Y., Koyama, S., Hayama, Y., Nojima, S., . . . Kumanogoh,

A. (2019). SEMA4A promotes eosinophil survival and contributes to eosinophil-

mediated allergic diseases. *Allergol Int, 68*(2), 274-276.

doi:10.1016/j.alit.2018.10.001

Naito, M., & Kumanogoh, A. (2023). The role of semaphorins in allergic diseases.

Allergol Int. doi:10.1016/j.alit.2023.08.006

Naito, Y., Koyama, S., Masuhiro, K., Hirai, T., Uenami, T., Inoue, T., . . .

Kumanogoh, A. (2023). Tumor-derived semaphorin 4A improves PD-1-blocking

antibody efficacy by enhancing CD8(+) T cell cytotoxicity and proliferation.

Sci Adv, 9(20), eade0718. doi:10.1126/sciadv.ade0718

Nakatsuji, Y., Okuno, T., Moriya, M., Sugimoto, T., Kinoshita, M., Takamatsu,

H., . . . Kumanogoh, A. (2012). Elevation of Sema4A implicates Th cell

skewing and the efficacy of IFN-β therapy in multiple sclerosis. *J Immunol,*

188(10), 4858-4865. doi:10.4049/jimmunol.1102023

Nakayama, H., Kusumoto, C., Nakahara, M., Fujiwara, A., & Higashiyama, S. (2018).

Semaphorin 3F and Netrin-1: The Novel Function as a Regulator of Tumor

Microenvironment. *Front Physiol, 9*, 1662. doi:10.3389/fphys.2018.01662

Ni, X., & Lai, Y. (2020). Keratinocyte: A trigger or an executor of psoriasis? *J*

Leukoc Biol, 108(2), 485-491. doi:10.1002/jlb.5mr0120-439r

Nojima, S., Toyofuku, T., Kamao, H., Ishigami, C., Kaneko, J., Okuno, T., . . .

Kumanogoh, A. (2013). A point mutation in Semaphorin 4A associates with

defective endosomal sorting and causes retinal degeneration. *Nat Commun, 4*,

1406. doi:10.1038/ncomms2420

Pan, J. X., Wang, F., & Ye, L. Y. (2016). Doxorubicin-induced epithelial-mesenchymal

transition through SEMA 4A in hepatocellular carcinoma. *Biochem Biophys Res*

Commun, 479(4), 610-614. doi:10.1016/j.bbrc.2016.09.167

Pasterkamp, R. J., & Kolodkin, A. L. (2003). Semaphorin junction: making tracks

toward neural connectivity. *Curr Opin Neurobiol, 13*(1), 79-89.

doi:10.1016/s0959-4388(03)00003-5

Polański, K., Young, M. D., Miao, Z., Meyer, K. B., Teichmann, S. A., & Park, J. E.

(2020). BBKNN: fast batch alignment of single cell transcriptomes.

Bioinformatics, 36(3), 964-965. doi:10.1093/bioinformatics/btz625

Prieto, C., & Barrios, D. (2019). RaNA-Seq: Interactive RNA-Seq analysis from FASTQ
files to functional analysis. *Bioinformatics*.
doi:10.1093/bioinformatics/btz854

Rajabinejad, M., Asadi, G., Ranjbar, S., Afshar Hezarkhani, L., Salari, F., Gorgin
Karaji, A., & Rezaiemanesh, A. (2020). Semaphorin 4A, 4C, and 4D: Function
comparison in the autoimmunity, allergy, and cancer. *Gene, 746*, 144637.
doi:10.1016/j.gene.2020.144637

Reitamo, S., Spuls, P., Sassolas, B., Lahfa, M., Claudy, A., & Griffiths, C. E.
(2001). Efficacy of sirolimus (rapamycin) administered concomitantly with a
subtherapeutic dose of cyclosporin in the treatment of severe psoriasis: a
randomized controlled trial. *Br J Dermatol, 145*(3), 438-445.
doi:10.1046/j.1365-2133.2001.04376.x

Ruf, M. T., Andreoli, A., Itin, P., Pluschke, G., & Schmid, P. (2014). Ribosomal
protein S6 is hyperactivated and differentially phosphorylated in epidermal
lesions of patients with psoriasis and atopic dermatitis. *Br J Dermatol,*
171(6), 1533-1536. doi:10.1111/bjd.13248

Serini, G., Maione, F., Giraudo, E., & Bussolino, F. (2009). Semaphorins and tumor
angiogenesis. *Angiogenesis, 12*(2), 187-193. doi:10.1007/s10456-009-9138-4

Suga, Y., Nagatomo, I., Kinehara, Y., Koyama, S., Okuzaki, D., Osa, A., . . .
Kumanogoh, A. (2021). IL-33 Induces Sema4A Expression in Dendritic Cells and
Exerts Antitumor Immunity. *J Immunol, 207*(5), 1456-1467.
doi:10.4049/jimmunol.2100076

Sun, T., Yang, L., Kaur, H., Pestel, J., Looso, M., Nolte, H., . . . Worzfeld, T.
(2017). A reverse signaling pathway downstream of Sema4A controls cell
migration via Scrib. *J Cell Biol, 216*(1), 199-215. doi:10.1083/jcb.201602002

Tan, Y., Wang, Y., Li, L., Xia, J., Peng, S., & He, Y. (2015). Chemokine-like factor
1-derived C-terminal peptides induce the proliferation of dermal
microvascular endothelial cells in psoriasis. *PLoS One, 10*(4), e0125073.
doi:10.1371/journal.pone.0125073

Tsoi, L. C., Rodriguez, E., Degenhardt, F., Baurecht, H., Wehkamp, U., Volks,
N., . . . Weidinger, S. (2019). Atopic Dermatitis Is an IL-13-Dominant
Disease with Greater Molecular Heterogeneity Compared to Psoriasis. *J Invest*
Dermatol, 139(7), 1480-1489. doi:10.1016/j.jid.2018.12.018

van der Fits, L., Mourits, S., Voerman, J. S., Kant, M., Boon, L., Laman, J. D., . . .
Lubberts, E. (2009). Imiquimod-induced psoriasis-like skin inflammation in
mice is mediated via the IL-23/IL-17 axis. *J Immunol, 182*(9), 5836-5845.

744    doi:10.4049/jimmunol.0802999

745 Vo, S., Watanabe, R., Koguchi-Yoshioka, H., Matsumura, Y., Ishitsuka, Y., Nakamura,
746    Y., . . . Fujimoto, M. (2019). CD8 resident memory T cells with interleukin
747    17A-producing potential are accumulated in disease-naïve nonlesional sites of
748    psoriasis possibly in correlation with disease duration. *Br J Dermatol,*
749    *181*(2), 410-412. doi:10.1111/bjd.17748

750 Vu, T. T., Koguchi-Yoshioka, H., & Watanabe, R. (2021). Skin-Resident Memory T Cells:
751    Pathogenesis and Implication for the Treatment of Psoriasis. *J Clin Med,*
752    *10*(17). doi:10.3390/jcm10173822

753 Walter, A., Schäfer, M., Cecconi, V., Matter, C., Urosevic-Maiwald, M., Belloni,
754    B., . . . van den Broek, M. (2013). Aldara activates TLR7-independent immune
755    defence. *Nat Commun, 4,* 1560. doi:10.1038/ncomms2566

756 Wataya-Kaneda, M., Nakamura, A., Tanaka, M., Hayashi, M., Matsumoto, S., Yamamoto, K.,
757    & Katayama, I. (2017). Efficacy and Safety of Topical Sirolimus Therapy for
758    Facial Angiofibromas in the Tuberous Sclerosis Complex : A Randomized
759    Clinical Trial. *JAMA Dermatol, 153*(1), 39-48.
760    doi:10.1001/jamadermatol.2016.3545

761 Wu, Y., Zhao, R. C., & Tredget, E. E. (2010). Concise review: bone marrow-derived
762    stem/progenitor cells in cutaneous repair and regeneration. *Stem Cells, 28*(5),
763    905-915. doi:10.1002/stem.420

764 Zhang, X., Lei, L., Jiang, L., Fu, C., Huang, J., Hu, Y., . . . Zeng, Q. (2023).
765    Characteristics and pathogenesis of Koebner phenomenon. *Exp Dermatol, 32*(4),
766    310-323. doi:10.1111/exd.14709

767 Zhou, X., Chen, Y., Cui, L., Shi, Y., & Guo, C. (2022). Advances in the pathogenesis
768    of psoriasis: from keratinocyte perspective. *Cell Death Dis, 13*(1), 81.
769    doi:10.1038/s41419-022-04523-3

770 Zhu, Z., Chen, J., Lin, Y., Zhang, C., Li, W., Qiao, H., . . . Wang, G. (2020). Aryl
771    Hydrocarbon Receptor in Cutaneous Vascular Endothelial Cells Restricts
772    Psoriasis Development by Negatively Regulating Neutrophil Recruitment. *J*
773    *Invest Dermatol, 140*(6), 1233-1243.e1239. doi:10.1016/j.jid.2019.11.022

Figures

Figure 1: Epidermal Sema4A expression is downregulated in psoriasis.

(**A**) UMAP plots, generated from single-cell RNA-sequencing data (GSE220116),

illustrate cell distributions from control (Ctl) and psoriatic lesion (L) samples ($n = 10$

for Ctl, $n = 11$ for L). (**B**) Subclustering of immune cells. (**C**) *SEMA4A* expression in

keratinocytes, dendritic cells, and macrophages. *****padj < 0.001. NS, not significant.

Analyzed using Python and cellxgene VIP. (**D**) Representative immunohistochemistry

and magnified views showing Sema4A expression in Ctl, psoriatic non-lesion (NL), and

L. Scale bar = 50 µm. (**E**) Mean epidermal (Epi) Sema4A intensity in

immunohistochemistry ($n = 10$ per group). Each dot represents the average intensity

from 5 unit areas per sample. (**F**) Relative *SEMA4A* expression in Epi ($n = 10$ for Ctl, L,

$n = 7$ for L and NL) and dermis (Derm, $n = 6$ per group). (**G**) Proportions of Sema4A-

expressing cells in blood CD4 T cells (left), CD8 T cells (middle), and monocytes

(right) from Ctl and psoriatic (Pso) patients. ($n = 13$ per group in CD4 and CD8, $n = 11$

for Ctl and $n = 13$ for Pso in Monocytes). (**H**) Serum Sema4A levels in Ctl ($n = 20$) and

Pso ($n = 60$). **E** to **H**: *$p < 0.05$, **$p < 0.01$, ****$p < 0.0001$. NS, not significant.

Figure 1-source data 1

Excel file containing quantitative data for Figure 1.

Figure 2: Imiquimod-induced Psoriasis-like dermatitis is augmented in Sema4AKO mice.

(**A**) Experimental scheme. Wild-type (WT, green) mice and Sema4A knockout (KO, black) mice were treated with 10 mg/ear of 5% Imiquimod (IMQ) for 4 consecutive days. Samples for flow cytometry analysis were collected on Day 4. (**B**) Ear thickness of WT mice and KO mice on Day 4 ($n = 15$ per group). (**C**) Relative expression of *Il17a* in epidermis ($n = 5$ per group). (**D** and **E**) The percentages of Vγ3, Vγ2, Vγ2-Vγ3-γδ

(DNγδ), CD4, and CD8 T cells (**D**) and those with IL-17A production (**E**) in CD3

fraction in the Epi (top) and Derm (bottom) of WT and KO ears (n = 6 per group, each

dot represents the average of 4 ear specimens). (**F**) The percentages of IL-17A-

producing γδ, CD4, and CD8 T cells in CD3 fraction in skin-draining lymph nodes

(dLN) (n = 9 per group). **B-F**: *p < 0.05, **p < 0.01, ****p < 0.0001. NS, not

significant.

Figure 2-source data 1

Excel file containing quantitative data for Figure 2.

Figure 3: Sema4A in keratinocytes may play a role in preventing murine psoriasis-

like dermatitis.

(**A**) Experimental scheme for establishing BM chimeric mice. (**B**) IMQ Day 4 ear

thickness in the mice with the indicated genotypes ($n = 14$ for WT→WT, $n = 13$ for WT

→KO, $n =9$ for KO→WT, $n = 9$ for KO→KO). (**C**) The percentages of IL-17A-

producing Vγ3, Vγ2, DNγδ, CD4, and CD8 T cells in CD3 fraction from IMQ Day 4

Epi (top) and Derm (bottom) of the ears from WT→ WT mice and WT→ KO mice ($n =$

6 per group). Each dot represents the average of 4 ear specimens. **B**-**C**: *$p < 0.05$, **$p <$

824   0.01, ****$p < 0.0001$. NS, not significant.

826   Figure 3-source data 1

827   Excel file containing quantitative data for Figure 3.

Figure 4: Naïve Sema4AKO epidermis is thicker than WT epidermis with increased γδ T17 infiltration.

(**A**) Ear thickness of WT mice and KO mice at week (Wk) 8 (*n* = 15 per group) and

representative images. (**B**) Left: representative Hematoxylin and eosin staining of Day 0

back and Wk 8 back and ear. Scale bar = 50 μm. Right: Epi and Derm thickness in Day

0 back (*n* = 5) and Wk 8 back (*n* = 5) and ear (*n* = 8). (**C**) Relative *Sema4a* expression

in WT Epi and Derm (*n* = 5 per group). (**D**) The percentages of the IL-17A-producing

Vγ3, Vγ2, DNγδ, CD4, and CD8 T cells in CD3 fraction (*n* = 4 per group) in Epi (top)

and Derm (bottom). Each dot represents the average of 4 ear specimens. (**E**) The graphs

showing the percentages IL-17A-producing γδ, CD4, and CD8 T cells in CD3 fraction

from draining LN (dLN) of WT mice and Sema4AKO mice (*n* = 6 per group). **A**-**E**: *p*

< 0.05, **p < 0.01, ****p < 0.0001. NS, not significant.

Figure 4-source data 1

Excel file containing quantitative data for Figure 4.

Figure 5: Sema4AKO skin shares the features of human psoriatic NL.

(**A** and **B**) The volcano plot (**A**) and Gene ontology (GO) analysis (**B**), generated from

RNA-sequencing data (GSE121212) using RaNA-seq, display changes in gene

expression in psoriatic NL compared to Ctl. (**C**) The difference in the expression of

epidermal differentiation markers between Ctl and NL ($n = 38$ for Ctl, $n = 27$ for NL)

was calculated with the transcripts per million values. **padj < 0.01. NS, not significant.

(**D**) Relative gene expression of epidermal differentiation markers between wk 8 Epi of

WT mice and KO mice ($n = 5$ for *Krt14* and *Krt16*, $n = 8$ for *Krt5*, *Krt10*, *Filaggrin,* and

Loricrin). (**E**) Left: Representative immunofluorescence pictures of Krt5, Krt10, Krt14,

and Krt16 (red) overlapped with DAPI. Scale bar = 50 μm. Right: Accumulated graphs

showing the numbers of Krt5, Krt10, Krt14, and Krt16 positive cells per 100 μm width

($n = 5$ per group) of wk 8 ear (right). Each dot represents the average from 5 unit areas

per sample. (**F**) Transepidermal water loss (TEWL) in back skin of WT mice and KO

mice at wk 8 ($n = 5$ per group). **D**-**F**: *p < 0.05, **p < 0.01. NS, not significant.

Figure 5-source data 1

Excel file containing quantitative data for Figure 5.

Figure 6: mTOR signaling is upregulated in the epidermis of psoriatic NL and Sema4AKO mice.

(**A**) Representative results of immunohistochemistry displaying cells positive for p-S6 (Ser235/236), S6, p-Akt (Ser473), and Akt in Ctl, NL, and L. The graphs of accumulated data show the mean intensity of p-S6 and S6 in the upper and lower epidermal layers ($n = 9$ per group). Scale bar = 100 μm. Each dot represents the average

mean intensity from 5 unit areas per sample. (**B**) The mean intensity of p-S6

(Ser235/236) and p-Akt (Ser473), detected by immunohistochemistry in the epidermis

of WT mice and KO mice, were analyzed. Scale bar = 50 μm. Each dot represents the

average intensity from 5 unit areas per sample (n = 8 per group). (**C** and **D**)

Immunoblotting of p-S6 (Ser235/236), S6, p-Akt (Ser473), and Akt in tissue lysates

from epidermis without treatment (**C**) and with IMQ treatment for consecutive 4 days

(**D**) (n = 5 per group, except for p-Akt and Akt in **C**, for which n = 4). **A-D**: *$p < 0.05$,

**$p < 0.01$. NS, not significant.

Figure 6-source data 1

Excel file containing quantitative data for Figure 6.

Figure 7: Inhibitors of mTOR signaling modulate the expression of cytokeratins in

Sema4AKO mice.

(**A** and **B**) Epidermal thickness of Sema4AKO mice treated intraperitoneally with

vehicle (Ctl) or rapamycin (**A**), and Ctl or JR-AB2-011 (**B**) ($n = 5$ per group). (**C** and **D**)

Relative expression of keratinocyte differentiation markers and *Il17a* in Sema4AKO Epi

(**C**), and the number of T cells in Epi and Derm under Ctl or rapamycin (**D**) ($n = 5$ per

group). (**E** and **F**) Relative expression of keratinocyte differentiation markers and *Il17a*

892   in Sema4AKO Epi (**E**), and the number of T cells in Epi and Derm under Ctl or JR-

893   AB2-011 (**F**) ($n = 5$ per group). **D** and **F**: Each dot represents the sum of numbers from

894   10 unit areas across 3 specimens. **A-F**: *$p < 0.05$, **$p < 0.01$. NS, not significant.

896   Figure 7-source data 1

897   Excel file containing quantitative data for Figure 7.

 SUPPLEMENTARY MARERIALS

A

Clinical information

Sample name	Title	Age	Gender	PASI	Site
Sample01	Control01	37	Male	0	Trunk
Sample02	Control02	37	Male	0	Trunk
Sample03	Control03	66	Male	0	Trunk
Sample04	Control04	40	Male	0	Leg (right)
Sample05	Control05	40	Male	0	Leg (right)
Sample06	Control06	41	Male	0	Leg (right)
Sample07	Control07	52	Female	0	Arm (left)
Sample08	Control08	52	Female	0	Arm (left)
Sample09	Control09	74	Male	0	Trunk
Sample10	Control10	74	Male	0	Trunk
Sample11	Psoriasis01_preTx_LS	52	Female	18	Arm (right)
Sample12	Psoriasis02_preTx_LS	63	Male	7.8	Back
Sample13	Psoriasis03_preTx_LS	68	Female	8.4	Leg (left)
Sample15	Psoriasis04_preTx_LS	41	Male	9.6	Trunk
Sample17	Psoriasis05_preTx_LS	46	Female	6.8	Arm (right)
Sample18	Psoriasis06_preTx_LS	39	Male	6	Leg (right)
Sample21	Psoriasis07_preTx_LS	39	Male	9.8	Trunk
Sample22	Psoriasis07_preTx_LS_02	40	Male	18	Trunk
Sample26	Psoriasis08_preTx_LS	48	Female	7.4	Arm (right)
Sample27	Psoriasis09_preTx_LS	45	Male	6.4	Leg (right)
Sample28	Psoriasis10_preTx_LS	38	Male	8.4	Leg (right)

B

Immune cell
Keratinocyte
Neural crest-like cell
fibroblast

ITGAM CD3E SOX10 KRT10 KRT14 KRT1 PTPRC MLANA S100B DCN COL1A1

Fraction of cells in group (%)
20 40 60 80 100
Mean expression in group
1 2

C

CD4 T cell
CD8 T cell
Dendritic cell
Macrophage
NK cell
Plasma cell

KLRB1 GNLY CD3D TRAC TRBC1 CD8A GZMK LAMP3 LY75 CIITA CD40 HLA-DQA1 HLA-DQB1 HLA-DRB1 HLA-DRA HLA-DRB5 CD14 CD68 CD163 CD79A MS4A1 IGLC2 IGKC

Fraction of cells in group (%)
20 40 60 80 100
Mean expression in group
0.0 2.5

D

Figure 1-figure supplement 1: Sema4A is downregulated in the keratinocytes of lesional psoriasis in the single-cell RNA-sequencing data.

(**A**) Sample information for specimens from Ctl and psoriatic L (GSE220116). (**B** and **C**) Clusters of cells were identified by their expression patterns of signature genes. (**D**) The volcano plot displays changes in gene expression in psoriatic L compared to Ctl.

907   Figure 1-figure supplement 1-source data 1

908   Excel file containing quantitative data for Figure 1-figure supplement 1.

Figure 1-figure supplement 2: Gating strategy in flow cytometry.

Gating strategy for human Sema4A expression in blood cells. Large and small cells

were distinguished using forward scatter (FCS) and side scatter (SSC) in a dot plot

panel, with dead cells being excluded. Monocytes were defined within the live large cell

population as CD11c positive. CD4 and CD8 T cells were identified within the live

small cell population as CD3 positive CD4 positive and CD3 positive CD8 positive

populations, respectively. The empty histogram represents the flow cytometry minus

one control for Sema4A.

Figure 2-figure supplement 1: Gating strategy in flow cytometry.

Gating strategy for murine T cells infiltrating the epidermis and dermis. After excluding

dead cells, TCRγδ positive T cells were evaluated for the expression of Vγ2. TCRγδ

positive Vγ2 negative population was further assessed the expression of Vγ3. The CD3

positive TCRγδ negative population was evaluated for the expression of CD4 and CD8.

Each population was analyzed for cytokine production. The empty histogram represents

the isotype control for IL-17A.

Figure 2-figure supplement 2: IL-23-mediated psoriasis-like dermatitis is

augmented in Sema4AKO mice.

(**A**) An experimental scheme involved intradermally injecting 20 μl of phosphate-

buffered saline containing 500 ng of recombinant mouse IL-23 into both ears of WT

mice and KO mice for 4 consecutive days. Samples for following analysis were

collected on Day 4. (**B** and **C**) Ear thickness (**B**) and Epi and Derm thickness (**C**) of WT

mice and KO mice on Day 4 (*n* = 12 per group). (**D** and **E**) The percentages of Vγ3, Vγ2,

DNγδ, CD4, and CD8 T cells (**D**) and those with IL-17A production (**E**) in CD3 fraction

in the Epi (top) and Derm (bottom) of WT and KO ears (*n* = 5 per group). Each dot

939   represents the average of 4 ear specimens. **B**-**E**: *$p < 0.05$, ***p < 0.01$. NS, not

940   significant.

942   Figure 2-figure supplement 2-source data 1

943   Excel file containing quantitative data for Figure 2-figure supplement 2.

Figure 3-figure supplement 1: T cells' fractions infiltrating in the chimeric mice ear.

(**A**) *Sema4a* expression in the Epi of WT→ WT mice and WT→ KO mice (*n* = 8 for

WT→ WT, *n* = 7 for WT→ KO). (**B**) The percentages of Vγ3, Vγ2, DNγδ, CD4, and

CD8 T cells in CD3 fraction from IMQ Day 4 Epi (top) and Derm (bottom) of the ears

from WT→ WT mice and WT→ KO mice (*n* = 6 per group). Each dot represents the

average of 4 ear specimens. **A**-**B**: *p < 0.05, ***p < 0.001. NS, not significant.

Figure 3-figure supplement 1-source data 1

954   Excel file containing quantitative data for Figure 3-figure supplement 1.

Figure 4-figure supplement 1: Naive Sema4AKO skin shows upregulation of

psoriasis related genes and an increase in resident memory T cells.

(**A**) Relative expression of psoriasis-associated genes in Epi (top) and Derm (bottom) of

WT mice and KO mice (*n* = 5 per group, #: not detected). (**B**) Representative dot plots

showing CD69 and CD103 expression in the indicated T cell fractions from whole skin.

The graphs show T cell counts per ear (top) and those with resident memory phenotype

(bottom) (*n* = 7 per group). Each dot represents the average of 4 ear specimens. **A**-**B**: *p

< 0.05, **p < 0.01. NS, not significant.

966   Figure 4-figure supplement 1-source data 1

967   Excel file containing quantitative data for Figure 1-figure supplement 1.

Figure 4-figure supplement 2: Expression of IFNγ and IL-4 is comparable between

naive WT and Sema4AKO skin.

(**A**) The percentages of Vγ3, Vγ2, DNγδ, CD4, and CD8 T cells in CD3 fraction from

naive WT and KO mice. (**B** and **C**) The graphs presenting the percentages of IFNγ (**B**)

and IL-4 (**C**) -producing Vγ2, DNγδ, CD4, and CD8 T cells in CD3 fraction in the Epi

(top) and Derm (bottom) of naive WT mice and KO mice ($n = 4$ per group). **A-C**: Each

976   dot represents the average of 4 ear specimens. *$p < 0.05$. NS, not significant.

978   Figure 4-figure supplement 2-source data 1

979   Excel file containing quantitative data for Figure 4-figure supplement 2.

Figure 4-figure supplement 3: Comparable T17 differentiation potential under

Th17-skewing conditions between WT mice and Sema4AKO mice.

Splenic T cells were cultured for 2 weeks, followed by flow cytometry analysis. The

accumulated data display the percentages of IL-17A-producing (right) and IFNγ-

producing (left) γδ, CD4, and CD8 T cells within CD3 fraction under various

conditions: IL-23 dependent Th17-skewing condition (top), IL-23 only Th17-skewing

condition (middle), and IL-23 independent Th17-skewing condition (bottom) . NS, not

significant.

Figure 4-figure supplement 3-source data 1

Excel file containing quantitative data for Figure 4-figure supplement 3.

995 **Figure 5-figure supplement 1: The epidermis of psoriatic non-lesion is thicker than**

996 **that of control skin.**

997 Epidermal thickness of Ctl and psoriatic NL ($n = 10$ per group). ***$p < 0.001$.

999 Figure 5-figure supplement1-source data1

1000 Excel file containing quantitative data for Figure 5-figure supplement1.

1001

1002

Figure 5-figure supplement 2: Upregulation of cytokeratin expression related to

psoriasis is not detected at birth in Sema4AKO mice.

Representative immunofluorescence pictures of Krt5, Krt10, Krt14, and Krt16 (red)

overlapped with DAPI, and the accumulated graphs showing the numbers of Krt5,

Krt10, Krt14, and Krt16 positive cells per 100 μm width ($n = 5$ per group) in the

epidermis of Day 0 back. Scale bar = 50 μm. Each dot represents the average from 5

unit areas per sample. NS, not significant.

1010

1011 Figure 5-figure supplement 2-source data 1

1012 Excel file containing quantitative data for Figure 5-figure supplement 2.

1013

Figure 7-figure supplement 1: Topical application of rapamycin gel yields partially similar results to intraperitoneal treatment.

(**A**) Comparison of Epi thickness between vehicle (Ctl) gel-treated right ears and rapamycin gel-treated left ears of Sema4AKO mice ($n = 10$ per group). (**B**) Relative expression of keratinocyte differentiation markers and *Il17a* in Sema4AKO Epi under Ctl gel or rapamycin gel treatments ($n = 5$ per group). (**C**) The number of T cells in the Epi (left) and Derm (right), under Ctl gel or Rapamycin gel treatments ($n = 5$ per group). Each dot represents the sum of numbers from 10 unit areas across 3 specimens. **A-C**: *$p < 0.05$, **$p < 0.01$. NS, not significant.

1025   Figure 7-figure supplement 1-source data 1

1026   Excel file containing quantitative data for Figure 7-figure supplement 1.

1027

1028

Figure 7-figure supplement 2: Rapamycin treatment reduced the epidermal

swelling observed in IMQ-treated Sema4AKO mice.

(**A**) Experimental scheme. (**B**) The Epi thickness on Day 18. ($n = 10$ for Ctl, $n = 12$ for

Rapamycin). (**C**) Relative expression of keratinocyte differentiation markers and *Il17a*

in Sema4AKO Epi ($n = 10$ for Ctl, $n = 12$ for Rapamycin). (**D**) The number of T cells in

the Epi (left) and Derm (right), under Ctl or rapamycin and IMQ treatments ($n = 10$ for

Ctl, $n = 12$ for Rapamycin). Each dot represents the sum of numbers from 10 unit areas

across 3 specimens. **A**-**C**: *$p < 0.05$, **$p < 0.01$. NS, not significant.

1037

Figure 7-figure supplement 2-source data 1

Excel file containing quantitative data for Figure 7-figure supplement 2.

Table S1   Patient information

Psoriasis severity is defined by the total body surface area (BSA) affected: <3% BSA for mild, 3%-10% BSA for moderate, and >10% BSA for severe disease.

Fig. 1, D and E   Analysis of Sema4A expression by immunohistochemistry

Fig. S4   Epidermal thickness

Psoriasis (n = 10)

Age	Gender	Body site	Severity
66	Male	Leg	Severe
55	Male	Trunk	Moderate
80	Female	Trunk	Severe
72	Male	Trunk	Moderate
30	Male	Trunk	Severe
72	Female	Leg	Moderate
63	Male	Arm	Mild
38	Male	Trunk	Mild
35	Male	Leg	Mild
71	Male	Trunk	Moderate

Control (n = 10)

Age	Gender	Body site
76	Male	Neck
52	Female	Leg
66	Female	Leg
33	Male	Leg
68	Female	Trunk
57	Female	Leg
65	Male	Trunk
41	Female	Trunk
51	Female	Trunk
66	Female	Trunk

Fig 1F   Analysis of Sema4A expression by qRT-PCR

Epidermis

Psoriasis (n = 7)

Age	Gender	Body site	Severity
71	Male	Leg	Moderate
76	Female	Neck	Mild
38	Female	Trunk	Moderate
49	Female	Trunk	Moderate
61	Male	Arm	Mild
68	Female	Trunk	Moderate
72	Male	Leg	Mild

Control (n = 10)

Age	Gender	Body site
57	Female	Leg
69	Female	Trunk

66	Female	Trunk
86	Female	Trunk
53	Female	Trunk
84	Male	Neck
37	Female	Trunk
46	Female	Trunk
51	Female	Trunk
46	Female	Trunk

Dermis

Psoriasis (n = 6)

Age	Gender	Body site	Severity
76	Female	Neck	Mild
38	Female	Trunk	Moderate
49	Female	Trunk	Moderate
61	Male	Arm	Severe
68	Female	Trunk	Moderate
72	Male	Leg	Severe

Control (n = 6)

Age	Gender	Body site
69	Female	Trunk
86	Female	Trunk
84	Male	Neck
37	Female	Trunk
51	Female	Trunk
46	Female	Trunk

Fig 1G Analysis of Sema4A expression by flow cytometry

	Cell type	Average of age	Male (n =)	Female (n =)
Psoriasis (n = 13)	CD4 T cell, CD8 T cell, and monocyte	56.1	9	4
Control (n = 13)	CD4 T cell and CD8 T cell	52.2	6	7
Control (n = 11)	Monocyte	54.1	6	5

Fig 1H Analysis of Serum Sema4A by ELISA

	Average of age	Male (n =)	Female (n =)
Psoriasis (n = 60)	55.3	43	17
Control (n = 20)	52.1	11	9

Fig 6A Pictures of p-S6, S6, p-Akt, and Akt staining

Psoriasis (n = 1)

Age	Gender	Body site	Severity
30	Male	Trunk	Severe

Control (n = 1)

Age	Gender	Body site
66	Female	Trunk

Analysis of p-S6, S6 expression by immunohistochemistry

Psoriasis (n = 9)

Age	Gender	Body site	Severity
66	Male	Leg	Severe
55	Male	Trunk	Moderate
80	Female	Trunk	Severe
72	Male	Trunk	Moderate
30	Male	Trunk	Severe
72	Female	Leg	Moderate
38	Male	Trunk	Mild
35	Male	Leg	Mild
71	Male	Trunk	Moderate

Control (n = 9)

Age	Gender	Body site
76	Male	Neck
66	Female	Leg
33	Male	Leg
68	Female	Trunk
57	Female	Leg
65	Male	Trunk
41	Female	Trunk
51	Female	Trunk
66	Female	Trunk

1040

1041

Table S2
Antibodies used for immunohistochemical, immunofluorescence, and western blot analyses

Target	Clone	Species	Supplier	Catalog	Dilution ratio		Incubation time	Incubation temperature
					IHC, IF	WB		
Sema4A	Polyclonal	Rabbit	Abcam	ab70178	100		30 min	RT
phospho-S6 (Ser235/236)	D57.2.2E	Rabbit	Cell Signaling	4858	400	2000	overnight	4 °C
S6	5G10	Rabbit	Cell Signaling	2217	100	1000	overnight	4 °C
phospho-Akt (Ser473)	D9E	Rabbit	Cell Signaling	4060	50	1000	overnight	4 °C
Akt	C67E7	Rabbit	Cell Signaling	4691	300	1000	overnight	4 °C
Keratin 5	Polyclonal	Rabbit	BioLegend	905503	800		overnight	4 °C
Keratin 10	Polyclonal	Rabbit	BioLegend	905403	400		overnight	4 °C
Keratin 14	Polyclonal	Rabbit	BioLegend	905303	400		overnight	4 °C
Cytokeratin 16	8L6R4	Rabbit	Invitrogen	MA5-42892	100		overnight	4 °C
CD3	CD3-12	Rat	Bio-Rad	MCA1477	100		overnight	4 °C
Rabbit IgG H&L (Alexa Fluor 555)	Polyclonal	Donkey	Abcam	ab150074	1000		30 min	RT
Rat IgG H&L (Alexa Fluor 555)	Polyclonal	Donkey	Abcam	ab150154	1000		30 min	RT
β-actin	AC-15	Mouse	Sigma-Aldrich	A5441		5000	overnight	4 °C
Anti-Mouse IgG, HRP-Linked Whole Ab	Monoclonal secondary	Sheep	Cytiva	NA931		10000	1 hour	RT
Anti-Rabbit IgG, HRP-Linked Whole Ab	Polyclonal secondary	Donkey	Cytiva	NA934		10000	1 hour	RT

RT: Room temperature
IHC: immunohistochemistry
IF: immunofluorescence
WB: western blot

1042

1043

Table S3

Primer sequences of real-time quantitative PCR used in human sample experiments

Gene Symbol	Forward primer (5'-3')	Reverse primer (5'-3')
GAPDH	GTCTCCTCTGACTTCAACAGCG	ACCACCCTGTTGCTGTAGCCAA
SEMA4A	TCTGCTCCTGAGTGGTGATG	AAACCAGGACACGGATGAAG

Primer sequences of real-time quantitative PCR in murine sample experiments

Gene Symbol	Forward primer (5'-3')	Reverse primer (5'-3')
Gapdh	TGTGTCCGTCGTGGATCTGA	TTGCTGTTGAAGTCGCAGGAG
Krt5	CAGAGCTGAGGAACATGCAG	CATTCTCAGCCGTGGTACG
Krt10	CGTACTGTTCAGGGTCTGGAG	GCTTCCAGCGATTGTTTCA
Krt14	CGTACTGTTCAGGGTCTGGAG	GCTTCCAGCGATTGTTTCA
Krt16	TGAGCTGACCCTGTCCAGA	TGAGCTGACCCTGTCCAGA
Filaggrin	GGAGGCATGGTGGAACTGA	TGTTTATCTTTTCCCTCACTTCTACATC
Loricrin	TCACTCATCTTCCCTGGTGCTT	GTCTTTCCACAACCCACAGGA
Sema4a	ATGGCCCTACCATCCCTGG	AGCAGCGTGTCAAAGTCTCG
Ccl20	ATGGCCTGCGGTGGCAAGCGT	CATCTTCTTGACTCTTAGGC
S100a8	TGCGATGGTGATAAAAGTGG	GGCCAGAAGCTCTGCTACTC
S100a9	CACCCTGAGCAAGAAGGAAT	TGTCATTTATGAGGGCTTCATTT
Ifnγ	ATCTGGAGGAACTGGCAAAA	TTCAAGACTTCAAAGAGTCTGAGGTA
Tnfα	GCCTCCCTCTCATCAGTTCT	CACTTGGTGGTTTGCTACGA
Il4	ACAGGAGAAGGGACGCCAT	GAAGCCCTACAGACGAGCTCA
Il17a	CTGTGTCTCTGATGCTGTTG	ATGTGGTGGTCCAGCTTTC
Il23	TCCCTACTAGGACTCAGCCAAC	GCTGCCACTGCTGACTAGAA

1044

1045

Table S4
Antibodies used for flow cytometry analysis

Target	Clone	Species	Supplier	Catalog
CD3	OKT3	Mouse	BioLegend	317335
CD4	RPA-T4	Mouse	BioLegend	300512
CD8a	RPA-T8	Mouse	eBioscience	47-0088-42
CD11c	3.9	Mouse	BioLegend	301628
SEMA4A	5E3/SEMA4A	Mouse	BioLegend	148404
CD3ε	145-2C11	Armenian Hamster	BioLegend	100328
CD4	GK1.5	Rat	BioLegend	100406
CD8a	53-6.7	Rat	BioLegend	100714
CD16/32	93	Rat	BioLegend	101301
CD69	H1.2F3	Armenian Hamster	BioLegend	104514
CD103	2E7	Armenian Hamster	BioLegend	121422
TCR Vγ2	UC3-10A6	Armenian Hamster	BioLegend	137705
TCR Vγ3	536	Syrian Hamster	BD Biosciences	743241
TCRγδ	GL3	Armenian Hamster	BioLegend	118124
IFNγ	XMG1.2	Rat	BioLegend	505813
IL-4	11B11	Rat	BD Biosciences	562915
IL-17A	TC11-18H10.1	Rat	BioLegend	506925

1046

1047

Table S5
Mouse recombinant cytokines

Cytokines	Supplier	Catalog
IL-1β	BioLegend	575102
IL-6	BioLegend	575702
IL-23	BioLegend	589002
TGFβ1	BioLegend	763102

1048

A 10 Gy

WT   KO   WT   KO

↓←WT-BM→↓   ↓←KO-BM→↓

WT→WT   WT→KO   KO→WT   KO→KO

IMQ application

B IMQ Day 4

Ear thickness (mm)

****
** **

0.30
0.25
0.20

WT→WT WT→KO KO→WT KO→KO

C IMQ Day 4

IL-17A production

Epi

| Vγ3 | Vγ2 | DNγδ | CD4 | CD8 |

NS — Vγ3: 0.25, 0.20, 0.15, 0.10, 0.05, 0.00 — % in CD3

NS — Vγ2: 3, 2, 1, 0

* — DNγδ: 4, 3, 2, 1, 0

NS — CD4: 5, 4, 3, 2, 1, 0

* — CD8: 4, 3, 2, 1, 0

WT→WT WT→KO

Derm

NS — 0.10, 0.08, 0.06, 0.04, 0.02, 0.00 — % in CD3

* — 8, 6, 4, 2, 0

NS — 10, 8, 6, 4, 2, 0

NS — 8, 6, 4, 2, 0

NS — 0.2, 0.1, 0.0

WT→WT WT→KO

A Wk 8 naïve

Ear thickness (mm)

WT KO

****

WT KO

B

WT KO

Day 0 Back

Wk 8 Back

Ear

Day 0 Back

Epi   Derm

NS   NS

WT KO   WT KO

Thickness (μm)

Wk 8 Back

Epi   Derm

*   NS

WT KO   WT KO

Thickness (μm)

Ear

**   NS

WT KO   WT KO

Thickness (μm)

C *Sema4a*

Relative expression

Epi   Derm

**

D IL-17A production

Epi

Vγ3   Vγ2   DNγδ   CD4   CD8

NS   *   *   NS   NS

% in CD3

WT KO   WT KO   WT KO   WT KO   WT KO

Derm

NS   NS   NS   NS   NS

% in CD3

WT KO   WT KO   WT KO   WT KO   WT KO

E IL-17A production

dLN

γδ   CD4   CD8

NS   NS   NS

% in CD3

WT KO   WT KO   WT KO

A

−log₁₀ *padj*

SPRR2F
SERPINB4
S100A7A
CXCL8
DEFB4B
PI3
SPRR2A
S100A8
S100A9
S100A7
DEFB4A
KRT14
KRT16
KRT5
KRT10
SEMA4A

log₂ Fold-change

B

GO

Peptide cross-linking
Defense response to virus
Keratinocyte differentiation
Type I interferon signaling pathway
Interferon-gamma-mediated signaling pathway
Inflammatory response
Viral process
Proteasomal protein catabolic process
Proteasomal ubiquitin-independent protein catabolic process
Response to virus

−log₁₀*P*

C

Expression

KRT5 ** | *KRT10* NS | *KRT14* ** | *KRT16* ** | *FILAGGRIN* NS | *LORICRIN* NS

Ctl NL

D

Epi   Relative expression

Krt5 NS | *Krt10* * | *Krt14* * | *Krt16* ** | *Filaggrin* ** | *Loricrin* NS

WT KO

E

Wk 8
Ear   WT   KO

Krt5
Krt10
Krt14
Krt16

Krt5 *
Krt10 *
/ unit
WT KO

Krt14 *
Krt16 *
/ unit
WT KO

F

TEWL NS
g/m²/h
WT KO

A

Mean intensity

Upper layer | Lower layer

p-S6 (Ser235/236) | S6 | p-Akt (Ser473) | Akt

Ctl
NL
L

B

p-S6 (Ser235/236) p-Akt (Ser473)

WT
KO

Mean intensity

C

p-S6 (Ser235/236)
S6
β-actin

p-Akt (Ser473)
Akt
β-actin

p-S6 (Ser235/236) p-Akt (Ser473)

S6 Akt

Relative band intensity

D

p-S6 (Ser235/236)
S6
β-actin

p-Akt (Ser473)
Akt
β-actin

p-S6 (Ser235/236) p-Akt (Ser473)

S6 Akt

Relative band intensity

A Rapamycin ip

Epi

NS

Thickness (μm)

Ctl / Rapamycin

B JR-AB2-011 ip

NS

Thickness (μm)

Ctl / JR-AB2-011

C Rapamycin ip

Epi

Krt5 ** / *Krt10* NS / *Krt14* NS / *Krt16* ** / *Il17a* NS

Relative expression

Ctl / Rapamycin

D Rapamycin ip

CD3 in Epi * # CD3 in Derm NS

Sum of 10 areas

Ctl / Rapamycin

E JR-AB2-011 ip

Epi

* / ** / ** / 0.0556 / NS

Relative expression

Ctl / JR-AB2-011

F JR-AB2-011 ip

NS   NS

Sum of 10 areas

Ctl / JR-AB2-011