A drought stress-responsive metabolite malate modulates stomatal responses through G-

protein-dependent pathway

Yoshiharu Mimata[1], Ruhai Gong[1,2], Xuanxuan Pei[1], Guochen Qin[1], Wenxiu Ye[1*]

[1]Peking University Institute of Advanced Agricultural Sciences, Weifang 261325, Shandong,

China

[2]College of Horticulture, Shanxi Agricultural University, Taigu 030801, Shanxi, China

*Corresponding author: wenxiu.ye@pku-iaas.edu.cn

Running title: TCA cycle metabolites regulate guard cell signaling

Abstract

Drought poses a significant environmental threat to global agricultural production and
distribution. Plant adaptation to dehydration stress involves intricate biological processes with
substantial changes in metabolite composition. In this study, we investigated the role of
tricarboxylic acid (TCA) cycle metabolites in drought tolerance in grapevine and Arabidopsis.
Metabolome analysis revealed that malate, citrate, and isocitrate were upregulated over time in
detached grapevine leaves. Five out of 10 organic acids tested, including TCA cycle metabolites,
elicited increases in cytosolic free Ca^{2+} concentration ($[Ca^{2+}]_{cyt}$) in guard cells. Eight out of 10
directly activated a major anion channel in guard cell SLAC1, suggesting that SLAC1 is a
general sensor of carboxylic acids. Furthermore, malate alone remarkably induced stomatal
closure, which required increases in $[Ca^{2+}]_{cyt}$ in guard cells and activation of SLAC1.
Pharmacological experiments suggested that cyclic ADP-ribose (cADPR), cAMP and inositol
trisphosphate (IP_3) act as second messengers essential for malate-induced $[Ca^{2+}]_{cyt}$ elevation
and stomatal closure. G-proteins, which regulate second messenger production, were identified
as components of malate signaling. These results indicate that malate plays a key role in
connecting metabolic regulation and drought tolerance. As TCA cycle intermediates are key
players in stress responses in mammals as well, we propose that they are common stress-
responsive signal molecules mediated by G-protein-dependent signal cascades in both the
animal and plant kingdoms.

Introduction

Drought constitutes a major environmental stress with global implications for crop survival and yields. Grapevine (*Vitis*) is one of the oldest domesticated crops and holds crucial economic importance for industries through the production of wine, brandy, juice, table grapes, and raisins. Despite the increasing demand for grapes and grape products, the global vineyard area is diminishing annually. In 2023, wine production was anticipated to reach its lowest levels in 60 years, primarily due to the impacts of global climate change, especially drought conditions (http://www.oiv.int/). In response to drought, plants synthesize a phytohormone abscisic acid (ABA) and close stomatal pores, formed by pairs of guard cells in the epidermis of leaves, to prevent excessive water loss through guard cell signaling (Murata et al., 2015). Thus, gaining a comprehensive understanding of the molecular mechanisms underlying stomatal closure is crucial for enhancing drought tolerance in plants.

Stomatal closure is initiated by the transport of anions across the plasma membrane of guard cells through the slow-type anion channel encoded by the *SLOW ANION CHANNEL-ASSOCIATED 1* (*SLAC1*) gene (Schroeder et al., 2001; Negi et al., 2008; Vahisalu et al., 2008). SLAC1 is phosphorylated and activated by cytosolic Ca^{2+} sensor kinases, CALCIUM-DEPENDENT PROTEIN KINASEs (CDPKs) (Brandt et al., 2015), leading to a decrease in turgor pressure and subsequent stomatal closure. Cytosolic free Ca^{2+} acts as a ubiquitous second messenger, and its concentration transiently increases in response to environmental, developmental, and growth signals (Luan and Wang, 2021). The increase in cytosolic free Ca^{2+} concentration ($[Ca^{2+}]_{cyt}$) results from the uptake of Ca^{2+} into the cell and the release of Ca^{2+} from internal stores through Ca^{2+} channels in response to a membrane potential shift (Hamilton et al., 2000) and second messengers such as cyclic adenosine diphosphate ribose (cADPR) (Leckie et al., 1998), cyclic adenosine monophosphate (cAMP) (Lemtiri-Chlieh and Berkowitz, 2004), inositol trisphosphate (IP_3) (Gilroy et al., 1990), reactive oxygen species (ROS) (Pei et al., 2000), nitric oxide (NO) (Garcia-Mata et al., 2003), cyclic guanosine monophosphate (cGMP) (Wang et al., 2013), and nicotinic acid adenine dinucleotide phosphate (NAADP) (Navazio et al., 2000). In guard cells, cytosolic Ca^{2+} binds to the EF hands of CDPKs, leading to activation of CDPKs, which then induces stomatal closure through the phosphorylation of

SLAC1 (Brandt et al., 2015).

Heterotrimeric G-proteins, composed of Gα, Gβ, and Gγ subunits, play pivotal roles in the

generation of second messengers, thereby participating in various biological processes such as

growth, development, and responses to environmental stimuli (Jin et al., 2013; Pandey 2020).

The genome of *A. thaliana* encodes one canonical Gα (GPA1), one Gβ (AGB1), and three Gγ

subunits (AGG1–AGG3). GPA1 and/or AGB1 are involved in regulating stomatal movements

and activating slow-type anion channels (Wang et al., 2001; Fan et al., 2008; Yu et al., 2018).

The regulation of metabolism stands out as a key mechanism for maintaining cell osmotic

potential during abiotic stress. Despite a comprehensive description of metabolic responses to

dehydration stress in *Arabidopsis thaliana* (Urano et al., 2009), the specific contributions of

these metabolites to stress tolerance remain unclear. The metabolic reprogramming triggered

by drought leads to elevated tricarboxylic acid (TCA) cycle intermediates in Arabidopsis leaves

(Pires et al., 2016). TCA cycle intermediates play an essential role in providing energy fuels

and metabolic precursors. Recent findings have demonstrated that TCA cycle intermediates also

function as signaling molecules in mammals. Succinate modulates systemic energy metabolism

as a metabolic signal "metabokine" *via* G-protein signaling cascades (Krzak et al., 2021).

Malate as well as succinate directly binds to a G-protein coupled receptor and causes rapid

increases in intracellular $[Ca^{2+}]$ and IP_3 accumulation (Trauelsen et al., 2017). In plants, malate

is crucial due to its significant associations with stomatal movements, aluminum toxicity, CO_2

fixation, ripening, and the taste of berries. Particularly in grapevines, malate is instrumental in

determining wine quality and facilitating the growth of microorganisms for vinification

(Sweetman et al., 2009; Fernie and Martinoia, 2009). The accumulation of malate is induced

by environmental changes and may be linked to physiological responses in various tissues such

as leaves, xylem, roots, and mesocarp (Van Kirk and Raschke, 1978; Kondo and Maruta, 1987;

Hedrich et al., 1994; Gabriel and Kesselmeier, 1999; Patonnier et al., 1999; Malcheska et al.,

2017; Delhaize et al., 1993; Wada et al., 2008). However, the regulatory mechanism of

physiological responses by TCA cycle metabolites remains unclear in plants.

Previous studies have shown that malate increases $[Ca^{2+}]_{cyt}$ and ROS production in guard cells

and directly activates SLAC1, leading to stomatal closure in *A. thaliana* (Mimata et al., 2022a;

2022b). In this study, we demonstrate that several TCA cycle metabolites accumulate in

grapevine leaves during dehydration stress, among which malate most effectively regulates

stomatal response *via* a G-protein signaling cascade. We propose that malate forms a hub

between energy homeostasis and stress response.

Results

Metabolic responses of TCA cycle metabolites to dehydration stress in grapevine leaves

To investigate metabolic changes in response to drought, grapevine (*Vitis vinifera*) leaves were

subjected to water-deficit stress. Detached leaves were sampled at 0–24 h of the dehydration

stress treatment and subjected to non-targeted metabolome analysis. Principal component

analysis (PCA) revealed that metabolite level changes were not pronounced between 0 and 1 h

but gradually increased thereafter (Figure 1A). Significant metabolite changes were classified

into three patterns: gradual decrease (subclass 1), gradual increase (subclass 2), and increase

followed by decrease (subclass 3) (Supplemental Figure 1A). The endogenous level of ABA,

an indicator of drought stress, increased after 2 h of dehydration stress and reached a plateau

after 6 h (Figure 1B). After 24 h of treatment, out of 2,407 metabolites, 436 were upregulated,

and 80 were downregulated (Figure 1C and Supplemental Data 1). Amino acids, lipids,

terpenoids, phenolic acids, alkaloids, and flavonoids were the primary categories exhibiting

changes (Figure 1D and Supplemental Figure 1B). Among the upregulated metabolites, we

specifically examined three TCA cycle metabolites—malate, isocitrate, and citrate—since

malate functions as a signaling molecule inducing stomatal closure (Mimata et al., 2022a).

Malate, isocitrate, and citrate increased after 12 or 24 h of dehydration stress (Figure 1E). In

contrast, *cis*-aconitate decreased immediately after 2–4 h and then returned to basal values.

Effects of TCA cycle metabolites on the $[Ca^{2+}]_{cyt}$ in guard cells

Since Ca^{2+} is a critical second messenger in guard cell signaling, we investigated the effects of

TCA cycle metabolites on the elevations of $[Ca^{2+}]_{cyt}$ through live cell imaging of *A. thaliana*

guard cells expressing a Ca^{2+} sensor fluorescent protein Yellow Cameleon 3.6. Exogenous

application of fumarate, malate, α-ketoglutarate, pyruvate, and acetate induced elevations in

$[Ca^{2+}]_{cyt}$, significantly increasing the frequency of $[Ca^{2+}]_{cyt}$ elevation (Figure 2 and

Supplementary Figure 2A). The malate-induced $[Ca^{2+}]_{cyt}$ increases were not observed in the

presence of a Ca^{2+} channel blocker La^{3+} (Figure 2A and 2D). These findings suggest that α-

hydroxy or α-keto acids (Supplementary Figure 2B) showed specificity in inducing $[Ca^{2+}]_{cyt}$

elevation in guard cells.

Interestingly, acetate caused transient long-term $[Ca^{2+}]_{cyt}$ increase (Figure 2A), likely due to its

higher pKa, leading to cytosolic acidification (Felle 1988), H^+-induced Ca^{2+} release from the

ER (Huang et al., 2023), and fluorescence quenching (Supplemental Figure 2A; Nagai et al.,

2004).

Effects of TCA cycle metabolites on the activation of SLAC1 expressed in Xenopus oocytes

Activation of the SLAC1 anion channel is crucial for stomatal closure (Vahisalu et al., 2008;

Negi et al., 2008). To explore the modulation of VvSLAC1 activity by TCA cycle metabolites,

we used a two-electrode voltage-clamp technique on Xenopus oocytes expressing VvSLAC1.

The negative currents in oocytes expressing VvSLAC1 were minimal (Supplementary Figure

3), resembling the characteristics of the inactive form of AtSLAC1 (Mimata et al., 2022a).

Constitutively active AtSLAC1 mutants, such as AtSLAC1T513D and AtSLAC1F450A,

exhibit substantial basal activity and their activity were further enhanced by malate in oocytes

(Chen et al., 2010; Maierhofer et al., 2014; Mimata et al., 2022a). Therefore, we expressed

VvSLAC1T503D, corresponding to AtSLAC1T513D, and VvSLAC1F440A, corresponding to

AtSLAC1F450A, in oocytes. These mutants exhibited 3.5- and 4.8-fold larger currents at −160

mV compared to oocytes expressing wild-type VvSLAC1, respectively (Supplementary Figure

3). We then analyzed VvSLAC1 activity during perfusion with bathing solution supplemented

with TCA cycle metabolites. Carboxylic acids did not increase the negative currents in oocytes

expressing wild-type VvSLAC1 (Figure 3A and 3C). On the other hand, all the carboxylic acids

except for acetate and pyruvate significantly increased the negative currents in oocytes

expressing VvSLAC1F440A (Figure 3A and 3C), similar to the results observed for AtSLAC1

(Figure 3B and 3D). Given the conservation of SLAC1 across various plant species (Hedrich

and Geiger, 2017), these findings suggest that SLAC1 may serve as a universal sensor for

carboxylic acids.

Malate emerges as a specific modulator in the regulation of stomatal responses

To assess how TCA cycle metabolites influence plant responses to drought stress, we measured

stomatal aperture in the presence of carboxylic acids. After stomata were fully open under light,

each carboxylic acid was applied, and stomatal aperture was measured. A significant reduction

in stomatal aperture was observed exclusively with malate treatment in *V. vinifera* (Figure 4A).

Consistent results were obtained using *A. thaliana* (Figure 4B). Furthermore, the malate-

induced stomatal closure was also observed in tomatoes and chili peppers (Supplemental Figure

4). These findings suggest that malate specifically acts as a modulator of stomatal closure, and

malate signaling is conserved across various plant species.

To further investigate the malate signaling pathway, we conducted a series of pharmacological

experiments (Supplemental Table 1). Malate-induced stomatal closure in both *V. vinifera* and *A.*

thaliana was abolished by anion channel blockers, 4,4′-diisothiocyano-2,2′-stilbenedisulfonic

acid (DIDS) and 9-anthracenecarboxylic acid (9-AC), an extracellular Ca^{2+} chelator 1,2-bis(2-

aminophenoxy)ethane-*N,N,N′,N′*-tetraacetic acid (BAPTA), and Ca^{2+} channel blockers,

nifedipine and La^{3+} (Figure 4C–4F; Mimata et al., 2022a). These results suggest anion channels

and Ca^{2+} signaling are essential for malate-induced stomatal closure in *V. vinifera* and *A.*

thaliana, consistent with the results for $[Ca^{2+}]_{cyt}$ and SLAC1 activity (Figure 2A, 2D and 3).

Malate stimulates Ca^{2+} signaling *via* second messengers including cADPR, cAMP, and IP₃

To further elucidate the malate-induced stomatal closure, the role of second messengers relevant

to Ca^{2+} signaling, including cADPR, cAMP, IP₃, ROS, NO, cGMP, NAADP, and PIP₃ (see

Introduction), was investigated by pharmacological approach. Nicotinamide, alloxan,

neomycin, and salicylhydroxamic acid (SHAM), which are inhibitors of cADPR, cAMP, IP₃

and peroxidase-catalyzed ROS production, respectively, completely abolished the malate-

induced stomatal closure in *V. vinifera* and *A. thaliana* (Figure 4C, 4D, 4F and 5A–C; Mimata

et al., 2022b). On the other hand, N-nitro-L-arginine methyl ester (L-NAME), 2-(4-

carboxyphenyl)-4,4,5,5-tetramethylimidazoline-1-oxyl-3-oxide (cPTIO), LY83583, Ned 19,

and wortmannin, which are an NO synthetase inhibitor, an NO scavenger, a guanylate cyclase

inhibitor, an antagonist of NAADP, and an inhibitor of PIP_3 production, respectively, have little

inhibitory effect on malate-induced stomatal closure (Supplementary Figure 5). Ca^{2+} imaging

experiments further showed that nicotinamide, alloxan, and neomycin but not SHAM inhibited

malate-induced $[Ca^{2+}]_{cyt}$ oscillations in Arabidopsis (Figure 5D and 5E). These results suggest

that Ca^{2+} signaling mediated by cADPR, cAMP, and IP_3 is required for malate-induced stomatal

closure.

G-proteins are master regulators in malate signaling

Since the generation of cADPR, cAMP, IP_3, and ROS is regulated by G-proteins (Zhang et al.,

2011; Jin et al., 2013), their roles in the stomatal response to malate were investigated with Gα

inhibitors YM-254890 and suramin, and Gβγ inhibitors gallein, guanosine 5′-O-(2-

thiodiphosphate) (GDPβS), and GRK2i. While YM-254890 did not suppress the malate-

induced stomatal closure, all other inhibitors completely abolished the malate effect in *V.*

vinifera and *A. thaliana* (Supplemental Figure 5E and 5F; Figure 6A and 6B). Furthermore,

stomata of *gpa1* and *agb1*, loss-of-function mutants of Arabidopsis canonical Gα subunit and

Gβ subunit, were insensitive to malate (Figure 6C). GDPβS also completely suppressed malate-

induced ROS production (Figure 6D). These data together suggest that malate signal is

transduced by G-proteins in guard cells.

Discussion

Drought has severe impacts on agricultural crops and results in metabolite fluctuations in plants.

In this study, we identified 436 upregulated and 80 downregulated metabolites in grapevine

leaves in response to dehydration stress (Figure 1C). These included modulators of stomatal

movements, such as ABA, phaseic acid, indole-3-acetic acid, jasmonic acid, 5-aminolevulinic

acid, γ-aminobutyric acid, adenosine-3'-5'-diphosphate, malic acid, spermidine, resveratrol,

cinnamic acid, gallic acid, and ascorbic acid (Supplemental Data 1). This study specifically

focused on the effect of the primary metabolite and clarified the mechanism underlying stomatal

closure triggered by a major TCA cycle metabolite malate.

Specific guard cell responses triggered by TCA cycle metabolites

TCA cycle metabolites, as carboxylic acids, elicit distinct guard cell responses, including

increases in $[Ca^{2+}]_{cyt}$, activation of SLAC1, and stomatal closure (Figure 2–4). The limited

specificity of the Ca^{2+} response to TCA cycle metabolites (Figure 2), compared to SLAC1

activation (Figure 3), suggests that SLAC1 activation does not necessarily lead to increases in

$[Ca^{2+}]_{cyt}$, and a carboxylate receptor independent from SLAC1 may exist. Despite both α-

ketoglutarate and oxaloacetate are α-keto acids, only α-ketoglutarate, with a carbon chain length

similar to glutamate, was effective in inducing elevation of $[Ca^{2+}]_{cyt}$ (Figure 2A and 2D),

indicating the importance of carbon chain length in perception. Dicarboxylate glutamate

induces $[Ca^{2+}]_{cyt}$ elevation in guard cells through a pathway dependent on glutamate receptor-

like channels (GLRs), while malate signaling is independently of GLRs (Mimata et al., 2022a).

Therefore, it is plausible that α-ketoglutarate is recognized by GLRs.

Although fumarate and α-ketoglutarate caused Ca^{2+} elevation and SLAC1 activation (Figure

2A, 2D and 3), they failed to close the stomata (Figure 4A and 4B), suggesting that Ca^{2+}

elevation and SLAC1 activation alone are insufficient for stomatal closure. It remains unclear

why only malate induce stomatal closure. Additionally, we observed enhanced ROS production

with fumarate and α-ketoglutarate, similar to malate (Supplemental Figure 6A). Therefore,

unknown factors specific to malate, rather than fumarate and α-ketoglutarate, require further

elucidation in future studies.

Malate plays a key role in stress responses

223 In a related study, the relative abundance of malate increased 6.78-fold in Arabidopsis aerial

224 parts after 10 hours of dehydration stress (Urano et al., 2009). Malate was observed to be

225 secreted during stomatal closure (Van Kirk and Raschke, 1978), with its content in guard cells

226 decreasing after ABA treatment (Kondo and Maruta, 1987; Jin et al., 2013). Notably, the malate

exporter AtALMT12, mainly expressed in guard cells and localized at the plasma membrane, is activated through the ABA signaling pathway (Sasaki et al., 2010; Meyer et al., 2010), and loss-of-function mutation of AtALMT12 have been shown to increase malate content in leaves and result in malate insensitivity (Medeiros et al., 2016; Sasaki et al., 2022). These findings suggest that malate accumulates in leaves under water deficit conditions and is secreted into the apoplast in response to ABA. Malate was the most potent TCA cycle metabolites which modulates stomatal aperture (Figure 4A and 4B). Feeding malate through the petiole decreases stomatal aperture and stomatal conductance (Patonnier et al., 1999; Malcheska et al., 2017; Rasulov et al., 2018). Therefore, its accumulation under dry conditions induces stomatal closure, minimizing water loss and facilitating adaptation to desiccation. Malate is not only closely related to drought stress but is also implicated in stomatal immunity. Some carboxylic acids associated with the TCA cycle, including malate, are detected in the apoplastic fluid from cells treated with an elicitor just before apoplastic oxidative burst (Bolwell et al., 2002). Blockers of malate-permeable channels repress antibacterial ROS production and pathogen resistance (Colcombet et al., 2009). Indeed, malate induced ROS production (Figure 6D; Mimata et al., 2022b). These findings highlight malate as a promising candidate for controlling disease resistance in plants.

Malate signaling is mediated by a specific set of second messengers

Stomatal measurements and Ca^{2+} imaging using inhibitors suggest the involvement of second messengers cADPR, cAMP, and IP_3 in malate signaling (Figure 4C, 4D, 4F, 5A, 5D and 5E). Malate-induced stomatal closure was also impaired by the inhibitor of peroxidase (Figure 5B and 5C). Malate promotes ROS production by peroxidases (Mimata et al., 2022b), and ROS activate plasma membrane Ca^{2+} channels and increase $[Ca^{2+}]_{cyt}$ (Pei et al., 2000) in *A. thaliana*. Thus, we hypothesized that malate accelerates Ca^{2+} influx by promoting ROS production. Contrary to our expectation, the inhibition of ROS production did not impair malate-induced Ca^{2+} responses (Figure 5D and 5E). Moreover, blocking Ca^{2+} influx did not affect ROS production (Supplemental Figure 6B), indicating that ROS production is independent of Ca^{2+} signaling in malate-induced stomatal closure.

Malate signaling is transduced by G-protein signaling cascades

Pharmacological and reverse genetics experiments suggest that G-proteins are master regulators in malate signaling (Figure 6). Based on the current results, we propose a model to summarize malate signaling in guard cells (Figure 6E). The external signal of malate is transmitted by G-proteins, which modulate the generation of second messengers, including cAMP, cADPR and IP$_3$. This cascade leads to increases in $[Ca^{2+}]_{cyt}$, resulting in the phosphorylation of SLAC1 by CDPKs. Additionally, G-proteins simultaneously modulate ROS production, which contributes to SLAC1 phosphorylation. Extracellular malate directly promotes anion transport through phosphorylated SLAC1, leading to a decrease in turgor pressure and subsequent stomatal closure.

In mammals, carboxylates are sensed by G-protein-coupled receptors (GPCRs). Succinate is released into the extracellular space from stimulated macrophages and injured tissues, with local concentrations reaching millimolar levels (Chouchani et al, 2014; Littlewood-Evans et al., 2016). Extracellular succinate activates GPCRs, stimulating IP$_3$ formation, inhibiting cAMP production, and increasing $[Ca^{2+}]_{cyt}$ *via* the G-protein signaling pathway (He et al., 2004; Trauelsen et al., 2021). Similarly, malate is perceived by the succinate receptor and causes rapid increases in intracellular $[Ca^{2+}]$ and IP$_3$ accumulation (Trauelsen et al., 2017). Since plant GPCRs differ significantly from those in animals, and GPCRs with guanine nucleotide exchange factor activity have not yet been identified, it is challenging to identify malate receptors. Nevertheless, based on the shared signaling components, we propose that TCA cycle metabolites are common stress-responsive signal molecules mediated by G-protein-dependent signaling cascades in both animal and plant kingdoms.

Materials and methods

Plants and growth conditions

Grapevine (*V. vinifera* L. cv. Chardonnay) was grown on a soil mixture of 1:1 = soil: vermiculite (v/v) in a growth room at 24 °C and 80% relative humidity under 16 h-light/8 h-dark

photoperiod with a photon flux density of 100 μmol m^{-2} s^{-1}. Arabidopsis (*A. thaliana* L. ecotype

Colombia-0) was grown on a soil mixture of 1:1 = soil: vermiculite (v/v) in a growth chamber

at 21 °C and 60% relative humidity under 16 h-light/8 h-dark photoperiod with a photon flux

density of 120 μmol m^{-2} s^{-1}. The T-DNA insertion lines, *gpa1-4* (SALK_001846) and *agb1*

(SALK_077086C), were obtained from Arashare (https://www.arashare.cn/index/).

Tomato (*Solanum lycopersicum* L. cv. Micro-Tom) was used. Tomato was grown on a soil

mixture of 1:1 = soil: vermiculite (v/v) in a glass greenhouse at 15 °C and 60% relative humidity.

Chili pepper (*Capsicum chinense* L. cv. Trinidad Moruga scorpion) was used. Chili pepper was

grown on a soil mixture of 1:1 = soil: vermiculite (v/v) in a glass greenhouse at 22 °C and 50%

relative humidity.

Dehydration stress treatment

293 Fully expanded leaves from 1 to 2-month-old grapevine plants were randomly detached.

294 Dehydration was performed as described previously (Urano et al., 2009) with a few

295 modifications. The detached leaves were exposed to dehydration stress on the paper at 26℃

296 and ambient humidity under light. At indicated time, the leaves were frozen by liquid nitrogen.

Sample preparation for LC-MS

The sample preparation, extract analysis, metabolite identification and quantification were

performed at Wuhan MetWare Biotechnology Co., Ltd (www.metware.cn) following their

standard procedures. The samples were freeze-dried in a lyophilizer (Scientz-100F; Scientz,

Zhejiang, China) and then homogenized (30 Hz, 1.5 min) into powder using a grinder (MM

400; Retsch, Dusseldorf, Germany). Next, 1200 μL of -20 °C pre-cooled 70% methanolic

aqueous internal standard extract added to 50 mg of sample powder. The sample was vortexed

once every 30 min for 30 s, for a total of 6 times. After centrifugation at 12000rpm for 3min,

the supernatant was aspirated, and the sample was filtered through a microporous membrane

(0.22 μm pore size) and stored in the injection vial for UPLC-MS/MS analysis.

Metabolite analysis using LC-MS

The sample extracts were analyzed using an UPLC-ESI-MS/MS system (UPLC, ExionLC™

AD: SCIEX, MA, USA; MS, Applied Biosystems 4500 Q TRAP: SCIEX). The analytical

conditions were as follows, UPLC: column, Agilent SB-C18 (1.8 μm, 2.1 mm * 100 mm); The

mobile phase was consisted of solvent A, pure water with 0.1% formic acid, and solvent B,

acetonitrile with 0.1% formic acid. Sample measurements were performed with a gradient

program that employed the starting conditions of 95% A, 5% B. Within 9 min, a linear gradient

to 5% A, 95% B was programmed, and a composition of 5% A, 95% B was kept for 1 min.

Subsequently, a composition of 95% A, 5.0% B was adjusted within 1.1 min and kept for 2.9

min. The flow velocity was set as 0.35 mL per minute; The column oven was set to 40°C; The

injection volume was 4 μL. The effluent was alternatively connected to an ESI-triple

quadrupole-linear ion trap (QTRAP)-MS.

The ESI source operation parameters were as follows: source temperature 550°C; ion spray

voltage (IS) 5500 V (positive ion mode)/-4500 V (negative ion mode); ion source gas I (GSI),

gas II (GSII), curtain gas (CUR) were set at 50, 60, and 25 psi, respectively; the collision-

activated dissociation (CAD) was high. QQQ scans were acquired as MRM experiments with

collision gas (nitrogen) set to medium. DP (declustering potential) and CE (collision energy)

for individual MRM transitions was done with further DP and CE optimization. A specific set

of MRM transitions were monitored for each period according to the metabolites eluted within

this period.

Sample preparation for GC-MS

The leaves subjected dehydration stress were ground to a powder in liquid nitrogen. 500 mg (1

mL) of the powder was transferred immediately to a 20 mL head-space vial (Agilent, CA, USA),

containing NaCl saturated solution, to inhibit any enzyme reaction. The vials were sealed using

crimp-top caps with TFE-silicone headspace septa (Agilent). At the time of SPME analysis,

each vial was placed in 60 °C for 5 min, then a 120 μm DVB/CWR/PDMS fiber (Agilent) was

exposed to the headspace of the sample for 15 min at 60 °C.

Metabolite analysis using GC-MS

After sampling, desorption of the VOCs from the fiber coating was carried out in the injection
port of the GC apparatus (Model 8890; Agilent) at 250°C for 5 min in the splitless mode. The
identification and quantification of VOCs was carried out using an Agilent Model 8890 GC and
a 7000D mass spectrometer (Agilent), equipped with a 30 m × 0.25 mm × 0.25 μm DB-5MS
(5% phenyl-polymethylsiloxane) capillary column. Helium was used as the carrier gas at a
linear velocity of 1.2 mL/min. The injector temperature was kept at 250°C and the detector at
280°C. The oven temperature was programmed from 40°C (3.5 min), increasing at 10°C/min
to 100°C, at 7°C/min to 180°C, at 25°C/min to 280°C, hold for 5 min. Mass spectra was
recorded in electron impact (EI) ionization mode at 70 eV. The quadrupole mass detector, ion
source and transfer line temperatures were set, respectively, at 150, 230 and 280°C. The MS
was selected ion monitoring (SIM) mode was used for the identification and quantification of
analytes.

Data analysis of the non-target metabolome

Relative metabolite abundances were calculated by the peak areas. Unsupervised PCA was
performed by statistics function prcomp within R (www.r-project.org). The relative contents of
all differential metabolites were processed by UV (unit variance scaling) followed by K-Means
cluster analysis. Identified metabolites were annotated using KEGG Compound database
(http://www.kegg.jp/kegg/compound/). Pathways with significantly regulated metabolites
mapped to were then fed into metabolite sets enrichment analysis (MSEA), their significance
was determined by hypergeometric test's p-values. Upregulated metabolites are defined VIP >
1, $\log_2$ (Fold change) $\geq$ 1 and P-value > 0.05, downregulated metabolites are defined VIP > 1,
$\log_2$ (Fold change) $\leq$ −1 and P-value > 0.05. P-value was calculated by Welch's t-test.

$[Ca^{2+}]_{cyt}$ imaging

Wild-type Arabidopsis plants expressing Yellow Cameleon 3.6 were used to measure $[Ca^{2+}]_{cyt}$
in guard cells as described previously (Mimata et al., 2022a). The abaxial side of an excised
rosette leaf was gently attached to a glass slide with a medical adhesive (stock no. 7730;
Hollister, IL, USA) and then mesophyll tissues were whittled away with a razor blade to keep
the abaxial epidermis intact on the slide. The remaining abaxial epidermis was immersed in

stomatal assay solution, comprising 5 mM KCl, 50 μM $CaCl_2$ and 10 mM MES/Tris (pH 5.6), in the light for 2 h to induce stomatal opening. The epidermis was treated with 10 mM organic acid in stomatal assay solution at 3 min after monitoring and with 10 mM $CaCl_2$ at 18 or 28 min after the measurement to check responsiveness. Inhibitors were added 5 min before starting imaging. The stock solution of organic acids was dissolved in stomatal assay solution and adjusted to a pH of 5.6 with Tris.

The images were acquired under a fluorescence microscope (ECLIPSE Ti2-E; NIKON). Excitation light was provided by a mercury arc lamp and a 436 nm filter (ET436/20x, Chroma Technology Corporation, VT, USA). Emission of the CFP was measured at 480 nm filter (ET480/40m, Chroma Technology Corporation) and of the YFP at 535 nm filter (ET535/30m, Chroma Technology Corporation) using a CMOS camera (ORCA-Fusion BT Digital CMOS camera C15440; HAMAMATSU, Shizuoka, Japan). Images were taken every 5 s.

Cloning and cRNA synthesis

All constructs were cloned into the oocyte expression vector pNB1u (Nour-Eldin et al., 2006) by ClonExpress II One Step Cloning Kit (Vazyme Biotech, Nanjing, China). The site-directed mutants were generated by FastCloning (Li et al., 2011). VvSLAC1 (LOC100244459) cDNA from *V. vinifera* was used for cloning, and all constructs were verified by sequencing. Primers used for cloning and site-directed mutagenesis are listed in Supplemental Table 2. cRNA was prepared using an mMESSAGE mMACHINE ™ T7 Transcription Kit (Thermo Fisher Scientific, MA, USA).

Two-electrode voltage-clamp

Xenopus laevis oocytes were injected with 50 nL cRNA (each 10 ng) and incubated in ND96 buffer at 18°C for a few days before voltage-clamp recordings (Mimata et al., 2022a). The bath solution contained 1 mM Mg-gluconate, 1 mM Ca-gluconate and 1 mM $LaCl_3$ ± 10 mM organic acid buffered with 10 mM MES/Tris to adjust the pH to 5.6. Osmolality was adjusted to 220 mM using D-sorbitol. The voltage pulse was commanded to clamp the membrane potential from +60 mV to −120 mV in ramp for 1.8 seconds or from +60 to −160 mV in 20 mV decrements in

390 step for 2.5 seconds with a holding potential of 0 mV. Voltage-clamp recordings for oocytes

391 were performed using an Axoclamp 900A amplifier (Molecular Devices, CA, USA), data were

392 acquired using a Digidata 1550B system (Molecular Devices) and analyzed using pCLAMP

393 11.2 software (Molecular Devices).

Measurement of stomatal aperture

Stomatal apertures were measured as described previously (Ye et al., 2020) with modifications.

Leaf discs (4 mm in diameter) obtained from fully expanded leaves were placed abaxial side

down on stomatal assay solution. The discs were exposed to light for 2 h to induce stomatal

opening and subsequently treated with 10 mM organic acid in stomatal assay solution for an

additional 2 h. Inhibitors were added 5 min before malate treatment. The abaxial epidermis was

captured under optical microscopes (ECLIPSE Ts-2R, ECLIPSE Ti2-E and ECLIPSE Ci;

NIKON, Tokyo, Japan) using NIS ELEMENTS software (NIKON). Stomatal apertures were

quantified using IMAGEJ software (NIH). We measured 30 stomatal apertures from a leaf disc

to calculate an average. This measurement was repeated four times using different plants, and

the overall average was calculated.

Measurement of ROS production

ROS production in guard cells was analyzed using the fluorescent dye 2',7'-

dihydrodichlorofluorescein diacetate (H_2DCF-DA) as described previously (Mimata et al.,

2022b) with modifications. The abaxial side of an excised rosette leaf was gently attached to a

glass slide with a medical adhesive and then mesophyll tissues were whittled away with a razor

blade. The remaining abaxial epidermis was immersed in stomatal assay solution in the light

for 2 h to induce stomatal opening. A total of 50 μM H_2DCF-DA was added to the stomatal

assay solution and the epidermal tissues were incubated in the dark for 30 min. After the dye

loading, the epidermal tissues were gently rinsed with stomatal assay solution. The epidermis

was treated with 10 mM organic acid ± inhibitor in stomatal assay solution. After the 30 min

incubation, fluorescent signals were captured using the fluorescence microscope with 480 ± 15

nm/ 535 ± 23 nm excitation/emission filters. We measured 30 guard cells from an epidermis to

calculate an average. This measurement was repeated three times using different plants, and the

overall average was calculated.

Acknowledgements

We thank Dr. Shangwei Zhong and Dr. Xu Wang (Peking University Institute of Advanced
Agricultural Sciences) for providing plant materials; Dr. Hussam Hassan Nour-Eldin
(University of Copenhagen) for providing pNB1u vector; Dr. Yoshiyuki Murata and Dr.
Shintaro Munemasa (Okamaya University) for providing Arabidopsis cDNA plasmid
constructs; Dongyue Wang (Peking University Institute of Advanced Agricultural Sciences) for
helping cloning. W.Y. laboratory is supported by Taishan Scholars Program of Shandong
Province and Shandong Provincial Science and Technology Innovation Fund.

Conflict of interests

428 The authors declare no competing interests.

Contributions

430 Y.M. and W.Y. designed the experiments. Y.M. analyzed all data and performed all experiments
431 except metabolomics. R.G. measured stomatal aperture and ROS production. X.P. conducted
432 metabolomics and G.Q. interpreted the data. Y.M. and W.Y. wrote the manuscript.

Data availability

434 The data underlying this article are available in the article and in its online supplementary
435 material.

References

Bolwell, G.P., Bindschedler, L.V., Blee, K.A., Butt, V.S., Davies, D.R., Gardner, S.L., Gerrish,
C., Minibayeva, F. (2002) The apoplastic oxidative burst in response to biotic stress in plants:
a three-component system. *J. Exp. Bot.* 10.1093/jexbot/53.372.1367.

Brandt, B., Munemasa, S., Wang, C., Nguyen, D., Yong, T., Yang, P.G., Poretsky, E., Belknap,
T.F., Waadt, R., Alemán F., et al. (2015) Calcium specificity signaling mechanisms in abscisic

acid signal transduction in Arabidopsis guard cells. *Elife* 10.7554/eLife.03599.

Chen, Y., Hu, L., Punta, M., Bruni, R., Hillerich, B., Kloss, B., Rost, B., Love, J., Siegelbaum,

S.A., Hendrickson, W.A. (2010) Homologue structure of the SLAC1 anion channel for closing

stomata in leaves. *Nature* 10.1038/nature09487.

Chouchani, E.T., Pell, V.R., Gaude, E., Aksentijević, D., Sundier, S.Y., Robb, E.L., Logan, A.,

Nadtochiy, S.M., Ord, E.N.J., Smith, A.C., et al. (2014) Ischaemic accumulation of succinate

controls reperfusion injury through mitochondrial ROS. *Nature* 10.1038/nature13909.

Colcombet, J., Mathieu, Y., Peyronnet, R., Agier, N., Lelièvre, F., Barbier-Brygoo, H., Frachisse,

J.-M. (2009) R-type anion channel activation is an essential step for ROS-dependent innate

immune response in *Arabidopsis* suspension cells. *Funct. Plant Biol.* 10.1071/FP09096.

Delhaize, E., Ryan, P.R., Randall, P.J. (1993) Aluminum Tolerance in Wheat (*Triticum aestivum*

L.) (II. Aluminum-Stimulated Excretion of Malic Acid from Root Apices). *Plant Physiol.*

10.1104/pp.103.3.695.

Fan, L.-M., Zhang, W., Chen, J.-G., Taylor, J.P., Jones, A.M., Assmann, S.M. (2008) Abscisic

acid regulation of guard-cell K^+ and anion channels in Gβ- and RGS-deficient *Arabidopsis* lines.

Proc. Natl. Acad. Sci. USA. 10.1073/pnas.0800980105.

Felle, H. (1988) Auxin causes oscillations of cytosolic free calcium and pH in *Zea mays*

coleoptiles. *Planta.* 10.1007/BF00634478.

Fernie, A.R., Martinoia, E. (2009) Malate. Jack of all trades or master of a few? *Phytochemistry*

10.1016/j.phytochem.2009.04.023.

Gabriel, R., Kesselmeier, J. (1999) Apoplastic solute concentrations of organic acids and

mineral nutrients in the leaves of several fagaceae. *Plant Cell Physiol.*

10.1093/oxfordjournals.pcp.a029583.

Garcia-Mata, C., Gay, R., Sokolovski, S., Hills, A., Lamattina, L., Blatt, M.R. (2003) Nitric

oxide regulates K^+ and Cl^- channels in guard cells through a subset of abscisic acid-evoked

467 signaling pathways. *Proc. Natl. Acad. Sci. USA.* 10.1073/pnas.1434381100.

468 Gilroy, S., Read, N.D., Trewavas, A.J. (1990) Elevation of cytoplasmic calcium by caged

469 calcium or caged inositol triphosphate initiates stomatal closure. *Nature* 10.1038/346769a0.

470 Hamilton, D.W.A., Hills, A., Köhler, B., Blatt, M.R. (2000) Ca^{2+} channels at the plasma

471 membrane of stomatal guard cells are activated by hyperpolarization and abscisic acid. *Proc.*

472 *Natl. Acad. Sci. USA.* 10.1073/pnas.080068897.

473 He, W., Miao, F.J.P., Lin, D.C.H., Schwandner, R.T., Wang, Z., Gao, J., Chen, J.-L., Tian, H.,

474 Ling, L. (2004) Citric acid cycle intermediates as ligands for orphan G-protein-coupled

475 receptors. *Nature* 10.1038/nature02488.

476 Hedrich, R., Marten, I., Lohse, G., Dietrich, P., Winter, H., Lohaus, G., Heldt, H.-W. (1994)

477 Malate-sensitive anion channels enable guard cells to sense changes in the ambient CO_2

478 concentration. *Plant J.* 10.1046/j.1365-313X.1994.6050741.x.

479 Hedrich, R., Geiger, D. (2017) Biology of SLAC1-type anion channels – from nutrient uptake

480 to stomatal closure. *New Phytol.* 10.1111/nph.14685.

481 Huang, S., Shen, L., Roelfsema, M.R.G., Becker, D., Hedrich, R. (2023) Light-gated

482 channelrhodopsin sparks proton-induced calcium release in guard cells. *Science*

483 10.1126/science.adj9696.

484 Jin, X., Wang, R.-S., Zhu, M., Jeon, B.W., Albert, R., Chen, S., Assmann, S.M. (2013) Abscisic

485 acid-responsive guard cell metabolomes of *Arabidopsis* wild-type and *gpa1* G-protein mutants.

486 *Plant Cell* 10.1105/tpc.113.119800.

487 Kondo, N., Maruta, I. (1987) Abscisic acid-induced stomatal closure in *Vicia faba* epidermal

488 strip. Excretion of solutes from guard cells and increase in elastic modulus of guard cell wall.

489 *Plant Cell Physiol.* 10.1093/oxfordjournals.pcp.a077303.

490 Krzak, G., Willis, C.M., Smith, J.A., Pluchino, S., Peruzzotti-Jametti, L. (2021) Succinate

491 Receptor 1: An Emerging Regulator of Myeloid Cell Function in Inflammation. *Trends*

492   *Immunol.* 10.1016/j.it.2020.11.004.

493   Leckie, C.P.,McAinsh, M.R., Allen, G.J., Sanders, D., Hetherington, A.M. (1998) Abscisic acid-
494   induced stomatal closure mediated by cyclic ADP-ribose. *Proc. Natl. Acad. Sci. USA.*
495   10.1073/pnas.95.26.15837.

496   Lemtiri-Chlieh, F., Berkowitz, G.A. (2004) Cyclic adenosine monophosphate regulates calcium
497   channels in the plasma membrane of Arabidopsis leaf guard and mesophyll cells. *J. Biol. Chem.*
498   10.1074/jbc.M400311200.

499   Lepetit, M., Brouquisse, R. (2023) Control of the rhizobium–legume symbiosis by the plant
500   nitrogen demand is tightly integrated at the whole plant level and requires inter-organ systemic
501   signaling. *Front. Plant Sci*. 10.3389/fpls.2023.1114840.

502   Li, C., Wen, A., Shen, B., Lu, J., Huang, Y., Chang, Y. (2011) FastCloning: a highly simplified,
503   purification-free, sequence- and ligation-independent PCR cloning method. *BMC Biotechnol*.
504   10.1186/1472-6750-11-92.

505   Littlewood-Evans, A., Sarret, S., Apfel, V., Loesle, P., Dawson, J., Zhang, J., Muller, A., Tigani,
506   B., Kneuer, R., Patel, S., et al. (2016) GPR91 senses extracellular succinate released from
507   inflammatory macrophages and exacerbates rheumatoid arthritis. *J. Exp. Med.*
508   10.1084/jem.20160061.

509   Luan, S., Wang, C. (2021) Calcium Signaling Mechanisms Across Kingdoms. *Annu. Rev. Cell*
510   *Dev. Biol.* 10.1146/annurev-cellbio-120219-035210.

511   Malcheska, F., Ahmad, A., Batool, S., Müller, H.M., Ludwig-Müller, J., Kreuzwieser, J.,
512   Randewig, D., Hänsch, R., Mendel, R.R., Hell, R., et al. (2017) Drought-Enhanced Xylem Sap
513   Sulfate Closes Stomata by Affecting ALMT12 and Guard Cell ABA Synthesis. *Plant Physiol.*
514   10.1104/pp.16.01784.

515   Maierhofer, T., Diekmann, M., Offenborn, J.N., Lind, C., Bauer, H., Hashimoto, K., Al-Rasheid,
516   K.A.S., Luan, S., Kudla, J., Geiger, D., et al. (2014) Site- and kinase-specific phosphorylation-
517   mediated activation of SLAC1, a guard cell anion channel stimulated by abscisic acid. *Sci.*

Signal. 10.1126/scisignal.2005703.

Medeiros, D.B., Martins, S.C.V., Cavalcanti, J.H.F., Daloso, D.M., Martinoia, E., Nunes-Nesi,

A., DaMatta, F.M., Fernie, A.R., Araújo, W.L. (2016) Enhanced Photosynthesis and Growth in

atquac1 Knockout Mutants Are Due to Altered Organic Acid Accumulation and an Increase in

Both Stomatal and Mesophyll Conductance. *Plant Physiol.* 10.1104/pp.15.01053.

Medeiros, D.B., Barros, K.A., Barros, J.A.S., Omena-Garcia, R.P., Arrivault, S., Sanglard,

L.M.V.P., Detmann, K.C., Silva, W.B., Daloso, D.M., DaMatta, F.M., et al. (2017) Impaired

Malate and Fumarate Accumulation Due to the Mutation of the Tonoplast Dicarboxylate

Transporter Has Little Effects on Stomatal Behavior. *Plant Physiol.* 10.1104/pp.17.00971.

Meyer, S., Mumm, P., Imes, D., Endler, A., Weder, B., Al-Rasheid, K.A.S., Geiger, D., Marten,

I., Martinoia, E., Hedrich, R. (2010) *At*ALMT12 represents an R-type anion channel required

for stomatal movement in Arabidopsis guard cells. *Plant J.* 10.1111/j.1365-313X.2010.04302.x.

Mimata, Y., Munemasa, S., Nakamura, T., Nakamura, Y., Murata, Y. (2022a) Extracellular

malate induces stomatal closure via direct activation of guard-cell anion channel SLAC1 and

stimulation of Ca^{2+} signalling. *New Phytol.* 10.1111/nph.18400.

Mimata, Y., Munemasa, S., Akter, F., Jahan, I., Nakamura, T., Nakamura, Y., Murata, Y. (2022b)

Malate induces stomatal closure via a receptor-like kinase GHR1- and reactive oxygen species-

dependent   pathway   in   *Arabidopsis   thaliana*.   *Biosci.   Biotechnol.   Biochem.*

10.1093/bbb/zbac122.

Murata, Y., Mori, I.C., Munemasa, S. (2015) Diverse Stomatal Signaling and the Signal

Integration Mechanism. *Annu. Rev. Plant Biol.* 10.1146/annurev-arplant-043014-114707.

Nagai, T., Yamada, S., Tominaga, T., Ichikawa, M., Miyawaki, A. (2004) Expanded dynamic

range of fluorescent indicators for Ca^{2+} by circularly permuted yellow fluorescent proteins.

Proc. Natl. Acad. Sci. USA. 10.1073/pnas.0400417101.

Navazio, L., Bewell, M.A., Siddiqua, A., Dickinson, G.D., Galione, A., Sanders, D. (2000)

Calcium release from the endoplasmic reticulum of higher plants elicited by the NADP

metabolite nicotinic acid adenine dinucleotide phosphate. *Proc. Natl. Acad. Sci. USA.*

10.1073/pnas.140217897.

Negi, J., Matsuda, O., Nagasawa, T., Oba, Y., Takahashi, H., Kawai-Yamada, M., Uchimiya, H.,

Hashimoto, M., Iba, K. (2008) CO_2 regulator SLAC1 and its homologues are essential for anion

homeostasis in plant cells. *Nature* 10.1038/nature06720.

Neill, S., Barros, R., Bright, J., Desikan, R., Hancock, J., Harrison, J., Morris, P., Ribeiro, D.,

Wilson, I. (2008) Nitric oxide, stomatal closure, and abiotic stress. *J. Exp. Bot.*

10.1093/jxb/erm293.

Nour-Eldin, H.H., Hansen, B.G., Nørholm, M.H.H., Jensen, J.K., Halkier, B.A. (2006)

Advancing uracil-excision based cloning towards an ideal technique for cloning PCR fragments.

Nucleic Acids Res. 10.1093/nar/gkl635.

Pandey S. (2020) Plant receptor-like kinase signaling through heterotrimeric G-proteins. *J. Exp.*

Bot. 10.1093/jxb/eraa016.

Patonnier, M.P., Peltier, J.P., Marigo, G. (1999) Drought-induced increase in xylem malate and

mannitol concentrations and closure of *Fraxinus excelsior* L. stomata. *J. Exp. Bot.*

10.1093/jxb/50.336.1223.

Pei, Z.M., Murata, Y., Benning, G., Thomine, S., Klüsener, B., Allen, G.J., Grill, E., Schroeder,

J.I. (2000) Calcium channels activated by hydrogen peroxide mediate abscisic acid signalling

in guard cells. *Nature* 10.1038/35021067.

Pires, M.V., Júnior, A.A.P., Medeiros, D.B., Daloso, D.M., Pham, P.A., Barros, K.A., Engqvist,

M.K.M., Florian, A., Krahnert, I., Maurino, V.G., et al. (2016) The influence of alternative

pathways of respiration that utilize branched-chain amino acids following water shortage in

Arabidopsis. *Plant Cell Environ.* 10.1111/pce.12682.

Prokic, L., Jovanovic, Z., McAinsh, M.R., Vucinic, Z., Stikic, R. (2006) Species-dependent

changes in stomatal sensitivity to abscisic acid mediated by external pH. *J. Exp. Bot.*

10.1093/jxb/erj057.

Rasulov, B., Talts, E., Bichele, I., Niinemets, Ü. (2018) Evidence That Isoprene Emission Is

Not Limited by Cytosolic Metabolites. Exogenous Malate Does Not Invert the Reverse

Sensitivity of Isoprene Emission to High [CO_2]. *Plant Physiol.* 10.1104/pp.17.01463.

Sasaki, T., Mori, I.C., Furuichi, T., Munemasa, S., Toyooka, K., Matsuoka, K., Murata, Y.,

Yamamoto, Y. (2010) Closing Plant Stomata Requires a Homolog of an Aluminum-Activated

Malate Transporter. *Plant Cell Physiol.* 10.1093/pcp/pcq016.

Sasaki, T., Ariyoshi, M., Yamamoto, Y., Mori, I.C. (2022) Functional roles of ALMT-type anion

channels in malate-induced stomatal closure in tomato and Arabidopsis. *Plant Cell Environ.*

10.1111/pce.14373.

Schroeder, J.I., Allen, G.J., Hugouvieux, V., Kwak, J.M., Waner, D. (2001) GUARD CELL

SIGNAL   TRANSDUCTION.   *Annu.   Rev.   Plant   Physiol.   Plant   Mol.   Biol.*

10.1146/annurev.arplant.52.1.627

Schwartz, A. (1985) Role of Ca^{2+} and EGTA on Stomatal Movements in *Commelina communis*

L.. *Plant Physiol.* 10.1104/pp.79.4.1003.

Shen, J.-L., Li, C.-L., Wang, M., He, L.-L., Lin, M.-Y., Chen, D.-H., Zhang, W. (2017)

Mitochondrial pyruvate carrier 1 mediates abscisic acid-regulated stomatal closure and the

drought response by affecting cellular pyruvate content in *Arabidopsis thaliana. BMC Plant*

Biol. 10.1186/s12870-017-1175-3.

Sweetman, C., Deluc, L.G., Cramer, G.R., Ford, C.M., Soole, K.L. (2009) Regulation of malate

metabolism   in   grape   berry   and   other   developing   fruits.   *Phytochemistry*

10.1016/j.phytochem.2009.08.006.

Trauelsen, M., Ulven, E.R., Hjorth, S.A., Brvar, M., Monaco, C., Frimurer, T.M., Schwartz,

T.W. (2017) Receptor structure-based discovery of non-metabolite agonists for the succinate

receptor GPR91. *Mol. Metab*. 10.1016/j.molmet.2017.09.005.

Trauelsen, M., Hiron, T.K., Lin, D., Petersen, J.E., Breton, B., Husted, A.S., Hjorth, S.A., Inoue,

A., Frimurer, T.M., Bouvier, M., et al. (2021) Extracellular succinate hyperpolarizes M2

macrophages through SUCNR1/GPR91-mediated Gq signaling. *Cell Rep*. 597 10.1016/j.celrep.2021.109246.

Urano, K., Maruyama, K., Ogata, Y., Morishita, Y., Takeda, M., Sakurai, N., Suzuki, H., Saito, 599 K., Shibata, D., Kobayashi, M., et al. (2009) Characterization of the ABA-regulated global 600 responses to dehydration in Arabidopsis by metabolomics. *Plant J.* 10.1111/j.1365- 601 313X.2008.03748.x

Vahisalu, T., Kollist, H., Wang, Y.-F., Nishimura, N., Chan, W.-Y., Valerio, G., Lamminmäki, 603 A., Brosché, M., Moldau, H., Desikan, R., et al. (2008) SLAC1 is required for plant guard cell 604 S-type anion channel function in stomatal signalling. *Nature* 10.1038/nature06608.

Van Kirk, C.A., Raschke, K. (1978) Release of malate from epidermal strips during stomatal 606 closure. *Plant Physiol.* 10.1104/pp.61.3.474.

Wada, H., Shackel, K.A., Matthews, M.A. (2008) Fruit ripening in *Vitis vinifera*: apoplastic 608 solute accumulation accounts for pre-veraison turgor loss in berries. *Planta* 10.1007/s00425- 609 008-0707-3.

Wang, X.-Q., Ullah, H., Jones, A.M., Assmann, S.M. (2001) G Protein Regulation of Ion 611 Channels and Abscisic Acid Signaling in *Arabidopsis* Guard Cells. *Science* 612 10.1126/science.1059046.

Wang, Y.-F., Munemasa, S., Nishimura, N., Ren, H.-M., Robert, N., Han, M., Puzõrjova, I., 614 Kollist, H., Lee, S., Mori, I., et al. (2013) Identification of Cyclic GMP-Activated Nonselective 615 Ca^{2+}-Permeable Cation Channels and Associated *CNGC5* and *CNGC6* Genes in Arabidopsis 616 Guard Cells. *Plant Physiol.* 10.1104/pp.113.225045.

Ye, W., Munemasa, S., Shinya, T., Wu, W., Ma, T., Lu, J., Kinoshita, T., Kaku, H., Shibuya, N., 618 Murata, Y. (2020) Stomatal immunity against fungal invasion comprises not only chitin-induced 619 stomatal closure but also chitosan-induced guard cell death. *Proc. Natl. Acad. Sci. USA.* 620 10.1073/pnas.1922319117.

Yu, Y., Chakravorty, D., Assmann, S.M. (2018) The G Protein β-Subunit, AGB1, Interacts with

FERONIA in RALF1-Regulated Stomatal Movement. *Plant Physiol.* 10.1104/pp.17.01277.

Zhang, W., Jeon, B.W., Assmann, S.M. (2011) Heterotrimeric G-protein regulation of ROS

signalling and calcium currents in Arabidopsis guard cells. *J. Exp. Bot.* 10.1093/jxb/erq424.

Figure legends

Figure 1. Metabolome analysis in grapevine leaves during dehydration treatment.

(A) PCA score plot of metabolomic datasets colored by the time of dehydration stress as clusters.

Dots represent biological replicates. Ellipse display 95% confidence regions of each cluster.

(B) Relative ABA levels. Fold changes were normalized to the values of 0 h. Data are the mean

± SE.

(C) Volcano plot for the differential metabolites. Red and green dots mark the metabolites with

significantly increased and decreased level in 0 h versus 24 h, respectively. Upregulated

metabolites are defined VIP > 1, $\log_2$(Fold change) $\geq$ 1 and P-value > 0.05, downregulated

metabolites are defined VIP > 1, $\log_2$(Fold change) $\leq$ −1 and P-value > 0.05. P-value was

calculated by Welch's t-test.

(D) Categorization of the differential metabolites in **(C)**. Upper panel shows upregulated

metabolites and lower panel shows downregulated metabolites.

(E) Relative TCA cycle metabolite levels at different time points. Heat maps represent

$\log_2$(Fold change). Fold changes were normalized to the values of 0 h. Data were obtained from

three independent biological replicates. Abbreviations: F6P, fructose-6-phosphate; F1,6BP,

fructose 1,6-bisphosphate; GADP, glyceraldehyde 3-phosphate; 3PG, 3-phosphoglycerate; PEP,

phosphoenolpyruvate; Pyr, pyruvate; Cit, citrate; *cis*-Aco, *cis*-aconitate; Isocit, isocitrate; Keto,

α-ketoglutarate; Suc, succinate; Fum, fumarate; Mal, malate; Oxal, oxalate; Ser, serine; Trp,

tryptophan; Tyr, tyrosine; Phe, phenylalanine; Val, valine; Leu, leucine; Ile, isoleucine; Thr,

threonine; Asp, aspartate; Asn, asparagine; Met, methionine; Lys, lysine; Glu, glutamate; Gln,

glutamine; His, histidine; Pro, proline; Arg, arginine; GABA, γ-aminobutyrate.

Figure 2. Ca²⁺ response to organic acids in guard cells.

(A and B) Representative traces of fluorescence emission ratios (535/480 nm) in *A. thaliana*

guard cells expressing the Ca^{2+} sensor Yellow Cameleon 3.6. Grey bars indicate the time point

when treatment was applied. The guard cells were treated with organic acids 3 min after the

measurement.

(C and D) Percentage of number of guard cells showing different numbers of transient $[Ca^{2+}]_{cyt}$

increases. An increase in $[Ca^{2+}]_{cyt}$ is defined by an increase in fluorescence ratio by ≥0.1 U from

the baseline ratio. Data were obtained from Mock (25), Isocit (16) for **C**; Mock (51 guard cells),

Suc (31), Fum (28), Mal (44) Oxal (27), Keto (28), Cit (30), *cis*-Aco (29), Ace (33), Pyr (33),

Mal+La³⁺ (21) for **D**. Suc, succinate; Fum, fumarate; Mal, malate; Oxal, oxalate; Keto, α-

ketoglutarate; Cit, citrate; *cis*-Aco, *cis*-aconitate; Ace, acetate; Pyr, pyruvate; Isocit, isocitrate.

P-value was calculated by Fisher's exact test.

Figure 3. SLAC1 activity in the presence of organic acids.

(A and B) Representative whole-cell negative current recordings in Xenopus oocytes

expressing **(A)** VvSLAC1 or **(B)** AtSLAC1 in the presence of external organic acids. The

whole-cell negative current recording in the presence of external malate showed as red trace.

The voltage pulse was commanded to clamp the membrane potential from +60 mV to −120 mV

in ramp for 1.8 seconds with a holding potential of 0 mV.

(C and D) Average steady-state negative currents at −120 mV in Xenopus oocytes expressing

(C) VvSLAC1 or **(D)** AtSLAC1. Data were obtained from six oocytes per condition. Data are

the mean ± SE. Different letters indicate statistical significances within condition based on one-

way ANOVA with Tukey's HSD test, P < 0.05.

Figure 4. Malate-induced stomatal closure is mediated by anion channels and Ca²⁺.

(A and B) Effects of organic acids on stomatal aperture in **(A)** *V. vinifera* or **(B)** *A. thaliana*

leaves. Male means maleate.

(C–F) Effects of inhibitors on malate-induced stomatal closure in **(C and D)** *V. vinifera* or **(E**

and F) *A. thaliana* leaves. Averages of stomatal apertures from four independent experiments

(n = 4) are shown. Data are the mean ± SE. Different letters indicate statistical significances

based on one-way ANOVA with Tukey's HSD test, $P < 0.05$.

Figure 5. Malate signaling is mediated by cADPR, cAMP and IP₃.

(A–C) Effects of inhibitors on malate-induced stomatal closure in **(A and C)** *A. thaliana* or **(B)**

V. vinifera leaves. Averages of stomatal apertures from four independent experiments (n = 4)

are shown. Data are the mean ± SE. Different letters indicate statistical significances based on

one-way ANOVA with Tukey's HSD test, $P < 0.05$.

(D) Representative traces of fluorescence emission ratios (535/480 nm) in *A. thaliana* guard

cells expressing the Ca^{2+} sensor Yellow Cameleon 3.6. Grey bar indicates the time point when

treatment was applied. The guard cells were treated with organic acids 3 min after the

measurement. Inhibitors were added 5 min before starting imaging.

(E) Percentage of number of guard cells showing different numbers of transient $[Ca^{2+}]_{cyt}$

increases. An increase in $[Ca^{2+}]_{cyt}$ is defined by an increase in fluorescence ratio by ≥0.1 U from

the baseline ratio. Data were obtained from Mock (10 guard cells), Mal (12), +Nicotinamide

(13), +Alloxan (10), +Neomycin (15), +SHAM (12). P-value was calculated by Fisher's exact

test.

Figure 6. Malate signaling is mediated by G-proteins.

(A and B) Effects of inhibitors on malate-induced stomatal closure in **(A)** *V. vinifera* or **(B)** *A.*

thaliana leaves.

(C) Effects of malate on stomatal aperture in *gpa1* and *agb1* mutants. Averages of stomatal

apertures from four independent experiments (n = 4) are shown. Data are the mean ± SE.

Different letters indicate statistical significances based on one-way ANOVA with Tukey's HSD

test, $P < 0.05$.

(D) Effects of GDPβS on malate-induced ROS production in *A. thaliana* guard cells. The ROS-

sensitive dye, 2',7'-dichlorodihydrofluorescein diacetate (H$_2$DCF-DA) was used for ROS

detection in guard cells. Fluorescence intensity was normalized to control values. Averages

from three independent experiments (n = 3) are shown. Data are the mean ± SE. Different letters

indicate statistical significances based on one-way ANOVA with Tukey's HSD test, P < 0.05.

(E) A proposed working model for malate-induced stomatal closure. The external signal of

malate is transmitted by G-proteins, which modulate the generation of second messengers,

including cAMP, cADPR and IP$_3$. This cascade leads to increases in [Ca^{2+}]$_{cyt}$, resulting in the

phosphorylation of SLAC1 by CDPKs. Additionally, G-proteins simultaneously modulate ROS

production, which contributes to SLAC1 phosphorylation. Extracellular malate directly

promotes anion transport through phosphorylated SLAC1, leading to a decrease in turgor

pressure and subsequent stomatal closure.

Supplemental Figure 1. Metabolome analysis in grapevine leaves during dehydration

treatment.

Supplemental Figure 2. Ca^{2+} response to acetate in guard cells.

Supplemental Figure 3. The ion channel activity of VvSALC1 mutants.

Supplemental Figure 4. Malate-induced stomatal closure in other plant species.

Supplemental Figure 5. Malate-induced stomatal closure in the presence of inhibitors.

Supplemental Figure 6. ROS production in the presence of organic acids.

Supplemental Table 1. List of inhibitors used in this work.

Supplemental Table 2. List of primers used in this work.

Supplemental Data 1. Data table of metabolome analysis.

Figure 1. Metabolome analysis in grapevine leaves during dehydration treatment.

Figure 2. Ca^{2+} response to organic acids in guard cells.

Figure 3. SLAC1 activity in the presense of organic acids.

Figure 4. Malate-induced stomatal closure is mediated by anion channels and Ca^{2+}.

Figure 5. Malate signaling is mediated by cADPR, cAMP and IP$_3$.

Figure 6. Malate signaling is mediated by on G-proteins.