Occasional and constant exposure to dietary ethanol shortens the lifespan of worker honey bees

Monika Ostap-Chec[1,2], Daniel Bajorek[3], Weronika Antoł[3], Daniel Stec[3], Krzysztof Miler[3]

[1] Doctoral School of Exact and Natural Sciences, Jagiellonian University, Kraków, Poland

[2] Institute of Environmental Sciences, Faculty of Biology, Jagiellonian University, Kraków, Poland

[3] Institute of Systematics and Evolution of Animals of the Polish Academy of Sciences, Kraków, Poland

Correspondence:

Monika Ostap-Chec, Institute of Environmental Sciences, Faculty of Biology, Jagiellonian University, Gronostajowa 7 St., 30-387 Kraków, Poland. E-mail: ostap.monika@gmail.com

Krzysztof Miler, Institute of Systematics and Evolution of Animals of the Polish Academy of Sciences, Sławkowska 17 St., 31-016 Kraków, Poland. E-mail: miler@isez.pan.krakow.pl

Funding: This work was supported by the National Science Centre, Poland [grant number Sonata UMO-2021/43/D/NZ8/01044].

Abstract

Honey bees (*Apis mellifera*) are one of the most crucial pollinators, providing vital ecosystem services. Their development and functioning depend on essential nutrients and substances found in the environment. While collecting nectar as a vital carbohydrate source, bees routinely encounter low doses of ethanol from yeast fermentation. Yet, the effects of repeated ethanol exposure on bees' survival and physiology remain poorly understood. Here, we investigate the impacts of constant and occasional consumption of food spiked with 1% ethanol on honey bee mortality and alcohol dehydrogenase (ADH) activity. This ethanol concentration might be tentatively judged close to that in natural conditions. We conducted an experiment in which bees were exposed to three types of long-term diets: constant sugar solution (control group that simulated conditions of no access to ethanol), sugar solution spiked with ethanol every third day (that simulated occasional, infrequent exposure to ethanol) and daily ethanol consumption (simulating constant, routine exposure to ethanol). The results revealed that both constant and occasional ethanol consumption increased the mortality of bees, but only after several days. These mortality rates rose with the frequency of ethanol intake. The ADH activity remained similar in bees from all groups. Our findings indicate that exposure of bees to ethanol carries harmful effects that accumulate over time. Further research is needed to pinpoint the exact ethanol doses ingested with food and exposure frequency in bees in natural conditions.

Keywords: *Apis mellifera*, alcohol dehydrogenase, ethanol, honey bees, survival

Introduction

Insect pollinators play a crucial role in the functioning of both natural and agricultural ecosystems (Klein et al. 2007, Gallai et al. 2009). During pollination services, they collect pollen, essential for growth, and nectar, a primary energy source. The quality and composition of their diet are critical for development, health, and survival, making the abundance and diversity of resources in the surrounding environment crucial. While collecting pollen and nectar, pollinators are exposed to substances that directly impact them. For instance, insecticides widespread in agricultural environments significantly increase mortality and adversely affect the development and foraging abilities of pollinators (e.g. in solitary bees, Boff et al. 2021, Mokkapati et al. 2021). Conversely, certain phytochemicals found in pollen and nectar, such as caffeine, can significantly reduce parasitisation, acting both preventatively and therapeutically on pollinators' health (e.g. in bumble and honey bees, Folly et al. 2021, Motta et al. 2023). Pollinators can be passively affected by environmental substances due to ingestion of contaminated resources, but they can also actively seek dietary components. For example, honey bees actively select food types advantageous to them in specific situations, such as opting for nectar with higher antibiotic potential during infection (Gherman et al. 2014) or gathering propolis to combat rising parasite levels (Pusceddu et al. 2021).

Honey bees (*Apis mellifera* L.), which are one of the key insect pollinators, likely encounter and consume ethanol in their natural environment. The main carbohydrate source for honey bees, floral nectar, is frequently colonised by yeasts capable of producing ethanol through fermentation (Lievens et al. 2015). Observations from tropical regions indicate that the concentration of ethanol in palm flower nectar can reach 6.9% (Hockings et al. 2015, Gochman et al. 2016). A concentration of about 1% ethanol, especially in more temperate climates, seems more realistic as a component of nectar in natural conditions (Ehlers and Olesen 1997, Jakubska et al. 2005, Goodrich et al. 2006, Wiens et al. 2008, Golonka et al. 2014, Rering et al. 2018). In any case, ethanol consumption must be significant enough to allow honey bees to biosynthesise ethyl oleate — a crucial pheromone that regulates colony demography by inhibiting the transition of workers into foragers (Leoncini et al. 2004, Castillo et al. 2012a, Castillo et al. 2012b). Only foragers, the type of individuals that leave the hive and explore the surroundings for food, possess dehydrogenase enzymes likely responsible for breaking down ethanol (Miler et al. 2021). Furthermore, despite the aversive taste of ethanol in water for honey bees, when dissolved in sugar, it is willingly consumed both in the laboratory and field, even in extreme concentrations as high as 20% (Abramson et

al. 2000, Abramson et al. 2004a, Maze et al. 2006, Sokolowski et al. 2012, Mustard et al. 2018). Moreover, foragers prefer feeding solutions containing about 1% or even 2.5% ethanol over pure sugar solutions (Mustard et al. 2018). All this suggests not only that ethanol frequently contaminates carbohydrate resources utilised by honey bees but also that they need this dietary component and are adapted for its encounter.

Nevertheless, ethanol has been observed to induce multiple negative effects on honey bees, both in social and non-social contexts. It disrupts social communication among individuals by directly influencing behaviours such as antennation or trophallaxis (Mixson et al. 2010, Wright et al. 2012) and through increasing aggression (Abramson et al. 2004b, Ammons and Hunt 2008). Moreover, ethanol consumption in foragers modifies proper dance communication (Bozic et al. 2006). Ethanol also impairs honeybees' locomotion, foraging, and learning (Abramson et al. 2005, Giannoni-Guzmán et al. 2014, Maze et al. 2006, Mustard et al. 2008, Black et al. 2021, Ahmed et al. 2022). The extent of these behavioural changes occurs in a dose-dependent manner, with higher doses of ethanol producing larger effects on behaviour (Maze et al. 2006, Bozic et al. 2007, Wright et al. 2012). However, the majority of the studies that document effects resulting from ethanol consumption focus on single-exposure impact, often with concentrations that are relatively high and unlikely of ecological relevance. Not only the effects of low ethanol levels are understudied but so are those related to repeated ethanol exposure. Thus far, it has been demonstrated that bees show tolerance, expressed as lower motor impairment in response to ethanol in individuals previously exposed to it than those exposed for the first time (Miler et al. 2018 but see Stephenson et al. 2021). Moreover, individuals fed on food spiked with ethanol for a prolonged time exhibit withdrawal symptoms upon discontinuation of access to such food (Ostap-Chec et al. 2021). These characteristics can be considered primary hallmarks of alcohol dependence development in conditions of repeated encounters with ethanol. Still very little is known about the survivorship and physiological effects associated with repeated exposure to relatively low concentrations of ethanol in the diet.

Here, we investigate the effects of occasional and constant low-level ethanol consumption on the survival and alcohol dehydrogenase (ADH) activity of forager-age honey bees. We hypothesise that there would be no significant increase in mortality compared to abstinent individuals. Moreover, we hypothesise that the ADH activity would be highest in individuals exposed to ethanol most frequently, lower in those with occasional exposure, and lowest in workers with no access to ethanol. Our expectations are based on current knowledge

regarding the likely routine presence of ethanol in food resources utilised by bees, their preferences for this dietary component, and adaptations towards its encounter.

Methods

General procedure

We performed the experiment in 4 bouts, with 2 days of delay between each bout. For each bout, we obtained newly emerged individuals that originated from 2 different colonies. For that purpose, we took a single bee-free frame with a capped brood from each colony participating in a bout and placed it in an incubator (KB53, Binder, Germany) at 32 °C overnight. We marked emerged workers with a coloured dot on the thorax using a non-toxic paint marker and then released them into an unrelated hive. We recollected marked workers when they were 7 days old and transported them to the laboratory. There, the workers were kept in wooden bee cages, 100 individuals per cage, 9 cages per bout, and with access to *ad libitum* water and food (2 M sucrose solution) in an incubator (KB400, Binder, Germany) at 32 °C. We applied a period of acclimation that lasted for 2 weeks, to be sure that all workers reached forager age (Dukas 2008) and remained naïve in terms of ethanol presence in their previous diet. We noted exactly how many individuals were in each cage as there was some mortality during the acclimation period (the number of workers ranged from 94-100 by the point of reaching 21 days of age). Then, we created 3 replicates in each bout, with each replicate comprising 3 groups, and started the diet. For the control group (CON), we provided the workers with a 2 M sucrose water solution. For the group with occasional exposure (EXP1), we provided the workers with the same solution, but every 3rd day, we switched it to a sucrose water solution with the addition of 1% ethanol. For the group with constant exposure (EXP2), we provided the workers with a sucrose water solution with the addition of 1% ethanol. In total, we had 12 cages per group (4 bouts and 3 group replicates in each). The food with the addition of ethanol was prepared iso-caloric relative to the 2 M sucrose solution to avoid the confounding effect of ethanol caloric value when digested. Hence, assuming 5.5 kcal per 1 mL of ethanol, we added 1.38 g less sucrose to the food with ethanol to balance out the calories. We switched water and food to fresh every morning for the duration of the study. During water and food renewal each day, we noted the number of dead individuals in each cage and removed the dead bodies. We exposed workers in each group to their respective diets for 21 days, and then the experiment was terminated.

On the last day of the experiment, we froze 6 individuals from a randomly selected group replicate in each bout to analyse the alcohol dehydrogenase (ADH) activity in their

bodies. For that purpose, each body was thoroughly homogenised in 400 µl of PBS in a Bead Ruptor Elite homogenizer with ceramic beads (Omni International, USA) and spun in a centrifuge at 13000 RPM for 7 min. Then, 10 µl of each supernatant was analysed using the ADH assay kit (ab102533, abcam, the Netherlands) as per the manufacturer's instructions. The kit enables the detection of type I ADH, which is present in bees (Miler et al. 2021). The optical density of the reaction product was read at 450 nm using the Multiskan FC microplate reader (Thermo Scientific, USA) after the initial incubation period (3 min) and in three measurement points between 7 and 35 min after the first measurement. The timeframe was determined to be within a range of linear reaction phase. The ADH activity (defined as nM NADH produced per min in 1 mL = mU/mL) was calculated based on the slope of the absorbance curve and translated into concentration values based on the prepared calibration curve slope. One sample was excluded from the analyses as the calculated activity had a slightly negative value (-0.087, 95% CI: -0.101-0.099), which was not biologically relevant.

Statistics

To analyse the data, we used R (R Core Team 2024). For the analysis of survivorship, we used the Cox mixed-effect regression ('coxme' and 'survival' packages, Therneau 2022, 2023) with a fixed factor of the group and random factors of the bout and replicate (coxme function). Individuals that remained alive when the experiment was terminated (day 21 of the diet) were censored. To analyse the ADH activity, we used the generalised linear mixed-effects model ('lme4' package, Bates et al. 2015) with a Gamma distribution and a fixed factor of the group and a random factor of the bout (lme function).

Results

Throughout the diet (21 days), the survival in the control groups (CON) dropped to 79% (95% CI: 81-77%). Compared to that, the groups with occasional ethanol exposure (EXP1; $z = 7.24$, $p < 0.001$) and with constant ethanol exposure (EXP2; $z = 18.02$, $p < 0.001$) both demonstrated decreased survival. In the case of occasional exposure, the survival dropped to 65% (95% CI: 68-62%) whereas for constant exposure – to 41% (95% CI: 44-38%). Groups exposed to ethanol started to diverge from the control groups after 10 and 6 days of the diet for groups EXP1 and EXP2, respectively (Fig 1). The ADH activity did not differ between groups ($\chi^2 = 1.029$, $p = 0.598$, Fig 2).

Fig 1. Survival plot for honey bees exposed to different types of diets for 21 days (CON = the control groups exposed to pure 2 M sucrose solution, EXP1 = the groups with occasional exposure to ethanol (addition of 1% ethanol in food every third day), EXP2 = the groups with constant exposure to ethanol (food with the addition of 1% ethanol)). There were 12 cages for each group, each with an initial number of individuals ranging from 94-100. Shadings show 95% CI.

Fig 2. Violin plot for the ADH activity (mU / mL) in honey bees exposed to different types of diets for 21 days (CON = the control groups exposed to pure 2 M sucrose solution (n = 24), EXP1 = the groups with occasional exposure to ethanol (addition of 1% ethanol in food every third day) (n = 24), EXP2 = the groups with constant exposure to ethanol (food with the addition of 1% ethanol) (n = 23)). Black dots indicate medians and coloured dots show individual data points.

Discussion

Honey bees surely encounter and consume ethanol in their natural environment. They rely on ethanol for synthesizing one of their crucial pheromones (Leoncini et al. 2004, Castillo et al. 2012a, Castillo et al. 2012b), possess enzymes to break down ethanol (Miler et al. 2021), and willingly consume it when mixed with sugars (Abramson et al. 2000, Abramson et al. 2004a, Maze et al. 2006, Sokolowski et al. 2012, Mustard et al. 2018). While previous research has focused on the effects of single and often high-concentration ethanol exposure on honey bees, this study investigates the consequences of repeated exposure to low levels of ethanol. Our findings reveal a significant decrease in lifespan due to ethanol exposure. Mortality rates in our study show that a significant drop in survival starts to appear after about 6 and 10 days of constant and occasional ethanol consumption, respectively. It is worth noting, however, that in natural conditions the average lifespan of forager-age bees is about a week (Dukas 2008). Therefore, while our results highlight the potentially harmful effects of even low ethanol concentrations and suggest cumulative impacts with each subsequent exposure, they need to be put in an ecological context. It seems likely that bees die for other reasons before the negative impact of routine, low-level ethanol exposure in their food appears. For comparison, in the study of Mustard et al. (2019), no discernible differences in mortality were observed in bees exposed daily to 1.25% ethanol and those abstinent. In their study, however, the authors used foragers of unknown age captured at hive entrances and a vast majority of all these bees died within a week. It is conceivable that in that study, the effects of diet had no chance to become manifested (Mustard et al. 2019). Rasmussen et al. (2021) utilized overwintering bees with a longer-lived phenotype and demonstrated that 1% ethanol in their diet significantly decreased survival, but with a drop in mortality occurring only after a few days of ethanol exposure. In our study, even if we consider mortality over the entire 3 weeks of measurement, well over a third of all individuals survived daily ethanol consumption (Fig 1). Together, these results highlight the extraordinary resilience to the potentially toxic effect of ethanol in honey bees. Previous research has demonstrated that ethanol induces an increase in heat shock protein concentration in honey bee brain tissue (Hranitz et al. 2010). Rasmussen et al. (2021) also showed that ethanol intake disrupts their DNA methylation patterns. These effects likely constitute the basis of the eventual harmfulness of ethanol consumption and its accumulative toxicity. Yet, it takes time for such effects to reflect on the survivorship of bees.

Bees likely ingest ethanol with their food as they collect floral nectar. Another potential source of ethanol is presented in spoiled or overripe fruits (e.g. Dudley 2002,

Dominy 2004, Sánchez et al. 2004) because bees often collect juices from such fruits, especially during periods without major flowering events. Our understanding of the sources of honey bee exposure to ethanol, as well as the frequency and concentration at which honey bees come into contact with this compound, remains a gap that requires extensive research. The same is true for other pollinating nectarivores as well as frugivores, all of which probably meet with ethanol routinely (Dudley and Maro 2021).

Our results revealed no differences in the ADH activity between the groups of individuals (Fig 2), suggesting that foragers consistently produce the enzyme, regardless of their ethanol exposure. Similar results to ours were obtained in fruit-feeding butterflies, in which a diet containing ethanol did not alter ADH levels (Beaulieu et al. 2017). Miler et al. (2021) demonstrated ADH presence in honey bee foragers but not in intranidal workers that stay inside the hive and care for brood. Additionally, another study showed that even repeated ethanol exposure failed to induce ADH production in intranidal workers (Miler et al. 2022). Our current findings complement this prior research (Beaulieu et al. 2017, Miler et al. 2021, Miler et al. 2022) indicating that ADH activity remains constant, regardless of ethanol intake. Notably, however, there are different types of ADH (Hernández-Tobías et al. 2011). In the case of honey bees, types I and II are present, but only the former is analysed here and in previous research (Miler et al. 2021, Miler et al. 2022). Short-chain type II ADH is more typical for insects and might show higher activity with ethanol in bees, similar to *Drosophila* (Hernández-Tobías et al. 2011). Hence, our results regarding ADH activity cannot be treated as entirely conclusive. The analysed type I ADH might not be restricted in its function to ethanol metabolism and this may confound the results. We recommend future research to focus also on *Drosophila*-type ADH activity in bees. At least in fruit flies, this type of ADH is connected to ethanol preference and tolerance (e.g. Ogueta et al. 2010).

Our findings stress the necessity for further research to find ecologically relevant levels of ethanol and the frequency of exposure to this substance in bees as well as their responses to this dietary component. The way for the comparative biology of ethanol exposure and response is open (Dudley and Maro 2021) and honey bees show promise in this context. Considering that ethanol is one of many neuroactive substances present in floral nectar that potentially mediate plant-pollinator interactions (Mustard 2020), this research field needs much more attention.

References

Abramson CI, Stone SM, Ortez RA, Luccardi A, Vann KL, Hanig KD, Rice J (2000) The development of an ethanol model using social insects I: behavior studies of the honey bee (*Apis mellifera* L). Alcohol Clin Exp Res 24:1153-1166. https://doi.org/10.1111/j.1530-0277.2000.tb02078.x

Abramson CI, Place AJ, Aquino IS, Fernandez A (2004b) Development of an ethanol model using social insects: IV. Influence of ethanol on the aggression of Africanized honey bees (*Apis mellifera* L). Psychol Rep 94:1107-1115. https://doi.org/10.2466/pr0.94.3c.1107-1115

Abramson CI, Sheridan A, Donohue D, Kandolf A, Božič J, Meyers JE, Benbassat D (2004a) Development of an ethanol model using social insects: III. Preferences for ethanol solutions. Psychol Rep 94:227-239. https://doi.org/10.2466/pr0.94.1.227-239

Abramson CI, Sanderson C, Painter J, Barnett S, Wells H (2005) Development of an ethanol model using social insects: V. Honeybee foraging decisions under the influence of alcohol. Alcohol 36:187-193. https://doi.org/10.1016/j.alcohol.2005.09.001

Ahmed I, Abramson CI, Faruque IA (2022) Honey bee flights near hover under ethanol-exposure show changes in body and wing kinematics. PLoS ONE 17:e0278916. https://doi.org/10.1371/journal.pone.0278916

Ammons AD, Hunt GJ (2008) Characterization of honey bee sensitivity to ethanol vapor and its correlation with aggression. Alcohol 42:129-136. https://doi.org/10.1016/j.alcohol.2007.12.005

Bates D, Mächler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models Using lme4. J Stat Softw 67:1-48. https://doi.org/10.18637/jss.v067.i01

Beaulieu M, Franke K, Fischer K (2017) Feeding on ripening and over-ripening fruit: interactions between sugar, ethanol and polyphenol contents in a tropical butterfly. J Exp Biol 220:3127-3134. https://doi.org/10.1242/jeb.162008

Black TE, Stauch KLN, Wells H, Abramson CI (2021) Effects of ethanol ingestion on aversive conditioning in honey bees (*Apis mellifera* L). J Comp Psychol 135:559–567. https://doi.org/10.1037/com0000296

Boff S, Scheiner R, Raizer J, Lupi D (2021) Survival rate and changes in foraging performances of solitary bees exposed to a novel insecticide. Ecotoxicol Environ Saf 211:111869. https://doi.org/10.1016/j.ecoenv.2020.111869

Bozic J, Abramson CI, Bedencic M (2006) Reduced ability of ethanol drinkers for social communication in honeybees (*Apis mellifera carnica* Poll). Alcohol 38:179-183. https://doi.org/10.1016/j.alcohol.2006.01.005

Bozic J, DiCesare J, Wells, H, Abramson CI (2007). Ethanol levels in honeybee hemolymph resulting from alcohol ingestion. Alcohol 41:281-284. https://doi.org/10.1016/j.alcohol.2007.04.003

Castillo C, Chen H, Graves C, Maisonnasse A, Le Conte Y, Plettner E (2012a) Biosynthesis of ethyl oleate, a primer pheromone, in the honey bee (*Apis mellifera* L). Insect Biochem Mol Biol 42:404-416. https://doi.org/10.1016/j.ibmb.2012.02.002

Castillo C, Maisonnasse A, Le Conte Y, Plettner E (2012b) Seasonal variation in the titers and biosynthesis of the primer pheromone ethyl oleate in honey bees. J Insect Physiol 58:1112-1121. https://doi.org/10.1016/j.jinsphys.2012.05.010

Dominy NJ (2004) Fruits, fingers, and fermentation: the sensory cues available to foraging primates. Integr Comp Biol 44:295-303. https://doi.org/10.1093/icb/44.4.295

Dudley R (2002) Fermenting fruit and the historical ecology of ethanol ingestion: is alcoholism in modern humans an evolutionary hangover? Addiction 97:381-388. https://doi.org/10.1046/j.1360-0443.2002.00002.x

Dudley R, Maro A (2021) Human evolution and dietary ethanol. Nutrients 13:2419. https://doi.org/10.3390/nu13072419

Dukas R (2008) Mortality rates of honey bees in the wild. Insect Soc 55:252-255. https://doi.org/10.1007/s00040-008-0995-4

Ehlers BK, Olesen JM (1997) The fruit-wasp route to toxic nectar in *Epipactis* orchids? Flora 192:223-229. https://doi.org/10.1016/S0367-2530(17)30787-9

Folly AJ, Koch H, Farrell IW, Stevenson PC, Brown MJ (2021) Agri-environment scheme nectar chemistry can suppress the social epidemiology of parasites in an important pollinator. Proc R Soc B 288:20210363. https://doi.org/10.1098/rspb.2021.0363

Gallai N, Salles JM, Settele J, Vaissière BE (2009) Economic valuation of the vulnerability of world agriculture confronted with pollinator decline. Ecol Econ 68:810-821. https://doi.org/10.1016/j.ecolecon.2008.06.014

Gherman BI, Denner A, Bobiş O, Dezmirean DS, Mărghitaş LA, Schlüns H, Moritz RFA, Erler S (2014) Pathogen-associated self-medication behavior in the honeybee *Apis mellifera*. Behav Ecol Sociobiol 68:1777-1784. https://doi.org/10.1007/s00265-014-1786-8

Giannoni-Guzmán MA, Giray T, Agosto-Rivera JL, Stevison BK, Freeman B, Ricci P, Brown EA, Abramson CI (2014) Ethanol-induced effects on sting extension response and punishment learning in the western honey bee (*Apis mellifera*). PLoS ONE 9:e100894. https://doi.org/10.1371/journal.pone.0100894

Gochman SR, Brown MB, Dominy NJ (2016) Alcohol discrimination and preferences in two species of nectar-feeding primate. R Soc Open Sci 3:160217. https://doi.org/10.1098/rsos.160217

Golonka AM, Johnson BO, Freeman J, Hinson DW (2014) Impact of nectarivorous yeasts on *Silene caroliniana*'s scent. East Biol 3:1-26.

Goodrich KR, Zjhra ML, Ley CA, Raguso RA (2006) When flowers smell fermented: the chemistry and ontogeny of yeasty floral scent in pawpaw (*Asimina triloba*: Annonaceae). Int J Plant Sci 167:33-46. https://doi.org/10.1086/498351

Hernández-Tobías A, Julián-Sánchez A, Piña E, Riveros-Rosas H (2011) Natural alcohol exposure: is ethanol the main substrate for alcohol dehydrogenases in animals? Chem Biol Interact 191:14-25. https://doi.org/10.1016/j.cbi.2011.02.008

Hockings KJ, Bryson-Morrison N, Carvalho S, Fujisawa M, Humle T, McGrew WC, Nakamura M, Ohashi G, Yamanashi Y, Yamakoshi G, Matsuzawa T (2015) Tools to tipple: ethanol ingestion by wild chimpanzees using leaf-sponges. R Soc Open Sci 2:150150. https://doi.org/10.1098/rsos.150150

Hranitz JM, Abramson CI, Carter RP (2010) Ethanol increases HSP70 concentrations in honeybee (*Apis mellifera* L.) brain tissue. Alcohol 44:275-282. https://doi.org/10.1016/j.alcohol.2010.02.003

Jakubska A, Anioł-Kwiatkowska J, Kadej M, Prząd D (2005) Why do pollinators become „sluggish"? Nectar chemical constituents from *Epipactis helleborine* (L.) Crantz (Orchidaceae). App Ecol Env Res 3:29-38. https://doi.org/10.15666/aeer/0302_029038

Klein AM, Vaissière BE, Cane JH, Steffan-Dewenter I, Cunningham SA, Kremen C, Tscharntke T (2007) Importance of pollinators in changing landscapes for world crops. Proc R Soc B 274:303-313. https://doi.org/10.1098/rspb.2006.3721

Leoncini I, Le Conte Y, Costagliola G, Plettner E, Toth AL, Wang M, Huang Z, Bécard JM, Crauser D, Slessor KN, Robinson GE (2004) Regulation of behavioral maturation by a primer pheromone produced by adult worker honey bees. Proc Natl Acad Sci 101:17559-17564. https://doi.org/10.1073/pnas.0407652101

Lievens B, Hallsworth JE, Pozo MI, Belgacem ZB, Stevenson A, Willems KA, Jacquemyn H (2015) Microbiology of sugar-rich environments: diversity, ecology and system constraints. Environ Microbiol 17:278-298. https://doi.org/10.1111/1462-2920.12570

Maze IS, Wright GA, Mustard JA (2006) Acute ethanol ingestion produces dose-dependent effects on motor behavior in the honey bee (*Apis mellifera*). J Insect Physiol 52:1243-1253. https://doi.org/10.1016/j.jinsphys.2006.09.006

Miler K, Kuszewska K, Privalova V, Woyciechowski M (2018) Honeybees show adaptive reactions to ethanol exposure. Sci Rep 8:8707. https://doi.org/10.1038/s41598-018-27117-6

Miler K, Stec D, Kamińska A, Pardyak L, Kuszewska K (2021) Alcohol intoxication resistance and alcohol dehydrogenase levels differ between the honeybee castes. Apidologie 52:230-241. https://doi.org/10.1007/s13592-020-00812-y

Miler K, Stec D, Pardyak L, Kamińska A, Kuszewska K (2022) No increase in alcohol dehydrogenase levels following repeated ethanol exposure in young honeybee workers. Physiol Entomol 47:110-116. https://doi.org/10.1111/phen.12380

Mixson TA, Abramson CI, Božič J (2010) The behavior and social communication of honey bees (*Apis mellifera carnica* Poll.) under the influence of alcohol. Psychol Rep 106:701-717. https://doi.org/10.2466/pr0.106.3.701-717

Mokkapati JS, Bednarska AJ, Laskowski R (2021) The development of the solitary bee *Osmia bicornis* is affected by some insecticide agrochemicals at environmentally relevant

concentrations. Sci Total Environ 775:145588.
https://doi.org/10.1016/j.scitotenv.2021.145588

Motta EV, Arnott RL, Moran NA (2023) Caffeine Consumption Helps Honey Bees Fight a
Bacterial Pathogen. Microbiol Spectrum 11:e00520-23.
https://doi.org/10.1128/spectrum.00520-23

Mustard JA (2020) Neuroactive nectar: compounds in nectar that interact with neurons.
Arthropod-Plant Interact 14:151-159. https://doi.org/10.1007/s11829-020-09743-y

Mustard JA, Edgar EA, Mazade RE, Wu C, Lillvis JL, Wright GA (2008) Acute ethanol
ingestion impairs appetitive olfactory learning and odor discrimination in the honey bee.
Neurobiol Learn Mem 90:633-643. https://doi.org/10.1016/j.nlm.2008.07.017

Mustard JA, Oquita R, Garza P, Stoker A (2019) Honey bees (*Apis mellifera*) show a
preference for the consumption of ethanol. Alcohol Clin Exp Res 43:26-35.
https://doi.org/10.1111/acer.13908

Ogueta M, Cibik O, Eltrop R, Schneider A, Scholz H (2010) The influence of Adh function
on ethanol preference and tolerance in adult *Drosophila melanogaster*. Chem Senses 35:813-
822. https://doi.org/10.1093/chemse/bjq084

Ostap-Chec M, Opalek M, Stec D, Miler K (2021) Discontinued alcohol consumption elicits
withdrawal symptoms in honeybees. Biol Lett 17:20210182.
https://doi.org/10.1098/rsbl.2021.0182

Pusceddu M, Annoscia D, Floris I, Frizzera D, Zanni V, Angioni A, Satta A, Nazii F (2021)
Honeybees use propolis as a natural pesticide against their major ectoparasite. Proc R Soc B
288:20212101. https://doi.org/10.1098/rspb.2021.2101

R Core Team (2023) R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. https://www.R-project.org/

Rasmussen EM, Seier KL, Pedersen IK, Kreibich C, Amdam GV, Münch D, Dahl JA (2021)
Screening bioactive food compounds in honey bees suggests curcumin blocks alcohol-
induced damage to longevity and DNA methylation. Sci Rep 11:19156.
https://doi.org/10.1038/s41598-021-98614-4

Rering CC, Beck JJ, Hall GW, McCartney MM, Vannette RL (2018) Nectar-inhabiting microorganisms influence nectar volatile composition and attractiveness to a generalist pollinator. New Phytol 220:750-759. https://doi.org/10.1111/nph.14809

Sánchez F, Korine C, Pinshow B, Dudley R (2004) The possible roles of ethanol in the relationship between plants and frugivores: first experiments with Egyptian fruit bats. Integr Comp Biol 44:290-294. https://doi.org/10.1093/icb/44.4.290

Sokolowski MB, Abramson CI, Craig DPA (2012) Ethanol Self-Administration in Free-Flying Honeybees (*Apis mellifera* L.) in an Operant Conditioning Protocol. Alcohol Clin Exp Res 36:1568-1577. https://doi.org/10.1111/j.1530-0277.2012.01770.x

Stephenson L, Chicas-Mosier A, Black T, Wells H, Abramson C (2021) Inducing ethanol tolerance in free-flying honey bees (*Apis mellifera* L.). Int J Comp Psychol 34:1-13. https://doi.org/10.46867/ijcp.2021.34.00.03

Therneau T (2022) coxme: mixed effects Cox models. R package version 2.2-18.1, https://CRAN.R-project.org/package=coxme

Therneau T (2023) survival: Survival Analysis. R package version 3.5-7, https://CRAN.R-project.org/package=survival

Wiens F, Zitzmann A, Lachance M-A, Yegles M, Pragst F, Wurst FM, von Holst D, Leng Guan S, Spanagel R (2008) Chronic intake of fermented floral nectar by wild treeshrews. Proc Natl Acad Sci 105:10426-10431. https://doi.org/10.1073/pnas.0801628105

Wright GA, Lillvis JL, Bray HJ, Mustard JA (2012) Physiological state influences the social interactions of two honeybee nest mates. PLoS ONE 7:e32677. https://doi.org/10.1371/journal.pone.0032677