Discovery of orally bioavailable SARS-CoV-2 papain-like protease inhibitor as a potential treatment for COVID-19

Yongzhi Lu[1,2#], Qi Yang [1,3#], Ting Ran[1#], Guihua Zhang[1], Wenqi Li[1], Peiqi Zhou[1], Jielin Tang[1], Minxian Dai[1], Jinpeng Zhong[1], Hua Chen[1], Pan He[1], Anqi Zhou[3], Bao Xue[1], Jiayi Chen[1], Jiyun Zhang[1], Sidi Yang[1], Kunzhong Wu[1], Xinyu Wu[1], Miru Tang[1], Wei K Zhang[1], Deyin Guo[1,3], Xinwen Chen[1,3*], Hongming Chen[1,2*] and Jinsai Shang[1,2,4*]

Affiliations

[1]Guangzhou National Laboratory, Guangzhou 510005, China

[2]School of Basic Medical Sciences, Guangzhou Laboratory, Guangzhou Medical University, Guangzhou 511436, China

[3]State Key Laboratory of Respiratory Disease, Guangzhou Medical University, Guangzhou, 511436, China

[4]Lead contact

#These authors contributed equally: Yongzhi Lu, Qi Yang and Ting Ran

*Correspondence should be addressed to X.C. (chen_xinwen@gzlab.ac.cn); H.C. (chen_hongming@gzlab.ac.cn) and J.S. (shang_jinsai@gzlab.ac.cn)

Abstract

The RNA-dependent RNA polymerase (RdRp), 3C-like protease (3CL[pro]), and papain-like protease (PL[pro]) are pivotal components in the viral life cycle of SARS-CoV-2, presenting as promising therapeutic targets. Currently, all FDA approved antiviral drugs against SARS-CoV-2 are RdRp or 3CL[pro] inhibitors. However, the mutations causing drug resistance have been observed in RdRp and 3CL[pro] from SARS-CoV-2, which makes it necessary to develop antivirals with novel mechanisms. Through the application of a structure-based drug design (SBDD) approach, we discovered a series of novel potent non-covalent PL[pro] inhibitors with remarkable *in vitro* potency and *in vivo* PK properties. The co-crystal structures of PL[pro] with leads revealed that the residues E164 and Q269 around the S2 site are critical for improving the inhibitor's potency. The lead compound GZNL-P36 not only inhibited SARS-CoV-2 and its variants at the cellular level with EC_{50} ranging from 58.2 nM to 306.2 nM, but also inhibited HCoV-NL63 and HCoV-229E with EC_{50} of 81.6 nM and 2.66 μM, respectively. Oral administration of the compound resulted in significantly improved survival and notable reductions in lung viral loads and lesions in SARS-CoV-2 infection mouse model, consistent with RNA-seq data analysis. Our results indicate that PL[pro] inhibitor is a promising SARS-CoV-2 therapy.

Introduction

Over 670 million people have been infected and over 6.8 million people have died in the worldwide pandemic caused by SARS-CoV-2 virus according to data from Johns Hopkins University. Despite the efficacy demonstrated by vaccines and targeted small molecule drugs in preventing and treating COVID-19, the ongoing emergence of viral mutations, such as Alpha, Beta, Gamma, Delta and Omicron presents escalating challenges [1-6]. The high mutation frequency of spike protein is responsible for the escape of SARS-CoV-2 from the vaccines. Unlike spike protein, the non-structural proteins (such as 3CL[pro](nsp5) and PL[pro](nsp3)) remain conserved among coronavirus functional proteins and show much lower mutation frequency in natural SARS-CoV-2 variants[7,8]. The high conservativeness makes 3CL[pro] and PL[pro] attractive drug targets. At present, there are several clinical available small molecule anti-SARS-CoV-2 drugs. Among these drugs, remdesivir, molnupiravir, and VV116 target RNA-dependent RNA polymerase (RdRp)[9-11], nirmatrelvir and ensitrelvir target chymotrypsin-like protease (3CL[pro], also referred as 3CL [pro]) [12,13]. Unfortunately, the clinical efficacy of remdesivir is controversial[14], and multiple reported cases have already outlined an increasing observed resistance to remdesivir in immuno-compromised patients undergoing treatment with the drug[15-17]. Molnupiravir is not authorized to be used for patients under the age of 18 due to its bone and cartilage toxicity, and also not applicable for pregnant patients due to the potential risk of major birth defects and miscarriage[18]. In order to increase the half-life and the *in vivo* concentration of nirmatrelvir, ritonavir is included as a boosting agent to inhibit the activity of cytochrome P450 3A4 (CYP3A4)[19]. Ensitrelvir, the 2nd generation 3CL [pro] inhibitor, showed favorable clinical antiviral efficacy, albeit having potent CYP3A4 inhibitory activity[13,20,21]. Although 3CL [pro] is known as a well conserved protein, Duan *et al*. reported several potential mutant sites by which SARS-CoV-2 might evolve the resistance to nirmatrelvir[22].

PL[pro], a major functional domain in SARS-CoV and SARS-CoV-2 non-structural protein 3 (nsp3), is an essential enzyme involved in viral replication and immune evasion[23-25]. PL[pro] plays an important role in viral transcription and replication by cleaving the peptide bonds in the viral polyprotein, while the deubiquitinating and deISGylating activity of PL[pro] is related to the immune evasion by antagonizing the host's innate immune response upon viral infection[24]. PL[pro]-mediated deubiquitination of STING disrupted the STING-IKKε-IRF3 complex by removing the K63-linked polyubiquitin chain from LYS[289] of STING[25]. Subsequently, the IFN-I signal pathway was inhibited. Hence, PL[pro] is a promising drug target against SARS-CoV-2, too.

GRL0617, a SARS-CoV PL[pro] inhibitor, also shows inhibition activity for SARS-CoV-2 PL[pro26,27]. In addition, several other SARS-CoV-2 PL[pro] inhibitors were reported[28,29,30-33]. GRL0617 prevents the substrate binding by inducing the conformation change of Y268 on the BL2 loop that closes the BL2 loop and narrows the binding cleft[34]. However, these reported compounds only show enzymatic activity from μM to sub-μM. Except for F0213, there is no drug-like PL[pro] inhibitors have reported *in vivo* antiviral efficacy in SARS-CoV-2 infected animal model[8,19,33]. In this study, we synthesized a series of novel PL[pro] inhibitors and evaluated their activities. The lead compound (GZNL-P36) showed excellent *in vitro* potency as well as decent

75   oral *in vivo* pharmacokinetic (PK) properties, more importantly, it also demonstrated similar *in*

76   *vivo* antiviral efficacy in SARS-CoV-2 infected mice with ensitrelvir.

Results

Structure based discovery and optimization of novel PL^{pro} inhibitors

As shown in the co-crystal structure of GRL0617, the compound is bound at a shallow

pocket on the protein surface (**Fig. 1A**), which has been recognized as the substrate binding site.

The naphthalene ring sandwiched in the BL2 groove (*i.e.*, S1 site) is a critical group for the

binding of GRL0617. In previous studies, replacing this group with alternative aromatic ring

systems can somewhat maintain its bio-activity rather than increase it[32]. However, we

hypothesize the potential enhancement of binding affinity through the substitution of

naphthalene with a bulkier substitute, as the naphthalene ring has not yet fully occupied the

entire surface of the groove, in particular the area corresponding to the residue P247. Thus, we

introduced the 1,2-dihydroacenaphthylene (DHAN) group as a replacement of naphthalene ring

to expand the hydrophobic contact surface with P247, and GZNL-P1 was synthesized (**Fig. 1C**).

Encouragingly, its inhibitory activity (IC_{50} = 2.83 µM) is better than GRL0617 (IC_{50} = 4.82 µM)

(**Fig. 1D**), indicating the beneficial effect of introducing a bulkier substitute in the BL2 groove.

What's even more interesting is that the activity can be further improved by over 10 folds for the

GZNL-P3 (IC_{50} = 185.80 nM) when we simultaneously changed the substituent in the linking

group (L) from methyl to cyclopropyl (**Fig. 1C and D**). Notably, cyclopropyl group is a much

better substituent than the di-methyl as the IC_{50} of GZNL-P2 is just around 6.90 µM. Therefore,

GZNL-P3 serves as a good starting point for further lead optimization.

Previous work done by Shen *et al*. has shown that it is possible to engage positively charged

amine groups on the benzene of GRL0617 to interact with E167 at the S2 site[32]. This interaction

could enhance activity by forming salt bridge and hydrogen bonding. A docking calculation

based library design was carried out to explore R_1 groups (**Fig. 1B, C**) which can form salt

bridge with E167. Around 41 library compounds were selected based on docking score for

synthesis and it was found that piperazine derivatives generally exhibit strong inhibition on

PL^{pro}). Meanwhile, compounds with 3-substituted azetidines also shows excellent activity against

PL^{pro}. Remarkably, GZNL-P17 is the most potent compound with IC_{50} of 2.91 nM, which is

more potent than the best piperazine derivative, *i.e.*, GZNL-P4 (IC_{50} = 36.29 nM). However,

modifying the methyl group on the para position of the benzene ring (*i.e.*, R_2 group in **Fig. 1C**)

extending to the recently identified important residue L162[35] leads to activity drop-off. At this

stage, we have successfully achieved potent enzymatic activity which is hundreds of times

improved comparing to GRL0617. The most active compounds GZNL-P4 and GZNL-P17 were

then selected to measure their antiviral activity against both wild-type SARS-CoV-2 and its two

epidemic variants (**Extended Data Fig. 4**) in infected VeroE6 cells and sub-micromolar anti-

viral potency were achieved which is much improved comparing with GRL0617 (EC_{50} = 23.64

µM [28]). To evaluate their ADME properties, *in vitro* liver stability of GZNL-P4 and GZNL-P17

(**Fig. 1F**) was measured. In rat liver microsomes, the half-life time ($T_{1/2}$) of both compounds is
lower than 15 minutes and their intrinsic clearance rate (Clint) is very high. Moreover, they have
worse stability in human liver microsomes. Metabolite identification work of GZNL-P4 indicates
that compound instability could partially be attributed to the oxidation of DHAN ring (see
metabolites analysis in **Extended Data Fig. 1**). To address the liver stability problem, we
changed the DHAN ring to the benzoindolone ring. To our delight, for GZNL-P35 and 36, the
liver stability was considerably improved (**Fig. 1F**), while their enzymatic activities were
maintained at the same level. The enzymatic inhibition activities of the finally designed
compound GZNL-P35 and GZNL-P36 were 8.15 nM and 6.45 nM (**Fig. 1D**), respectively. The
inhibitors can stabilize and increase the melt temperature (T_m) of SARS-CoV-2 PL^{pro} (data was
not shown). The cellular antiviral activity for benzoindolone compounds were examined, where
GZNL-P31, 35, and 36 exhibit better potency than 100 nM against XBB.1 strains, and their
toxicity to normal cells (HEK293T CC_{50} = 157.4, 67.67, 88.41 μM, respectively) is higher than
60 μM (Extended Data Table 1). Overall, GZNL-P35 and 36 demonstrate the most favorable
profile in terms of both potency and stability. The overall workflow for lead optimization is
shown in **Fig. 1B, C**. Bioactivity data of selected compounds are listed in **Fig. 1F** and **Extended
Data Table S1**.

An *in vivo* PK study was carried out for GZNL-P35 and 36 using 3 male CD1 mice (SPF level)
per group with a dosage of 10 mg/kg, both compounds can reach the maximum plasma
concentration at 1.58 and 1.67 h (T_{max}), respectively with a peak plasma concentration (C_{max}) of
227 and 549 ng/mL (**Fig. 1E**). However, the clearance of GZNL-P35 ($T_{1/2}$ = 0.96 h) is much faster
than GZNL-P36 ($T_{1/2}$ = 1.45 h). This results in an enhancement of the performance of GZNL-P36
on drug blood exposure. Particularly, the bioavailability (F%) of GZNL-P36 is much higher than
GZNL-P35. Further profiling of PK properties demonstrated that GZNL-P36 has weak inhibition
on major metabolic enzymes in liver (**Fig. 1G**). Its inhibition on CYP 1A2, 2C9, 2C19, 2D6, 3A4
all are very weak. Additionally, its hERG toxicity is within acceptable limits. In summary, the
strong *in vitro* activity and good PK properties of GZNL-P36 make it suitable to move forward to
in vivo efficacy study.

X-ray crystal structures of SARS-CoV-2 PL^{pro} with inhibitor

To clarify the binding mechanism of inhibitors, the X-ray complex crystal structures of SARS-
CoV-2 PL^{pro} with GZNL-P4, GZNL-P28, GZNL-P31, and GZNL-P35 were determined
(resolution range of 1.7 to 2.6 Å; **Fig. 2, Extended Data Table 2**). These compounds have similar
binding patterns, and the unbiased electron density for PL^{pro} inhibitors GZNL-P4, GZNL-P28,
GZNL-P31, and GZNL-P35 are unambiguous. The amide structures of GZNL-P4, GZNL-P28,
and GZNL-P31 make two hydrogen bonds with Q269 and E164. While GZNL-P35 only formed
one hydrogen bond using its amide structure with E167. Compared to the naphthalene ring of
GRL0617, its characteristic DHAN group at the BL2 groove makes a similar π-π stacking
interaction with Y268 and further expands its contact surface with P247 and P248 (**Fig. 2**). For

these compounds, the substitution with methyl-substituted piperazine at R^1 makes additional
hydrogen bond and salt bridge with the residue E167 and Q269 that is responsible for the
significant potency enhancement (**Fig. 2**). Based on the remarkable enhancement of enzymatic
activity and the structure comparison (Extended Data Table 1) of GZNL-P1 and GZNL-P3, it is
clear that the cyclopropyl group is critical for the enzymatic inhibition activity improvement.
Comparison among co-crystal structures of PL^{pro} with GRL0617 (PDB: 7JRN), GZNL-P4 and
GZNL-P35 showed that the cyclopropyl group makes the plane of amido bond rotate 42.7 degrees
for GZNL-P4 and 34.2 degree for GZNL-P35 (**Extended Data Fig. 2J, 2K**) comparing to that of
GRL0617. This conformation change resulted that GZNL-P4 and GZNL-P35 form a hydrogen
bond with the residue Y264 (**Extended Data Fig. 2D-2F**). In addition, the H-π interaction between
the cyclopropyl group and the aromatic side chain of Y264 is another favorable factor.

The mechanism of GZNL-P36 in inhibiting SARS-CoV-2 PL^{pro}

It is known that PL^{pro} plays a key role in the proteolytic processing of viral polyproteins and
the dysregulation of the host immune response. The deubiquitylation and de-ISGylation activity
of PL^{pro} is related with the host innate immune pathways and the innate immune evasion of SARS-
CoV-2[24]. To characterize the enzymatic inhibition of the designed compounds, we performed the
PL^{pro} enzymatic assay using the labeled peptide substrate RLRGG-AMC (GLPBIO, GA23715).
The final selected candidate compound GZNL-P36 showed potent enzymatic inhibition with IC_{50}
value of 6.4 nM, compared to 4.8 μM for reference compound GRL0617 and 36.3 nM for the lead
compound GZNL-P4 (**Fig.1D**). Correspondingly, the binding affinity of GRL0617, GZNL-P35
and GZNL-P36 to PL^{pro} are 2.59 μM, 4.43 nM and 21.8nM in ITC experiment (**Extended Data**
Fig. 3A-3D). The enthalpy (ΔH), entropy (−TΔS) and Gibbs free energy (ΔG) are -14 kcal/mol,
2.65 kcal/mol and -11.4 kcal/mol for GZNL-P35; -15kcal/mol, 4.52 kcal/mol and -10.5 kcal/mol
for GZNL-P36, but -8.6 kcal/mol, 0.964 kcal/mol and -7.62 kcal/mol for the reference compound
GRL0617. So, the structure optimization from GRL0617 to GZNL-P35 and GZNL-P36 is mainly
driven by enthalpy (**Extended Data Fig. 3A-3D**). The improvement of Gibbs free energy from
GRL0617 to GZNL-P4/GZNL-P17 and GZNL-P35/GZNL-P36 is benefited from the substitution
of naphthalene by acenaphthylene and the additional N-methyl-substituted piperazine group
(GZNL-P4) or bridged piperazine (GZNL-P35) that contribute bigger interaction area with PL^{pro}
(**Extended Data Fig. 2A-2C**). In addition, the piperazine group also forms hydrogen bond with
residue E167 (**Fig. 2**). The increase of van der Waals interactions and the additional hydrogen
bond formation are responsible for the decrease of enthalpy in the structure optimization[36]. To
investigate the binding kinetic properties between ligand and receptor, biolayer interferometry
(BLI) experiment was performed to get the association constant (K_{on}), dissociation constant (K_{off})
and equilibrium dissociation constant (K_d) of GRL0617, GZNL-P4, GZNL-P36 binding to PL^{pro}.
The K_d tested by BLI are consistent with that from ITC, with the value of 8.1 μM, 114.0 nM and
13.9 nM for GRL0617, GZNL-P4 and GZNL-P36, respectively (**Extended Data Fig. 3E-3G**).
The association constant (K_{on}, 1/Ms) are 1.175E5, 8.459E5 and 4.350E6 and the dissociation
constant (K_{off}, 1/s) are 9.562E-1, 9.647E-2 and 6.045E-2, respectively (**Extended Data Fig. 3E，**

F). The increasing K_{on} from GRL0617 to GZNL-P4 and GZNL-P36 are contributed by the
increasing hydrophobic force due to the substitution of naphthalene by a bigger acenaphthylene
group and the additional piperazine group, by the increasing polar contact due to the formation of
hydrogen bond between the piperazine group of GZNL-P4/GZNL-P36 and the residue E167[36].
The change of melt temperature (Tm, data was not shown) determined by differential scanning
fluorimetry (DSF) assay is consistent with K_d determined by ITC and BLI. The enzymatic
inhibition activity is related with the blockade of substrate binding to PL[pro] by inhibitors. GRL0617
bound to PL[pro] mainly hinders the residue L73 of substrate (72RLRGG76), but for GZNL-P4 and
GZNL-P35, piperazine group also impedes the residue R72 of the substrate (**Extended Data Fig.**
2G-2I).

Evaluation of *in vitro* antiviral activity cross coronavirus family

To test whether GZNL-P36 could effectively inhibit PL[pro] across multiple coronavirus
subtypes, we performed a fluorescence resonance energy transfer (FRET) inhibition assay against
PL[pro] proteins from different species coronaviruses (**Fig. 3A, B**). GZNL-P36 exhibited broad
inhibition efficacy against PL[pro] derived from all coronaviruses tested (**Fig. 3A**). The cellular
antiviral activity of GZNL-P36 was examined by a cell protection assay. In this assay, the
cytopathic effect (CPE) of SARS-CoV-2-infected Vero E6 cells with or without treatment by the
compounds was assessed using Celigo Image Cytometer[37]. The cells were challenged with WT
SARS-CoV-2 and two other variants Omicron BA.5 and XBB.1. GZNL-P36 dose-dependently
protected cells from death with 50% effective concentration (EC_{50}) values for wild type (WT),
Omicron BA.5 and XBB.1 is 111.0 nM, 306.2 nM and 58.6 nM, respectively (**Fig. 3C-3E**).
Compared with S-217622 (Ensitrelvir), GZNL-P36 possessed similar effective anti-viral activity
for SARS-CoV-2 and its variants (**Fig. 3C-3E**). Besides SARS-CoV-2, GZNL-P36 also illustrated
anti-viral activity for the other coronaviruses, such as HCoV-NL63 (EC_{50}: 81.6 nM), HCoV-229E
(EC_{50}: 2.66 μM) and HCoV-OC43 (EC_{50}: 46.3 μM) (**Fig. 3F**). Together, our data demonstrated
that GZNL-P36 rendered superb cross-protection against SARS-CoV-2, HCoV-OC43, HCoV-
229E, and HCoV-NL63.

***In vivo* antiviral efficacy of GZNL-P36**

To assess the *in vivo* anti-viral activity of GZNL-P36, we treated the model mice infected with
SARS-CoV-2 by oral administration (**Fig. 4A**). K18-hACE2 transgenic mice aged 8 weeks were
used as our mouse model, forty eight female hACE2 transgenic mice were divided into six groups
with eight mice in each group to evaluate the efficacy of mock, vehicle, positive comparator S-
217622 of 25 milligrams per kilograms (mpk), and GZNL-P36 of 25 mpk, 50 mpk and 100 mpk
in the therapeutic treatment. The weight loss plot shows the about 15% loss of the vehicle group,
but the weight loss is less than 10% for the treated groups (**Fig. 4B**). The lung live viral titers
cannot be detected (**Fig. 4C**) for the group treated with GZNL-P36 at 100 mpk. The groups treated
by GZNL-P36 at the dose of 25 mpk or 50 mpk also showed significant viral titer decrease at 2
days post-infection. The anti-viral efficiency of GZNL-P36 is slightly weaker than the same dose

of positive drug S-217622. Immunohistochemistry assays with SARS-CoV-2 nucleocaspid protein
antibody revealed that abundant expression of viral antigen was identified in the lung of vechile-
treated mice at 4 days post-infection (**Fig. 4D**). In contrast, GZNL-P36 treatment, even when
administered after the virus challenge, markedly suppressed viral nucleocapsid protein expression
in the lung (**Fig. 4D**). Next, haematoxylin and eosin (H&E) stained lungs indicate that the lung
was significantly protected from the GZNL-P36 treatment (**Fig. 4E**).

PL[pro] can dysregulate the host inflammation and antiviral response due to its deubiquitinating
activity. The transcription levels of inflammatory genes, including CXCL10, IFNB1, and IFNγ1,
were determined with the lungs of the mice collected at 2 d.p.i. Compared to the vehicle group,
the transcription level of these pro-inflammatories in GZNL-P36 treated groups was significantly
decreased. Furthermore, the transcription level of CXCL10 and IFNγ1 in the GZNL-P36 treated
groups was also lower than that in the S-217622 treated group (**Extended Data Fig. 6**). These
results indicated that PL[pro] inhibitors may have more benefits on the anti-inflammation properties.

We further perform bulk RNA sequencing on lung samples of all SARS-CoV-2 infected mice.
GSVA analysis results revealed that GZNL-P36 successfully reversed most of SARS-CoV-2-
induced changes, identical to the 3CL[pro] inhibitor S-217622 (**Fig. 5**). More importantly, high dose
of GZNL-P36 greatly reversed the GSVA scores of both WP_FOXP3_IN_COVID19 and
WP_PATHOGENESIS_OF SARSCOV2_MEDIATED_BY_NSPINSP10_COMPLEX genesets,
while S-217622 failed, suggesting our GZNL-P36 might possess synergistic effect on the recovery
of SARS-CoV-2 infected mice comparing with the 3CL[pro] inhibitor S-217622 (**Extended Data
Fig. 7**).

Discussion

Vaccines and neutralizing antibodies cannot provide complete protection against the
continuously emerging variants of SARS-CoV-2. Small molecule antiviral drugs targeting the
conserved viral proteases are particularly important. At present, there are several clinically
available drugs targeting RdRp and 3CL[pro]. Unfortunately, the resistance mutations of RdRp[16,17]
and 3CL[pro22,38-40] have been reported. For PL[pro], another important potential antiviral drug target,
still no targeted drug has been reported. Starting from a weak PL[pro] inhibitor GRL0617, a novel
benzoindolone series of PL[pro] inhibitors was discovered through the utilization of docking-based
library design. Our lead compound GZNL-P36 shows excellent PL[pro] inhibitory potency and
decent pharmacokinetics and *in vitro* safety profile. Furthermore, this compound shows strong *in
vivo* anti-viral efficacy in antiviral mice model suggesting its potential as a COVID-19 therapy
candidate. Interestingly, compared to S-217622, GZNL-P36 treatment showed lower expression
of the pro-inflammatory genes. Our results demonstrate that PL[pro] is an attractive druggable
antiviral target and PL[pro] inhibitor is a class of promising antiviral drug with dual-effect on antiviral
and anti-inflammation.

Methods and Materials

Detailed descriptions of the in vitro pharmacology studies, *in vivo* pharmacology studies, transcriptomics studies, X-ray crystallography, computational study, and synthetic methods can be found in supplementary materials.

Reference

Planas, D. *et al.* Reduced sensitivity of SARS-CoV-2 variant Delta to antibody neutralization. *Nature* **596**, 276-280, doi:10.1038/s41586-021-03777-9 (2021).

Lopez Bernal, J. *et al.* Effectiveness of Covid-19 Vaccines against the B.1.617.2 (Delta) Variant. *N Engl J Med* **385**, 585-594, doi:10.1056/NEJMoa2108891 (2021).

Tuekprakhon, A. *et al.* Antibody escape of SARS-CoV-2 Omicron BA.4 and BA.5 from vaccine and BA.1 serum. *Cell* **185**, 2422-2433 e2413, doi:10.1016/j.cell.2022.06.005 (2022).

Cao, Y. *et al.* BA.2.12.1, BA.4 and BA.5 escape antibodies elicited by Omicron infection. *Nature* **608**, 593-602, doi:10.1038/s41586-022-04980-y (2022).

Yao, L. *et al.* Omicron subvariants escape antibodies elicited by vaccination and BA.2.2 infection. *Lancet Infect Dis* **22**, 1116-1117, doi:10.1016/S1473-3099(22)00410-8 (2022).

Andrews, N. *et al.* Covid-19 Vaccine Effectiveness against the Omicron (B.1.1.529) Variant. *N Engl J Med* **386**, 1532-1546, doi:10.1056/NEJMoa2119451 (2022).

Wang, H., Xue, S., Yang, H. & Chen, C. Recent progress in the discovery of inhibitors targeting coronavirus proteases. *Virol Sin* **31**, 24-30, doi:10.1007/s12250-015-3711-3 (2016).

Li, X. & Song, Y. Targeting SARS-CoV-2 nonstructural protein 3: Function, structure, inhibition, and perspective in drug discovery. *Drug Discov Today* **29**, 103832, doi:10.1016/j.drudis.2023.103832 (2024).

Kokic, G. *et al.* Mechanism of SARS-CoV-2 polymerase stalling by remdesivir. *Nat Commun* **12**, 279, doi:10.1038/s41467-020-20542-0 (2021).

Jayk Bernal, A. *et al.* Molnupiravir for Oral Treatment of Covid-19 in Nonhospitalized Patients. *N Engl J Med* **386**, 509-520, doi:10.1056/NEJMoa2116044 (2022).

Cao, Z. *et al.* VV116 versus Nirmatrelvir-Ritonavir for Oral Treatment of Covid-19. *N Engl J Med* **388**, 406-417, doi:10.1056/NEJMoa2208822 (2023).

Arbel, R. *et al.* Nirmatrelvir Use and Severe Covid-19 Outcomes during the Omicron Surge. *N Engl J Med* **387**, 790-798, doi:10.1056/NEJMoa2204919 (2022).

Mukae, H. *et al.* Efficacy and Safety of Ensitrelvir in Patients With Mild-to-Moderate Coronavirus Disease 2019: The Phase 2b Part of a Randomized, Placebo-Controlled, Phase 2/3 Study. *Clin Infect Dis* **76**, 1403-1411, doi:10.1093/cid/ciac933 (2023).

14   Consortium, W. H. O. S. T. Remdesivir and three other drugs for hospitalised patients
with COVID-19: final results of the WHO Solidarity randomised trial and updated meta-
analyses. *Lancet* **399**, 1941-1953, doi:10.1016/S0140-6736(22)00519-0 (2022).

15   Heyer, A. *et al.* Remdesivir-induced emergence of SARS-CoV2 variants in patients with
prolonged infection. *Cell Rep Med* **3**, 100735, doi:10.1016/j.xcrm.2022.100735 (2022).

16   Hogan, J. I. *et al.* Remdesivir Resistance in Transplant Recipients With Persistent
Coronavirus Disease 2019. *Clin Infect Dis* **76**, 342-345, doi:10.1093/cid/ciac769 (2023).

17   Gandhi, S. *et al.* De novo emergence of a remdesivir resistance mutation during treatment
of persistent SARS-CoV-2 infection in an immunocompromised patient: a case report.
Nat Commun **13**, 1547, doi:10.1038/s41467-022-29104-y (2022).

18   Saravolatz, L. D., Depcinski, S. & Sharma, M. Molnupiravir and Nirmatrelvir-Ritonavir:
Oral Coronavirus Disease 2019 Antiviral Drugs. *Clin Infect Dis* **76**, 165-171,
doi:10.1093/cid/ciac180 (2023).

19   Tan, H., Hu, Y., Jadhav, P., Tan, B. & Wang, J. Progress and Challenges in Targeting the
SARS-CoV-2 Papain-like Protease. *J Med Chem* **65**, 7561-7580,
doi:10.1021/acs.jmedchem.2c00303 (2022).

20   Shimizu, R. *et al.* Evaluation of the Drug-Drug Interaction Potential of Ensitrelvir
Fumaric Acid with Cytochrome P450 3A Substrates in Healthy Japanese Adults. *Clin*
Drug Investig **43**, 335-346, doi:10.1007/s40261-023-01265-8 (2023).

21   Shimizu, R. *et al.* Safety, Tolerability, and Pharmacokinetics of the Novel Antiviral
Agent Ensitrelvir Fumaric Acid, a SARS-CoV-2 3CL Protease Inhibitor, in Healthy
Adults. *Antimicrob Agents Chemother* **66**, e0063222, doi:10.1128/aac.00632-22 (2022).

22   Duan, Y. *et al.* Molecular mechanisms of SARS-CoV-2 resistance to nirmatrelvir. *Nature*
622, 376-382, doi:10.1038/s41586-023-06609-0 (2023).

23   Baez-Santos, Y. M., St John, S. E. & Mesecar, A. D. The SARS-coronavirus papain-like
protease: structure, function and inhibition by designed antiviral compounds. *Antiviral*
Res **115**, 21-38, doi:10.1016/j.antiviral.2014.12.015 (2015).

24   Shin, D. *et al.* Papain-like protease regulates SARS-CoV-2 viral spread and innate
immunity. *Nature* **587**, 657-662, doi:10.1038/s41586-020-2601-5 (2020).

25   Cao, D. *et al.* The SARS-CoV-2 papain-like protease suppresses type I interferon
responses by deubiquitinating STING. *Sci Signal* **16**, eadd0082,
doi:10.1126/scisignal.add0082 (2023).

26   Ratia, K. *et al.* A noncovalent class of papain-like protease/deubiquitinase inhibitors
blocks SARS virus replication. *Proc Natl Acad Sci U S A* **105**, 16119-16124,
doi:10.1073/pnas.0805240105 (2008).

27   Yan, H. *et al.* A robust high-throughput fluorescence polarization assay for rapid
screening of SARS-CoV-2 papain-like protease inhibitors. *Virology* **574**, 18-24,
doi:10.1016/j.virol.2022.07.006 (2022).

28   Ma, C. *et al.* Discovery of SARS-CoV-2 Papain-like Protease Inhibitors through a
339   Combination of High-Throughput Screening and a FlipGFP-Based Reporter Assay. *ACS*
340   *Cent Sci* **7**, 1245-1260, doi:10.1021/acscentsci.1c00519 (2021).

341   29   Shan, H. *et al.* Development of potent and selective inhibitors targeting the papain-like
342   protease of SARS-CoV-2. *Cell Chem Biol* **28**, 855-865 e859,
343   doi:10.1016/j.chembiol.2021.04.020 (2021).

344   30   Zang, Y. *et al.* High-throughput screening of SARS-CoV-2 main and papain-like protease
345   inhibitors. *Protein Cell* **14**, 17-27, doi:10.1093/procel/pwac016 (2023).

346   31   Wang, Q. *et al.* Structure-Based Design of Potent Peptidomimetic Inhibitors Covalently
347   Targeting SARS-CoV-2 Papain-like Protease. *Int J Mol Sci* **24**,
348   doi:10.3390/ijms24108633 (2023).

349   32   Shen, Z. *et al.* Design of SARS-CoV-2 PLpro Inhibitors for COVID-19 Antiviral
350   Therapy Leveraging Binding Cooperativity. *J Med Chem* **65**, 2940-2955,
351   doi:10.1021/acs.jmedchem.1c01307 (2022).

352   33   Yuan, S. *et al.* Targeting papain-like protease for broad-spectrum coronavirus inhibition.
353   *Protein Cell* **13**, 940-953, doi:10.1007/s13238-022-00909-3 (2022).

354   34   Gao, X. *et al.* Crystal structure of SARS-CoV-2 papain-like protease. *Acta Pharm Sin B*
355   **11**, 237-245, doi:10.1016/j.apsb.2020.08.014 (2021).

356   35   Shiraishi, Y. & Shimada, I. NMR Characterization of the Papain-like Protease from
357   SARS-CoV-2 Identifies the Conformational Heterogeneity in Its Inhibitor-Binding Site. *J*
358   *Am Chem Soc* **145**, 16669-16677, doi:10.1021/jacs.3c04115 (2023).

359   36   Liu, W., Jiang, J., Lin, Y., You, Q. & Wang, L. Insight into Thermodynamic and Kinetic
360   Profiles in Small-Molecule Optimization. *J Med Chem* **65**, 10809-10847,
361   doi:10.1021/acs.jmedchem.2c00682 (2022).

362   37   Guo, Y. *et al.* Discovery and characterization of potent pan-variant SARS-CoV-2
363   neutralizing antibodies from individuals with Omicron breakthrough infection. *Nat*
364   *Commun* **14**, 3537, doi:10.1038/s41467-023-39267-x (2023).

365   38   Focosi, D. *et al.* Nirmatrelvir and COVID-19: development, pharmacokinetics, clinical
366   efficacy, resistance, relapse, and pharmacoeconomics. *Int J Antimicrob Agents* **61**,
367   106708, doi:10.1016/j.ijantimicag.2022.106708 (2023).

368   39   Iketani, S. *et al.* Multiple pathways for SARS-CoV-2 resistance to nirmatrelvir. *Nature*
369   **613**, 558-564, doi:10.1038/s41586-022-05514-2 (2023).

370   40   Kiso, M. *et al.* In vitro and in vivo characterization of SARS-CoV-2 resistance to
371   ensitrelvir. *Nat Commun* **14**, 4231, doi:10.1038/s41467-023-40018-1 (2023).

Acknowledgements

We are grateful for support from Shanghai Synchrotron Radiation Facility (SSRF) beamlines (BL19U1 and BL02U1). This study was supported by grants from the National Natural Science Foundation of China (821704730), the startup and R&D Program of Guangzhou National Laboratory (YW-YWYM0202, GZNL2023A02012, GZNL2023A01008, SRPG22-002, SRPG22-003), and the Guangdong Natural Science Foundation (2021QN020451, 2021CX020227), and the Basic and Applied Basic Research Projects of Guangzhou Basic Research Program (202201011795, 2023A04J0161).

Author contributions

Y.L., Q.Y., W.L., M.D., J.T., J.C., J.Z., A.Z., S.Y., K.W., and X.W. performed the cellular and biochemical assays. Y.L. and J.S. performed crystallography. G.Z., P.Z., H.C., P.H., and T.R. performed the medicinal chemistry. Q.Y., W.L., B.X., and J.T. performed the animal models. T.R., M.T., and C.H. performed the computational studies. Y.L., W.L., M.D., J.C., J.Z., K.W., and X.W. performed cloning and/or purified proteins. Y.L., W.L., J.S., and W.K.Z. performed the transcriptome study. Y.L., Q.Y., T.R. X.C., C.H., and J.S. conceived the experiments and wrote the manuscript with input from all authors.

Competing interests

The authors declare no competing interests.

Fig. 1. Rational design of SARS-CoV-2 PL^pro inhibitors and PK profiling. (**A**) Analysis of
potential ligand binding sites S1 and S2 (PDB: 7JRN). The critical hydrogen bonds between
GRL0617 and PL^pro are shown as marine dash lines. GRL0617 and the key residues of the ligand
binding pocket are shown as cyan sticks and wheat sticks, respectively. (**B**) Strategies of structure-
guided compound design. The optimized groups are shown as colored circles, bright orange for
hydrophobicity, and split pea for hydrophilicity. (**C**) Procedures of activity optimization and *in
vitro* PK optimization indicated by the representative compounds. (**D**) The inhibition activity on
SARS-CoV-2 PL^pro of the representative compounds. (**E**)   The *in vivo* PK profiling of GZNL-P35
and GZNL-P36 at 10 mg/kg and 20 mg/kg. (**F**) The *in vitro* stability in liver microsome of the
representative compounds. (**G**) Summary of *in vivo* PK, metabolism, distribution, and toxicity
properties of GZNL-P36.

409
Fig. 2. X-ray crystal structures with SARS-CoV-2 PL^pro inhibitors. X-ray co-crystal structure
of SARS-CoV-2 PL^pro with GZNL-P4 (**A**), GZNL-P28 (**B**), GZNL-P31 (**C**), and GZNL-P35(**D**).
The residues interacting with the ligand are shown as yellow sticks, GZNL-P4 (PDB: 8YX2),
GZNL-P28 (PDB: 8YX3), GZNL-P31 (PDB: 8YX4), and GZNL-P35 (PDB: 8YX5) are shown as
brown sticks. Hydrogen bonds are shown as black dashed lines and the water molecules are shown
as small red spheres. The distances of hydrogen bonds and the residues are labeled.
416

Fig. 3. Pan-antiviral activity of GZNL-P36 against SARS-CoV-2 variants and other coronavirus. (**A**) Enzymatic inhibition efficacy of GZNL-P36 against coronavirus PL^{pro}. (**B**) The phylogenetic tree of coronavirus PL^{pro}s used in this experiment. Antiviral activity of GZNL-P36 against SARS-CoV-2 wild type (WT) (**C**), variants Omicron BA.5 (**D**), and XBB.1 (**E**). Vero E6 cells were pre-treated with indicated compounds with different concentrations for 1 h and then infected with SARS-CoV-2 wild type (WT), variants Omicron BA.5, and XBB.1 at an MOI of 0.01. The EC$_{50}$ was assessed after being cultured for three days. (**F**) The representative inhibition curves of GZNL-P36 against HCoV-229E and HCoV-OC43 in Huh-7 cells. The EC$_{50}$ was assessed after being cultured for two days. Three independent experiments were performed on infections and one representative is shown.

Fig. 4. *In vivo* antiviral activity of PL^pro inhibitor GZNL-P36. (**A**) Experimental design for the
4-day experiment in K18-ACE2 mouse. (**B**) Body weight loss of mice from different groups. (**C**)
Live viral titers in lungs collected at 2 d.p.i. (**D**) and (**E**) Lungs collected at 4 d.p.i. from different
groups were immunostained with SARS-CoV-2 nucleocaspid protein antibody (**D**) or stained with
haematoxylin and eosin (H&E) (**E**). Each dot represents one mouse at the indicated time point.
The data are representative of at least two experiments. The error bars are mean ± SD. Statistical
differences were determined by two-way ANOVA in **C**. ***$P < 0.001$, ****$P < 0.0001$; ns, not
significant.

Fig. 5. RNAseq analysis of GZNL-P36 in SARS-CoV-2 infected Mice. GSVA scores of selected genesets on bulk RNAseq data of GZNL-P36 in SARS-CoV-2 infected Mice. Genesets were obtained from MSigDB and curated based on a criterion of including keywords such as "SARS" or "COVID". The data were then scaled, GSVA scores ranging from -1 (blue, down-regulated) to 1 (red, up-regulated).

Extended Data Table S1 | Representative compounds with enzymatic, antiviral, cell toxicity activity.

Cpd. No.[a]	L	A	R₁	R₂	PLpro IC$_{50}$ (μM)	VeroE6 WT EC$_{50}$ (μM)[b]	VeroE6 XBB EC$_{50}$ (μM)[b]	VeroE6 Omic-BA.5 EC$_{50}$ (μM)[b]	HEK293T CC$_{50}$ (μM)[b]
GRL0617	(R)	naphthalene	–NH₂	–CH₃	4.8200				
GNZL-P1	(R)		–NH₂	–CH₃	2.8310	-	-	-	-
GNZL-P2			–NH₂	–CH₃	6.9000	-	-	-	-
GNZL-P3			–NH₂	–CH₃	0.1858	-	-	-	-
GNZL-P4			piperazine	–CH₃	0.0363	0.4	0.45	0.86	24.57
GNZL-P5			piperazine-NH	–CH₃	0.0210	-	-	-	-
GNZL-P13			N-methyl diazepane	–CH₃	0.0900	-	-	-	-
GNZL-P17			azetidine-piperidine-OH	–CH₃	0.0030	0.233	0.169	0.409	36.68
GNZL-P20			piperidine-N(CH₃)₂	–CH₃	0.1150	-	-	-	-
GNZL-P25			piperazine	–CH₂CH₃	0.4700	-	-	-	-

GNZL-P28				—CH₃	0.0146	-	-	-	-
GNZL-P29				—CH₃	0.0420	0.22	0.24	0.8	12.23
GNZL-P30				—CH₃	0.0440	-	-	-	-
GNZL-P31				—CH₃	0.0085	0.035	0.042	0.174	157.4
GNZL-P35				—CH₃	0.0081	0.04	0.043	0.2	67.67
GNZL-P36				—CH₃	0.0064	0.111	0.058	0.306	88.41

[a] Compound number; [b] "-" represents no data

Extended Data Table S2 | X-ray data collection and refinement statistics

	PL^{pro} in complex GZNL-P4	PL^{pro} in complex GZNL-P28	PL^{pro} in complex GZNL-P31	PL^{pro} in complex GZNL-P35
Data collection	SSRF-BL02U1	SSRF-BL19U1	SSRF-BL19U1	SSRF-BL19U1
Space group	P 2$_1$ 2$_1$ 2$_1$	P 2$_1$ 2$_1$ 2$_1$	I 4$_1$ 2 2	P 1 2$_1$ 1
Cell dimensions				
a, b, c (Å)	60.63, 98.32, 145.69	60.63, 95.44, 145.77	112.5, 112.5, 219.8	46.87, 144.97, 60.1
α, β, γ (°)	90, 90, 90	90, 90, 90	90, 90, 90	90, 99.24, 90
Resolution	49.16 – 2.31	79.85 – 2.60	50.07 – 2.28	25.32 – 1.74
	(2.39 – 2.31)	(2.69 – 2.60)	(2.36 – 2.28)	(1.802 – 1.74)
R_{pim}	0.047 (0.248)	0.035 (0.174)	0.049 (0.51)	0.027 (0.443)
I / σ(I)	6.24 (2.37)	10.61 (3.66)	7.84 (1.49)	15.19 (1.70)
CC1/2 in highest shell	0.933	0.977	0.596	0.772
Completeness (%)	90.10 (98.39)	99.14 (98.60)	99.87 (99.91)	96.00 (73.35)
Redundancy	2.0 (2.0)	2.0 (2.0)	2.0 (2.0)	2.0 (2.0)
Refinement				
Resolution (Å)	2.31	2.60	2.28	1.74
No. of unique reflections	35139	26381	32524	26381
R_{work}/R_{free} (%)	21.04/25.38	21.31/27.51	20.17/23.37	17.27/19.50
No. of atoms				
Protein	4974	4860	2461	4975
Water	185	59	118	539
B-factors				
Protein	44.49	49.09	54.24	28.97
Ligand	35.30	34.90	44.77	43.62
Water	41.04	40.09	45.77	37.98
RMSD				
Bond lengths (Å)	0.013	0.012	0.015	0.011
Bond angles (°)	1.31	1.32	1.30	1.13
Ramachandran favored (%)	94.52	96.65	96.76	97.38
Ramachandran outliers (%)	0.16	0.00	0.00	0.00
PDB accession code	8YX2	8YX3	8YX4	8YX5

Values in parentheses indicate highest resolution shell.

450
451 **Extended Data Fig. 1. Metabolism analysis of GZNL-P4.** (**A**) Metabolic pathway analysis.
452 Sites of Metabolism are highlighted by red color. (**B**) MS detection of metabolites by retention
453 time. (**C**) MS characterization of metabolites by mass-over-charge ratio (m/z). (**D**) UV spectrum
454 detection of metabolites.

Extended Data Fig. 2. The comparison of binding conformations between GRL0617 and GZNL-P4 or GZNL-P35. (**A**) - (**C**) The pattern of GRL0617 (PDB: 7JRN) (**A**), GZNL-P4 (PDB: 8YX2) (**B**), and GZNL-P35 (PDB: 8YX5) (**C**) binding to SARS-CoV-2 PL^{pro}. PLpro is shown as electrostatics surface and the inhibitors (GRL0617, GZNL-P4, GZNL-P35) are shown as sticks. (**D**) - (**F**) The comparison of GRL0617 (**D**), GZNL-P4 (**E**), and GZNL-P35 (**F**) binding to SARS-CoV-2 PL^{pro}. (**G**) - (**I**) Comparison of the PL^{pro} substrate and GRL0617 (**G**), GZNL-P4 (**H**), and GZNL-P35 (**I**) binding to SARS-CoV-2 PL^{pro}. The structures are superimposed using the PL^{pro} protein structure of the co-crystal structures. The substrate peptide RLRGG (PDB: 6YVA), GRL0617 (PDB: 7JRN), GZNL-P4, and GZNL-P35 are shown as green, cyan, yellow, and purple sticks, respectively. The protein is shown as cartoon with the same color as the ligand. (**J**) - (**K**) Comparison of the binding conformation between GRL0617 and GZNL-P4 (**J**) or GZNL-P35 (**K**). Hydrogen bonds are shown as black dashed lines. The distances of hydrogen bonds and the residues are labeled.

Extended Data Fig. 3. The binding constant determination by ITC and BLI. (**A**) - (**C**) ITC
binding curve for the interactions of SARS-CoV-2 PL^pro with GRL0617 (**A**), GZNL-P35 (**B**), and
GZNL-P36 (**C**). (**D**) The ΔG, ΔH, and -TΔH for (**A**) - (**C**). (**E**) - (**G**) BLI binding curve for the
interactions of SARS-CoV-2 PL^pro with GRL0617 (**E**), GZNL-P4 (**F**) and GZNL-P36 (**G**).

Extended Data Fig. 4. The anti-viral activity of the designed compounds against SARS-CoV-

2. Vero E6 cells were pre-treated with indicated compounds with different concentrations for 1 h

and then infected with SARS-CoV-2 WT (**A**), variants Omicron BA.5 (**B**), and XBB.1 (**C**) at an

MOI of 0.01. The Y-axis of the graphs represents the mean inhibition (%) of virus yield of the

compounds. The EC_{50} was assessed after being cultured for three days. Three independent

experiments were performed on infections and one representative is shown.

Extended Data Fig. 5. Evaluation of enzymatic inhibition efficacy of GZNL-P36 against PL^pro

from different species coronaviruses by a fluorescence resonance energy transfer (FRET)

inhibition assay. PL^pros tested in this experiment are from alpha-, beta-, delta-, and gamma-

coronavirus sub-family.

487
488 **Extended Data Fig. 6. The effects of GZNL-P36 on the transcription level of anti-**
489 **inflammatory genes in SARS-CoV-2 infected mice.** Relative mRNA expression of CXCL10 (**A**),
490 IFNB1 (**B**), and IFNγ1 (**C**) of the lungs collected at 2 d.p.i.. Each dot represents one mouse at the
491 indicated time point. The error bars are mean ± SD. Statistical differences were determined by
492 two-way ANOVA. **$P < 0.01$, ***$P < 0.001$; ns, not significant.
493

494
Extended Data Fig. 7. RNAseq results of GZNL-P36 in SARS-CoV-2 infected mice. (A)
Principal component analysis (PCA) of all datasets on count matrix. (**B**) GSVA scores of different
treatment groups revealed a better efficacy of GZNL-P36 on these specific genesets comparing
with the positive control S-217622. P value represented significance among GSVA scores of
indicated datasets in the same geneset. (**C**) Relative expression levels of genes included in genesets
demonstrated in (B). Scale, row-scaled TPM of certain gene (row) in certain sample (column).