TITLE PAGE

ARTICLE TITLE*:* VENTRAL HIPPOCAMPAL AND BASOLATERAL AMYGDALAR INPUTS TO DEEP, BUT NOT SUPERFICIAL LAYERS OF MEDIAL PREFRONTAL CORTEX UNDERLIE STRESS SUSCEPTIBILITY

SHORT TITLE: Synapses Formed by Long-Range Afferents to mPFC Deep Layers Underlie Stress Susceptibility

Keywords: Medial prefrontal cortex, stress susceptibility, ventral hippocampus, basolateral amygdala, synaptic reorganization, eGRASP

Full names of Authors and their Affiliations:

Sabahaddin Taha Solakoğlu[1], Şefik Evren Erdener[1], Olga Gliko, Alp Can, Uygar Sümbül, Emine Eren-Koçak[1,4,5]

1. Institute of Neurological Sciences and Psychiatry, Hacettepe University, Ankara, Turkey
2. Allen Institute for Brain Science, Seattle, Washington, USA
3. Department of Histology and Embryology, Ankara University Faculty of Medicine, Ankara, Turkey
4. Department of Psychiatry, Faculty of Medicine, Hacettepe University, Ankara, Turkey
5. Lead Contact

Information for Lead Contact:

Mailing Address: Hacettepe Universitesi, Norolojik Bilimler ve Psikiyatri Enstitusu, Tip Fakultesi Binasi, Asma kat, Sihhiye, 06100 / Ankara/ Turkey

Phone number: +90 312 305 2130

Email: erenemi@hacettepe.edu.tr

ABSTRACT

Stress response is essential for adapting to an ever-changing environment. However, the mechanisms that render some individuals susceptible to stress are poorly understood. While chronic stress is known to induce dendritic atrophy and spine loss in medial prefrontal cortex (mPFC), its impact on synapses made by long-range projections terminating on the mPFC remains unknown. Here, we labeled synapses on mouse mPFC dendrites formed by ventral hippocampus (VH), basolateral amygdala (BLA) and ventral tegmental area (VTA) long-range afferents using different-colored eGRASP constructs. We obtained multispectral 3D images of the mPFC covering all cortical laminae, and automatically segmented the dendrites and synapses. We investigated the relative abundances and spatial organizations of VH-mPFC, BLA-mPFC and VTA-mPFC synapses on a laminar basis in stress naïve, stress resilient and stress susceptible mice. Our findings demonstrate afferent- and lamina-specific differential reorganization of synapses between different stress phenotypes, suggesting specific roles for different long-range projections in mediating the stress response.

GRAPHICAL ABSTRACT

HIGHLIGHTS

- Synapses between mPFC and its long-range BLA, VH and VTA afferents were mapped separately across cortical laminae, and compared among stress naïve, stress resilient and stress susceptible mice.
- In stress susceptible mice, there was an increase in the abundance of BLA synapses and a decrease in the abundance of VH synapses within the deep layers of mPFC.
- In stress susceptible mice, the number of synapses that originate from different brain regions and converge onto the same dendritic segment in deep layers of mPFC increased.
- In stress susceptible mice, the distances between VH and BLA synapses in deep layers of mPFC were altered.
- This laminae- and afferent- specific reorganization of synapses in stress susceptible mice alter social approach behaviors, presumably by overgeneralizing threat cues.

INTRODUCTION

Stress is an essential adaptive response to imminent or perceived changes in the environment to restore homeostasis. In some individuals, however, stress response can go awry resulting in mental disorders like depression, anxiety disorders and posttraumatic stress disorders for yet unknown reasons. Understanding mechanisms underlying stress susceptibility, therefore, is important for developing prevention and early intervention strategies against stress-related mental disorders.

A key region mediating the stress response is the medial prefrontal cortex (mPFC). Chronic stress causes atrophy of apical dendrites and loss of spines in layer II/III and layer V of mPFC pyramidal neurons without changes in basal dendrites (1-6). These effects of chronic stress are circuit specific: entorhinal cortex-projecting pyramidal neurons show dendritic atrophy in response to chronic stress, whereas basolateral amygdala (BLA)-projecting pyramidal neurons are resilient to stress-induced dendritic remodeling (7). Whether such dendritic resilience to remodeling also applies to afferent projections to the mPFC remains to be answered. Chronic stress-induced dendritic atrophy would be expected to result in the loss of synaptic connections that these dendrites make. If long-range projections to the mPFC from different brain regions are affected differently by this synaptic loss, it will alter the relative weight of inputs from different brain regions on the mPFC and thus, alter the brain circuitry involved in stress response. To date, however, **which mPFC-afferents are affected by dendritic retraction or spine loss is unknown**. Similarly, **whether the reported morphological changes in the dendritic arbor are adaptive or maladaptive remains unknown, because susceptibility to stress was not evaluated in the previous literature**.

mPFC receives input from ventral hippocampus (VH), basolateral amygdala (BLA) and ventral tegmental area (VTA); brain regions indicated in the regulation of the stress response (8, 9). Optogenetic or chemogenetic activation of mPFC or mPFC-projecting VTA, VH and BLA neurons were reported to alter depression- and anxiety-like behaviors (10-19). Unlike sensory cortices, mPFC is agranular, lacks a thalamus projecting layer IV, and receives inputs across all layers (8). Information from various long-range afferents into different layers is integrated by the local circuits within the same layer (intralaminar) or across different layers (interlaminar) to generate an optimal response to adapt to everchanging conditions. **Little is known on how chronic stress alters mPFC circuitry across cortical laminae,** as earlier studies mostly

focused on single layers (20). Evidence indicates that long-range inputs may regulate local circuits selectively since they preferentially target certain cortical layers: Layer II/III and layer V for BLA and VH, respectively (8, 21-23). Distinct mPFC projecting neuron populations identified in both BLA (fear and extinction neurons) and VH (approach and avoidance neurons) regulate stress response and related fear in opposite directions; promoting or suppressing it by integrating the perceptual, emotional and spatial data arriving at mPFC (24, 25). Whether these neurons with opposite functions project to the same or different layers of mPFC is not known. Examining chronic stress-induced changes in specific synaptic inputs across all layers of the mPFC is therefore, crucial to understanding how stress alters brain circuits to produce adaptive responses, as well as what goes awry in stress-related mental disorders.

Here we used the enhanced green fluorescent protein reconstitution across synaptic partners (e-GRASP) system to map synapses between glutamatergic neurons of VH, BLA and VTA and mPFC pyramidal neurons across all layers of the cortical column (26). We obtained multispectral, high resolution 3D images of a ~200 $\times$ 1500 $\times$ 30 μm^3 volume from the mPFC, covering all layers, from stress naïve (SN), stress resilient (SR) and stress susceptible (SS) mice. Mice were divided into SR and SS groups based on their social interaction scores following 10 days of chronic social defeat stress (CSDS). Analysis of the data revealed laminar differences: in superficial layers of mPFC, chronic stress exposure induced similar changes to the relative abundances of BLA-mPFC and VH-mPFC synapses in both SS and SR groups compared to SN mice. In deep layers of mPFC, on the other hand, the abundance of BLA-mPFC synapses relative to all synapses increased, whereas the relative abundance of VH-mPFC synapses relative to all synapses decreased in SS mice compared to SN and SR mice. There were also disruptions to the spatial organization of VH and BLA synapses in deep layers of the mPFC in the SS group. In line with this reorganization of synaptic inputs to mPFC, the percentage of dendritic segments in layer VI that receive input from multiple brain regions was higher in SS mice than the other two groups. Although the percentage of VTA-mPFC synapses was similar across all cortical layers in all groups, we find that the co-presence of a VTA synapse with either BLA or VH synapse on a dendritic segment can predict the stress phenotype. **Our findings reveal an afferent- and lamina-specific differential reorganization of the mPFC synapses in the SS group**. This suggests that disruptions in the long-range afferents to deep mPFC layers may interfer with information processing within the involved circuits, possibly disrupting the discrimination of safety versus threat cues, and thereby play a crucial role in mediating the stress response in health and disease.

RESULTS

Labeling and detection of long-range inputs from VH, BLA and VTA to mPFC pyramidal neurons across the cortical column

To differentially label synapses between VH, VTA and BLA excitatory projections and mPFC pyramidal neurons, AAVs expressing post-eGRASP were injected into mPFC and AAVs expressing cyan, green and yellow pre-eGRASP were injected into VTA, VH and BLA, respectively (Fig. 1a-c). Three weeks after the injections, mice were exposed to 10-day chronic social defeat stress (CSDS) (Fig. 1d). Stress-naive control mice were kept under similar conditions, but with the cage mates kept on the other side of the plexiglass barrier instead of the aggressor mice (Fig. 1e). On day 11, stressed animals were divided into stress susceptible (SS)(n=3) and stress resilient (SR) (n=4) groups according to their social interaction ratio scores (SIR) (Fig. 1f). Stress naïve (SN) mice (n=3) had SIR scores higher than 1, i.e. spent more time around the cage when a social target was present (Fig. 1f)(27). Multispectral, high resolution 3D images were acquired from mPFC across the full length of the cortical column (200 x 1500 x 30 µm^3) (Fig. 1g,h).

Since the emission wavelengths of the synaptic signals overlap, we unmixed the signals with the Learning Unsupervised Means of Spectra (LUMoS) algorithm (Fig. 2, Suppl. Fig. 1) (28). Both dendritic and synaptic signals were then deconvolved, and dendritic signals were segmented with an artificial neural network model (29) and post-processed to obtain reconstructed dendritic segments (Fig. 1f). Reconstructed dendritic tree and synapse locations strongly aligned with their fluorescent signals in 3D (Fig. 2, Suppl. Fig. 2, Suppl. Vid.1-2). The Synapse Detector Algorithm (30) was used for detection and quantification of the synapses between VTA, VH and BLA projections and the mPFC pyramidal neuron dendrites, as manual detection of synapses in such a large 3D volume was not practical. We confirmed that automated synapse detection is highly accurate by comparing the numbers of detected synapses by the Synapse Detector Algorithm with those annotated manually (Suppl. Fig. 3).

The median total numbers of detected synapses were 2,501 (IQR = 1,077 – 6,293; n=3 mice); 7,509 (IQR = 2,560 – 9,177; n=4 mice) and 5,845 (IQR = 789 – 8,598; n=3 mice) in SN, SR and SS mice, respectively. In the SN group, VH-mPFC synapses constituted the highest fraction of all synapses (70.33 ± 16.82%) followed by BLA-mPFC synapses (16.09 ± 11.25%) and VTA-

mPFC synapses (13.57 ± 6.09%). In the SR mice, on the other hand, fractions of VH-mPFC and BLA-mPFC were similar (49.62 ± 17.80% and 38.17 ± 20.37%, respectively; p = 0.69, MWU) followed by VTA-mPFC synapses (12.19 ± 4.32%). The percentages of BLA-mPFC, VH-mPFC and VTA-mPFC synapses were similar in SR and SN groups (p = 0.09, p = 0.15 and p = 0.69, MWU, respectively). In the SS group, the percentage of BLA-mPFC synapses tend to be higher than that of VH-mPFC synapses (54.27 ± 17.18% vs. 25.74 ± 15.12%, respectively; p = 0.055, MWU). The percentage of VTA-mPFC synapses in SS mice was similar to that of the other two groups (19.97 ± 9.79%; p = 0.30, MWU).

Layer and stress phenotype-dependent changes to the ratio of BLA synapses

We then examined the distribution of synapses across each cortical layer. Previous literature showed that BLA and VH axonal projections to the mPFC preferentially target layer II/III and layer V, respectively (8, 21-23). In line with the previous literature, we found that the percentage of BLA-to-all detected synapses (BLA/(BLA+VH+VTA)*100) was higher in layer II/III than layer Vb and VI in SN group (p=0.031, p=0.0079, respectively, MWU test). Chronic stress exposure increased the percentage of BLA synapses in layer II/III compared to SN controls (p= 0.019, MWU test) (Supp Fig. 3). The percentage of BLA synapses in layer II/III was similar in SS and SR groups (Fig. 3b). On the other hand, the percentage of BLA synapses in layer Vb and VI increased only in the SS group and was significantly higher than those in the SN and SR groups. The percentage of BLA synapses in layer Vb and VI showed a significant negative correlation with SIR (Fig. 3f-i).

Layer and stress phenotype-dependent changes to the ratio of VH synapses

Changes observed in VH-mPFC synapses were similar to BLA-mPFC synapses but in the opposite direction. Mice exposed to chronic social stress showed a decrease in the percentage of VH-to-all synapses [VH/(BLA+VH+VTA)*100] in layer II/III compared to SN controls (p=0.019, MWU test) (Supp Fig. 3). We did not observe any differences in the the percentage of VH synapses between SS and SR groups. In contrast, only SS mice showed a significantly decreased percentage of VH synapses in layer Vb and VI (Fig. 3c). The percentages of VH synapses in layer Vb and VI were significantly correlated with SIR (Fig. 3j-m).

Invariance of the ratio of VTA synapses to the stress phenotype

The VTA-to-all synapses percentage (VTA/(BLA+VH+VTA)*100), was similar across all layers of the mPFC in all groups (Fig. 3d). We did not find statistically significant correlations between SIR and the percentages of VH synapses across different layers (Fig. 3n-r).

A low dimensional representation of the relative abundances of VH, BLA and VTA synapses across the cortical column visually differentiates SS mice from SR mice.

To further examine the dissociation of groups according to their stress susceptibility, we examined the low-dimensional structure of synaptic distributions across all layers using principal components analysis (PCA) in 13 cortical columns with an intact layer I (2 columns lacking layer I were excluded from the analysis, due to damage, presumably from the AAV injections). We converted the 3x13x5 (synapse type count x column count x layer count) synapse percentage matrix into a 13x15 matrix by arranging the synaptic percentage data from VTA, VH and BLA from each layer into a single row. We then applied PCA to this matrix and visualized the distribution of cortical columns in the 2-dimensional plane based on the top two principal components (PCs). Approximately 77.44% of the variance in the synaptic percentages across the cortical column was captured by these two PCs. Notably, the cortical columns associated with stress susceptibility occupied distinct locations in this low dimensional space (Fig. 3s). This supports the finding that relative synaptic abundances of BLA, VH and VTA across the mPFC laminae can explain the behavioral differences between SR and SS mice.

The abundance of dendritic segments receiving input from multiple brain regions increases in SS mice.

To gain a deeper insight into differences of SS mice from others, we investigated the percentage of reconstructed dendritic segments that have 2 or more synapses in layer Vb and VI (Fig. 4). In both layer Vb and layer VI, the percentage of dendritic segments receiving input from only VH was significantly lower and those receiving input from only BLA tended to be higher in the SS mice than the other two groups. There was no difference in the percentage of dendritic segments receiving input from only VTA among groups (Suppl. Fig. 4).

In layer Vb, the percentage of dendritic segments receiving input from multiple brain regions was similar between groups (Fig. 4b,c). When analyzed separately for each pair of multi-brain region originated synapses on individual dendritic segments, however, we observed that the percentage of BLA-VTA synapses increased in both stressed groups; the rise in the SS group was significantly higher than that of the SR group (Fig. 4d,e). In layer VI, on the other hand, the

percentage of dendritic segments receiving input from multiple brain regions was significantly higher in the SS group than the other two groups (Fig. 4g,h). When analyzed separately for each pair, we observed a reduction in the percentages of VTA-VH synapses in the SS group (Fig. 4i,j). Interestingly, we did not observe any difference in the percentage of BLA-VH synapses among groups in either layer Vb or layer VI (Fig. 4e,j).

These findings suggest that under physiological conditions (i.e. in case of an adaptive stress response) stress response alters synapses in deep layers by changing the number of synapses originating from the same brain region on a single mPFC dendritic segment, and that increases in the percentage of multi-origin-synaptic pairs on the same dendritic segment in SS mice may cause a disorganization in the outputs of these layers.

The spatial organization of VH and BLA synapses on mPFC deep layers is disrupted in SS mice.

To elucidate whether there is a general innervation pattern of the mPFC by VH and BLA in layer Vb and VI, we measured the Euclidean distances between VH and BLA synapses and determined the nearest VH and BLA synapses for each VH and BLA synapse in the reconstructed ~200 x 700 x 30 μm^3 mPFC volume (Fig. 5a, b). Among the three groups studied, there were no differences in either the distances of BLA synapses to their corresponding closest BLA synapses or the distances of VH synapses to their corresponding closest VH synapses (Fig. 5c, d). We observed differential spatial organization of VH and BLA synapses relative to each other in SN/SR mice vs. SS mice: In the SN and SR mice, the distances of VH synapses to their corresponding nearest BLA synapses were distributed relatively uniformly across a wide range (Fig-5e), whereas the BLA synapses were preferentially located in close proximity (within 5 um) to VH synapses (Fig. 5d, f). In contrast, in the SS group, we observed a statistically significant decrease in the distances of VH synapses to their corresponding closest BLA synapses and a statistically significant increase in the distances of BLA synapses to their corresponding closest VH synapses (Fig. 5e-h).

These results suggest that the long-range projections follow structured, differential innervation patterns and are not random. We quantified this observation by training a random forest classifier (RFC) on a 42,317 x 8 matrix of nearest synaptic distances, where each row corresponds to a synapse. In each row, the first four columns denote the Euclidean distances of that synapse to its four closest VH synapses, and the last four columns denote the distances to

its four closest BLA synapses (Figure 5). This RFC had 76% accuracy in differentiating all three groups (Fig. 5i). Examining the confusion matrix reveals that the RFC can predict the SS group with ~88% accuracy and the SR group with ~83% accuracy. However, the prediction accuracy of the SN group is low (~47%) (Fig. 5j). We observed that the classification accuracy and the F1 score (harmonic mean of precision and recall) decreases significantly when the RFC is trained on the same dataset with shuffled labels, simulating a scenario where synapses from the different regions are randomly distributed in the mPFC over the observed synaptic locations (Fig. 5i-k). Finally, we visualized this result using uniform manifold approximation and projection (UMAP) (31), a non-linear dimensionality reduction method: the synapses associated with the SR and SS phenotypes occupy different regions of the two-dimensional UMAP map of the same data matrix (Fig. 5l,m).

DISCUSSION

While the stress response is a fundamental component of the behavioral repertoire of animals, a mechanistic, circuit-level understanding of it in health and disease remains incomplete. This is, in part, due to the complexity of the distributed circuitry that governs the behavioral responses, involving numerous synaptic connections between multiple brain regions. Here, by using multiple eGRASP constructs, multispectral volumetric imaging, computational tools for dendrite and synapse identification, and statistical methods for analyzing the resulting large-scale dataset, we aimed to elucidate this circuitry in the mouse. To our knowledge, this is the first study that mapped VH-mPFC, VTA-mPFC and BLA-mPFC synaptic connections separately across all cortical layers and compared them between SN, SS and SR mice. This multi-pronged approach revealed distinct patterns of organization of synaptic inputs to deep and superficial mPFC layers in the stress response and susceptibility.

Previous literature showed that BLA projections preferentially target layer II (21, 22, 32), whereas, VH projections are largely restricted to layer V of mPFC (22, 23). Consistent with these findings, we observed a similar organization of long-range synaptic inputs to the cortical layers of SN mice. CSDS altered the ratios of BLA-to-all synaptic inputs [BLA/(BLA+VH+VTA)] and VH-to-all synaptic inputs [VH/(BLA+VH+VTA)] in mPFC superficial layers, regardless of stress susceptibility of the mice. Organization of long-range inputs to deep layers of mPFC, on the other hand, was disrupted only in the SS group: BLA-mPFC synapses predominated over VH-mPFC synapses in layer Vb and VI suggesting that the reorganization of long-range synaptic inputs to mPFC deep layers may underlie stress susceptibility. In contrast, the ratio of VTA-to all synaptic inputs [VTA/(BLA+VH+VTA)] was similar across all cortical layers and groups.

All mice exposed to CSDS showed an increase in the abundance of BLA-mPFC synapses and a reduction in the abundance of VH-mPFC synapses in layer II/III, indicating that changes observed in this layer are related to stress exposure rather than stress susceptibility. Previous studies reported reductions in dendritic length and spine numbers in mPFC layer II/III after chronic stress exposure (1-5). Our results, when considered together with these findings, suggest that mPFC neurons that receive input from VH were preferentially atrophied by chronic stress resulting in a decrease in VH inputs, whereas those that receive input from BLA were unaffected. Supporting this suggestion, Shansky et al (2009) demonstrated that pyramidal neurons projecting to entorhinal cortex showed dendritic atrophy in response to chronic stress,

whereas those projecting to amygdala were resilient to stress-induced dendritic remodeling (7). Considering that there is a reciprocal connection between BLA and mPFC (i.e. BLA inputs preferentially innervate BLA-projecting mPFC neurons), one may speculate that BLA input receiving mPFC neurons are also resilient to atrophy (21, 22, 32, 33).

Brain networks connecting mPFC, BLA, and hippocampus are involved in the regulation of fear-related and social behaviors(14, 17, 34-37). The interplay between these three regions in the control of anxiogenic/social behaviors is complex, and may both promote and suppress fear or social behavior (17, 22, 24, 33, 38). Input from BLA and VH to the mPFC were reported to have opposite effects on fear-related and social behaviors(14, 17, 39). Moreover, studies have demonstrated that within each of these brain regions there are distinct populations of neurons with distinct connections that encode "safe" and "aversive" cues. This allows the organism to react optimally to various stimuli (24, 25, 33). Selective activation of distinct neuronal circuits may serve as a mechanism to switch between alternative behavioral responses by integrating sensory, contextual and emotional information, fine-tuning the expression and suppression of fear-related and social responses (24, 38, 40).

Mice are social animals and they tend to spend more time with social targets (stranger mouse) compared to non-social targets (i.e. objects) (41). Following CSDS, SS and SR mice demonstrate opposite behavioral responses when encountering a novel mouse in a novel context: SS mice exhibit social avoidance, while SR mice demonstrate social approach. This suggests that SR mice can discriminate threat-related stimuli or contexts from novel stimuli or contexts, whereas SS mice tend to overgeneralize the threat to all novel stimuli or contexts they encounter. Overgeneralization of fear is a characteristic feature of anxiety disorders, and thought to be one of the mechanisms underlying posttraumatic stress disorder, panic disorder and generalized anxiety disorder (42-45). The increase in the relative abundance of long-range inputs from BLA and the decrease in the relative abundance of long-range inputs from VH to deep layers of the mPFC pyramidal neurons in SS mice, as reported in the current study, could potentially interfere with the discrimination between threat-associated and non-threat associated stimuli. This interference may then disrupt the ability of the mPFC to produce an adaptive response. Intriguingly, a previous study showed that the discrimination of aversive cues from safe cues was associated with synchrony between mPFC and BLA activities in the theta frequency range (38). The entrainment of BLA activity by mPFC activity in the theta frequency range was reported in only those mice that can discriminate conditioned auditory tones from unconditioned tones (discriminators). Specifically, these mice demonstrated a significantly

higher degree of freezing in response to conditioned auditory tones compared to unconditioned tones (38). Additionally, the VH also synchronizes with mPFC activity at the theta frequency range during the exploration of anxiogenic novel environments, facilitating the discrimination of safe and aversive components of the environment (18, 19, 40). VH is also implicated in the formation and storage of social memory (36). VH neurons are more strongly activated in response to a familiar mouse compared to a stranger mouse, and their optogenetic inhibition interferes with the discrimination of familiar and unfamiliar mice (36). In parallel with these findings, our results suggest that the balance between BLA and VH inputs to the mPFC is crucial in associating social cues with aversion, and in discriminating safe cues from aversive cues to produce an adaptive response (Please see graphical abstract). As we did not investigate the efferent connections of mPFC, it remains unclear which brain region is impacted by the reported disorganization of long-range inputs to deep layers of mPFC. However, existing literature indicated the involvement of mPFC projections to nucleus accumbens (NAc) in social approach behavior (46). Considering that the organization of long-range inputs in SS mice was impaired only in the deep layers of the mPFC, which predominantly project to subcortical regions, it is possible that imbalance in BLA and VH inputs to mPFC deep layers may disrupt the activity in mPFC-NAc circuit. This issue warrants further investigation in future studies.

How excitatory inputs from different brain regions onto mPFC contribute to different stress phenotypes is an intriguing question. One possibility is that these inputs may selectively target specific neuronal ensembles, each with distinct roles in fear-related behavior. Supporting this suggestion, our findings showed that inputs from the same brain region tend to make synapses on the same dendritic segments in SN and SR mice. In SS mice, however, we observed increase in the number of inputs from different brain regions converging onto the same dendritic segment. Additionally, our data indicates that under normal physiological conditions, inputs from the BLA tend to terminate in close proximity to VH inputs, but it appears crucial for some VH-mPFC synapses to be located farther from BLA-mPFC synapses. In SS mice, this innervation pattern is disrupted with almost all VH inputs terminating close to BLA-inputs. One may speculate that proximity of BLA inputs to VH inputs play a role in integrating threat-signals into ongoing information processing within those neuronal ensembles, while VH inputs distant from BLA inputs may be involved in processing non-threat-related information. Whether these distinct inputs originate from distinct mPFC-projecting VH neuronal populations, each with opposite roles on fear-related behaviors, remains to be addressed in future studies(25). Interestingly the disorganization in VH-BLA innervation of deep mPFC layers was not observed at the level of

single dendritic segments, as there were no differences in the percentage of VH-BLA inputs on single dendritic segments among all three groups.

Although previous studies have indicated the role of VTA in regulating both stress response and social behavior, we did not observe any changes in the ratio of VTA-to-total synaptic inputs either following stress exposure or as a result of stress susceptibility. This discrepancy may be attributed to the fact that most of these studies focused on dopaminergic neurons, whereas we studied glutamatergic VTA projections to the mPFC. Another plausible explanation is that VTA projections may have synapsed preferentially on interneurons, which we did not study. Long-range projections on mPFC interneurons may also contribute to fine-tuning the stress response, given various reports of altered excitatory:inhibitory balance in stress-related mental disorders (47-51). Intriguingly, we observed that in dendritic segments receiving input from multiple brain regions, change in VTA input-containing synaptic pairs correlated with stress susceptibility. This suggests that VTA inputs may also play a role in the regulation of stress response. Further investigation of the involvement of VTA glutamatergic projections in stress response, particularly focusing on VTA-mPFC-synapses on interneurons, is warranted in future studies.

One limitation of our study is that it was conducted only in male mice, thereby restricting the generalizability of our findings to females. CSDS is generally used in identification of stress susceptibility in male mice due to challenges in implementing it in females. Resident mice of either sex do not attack the intruder females, so, unlike male mice, females do not naturally face such social attacks. Although a CSDS paradigm has been developed to induce social defeat in females by applying another CD1 male mouse's urine to vaginal orifices of female intruders (52), we chose to conduct our study in male mice to explore the mechanisms underlying stress susceptibility and resilience, given its higher ethological relevance in males.

Additionally, our study primarily focused on long-range projections to pyramidal mPFC neurons, and we did not investigate projections terminating on mPFC interneurons. This decision was influenced by the challenges associated with confocal multicolor imaging, particularly the quantitative limit for visualizing multiple fluorophores simultaneously. However, future studies could explore the role of long-range projections to mPFC interneurons to provide a more comprehensive understanding of the neural circuitry underlying stress response and resilience.

In conclusion, the findings of the current study enhance our comprehension of chronic stress effects on brain circuitry in both physiological and pathological states by focusing on the

synapses formed by long-range projections on mPFC. We have demonstrated laminae- and afferent- specific changes in stressed groups depending on their stress phenotype (resilient vs susceptible), highlighting potential avenues for developing preventive or therapeutic strategies in stress susceptible individuals following a stressful or traumatic event.

METHODS

Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Bacterial and virus strains		
AAV2/1-CWB-pre-eGRASP(p32)	Charite – Universtatmedizin Berlin Viral Core Facility	#VCA-340a
AAV2/1-EWB-DIO-myrTagRFP-T-P2A-post-eGRASP	Charite – Universtatmedizin Berlin Viral Core Facility	#VCA-338a
AAV2/1-CWB-cyan pre-eGRASP(p30)	Charite – Universtatmedizin Berlin Viral Core Facility	#VCA-339a
AAV2/1-CWB-yellow pre-eGRASP(p32)	Charite – Universtatmedizin Berlin Viral Core Facility	#VCA-337a
AAV2/1 – CaMKIIα - iCre	Charite – Universtatmedizin Berlin Viral Core Facility	#BA-257b
Critical commercial assays		
QIAGEN Plasmid Mega Kit	Qiagen	Cat. #12181; RRID:SCR_008539
Experimental models: Organisms/strains		
C57BL/6	Hacettepe University Laboratory Animals Research and Application Centre (HUDHAM)	N/A
CD-1	Kobay Deney Hayvanları Laboratuvarı	N/A
Recombinant DNA		
pAAV-CWB-pre-eGRASP(p32)	Choi et al.(26)	Addgene #111598; RRID:Addgene_111598
pAAV-EWB-DIO-myrTagRFP-T-P2A-post-eGRASP	Choi et al.(26)	Addgene #111581; RRID:Addgene_111581
pAAV-CWB-cyan pre-eGRASP(p30)	Choi et al.(26)	Addgene #111586; RRID:Addgene_111586
pAAV-CWB-yellow pre-eGRASP(p32)	Choi et. al.(26)	Addgene #111580; RRID:Addgene_111580
Software and algorithms		

Fiji is just ImageJ 2.1	Schendelin et al.(53)	https://fiji.sc
Vaa3D – x 1.1.2	Peng et al.(54)	https://github.com/Vaa3D; RRID:SCR_002609
BigStitcher	Hörl et al.(55)	
ZEN 3.2 Black Edition	Zeiss	https://www.zeiss.com/
LUMoS	McRae et al.(28)	https://imagej.net/plugins/lumos-spectral-unmixing
RedLionfish (RL) deconvolution	Perdigao et al.(56)	https://github.com/rosalindfranklininstitute/RedLionfish
Neuron Reconstruction Algorithm	Gliko et. al.(29)	N/A
IS Detector (Synapse Detector)	Iascone et al(30)	https://github.com/Vaa3D
Other		
LSM810 laser-scanning confocal microscope	Zeiss	https://www.zeiss.com/
Leica SP8 confocal microscope	Leica	https://www.leica-microsystems.com/
Allen Brain Atlas	Allen Institute for Brain Science	http://mouse.brain-map.org

Animals

Ten 12–16-week-old male C57BL/6 wild-type and 4–20-month-old CD-1 male wild-type mice were used for the study. Mice were maintained at 12:12 light:dark cycle (07:00-19:00) and stable temperature (19-22 ^{o}C) with *ad libitum* access to food and water. Mice were purchased from Kobay AS and Hacettepe University Laboratory Animals Research and Application Centre, and housed at the Animal Unit of the Institute of Neurological Sciences and Psychiatry during the experiments. All procedures were done in accordance with Animal Experimentations Local Ethics Board of Hacettepe University (Ethics number: 2019/78).

Construction of Adeno Associated Virus

eGRASP plasmids to label synaptic connections from 3 different brain regions onto mPFC dendritic tree; specifically, pAAV-CWB-yellow-pre-eGRASP(p32), pAAV-CWB-cyan-pre-eGRASP(p30), pAAV-CWB-green-pre-eGRASP(p32), pAAV-EWB-DIO-myrTagRFP-T-P2A-post-eGRASP were obtained from Addgene (#111580, #111581, #111586, #111598, gifts from Dr. Bong-Kiun Kaang)(26). Plasmids were purified by Qiagen Plasmid Mega kit and then sent to Viral Core Facility of Universtatmedizin Berlin Charite to be packaged into AAV 2/1. AAV2/1-CaMKIIa-iCre was obtained from the same facility.

Intracranial Virus Injections

Mice were anesthetized with intraperitoneal (ip) injection of 5-10 mg/kg xylazine and 50-100 mg/kg ketamine, then maintained under 1-2% isoflurane or sevoflurane anesthesia. AAV2/1-pAAV-CWB-yellow-pre-eGRASP(p32), AAV2/1-pAAV-CWB-green-pre-eGRASP(p32) and AAV2/1-pAAV-CWB-cyan-pre-eGRASP(p30) were injected into BLA, VH and VTA, respectively. AAV2/1-CaMKIIa-iCre and AAV2/1-pAAV-EWB-DIO-myrTagRFP-T-P2A-post-eGRASP were injected into mPFC. Injections were performed by lowering 28 gauge microinjectors unilaterally into right VH, VTA, BLA and mPFC sequentially (coordinates from bregma are AP: -1.2, ML: -3.42, DV: -5.4 **for BLA**; AP: +1.3, ML: -0.3, DV: -2.3 **for mPFC**; AP: -3.05, ML: -3.35, DV: -4.2 **for VH** and AP: -2.87, ML: -0.35, DV: -5.1 **for VTA**). AAV2/1 injections were carried out at a rate of 0.1 µL/min, details of viral genome densities and volumes injected in each region are summarized in Supplemental Table-1. Microinjectors were kept in the injection sites for 10 minutes to allow the diffusion of AAVs. Craniotomies were then filled with bone wax; scalp was sutured and subcutenous flunixin meglumine (2.5 mg/kg) and ip saline (0.3 ml) was administered. Eye ointment was frequently applied and rectal temperature was monitored during the procedure.

Experimental Timeline

After AAV2/1 injections, stress exposed mice were left undisturbed for 3 weeks to allow for optimal expression of the virus. After 10-day chronic social defat stress (CSDS), social interaction (SI) test was performed to determine stress phenotype (susceptible vs resilient). Stress-naïve mice were kept under similar conditions for 4 weeks except with another C57BL/6 mice separated by a plexiglass barrier. After the SI test (within ~ 1h), the brains were removed for processing.

Chronic Social Defat Stress and Social Interaction Test

Chronic Social Defat Stress (CSDS) was carried out as Golden et al described (27). Briefly, an intruder C57BL/6 male mouse was placed in the cage of an aggressive CD-1 mouse. Duration of physical interaction was determined with an algorithm for each mouse based on the occurrence of bleeding to prevent injury. Then the C57BL/6 mouse was physically separated from the CD1 mouse by dividing the cage with a plexiglass barrier that allowed for visual, olfactory and vocal interaction. They were housed in the divided cage for 24 hours, then the stress protocol was repeated for 10 days with a different CD-1 mice. On the 11[th] day social

interaction test (SI test) was carried out. Briefly, the interaction time of C57BL/6 mice with an interaction cage (9.5 cm in diameter) placed at one edge of a 45 x 45 cm arena was recorded with Ethovision XT 8.0. The SI Test consisted of two stages. During the first stage, mice explored the arena that contains an empty interaction cage for 5 minutes. During the second stage, an unfamiliar novel CD-1 male mouse was put in the interaction cage and C57BL/6 mice explored the same arena for 5 minutes.

The interaction zone was defined as the circular area with a 23.75 cm diameter sharing the same center with interaction cage. Social interaction ratio was calculated by dividing the time spent around the interaction cage when CD-1 mouse was present to the time spent around the interaction cage when interaction cage was empty. Mice were classified as stress-resilient if the SI ratio is above 1 and stress-susceptible if the SI ratio is below 1.

Tissue Preparation

After the SI test, mice were anesthetized with %10 chloral hydrate-saline solution and perfused with heparin and 4% PFA. Brains were extracted and kept at $+4^{o}C$ in 4% PFA solution overnight. To minimize PFA-induced autofluorescence, the brains were washed in cold phosphate buffered saline (PBS) for 3 days. Then the brains were immersed in 30% sucrose solution for 2-5 days and kept at -80°C until sectioning. Free floating 60 μm thick sagittal sections were obtained and mounted with 1/1 : glycerol/PBS solution.

Imaging of Synapses and Dendrites

High resolution, multispectral, 3D tile scans were obtained by a Zeiss LSM810 laser-scanning confocal microscope, using an oil objective (40✕, NA: 1.3). x, y pixel and z-step sizes were determined for adequate acquisition of cyan eGRASP signal which has the lowest emission wavelength among other signals. Therefore, we adjusted the pixel size as 0.103 μm. Then, 30 μm-thick overlapping (10%) tile scans with field of view (FOV) of 205 ✕ 205 $μm^2$ were obtained, spanning the full cortical thickness from surface to corpus callosum. For multispectral imaging, the same FOVs were imaged using 453, 488, 543 and 633 nm laser excitations. Emissions of fluorescent proteins were bandpass-filtered at 465-490 nm for cyan eGRASP under 453 nm excitation; 495-515, 515-535 and 535-555 nm for green and yellow eGRASP under 488 nm excitation; 550-630 nm for TagRFP-T under 543 nm excitation and 645-750 nm for autofluorescent signals under 633 nm excitation.

Reflectance Imaging

Confocal reflectance imaging was performed to determine the boundary between layer VI and corpus callosum. Images were obtained in tile scans by a Leica SP8 confocal microscope. Briefly, a narrow band reflectance signal was obtained by excitation at 488 nm and emission at ~483-493 nm, using a RT15/85 beamsplitter. Z-stacks were obtained with 2-3 μm step size. The pixel size for this setting was ~0.6 μm. Dendritic TagRFP-T labelled signals detected by 552 nm excitation were overlaid on the reflectance signal. Finally, images were stitched via stitching algorithm of LASX v3.5.5.19976.

Image Processing

Stitching: Acquired tiles were stitched together using the Zeiss Stitching algorithm (ZEN 3.2 Black Edition) or BigStitcher plug-in of FIJI (55). The correlation threshold was kept above "0" for Zeiss Stitching algorithm. For BigStitcher application, x1 downscaling was applied to prevent loss or duplication of signals in the overlapping regions.

Unmixing: Signals from dendrites and synapses overlapped in the acquired images. Unmixing both signals at the same time can cause interruptions to dendritic continuity, which will interfere with reconstruction. Therefore, we unmixed dendritic and synaptic signals separately by LUMoS plug-in of FIJI (28). Dendritic TagRFP-T signals were unmixed from the autofluorescent signals, whereas the synaptic yellow/green/cyan-eGRASP signals were unmixed from each other and auto-fluorescent signals (Fig.1-a,c). The unmixing process was repeated at least 50 times with 100 iterations. The reference image was selected from tile images and axial slices, where all signal subgroups (from all fluorophores as well as nonspecific autofluorescence) were observable within the field of view.

Deconvolution: To be able to generate a 3D reconstruction of the dendrites and synapses, deconvolution of the synaptic and dendritic signals was performed using GPU-based Richarson-Lucy algorithm (56). First, we prepared images by making the voxels isotropic via Reslice Z function in FIJI. Then point spread function (PSF) with the size of 256 x 256 x N, where N is the number of z-stacks of the tile image, was prepared by using Diffraction PSF 3D plug-in of FIJI. Optimum wavelengths of eGRASP proteins and TagRFP-T were used for PSF generation (57). Finally, Richardson-Lucy deconvolution algorithm with 20 iterations was applied for all signals. Synaptic signals were further processed by Gaussian filtering with a window size of 3 pixels. Stitched (BigStitcher) and unmixed (LUMoS) images were processed in Fiji.

Reconstruction: Synaptic and dendritic images were merged according to their tile locations. We reconstructed the dendrites using the automated approach in Gliko et al (2024) that combines U-Net convolutional neural network segmentation with post-processing to produce a digital reconstruction. The segmentation model that was trained with a set of brightfield images of mouse cortical excitatory neurons (29) was used without any tuning. Segmentation was post-processed by thresholding the background, connected component analysis to remove short segments, skeletonization, and conversion to the .swc format, followed by pruning and down sampling. We did not determine the diameters of the dendritic branches. We also did not classify the dendrites into apical, basal, and oblique compartments since the directionality with respect to the soma could not be determined during imaging.

Definition of Dendritic Segments

Dendrites of all mPFC pyramidal neurons across all layers were labelled with TagRFP-T. Dendritic signals were intermingled and it was hard to decide if they belonged to the same dendritic tree or not, owing in part to the neuronal organization of mPFC and diffraction limit (0.25 - 0.35 μm) of confocal microscopy, and in part to the erroneous linkages between dendritic segments introduced by dendritic reconstruction. Therefore, we followed a conservative strategy and defined dendritic segments as part of the reconstructed tree between two adjacent branching or leaf nodes and evaluated the synapses only on these short dendritic segments.

We excluded the segments resulting from image edge artefacts during quality control. Additional quality control was applied to remove (spurious) segments of 0 μm length.

Synapse Detection

BLA-mPFC, VH-mPFC and VTA-mPFC synapses were detected by the Synapse Detector plug-in of Vaa3D (30), which performs synapse detection via adaptive watershed segmentation within a given distance from the dendritic reconstruction. We ran this plug-in by the following five steps:

1- We thresholded deconvolved synaptic and segmented dendritic signals. Threshold values for synapses were determined by comparing raw synapse data and deconvolved synapse data (Fig. 2g, Suppl. Fig. 1). Threshold values for dendritic images were determined to avoid discontinuities along dendritic branches.

2-   We prepared the 3D 8-bit RGB stacks that the Synapse Detector requires as input as follows:

$$I = [I_R, I_G, I_B] \#(1)$$

$$I_R = I_{dendrite} \cup I_{synapse} \#(2)$$

$$I_G = I_{synapse} \#(3)$$

$$I_B = 0 \#(4)$$

I_R, I_G and I_B indicate 8-bit images, sized H x W x Z (height x width x depth of the image stack). $I_{dendrite}$ indicates the segmented dendritic image stack, $I_{synapse}$ indicates the synapse image of size H x W x Z. I_R is the union of dendritic ($I_{dendrite}$) and synaptic image ($I_{synapse}$) stacks. I_B is a stack of all zeros since the the algorithm writes the detected synapse information on this third channel after detection.

3-   Synapse Detector has 4 main parameters: minimum signal density threshold values for dendritic and synaptic images, minimum synapse volume and maximum search distance from .swc tree. We did not use the signal density threshold values for dendritic and synaptic images as we applied thresholding in step 1. We limited the analysis to those synapses detected within ≤ 20 pixels (~ 2 µm) distance around the swc tree[58] and those occupying at least 30 voxels (~ 0.030 µm³) as the diffraction limit (0.25 - 0.35 µm) requires a detectable signal to occupy at least 3 pixels space (~ 0.3 µm) along each dimension.

4-   The Graphical User Interface (GUI) of the Synapse Detector allows the user to examine the synapses segment-by-segment, include/exclude detected synapses from the analysis, adjust the detected synapse volumes, and annotate synaptic positions as dendrite-targeting or spine-targeting. While human-annotated synaptic mapping is the gold standard way of input mapping, such manual inspection is infeasible for large datasets. Therefore, we visually examined the automatically detected synapses using only a zoomed-out view within the GUI of Synapse Detector. If most of the synapses have been tagged as detected, we have extracted the outputs of the algorithm in .csv format.

5-   We repeated the above 4 steps for each of the three different colored synapse separately.

The outputs of Synapse Detector Used in this study are:

-       3D spatial locations of detected synapses

-       Volumes of the detected synapses

-       For each synapse, the closest node on the dendritic segment from the detected synapse, which also includes dendritic segment and tree information.

Comparison Between Automatic and Manual Counting Results

We checked the accuracy of the Synapse Detector in automatic synapse detection by two means:

1- In randomly selected 3D stacks (n = 49 images with a size of ~600 x 2000 x 300 from 15 cortical columns) from different layers, we confirmed each automatically detected synapse by visually inspecting and locating it in the unmixed images. Visual comparison was performed in Fiji by comparing detected synapses and the unmixed synaptic signals. Subsequently, the percentage of matched synapses among the total number of automatically detected synapses was calculated (Suppl. Fig. 3-a, b). This method suggests that 95.7% of automatically labeled synapses were truly synaptic signals.

2- We manually counted all three types of synapses (formed by projections from BLA, VH and VTA) in randomly selected 3D frames from different layers on 3D visualization module of Vaa3D. Only signals within ~2 μm from the reconstruction tree were manually counted, as Synapse Detector was set to detect synapses within ~2 μm of a reconstructed dendrite. After manual labeling of synapses, spatial locations of the synapses were saved as a ".marker" file. We then computationally matched the spatial locations of manually detected synapses with automatically detected synapses using the Hungarian algorithm. The Hungarian algorithm creates match results for the group with the fewer number of synapses. We declared that manually detected and automatically detected synapses are matched when the match distance is less than 2 μm. Then, we quantified the accuracy of synapse detection separately for different synapse types and behavior phenotypes (Supp. Fig. 3c-f).

In all cases, we found that automated synapse detection is highly accurate: First, we found that, on average, 76% of the synapses annotated manually by an expert were detected by the Synapse Detector software (Supp. Fig. 3c). On the flip side, we initially found that, on average, 20% of the synapses detected by the Synapse Detector were not annotated by the expert (Supp. Fig. 3d). Upon closer inspection, however, we found that these are overwhelmingly synapses with small volumes (Supp. Fig. 3e), making them more likely to be missed by the human. Indeed, when the expert was asked to inspect these locations, 93% of those detections were retrospectively verified as genuine synapses by the expert (Supp. Fig. 3f). We also observed more than half of the labeled synapses in 3D volume would be falsely labeled in 3D viewer of Vaa3D (Suppl. Fig. 3g). After exclusion of mislabeled synapses and inclusion of smaller synapses, the overall synapse detection performance was higher than 90% (Suppl. Fig. 3h).

Definition of Cortical Layers and Preparation of Their Reconstructions

Boundaries between cortical layers were identified by using the parcellation scheme of Common Coordinate Framework of Allen Brain Atlas (59). Accordingly, layer I, II/III, V and VI have been expected to cover 8%, 14%, 61% and 17% of the cortical column, respectively. Layer V was then divided into 2 equal halves: layer Va and layer Vb. Then 3D cortical reconstructions were divided into laminae according to the identified vertical coordinates. Segments straddling two neighboring layers were excluded from the analysis.

Calculation of Percentages of Each Synapse-to-all Detected Synapses

After dividing cortical columns into cortical layers, we calculated the ratios of synapses made by each long-range projections originated from certain brain region (represented by color in the formula) to all detected synapses from all three brain regions. Let n_{layer}^{color} indicate number of synapses from BLA, VH or VTA in the layer I, II/III, Va, Vb or VI:

$$N_{layer} = n_{layer}^{cyan} + n_{layer}^{green} + n_{layer}^{yellow} \#(5)$$

$$P_{layer}^{color} = \frac{n_{layer}^{color}}{N_{layer}} \times 100; \; color \in \{cyan, green, yellow\} \#(6)$$

N_{layer} indicates numbers number of total detected synapses in a certain cortical layer. P_{layer}^{color} indicates the percentage of certain detected synapses color in given layer. We than created a color percentage matrix M^{color} for each color:

$$M_{i,layer}^{color} = P_{layer}^{color} \#(7)$$

$M_{i,layer}^{color}$ indicates the percentage value of cyan, green or yellow eGRASP labeled synapses in certain $layer$ of cortical column i where i is the identity of cortical column. The expected size of the color percentage matrix is 5 x 13 (parts of cortical columns x number of mice) for each brain region projecting to mPFC.

Dimensionality Reduction Matrix for Principal Component Analysis

To apply dimension reduction first, M^{color} matrices belonging different brain regions were concatenated in 2 dimensions.

$$M^P = [M^{cyan}, M^{green}, M^{yellow}] \#(8)$$

The final size of dimension reduction matrix (M^P) was 13 x 15. Then, principal component analysis was performed. The first and second principal components have been used for plotting low dimensional projection of percentages of "synapses made by single brain region-to-all detected synapses matrix.

Formation and Analysis of Closest Spatial Distance Matrix of BLA and VH Synapses

For BLA and VH synapses in layer Vb and layer VI, the first four closest spatial distances to BLA or VH synapses were measured. A 42,317 x 8 sized matrix was formed by appending closest VH and BLA synapse distances, respectively. Then Uniform Manifold Approximation and Projection (UMAP) was used to visualize synapses, behavioral profiles, social interaction scores, mouse IDs (31). Default values (version 0.5.5) were used for the hyperparameters of the UMAP code.

Random Forest Classifiers were formed using 10 forests with 75 trees. K-fold cross validation scheme with 10 independent parts was applied to each forest. In each fold, 10% of the dataset has been used as testing dataset. Only data labels have been randomized to train RF classifier to compare RF classification performance for structured and randomized datasets. Parametric measures of accuracy and F1 scores of RF classifier have been used for comparison between

main data and randomly shuffled data. RF classifiers have been trained using sklearn v1.4.0 inside Python 3.9.

Statistical Analysis

Data are presented as median with interquartile ranges, mean +/- S.E.M or mean with 95% confidence interval. Since the number of the samples in a group is low, non-parametric tests were used: Kruskal-Wallis (KW) test was used for multiple group comparisons and followed by pairwise comparisons by Mann Whitney U (MWU) tests, p values for MWU were adjusted by Bonferroni correction. All analysis were two tailed. p value less than 0.05 and 0.0167 were considered as significant in KW and post-hoc MWU tests, respectively. All statistical analysis was performed using Scipy v1.11.1 library of Python 3.9.

COMPETING INTERESTS

The authors declare that they have no competing interests.

AUTHOR CONTRIBUTIONS

STS, SEE and EEK were responsible for the design and conception of the study. STS, SEE and AC were responsible for the acquisition of data, STS, OG, SEE, AC, US and EEK were responsible for the analysis and interpretation of data. STS, US and EEK drafted the manuscript.

ACKNOWLEDGEMENTS

This study is supported by Research Fund of Hacettepe University, Grant number: TSA-2020-18753 to EEK. We thank Dr. Turgay Dalkara for his valuable opinions on the manuscript; Dr. Nevin Belder, Dr. Aslıhan Bahadır-Varol, Mesut Fırat and Anke Schönherr for their technical support. Graphical depictions inside Figure-1 and Graphical Abstract were prepared with Biorender.

MAIN FIGURE DESCRIPTIONS

Figure – 1: Description of the Experimental Strategy. a: AAV constructs used in the study. **b:** AAV injection sites, colored according to the e-GRASP constructs AAVs express. **c:** Schematic of a pyramidal mPFC neuron dendrite, tagged with TagRFP-T, that makes synapses with afferents from BLA, VH and VTA, each labelled with a different color. **d:** Timeline of the experimental procedure. **e:** Representative tracking images of SS (top) and SR (bottom) mice during the social interaction test. **f:** Social interaction scores. Data is shown as mean +/- SEM. **g:** Tiled imaging scheme across an mPFC column covering all layers. **h:** Emission and excitation wavelengths used for multispectral confocal imaging. Emission wavelengths taken from *fpbase.org(60).*

Figure – 2: Labeling and detection of long-range inputs from VH, BLA and VTA to mPFC pyramidal neurons across the cortical column. a: Image processing steps: Left: Schematic of tiled acquisition. Right: Processing of individual tiles. Processing of the dendritic images is shown at the top, processing of the synaptic signals is shown at the bottom. **b-h:** Representative images from different steps of the image processing pipeline. **b:** raw image stack; **c:** unmixed image; **d:** deconvolved image; **e:** segmented image; **f:** top: dendritic reconstruction of the sample, bottom: image of the segmented stack overlaid on dendritic reconstruction; **g:** top: thresholded image stack, bottom: alpha blended version of the thresholded image; **h:** synapses detected by Synapse Detector (Vaa3D), left: alpha blended version of detected synapses on dendritic segments, right: ball and stick version of dendritic reconstruction and detected synapses. Scale bar: 25 μm. **i:** representative image of synapses on a dendritic segment; left: deconvolved stack sample, right: zoomed-in area shown in the left image. Scaler bar 20 μm. **i.1**: reconstruction of dendritic segment overlaid on the deconvolved image, **i.2:** ball and stick version of dendritic segment and its synapses. Scale bar: 5 μm.

Figure – 3: Layer and stress phenotype-dependent changes of BLA, VH and VTA synapses. a: Representative unmixed image of a cortical column covering all layers of mPFC. Scale bar: 200 μm. **b:** Percentage of BLA-to-all detected synapses in mPFC layers I-VI. From top to bottom: Layer I (Kruskal-Wallis (KW) test: H = 4.02, p = 0.13); layer II/III (KW test: H = 5.53, p = 0.06); layer Va (KW test: H = 4.87, p = 0.08); layer Vb (KW test: H = 8.78, p = 0.012; Mann-Whitney U (MWU) test: p = 0.0158, SS vs SR ; p = 0.0158, SS vs SN ; p = 0.22, SR vs SN); layer VI (KW test: H = 10.50, p = 0.0052; MWU test: p = 0.0079, SS vs SR ; p = 0.0079, SS vs SN ; p = 0.15, SR vs SN). **c:** Percentage of VH-to-all detected synapses in mPFC layers

I-VI. From top to bottom: Layer I (KW test: H = 3.75, p = 0.15); layer II/III (KW test: H = 6.02, p = 0.049; MWU test: p = 0.54, SS vs SR ; p = 0.15, SS vs SN ; p = 0.0158, SR vs SN); layer Va (KW test: H = 7.57, p = 0.022; MWU test: p = 0.69, SS vs SR ; p = 0.0079, SS vs SN ; p = 0.055, SR vs SN, MWU test); layer Vb (KW test: H = 10.82, p = 0.0044; MWU test: p = 0.0079, SS vs SR ; p = 0.0079, SS vs SN ; p = 0.09, SR vs SN); layer VI (KW test: H = 9.50, p = 0.0086; MWU test: p = 0.0079, SS vs SR ; p = 0.0079, SS vs SN ; p = 0.69, SR vs SN). **d:** Percentage of VTA-to-all detected synapses in mPFC layers I-VI. From top to bottom: Layer I (KW test: H = 0.11, p = 0.94); layer II/III (KW test: H = 2.47, p = 0.28); layer Va (KW test: H = 1.26, p = 0.53); layer Vb (KW test: H = 4.5, p = 0.10); layer VI (KW test: H = 4.34, p = 0.11). **e:** Pooled data from each group showing percentages of BLA, VH and VTA synapses. **f:** Cortical depth vs percentage of BLA-to-all synapses, in all three groups. **g:** Correlation of BLA percentages in each layer with the social interaction (SI) ratio. **h:** Correlation of layer Vb's BLA percentages with the SI ratio (Spearman correlation: rho = - 0.55, p = 0.030; linear regression: F(1,13) = 4.96, R^2 = 0.28, p = 0.044). **i:** Correlation of layer VI's BLA percentages with the SI ratio (rho = - 0.74, p = 0.001, Spearman correlation; F(1,13) = 12.97, R^2 = 0.50, p = 0.0032, Linear regression). **j:** Cortical depth vs percentage of VH-to-all synapses, in all three groups. **k:** Correlation of VH percentages in each layer with the SI ratio. **l:** Correlation of layer Vb's VH percentages with the SI ratio (Spearman correlation: rho = 0.62, p = 0.012; Linear regression: F(1,13) = 8.33, R^2 = 0.39, p = 0.0127). **m:** Correlation of layer VI's VH percentages with the SI ratio (Spearman correlation: rho = 0.69, p = 0.0040; Linear regression: F(1,13) = 9.80, R^2 = 0.43, p = 0.0080). **n:** Cortical depth vs percentage of VTA-to-all synapses, in all three groups. **o:** Correlation of VTA percentages in each layer with the social interaction (SI) ratio. **p:** Correlation of layer Vb's VTA percentages with the SI ratio (Spearman correlation: rho = -0.34, p = 0.21; Linear regression: F(1,13) = 2.60, R^2 = 0.17, p = 0.13,). **r:** Correlation of layer VI's VTA percentages with the SI ratio (Spearman correlation: rho = - 0.16, p = 0.56; Linear regression: F(1,13) = 1.58, R^2 = 0.11, p = 0.23). **s:** Projection onto top 2 PCs of the synapse count data (synapse count per layer and afferent), for different behavioral groups. Data is presented as median and IQR for figure 4b,c,d; mean and 95% CI for figure 4f,j,n. Asterisks in the figures stand for ∗∗: p ≤ 0.01, and ∗: p < 0.05.

Figure – 4: The number of segments forming synapses with multiple brain regions increased in Layer VI of SS mice

a-e: Analysis of dendritic segments receiving inputs from a single brain region (monocolor) and from multiple brain regions (multicolor) in layer Vb. **a:** Pooled data from each group showing

total numbers and percentages of dendritic segments receiving inputs from a single brain region (monocolor) and from multiple brain regions (multicolor) in layer Vb. Number of monocolor segments in SN:SR:SS = 1,082:1,568:1,020. Number of multicolor segments in SN:SR:SS = 604:1,317:1,237. **b:** Percentage of dendritic segments receiving input from at least 2 different brain regions in layer Vb (Kruskal-Wallis (KW) test: H = 2.96, p = 0.22). **c:** Correlation of the ratio of multicolor dendritic segments in layer Vb with the SI ratio. (Spearman correlation: rho = - 0.18, p = 0.49; linear regression: $F_{(1,13)}$ = 0.02, R^2 = 0.0016, p = 0.88). **d:** Pooled data from each group showing total number and percentages of dendritic segments receiving input from multiple brain areas in layer Vb. **e:** Comparisons of percentages of dendritic segments receiving input from VTA-BLA (KW test: H = 12.58, p = 0.0018; Mann Whitney-U (MWU) test: p= 0.0079, SS vs SR; p = 0.011, SS vs SN; p = 0.011, SR vs SN), VTA-VH (KW test: H = 3.88, p = 0.14), VH-BLA (KW test: H = 1.93, p = 0.37) and VTA-VH-BLA (KW test: H = 3.31, p = 0.19). **f-j:** Same as a-e for layer VI. **f:** For layer VI, number of monocolor segments in SN:SR:SS = 352:1,589:465, number of multicolor segments in SN:SR:SS = 197:628:1,015. **g:** Percentage of dendritic segments receiving input from at least 2 different brain regions in layer VI (KW test: H = 8.66, p = 0.013; MWU test: p = 0.015, SS vs SR; p = 0.0079, SS vs SN; p = 0.84, SN vs SS). **h:** Correlation of multicolor dendritic segments in layer VI with SI ratio (Spearman correlation : rho = - 0.63, p = 0.010; linear regression: $F_{(1,13)}$ = 7.31, R^2 = 0.36, p = 0.018). **i:** Same as d for layer VI. **j:** Comparisons of percentages of dendritic segments receiving input: VTA-BLA (KW test: H = 2.61, p = 0.27); VTA-VH (KW test: H = 9.08, p = 0.010; MWU test: p = 0.11, SS vs SR ; p = 0.0079, SS vs SN ; p = 0.055, SR vs SN); VH-BLA (KW test: H = 3.46, p = 0.17); VTA-VH-BLA (KW test: H = 3.31, p = 0.19). Data shown as median and IQR. Asterisks in the figures stand for ∗∗: p ≤ 0.01, and ∗: p < 0.05.

Figure – 5: Euclidean Distances Between Unidentical, but not Identical Synapses were Distinct in SS mice

a-b: Cartoon illustration of spatial organization of BLA and VH synapses in layer Vb and VI of **a.** SN/SR and **b.** SS mice. **c:** Histograms of distances between each VH synapses and its closest VH synapse for different groups. **d:** Histograms of distances between each BLA synapse and its closest BLA synapse for different groups. **e:** Histograms of distances between each VH synapse and its closest BLA synapse for different groups. **f:** Histograms of distances between each BLA synapse and its closest VH synapse for different groups. **g:** Histogram of the difference between the distance to the closest VH synapse and the distance to the closest BLA synapse for each VH synapse, for different groups. **h:** Histogram of the difference between the distance to the

closest VH synapse and the distance to the closest BLA synapse for each BLA synapse, for different groups. **i:** Comparison of accuracy after Random Forest Classification between observed and randomized data (p = 5.16 x 10^{-5}, independent samples t-test). **j:** Confusion matrices of observed data (left) and randomized data (right). Values were normalized for the number of true labels. **k:** F1 scores for behavioral groups. For SN mice: p = 1.83 x 10^{-24}; for SR mice: p = 2.34 x 10^{-23}; for SS mice: p = 2.12 x 10^{-30}, independent samples t-test. **l-m:** 2D UMAP projection of Euclidean distances to four closest VH and four closest BLA synapses. **l:** colored by behavioral groups, **m:** colored by SI Ratio. Data is presented as mean and 95% CI. ***: p < 0.001; **: p < 0.01; *: p < 0.05.

Supplemental Table 1. Characteristics of the Adeno-associated-virus, their injection areas and volumes

Virus	Density (vg/µL)	Volume (µL)	Area
AAV2/1-CaMKIIa-yellow pre-eGRASP (p32)	$6,67 \times 10^9$	1	BLA
AAV2/1-CaMKIIa-green pre-eGRASP (p32)	$6,65 \times 10^9$	1	VH
AAV2/1-CaMKIIa- cyan pre-eGRASP (p30)	$6,8 \times 10^9$	1	VTA
AAV2/1-DIO-TagRFP-T -p2A-post-eGRASP	$6,79 \times 10^8$	0,25	mPFC
AAV2/1-CaMKIIa-iCre	4×10^8	0,25	mPFC

SUPPLEMENTAL FIGURE DESCRIPTIONS

Supplemental Figure – 1: Sample Image Processing for Each Signal

a: Sample image processing for dendritic signals. **b:** Sample image processing for synaptic signals from top to bottom: VTA (cyan eGRASP), VH (green eGRASP) and BLA (yellow eGRASP). Scale bar: 25 µm

Supplemental Figure – 2: Accuracy of synapse detection for different synapse types and behavior phenotypes

a: Percentages of the automatically detected synapses that are confirmed by an expert. **b:** Percentage of retrospectively confirmed unmixed synapses among automatically detected synapses distributed for different brain regions and behavioral phenotypes. **c:** Initial percentage of matched synapses among the manually detected synapses. **d:** Initial percentage of matched synapses among the automatically detected synapses. **e:** Comparison of mean synaptic volumes of spatially matched and unmatched synapses which were detected automatically. (p = 2.57 x 10-8, Mann-Whitney U test). Spatially unmatched synapses had lower mean synapse volume than spatially matched synapses. **f:** Retrospectively determined percentage of true synapses among automatically detected synapses with smaller volumes **g:** Percentage of manually mislabeled synapses among unmatched but manually labeled synapses. **h:** Overall performance of Synapse Detector. Data is presented as mean and 95% CI.

Supplemental Figure – 3: Chronic stress increased and decreased the percentages of BLA--to-all detected synapses and VH-to-all detected synapses, respectively in superficial layers of mPFC.

a: Comparison of percentages of BLA-to-all detected synapses in mPFC layers I and II/III between stressed and stress naïve groups. Top: Layer I, p = 0.06; bottom: Layer II/III, p = 0.019, Mann-Whitney U test. **b:** Comparison of percentages of VH-to-all detected synapses in mPFC layers I and II/III between stressed and stress naïve groups. Top: Layer I, p = 0.17; bottom: Layer II/III, p = 0.019, Mann-Whitney U test. **c:** Comparison of percentages of VTA-to-all detected synapses in mPFC layers I and II/III between stressed and stress naïve groups. Top: Layer I, p = 0.83; bottom: Layer II/III, p = 0.85, Mann-Whitney U test. Data shown as median and IQR.

Supplemental Figure – 4: The number of dendritic segments forming synapses with afferents from a single brain region in Layer Vb and VI.

a: Pooled data from each group showing total number and percentages of dendritic segments forming at least two synapses with afferents from only one brain region (yellow colored for BLA-only, green colored for VH-only and cyan colored for VTA-only) and from multiple brain regions (purple colored) in layer Vb. **b:** Percentages of BLA-only-to-all segments with at least 2 synapses in layer Vb. Kruskal – Wallis (KW) Test: H= 7.79, p = 0.020; Mann-Whitney U (MWU) test: SS vs SR , p = 0.055; SS vs SN , p = 0.021; SR vs SN , p = 0.20. **c:** Percentages of VH-only-to-all segments with at least 2 synapses in layer Vb. KW test: H = 9.62, p = 0.0081; MWU test: SS vs SR , p = 0.0079; SS vs SN , p = 0.0079; SR vs SN , p = 0.54. **d:** Percentages of VTA-only-to-all segments with at least 2 synapses in layer Vb. KW test: H = 0.96, p = 0.61. **e:** Correlation of the percentage of BLA-only-to-all segments with at least 2 synapses with the SI ratio in layer Vb. (Spearman correlation: rho = - 0.55, p = 0.029; Linear regression: $F(1,13)$ = 3.23, R^2 = 0.20, p = 0.09). **f:** Correlation of the percentage of VH-only-to-all segments with at least 2 synapses with the SI ratio in layer Vb. (Spearman correlation: rho = 0.57, p = 0.025; Linear regression: $F(1,13)$ = 2.75, R^2 = 0.17, p = 0.12). **g:** Correlation of the percentage of VTA-only-to-all segments with at least 2 synapses with the SI ratio in layer Vb. (Spearman correlation: rho = - 0.15, p = 0.57; Linear regression: $F(1,13)$ = 1.89, R^2 = 0.13, p = 0.19). **h:** Pooled data from each group showing total number and percentages of dendritic segments forming at least two synapses with afferents from only one brain region (yellow colored for BLA-only, green colored for VH-only and cyan colored for VTA-only) and from multiple brain regions (purple colored) in layer VI. **i:** Percentages of BLA-only-to-all segments with at least 2 synapses in layer VI. (KW test: H = 5.82, p = 0.054). **j:** Percentages of VH-only-to-all segments with at least 2 synapses in layer VI. (KW test: H = 9.42, p = 0.0090; MWU test: SS vs SR , p = 0.0079; SS vs SN , p = 0.0079; SR vs SN , p = 0.84). **k:** Percentages of VTA-only-to-all segments with at least 2 synapses in layer VI. (KW test: H = 0.28, p = 0.86). **l:** Correlation of the percentage of BLA-only-to-all segments with at least 2 synapses with the SI ratio in layer VI. (Spearman correlation: rho = - 0.61, p = 0.013; Linear regression: $F(1,13)$ = 3.92, R^2 = 0.23, p = 0.06). **m:** Correlation of the percentage of VH-only-to-all segments with at least 2 synapses with the SI ratio in layer VI. (Spearman correlation: rho = 0.63, p = 0.010; Linear regression: $F(1,13)$ = 9.32, R^2 = 0.42, p = 0.0092). **n:** Correlation of the percentage of VTA-only-to-all segments with at least 2 synapses with the SI ratio in layer VI. (Spearman correlation: rho = - 0.02, p = 0.94; Linear regression: $F(1,13)$ = 0.16, R^2 = 0.01, p = 0.69)

SUPPLEMENTAL VIDEO DESCRIPTIONS

Supplemental Video – 1: Synapses aligned with dendritic reconstruction.

Supplemental Video – 2: Dendritic reconstruction and BLA, VH and VTA synapses throughout mPFC laminae.

REFERENCES

1.   Cook SC, Wellman CL. Chronic stress alters dendritic morphology in rat medial prefrontal cortex. J Neurobiol. 2004;60(2):236-48. doi: 10.1002/neu.20025. PubMed PMID: 15266654.

2.   Liston C, Miller MM, Goldwater DS, Radley JJ, Rocher AB, Hof PR, et al. Stress-induced alterations in prefrontal cortical dendritic morphology predict selective impairments in perceptual attentional set-shifting. J Neurosci. 2006;26(30):7870-4. doi: 10.1523/JNEUROSCI.1184-06.2006. PubMed PMID: 16870732; PubMed Central PMCID: PMC6674229.

3.   Radley JJ, Arias CM, Sawchenko PE. Regional differentiation of the medial prefrontal cortex in regulating adaptive responses to acute emotional stress. J Neurosci. 2006;26(50):12967-76. doi: 10.1523/JNEUROSCI.4297-06.2006. PubMed PMID: 17167086; PubMed Central PMCID: PMC6674963.

4.   Goldwater DS, Pavlides C, Hunter RG, Bloss EB, Hof PR, McEwen BS, Morrison JH. Structural and functional alterations to rat medial prefrontal cortex following chronic restraint stress and recovery. Neuroscience. 2009;164(2):798-808. Epub 20090829. doi: 10.1016/j.neuroscience.2009.08.053. PubMed PMID: 19723561; PubMed Central PMCID: PMC2762025.

5.   Hercher C, Canetti L, Turecki G, Mechawar N. Anterior cingulate pyramidal neurons display altered dendritic branching in depressed suicides. J Psychiatr Res. 2010;44(5):286-93. Epub 20090917. doi: 10.1016/j.jpsychires.2009.08.011. PubMed PMID: 19765721.

6.   Moda-Sava RN, Murdock MH, Parekh PK, Fetcho RN, Huang BS, Huynh TN, et al. Sustained rescue of prefrontal circuit dysfunction by antidepressant-induced spine formation. Science. 2019;364(6436). doi: 10.1126/science.aat8078. PubMed PMID: 30975859; PubMed Central PMCID: PMC6785189.

7.   Shansky RM, Hamo C, Hof PR, McEwen BS, Morrison JH. Stress-induced dendritic remodeling in the prefrontal cortex is circuit specific. Cereb Cortex. 2009;19(10):2479-84. Epub 20090204. doi: 10.1093/cercor/bhp003. PubMed PMID: 19193712; PubMed Central PMCID: PMC2742599.

8.   Anastasiades PG, Carter AG. Circuit organization of the rodent medial prefrontal cortex. Trends Neurosci. 2021;44(7):550-63. Epub 20210507. doi: 10.1016/j.tins.2021.03.006. PubMed PMID: 33972100; PubMed Central PMCID: PMC8222144.

9.   Fillinger C, Yalcin I, Barrot M, Veinante P. Afferents to anterior cingulate areas 24a and 24b and midcingulate areas 24a' and 24b' in the mouse. Brain Struct Funct. 2017;222(3):1509-32. Epub 20160818. doi: 10.1007/s00429-016-1290-1. PubMed PMID: 27539453.

10.   Covington HE, 3rd, Lobo MK, Maze I, Vialou V, Hyman JM, Zaman S, et al. Antidepressant effect of optogenetic stimulation of the medial prefrontal cortex. J Neurosci. 2010;30(48):16082-90. doi: 10.1523/JNEUROSCI.1731-10.2010. PubMed PMID: 21123555; PubMed Central PMCID: PMC3004756.

11.   Chaudhury D, Walsh JJ, Friedman AK, Juarez B, Ku SM, Koo JW, et al. Rapid regulation of depression-related behaviours by control of midbrain dopamine neurons. Nature. 2013;493(7433):532-6. Epub 20121212. doi: 10.1038/nature11713. PubMed PMID: 23235832; PubMed Central PMCID: PMC3554860.

12.   Friedman AK, Walsh JJ, Juarez B, Ku SM, Chaudhury D, Wang J, et al. Enhancing depression mechanisms in midbrain dopamine neurons achieves homeostatic resilience. Science. 2014;344(6181):313-9. doi: 10.1126/science.1249240. PubMed PMID: 24744379; PubMed Central PMCID: PMC4334447.

13.   Carreno FR, Donegan JJ, Boley AM, Shah A, DeGuzman M, Frazer A, Lodge DJ. Activation of a ventral hippocampus-medial prefrontal cortex pathway is both necessary and sufficient for an antidepressant response to ketamine. Mol Psychiatry. 2016;21(9):1298-308. Epub 20151201. doi: 10.1038/mp.2015.176. PubMed PMID: 26619811.

14.   Felix-Ortiz AC, Burgos-Robles A, Bhagat ND, Leppla CA, Tye KM. Bidirectional modulation of anxiety-related and social behaviors by amygdala projections to the medial prefrontal cortex. Neuroscience. 2016;321:197-209. Epub 20150721. doi: 10.1016/j.neuroscience.2015.07.041. PubMed PMID: 26204817; PubMed Central PMCID: PMC4721937.

15.   Burgos-Robles A, Kimchi EY, Izadmehr EM, Porzenheim MJ, Ramos-Guasp WA, Nieh EH, et al. Amygdala inputs to prefrontal cortex guide behavior amid conflicting cues of reward and punishment. Nat Neurosci. 2017;20(6):824-35. Epub 20170424. doi: 10.1038/nn.4553. PubMed PMID: 28436980; PubMed Central PMCID: PMC5448704.

16.   Klavir O, Prigge M, Sarel A, Paz R, Yizhar O. Manipulating fear associations via optogenetic modulation of amygdala inputs to prefrontal cortex. Nat Neurosci. 2017;20(6):836-44. Epub 20170313. doi: 10.1038/nn.4523. PubMed PMID: 28288126.

17.   Sotres-Bayon F, Sierra-Mercado D, Pardilla-Delgado E, Quirk GJ. Gating of fear in prelimbic cortex by hippocampal and amygdala inputs. Neuron. 2012;76(4):804-12. doi: 10.1016/j.neuron.2012.09.028. PubMed PMID: 23177964; PubMed Central PMCID: PMC3508462.

18.   Adhikari A, Topiwala MA, Gordon JA. Synchronized activity between the ventral hippocampus and the medial prefrontal cortex during anxiety. Neuron. 2010;65(2):257-69. doi: 10.1016/j.neuron.2009.12.002. PubMed PMID: 20152131; PubMed Central PMCID: PMC2822726.

19.   Adhikari A, Topiwala MA, Gordon JA. Single units in the medial prefrontal cortex with anxiety-related firing patterns are preferentially influenced by ventral hippocampal activity. Neuron. 2011;71(5):898-910. doi: 10.1016/j.neuron.2011.07.027. PubMed PMID: 21903082; PubMed Central PMCID: PMC3201792.

20.   Woo E, Sansing LH, Arnsten AFT, Datta D. Chronic Stress Weakens Connectivity in the Prefrontal Cortex: Architectural and Molecular Changes. Chronic Stress (Thousand Oaks). 2021;5:24705470211029254. Epub 20210829. doi: 10.1177/24705470211029254. PubMed PMID: 34485797; PubMed Central PMCID: PMC8408896.

21.   Little JP, Carter AG. Subcellular synaptic connectivity of layer 2 pyramidal neurons in the medial prefrontal cortex. J Neurosci. 2012;32(37):12808-19. doi: 10.1523/JNEUROSCI.1616-12.2012. PubMed PMID: 22973004; PubMed Central PMCID: PMC3490687.

22.   Little JP, Carter AG. Synaptic mechanisms underlying strong reciprocal connectivity between the medial prefrontal cortex and basolateral amygdala. J Neurosci. 2013;33(39):15333-42. doi: 10.1523/JNEUROSCI.2385-13.2013. PubMed PMID: 24068800; PubMed Central PMCID: PMC3782616.

23.   Liu X, Carter AG. Ventral Hippocampal Inputs Preferentially Drive Corticocortical Neurons in the Infralimbic Prefrontal Cortex. J Neurosci. 2018;38(33):7351-63. Epub 20180629. doi: 10.1523/JNEUROSCI.0378-18.2018. PubMed PMID: 29959235; PubMed Central PMCID: PMC6096040.

24.   Herry C, Ciocchi S, Senn V, Demmou L, Muller C, Luthi A. Switching on and off fear by distinct neuronal circuits. Nature. 2008;454(7204):600-6. Epub 20080709. doi: 10.1038/nature07166. PubMed PMID: 18615015.

25.   Sanchez-Bellot C, AlSubaie R, Mishchanchuk K, Wee RWS, MacAskill AF. Two opposing hippocampus to prefrontal cortex pathways for the control of approach and avoidance behaviour. Nat Commun. 2022;13(1):339. Epub 20220117. doi: 10.1038/s41467-022-27977-7. PubMed PMID: 35039510; PubMed Central PMCID: PMC8763938.

26.   Choi JH, Sim SE, Kim JI, Choi DI, Oh J, Ye S, et al. Interregional synaptic maps among engram cells underlie memory formation. Science. 2018;360(6387):430-5. doi: 10.1126/science.aas9204. PubMed PMID: 29700265.

27.   Golden SA, Covington HE, 3rd, Berton O, Russo SJ. A standardized protocol for repeated social defeat stress in mice. Nat Protoc. 2011;6(8):1183-91. Epub 20110721. doi: 10.1038/nprot.2011.361. PubMed PMID: 21799487; PubMed Central PMCID: PMC3220278.

28.   McRae TD, Oleksyn D, Miller J, Gao YR. Robust blind spectral unmixing for fluorescence microscopy using unsupervised learning. PLoS One. 2019;14(12):e0225410. Epub 20191202. doi: 10.1371/journal.pone.0225410. PubMed PMID: 31790435; PubMed Central PMCID: PMC6886781.

29.   Gliko O, Mallory M, Dalley R, Gala R, Gornet J, Zeng H, et al. High-throughput analysis of dendritic and axonal arbors reveals transcriptomic correlates of neuroanatomy. bioRxiv. 2024:2022.03.07.482900. doi: 10.1101/2022.03.07.482900.

30.   Iascone DM, Li Y, Sumbul U, Doron M, Chen H, Andreu V, et al. Whole-Neuron Synaptic Mapping Reveals Spatially Precise Excitatory/Inhibitory Balance Limiting Dendritic and Somatic Spiking. Neuron. 2020;106(4):566-78 e8. Epub 20200312. doi: 10.1016/j.neuron.2020.02.015. PubMed PMID: 32169170; PubMed Central PMCID: PMC7244395.

31.   McInnes L, Healy J, Melville J. Umap: Uniform manifold approximation and projection for dimension reduction. arXiv preprint arXiv:180203426. 2018.

32.   Sesack SR, Deutch AY, Roth RH, Bunney BS. Topographical organization of the efferent projections of the medial prefrontal cortex in the rat: an anterograde tract-tracing study with Phaseolus vulgaris leucoagglutinin. J Comp Neurol. 1989;290(2):213-42. doi: 10.1002/cne.902900205. PubMed PMID: 2592611.

33.   McGarry LM, Carter AG. Prefrontal Cortex Drives Distinct Projection Neurons in the Basolateral Amygdala. Cell Rep. 2017;21(6):1426-33. doi: 10.1016/j.celrep.2017.10.046. PubMed PMID: 29117549; PubMed Central PMCID: PMC5714295.

34.   Tanimizu T, Kenney JW, Okano E, Kadoma K, Frankland PW, Kida S. Functional Connectivity of Multiple Brain Regions Required for the Consolidation of Social Recognition Memory. J Neurosci. 2017;37(15):4103-16. Epub 20170314. doi: 10.1523/JNEUROSCI.3451-16.2017. PubMed PMID: 28292834; PubMed Central PMCID: PMC6596582.

35.   Hitti FL, Siegelbaum SA. The hippocampal CA2 region is essential for social memory. Nature. 2014;508(7494):88-92. Epub 20140223. doi: 10.1038/nature13028. PubMed PMID: 24572357; PubMed Central PMCID: PMC4000264.

36.   Okuyama T, Kitamura T, Roy DS, Itohara S, Tonegawa S. Ventral CA1 neurons store social memory. Science. 2016;353(6307):1536-41. doi: 10.1126/science.aaf7003. PubMed PMID: 27708103; PubMed Central PMCID: PMC5493325.

37.   Lee E, Rhim I, Lee JW, Ghim JW, Lee S, Kim E, Jung MW. Enhanced Neuronal Activity in the Medial Prefrontal Cortex during Social Approach Behavior. J Neurosci. 2016;36(26):6926-36. doi: 10.1523/JNEUROSCI.0307-16.2016. PubMed PMID: 27358451; PubMed Central PMCID: PMC6604896.

38.   Likhtik E, Stujenske JM, Topiwala MA, Harris AZ, Gordon JA. Prefrontal entrainment of amygdala activity signals safety in learned fear and innate anxiety. Nat Neurosci. 2014;17(1):106-13. Epub 20131117. doi: 10.1038/nn.3582. PubMed PMID: 24241397; PubMed Central PMCID: PMC4035371.

39.   Silverstein SE, O'Sullivan R, Bukalo O, Pati D, Schaffer JA, Limoges A, et al. A distinct cortical code for socially learned threat. Nature. 2024;626(8001):1066-72. Epub 20240207. doi: 10.1038/s41586-023-07008-1. PubMed PMID: 38326610.

40.   Padilla-Coreano N, Canetta S, Mikofsky RM, Alway E, Passecker J, Myroshnychenko MV, et al. Hippocampal-Prefrontal Theta Transmission Regulates Avoidance Behavior. Neuron. 2019;104(3):601-10 e4. Epub 20190911. doi: 10.1016/j.neuron.2019.08.006. PubMed PMID: 31521441; PubMed Central PMCID: PMC6842114.

41.   Kaidanovich-Beilin O, Lipina T, Vukobradovic I, Roder J, Woodgett JR. Assessment of social interaction behaviors. J Vis Exp. 2011(48). Epub 20110225. doi: 10.3791/2473. PubMed PMID: 21403628; PubMed Central PMCID: PMC3197404.

42.   Lissek S, Rabin S, Heller RE, Lukenbaugh D, Geraci M, Pine DS, Grillon C. Overgeneralization of conditioned fear as a pathogenic marker of panic disorder. Am J Psychiatry. 2010;167(1):47-55. Epub 20091116. doi: 10.1176/appi.ajp.2009.09030410. PubMed PMID: 19917595; PubMed Central PMCID: PMC2806514.

43.   Jovanovic T, Kazama A, Bachevalier J, Davis M. Impaired safety signal learning may be a biomarker of PTSD. Neuropharmacology. 2012;62(2):695-704. Epub 20110304. doi: 10.1016/j.neuropharm.2011.02.023. PubMed PMID: 21377482; PubMed Central PMCID: PMC3146576.

44.   Lissek S, Kaczkurkin AN, Rabin S, Geraci M, Pine DS, Grillon C. Generalized anxiety disorder is associated with overgeneralization of classically conditioned fear. Biol Psychiatry. 2014;75(11):909-15. Epub 20130831. doi: 10.1016/j.biopsych.2013.07.025. PubMed PMID: 24001473; PubMed Central PMCID: PMC3938992.

45.   Kaczkurkin AN, Burton PC, Chazin SM, Manbeck AB, Espensen-Sturges T, Cooper SE, et al. Neural Substrates of Overgeneralized Conditioned Fear in PTSD. Am J Psychiatry. 2017;174(2):125-34. Epub 20161031. doi: 10.1176/appi.ajp.2016.15121549. PubMed PMID: 27794690; PubMed Central PMCID: PMC7269602.

46.   Fetcho RN, Hall BS, Estrin DJ, Walsh AP, Schuette PJ, Kaminsky J, et al. Regulation of social interaction in mice by a frontostriatal circuit modulated by established hierarchical relationships. Nat Commun. 2023;14(1):2487. Epub 20230429. doi: 10.1038/s41467-023-37460-6. PubMed PMID: 37120443; PubMed Central PMCID: PMC10148889.

47.   Yizhar O, Fenno LE, Prigge M, Schneider F, Davidson TJ, O'Shea DJ, et al. Neocortical excitation/inhibition balance in information processing and social dysfunction. Nature. 2011;477(7363):171-8. Epub 20110727. doi: 10.1038/nature10360. PubMed PMID: 21796121; PubMed Central PMCID: PMC4155501.

48.   Sun Q, Li X, Ren M, Zhao M, Zhong Q, Ren Y, et al. A whole-brain map of long-range inputs to GABAergic interneurons in the mouse medial prefrontal cortex. Nat Neurosci. 2019;22(8):1357-70. Epub 20190708. doi: 10.1038/s41593-019-0429-9. PubMed PMID: 31285615.

49.   Lee AT, Cunniff MM, See JZ, Wilke SA, Luongo FJ, Ellwood IT, et al. VIP Interneurons Contribute to Avoidance Behavior by Regulating Information Flow across Hippocampal-Prefrontal Networks. Neuron. 2019;102(6):1223-34 e4. Epub 20190430. doi: 10.1016/j.neuron.2019.04.001. PubMed PMID: 31053407; PubMed Central PMCID: PMC6800223.

50.   Soumier A, Sibille E. Opposing effects of acute versus chronic blockade of frontal cortex somatostatin-positive inhibitory neurons on behavioral emotionality in mice. Neuropsychopharmacology. 2014;39(9):2252-62. Epub 20140401. doi: 10.1038/npp.2014.76. PubMed PMID: 24690741; PubMed Central PMCID: PMC4104344.

51.   Perova Z, Delevich K, Li B. Depression of excitatory synapses onto parvalbumin interneurons in the medial prefrontal cortex in susceptibility to stress. J Neurosci. 2015;35(7):3201-6. doi: 10.1523/JNEUROSCI.2670-14.2015. PubMed PMID: 25698754; PubMed Central PMCID: PMC4331634.

52.   Harris AZ, Atsak P, Bretton ZH, Holt ES, Alam R, Morton MP, et al. A Novel Method for Chronic Social Defeat Stress in Female Mice. Neuropsychopharmacology. 2018;43(6):1276-83. Epub 20171101. doi: 10.1038/npp.2017.259. PubMed PMID: 29090682; PubMed Central PMCID: PMC5916350.

53.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82. Epub 20120628. doi: 10.1038/nmeth.2019. PubMed PMID: 22743772; PubMed Central PMCID: PMC3855844.

54.   Peng H, Bria A, Zhou Z, Iannello G, Long F. Extensible visualization and analysis for multidimensional images using Vaa3D. Nat Protoc. 2014;9(1):193-208. Epub 20140102. doi: 10.1038/nprot.2014.011. PubMed PMID: 24385149.

55. Horl D, Rojas Rusak F, Preusser F, Tillberg P, Randel N, Chhetri RK, et al. BigStitcher: reconstructing high-resolution image datasets of cleared and expanded samples. Nat Methods. 2019;16(9):870-4. Epub 20190805. doi: 10.1038/s41592-019-0501-0. PubMed PMID: 31384047.

56. Perdigão LMA, Yee NBy, Ho EML, Pennington AH, King ONF, Darrow MC, Basham M. Computing for Optimized Biological Microscopy Data Processing and Analysis at The Rosalind Franklin Institute. Microscopy and Microanalysis. 2022;28(S1):1462-4. Epub 2022/07/22. doi: 10.1017/S1431927622005931.

57. Choi DI, Kaang BK. Interrogating structural plasticity among synaptic engrams. Curr Opin Neurobiol. 2022;75:102552. Epub 20220519. doi: 10.1016/j.conb.2022.102552. PubMed PMID: 35598549.

58. Shao LX, Liao C, Gregg I, Davoudian PA, Savalia NK, Delagarza K, Kwan AC. Psilocybin induces rapid and persistent growth of dendritic spines in frontal cortex in vivo. Neuron. 2021;109(16):2535-44 e4. Epub 20210705. doi: 10.1016/j.neuron.2021.06.008. PubMed PMID: 34228959; PubMed Central PMCID: PMC8376772.

59. Wang Q, Ding SL, Li Y, Royall J, Feng D, Lesnar P, et al. The Allen Mouse Brain Common Coordinate Framework: A 3D Reference Atlas. Cell. 2020;181(4):936-53 e20. Epub 20200507. doi: 10.1016/j.cell.2020.04.007. PubMed PMID: 32386544; PubMed Central PMCID: PMC8152789.

60. Lambert TJ. FPbase: a community-editable fluorescent protein database. Nat Methods. 2019;16(4):277-8. doi: 10.1038/s41592-019-0352-8. PubMed PMID: 30886412.

a

VH — CaMKIIα — Green pre-eGRASP

BLA — CaMKIIα — Yellow pre-eGRASP

VTA — CaMKIIα — Cyan pre-eGRASP

mPFC (ACC) — CaMKIIα — iCre

EF1α — post-eGRASP — P2A — myr_TagRFP-T

b

mPFC BLA VH VTA

c

VTA VH BLA

d

Virus Injections — Physical Stress 3-10 mins — Sensory Stress 24 h — Social Interaction Test

Day 1 — Day 21 — Day 30 — Day 31

CSDS x 10 days

e

Empty   with CD-1

Stress Susceptible

Stress Resilient

f

Naive

Resilient

Susceptible

Social Interaction Ratio

g

~200 μm
~20 μm
1
2

h

453 nm   488 nm   543 nm   633 nm

1   2   3   4   5   6

450   500   550   600   650   700   750

a

b

Raw

c

Unmixed

d

Deconvolved

e

Segmented

f

Reconstructed

g

Thresholded

h

VTA

VH

BLA

mPFC

Detected with Synapse Detector

i