# Heterologous HSPC transplantation rescues neuroinflammation and ameliorates peripheral manifestations in the mouse model of lysosomal transmembrane enzyme deficiency, MPS IIIC.

Xuefang Pan[1], Antoine Caillon[1], Shuxian Fan[1,2], Shaukat Khan[3], Shunji Tomatsu[3], and Alexey V. Pshezhetsky[1,2]

[1]Department of Pediatrics and Centre Hospitalier Universitaire Sainte-Justine Research Centre, University of Montreal, Montreal, QC, Canada;

[2]Department of Anatomy and Cell Biology, McGill University, Montreal, QC, Canada;

[3]Nemours/Alfred I. duPont Hospital for Children, Wilmington, DE, USA.

Corresponding author:
Alexey V. Pshezhetsky
CHU Sainte-Justine
3175 Côte Ste-Catherine, Montréal (QC) H3T 1C5, Canada
Tel. (514) 345-4931/2736,
E-Mail: alexei.pchejetski@umontreal.ca

**Running title:**

HSPC transplantation rescues neuroinflammation in MPS IIIC mice

**Key words:**

Mucopolysaccharidosis, Sanfilippo disease, neuroinflammation, heparan sulfate, HSPC, allogenic

HSPC transplantation, microglia.

**Abstract**

Mucopolysaccharidosis III type C (MPS IIIC) is an untreatable neuropathic lysosomal storage disease caused by a genetic deficiency of the lysosomal N-acetyltransferase, HGSNAT, catalyzing a transmembrane acetylation of heparan sulfate. HGSNAT is a transmembrane enzyme uncapable of free diffusion between the cells and their cross-correction which limits development of therapies based on the enzyme replacement and gene correction. Since our previous work identified neuroinflammation as a hallmark of the CNS pathology in MPS IIIC, we tested whether it can be corrected by replacement of activated brain microglia with neuroprotective macrophages/microglia derived from a heterologous HSPC transplant.

Eight-week-old MPS IIIC ($Hgsnat^{P304L}$) mice were transplanted with HSPC from congenic WT mice after myeloablation with Busulfan and studied by behaviour test battery staring from the age of 6 months. At the age of ~8 months, mice were sacrificed to study pathological changes in the brain, heparan sulfate storage and other biomarkers of the disease. We found that the treatment corrected several behaviour deficits including memory decline, and reduction of socialization, as well as several features of CNS pathology such as microastroglyosis, expression of pro-inflammatory cytokine IL-1β, and accumulation of misfolded amyloid aggregates in cortical neurons. At the periphery, the treatment delayed development of terminal urinary retention, potentially increasing longevity, and reduced blood levels of heparan sulfate. However, no correction of lysosomal storage phenotype in neurons and brain levels of heparan sulfate was observed.

Together, our results demonstrate that neuroinflammation in a neurological lysosomal storage disease, caused by defects in a transmembrane enzyme, can be effectively ameliorated by replacement of microglia bearing the genetic defect with the cells from normal healthy donor.

They also suggest, that heterologous HSPC transplant if used together with other methods, such as chaperone therapy or substrate reduction therapy, may constitute an effective combination therapy for MPS IIIC and other disorders with a similar etiology.

## Background

Lysosomal storage disorders (LSDs) are conditions caused by the accumulation (storage) of biological macromolecules due to genetic defects in lysosomal catabolism. More than 70 LSDs are currently described that, together, have frequency of 1:4000 live births [1]. About 30 % of children with LSDs are affected by mucopolysaccharidoses (MPS) caused by impaired lysosomal degradation of glycosaminoglycans (GAGs). Most MPS affect the central nervous system (CNS) early in childhood and result in severe progressive neurodegenerative changes, eventually leading to handicap and premature death. Non-neurological forms of MPS and other LSDs are partially treatable by enzyme replacement therapy (ERT), but this approach is ineffective for the neurological forms due to inability of recombinant enzymes to cross the blood-brain barrier (BBB). In MPS III type C, caused by genetic deficiency of lysosomal membrane enzyme, heparan sulfate acetyl-CoA: α-glucosaminide N-acetyltransferase (HGSNAT), which catalyzes transmembrane acetylation of glucosamine residues of heparan sulfate (HS), an additional constraint is the absence of cross-correction between the cells. Delivering replacement enzyme intrathecally or intraventricularly, thus bypassing the BBB [2], is not possible for membrane enzymes and proteins such as HGSNAT and difficult to implement clinically due to the invasive nature of the procedure.

Hematopoietic stem and progenitor cell (HSPC) transplant is the clinically recommended therapy for severe neuropathic forms of MPS I, where the missing enzyme is soluble and can be

effectively secreted by donor cells [3]. The treatment restores catabolism in HSPC which, in combination with cross-correction of other cell types through secretion of the enzyme, reduces storage in the tissues of peripheral organs. Importantly, the donor-derived monocytes also cross the BBB and differentiate into microglial cells, thus ameliorating the neuropathology [4, 5].

Emerging data demonstrate that lentivirus (LV)-mediated gene correction of HSPC, followed by their transplantation, can treat neurological LSDs by soluble enzyme secreted from HSPC-derived brain microglia and macrophages [5-10]. In LSDs, when the CNS damage occurs, cytokines and chemokines are generated by activated macrophages/microglia and astrocytes, which can stimulate the upregulation of adhesion molecules on brain endothelial cells, resulting in enhanced trans-endothelial migration of monocytes from the blood into perivascular regions. These enzyme-deficient cells migrate into the parenchyma to sites of damage, where they can secrete potentially cytotoxic pro-inflammatory cytokines, leading to further damage and cell death. However, if the hematopoietic precursors are gene-corrected, the macrophages and microglia gain the ability to transfer the missing lysosomal enzyme to the CNS and catabolize stored metabolites. This approach, now approved for patients with metachromatic leukodystrophy, was shown to improve or completely treat the disease in several animal MPS models including the mouse models of MPS IIIA and IIIB. Clinical trials for presymptomatic MPS IIIA patients are currently in progress (Manchester Hospital Clinical Trial 2023, Dr. Simon Johns, PI), but it is not clear whether this methodology would show efficacy in MPS IIIC where the defective enzyme is a transmembrane protein.

However, it is possible to hypothesize that the transplantation of gene-corrected or wild-type microglia, could potentially also mitigate neuroinflammatory response, which is described for all neurological MPS spectrum diseases, including MPS IIIC. In our MPS IIIC mouse model *Hgsnat-*

*Geo*, a functional knockout (KO) of the *Hgsnat* locus in mice, storage of HS in microglia leading to their activation was the initial pathological event in the brain, present already at 2 months and gradually aggravating with age [11]. Microglia activation resulted in release of pro-inflammatory cytokines, including IL-1β, that cause mitochondrial damage [12-14], and, together with accumulation of misfolded toxic protein aggregates, eventually lead to neuronal death dominant in all brain regions. More recently, we confirmed that similar pathophysiological mechanism also manifests in a knock-in MPS IIIC mouse strain *Hgsnat^{P304L}*, that expresses mutant misfolded HGSNAT protein [15]. This study defined CD68-positive microglia, GFAP-positive activated astrocytes, as well as subunit C mitochondrial ATPase (SCMAS)-positive, amyloid-β-positive, and Thioflavin-S-positive aggregates in cortical neurons as the most robust biomarkers of the disease progression [11, 16]. We also confirmed that similar biomarkers are present in *post-mortem* cortical tissues of human Sanfilippo patients, which suggested that the described pathophysiological mechanism is common for mice and humans [17].

In the current study, we show that neuroinflammation in the brain of *Hgsnat^{P304L}* mice is effectively corrected by allogenic transplantation of HSPC from WT mice, despite the absence of a rescue of HGSNAT activity or HS storage in the brain. Transplantation conducted in 2-month-old *Hgsnat^{P304L}* mice also prevented the development or urinary retention thus extending the life span of animals beyond the age of 8 months. Together, our results demonstrate that neuroinflammation is the crucial underlying pathological mechanism in MPS IIIC that can be effectively treated by replacement of microglia bearing the genetic defect with the cells from a normal healthy donor. Still, behavioral deficits and pathological changes in the neurons showed only partial improvement, suggesting that HSPC transplantation needs to be coupled with other interventions for development of effective therapy

## Materials and Methods

### Animals

The knock-in mouse model of MPS IIIC, *Hgsnat*[P304L], expressing the HGSNAT enzyme with human missense mutation P304L, on a C57BL/6J genetic background, has been previously described [15]. C57BL/6 congenic mouse strain B6.SJL-Ptprca Pepcb/BoyJ, carrying the differential *Ptprc* pan leukocyte marker CD45.1 (Ly5.1), was purchased from The Jackson Laboratory (Stock No. 033076). The animals were housed in the Animal facilities of CHU Ste-Justine, following the guidelines of the Canadian Council on Animal Care (CCAC). The animals were kept in an enriched environment with a 12 h light/dark cycle, fixed temperature, humidity and continuous access to water and food.

Equal cohorts of male and female mice were studied separately for each experiment, and statistical methods were used to test whether the progression of the disease, levels of biomarkers or response to therapy were different for male and female animals. Since differences between sexes were not detected, the data for male and female mice were pooled together.

### Isolation and transplantation of HSPC.

Tibia, femur, and iliac bones were dissected and washed with 70% (v/v) ethanol, followed by 3 washes with PBS containing 1% (w/v) penicillin and streptomycin. Under sterile conditions, bone marrow was flushed with ice-cold SCGM media using 25G syringe, and cells were dissociated with a glass pipette, strained through 70 μm nylon mesh and centrifuged at 380 x g for 10 min at 4°C. The pellet was resuspended in 10 mL of red blood cells (RBC) lysis buffer (155 mM $NH_4Cl$, 12 mM $NaHCO_3$, 0.1 mM EDTA) and incubated for 1 min followed by addition of 20 mL of ice-cold SCGM media. To isolate HSPC, cells were harvested by centrifugation at 380 x g for 10 min at 4°C and resuspended in EasySepT Buffer (Stem Cell Technologies). HSPCs

were purified by negative selection using the EasySep Mouse Hematopoietic Progenitor Cell Isolation Kit (Stem Cell Technologies), according to the manufacturer's instructions, and maintained in culture in 10 mL of SCGM media supplemented with 10 ng/ml rmIL-6, 100 ng/ml rmFlt-3L, 100 ng/ml rmTPO and 100 ng/ml rmSCF at 37°C with 5% $CO_2$. After 24 h, cells were resuspended in PBS, counted and injected in the tail vein of $Hgsnat^{P304L}$ mice ($2.4$-$2.8 \times 10^6$ cells per mouse in 100 μL of PBS). Prior to HSPC transplantation, the recipient mice were treated for 5 consecutive days with busulfan (IP, 25 mg/kg BW/day). During the myeloablation and for 15 days following the transplantation, mice were housed in pathogen-free conditions, fed with food sterilized by irradiation and treated with Baytril® (enrofloxacin, 50 mg/ml in drinking water).

**Flow cytometry analysis of blood leucocytes.**

Blood from the mouse mandibular vein (50 μl-100 μl) was collected in EDTA-coated capillary and was stained for 15 min in the dark at room temperature with APC labelled anti-mouse CD45.1 antibody (Biolegend, 1:20) and PE labelled anti-mouse CD45.2 antibody (Biolegend, 1:20). Erythrocytes were lysed with RBC lysis buffer for 5 min, as described above, and white cells were collected by centrifugation for 5 min at 380 x g and washed twice with PBS containing 1% FBS. Fluorescence compensation was performed with anti-mouse Ig, κ/Negative Control (FBS) Compensation Particles Set (BDTM CompBeads) according to manufacturer's instructions, and flow cytometry analysis was performed using BD FACSCanto II instrument.

**Enzyme activity assays**

Enzymatic activities of HGSNAT and total β-hexosaminidase, were assayed essentially as previously described [18-20]. Tissues extracted from mice were snap-frozen in liquid nitrogen before storage at -80°C. Fifty mg samples were homogenized in 250 μl of $H_2O$ using a sonic homogenizer (Artek Systems Corporation). For HGSNAT assays, 20 μl aliquots of the

homogenates were combined with 10 µl of McIlvain Buffer (pH 5.5), 10 µl of 3 mM 4-methylumbelliferyl-β-D-glucosaminide (Moscerdam), and 10 µl of 5 mM acetyl-coenzyme A in $H_2O$. The reaction was carried on for 3 h at 37°C, stopped with 200 µl of 0.4 M glycine buffer (pH 10.4), and fluorescence was measured using a ClarioStar plate reader (BMG Labtech). Blank samples were incubated without homogenates which were added after the glycine buffer.

The activity of β-hexosaminidase was measured by combining 2.5 µl of 10-times diluted homogenate (~ 2.5 ng of protein) with 15 µl of 0.1 M sodium acetate buffer (pH 4.2), and 12.5 µl of 3 mM 4-methylumbelliferyl N-acetyl-β-D-glucosaminide (Sigma-Aldrich) followed by incubation for 15 min at 37ºC. The reaction was stopped with 200 µl of 0.4 M glycine buffer (pH 10.4), and fluorescence was measured as above.

**Real-time qPCR.**

RNA was isolated from snap-frozen brain tissues using the TRIzol reagent (Invitrogen) and reverse-transcribed using the Quantitect Reverse Transcription Kit (Qiagen # 205311) according to the manufacturer's protocol. qPCR was performed using a LightCycler® 96 Instrument (Roche) and SsoFast™ EvaGreen® Supermix with Low ROX (Bio RAD #1725211) according to the manufacturer's protocol. RLP32 mRNA was used as a reference control. The following forward (F) and reverse (R) primers have been used:

| Gene | Sequence |
|---|---|
| F-IL-1β | TGAAATGCCACCTTTTGACA |
| R-IL-1β | GTAGCTGCCACAGCTTCTCC |
| F-TGF-β1 | GTCAGACATTCGGGAAGCAG |
| R-TGF-β1 | CTGCCGTACAACTCCAGTGA |
| F-RPL32 | TTCTTCCTCGGCGCTGCCTACGA |

| R-RPL32 | AACCTTCTCCGCACCCTGTTGTCA |
| --- | --- |

**Behavioral analysis**

Short-term recognition memory was assessed by Novel object recognition test [21, 22]. Mice were placed individually in a 44 × 45 × 40 cm (length x width x height) testing chamber with white Plexiglas walls for 10 min habituation period and returned to their home cage. The next day, mice were placed for 10 min in the same testing chamber containing two identical objects (red plastic towers, 3 x 1.5 x 4.5 cm), returned to the home cages, and 1 h later, placed back for 10 min into the testing chamber containing one of the original objects and a novel object (a blue plastic base, 4.5 x 4.5 x 2 cm). After each mouse, the arena and the objects were cleaned with 70% ethanol to avoid olfactory cue bias. The discrimination index (DI) was calculated as the difference in the exploration time between the novel and old object divided by the total exploration time. A preference for the novel object was defined as a DI significantly higher than 0 [23]. Mice who showed a side preference, noted as a DI of ± 0.20 during familiarization period, and those who had a total exploration time lower than 3 seconds were excluded from the analysis.

The open-field test was performed as previously described [24]. Briefly, mice were habituated in the experimental room for 30 min before the commencement of the test. Each mouse was, then, gently placed in the center of the open-field arena and allowed to explore for 20 min. The mouse was removed and transferred to its home cage after the test, and the arena was cleaned with 70% ethanol before the next test. The behavioral activity was analysed using the Smart video tracking software (v3.0, Panlab Harvard Apparatus); the total distance traveled, and percent of time spent in the center zone were measured for hyperactivity and anxiety assessment, respectively.

Contextual fear conditioning used to analyze the ability of mice to associate an environment (context) with a fear-inducing stimulus was performed as previously described [25]. Briefly, one day prior to fear conditioning (Day 0), mice were allowed to freely explore the context for habituation for 3 min. On Day 1, mice were trained using a fear-inducing stimulus (5.5 min in total, 3 random 0.6 mA electric shocks, 2 sec each time). On Day 2 and Day 7, conditioned mice were placed in the context chamber for 5 min. The freezing time was calculated based on the video records.

Social interaction behavior was studied using Crawley's sociability and preference for social novelty test as described [26]. Briefly mice were allowed to freely explore a rectangular, three-chamber box with each chamber measuring 19 x 45 cm and the dividing walls made from clear Plexiglas, with an open middle section and a free access to each chamber. Side chambers contained two identical, wire cup containers, one empty and one holding a naïve/unfamiliar mouse. All chambers were cleaned with 70% ethanol after each trial. Mice were habituated for 5 min, after which the walls between the compartments were removed to allow free access for the subject mouse to explore each of the three chambers for the duration of 10 min. The time spend in each chamber and the time spend interacting with wire containers were calculated based on the video records.

**Immunohistochemistry.**

Mouse brains were collected from animals, perfused with 4% PFA in PBS and post-fixed in 4% PFA in PBS overnight. Brains were cryopreserved in 30% sucrose for 2 days at 4°C, embedded in Tissue-Tek® OCT Compound, cut in 40 μm-thick sections and stored in cryopreservation buffer (0.05 M sodium phosphate buffer, pH 7.4, 15% sucrose, 40% ethylene

glycol) at –20°C pending immunohistochemistry. Mouse brain sections were washed 3 times with PBS and permeabilized/blocked by incubating in 5% bovine serum albumin (BSA), 0.3% Triton X-100 in PBS for 1 h at room temperature. Incubation with primary antibodies, diluted in 1% BSA, 0.3% Triton X-100 in PBS, was performed overnight at 4°C. The following antibodies and working concentrations were used:

| Antigen | Host/Target species | Dilution | Manufacturer |
|---|---|---|---|
| GFAP | Rabbit anti-mouse | 1:300 | DSHB (8-1E7-s) |
| Lysosomal-associated membrane protein 2 (LAMP-2) | Rat anti-mouse | 1:200 | DSHB (ABL-93-s) |
| NeuN | Rabbit anti-mouse | 1:200 | Millipore Sigma (MABN140) |
| CD68 | Rabbit polyclonal to CD68 | 1:200 | Abcam (ab125212) |
| $G_{M2}$ ganglioside | Mouse humanized | 1:400 | KM966 |
| β-Amyloid (D54D2) | Rabbit anti-mouse | 1:200 | Cell Signaling (8243S) |
| Recombinant Anti-ATP synthase C antibody (SCMAS) | Rabbit anti-mouse | 1:200 | Abcam (ab181243) |

| IgG | Goat anti-rabbit, anti-mouse, or anti-rat Alexa Fluor 488-, Alexa Fluor 555- or Alexa Fluor 633-conjugated | 1:400 | Thermo Fisher Scientific |
|---|---|---|---|

The mouse brain sections were washed 3 times with PBS and counterstained with Alexa Fluor-labeled secondary antibodies (dilution 1:400) for 2 h at room temperature. After washing 3 times with PBS, the mouse brain sections were treated with TrueBlack® Lipofuscin Autofluorescence Quencher (Biotium, 23007, dilution 1:20) for 1 min, and then, again washed 3 times with PBS. The slides were mounted with Prolong Gold Antifade mounting reagent with DAPI (Invitrogen, P36935) and analyzed using Leica DM 5500 Q upright confocal microscope (10x, 40x, and 63x oil objectives, N.A. 1.4). Images were processed and quantified using ImageJ 1.50i software (National Institutes of Health, Bethesda, MD, USA) in a blinded fashion. Panels were assembled with Adobe Photoshop.

**Flow cytometry analysis of dissociated brain cells and splenocytes.**

The dissociated brain and spinal cord immune cells were obtained as described [27]. Briefly, an intact mouse spinal cord or the dorsal part of the left brain hemisphere, obtained from mice transcardially perfused with saline, were weighted, cut to very small pieces using a scalpel blade and digested using Collagenase D (Roche, Cat No. 11088858001, 1 mg/ml) /DNase I (Thermo Fisher, Cat No. 11875093, 50 μg/ml) solution in RPMI for 30 min at 37°C. Cells were further purified, as described, by centrifugation in a 40-60% Percoll gradient in HBSS to remove debris and myelin [27]. Dissociated splenocytes were obtained from ~1/3 of the entire mouse spleen, as

described [28], by forcing tissue pieces through a 100 μm nylon mesh cell strainer (BD Biosciences, Durham, NC, USA) submerged in ice-cold RPMI 1640. Splenocytes were collected by centrifugation at 300 x g for 5 min at 4°C.

The dissociated cells were labeled with anti-mouse CD45-BV785, CD11b-FITC, CX3CR1-BV605, CD68-BV421, CD63-Pe-Cy7, CD45.1-APC, CD45.2-Pe (all from Biolegend) and CD3-BUV395 (Becton Dickinson) antibodies. The cells were co-stained with the Aqua-Live/Dead fixable dye (LIVE/DEAD™ Fixable Aqua Dead Cell, Invitrogen Thermofisher) and fixed using 1% (v/v) PFA. Flow cytometry analysis was conducted using a Fortessa II instrument (Becton Dickinson) within 48 h following cell labeling and fixation. Data were analyzed using FlowJo X and GraphPad Prism 9.5.0. software.

**Analysis of glycosaminoglycans by LC-MS/MS**

Analysis of glycans in dry blood spots, plasma, urine and brain tissues was conducted as previously described [17]. Briefly, 30-50 mg of mouse tissues were homogenized in ice-cold acetone and centrifuged at 12,000 × g for 30 min at 4°C. The pellets were dried, resuspended in 0.5 N NaOH and incubated for 2 h at 50°C. Then, the pH of the samples was neutralized with 1 N HCl, and NaCl was added to the reaction mix at a final concentration of 3 M. After centrifugation at 10,000 × g for 5 min at room temperature, the supernatants were collected and acidified using 1 N HCl. Following another centrifugation at 10,000 × g for 5 min at room temperature, the supernatants were collected and neutralized with 1 N NaOH to a pH of 7.0. The samples were diluted at a ratio of 1:2 with 1.3% potassium acetate in absolute ethanol and centrifuged at 12,000 × g and 4°C for 30 min. The pellets were washed with ice-cold 80% ethanol, dried at room temperature, and dissolved in 50 mM Tris-HCl buffer. The samples were placed in AcroPrep™ Advance 96-Well

Filter Plates with Ultrafiltration Omega 10 K membrane filters (PALL Corporation, USA) and digested with chondroitinase B, heparitinase, and keratanase II, overnight at 37°C. The samples were analysed by mass spectrometry using a 6460 Triple Quad instrument (Agilent technologies) with Hypercarb columns, as described [17].

**Statistical analysis**

Statistical analyses were performed using Prism GraphPad 9.3.0. software (GraphPad Software San Diego, CA). The normality for all data was verified using the D'Agostino & Pearson omnibus normality test. Significance of the difference was determined using t-test (normal distribution) or Mann-Whitney test, when comparing two groups. One-way ANOVA or Nested ANOVA tests, followed by Tukey multiple comparison test (normal distribution), or Kruskal-Wallis test, followed by Dunn multiple comparisons test, were used when comparing more than two groups. Two-way ANOVA followed by Tukey post hoc test was used for two-factor analysis. A P-value of 0.05 or less was considered significant.

## Results.

### Transplantation of MPS IIIC mice with WT HSPC results in a partial amelioration of the disease symptoms.

To test whether allogeneic hematopoietic cell transplantation provides potential therapeutic benefits for MPS IIIC, we conducted a transplantation of HSPC from WT mice to $Hgsnat^{P304L}$ MPS IIIC mice. This strain expressing a misfolded mutant HGSNAT enzyme shows a faster disease progression, compared to knockout $Hgsnat$-$Geo$ mice, and recapitulates the clinical phenotype of early-onset human patients [16]. The study design is shown in Fig. 1A. HSPC (CD34+) cells were isolated from bone marrow of male B6.SJL-Ptprca Pepcb/BoyJ mice carrying the differential $Ptprc$ pan leukocyte marker CD45.1 (Ly5.1) using EasySep Mouse Hematopoietic Progenitor Cell Isolation kit. Sixteen randomly selected male and female 8-9-week-old $Hgsnat^{P304L}$ mice (Supplementary Table S1) were transplanted with 2.4-2.8x10$^6$ HSPC/mouse after 5-day myeloablation procedure involving daily busulfan injections (IP 25 mg/kg BW/day). Two male mice died on the 4$^{th}$ and 21$^{st}$ days after transplantation. The exact cause of death was not determined, and we cannot exclude that it occurred due to complications related to the transplantation procedure. In addition, one female and one male mouse died 10 and 12 weeks after transplantation due to a malfunction of a water bottle and malocclusion, respectively. Other mice survived until the end of the experiment.

Six weeks after transplantation, engraftment of HSPC was measured by flow cytometry analyses of CD45.1 expression in blood cell lineages. All mice showed engraftment between 82% and 94% (Supplementary Table S1). For 6 untransplanted and 9 transplanted $Hgsnat^{P304L}$ mice as well as for 4 untransplanted WT mice, dry blood spots (DBS) were obtained during the same blood collection procedure and used to analyze GAG levels by LC-MS/MS. This analysis determined

drastically increased levels of HS-derived O-sulfated (ΔDiHS-OS) and N-sulfated (ΔDiHS-NS) disaccharides in the DBS of untreated *Hgsnat^P304L* mice of the same age, while in the transplanted mice the levels of both disaccharides were not significantly different from the normal levels (Supplementary Fig. S1). This suggested that the transplantation completely or almost completely normalized HS levels in the plasma and blood cells of *Hgsnat^P304L* mice.

Between the ages of 6 and 6.5 months (16 and 18 weeks post transplantation) the behaviour of mice was analysed using Open Field (OF, activity and anxiety), Novel Object Recognition (NOR, short-term memory), Contextual Fear Conditioning (CFC, associative learning) and Three-Chamber Sociability (general sociability and autistic-type behaviour) tests. In the OF test, untreated *Hgsnat^P304L* mice showed increased distance traveled in the field and number of entries to the central part of the arena (Fig. 1B). These results, consistent with our previous data [16], are suggestive of hyperactivity and reduced anxiety, respectively. The total distance traveled for the transplanted group was similar to those of the WT group and statistically different from the untreated *Hgsnat^P304L* group showing a rescue of hyperactivity. The transplanted mice also showed a non-significant trend in towards reduction in the number of entries to the central area as compared to untreated group and were not statistically different from the WT group. The NOR test with an interval of 1 h between encoding and retrieval, revealed that *Hgsnat^P304L* mice spend much less time exploring the novel object than WT mice and showed the discrimination index close to zero (Fig. 1C) consistent with impaired short-term memory. For transplanted *Hgsnat^P304L* mice, both parameters showed a non-significant trend for increase and were not statistically different to those of WT mice, suggesting amelioration of the memory deficit (Fig. 1C). In CFC test, 2 and 7 days after the end of the fear-training period, both transplanted and untransplanted *Hgsnat^P304L* mice showed significantly reduced freezing time revealing a problem to associate an

environment with a fear-inducing stimulus (Fig. 1D). Still, transplanted mice showed a trend towards the increase of freezing time compared to untreated *Hgsnat^P304L* mice consistent with a partial recovery of associative learning. Finally, when mice were studied by Three-Chamber Sociability Test that assesses cognition in the form of general sociability, WT mice spent significantly more time in a chamber with an access to a never-before-met intruder mouse than in the identical empty chamber showing a normal sociability typical for rodents (Fig. 1E). In contrast, no statistically significant difference in the time spent in the chambers was detected for untreated *Hgsnat^P304L* mice consistent with MPS IIIC mice exhibiting deficits in social behavior and resembling autistic traits in human Sanfilippo patients. Transplanted mice showed a behaviour similar to WT animals suggesting that they retained a normal sociability (Fig. 1E).

**Figure 1. Transplantation of WT HSPC rescues several behaviour abnormalities in *Hgsnat^P304L* mice.**

**(A)** Experimental design. **(B)** Transplanted *Hgsnat*[P304L] mice show rescue of hyperactivity (total distance traveled) and a trend for recovery of reduced anxiety (number of entries to the center of arena) in the OF test. **(C)** Transplanted mice show a trend for improvement of short-term memory deficit (new object exploration time, discrimination index) in NOR. **(D)** Transplanted mice reveal a trend for improvement of associative learning in the CFC test as compared to untreated mice, but still show a deficit compared with the WT group. **(E)** Transplanted mice demonstrate a recovery of general sociability deficit. Individual results, means and SD of experiments performed with 10-12 male and female mice per genotype, per treatment are shown. P values were calculated using one-way ANOVA with Tukey post hoc test.

Starting from ~7 months (30 weeks) of age, mice were daily evaluated by a veterinarian for the signs of urinary retention (none, mild, moderate, severe or very severe). Urinary retention is rarely seen in the patients, but it is developed by MPS III mice with all genetic subtypes [11, 15, 29]. This trait leads to distortion of the bladder and peritonitis and requires humane sacrifice of severely affected mice thus limiting their lifespan. At the time of first observation, no mice in transplanted group showed urinary retention, while one mouse in the untreated group showed mild and one mouse, moderate signs (Fig. 2A). The symptoms progressed with time and, by the age of 34 weeks, all untreated *Hgsnat*[P304L] mice developed urinary retention (1 very severe, 5 severe and 5 moderate) requiring their immediate sacrifice. In the treated group, only 1 of 11 mice of similar age had moderate signs and 5 mice had no signs of urinary retention (Fig. 2A). Thus allogeneic hematopoietic cell transplantation normalized or partially improved behavior deficits in MPS IIIC mice at the age of 6-6.5 months and delayed the onset of urinary retention, potentially increasing longevity.

All mice (both with and without urinary retention) were sacrificed by the age of 36 weeks (on average 28 weeks after the transplant) to analyze pathological changes in the CNS and peripheral tissues. At sacrifice, we measured the volume of urine retained in the bladder of mice. Untreated *Hgsnat*[P304L] mice had on average 1.5 ml of urine remaining in the bladder, while WT had none or less than 0.3 ml. The amount of urine in the bladders of transplanted *Hgsnat*[P304L] mice was on average equal to 0.5 ml and not significantly different from WT mice, consistent with

the rescue of urinary retention (Fig. 2B). We also measured the wet weight of the visceral organs (liver, kidney and spleen) to assess the extent of organomegaly, reported for the $Hgsnat^{P304L}$ model [15]. Both treated and untreated $Hgsnat^{P304L}$ mice showed drastically increased relative weights of all three organs compared to WT mice (Fig. 2C-E).

**Figure 2. Transplantation of WT HSPC rescues urinary retention in $Hgsnat^{P304L}$ mice but does not alter organomegaly.**
**(A-B)** Development of urinary retention in untransplanted and transplanted $Hgsnat^{P304L}$ and WT mice. (**A**) Urinary retention score (0, none; 1, mild; 2, moderate; 3, severe; 4, very severe) was assessed for untransplanted and transplanted $Hgsnat^{P304L}$ male and female mice, and their WT counterparts (n=16 for each group). The significance of differences between untreated and treated groups was determined by two-

way ANOVA (P<0.0001). By the age of 34 weeks, all untreated *Hgsnat*[P304L] mice but none of transplanted *Hgsnat*[P304L] mice had severe urinary retention requesting euthanasia. (**B**) Volume of urine in the bladder of mice at sacrifice. The amount of urine in the bladders of transplanted *Hgsnat*[P304L] mice is not significantly different from that in WT mice. (**C-E**) Wet organ weight of mice at sacrifice (34-36 weeks). Enlargement of liver (**C**), kidney (**D**) and spleen (**E**) compared to age-matched WT controls, consistent with the lysosomal storage and inflammatory cell infiltration, is detected in both untreated and transplanted *Hgsnat*[P304L] mice. Graphs shows individual data, means and SD. P values were calculated using ANOVA with Tukey post hoc test.

**Transplantation partially increases HGSNAT activity in peripheral tissues and reduces plasma levels of HS.**

The levels of HGSNAT activity and total β-hexosaminidase activity (a marker of lysosomal biogenesis) were measured in the bone marrow, peripheral tissues (liver, spleen, lungs, and kidney) and brain of untransplanted and transplanted *Hgsnat*[P304L] mice. As expected, the HGSNAT activity was fully corrected in the bone marrow and the spleen. In the lungs and liver, the activity was increased on average to about 20% of normal, while in the kidney and the brain HGSNAT activity showed only a minor increase and remained below 10% of normal (Fig. 3A). These results indicate that transplantation of WT HSPC restored levels of the deficient enzyme in bone marrow and hematopoietic cells but not in other peripheral tissues or in the brain. In accordance with our previous findings [16], levels of total β-hexosaminidase activity, measured in the spleen, lungs, liver, kidney, and brain of untreated *Hgsnat*[P304L] mice, showed elevation compared to those in WT mice, consistent with increased levels of lysosomal biogenesis and lysosomal storage. In transplanted *Hgsnat*[P304L] mice, we observed a partial normalization of β-hexosaminidase activity in the lungs, liver, and the brain, but not in the spleen or kidney (Fig. 3B). In the bone marrow, total β-hexosaminidase activity was similar for WT and transplanted or untransplanted mice (Fig. 3B).

The results of HGSNAT and β-hexosaminidase assays were corroborated by the analysis of GAG levels in tissues, blood plasma and urine by LC-MS/MS [15]. Levels of O-sulfated

(ΔDiHS-OS) and N-sulfated (ΔDiHS-NS) disaccharides produced by enzymatic digestion of HS were drastically increased in all tissues as well as in the urine and plasma of untransplanted *Hgsnat*$^{P304L}$ compared to WT mice. In transplanted mice, levels of both disaccharides were reduced approximately two-fold in plasma but remained at the levels similar to those of untreated group in urine, spleen, lungs, liver, kidney, and brain (Fig. 3C, and D). The levels of ΔDi-OS/4S (dermatan sulfate-derived disaccharide), as well as mono-sulfated (KS) and di-sulfated keratan sulfate (DiS-KS) were similar for untreated and treated *Hgsnat*$^{P304L}$ mice and their WT counterparts (not shown) with the exception of spleen, liver, and plasma where KS was increased in untreated *Hgsnat*$^{P304L}$ mice and further normalized by treatment (Supplementary Fig. S2). The increase of KS in the MPS IIIC mice could be potentially associated with a partial inhibition of KS-catabolizing enzyme GALS by HS, but this requires further experimental confirmation.

**Figure 3. Transplanted *Hgsnat^P304L* mice show normalized HS levels in plasma and partial rescue of HGSNAT deficiency and increased lysosomal biogenesis in bone marrow and peripheral tissues, but not in the CNS.**

**(A-B)** HGSNAT activity **(A)** and total lysosomal β-hexosaminidase activity **(B)** in the tissues of 8-month-old WT, *Hgsnat^P304L* and transplanted *Hgsnat^P304L* mice. **(C-D)** Levels of disaccharides produced by enzymatic digestion of HS, ΔDiHS-OS **(C)** and ΔDiHS-NS **(D)** measured by tandem mass spectrometry in blood serum, urine and tissues of WT, *Hgsnat^P304L* and transplanted *Hgsnat^P304L* mice. All graphs show individual data, means and SD of experiments performed using tissues from 5 male and female mice per genotype per treatment except for the bone marrow, where results for 3 samples, each containing pooled tissues of two mice, are shown. P values were calculated by one-way ANOVA with Tukey post hoc test. Only p values <0.05 are shown.

**Transplantation of MPS IIIC mice with WT HSPC reduces neuroinflammation.**

Microglia and tissue macrophages in brain, spinal cord, and spleen of WT, *Hgsnat^P304L* and

transplanted *Hgsnat^P304L* mice were analyzed by flow cytometry of dissociated cells to access their origin (host or transplant) and inflammatory phenotype. The gating strategy for dissociated brain/spinal cord cells and splenocytes is shown in Supplementary Fig. S3.

This analysis revealed that virtually all CD45/CD11b/CX3CR1-positive macrophages in the spleen of transplanted *Hgsnat^P304L* mice showed CD45.1 phenotype indicating that they were derived from transplanted HSPC (Supplementary Fig. S4). In both brain and the spinal cord, approximately 50% of CD45/CD11b/CX3CR1-positive macrophages/microglia cells were also CD45.1-positive (Supplementary Fig. S5) demonstrating that our strategy resulted in effective population of the CNS by macrophages derived from the transplant.

To assess the inflammatory phenotype of macrophages/brain microglia, we estimated fractions of cells expressing high levels of CD63 and CD68, robust markers of activation, in the total population of CX3CR1/CD11b-positive cells. In the spleen of WT mice, only 40% of CX3CR1/CD11b-positive cells expressed CD63 or CD68, while in the untreated *Hgsnat^P304L* mice this number was above 60%. In contrast, in the spleens of transplanted *Hgsnat^P304L* mice the percentage of CD63 or CD68-positive cells was not different from that in the WT mice indicating that the transplantation normalized activation of the resident splenic macrophages (Fig. 4A-C).

Similarly in the brain, the fractions of CD45/CD11b/CX3CR1-positive cells expressing elevated levels of CD63 and CD68 were less than 10% for WT mice, over 80% for untreated *Hgsnat^P304L* mice and about 50% for treated *Hgsnat^P304L* mice (Fig. 4D, E). Moreover, treatment also normalized total number of cells expressing elevated levels of CD63 or CD68 per mg of brain tissue that was drastically increased in the untreated *Hgsnat^P304L* mice (Fig. 4D, E). Consistently with the results of flow cytometry, the mRNA brain levels of pro-inflammatory cytokine IL-1β, expressed by activated microglia/macrophages, were drastically increased in untreated

*Hgsnat^{P304L}* mice compared to controls (Fig. 4F). In the brains of transplanted *Hgsnat^{P304L}* mice, IL-1β expression was reduced compared to untreated *Hgsnat^{P304L}* mice although remained elevated compared to WT controls (Fig. 4F). In contrast, the expression levels of an anti-inflammatory cytokine TGF-β1 showed a non-significant trend towards an increase in the brains of transplanted compared to untransplanted *Hgsnat^{P304L}* mice (Fig. 4G).

**Figure 4. Systemic and brain inflammation is reduced in *Hgsnat^{P304L}* mice transplanted with WT HSPC.**
To estimate inflammatory activation of spleen macrophages, frequency of CX3CR1 labeling on CD11b-positive cells **(A)** and CD63 **(B)** or CD68 **(C)** labeling on CX3CR1/CD11b-positive cells was evaluated in

populations of dissociated splenocytes from WT, *Hgsnat^P304L*, and *Hgsnat^P304L* mice transplanted with WT HSPC (*Hgsnat^P304L–BM)*. To estimate the level of neuroinflammation, frequency for CD63 **(D)** and CD68 **(E)** was evaluated on CD45/CD11b/CX3CR1-positive cells in populations of dissociated brain cells of WT, *Hgsnat^P304L*, and *Hgsnat^P304L–BM* mice. Data are shown as % of CD63+ or CD68+ cells among CD45+/CD11b+/CX3CR1+ cells (graphs on the left) or as total numbers of of CD63+ or CD68+ cells per mg of brain tissue (graphs on the right). **(F-G)** Expression levels of pro-inflammatory cytokine IL-1β **(F)** were reduced in treated compared to untreated *Hgsnat^P304L* mice while the expression levels of an anti-inflammatory cytokine TGF- β1 **(G)** showed a trend for an increase.
Graphs show individual data, means and SD for 5 mice per group (**A-E**) and 3 mice per group (**F-G**). P values were calculated using one-way ANOVA with Dunn post hoc test. Only p values <0.05 are shown.

We further assessed the levels of CD63 and CD68 specifically in the fractions of CD45.1/CD11b/CX3CR1-positive and CD45.2/CD11b/CX3CR1-positive cells to estimate the activation levels of host microglia and those derived from the transplant (Fig. 5A). CD45.1-positive microglia cells in the brains of transplanted *Hgsnat^P304L* mice showed CD63 levels similar to those in the brains of WT mouse donors (Fig. 5B). Moreover, the expression of CD63 in the CD45.2-positive (host) microglia was also significantly reduced compared to untreated *Hgsnat^P304L* mice (Fig. 5B). CD45.1-potitive brain macrophages/microglia also showed ~50% reduction in the CD68 levels, while CD45.2-positive cells showed a trend towards a reduction (Fig. 5C).

**Figure 5. Reduction of neuroinflammation is mediated by infiltration of WT donor derived CX3CR1/CD45.1-positive cells.**
**(A)** Frequency for CD45.1 and CD45.2 was evaluated on CD45/CD11b/CX3CR1-positive dissociated brain cells of WT, untransplanted *Hgsnat^{P304L}*, and transplanted *Hgsnat^{P304L}–BM* mice. **(B-C)** MFI for CD63 **(B)** and CD68 **(C)** was evaluated on CD45/CD11b/CX3CR1-positive dissociated brain cells of WT, *Hgsnat^{P304L}*, and transplanted *Hgsnat^{P304L}–BM* mice. Graphs show individual data, means and SD for 5 mice per group. P values were calculated using one-way ANOVA with Dunn post hoc test.

Together, these data demonstrate that HSPC transplantation resulted in general reduction of neuroinflammation in MPSIIIC mice.

**Transplantation partially normalises the levels of biomarkers of CNS pathology.**

The levels of activated CD68-positive microglia and GFAP-positive astrocytes measured by immunofluorescent analysis of sagittal brain sections were significantly increased in the hippocampi and somatosensory (layers 4-5) cortices of untreated *Hgsnat^{P304L}* compared to WT mice (Fig. 6A-B). The treatment reduced the levels of CD68-positive cells in both brain areas, while levels GFAP-positive cells were normalized in the hippocampus only.

Previously, we have demonstrated that cortical pyramidal neurons accumulate autofluorescent materials that can be labeled with Thioflavin S and antibodies against amyloid-β protein [17]. The same cells also showed labeling with antibodies against misfolded subunit C of mitochondrial ATP synthase (SCMAS) suggestive of mitophagy block and a general impairment of proteolysis. When the levels of all 4 biomarkers were assessed in cortices of untransplanted and transplanted $Hgsnat^{P304L}$ mice, we found that, while the levels of neurons showing autofluorescent inclusions, amyloid deposits and SCMAS were ameliorated in the treated mice, they still remained considerably higher than in the WT cells, suggesting that transplantation could not completely correct this pathology (Fig. 6C-E). We further analyzed levels of a $G_{M2}$ ganglioside, known to be drastically increased in the brains of MPS IIIA-D patients [17] and mouse models [11, 15], using the human-mouse chimeric monoclonal antibody, KM966 [30]. Numerous KM966-positive neurons were observed in the somatosensory cortex layers 4-5 of both treated and untreated $Hgsnat^{P304L}$ mice (Fig. 6F). However, the amounts of $G_{M2}$-positive cells showed a trend towards a decrease in the treated compared to untreated mice. Consistent with unchanged levels of the primary storage product HS and total β-hexosaminidase activity in the brains of treated mice, we found similarly increased size and abundance of LAMP-2 labeled puncta in NeuN-positive neurons in the brains of both treated and untreated $Hgsnat^{P304L}$ mice (Supplementary Fig. S6 A).

**Figure 6.** *Hgsnat$^{P304L}$* **mice transplanted with WT HSPC reveal recovery of astrogliosis and amelioration of levels of biomarkers of CNS pathology in the somatosensory cortex and hippocampus.**

**(A-B)** Levels of activated CD68-positive microglia are reduced in the somatosensory cortex **(A)** and hippocampus **(B)** of 8-month-old transplanted compared to untransplanted *Hgsnat$^{P304L}$* mice. Levels of GFAP-positive astrocytes are reduced in the hippocampus, but not in the cortex. Panels show representative images of somatosensory cortex (layers 4-5) and CA1 region of the hippocampus labeled for GFAP (green) and CD68 (red). DAPI was used as a nuclear counterstain. Bar graphs show quantification of CD68 and GFAP-positive areas with ImageJ software. **(C)** Reduction of thioflavin S-positive β-amyloid aggregates in cortical neurons of transplanted *Hgsnat$^{P304L}$* mice. Panels show representative images of brain cortex (layers 4-5) labeled for β-amyloid (red) and misfolded proteins (Thioflavin-S, green). Graphs show quantification of β-amyloid and Thioflavin-S staining with ImageJ software. **(D)** Levels of misfolded SCMAS aggregates in cortical neurons are reduced in transplanted *Hgsnat$^{P304L}$* mice. Panels show representative images of somatosensory cortex (layers 4-5), labeled for SCMAS (red) and neuronal marker NeuN (green). DAPI (blue) was used as a nuclear counterstain. The bar graph shows quantification of SCMAS staining with ImageJ software. **(E)** Reduction of granular autofluorescent ceroid material in cortical neurons. Panels display representative images of brain cortex showing autofluorescent ceroid inclusions in the neurons (green). The graph shows quantification of autofluorescence with ImageJ software. **(F)** Levels of G$_{M2}$ ganglioside in cortical neurons are not reduced by transplantation. Panels show representative images of somatosensory cortex (layers 4-5) immunolabeled for G$_{M2}$ ganglioside (green). The bar graphs show quantification of G$_{M2}$ staining with ImageJ software.

In all panels scale bars equal 25 μm. All graphs show individual results, means and SD from experiments conducted with 5 mice (three panels per mouse) per genotype per treatment. P values were calculated using Nested one-way ANOVA test with Tukey post hoc test. Only p values <0.05 are shown.

**Discussion.**

MPS IIIC is one of multiple LSDs caused by defects in transmembrane proteins. Since such proteins are uncapable of a free diffusion between the cells and their cross-correction, it was assumed that heterologous HSPC transplantation or autologous HSPC gene therapy would not be effective for this disease. However, since the brain macrophages/microglia derived from the transplanted HSPC are capable of digestion of accumulating substrates, they are expected to develop a neuroprotective M2 phenotype and ameliorate the general levels of neuroinflammation in the recipient patients or animal models by secreting anti-inflammatory cytokines. In the current

work, we tested this hypothesis by conducting a heterologous WT HSPC transplantation of *Hgsnat*$^{P304L}$ MPS IIIC mice homozygous for a pathological missense variant of the HGSNAT enzyme. Since HGSNAT is a multi-spanning transmembrane enzyme, we did not expect a neuronal cross-correction by donor-derived brain macrophages, so all improvements in CNS pathophysiology, if any, would result from the presence of macrophages with a "repair" M2 phenotype.

Our results established that heterologous transplantation of WT HSPC results in substantial amelioration of central manifestations of the disease and rescues urinary retention, the most severe life-threatening peripheral symptom. Unlike untreated, transplanted *Hgsnat*$^{P304L}$ mice did not develop severe urinary retention by the age of 8 months, suggesting that the transplantation corrected inflammatory changes in the bladder walls and/or urethra, thought to be responsible for this trait [31]. Likewise, the levels of HS in the blood, measured 6 weeks after the transplantation, were completely normalized and remained 50% lower than those in the untreated mice at least until the age of 8 months. Surprisingly, the GAG levels in the liver, lungs, spleen, kidney, and urine remained similar to those in the untreated mice. For the spleen, these results were somewhat unexpected considering that the vast majority of spleen macrophages originated from the donor HSPC, and HGSNAT activity levels were normalized. It's tempting to speculate that HS level in the spleen reflects the accumulation of this substrate, that occurred before the transplant and was located in the areas not accessible to the infiltrating WT macrophages. Further studies are necessary to confirm this hypothesis. In other peripheral tissues, levels of HGSNAT activity showed a variable recovery to the levels from <3% of the WT in the kidney to 18-22% of the WT in lungs and liver. In the last two organs, we also observed a significant reduction in the levels of total β-hexosaminidase, consistent with partial normalization of lysosomal biogenesis but, like in

the spleen, we did not observe any reduction of HS stored in these tissues.

In the brain of transplanted mice, at least 50% of microglia/macrophages were derived from the donor HSPC, which was consistent with the numbers observed in other studies that used Busulfan conditioning [7, 9]. At the same time, HGSNAT activity level in the brain on average remained below ~10% of the WT level, reflecting the contribution of the microglia to the total pool of this enzyme in the CNS. The absence of the enzyme deficiency correction was consistent with unchanged brain levels of HS and lysosomal storage/lysosomal biogenesis markers, LAMP-2 and total β-hexosaminidase. These results are highly consistent with those obtained for the mouse models of MPS IIIA and MPS IIIB, transplanted with WT HSPC [32, 33]. Although both diseases are caused by defects in soluble enzymes, their levels of secretion from WT HSPC-derived brain macrophages were not sufficient for cross-correction of neurons or reduction of HS [32, 33].

Other CNS disease biomarkers showed a diverse response to the therapy. The secondary storage materials, misfolded β-amyloid, SCMAS and autofluorescent ceroid inclusions in the pyramidal cortical neurons, all associated with the blockage of autophagy, were reduced while the levels of $G_{M2}$ ganglioside remained unchanged reflecting the absence of cross-correction of neurons by macrophages. At the same time, levels of GFAP-positive astrocytes were drastically decreased in hippocampus, and the levels of CD68-positive activated microglia decreased in both cortex and hippocampus suggestive of reduced neuroinflammation. This was further confirmed by decrease in expression levels of proinflammatory cytokine IL-1β but not anti-inflammatory cytokine TGF-β1. Analysis of immune cells in the brain by flow cytometry provided an explanation for these observations: at least 50% of microglia in the brain of transplanted mice were CD45.1-positive and showed levels of activation markers similar to those in the brains of the WT mice. Moreover, activation of the resident CD45.2-positive macrophages was also reduced

compared to the brains of untreated $Hgsnat^{P304L}$ mice, indicating that the donor-derived microglia could mitigate the neuroimmune response.

Most importantly, transplanted $Hgsnat^{P304L}$ mice showed a normalization of hyperactivity and sociability, and amelioration of reduced anxiety and short-term memory deficit, while the deficient associative learning showed only a partial improvement. These positive effects were in general not expected, since the overwhelming majority of neurons remained devoid of the defective enzyme. We speculate, therefore, that the observed effects are mainly due to non-cell-autonomous effects of transplanted cells on neuroimmune response. This is also supported by published results showing that correction of neuroinflammation by hematopoietic stem cell gene therapy with IL-1Ra completely prevented the development of hyperactivity and spatial memory loss in MPS IIIA mice [34]. In contrast this report, however, our data provide no evidence for complete rescue of neuronal pathology and behavior deficits. One explanation for this could be that, in our study, HSPC transplantation was conducted at 2 months (8 weeks) of age. According to our published data [35], at this age, pyramidal neurons of MPS IIIC mice already show synaptic deficits and significantly reduced density of dendritic spines. Similarly, levels of activated microglia show a drastic increase already at P25 [36]. Further studies are required to determine whether transplantation of mice at an earlier age would show efficacy in preventing CNS deficits, however since most of MPS IIIC patients are diagnosed post-symptomatically, we believe that our experimental design reflects the current patient situation. Also, it remains to be determined, if transplantation of gene-corrected autologous HSPC overexpressing supraphysiologic HGSNAT levels would correct the brain pathology as it did for the mouse models of MPS IIIA and MPS IIIB, two MPS III subtypes caused by deficiencies of soluble enzymes [6, 33]. Interestingly, HGSNAT has been detected in the extracellular vesicles (exosomes) isolated from culture medium

conditioned with HEK 293T cells overexpressing the enzyme [37]. Thus, HGSNAT-loaded

exosomes derived by microglia overexpressing the enzyme could potentially provide for a certain

level of neuronal cross-correction. Of note, our recent study [15] revealed that MPS IIIC mice

treated by a HGSNAT chaperone, glucosamine, showed an improvement of CNS pathology,

except for neuroinflammation that, as we show in the current work, is effectively treated by the

HSPC transplant. It is tempting to speculate, therefore, that heterologous HSPC transplant used

together with small molecule drugs either stabilizing a mutant HGSNAT enzyme and increasing

its activity levels in the neurons (chaperone therapy) or reducing HS production (substrate

reduction therapy) may be an effective component of a combination therapy for MPS IIIC.

**Abbreviations** BSA = bovine serum albumin; DBS = dry blood spots; DAPI = 4′,6-diamidino-2-phenylindole; CNS = central nervous system; GAGs = glycosaminoglycans; HGSNAT = heparan sulfate acetyl-CoA: α-glucosaminide N-acetyltransferase**;** HS = heparan sulfate; HSPC = hematopoietic stem and progenitor cells; KS = keratan sulfate; LAMP-2 = lysosome-associated membrane protein 2; LC-MS/MS = liquid chromatography tandem mass spectrometry; LSDs = lysosomal storage disorders; MFI = mean fluorescence intensity; MPS = mucopolysaccharidosis; NOR = novel object recognition; OF = open field; PBS = phosphate-buffered saline; SCMAS = subunit C of mitochondrial ATP synthase; YM = Y-maze.

**Acknowledgements**
The authors thank Dr. Elke Küster-Schöck and the Plateforme d'Imagerie Microscopique (PIM – CHU Sainte Justine) for the help with life imaging microscopy, and Dr. Mila Ashmarina for critically reading the manuscript and helpful advice.

**Funding**
This work has been partially supported by operating grants PJT-156345 and PJT-180546 from the Canadian Institutes of Health Research and a research grant ND-1 from the Canadian Glycomics Network to A.V.P., Elisa Linton Research Chair in Lysosomal Diseases and gifts from Cure Sanfilippo Foundation and Sanfilippo Children's Foundation (Australia) to A.V.P.

**Availability of data and materials**
The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

**Authors' contributions**
Conducted experiments and acquired data: X.P.; A.C., S.F., S.K.; analyzed data: X.P.; A.C., S.F.,

S.K., S.T. and A.V.P.; provided essential resources: S.T. and A.V.P., wrote the manuscript (first draft): A.V.P., X.P., A.C.; wrote the manuscript (editing): S.T. and A.V.P. All authors read and approved the final manuscript.

**Ethics approval**
Approval for the animal experimentation was granted by the Animal Care and Use Committee of the CHU Sainte-Justine.

**Consent for publication**
All authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work. All authors gave their consent for this publication.

**Competing interests**
A.V.P. received honoraria and research contracts from Phoenix Nest Inc involved in development of therapies for MPS IIID and IIIC. Authors have no additional financial interests.

# References

1.    Giugliani, R., *Newborn screening for lysosomal diseases: current status and potential interface with population medical genetics in Latin America.* J Inherit Metab Dis, 2012. **35**(5): p. 871-7.
2.    Mader, K.M., et al., *Effect of high dose, repeated intra-cerebrospinal fluid injection of sulphamidase on neuropathology in mucopolysaccharidosis type IIIA mice.* Genes Brain Behav, 2008. **7**(7): p. 740-53.
3.    Bonney, D.K., et al., *Successful allogeneic bone marrow transplant for Niemann-Pick disease type C2 is likely to be associated with a severe 'graft versus substrate' effect.* J Inherit Metab Dis, 2010. **33 Suppl 3**: p. S171-3.
4.    Krivit, W., et al., *Microglia: the effector cell for reconstitution of the central nervous system following bone marrow transplantation for lysosomal and peroxisomal storage diseases.* Cell Transplant, 1995. **4**(4): p. 385-92.
5.    Biffi, A., et al., *Correction of metachromatic leukodystrophy in the mouse model by transplantation of genetically modified hematopoietic stem cells.* J Clin Invest, 2004. **113**(8): p. 1118-29.
6.    Sergijenko, A., et al., *Myeloid/Microglial driven autologous hematopoietic stem cell gene therapy corrects a neuronopathic lysosomal disease.* Mol Ther, 2013. **21**(10): p. 1938-49.
7.    Wilkinson, F.L., et al., *Busulfan conditioning enhances engraftment of hematopoietic donor-derived cells in the brain compared with irradiation.* Mol Ther, 2013. **21**(4): p. 868-76.
8.    Biffi, A., et al., *Lentiviral hematopoietic stem cell gene therapy benefits metachromatic leukodystrophy.* Science, 2013. **341**(6148): p. 1233158.
9.    Capotondo, A., et al., *Brain conditioning is instrumental for successful microglia reconstitution following hematopoietic stem cell transplantation.* Proc Natl Acad Sci U S A, 2012. **109**(37): p. 15018-23.
10.   Sessa, M., et al., *Lentiviral haemopoietic stem-cell gene therapy in early-onset metachromatic leukodystrophy: an ad-hoc analysis of a non-randomised, open-label, phase 1/2 trial.* Lancet, 2016. **388**(10043): p. 476-87.
11.   Martins, C., et al., *Neuroinflammation, mitochondrial defects and neurodegeneration in mucopolysaccharidosis III type C mouse model.* Brain, 2015. **138**(Pt 2): p. 336-55.
12.   Baregamian, N., et al., *Tumor necrosis factor-alpha and apoptosis signal-regulating kinase 1 control reactive oxygen species release, mitochondrial autophagy, and c-Jun N-terminal kinase/p38 phosphorylation during necrotizing enterocolitis.* Oxid Med Cell Longev, 2009. **2**(5): p. 297-306.
13.   Chen, X.H., et al., *TNF-alpha induces mitochondrial dysfunction in 3T3-L1 adipocytes.* Mol Cell Endocrinol, 2010. **328**(1-2): p. 63-9.
14.   Vitner, E.B., et al., *Contribution of brain inflammation to neuronal cell death in neuronopathic forms of Gaucher's disease.* Brain, 2012. **135**(Pt 6): p. 1724-35.
15.   Pan, X., et al., *Glucosamine amends CNS pathology in mucopolysaccharidosis IIIC mouse expressing misfolded HGSNAT.* Journal of Experimental Medicine, 2022. **219**(8): p. e20211860.
16.   Pan, X., et al., *Glucosamine amends CNS pathology in mucopolysaccharidosis IIIC mouse expressing misfolded HGSNAT.* J Exp Med, 2022. **219**(8).
17.   Viana, G.M., et al., *Brain Pathology in Mucopolysaccharidoses (MPS) Patients with Neurological Forms.* J Clin Med, 2020. **9**(2).
18.   Seyrantepe, V., et al., *Mice deficient in Neu4 sialidase exhibit abnormal ganglioside catabolism and lysosomal storage.* Hum Mol Genet, 2008. **17**(11): p. 1556-68.
19.   Potier, M., et al., *Fluorometric assay of neuraminidase with a sodium (4-methylumbelliferyl-α-d-N-acetylneuraminate) substrate.* Analytical Biochemistry, 1979. **94**(2): p. 287-296.
20.   Guo, T., et al., *Selective Inhibitors of Human Neuraminidase 3.* Journal of Medicinal Chemistry, 2018. **61**(5): p. 1990-2008.
21.   Antunes, M. and G. Biala, *The novel object recognition memory: neurobiology, test procedure, and its modifications.* Cogn Process, 2012. **13**(2): p. 93-110.
22.   Akkerman, S., et al., *Object recognition testing: methodological considerations on exploration and discrimination measures.* Behav Brain Res, 2012. **232**(2): p. 335-47.
23.   Ennaceur, A. and J. Delacour, *A new one-trial test for neurobiological studies of memory in rats. 1: Behavioral data.* Behav Brain Res, 1988. **31**(1): p. 47-59.
24.   Amegandjin, C.A., et al., *Sensitive period for rescuing parvalbumin interneurons connectivity and social behavior deficits caused by TSC1 loss.* Nat Commun, 2021. **12**(1): p. 3653.
25.   Lavertu-Jolin, M., et al., *Acan downregulation in parvalbumin GABAergic cells reduces spontaneous recovery of fear memories.* Mol Psychiatry, 2023. **28**(7): p. 2946-2963.
26.   Kaidanovich-Beilin, O., et al., *Assessment of social interaction behaviors.* J Vis Exp, 2011(48).
27.   Manglani, M., S. Gossa, and D.B. McGavern, *Leukocyte Isolation from Brain, Spinal Cord, and Meninges for Flow Cytometric Analysis.* Curr Protoc Immunol, 2018. **121**(1): p. e44.
28.   Comeau, K., et al., *Determination of Interleukin-17A and Interferon-gamma Production in gammadelta,*

*CD4(+), and CD8(+) T Cells Isolated from Murine Lymphoid Organs, Perivascular Adipose Tissue, Kidney, and Lung.* Bio Protoc, 2023. **13**(10): p. e4679.

29.	Li, H.H., et al., *Mouse model of Sanfilippo syndrome type B produced by targeted disruption of the gene encoding alpha-N-acetylglucosaminidase.* Proc Natl Acad Sci U S A, 1999. **96**(25): p. 14505-10.

30.	Nakamura, K., et al., *Chimeric anti-ganglioside GM2 antibody with antitumor activity.* Cancer Res, 1994. **54**(6): p. 1511-6.

31.	Marco, S., et al., *Progressive neurologic and somatic disease in a novel mouse model of human mucopolysaccharidosis type IIIC.* Dis Model Mech, 2016. **9**(9): p. 999-1013.

32.	Langford-Smith, A., et al., *Hematopoietic stem cell and gene therapy corrects primary neuropathology and behavior in mucopolysaccharidosis IIIA mice.* Mol Ther, 2012. **20**(8): p. 1610-21.

33.	Holley, R.J., et al., *Macrophage enzyme and reduced inflammation drive brain correction of mucopolysaccharidosis IIIB by stem cell gene therapy.* Brain, 2018. **141**(1): p. 99-116.

34.	Parker, H., et al., *Haematopoietic stem cell gene therapy with IL-1Ra rescues cognitive loss in mucopolysaccharidosis IIIA.* EMBO Mol Med, 2020. **12**(3): p. e11185.

35.	Para, C., et al., *Early defects in mucopolysaccharidosis type IIIC disrupt excitatory synaptic transmission.* JCI Insight, 2021.

36.	Taherzadeh, M., et al., *Severe central nervous system demyelination in Sanfilippo disease.* Front Mol Neurosci, 2023. **16**: p. 1323449.

37.	Fedele, A.O., et al., *Lysosomal N-acetyltransferase interacts with ALIX and is detected in extracellular vesicles.* Biochim Biophys Acta Mol Cell Res, 2018. **1865**(10): p. 1451-1464.