Can ecological interactions drive evolutionary outcomes? Evidence from insect host shifts between parasitic and non-parasitic plants

Zelvelder B.1,2,*, Kergoat G.J.2, Benoit L.3, Tsuchida T.4, Haran J.3 & Allio R.2,*

1 ISEM, Univ Montpellier, CNRS, IRD, Montpellier, France. ORCID: 0009-0007-8486-2511 (BZ; benjamin.zelvelder@umontpellier.fr)

2 Centre de Biologie pour la Gestion des Populations, INRAE, IRD, CIRAD, Institut Agro, Univ. Montpellier, Montpellier, France. ORCID: 0000-0002-8284-6215 (GJK; gael.kergoat@inrae.fr); 000-0003-3885-5410 (RA; remi.allio@inrae.fr)

3 Centre de Biologie pour la Gestion des Populations, CIRAD, INRAE, IRD, Institut Agro, Univ. Montpellier, Montpellier, France. ORCID: 0000-0003-3740-5346 (LB; laure.benoit@cirad.fr); 0000-0001-9458-3785 (JH; julien.haran@cirad.fr)

4 Faculty of Science, Academic Assembly, University of Toyama, 3190 Gofuku, Toyama, Toyama 930-8555, Japan. ORCID: 0000-0001-6065-4469 (TT; tsuchida@sci.u-toyama.ac.jp)

*Corresponding authors: Rémi Allio (+33689908969, remi.allio@inrae.fr) & Benjamin Zelvelder (+33652599011, benjamin.zelvelder@umontpellier.fr), 755 avenue du campus Agropolis, 34988, Montferrier-sur-lez, France

Key words: coevolution, *Cuscuta*, Curculionidae, host plant shift, parasitic plants, phylogenomics, phytophagous insects, plant-insect interactions, plant-plant interactions, target capture

Abstract

Phytophagous insects have specialized on virtually every plant lineage. Parasitic plants, however, are uncommon hosts. Among insects, only a single lineage of weevils, the Smicronychini, has successfully radiated on both parasitic and non-parasitic plants in a large panel of distantly related Asterid families. This unusual pattern suggests that major host plant shifts have occurred over the course of their diversification. Through the analysis of a phylogenomic dataset, we reconstruct for the first time their evolutionary history and ancestral host plant associations. Our results show that independent host plant shifts occurred both from parasitic to non-parasitic hosts and between distinct parasitic lineages. These results suggest that host shift mechanisms can be driven by ecological opportunities provided by plant-plant interactions. This first evidence of extreme insect host plant shifts apparently mediated by parasitic plant-plant interactions emphasizes the core importance of ecological interactions as driving forces behind insect host plant shifts.

Introduction

Plant-insect interactions are one of the main components shaping terrestrial ecosystems and have therefore been extensively studied throughout all major fields of Ecology and Evolution. Indeed, studying these interactions offer a promising opportunity for bridging micro- and macroevolutionary processes [1]. To apprehend the fundamental processes associated with the evolution of plant-insect interactions, considering the notion of host repertoire (*sensu* [2]) is particularly relevant. Echoing the Hutchinsonian concept of the ecological niche [3] (see also [4]), phytophagous insects have a *fundamental* repertoire of suitable hosts and a *realized* repertoire of hosts actually used by the insect in its environment [2]. The *realized* repertoire of hosts can either remain the same (i.e. niche conservatism [5]) or be subject to evolution via: specialization (i.e. fewer hosts [6-7]), diet expansion (i.e. hosts acquisition without the loss of ancestral hosts [8]) or host plant shifts (i.e. hosts acquisition with the loss of ancestral hosts [9]; see also [2]). With this point of view, host plant shifts are often seen as the most extreme scenarios as they can be seen as a shift to a distinct biotic environmental space [10-11]. Adaptation to new host plants requires the adaptation to new structures, plant defenses, phenologies, micro-climates and so on (e.g. [12-18]). Altogether, this multidimensional nature of host plant shifts is *de facto* much more challenging to study than other processes. As a result, underlying forces that enable such changes in the host repertoire of phytophagous insects remain poorly understood.

The evolution of host repertoire in phytophagous insects' phylogenies follow these general patterns: *(i)* host plant shifts are ubiquitous events at large taxonomic scales, *(ii)* phytophagous insect phylogenies do not match host plant phylogenies in most cases, and *(iii)* phylogenetic niche conservatism of host associations is more common at shallower taxonomic scales (see the reviews of [2,19-21]). The description of these patterns led to the formulation of several macroevolutionary scenarios such as the classical 'escape & radiate' arms race by Ehrlich and Raven [22], the oscillation hypothesis [23], and the musical chairs hypothesis [24]. All these scenarios posit that host plant shifts are diversification drivers, a claim that has often been demonstrated on empirical datasets [24-28] (but see [21,29-31]). Host plant shifts have also been shown to be at the origin of genetic divergence between lineages, sometimes leading to reproductive isolation and even to ecological speciation (see the review of [32]). Overall, the signature of host plant shift and adaptations to new hosts have been extensively studied, from complex genetic changes [26,28,33] (see also the reviews of [34-35]) to transcriptomic signatures [11,36-37] and even morphological plasticity [38]. However, although the importance of this phenomenon in the evolution of arthropods has been widely acknowledged, how host plant shifts occur in the first place and what drives the directionality of these events has received very little attention.

Under the right conditions, it appears that phytophagous insects can easily make host plant shifts, at least towards plants in their fundamental repertoire. For example, through a process called ecological fitting (*sensu* [39]), numerous host plant shifts have been documented following geographical opportunities, such as new potential hosts introduction or parasites invasion in a new geographical range [40-43]. For those species, it seems that the only barrier to host shifts was the host's geographical range. Other barriers associated with the plant as a niche, such as the plant's phenology [44-45], physical structures (e.g. [46-47]), detectability [48], and physiology or chemistry (e.g. [12,49]), can also be overcome and ultimately lead to host plant shifts. These barriers therefore hold an important role in the respective diversification of plants and insects [22,26,50-53]. The phylogenetic proximity between related plants that, on average, share most of these characteristics, may in turn explain the ubiquitous 'minor' host plant shifts observed at shallow taxonomic scales [39,54-55]. Nevertheless, several experiments have succeeded in forcing host shifts towards plants supposedly outside of the insect's host repertoire, showing that, under the right conditions (i.e. a consistent exposition to a new host), host preference is a relatively labile trait (e.g. [38,56]). Therefore, the key factors that explain 'major' insect host plant shifts - between distantly related plant species - may rely on rarer ecological opportunities that

attenuate those barriers, allowing insects to overcome them. Interestingly, one such opportunity can be found *in natura* thanks to the interactions between parasitic plants and their hosted phytophagous insects.

Parasitic plants (*ca.* 4,500 species from 28 families) have a unique heterotrophic lifestyle associated with the development of a multicellular organ - the haustorium - which allows them to absorb water and nutrients but also proteins, nucleotides or pathogens [57]. Phytophagous insects that live on parasitic plants are directly impacted by the nature of the compounds exchanged with the parasitized plants [58-59]. As a result, phytophagous insects that feed on parasitic plants evolve in unique circumstances of physiological proximity between phylogenetically distant plants. To date, only one group of insects is known to have radiated on parasitic plants, that being the weevils of the tribe Smicronychini (Coleoptera: Curculionoidea; [60-67]).

Smicronychini weevils form a widespread tribe known from all continents except Antarctica that consist of *ca*. 200 species classified in six genera, the genus *Smicronyx* being by far the most diverse with 156 described species [63]. They are essentially monophagous or narrowly oligophagous on closely related plants and their endophagous larvae develop either in stems, roots, seeds or fruits, sometimes inducing galls in their host (see figure 1 and Table S1 for details, [63]). Although this tribe is known for its association with parasitic plants, it has a diverse repertoire of phylogenetically unrelated hosts plants, belonging to four Asterid families: some Smicronychini species specifically develop in holoparasitic plants (obligatory parasites with no or little photosynthetic activity [68]), like dodders (*Cuscuta* - Convolvulaceae) or broomrapes (*Orobanche* - Orobanchaceae), others develop in hemiparasitic plants (obligatory or facultative parasites that still have some degree of photosynthetic activity [68]; such as *Euphrasia* or *Striga* - Orobanchaceae), while the rest develop in non-parasitic plants belonging to the Asteraceae, Convolvulaceae and Gentianaceae families (figure 1; Table S1; [69]). At least one species from North America, *Smicronyx quadrifer,* is known to feed in both parasitic and non-parasitic plants, starting its larval development in a *Cuscuta* species and completing it in one of a few Asteraceae hosts parasitized by the *Cuscuta* [61]. Based on this striking ecological observation and because Smicronychini have a remarkably diverse host repertoire compared to other tribes of Curculioninae [70-72], we hypothesize that the close physiological proximity between distantly related plants provided by parasitic plant-plant interactions may give unique host plant shift opportunities to insects feeding on them.

Figure 1: Host repertoire of Smicronychini weevils. From top to bottom: diagrams represent parasitic interactions between Smicronychini weevils and their host plants. Host plant families that make up the host repertoire of Smicronychini are shown below each corresponding diagram, with examples of associated weevils and known galls. Credits: B.Z., J.H., except for Orobanchaceae (Clémence Massard) and Asteraceae (@sinaloasilvestre, iNaturalist) pictures; diagrams with ©BioRender.

In this study, we address this hypothesis at the macroevolutionary scale by reconstructing ancestral host plant associations of the Smicronychini and propose a possible evolutionary scenario that may explain the host plant shift patterns inferred. We provide the first molecular phylogeny of the tribe following a two-stage approach. Firstly, we gathered and assembled hundreds of genome-wide markers using Anchored Hybrid Enrichment (AHE) capture, thus taking advantage of specimens held in museum collections to sample a representative dataset of the tribe. Secondly, we expanded our taxon sampling by grafting additional Smicronychini specimens/species for which mitochondrial markers (subunit I of the cytochrome *c* oxidase: COI) data was available. We then used the resulting phylogeny in combination with species-level ecological information to reconstruct the evolution of ancestral associations between Smicronychini and their host plants.

Materials and methods

Taxon sampling

The taxon sampling consists of 67 specimens intended for target capture (hereafter *AHE dataset*) and 205 specimens for which COI data were already available or newly generated (hereafter *COI dataset*).

AHE dataset - Smicronychini weevils were collected throughout ten years of field work carried out by J. Haran and by collecting specimens from Museum collections (Table S2). This sampling covers the Afrotropical, Palearctic and Oriental bioregions, representing all major genera encountered in this geographical range (i.e. *Afrosmicronyx*, *Sharpia*, *Smicronyx* and *Topelatus*), with the exception of the monospecific genus *Hedychrous*. Additional individuals were sampled in seven tribes (Anthonomini, Cionini, Curculionini, Derelomini, Erirhinini, Rhamphini and Styphlini) related to Smicronychini to serve as outgroups for phylogenetic inferences. These tribes were selected based on the most recent and comprehensive phylogenomic study of the Curculioninae subfamily [72].

162 *COI dataset* - COI sequences of individuals from the *AHE dataset* have been extracted
163 using MitoFinder v1.4.1 [73]. When no COI sequence was found with MitoFinder, when
164 available, COI sequences generated from the same sample or from different individuals of
165 the same species were assigned to these individuals. This dataset allowed us to expand our
166 taxon sampling and include taxa native to the Nearctic bioregion (poorly represented in the
167 *AHE dataset*). Species from the genus *Promecotarsus* and the *Smicronyx* subgenera
168 *Pachyphanes*, *Pseudosmicronyx* and *Smicronyx* were thus added, along with additional COI
169 sequences of species from the *AHE dataset*. Those were either generated by our research
170 group using specimens from museum collections or downloaded from GenBank (NCBI) or
171 private and public data from BOLD Systems v4 [74] (see Table S2 for specimen details).

172 This sampling covers the entire morphological diversity and known host repertoires of the
173 Smicronychini, considering both botanical families and the parasitic ecology of some of
174 their hosts from the literature (Table S1), making it the most comprehensive sampling ever
175 assembled for this tribe.

176 **Target capture and DNA sequencing**
177 Specimens from the *AHE dataset* were prepared following the target enrichment protocol of
178 [75]. Tissues from ethanol- and dry-preserved specimens were extracted non-destructively
179 using the EZ-10 Spin Column Animal Genomic DNA Miniprep Kit (Biobasic Inc., Canada)
180 with an overnight lysis step. DNA concentration was normalized (except for low quality
181 samples) and mechanically fragmented to 300-600 bp fragments using a Bioruptor Pico
182 (Diagenode; 15s ON, 90s OFF, eight cycles). Low quality DNA were then reconcentred with
183 AMPure XP beads and libraries were constructed according to the NEBNext® Ultra II DNA
184 Library Prep for Illumina® kit recommendations (New England Biolabs, Ipswich, MA,
185 USA). Each library was individually labeled and subjected to 12 PCR cycles for final
186 enrichment and pooled in equal quantities, grouping 24 libraries. The myBaits Hyb Capture
187 for AHE kit (Arbor Biosciences, Ann Arbor, MI, USA) was applied to these two pools
188 following the instructions of the manufacturer and assayed by qPCR. The resulting AHE
189 library was sequenced paired-end, on a 2x150 bp SP line on an Illumina NovaSeq sequencer
190 (Illumina, San Diego, CA, USA) at the GenomiX platform in Montpellier, France (MGX).

191
192 **Adapting probes to our dataset**
193 The set of probes used during target enrichment was designed on 941 coding loci anchored
194 in informative flanking regions within the genomes of distantly related coleopteran groups
195 [76-77]. To improve the recovery of targeted loci on a shallower taxonomic level, we

generated a new set of AHE reference sequences based on the complete genome of the boll weevil *Anthonomus grandis* (Curculionidae: Curculioninae; GCF_022605725.1, NCBI), the closest genome to Smicronychini at the time of this study (based on [72]). We mapped the entire probe set made available by Haddad et al. [76] on gene sequences of *A. grandis* using the *tblastx* command from BLAST+ 2.9.0 [78], which successfully retrieved all 941 genes targeted by the AHE probe set. Resulting matches were then filtered manually prior to phylogenomic analyses to remove 40 duplicated genes that mapped on multiple loci and 69 loci containing at least one insert more than 500 nucleotides long (out of 137 loci containing inserts in the putatively conserved region). The resulting set of 832 AHE reference sequences was further used to conduct assembly and alignment.

AHE assemblies and alignments

A specific pipeline was developed to process AHE data (figure S1). Reads were first demultiplexed (i.e. reassigning reads to each individual on the basis of their tags used during target capture) with cutadapt v2.8 [79] and cleaned with Trimmomatic v0.39 [80]. Resulting reads were then assembled using a custom version of IBA [81]. This program follows an iterative approach to map reads on given reference sequences. Loci obtained from IBA assembly were then aligned to their corresponding AHE reference sequences using MAFFT v7.453 with *--add* and *--adjustdirectionaccurately* options [82-83]. We used the L-INS-I model, designed to be robust to divergent flanking regions around a core easier to align, which corresponds exactly to the AHE structure. To avoid any bias linked to missing data [84], we only retained loci represented by more than 70% of the individuals in our dataset, a value established as a good ratio between the quantity of loci retained and total percentage of missing data. Although AHE probes target coding regions, assembled sequences can be surrounded by non-coding flanking regions. Alignments were therefore cleaned up differently depending on their coding or non-coding nature. Firstly, partitions were defined based on the region targeted by the probes (custom script: *make_gapped_parts.py*). Secondly, using the annotation of the aforementioned genome of *A. grandis*, we were able to refine the boundaries of coding and non-coding regions more precisely (custom R script: *improve_parts.R*). This process led to the identification of potential overlapping sequences between AHE loci that were consequently pruned with cutadapt. Thirdly, coding and non-coding partitions were split using AMAS [85] and processed with the OMM_MACSE pipeline [86] or HMMCleaner [87] followed by PaulBlocks (provided by Paul Simion; to remove positions with more than 50% of missing data), respectively. All of these alignments were then concatenated into one supermatrix and translated into usable formats for

phylogenetic inference using *seqCat.pl* and *seqConverter.pl*. This pipeline (scripts and details) is available at Zenodo (*https://doi.org/10.5281/zenodo.15098685*).

Combined phylogenetic approach

Phylogenomic inferences based on AHE dataset - Phylogenetic inferences were conducted on the CIPRES Gateway cluster [88] within a Maximum Likelihood (ML) framework using IQTREE v2.3.2 [89-90]. The supermatrix was partitioned so that each flanking region, each inserted region and each coding position (figure 2B) was allowed to have its own evolution rate using the -*p* option [91]. Model estimation was systematically conducted using the greediest and most accurate method available in IQ-TREE using ModelFinder (-*MFP+MERGE* option) with -*rclusterf 20* to improve partition merging performance and the Bayesian information criterion (BIC) as a metric for model selection [92-93]. A perturbation level of heuristics of 0.2 (-*pers 0.2* option; recommended for a phylogenomic dataset according to IQTREE documentation) was implemented and symmetry tests were performed using the --*symtest-remove-bad* option [94]. The robustness of the nodes was estimated with ultra fast bootstraps (UFBS; [95]), with the -*bnni* option to compensate for model violations and the Shimodaira-Hasegawa approximate likelihood-ratio test (SH-aLRT; [96]), supposed to be more robust to the latter. A node was considered highly-supported for values of SH-aLRT $\geq$ 80 and UFBS $\geq$ 95% (following IQTREE authors' recommendations).

Extending the phylogeny with the COI dataset - The tree obtained from the *AHE dataset* was pruned to contain only ingroup individuals (from the Smicronychini tribe). This tree was then used to constrain a second ML phylogenetic inference (-*g* option) on the *extended dataset* (i.e. including both *AHE* and *COI datasets*, excluding outgroups). This approach allows to densify the taxon sampling while retaining high confidence in the relationships at the species-level [97] (see [98-100] for empirical examples).

Ancestral character state reconstructions

Ancestral character state reconstructions (ASR) of host plants were carried out using the R package *Phytools* v2.5.2 [101] to deal with polytomies and the R package *ape* v5.8 [102] to deal with unknown character states. Characters encoded in the main model correspond to known host plant families of Smicronychini (Asteraceae, Convolvulaceae, Gentianaceae and Orobanchaceae) associated with their parasitic interactions (non-parasitic, hemiparasitic and parasitic; figure 1; Table S1). Six categories and one unknown state were thus defined as follows: non-parasitic Asteraceae (A), non-parasitic Convolvulaceae (C), parasitic

Convolvulaceae (pC), non-parasitic Gentianaceae (G), parasitic Orobanchaceae (pO) and hemiparasitic Orobanchaceae (hO). Alternative character states combinations were also tested using only host plant families or only parasitic types of host plants. ASR were carried out by comparing the likelihood of three Markov models (Mk), based on three different character state transition matrices, one with equal rates (ER), one with symmetrical rates (SYM) and one with all rates different (ARD). These models were then applied to the *extended dataset* phylogram to avoid biases linked to ultrametric trees [103-105], and reduced to one representative per species to limit ancestral trait estimation biases with the *fitMk* function. AIC values were compared using the *aic.w* function. Based on the best fitting model, each character state was assigned, at each internal node of the phylogeny, either a posterior probability using the *make.simmap* function over 1,000 simulations by a continuous-time reversible Markov process with *Phytools*, or a likelihood (L) directly estimated by the *ace* function with *ape*.

Results

Dataset filtration

AHE dataset - Prior to read assembly, the demultiplexing step highlighted the existence of sequences mapped to unused tag combinations due to their redundancy in target capture pools. These chimeric sequences originate from accidental "tag jumping" events described by [106], which occurred during the target capture protocol. Thus, three specimens with a number of reads falling below an arbitrary threshold of 2 Mpb were excluded from the dataset because their genetic information was considered indistinguishable from that of potential chimeras. For the same reason, two additional specimens were removed from the dataset as 0 ng/µL of DNA was detected following DNA extraction, even if DNA reads have been sequenced afterwards. Two cross-contamination events were also highlighted with MitoFinder and removed from the dataset. Finally, following the phylogenomic pipeline, eight individuals with less than 50% DNA information were filtered out, resulting in a final supermatrix of 52 taxa.

COI dataset - Out of 202 initial COI sequences, 153 were kept following filtration of sequences merged with missing COI from the AHE dataset, misidentifications, pseudogenes, contaminations and too short sequences.

Phylogenomic pipeline

Following dataset filtration, the *extended dataset* consists of 195 specimens, representing 69 different species of Smicronychini. Interestingly, among the 62 individuals we know were sampled in dry museum collections, 40 were included in the final sampling (11 out of 19 in the *AHE dataset* and 29 out of 43 in the *COI dataset*). Specimens selected for the *AHE dataset* then followed our pipeline (figure S1). From 832 AHE reference sequences used for the iterative assembly 230 loci were filtered out mostly because of their low sampling coverage (less than 70% of the individual captured), resulting in 602 selected loci for the final supermatrix. The supermatrix built from the *AHE dataset* comprises 228,248 nucleotides from 52 individuals including outgroups. Interestingly, the elongation of the coding regions increases the coding part of the supermatrix from 50.3% to 61.3% of its total length. Overall, the concatenation of partitions carried out by ModelFinder reduced the number of partitions from 2703 to only 58, which is consistent with the high degree of coding partitions in the sequences obtained with AHE.

Tree inferences

The phylogenomic tree of the *AHE dataset* is shown in figure 2A. It recovers the Smicronychini as a monophyletic tribe with maximum support, their most closely related outgroup is inferred to be from the genus *Pachytychius* (Curculioninae *uncert. sed.*; sometimes included in Smicronychini, e.g. by [107]). All but two nodes of the ingroup are inferred with maximum support (SH-aLRT > 99, UFBS > 98%). The genera *Sharpia* and *Topelatus* are recovered as sister to all remaining Smicronychini, but their monophyly is uncertain considering branch support (SH-aLRT = 5.3, UFBS = 53%). The most speciose genus *Smicronyx* is however found paraphyletic with the genus *Afrosmicronyx* in apical position embedded among *Smicronyx* species.

The phylogenomic tree of the *extended dataset*, consisting of both AHE and COI data, is shown in figure S2 and rerun pruned to one individual per species in figure 3. The addition of the genus *Promecotarsus* supports its monophyly (SH-aLRT = 90.1, UFBS = 95%) and reveals once again the paraphyly of *Smicronyx*, even though their relative position among *Smicronyx* species is poorly supported and variable between runs (SH-aLRT = 54.6, UFBS = 62%; figure S2-3). Consequently, eight major Smicronychini clades can be drawn from this topology: one non-parasitic Convolvulaceae-feeding *Sharpia* species, three independent clades of parasitic *Cuscuta*-feeding *Smicronyx* species, one non-parasitic

Asteraceae-feeding *Smicronyx* species, two independent hemi- and parasitic
Orobanchaceae-feeding clades of *Smicronyx* and *Afrosmicronyx* species and one clade of
non-parasitic Gentianaceae-feeding *Smicronyx* species. However, surely because it mainly
consists of COI grafted species, the Asteraceae feeding clade remains doubtful. Topology
tests conducted between three topologies assuming (i) its paraphyly, (ii) its monophyly or
(iii) its monophyly including *Promecotarsus* species revealed no significant differences in
likelihood but ASR conducted over these alternative topologies differ slightly for this clade
(see supplementary results on Zenodo at *https://doi.org/10.5281/zenodo.15098685*). Thus,
the topology presented in figure 3 is the most parsimonious one of the equally likely
topologies tested. In addition to *Promecotarsus* spp., four species remain difficult to assign
to any of these clades, namely: *Topelatus* sp., *Smicronyx kiboanus*, *Smicronyx zonatus* and
Smicronyx sp. nov. 2.

Figure 2: Phylogenomic tree based on the AHE dataset. A. Maximum likelihood tree of the AHE dataset. Nodes poorly supported (SH-aLRT < 80 or UFBS < 95%) are respectively highlighted with gray triangles and circles. For graphical purposes, some branches were shortened to half of their length (//). Pictures of weevils are linked to corresponding species. **B.** Partitioning scheme of a theoretical AHE loci. Flanking and inserted partitions (f1-3) are highlighted in grey and coding partitions (c1-3) are highlighted in green. The region targeted by the probes (P) is shown in red but does not appear in the final supermatrix. Credits for weevil pictures: J.H..

Ancestral host plant estimations

ASR analyses of host-plant associations were conducted on the *extended dataset* pruned to one sample per species. Because chronograms are commonly used in ASR analyses, we additionally performed ASR on an ultrametric tree as a robustness check to assess whether our inferences were sensitive to tree type. ASR conducted on ultrametric trees or character state combinations restricted to plant families or parasitic types all agree with the main analysis presented in figure 3, that consists in a phylogram with plant families and parasitic types as character states (but see alternative ASR results on Zenodo at *https://doi.org/10.5281/zenodo.15098685*). The best fitting model is estimated to be the equal transition rate matrix (AICw: ER = 0.9998722; SYM = 0.0001278; ARD = 0). Results obtained from *Phytools* and *ape* packages being identical (Pearson's correlation between matrices = 0.9999622), results of *ape* function that take into account unknown character states are presented in figure 3 to show additional species from the *extended dataset*. The most recent common ancestor of Smicronychini has an estimated likelihood of $L_{pC} = 0.51$ in favor of parasitic Convolvulaceae, $L_C = 0.42$ in favor of non-parasitic Convolvulaceae and less than 0.05 for every other character state. The parasitic Convolvulaceae character state is well supported as the ancestral state of all *Smicronyx* spp. *sensu lato* (including *Afrosmicronyx* and *Promecotarsus* regarding present analyses) with an estimated likelihood of $L_{pC} = 0.92$ or $L_{pC} = 0.99$ if we ignore *S. kiboanus*. Consequently, we infer several shifts from parasitic Convolvulaceae to other plant lineages such as non-parasitic Asteraceae ($L_{pC} = 0.96$) and parasitic Orobanchaceae ($L_{pC} = 0.46$, $L_G = 0.25$, $L_{hO} = 0.23$), without implying directional differences in transition rates or a causal driving role under the ER model. However, the ancestral host plant of weevils feeding on non-parasitic Gentianaceae and those feeding on hemiparasitic Orobanchaceae remains uncertain ($L_{pC} = 0.26$, $L_G = 0.38$, $L_{hO} = 0.32$).

376
377
378 **Figure 3: Ancestral host plant reconstruction (ASR) of Smicronychini weevils and their**

379 **current ecologies.** The phylogenetic tree topology corresponds to the extended dataset ML

380 tree pruned to one branch per species. The AHE backbone tree is represented with plain

381 branches while less robust COI data are represented with dashed branches. Reported gall-

382 inducing species are marked with a "x" between branch tips and species names. Pie charts

383 represented on each node of the phylogeny correspond to the estimated likelihood of each

384 character state computed by the ace function. The colors of pie charts and background

385 shading correspond to host plant character states used in the ASR as detailed in the bottom-

386 left table. Colored shading represents the most probable ancestral state estimated at each node

387   and left blank for unknown states and uncertain ancestral states. Photo credits: B.Z. & J.H.

388
389   **Discussion**
390
This study provides the first robust phylogenetic hypothesis for the Smicronychini and
reveals a remarkable pattern in the evolution of their host plant use. Phylogenomic
inferences recover the monophyly of the tribe, but the paraphyly of the most speciose genus
Smicronyx indicates that the current genus-level classification based on morphology is not
satisfactory and requires a detailed re-investigation. That being said, the inferred *Smicronyx*
clades consist of species with consistent morphologies and host plant associations, agreeing
with the expectation of niche conservatism generally observed at a shallow taxonomic level
in weevils (see the review of [70]).

The evolution of host plant preferences inferred herein reveals that the ancestral host
plant of the tribe Smicronychini belongs with a high probability to the Convolvulaceae
family. The most speciose lineage, represented by species belonging to the genus *Smicronyx*
or closely related to it, are ancestrally associated with holoparasitic Convolvulaceae with
great confidence. Interestingly, this lineage has experienced extreme secondary shifts toward
phylogenetically distant botanical families encompassing both parasitic (Orobanchaceae)
and non-parasitic Asterids (Asteraceae and perhaps Gentianaceae, both of which are known
hosts of *Cuscuta* parasites). To our knowledge, this pattern is unique in phytophagous
insects and is consistent with the hypothesis formulated in the introduction, which proposes
that interactions between parasitic plants and their hosts may create ecological contexts
conducive to major host plant shifts. The specific underlying molecular mechanism required
to formally express this hypothesis has not been explored in the context of this study.
However, the case of *S. quadrifer,* performing its development from a dodder to the host
Asteraceae shows that such a transition is physically and physiologically possible.
Based on this, we can formulate putative evolutionary explanations to the general host plant
shift patterns that are observed. It has been shown that host-parasite interactions imply an
intricate spatial proximity of tissues and allow exchanges of primary and secondary
compounds between plants [108-109]. As such, the Smicronychini developing as larvae in
the stems of parasitic plants are directly exposed to a physicochemical environment close to
that of a dodder's host. This shared environment can be seen as the ecological opportunity
that led weevils feeding on dodders to evolve adaptations to their parasitic plant's host, even
though Asteraceae and Convolvulaceae have few secondary compounds in common (for

Asteraceae see [110] ; for Convolvulaceae see [111]). Formulating hypotheses regarding
shifts to other non parasitic plant lineages is more difficult, as it is unclear whether
Gentianaceae feeding weevils shifted from parasitic Convolvulaceae or hemiparasitic
Orobanchaceae.

Another intriguing host plant shift pattern in Smicronychini is the shift inferred
between phylogenetically distant holoparasitic plant lineages, namely from Convolvulaceae
to Orobanchaceae. This pattern suggests that a convergent parasitic lifestyle between
phylogenetically distant lineages of plants could have promoted the shifts. One feature of
parasitic plants that could explain this pattern is the existence of similar chemical
compounds between dodders and broomrapes [112], like those involved in haustorium
formation (e.g. [113-115]) that potentially facilitated the detection of these distant plants by
one or a few Smicronychini lineages towards holoparasitic Orobanchaceae (although their
detectability by Smicronychini have never been put to the test). Additionally, a second
feature of parasitic plants is that they may share similar host plants and therefore show some
degree of chemical similarity according to the proximity of their host plants. However, the
host plant repertoire of Orobanchaceae-feeding weevils is not yet known with sufficient
precision to corroborate this second hypothesis.

Phytophagous insect diversity has been widely associated with major shifts between
distant plant lineages leading to bursts of diversification [19,22]. Yet, few papers have
questioned how these major shifts occur in the first place. Although the patterns inferred in
this study are specific to plant-plant parasitic interactions, the underlying scenario proposed
here might provide valuable insights on the processes leading to major host plant shifts in
phytophagous insects. From the insect perspective, plants with overlapping niches, for
example with similar physiology, structure or chemistry, could provide ecological
opportunities at the basis of host shifting events. In other words, overlapping plant niches
can make up "facilitating bridges" that can be crossed by insects toward new potential hosts,
as long as they are spatially and temporally consistent. Insects that shift from parasitic plants
to their hosts make these opportunities more obvious in all of its multifaceted parts. It is
likely that a large share of historical shifts has been driven by these kinds of opportunities,
but the scarcity of ancestral data, intermediate states and our limited ability to link
ecological niches in their multifaceted aspects to host plant shifts make it difficult to
conceptualize. In this context, "minor" shifts that typically occur within a plant lineage -
generally described as niche conservatism - may also derive from such ecological

opportunities provided by the general relatedness (chemical, physiological, etc) between
species in a given lineage. The decorrelation between phylogeny and niche relatedness
inferred in the host repertoire of Smicronychini emphasizes the preeminence of ecological
factors as prior components driving host shifts in phytophagous insects [54,116]. Similarly,
many events of independent host plant shifts can be proposed as a result of ecological niche
similarities at a broader scale. For example, the structural and physiological similarities in
flower structures were suggested to be at the origin of major host shifts from monocots to
dicots in derelomine flower weevils [117]. Another example is that of nematine sawflies
(Hymenoptera: Tenthredinidae), where major host shifts occurred between phylogenetically
distant, but ecologically similar hosts [118]. Altogether, such examples are consistent with a
unified view of host shift mechanisms based on ecological opportunities arising from
overlapping niches provided by plants.

Ethics statement

The following services are acknowledged for providing permits: Western Cape Province
(Western Cape Nature Conservation Board [permit No. CN44-30-4229], the Cape Research
Centre [Republic of South African National Parks, CRC/2019-2020/012--2012/V1]),
KwaZulu-Natal Province (Ezemvelo KZN Wildlife permits office, Collecting Permit KZN:
OP1382-2019), the Direction de la faune de la chasse des Parcs et des Réserves (Niger:
permit No. 1778177) and the Ministry of environment and sustainable development
(Senegal: Authorization No. 231). Museum samples accession numbers can be found as part
of supplementary material.

Data accessibility statement

The dataset supporting this article is openly available in Zenodo at
https://doi.org/10.5281/zenodo.15098685 [119]. Raw target capture data and assembled
AHE results are publicly available in GenBank (NCBI) under the accession PRJNA1244829:
https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA1244829. COI sequences generated
for the present study are publicly available on BOLD systems under the project name
SCRYX and are accessible on Zenodo as well.

Authors' contributions

R.A., J.H., G.J.K. and B.Z. designed the study; J.H. collected data; L.B. and B.Z. conducted the experiments; B.Z. wrote the pipeline and performed all molecular analyses, supervised by R.A. and G.J.K.; B.Z. wrote the first draft; all authors substantially contributed to the writing of the final draft.

Conflicts of interest

We declare we have no competing interests.

AI declaration

We have not used AI-assisted technologies in creating this article.

Fundings

This project was supported by the ANR Project ANR-21-CE32-0006 AgriBioDiv, coordinated by Christine Meynard.

Acknowledgements

The following colleagues are acknowledged for providing some of the specimens used in this study: Robert Anderson (Canadian Museum of Nature, Gatineau, Canada), Patrice Bouchard (Ottawa Research and Development Centre, Ottawa, Canada), Lourdes Chamorro (USDA, Smithsonian Institution - National Museum of Natural History, Washington DC, USA), Desmond Edward Conlong (Stellenbosch University, South Africa), Madougou Garba (Direction Générale de la Protection des Végétaux, Ministère de l'Agriculture, Niamey, Niger), Hiroaki Kojima (Laboratory of Entomology, Tokyo University of Agriculture, Japan), Jean Pelletier (Monnaie, France) and Patrick Weil (Maison de la Nature et de l'Environnement, Pau, France). The analyses benefited from the GenoToul Bioinformatics platform (Toulouse, France) platform services. We also thank Christine Meynard for funding this project. This is contribution ISEM 2026-023 of the Institut des Sciences de l'Evolution de Montpellier.

References

1. Giron D et al. 2018 Promises and challenges in insect–plant interactions. Entomologia Exp
Applicata 166, 319–343. (doi:10.1111/eea.12679)

2. Braga MP, Janz N. 2021 Host repertoires and changing insect–plant interactions.
Ecological Entomology 46, 1241–1253. (doi:10.1111/een.13073)

3. Hutchinson GE. 1957 Concluding Remarks. Cold Spring Harbor Symposia on Quantitative
Biology 22, 415–427. (doi:10.1101/SQB.1957.022.01.039)

4. Holt RD. 2009 Bringing the Hutchinsonian niche into the 21st century: Ecological and evolutionary
perspectives. Proc. Natl. Acad. Sci. U.S.A. 106, 19659–19665.
(doi:10.1073/pnas.0905137106)

5. Peterson AT, Soberón J, Sánchez-Cordero V. 1999 Conservatism of ecological niches in
evolutionary time. Science 285, 1265–1267. (doi:10.1126/science.285.5431.1265)

6. Futuyma DJ, Moreno G. 1988 The evolution of ecological specialization. Annu. Rev. Ecol. Syst. 19,
207–233. (doi:10.1146/annurev.es.19.110188.001231)

7. Jaenike J. 1990 Host specialization in phytophagous insects. Annu. Rev. Ecol. Syst. 21, 243–
273. (doi:10.1146/annurev.es.21.110190.001331)

8. Janz N, Nylin S, Wahlberg N. 2006 Diversity begets diversity: host expansions and the
diversification of plant-feeding insects. BMC Evol Biol 6, 4. (doi:10.1186/1471-2148-6-4)

9. Holloway JD, Hebert PDN. 1979 Ecological and taxonomic trends in macrolepidopteran host
plant selection. Biological Journal of the Linnean Society 11, 229–251. (doi:10.1111/j.1095-
8312.1979.tb00036.x)

10. Merrill RM, Naisbit RE, Mallet J, Jiggins CD. 2013 Ecological and genetic factors influencing the
transition between host-use strategies in sympatric *Heliconius* butterflies. J of
Evolutionary Biology 26, 1959–1967. (doi:10.1111/jeb.12194)

11. Singh KS et al. 2020 The genetic architecture of a host shift: An adaptive walk protected an
aphid and its endosymbiont from plant chemical defenses. Sci. Adv. 6, eaba1070.
(doi:10.1126/sciadv.aba1070)

12. Berenbaum MR, Favret C, Schuler MA. 1996 On defining 'key innovations' in an adaptive
radiation: cytochrome P450S and Papilionidae. The American Naturalist 148, S139–
S155. (doi:10.1086/285907)

13. Gatehouse JA. 2002 Plant resistance towards insect herbivores: a dynamic interaction.
New Phytologist 156, 145–169. (doi:10.1046/j.1469-8137.2002.00519.x)

14. Connor EF. 2006 Effects of the light environment on oviposition preference and survival of a leaf-
mining moth, *Cameraria hamadryadella* (Lepidoptera: Gracillariidae), on *Quercus alba* L.
Ecological Entomology 31, 179–184. (doi:10.1111/j.0307-6946.2006.00779.x)

15. Yarnes CT, Boecklen WJ, Salminen J-P. 2008 No simple sum: seasonal variation in
tannin phenotypes and leaf-miners in hybrid oaks. Chemoecology 18, 39–51.
(doi:10.1007/s00049-007-0391-y)

16. Mithöfer A, Boland W. 2012 Plant defense against herbivores: chemical aspects. Annu. Rev.
Plant Biol. 63, 431–450. (doi:10.1146/annurev-arplant-042110-103854)

17. Heidel-Fischer HM, Vogel H. 2015 Molecular mechanisms of insect adaptation to plant
secondary compounds. Current Opinion in Insect Science 8, 8–14.
(doi:10.1016/j.cois.2015.02.004)

18. Brand P, Hinojosa-Díaz IA, Ayala R, Daigle M, Yurrita Obiols CL, Eltz T, Ramírez SR. 2020 The
evolution of sexual signaling is linked to odorant receptor tuning in perfume-collecting
orchid bees. Nat Commun 11, 244. (doi:10.1038/s41467-019-14162-6)

19. Janz N. 2011 Ehrlich and Raven revisited: mechanisms underlying codiversification of plants
and enemies. Annu. Rev. Ecol. Evol. Syst. 42, 71–89. (doi:10.1146/annurev-ecolsys-102710-
145024)

20. Kergoat GJ, Meseguer AS, Jousselin E. 2017 Evolution of plant–insect interactions. In Advances
in Botanical Research, pp. 25–53. Elsevier. (doi:10.1016/bs.abr.2016.09.005)

21. Jousselin E, Elias M. 2019 Testing host-plant driven speciation in phytophagous insects :
a phylogenetic perspective. Peer Community in Evolutionary Biology
(doi:10.20944/preprints201902.0215v2)

22. Ehrlich PR, Raven PH. 1964 Butterflies and plants: a study in coevolution. Evolution 18, 586–
608. (doi:10.1111/j.1558-5646.1964.tb01674.x)

23. Janz N, Nylin S. 2008 The oscillation hypothesis of host-plant range and speciation. In
Specialization, Speciation, and Radiation: The Evolutionary Biology of Herbivorous
Insects, University of California Press. (doi:10.1525/california/9780520251328.003.0015)

24. Hardy NB, Otto SP. 2014 Specialization and generalization in the diversification of phytophagous
insects: tests of the musical chairs and oscillation hypotheses. Proc. R. Soc. B. 281,
20132960. (doi:10.1098/rspb.2013.2960)

25. Nylin S, Slove J, Janz N. 2014 Host plant utilization, host range oscillations and diversification
in nymphalid butterflies: a phylogenetic investigation. Evolution 68, 105–124.
(doi:10.1111/evo.12227)

26. Edger PP et al. 2015 The butterfly plant arms-race escalated by gene and genome duplications.
Proc. Natl. Acad. Sci. U.S.A. 112, 8362–8366. (doi:10.1073/pnas.1503926112)

27. Wang H, Holloway JD, Janz N, Braga MP, Wahlberg N, Wang M, Nylin S. 2017 Polyphagy
and diversification in tussock moths: Support for the oscillation hypothesis from
extreme generalists. Ecology and Evolution 7, 7975–7986. (doi:10.1002/ece3.3350)

28. Allio R et al. 2021 Genome-wide macroevolutionary signatures of key innovations in
butterflies colonizing new host plants. Nat Commun 12, 354. (doi:10.1038/s41467-020-
20507-3)

29. Nyman T, Vikberg V, Smith DR, Boevé J-L. 2010 How common is ecological speciation
in plant-feeding insects? A 'Higher' Nematinae perspective. BMC Evol Biol 10,
266. (doi:10.1186/1471-2148-10-266)

30. Imada Y, Kawakita A, Kato M. 2011 Allopatric distribution and diversification without niche shift
in a bryophyte-feeding basal moth lineage (Lepidoptera: Micropterigidae). Proc. R. Soc. B.
278, 3026–3033. (doi:10.1098/rspb.2011.0134)

31. Althoff DM, Segraves KA, Johnson MTJ. 2014 Testing for coevolutionary diversification:
linking pattern with process. Trends in Ecology & Evolution 29, 82–89.
(doi:10.1016/j.tree.2013.11.003)

32. Forbes AA, Devine SN, Hippee AC, Tvedte ES, Ward AKG, Widmayer HA, Wilson CJ. 2017
Revisiting the particular role of host shifts in initiating insect speciation. Evolution 71, 1126–
1137. (doi:10.1111/evo.13164)

33. Smadja CM, Canbäck B, Vitalis R, Gautier M, Ferrari J, Zhou J-J, Butlin RK. 2012 Large-scale
candidate gene scan reveals the role of chemoreceptor genes in host plant specialization
and speciation in the pea aphid: large-scale candidate gene scan and speciation. Evolution
66, 2723–2738. (doi:10.1111/j.1558-5646.2012.01612.x)

34. Simon J-C, d'Alencon E, Guy E, Jacquin-Joly E, Jaquiery J, Nouhaud P, Peccoud J, Sugio A, Streiff
R. 2015 Genomics of adaptation to host-plants in herbivorous insects. Briefings in
Functional Genomics 14, 413–423. (doi:10.1093/bfgp/elv015)

35. Vertacnik KL, Linnen CR. 2017 Evolutionary genetics of host shifts in herbivorous insects:
insights from the age of genomics. Annals of the New York Academy of Sciences 1389, 186–
212. (doi:10.1111/nyas.13311)

36. Hou Z, Shi F, Ge S, Tao J, Ren L, Wu H, Zong S. 2021 Comparative transcriptome analysis of the
newly discovered insect vector of the pine wood nematode in China, revealing putative
genes related to host plant adaptation. BMC Genomics 22, 189. (doi:10.1186/s12864-021-
07498-1)

37. Celorio-Mancera MDLP, Steward RA, Pruisscher P, Smialowska A, Pires Braga M, Janz N,
Wheat CW, Nylin S. 2023 Larval transcriptomes reflect the evolutionary history of plant–
insect associations. Evolution 77, 519–533. (doi:10.1093/evolut/qpac049)

38. Savković U, ĐorĐević M, Šešlija Jovanović D, Lazarević J, Tucić N, Stojković B. 2016

613   Experimentally induced host-shift changes life-history strategy in a seed beetle. J
614   of Evolutionary Biology 29, 837–847. (doi:10.1111/jeb.12831)

615 39. Agosta SJ. 2006 On ecological fitting, plant–insect associations, herbivore host shifts, and
616   host plant selection. Oikos 114, 556–565. (doi:10.1111/j.2006.0030-1299.15025.x)

617 40. Graves SD, Shapiro AM. 2003 Exotics as host plants of the California butterfly fauna.
618   Biological Conservation 110, 413–433. (doi:10.1016/S0006-3207(02)00233-1)

619 41. Singer MS. 2008 Evolutionary ecology of polyphagy. In Specialization, speciation, and
620   radiation: The evolutionary biology of herbivorous insects, University of California Press.
621   (doi:10.1525/california/9780520251328.003.0003)

622 42. Gougherty AV, Davies TJ. 2021 Towards a phylogenetic ecology of plant pests and pathogens.
623   Phil. Trans. R. Soc. B 376, 20200359. (doi:10.1098/rstb.2020.0359)

624 43. Braga MP. 2023 Are exotic host plants a life raft or a trap for butterflies? Current Opinion in
625   Insect Science 58, 101074. (doi:10.1016/j.cois.2023.101074)

626 44. Wood TK, Olmstead KL, Guttman SI. 1990 Insect phenology mediated by host-plant
627   water relations. Evolution 44, 629–636. (doi:10.1111/j.1558-5646.1990.tb05943.x)

628 45. Whipple AV, Abrahamson WG, Khamiss MA, Heinrich PL, Urian AG, Northridge EM. 2009. Host-
629  race formation: promoted by phenology, constrained by heritability. J of Evolutionary Biology 22,
630  793–804. (doi:10.1111/j.1420-9101.2009.01690.x)

631 46. Center TD, Johnson CD. 1974 Coevolution of some seed beetles (Coleoptera: Bruchidae) and
632   their hosts. Ecology 55, 1096–1103. (doi:10.2307/1940359)

633 47. Jaime R, Rey PJ, Alcántara JM, Bastida JM. 2013 Glandular trichomes as an inflorescence
634   defence mechanism against insect herbivores in Iberian columbines. Oecologia 172, 1051–
635   1060. (doi:10.1007/s00442-012-2553-z)

636 48. Li N, Li S, Ge J, Schuman MC, Wei J, Ma R. 2017 Manipulating two olfactory cues causes
637   a biological control beetle to shift to non-target plant species. Journal of Ecology
638   105, 1534–1546. (doi:10.1111/1365-2745.12778)

639 49. Groen SC, Whiteman NK. 2021 Convergent evolution of cardiac-glycoside resistance in
640   predators and parasites of milkweed herbivores. Current Biology 31, R1465–R1466.
641   (doi:10.1016/j.cub.2021.10.025)

642 50. Farrell BD, Mitter C. 1998 The timing of insect/plant diversification: might *Tetraopes*
643   (Coleoptera: Cerambycidae) and *Asclepias* (Asclepiadaceae) have co-evolved? Biol. J. Linn.
644   Soc. 63,
645   553–557.

646 51. Becerra JX. 1997 Insects on plants: macroevolutionary chemical trends in host use. Science
647   276, 253–256. (doi:10.1126/science.276.5310.253)

648 52. Kergoat GJ, Delobel A, Fédière G, Rü BL, Silvain J-F. 2005 Both host-plant phylogeny and
649   chemistry have shaped the African seed-beetle radiation. Molecular Phylogenetics and
650   Evolution 35, 602–611. (doi:10.1016/j.ympev.2004.12.024)

651 53. Van Der Linden CFH, WallisDeVries MF, Simon S. 2021 Great chemistry between us: The
652   link between plant chemical defenses and butterfly evolution. Ecology and Evolution
653   11, 8595–8613. (doi:10.1002/ece3.7673)

654 54. Brooks DR, McLennan DA. 2002 The nature of diversity: an evolutionary voyage of
655   discovery. University of Chicago Press. See http://www.bibliovault.org/BV.landing.epl?
656   ISBN=9780226075907.

657 55. Celorio-Mancera MDLP, Wheat CW, Huss M, Vezzi F, Neethiraj R, Reimegård J, Nylin S, Janz N.
658   2016 Evolutionary history of host use, rather than plant phylogeny, determines gene
659   expression in a generalist butterfly. BMC Evol Biol 16, 59. (doi:10.1186/s12862-016-0627-
660   y)

661 56. Messina FJ, Jones JC. 2009 Does rapid adaptation to a poor-quality host by *Callosobruchus*
662   *maculatus* (F.) cause cross-adaptation to other legume hosts? Journal of Stored
663   Products Research 45, 215–219. (doi:10.1016/j.jspr.2009.02.004)

664 57. Yoshida S, Cui S, Ichihashi Y, Shirasu K. 2016 The haustorium, a specialized invasive organ

665   in parasitic plants. Annu. Rev. Plant Biol. 67, 643–667. (doi:10.1146/annurev-arplant-
666   043015-111702)

58. Adler L. 2002 Host effects on herbivory and pollination in a hemiparasitic plant.

59. Rowntree JK, Fisher Barham D, Stewart AJA, Hartley SE. 2014 The effect of multiple host species on a keystone parasitic plant and its aphid herbivores. Functional Ecology 28, 829–836. (doi:10.1111/1365-2435.12281)

60. Anderson DM. 1962 The weevil genus *Smicronyx* in America North of Mexico (Coleoptera: Curculionidae). Proceedings of the United States National Museum 113, 185–372. (doi:10.5479/si.00963801.113-3456.185)

61. Anderson DM. 1970 Dodder weevils in simultaneous association with parasitic plants and their hosts. Science 168, 132–133. (doi:10.1126/science.168.3927.132)

62. Anderson RS. 1993 Weevils and plants: phylogenetic versus ecological mediation of evolution of host plant associations in Curculioninae (Coleoptera: Curculionidae). Mem. Entomol. Soc. Can. 125, 197–232. (doi:10.4039/entm125165197-1)

63. Caldara R, Franz NM, Oberprieler RG. 2014 Curculioninae Latreille, 1802. In Morphology and Systematics, pp. 589–628. Germany: de Gruyter.

64. Haran J, Schütte A, Friedman A-L-L. 2017 A review of *Smicronyx* Schoenherr (Coleoptera, Curculionidae) of Israel, with description of two new species. Zootaxa 4237, 17. (doi:10.11646/zootaxa.4237.1.2)

65. Haran J. 2018 A review of the genus *Smicronyx* Schoenherr (Coleoptera, Curculionidae, Curculioninae) in tropical Africa. Zootaxa 4508. (doi:10.11646/zootaxa.4508.2.9)

66. Haran JM. 2021 The Smicronychini of southern Africa (Coleoptera, Curculionidae): Review of the tribe and description of 12 new species. EJT 735. (doi:10.5852/ejt.2021.735.1239)

67. Haran J, Perrin H. 2017 Revision of the genus *Afrosmicronyx* Hustache (Coleoptera, Curculionidae, Curculioninae). Zootaxa 4365, 132. (doi:10.11646/zootaxa.4365.2.2)

68. Irving LJ, Cameron DD. 2009 Chapter 3 You are what you eat: Interactions between root parasitic plants and their hosts. pp. 87–138. Academic Press. (doi:https://doi.org/10.1016/S0065-2296(08)00803-3)

69. The Angiosperm Phylogeny Group. 2016 An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. Bot. J. Linn. Soc. 181, 1–20. (doi:10.1111/boj.12385)

70. Oberprieler RG, Marvaldi AE, Anderson RS. 2007 Weevils, weevils, weevils everywhere*. Zootaxa 1668, 491–520. (doi:10.11646/zootaxa.1668.1.24)

71. Hernández-Vera G, Toševski I, Caldara R, Emerson BC. 2019 Evolution of host plant use and diversification in a species complex of parasitic weevils (Coleoptera: Curculionidae). PeerJ 7, e6625. (doi:10.7717/peerj.6625)

72. Haran J et al. 2023 Phylogenomics illuminates the phylogeny of flower weevils (Curculioninae) and reveals ten independent origins of brood-site pollination mutualism in true weevils. Proc. R. Soc. B. 290, 20230889. (doi:10.1098/rspb.2023.0889)

73. Allio R, Schomaker-Bastos A, Romiguier J, Prosdocimi F, Nabholz B, Delsuc F. 2020 MitoFinder: Efficient automated large-scale extraction of mitogenomic data in target enrichment phylogenomics. Molecular Ecology Resources 20, 892–905. (doi:10.1111/1755-0998.13160)

74. Ratnasingham S, Hebert PDN. 2007 BOLD: The Barcode of Life Data system (http://www.barcodinglife.org). Molecular Ecology Notes 7, 355–364. (doi:10.1111/j.1471-8286.2007.01678.x)

75. Faircloth BC, Branstetter MG, White ND, Brady SG. 2015 Target enrichment of ultraconserved elements from arthropods provides a genomic perspective on relationships among Hymenoptera. Mol Ecol Resour 15, 489–501. (doi:10.1111/1755-0998.12328)

76. Haddad S, Shin S, Lemmon AR, Lemmon EM, Svacha P, Farrell B, Ślipiński A, Windsor D, Mckenna
DD. 2018 Anchored hybrid enrichment provides new insights into the phylogeny and evolution of
longhorned beetles (Cerambycidae). Syst Entomol 43, 68–89. (doi:10.1111/syen.12257)
77. Shin S et al. 2018 Phylogenomic data yield new and robust insights into the phylogeny
and evolution of weevils. Molecular Biology and Evolution 35, 823–836.
(doi:10.1093/molbev/msx324)
78. Altschul S. 1997 Gapped BLAST and PSI-BLAST: a new generation of protein database
search programs. Nucleic Acids Research 25, 3389–3402.
(doi:10.1093/nar/25.17.3389)
79. Martin M. 2011 Cutadapt removes adapter sequences from high-throughput sequencing
reads. EMBnet j. 17, 10. (doi:10.14806/ej.17.1.200)
80. Bolger AM, Lohse M, Usadel B. 2014 Trimmomatic: a flexible trimmer for Illumina sequence
data. Bioinformatics 30, 2114–2120. (doi:10.1093/bioinformatics/btu170)
81. Breinholt JW, Earl C, Lemmon AR, Lemmon EM, Xiao L, Kawahara AY. 2018 Resolving
relationships among the megadiverse butterflies and moths with a novel pipeline for
anchored phylogenomics. Systematic Biology 67, 78–93. (doi:10.1093/sysbio/syx048)
82. Katoh K, Frith MC. 2012 Adding unaligned sequences into an existing alignment using MAFFT
and LAST. Bioinformatics 28, 3144–3146. (doi:10.1093/bioinformatics/bts578)
83. Katoh K, Standley DM. 2013 MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in performance and usability. Molecular Biology and Evolution 30, 772–
780. (doi:10.1093/molbev/mst010)
84. Xi Z, Liu L, Davis CC. 2016 The impact of missing data on species tree estimation. Mol Biol Evol
33, 838–860. (doi:10.1093/molbev/msv266)
85. Borowiec ML. 2016 AMAS: a fast tool for alignment manipulation and computing of summary
statistics. PeerJ 4, e1660. (doi:10.7717/peerj.1660)
86. Ranwez V, Chantret N, Delsuc F. 2021 Aligning protein-coding nucleotide sequences with
MACSE. In Multiple Sequence Alignment (ed K Katoh), pp. 51–70. New York, NY: Springer
US.
87. Di Franco A, Poujol R, Baurain D, Philippe H. 2019 Evaluating the usefulness of alignment
filtering methods to reduce the impact of errors on evolutionary inferences. BMC Evol Biol
19, 21. (doi:10.1186/s12862-019-1350-2)
88. Miller MA, Pfeiffer W, Schwartz T. 2010 Creating the CIPRES Science Gateway for inference of
large phylogenetic trees. In 2010 Gateway Computing Environments Workshop (GCE), pp. 1–
8. New Orleans, LA, USA: IEEE. (doi:10.1109/GCE.2010.5676129)
89. Nguyen L-T, Schmidt HA, Von Haeseler A, Minh BQ. 2015 IQ-TREE: A fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. Molecular Biology
and Evolution 32, 268–274. (doi:10.1093/molbev/msu300)
90. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Von Haeseler A, Lanfear
R. 2020 IQ-TREE 2: New models and efficient methods for phylogenetic inference in the
genomic era. Molecular Biology and Evolution 37, 1530–1534.
(doi:10.1093/molbev/msaa015)
91. Chernomor O, Von Haeseler A, Minh BQ. 2016 Terrace aware data structure for
phylogenomic inference from supermatrices. Syst Biol 65, 997–1008.
(doi:10.1093/sysbio/syw037)
92. Kalyaanamoorthy S, Minh BQ, Wong TKF, Von Haeseler A, Jermiin LS. 2017 ModelFinder:
fast model selection for accurate phylogenetic estimates. Nat Methods 14, 587–589.
(doi:10.1038/nmeth.4285)
93. Lanfear R, Frandsen PB, Wright AM, Senfeld T, Calcott B. 2017 PartitionFinder 2: New Methods
for Selecting Partitioned Models of Evolution for Molecular and Morphological Phylogenetic
Analyses. Mol Biol Evol 34, 772–773. (doi:10.1093/molbev/msw260)
94. Naser-Khdour S, Minh BQ, Zhang W, Stone EA, Lanfear R. 2019 The prevalence and impact of
model violations in phylogenetic analysis. Genome Biology and Evolution 11, 3341–

768   3352. (doi:10.1093/gbe/evz193)

769 95. Hoang DT, Chernomor O, Von Haeseler A, Minh BQ, Vinh LS. 2018 UFBoot2: Improving
770   the Ultrafast Bootstrap Approximation. Molecular Biology and Evolution 35, 518–
771   522. (doi:10.1093/molbev/msx281)

772 96. Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010 New algorithms and
773   methods to estimate maximum-likelihood phylogenies: assessing the performance of
774   PhyML 3.0. Systematic Biology 59, 307–321. (doi:10.1093/sysbio/syq010)

775 97. Talavera G, Lukhtanov V, Pierce NE, Vila R. 2022 DNA barcodes combined with multilocus data
776   of representative taxa can generate reliable higher-level phylogenies. Systematic Biology
777   71, 382–395. (doi:10.1093/sysbio/syab038)

778 98. Upham NS, Esselstyn JA, Jetz W. 2019 Inferring the mammal tree: Species-level sets of
779   phylogenies for questions in ecology, evolution, and conservation. PLoS Biol 17,
780   e3000494. (doi:10.1371/journal.pbio.3000494)

781 99. Azevedo GHF, Bougie T, Carboni M, Hedin M, Ramírez MJ. 2022 Combining genomic, phenotypic
782   and Sanger sequencing data to elucidate the phylogeny of the two-clawed spiders
783   (*Dionycha*). Molecular Phylogenetics and Evolution 166, 107327.
784   (doi:10.1016/j.ympev.2021.107327)

785 100. Espeland M et al. 2023 Combining target enrichment and Sanger sequencing data to clarify
786   the systematics of the diverse Neotropical butterfly subtribe Euptychiina (Nymphalidae,
787   Satyrinae). Systematic Entomology , syen.12590. (doi:10.1111/syen.12590)

788 101. Revell LJ. 2012 phytools: an R package for phylogenetic comparative biology (and other
789   things). Methods in Ecology and Evolution 3, 217–223. (doi:10.1111/j.2041-
790   210X.2011.00169.x)

791 102. Paradis E, Schliep K. 2019 ape 5.0: an environment for modern phylogenetics and
792   evolutionary analyses in R. Bioinformatics 35, 526–528.
793   (doi:10.1093/bioinformatics/bty633)

794 103. Cusimano N, Renner SS. 2014 Ultrametric trees or phylograms for ancestral state
795   reconstruction: Does it matter? TAXON 63, 721–726. (doi:10.12705/634.14)

796 104. Cascini M, Mitchell KJ, Cooper A, Phillips MJ. 2019 Reconstructing the evolution of giant extinct
797   kangaroos: comparing the utility of DNA, morphology, and total evidence. Systematic
798   Biology 68, 520–537. (doi:10.1093/sysbio/syy080)

799 105. Wilson JD, Mongiardino Koch N, Ramírez MJ. 2022 Chronogram or phylogram for ancestral
800   state estimation? Model-fit statistics indicate the branch lengths underlying a binary
801   character's evolution. Methods Ecol Evol 13, 1679–1689. (doi:10.1111/2041-210X.13872)

802 106. Schnell IB, Bohmann K, Gilbert MTP. 2015 Tag jumps illuminated - reducing sequence-to-
803   sample misidentifications in metabarcoding studies. Mol Ecol Resour 15, 1289–1303.
804   (doi:10.1111/1755-0998.12402)

805 107. Caldara R, Skuhrovec J, Gosik R, Marvaldi AE. 2018 On the affinities and systematic position of
806   *Pachytychius* Jekel, a genus currently incertae sedis in Coleoptera, Curculionidae: Evidence
807   from immature stages and the COI gene support its placement in Smicronychini.
808   Zoologischer Anzeiger 277, 218–230. (doi:10.1016/j.jcz.2018.08.007)

809 108. Smith JD, Woldemariam MG, Mescher MC, Jander G, De Moraes CM. 2016 Glucosinolates
810   from host plants influence growth of the parasitic plant Cuscuta gronovii and its
811   susceptibility to aphid feeding. Plant Physiol. 172, 181–197. (doi:10.1104/pp.16.00613)

812 109. Kokla A, Melnyk CW. 2018 Developing a thief: haustoria formation in parasitic
813   plants. Developmental Biology 442, 53–59. (doi:10.1016/j.ydbio.2018.06.013)

814 110. Heywood VH, Harborne JB, Turner BL, Linnean Society of London, Phytochemical Society,
815   editors. 1977 The biology and chemistry of the Compositae. London ; New York: Academic
816   Press.

817 111. Eich E. 2008 Solanaceae and Convolvulaceae: secondary metabolites. Berlin,
818   Heidelberg: Springer Berlin Heidelberg. (doi:10.1007/978-3-540-74541-9)

112. Rehker J, Lachnit M, Kaldenhoff R. 2012 Molecular convergence of the parasitic plant species

Cuscuta reflexa and *Phelipanche aegyptiaca*. Planta 236, 557–566. (doi:10.1007/s00425-012-
1626-x)

113. Sun G et al. 2018 Large-scale gene losses underlie the genome evolution of parasitic plant
Cuscuta australis. Nat Commun 9, 2683. (doi:10.1038/s41467-018-04721-8)

114. Qiu S, Bradley JM, Zhang P, Chaudhuri R, Blaxter M, Butlin RK, Scholes JD. 2022 Genome-
enabled discovery of candidate virulence loci in *Striga hermonthica*, a devastating parasite of
African cereal crops. New Phytologist 236, 622–638. (doi:10.1111/nph.18305)

115. Westwood JH. 2022 Cracking open the witch's spell book: the witchweed genome provides
clues to plant parasitism. New Phytologist 236, 316–318. (doi:10.1111/nph.18398)

116. Nyman T. 2010 To speciate, or not to speciate? Resource heterogeneity, the subjectivity of
similarity, and the macroevolutionary consequences of niche-width shifts in plant-
feeding insects. Biological Reviews 85, 393–411. (doi:10.1111/j.1469-185X.2009.00109.x)

117. Haran J, Procheş Ş, Benoit L, Kergoat GJ. 2022 From monocots to dicots: host shifts in
Afrotropical derelomine weevils shed light on the evolution of non-obligatory
brood pollination mutualism. Biological Journal of the Linnean Society 137, 15–29.
(doi:10.1093/biolinnean/blac069)

118. Nyman T, Farrell BD, Zinovjev AG, Vikberg V. 2006 Larval habits, host-plant associations,
and speciation in nematine sawflies (Hymenoptera: Tenthredinidae). Evol 60, 1622.
(doi:10.1554/05-674.1)

119. Zelvelder B, Kergoat GJ, Benoit L, Tsuchida T, Haran J, Allio R. 2026 Can ecological interactions
drive evolutionary outcomes? Evidence from insect host shifts between parasitic and non-
parasitic plants. *Zenodo*. (doi:10.5281/zenodo.15098685)