Atypical epigenetic and small RNA control of transposons in clonally reproducing *Spirodela polyrhiza.*

Rodolphe Dombey[1,2], Verónica Barragán-Borrero[1], Daniel Buendía-Ávila[1,2], Arturo Ponce-Mañe[1], José Mario Vargas-Guerrero[1,3], Rana Elias[1] and Arturo Marí-Ordóñez[1].*

[1] Gregor Mendel Institute of Molecular Plant Biology (GMI) of the Austrian Academy of Sciences, Vienna, 1030, Austria.

[2] Vienna BioCenter PhD Program, Doctoral School of the University of Vienna and Medical University of Vienna, Austria.

[3] Current address: Biology School, Costa Rica University, San José, 11501-2060, Costa Rica.

* The author responsible for correspondence and distribution of materials integral to the findings presented in this article is: Arturo Marí-Ordóñez (arturo.mari-ordonez@gmi.oeaw.ac.at)

Abstract

Background: A handful of model plants have provided insight into silencing of transposable elements (TEs) through RNA-directed DNA methylation (RdDM). Guided by 24-nt long small-interfering RNAs (siRNAs), this epigenetic regulation installs DNA methylation and histone modifications like H3K9me2, which can be subsequently maintained independently of siRNAs. However, the genome of the clonally propagating duckweed *Spirodela polyrhiza* (*Lemnaceae*) has low levels of DNA methylation, very low expression of RdDM components, and near absence of 24-nt siRNAs. Moreover, some genes encoding RdDM factors, DNA methylation maintenance, and RNA silencing mechanisms are missing from the genome.

Results: We investigated the distribution of TEs and their epigenetic marks in the Spirodela genome. While abundant degenerated TEs have largely lost DNA methylation and H3K9me2 is low, they remain transcriptionally silenced and are marked by H3K9me1. By contrast, we found high levels of DNA methylation and H3K9me2 in the relatively few intact TEs which are source of 24-nt siRNAs like RdDM-controlled TEs in other angiosperms. Some intact TEs are also regulated by post-transcriptional gene silencing (PTGS), producing 22-nt siRNAs despite the absence of a DCL2, required for their biogenesis in other angiosperms.

Conclusions: The data suggest that, potentially as adaptation to vegetative propagation, RdDM extent, components, and targets are different from other angiosperms, preferentially focused on potentially intact TEs. It also provides evidence for heterochromatin maintenance independently of DNA methylation. These discoveries highlight the diversity of silencing mechanisms that exist in plants and the importance of using disparate model species to discover these mechanisms.

Keywords

 epigenetics, transposons, DNA methylation, small RNAs, heterochromatin, angiosperms, clonal reproduction, duckweeds.

Introduction

Transposable elements (TEs) are mobile genetic elements which can change location and generate new copies in their host genomes. As a result, TEs populate eukaryotic genomes and can contribute to a significant fraction of their nuclear DNA [1]. Owing to their mobile nature, TEs have an important role in evolution and genetic innovation, but also pose a threat to the genome integrity [2,3]. To control TEs, various mechanisms have evolved to prevent or reduce their mobility.

In eukaryotes, transcriptional gene silencing (TGS) of TEs relies on the deposition of chromatin modifications leading to the formation of compact chromatin states, collectively known as heterochromatin, incompatible with canonical Pol II transcription [4]. Heterochromatin is generally characterized by repressive epigenetic marks, prominently DNA methylation (5'methylcytosine; 5mC), histone modifications such as histone H3 lysine-9 mono- and dimethylation (H3K9me1/H3K9me2, constitutive heterochromatin) or histone H3 lysine-27 triple methylation H3K27me3 (facultative or developmentally regulated heterochromatin), and the presence of specific histone variants [5,6]. Upon silencing, transposons accumulate mutations and degenerate into non-autonomous TEs and fragmented remnants or relics [7]. Despite their inability to mobilize, degenerated TEs provide alternative DNA and RNA regulatory sequences (e.g., transcription factor binding sites, splicing information, transcriptional start or termination sites) or induce recombination due to their repetitiveness [8–12]. Therefore, maintaining epigenetic control is important even for degenerated TEs, and in consequence, TE abundance and location largely shapes the epigenetic landscape of genomes [13–17].

Often involved and largely conserved in mechanisms to suppress TEs and maintain epigenetic silencing are ~18-30-nt long small RNAs (sRNAs) and members of the ARGONAUTE (AGO) protein family. Loaded into AGOs, TGS-associated sRNAs guide chromatin modifying complexes to transposons through sequence complementarity [18,19]. In flowering plants (angiosperms), where TE silencing is mostly associated with DNA methylation and H3K9me2, this mechanism is known as RNA-directed DNA methylation (RdDM). Best studied in *Arabidopsis thaliana,* during RdDM, H3K9me2 and DNA methylation recruit the plant-specific RNA Polymerases IV and V (Pol IV, V) to respectively provide the source and target for 24-nt siRNAs,

loaded into members of the AGO4/6 clade, to deposit DNA methylation in all cytosine contexts (CG, CHG and CHH, where H is any nucleotide but G). Once stablished, DNA methylation can reinforce the deposition of H3K9me2 through the action of SUPRESSOR OF VARIEGATION 3-9 HOMOLOGUE (SUVH) family H3K9 methyltransferases, which can read DNA methylation. In turn, CHROMO-METHYL TRANSFERASE 3 and 2 (CMT3, CMT2) can bind to H3K9me2 to mediate CHG and CHH methylation respectively. This results in a self-reinforcing loop between non-CG methylation and H3K9me2 independent of siRNAs, although in Arabidopsis most of the CHH methylation depends on the RdDM pathway. Symmetrical CG methylation is maintained across cell divisions through the action of METHYL-TRANSFERASE 1 (MET1) (Reviewed in [20]). Loss of RdDM, 5mC or H3K9me2 results in the transcriptional reactivation of TEs [21–25]. Therefore, DNA methylation, together with H3K9me2 and in cooperation with sRNAs, keep TEs silenced.

Loss of TE control in somatic tissue is likely less deleterious as long as it does not affect the germline to the next generation in which stability of the genome is highly relevant. In accordance, small RNA-guided TGS pathways are prominently active in reproductive tissues and silence active TEs encountered during fertilization [26–28]. Nevertheless, as opposed to metazoans, RdDM is not constrained to reproductive organs in most flowering plants: 24-nt siRNAs are also present in vegetative, somatic organs such as leaves or roots across all developmental stages [29–31]. With interesting exceptions: besides the aquatic carnivorous plant *Utricularia gibba*, the monocot *Spirodela polyrhiza* (L.) Scheid. contains much less 24-nt siRNAs and reduced genome-wide levels of DNA methylation [32–34]. In contrast to the small *U. gibba* genome, where repeats represent less than 3% of the genome [35], the TE content in the genome of Spirodela (~139 Mbp) is similar to Arabidopsis [36] excluding the reduced genomic TE load as explanation for the low levels of DNA methylation and small RNAs. Furthermore, several sRNA silencing components including some involved in RdDM have been shown to be missing, low or not expressed. In addition, CMT2, which maintains CHH methylation, is lost [37–39].

The *Spirodela* genus is the most basal of the *Lemnaceae* family (order Alismatales), a monophyletic group of free-floating freshwater monocot plants, commonly known as duckweeds, composed of 5 genera (*Spirodela*, *Landoltia*, *Lemna*, *Wolfiella* and *Wolffia*) [40,41] (**Fig. 1a, Supplemental Fig. S1a, b**). Comprised of a leaf-like structure known as a frond, ranging from ~2 cm to 1 mm, duckweeds display

an evolutionary reduction and simplification of their body plan compared to most

angiosperms, including vestigiality and loss of organs such as roots and vasculature

[42–45] (**Supplemental Fig. S1c**). Although flowering and seed production has been

reported, it is rare in most species [42,46,47]. Duckweeds mostly propagate through

rapid clonal reproduction (doubling of individuals ~1-5 days), known as vegetative

budding: new fronds originate from budding pockets where meristems containing stem

cells are located [42,48]. Mother fronds (MF) continuously give rise to daughter fronds

(DF), which carry grand-daughter fronds (GDF) in their budding pockets even before

detaching from the mother frond [42,43] (**Fig. 1b**). This extremely efficient amplification

of somatic tissue, omitting regular sexual propagation, and the free-floating way of

living, make Spirodela a unique model to explore if the differences in epigenetic

parameters are connected with these differences in lifestyle compared to other

angiosperms.

Our results expand the notion of simplified silencing pathways in Spirodela.

DNA methylation and H3K9me2 profiles indicate that maintenance and feedback loops

of RdDM are still functioning at the few long, intact TEs but are weakened at the

majority of degenerated TEs, explaining the generally poorly methylated Spirodela

genome. Nonetheless, H3K9me1 remains at demethylated TE sequences. Despite

the absence of an obvious DCL2 ortholog in the genome, the siRNA fraction contains

22-nt siRNAs, produced from TEs as well as other endogenous and exogenous

sources of dsRNA. Our data document the variability of TE control and open the

possibility to study tissue-specific or developmentally regulated mechanisms, in

connection with the lifestyle, and reproduction strategy of Spirodela.

Results

Genome sequencing of *S. polyrhiza* #9509 from the Landolt collection.

The genomes of several Spirodela polyrhiza accessions (clones) have been

sequenced [32,36,49], including a high-resolution reference genome assembly of its

20 chromosomes from the *S. polyrhiza* 9509 and 7498 clones (Sp9509, Sp7498)

[32,50,51]. For our study, we retrieved Sp9509 from the Landolt duckweed collection

(Zurich, Switzerland), now managed at the Institute of Agriculture Biology and

Biotechnology (IABB, CNR, Milan, Italy). We performed 88X coverage short-read

Illumina genome sequencing to account for potential genetic differences due to

independent clonal propagation over thousands generations between this accession

and that of the Rutgers Duckweed Stock Cooperative (RDSC, Rutgers University, NJ,

USA) used to generate the reference genome [50]. The Landolt Sp9509 genome

showed ~163k indels and ~512k SNPs (~0.4% of the Spirodela genome) compared to

the Sp9509 reference (**Supplemental Fig. S2**). Given our special interest on TEs, we

performed a *de novo* TE annotation using a combination of EDTA and

RepeatModeler2 to identify repeats that were then further classified using DeepTE

[52–54].

Silencing pathways and epigenetic mechanisms in general are well

characterized by molecular and genetic means in Arabidopsis. Similar genome size

and TE content of Spirodela and Arabidopsis (**Fig. 1c**) allow therefore valid

comparisons despite the evolutionary distance between the species (**Fig. 1a,**

Supplemental Fig. S1a).

Small RNA and DNA methylation patterns in Spirodela imply low RdDM activity

in Spirodela.

We compared the AGO-loaded small RNAs, using TraPR [55], of Spirodela

confluent cultures (including fronds in all developmental stages) to those of 10 day-old

Arabidopsis seedlings as the closest equivalent developmental stage. In contrast to

Arabidopsis, where TE-derived 24-nt siRNAs are highly abundant, Spirodela displays

very low levels of 24-nt siRNAs and its sRNA profile is dominated by 21-22-nt sRNAs

(**Fig. 1d, Supplemental Fig. S3**), predominantly contributed by regulatory microRNAs

(miRNA). The lack of 24-nt siRNAs in Spirodela is more pronounced than their loss in

Arabidopsis RdDM siRNA biogenesis defective Pol IV mutant (*nrpd1*) (**Supplemental**

Fig. S3). Furthermore, the reduced amounts of 24-mers is not compensated by TE-

derived 21-22-nt siRNAs as in DNA methylation-deficient Arabidopsis mutants [56–

58]. Hence, low levels of 24-nt siRNAs in Spirodela are compatible with the hypothesis

of reduced or lack of RdDM activity at TEs (**Supplemental Fig. S4**).

Methylome analysis by Enzymatic Methyl-seq (EM-seq) [59] showed similar

results to those previously published using bisulfite sequencing [32]. Transposon 5mC

in all contexts is lower in Spirodela compared to Arabidopsis, with very low mCHG

levels and hardly any mCHH (**Fig. 1e**). This is in agreement with the partial

dependency of mCHH on 24-nt siRNAs and seconds a weak RdDM in *S. polyrhiza*. In

Figure 1: *Spirodela polyrhiza* general developmental, small RNA and DNA methylation patterns.
a Simplified phylogenetic relationship of Spirodela and other duckweeds relative to Arabidopsis. **b** Representative images of Spirodela fronds in culture and schematic representation of the organization of clonal generations in a group of fronds during vegetative growth. **c** Genome size, number and genomic occupancy of genes and TEs in Spirodela and Arabidopsis. **d** Size distribution and genomic feature annotation of TraPR-purified small RNA from Arabidopsis and Spirodela. **e** Metaplots of averaged weighted DNA methylation in all three contexts (CG, CHG and CHH, where H is any but G) over transposons (≥100 bp) and 2 kb flanking regions in Arabidopsis and Spirodela. **f** number and percentage of transposon CG sites displaying methylation levels equal or above 1% in Arabidopsis and Spirodela. **g** DNA methylation levels at CG sites from (f). **h** number and percentage of transposon CHG sites displaying methylation levels equal or above 1% in Arabidopsis and Spirodela. **i** DNA methylation levels at CCG and CWG in Arabidopsis and Spirodela. **h** Inner 5mC (CmCG) DNA methylation levels at CCG sites with outer 5mC (mCCG) equal or higher than 1% in Arabidopsis and Spirodela. (**) indicates p-value <2.22 x 10⁻¹⁶ (Wilcoxon rank sum test). In all boxplots: median is indicated by solid bar, the boxes extend from the first to the third quartile and whiskers stretch to the furthest value within 1.5 times the interquartile range. At: Arabidopsis, Sp: Spirodela.

186   addition, Spirodela lacks gene body CG methylation (gbM) [32] (**Supplemental**

187   **Fig. S4a**). Although the role of gbM is unclear, it generally associates with highly

188   expressed genes. In contrast, mCG near the transcription start site (TSS) is associated

189   with gene silencing [60,61]. Like in Arabidopsis, mCG around the TSS is higher in non-

expressed genes of Spirodela (**Supplemental Fig. S4b, c**), indicating that mCG in gene regulatory sequences influences transcription also in Spirodela.

CG and CHG methylation signatures are compatible with functional maintenance of DNA methylation independently of small RNAs.

Once established, DNA methylation can be further maintained independently of small RNAs through several pathways depending on the sequence context. CG methylation is maintained across cell divisions through the action of MET1 which, after DNA replication, copies mCG patterns in daughter strands [62–64] . In Spirodela, the ratio of mCG sites at TEs (51,9%) is nearly half of that in Arabidopsis (90,9%) (**Fig. 1f**). However, at CG sites with 5mC levels equal or higher than 1%, mCG is similarly stable as in Arabidopsis, displaying median methylation levels above 80% (**Fig. 1g**). Hence, the reduced global mCG level at TEs in Spirodela is the result of less individual CG sites being methylated. However, a higher variation in Spirodela than in Arabidopsis (**Fig. 1g**) does not exclude that MET1 might be less efficient, not active in all cells, or not all CG sites being as faithfully maintained.

As for mCG, the number of mCHG sites at TEs is much lower in Spirodela (36,1%) than in Arabidopsis (90,3%) (**Fig. 1h**). CHG sites can be further divided into CCG and CWG (CAG/CTG) sub-contexts. In Arabidopsis, mCWG is higher than mCCG (**Fig. 1i**) due to the combined differential affinities of SUVH4/5/6 and CMT3 to read and methylate individual cytosine sequence contexts [63,65–67]. This applies also to Spirodela, despite lower mCHG in both sub-contexts (**Fig. 1i**). mCCG maintenance relies additionally on MET1, as the sites have a C in both CG and CHG contexts, are generally observed as mCmCG due to the outer cytosine methylation (mCCG) being largely dependent on the internal CG methylation (CmCG) mediated by MET1 [65,66,68,69]. Consequently, mCCG sites in Arabidopsis display a high internal CG methylation level, a pattern also observed in Spirodela (**Fig. 1j**). The similar mCHG patterns between the two species indicate that the responsible enzymes, SUVHs and ZMET, the CMT3 homolog in monocots [70] probably share the same sequence preference albeit their activity seems to be lower than in Arabidopsis. Nonetheless, given that CMT3/ZMET has been shown to be involved in gbM [60,71,72], lower mCHG and lack of gbM in Spirodela might be related.

Altogether, our analysis indicates that mCG and mCHG maintenance mechanisms in Spirodela might operate in a similar fashion as in Arabidopsis.

However, while low mCG is a consequence of reduced number of CG sites being methylated, low mCHG appears to be the combined result of lower number of targeted CHG sites and decreased maintenance activity.

Genetic composition and expression of silencing pathways partially account for sRNAs and DNA methylation features in Spirodela.

In addition to low signatures of RdDM in Spirodela, coding potential for several of its components have been shown to be missing [37,39,73]. To gain further insights why levels of 24-nt siRNA and DNA methylation are reduced, we set to identify and examine the expression of known core components of angiosperm silencing pathways in *S. polyrhiza* (**Fig. 2, Supplemental Fig. S5**). The identification of orthologues and paralogues in the genome of *S. polyrhiza* 9509 was based on the presence of conserved functional domains. Annotations were further curated based on phylogeny trees built with proteins from several angiosperms and the prediction of known key domains and their organization (**Supplemental Fig. S6-S16**). For gene expression, we performed short-read transcriptome analysis in Spirodela and compared it to that of recently published Arabidopsis seedlings [74] (**Fig. 2**). We further performed long-read RNAseq (PacBio Isoform-Sequencing, Iso-Seq) in Spirodela to validate and eventually improve gene model annotations.

As expected from the prevalence of miRNAs in Spirodela TraPR analysis, conserved plant miRNA biogenesis factors [75] are expressed at similar levels as those in Arabidopsis (**Fig. 2a**). Likewise, genes required for the production of regulatory 21-nt siRNA trans-acting (ta)siRNA [76,77]: SUPRESSOR OF GENE SILENCING 3 (SGS3), RNA-DEPENDENT RNA POLYMERASE 6 (RDR6) and DICER-LIKE 4 (DCL4), are present and expressed (**Fig. 2b**). Remarkably, Spirodela lacks many components, effectors and regulators of 21-22-nt siRNA pathways, such as RNASE THREE-LIKE 1 (RTL1), DOUBLE-STRANDED RNA BIDING PROTEIN 7 (DRB7) [78–80], and orthologues of the RDRγ clade highly conserved in plants and fungi [81,82] (**Fig. 2b,c, Supplemental Fig. S6, S7**). Gene absences include the antiviral factors DICER-LIKE 2 (DCL2) and ARGONAUTE 2 (AGO2) [83–86] (**Fig. 2b, c, Supplemental Fig. S8, S9**). In contrast, two RDR1 and eight tandemly repeated copies of AGO5 are present (**Fig. 2b, c, Supplemental Fig. S7, S9, S10**), which have been implicated in virus exclusion and TE silencing in meristems [87–90].

Figure 2: Genetic composition and expression of silencing pathways in Spirodela. a-f Expression in transcripts per million (TPM) of silencing components in Arabidopsis and their identified orthologues and paralogues in the *Spirodela polyrhiza* #9509 (Sp9509) genome. Gene IDs for both Arabidopsis and Spirodela can be found in Supplementary Figure 5.

Regarding the RdDM pathway, Spirodela contains the minimal set of its components, either with no or with a reduced number of paralogues compared to Arabidopsis (**Fig. 2c, d**). In Spirodela, SAWADEE HOMEODOMAIN HOMOLGUE 1 (SHH1), a dual reader for repressive H3K9me2 and unmethylated H3K4(me0), and the chromatin remodeler CLASSY 1/2 (CLSY1/2) are absent (**Fig. 2d, Supplemental Fig. S11, S12**). However, a single orthologue of CLSY3 was found in Spirodela (**Fig. 2d, Supplemental Fig. S11**). In Arabidopsis, the SHH1-CLSY1/2 module mainly recruits Pol IV to short TEs within gene-rich regions, while CLSY3 and its paralogue CLSY4 recruit Pol IV to TE-rich pericentromeric regions in a SHH1-idependent but DNA methylation-dependent manner [31,91–93]. However, Spirodela has two SHH2 orthologues (**Supplemental Fig. S11**). Part of a nucleosome positioning complex during transcription initiation in Arabidopsis [94], in maize, SHH2 binds preferentially

to H3K9me1 *in vitro* and interacts *in vivo* with both Pol IV and V in maize, possibly connecting RdDM to H3K9me1 installation [95,96].

Similarly, no AGO6 orthologue was found, yet two AGO4, both required non-redundantly to maintain DNA methylation at most RdDM target loci in Arabidopsis [97] (**Fig. 2d, Supplemental Fig. S9**). A single SUVH2 orthologue is present in Spirodela (**Fig. 2d, Supplemental Fig. S13**), which together with its paralogue SUVH9 redundantly recruit Pol V to RdDM targets by binding to methylated DNA in Arabidopsis [98–100]. Regarding the RdDM-associated DNA methyltransferases, DOMAIN REARRANGED METHYLTRASFERASE 2 (DRM2) but no DRM3, a conserved catalytically inactive paralogue that acts as cofactor of DRM2 [101–105], orthologues were found (**Fig. 2d, Supplemental Fig. S14**). Additionally, although NRPE1 (Pol V) and its partner SUPPRESSOR OF TY INSERTION-LIKE (SPT5L) (a paralogue of the general transcription elongation factor SPT5) are conserved in Spirodela; we noticed that both contain fewer tryptophan/glycine (GW/WG) repeats than those of other angiosperms (**Fig. 2d, Supplemental Fig. S15, S16**) [106–110]. GW/WG repeats, or AGO-hooks, allow AGO4/6 binding to Pol V to scan its nascent transcripts for positive siRNA matches to recruit DRM2. These, together with NRPD1, RDR2, DCL3, and the rest of conserved RdDM components, might suffice for a functional pathway in Spirodela (**Fig. 2, Supplemental Fig. S7, S8, S12, S15**). Nonetheless, their low expression levels (**Fig. 2 c, d**), rationalize the low abundance of 24-nt siRNAs and associated CHH methylation in Spirodela (**Fig. 1d, e**).

Lastly, we inspected DNA methylation maintenance pathways. As predicted from the patterns of mCG and mCHG (**Fig. 1f-j**), MET1, CMT3/ZMET, SUVH4, SUVH5 and DECREASE IN DNA METHYLATION 1 (DDM1) orthologues were found present and expressed in Spirodela, albeit at a lower degree than in Arabidopsis. On the contrary, CMT2, required to maintain mCHH together with SUVH4 [23], is missing as previously reported [73] (**Fig. 2e, Supplemental Fig. S13, S14**).

Thus, reduced 24-nt siRNA and mCHH levels in Spirodela can be explained by the combined reduction in RdDM composition and expression in addition to the absence of CMT2. However, it does not account for the global decrease in mCG and mCHG as their maintenance pathways seem intact and expressed in Spirodela.

Few transposable elements from different superfamilies display high levels of DNA methylation in Spirodela.

To further gain insight into the patterns of TE DNA methylation, we examined its distribution along the Spirodela genome. To exclude any confounding effects of gbM, we calculated mean DNA methylation levels exclusively on TEs and their density in 100-kb windows. At gene-rich regions, DNA methylation is very scarce (**Fig. 3a, Supplemental Fig. S17a**). As in Arabidopsis, Spirodela DNA methylation is concentrated at TE-dense regions (presumably centromeric and pericentromeric domains). However, not all TE-dense regions are highly methylated (**Supplemental Fig. S17b**). Hence, we investigated if local DNA methylation levels correlate with TE density as in other angiosperms [14,16,111,112]. At equal TE-content, DNA methylation is lower in all contexts in Spirodela compared to Arabidopsis, especially at CHG and CHH (**Fig. 3b, Supplemental Fig. S18a**). Moreover, at the CG context, Spirodela displays a broad variation range for a given TE content: even at TE densities above 50% or higher, average mCG still spans between 0 and 75% (**Fig. 3b**).

To address the origin of such variation, we investigated the DNA methylation levels of individual transposons. This revealed that the majority of TEs in Spirodela display low or very low levels of DNA methylation in all contexts, especially CHG and CHH (**Fig. 3c, Supplemental Fig. S18b**). Nonetheless, despite this substantially reduced DNA methylation, increased expression at TEs was not observed (**Fig. 3d**). TEs with high or low DNA methylation levels did not belong to any specific transposon class or superfamily. For all the main plant TE superfamilies [113] (TE classification and codes used here are summarized in **Supplemental Fig. S19**), DNA methylation was nearly absent except for a few elements (**Fig. 3e**). Thus, low genome-wide methylation levels in Spirodela result from only a few methylated TEs, although these can display high mCG levels similar to those found in Arabidopsis (**Fig. 3e**).

Figure 3: Distribution of DNA methylation in Spirodela chromosomes and transposable elements. a Circos plot of genome-wide distribution of DNA methylation densities by chromosome in Spirodela. From outside to inside: gene density, TE density, CG, CHG and CHH methylation. Mean DNA methylation, gene and TE densities values within the indicated ranges are displayed for 100 kbp intervals in 50 kbp sliding windows. Arabidopsis chromosome 3 is shown for comparison. **b** Scatter plot and best fit line of average TE CG and CHG DNA methylation against TE content per 100 kb genomic bins in Arabidopsis and Spirodela. Colored shadows indicate the standard error of the mean. **c** Bar plot of the distribution (%) of TE annotations (≥100 bp) within 10% incremental intervals of CG and CHG methylation in Arabidopsis and Spirodela. **d** Analysis of Spirodela TE expression within 25% incremental intervals of CG methylation. Global gene expression values are shown for comparison. **e** Distribution of CG, CHG and CHH methylation across the main TE superfamilies in Arabidopsis and Spirodela. The number of TE annotations belonging to each superfamily by species is indicated (n). (**) indicates p-value <2.22 x 10^-16 (Wilcoxon rank sum test). In boxplots, median is indicated by solid bar, the boxes extend from the first to the third quartile and whiskers stretch to the furthest value within 1.5 times the interquartile range. At: Arabidopsis, Sp: Spirodela.

Potentially intact transposons are the conspicuous targets of DNA methylation in Spirodela.

Although TE location and density influences their epigenetic regulation [15], in Arabidopsis and tomato DNA methylation of longer, evolutionary younger and therefore potentially mobile TE insertions are differentially regulated than often fragmented TE relics resulting from the accumulation of mutations and truncations over evolutionary times following their silencing [7,15,114]. On one hand, DNA methylation in long TEs is maintained by a combination of RdDM acting on their boundaries (mainly in young active TEs) and CMT activity along their body. On the other hand, short, degenerated TE remnants do not need CMT to maintain their methylation [115–117]. To explore if transposon length influences DNA methylation in Spirodela, we sorted TEs into short TE fragments (100 to 500 bp), intermediate TEs (0.5 to 4 kb) and long TEs (above 4 kb). This revealed that only long TEs are heavily methylated at CG and, to a lesser extent, in CHG. Intermediate TEs display a wide range of mCG levels, whereas TE fragments are minimally or not methylated (**Fig. 4a**).

To further explore the connection between TE length and DNA methylation we focused on the two superfamilies of long terminal repeat (LTR) retroelements (RTEs): *Ty1/Copia* (RLC) and *Ty3/Gypsy* (RLG). LTR-RTEs are the most abundant class of TEs in the Spirodela genome, collectively representing more than 50% of the transposon fraction of the genome and largely contributing as well to that of Arabidopsis (**Fig. 4b, Supplemental Fig. S20**). In addition, the genetic structure and domain organization of complete functional elements are well known [118–120]. Thus, LTR-RTEs offer a good case study to investigate any link between DNA methylation and TE length, using the latter as a proxy for their completeness and potential autonomy and mobility. For better resolution, we split TE sizes into further size categories in the intermediary range between 0.5 and 4 kb. Within each size bin, Spirodela DNA methylation is lower than in Arabidopsis. However, in Spirodela, average mCG and mCHG gradually increase with TE size, although mCHG does not

Figure 4: Length-dependent TE DNA methylation in *Spirodela polyrhiza*. a CG, CHG and CHH methylation levels (%) across all TEs (≥100 bp) in Arabidopsis and Spirodela split by size into short (≥100 bp, <500 bp), intermediate (>500 bp, ≤ 4 kbp) and long (> 4kbp). **b** Waffle chart of the relative contribution of different TE superfamilies over the total genomic space occupied by TEs in Spirodela. **c-d** CG and CHG methylation levels (%) between different length range groups for RLC: *Ty1/Copia* (c) and RLG: *Ty3/Gypsy* (d) in Arabidopsis and Spirodela. **e-f** Presence and number of RCL (e) and RLG (f) TE protein domains across TE annotations of varying. Violin and box plots depict domain count distributions for TEs with at least one domain. Numbers above denote TE annotations in each size category with at least one domain, with percentages shown in brackets relative to total TE annotations per size category (g-h). Schematic representation of RLC and RLG protein and domain structure is included on top of each plot. GAG: Nucleocapsid; PR: Protease; IN: Integrase; RT: Reverse transcriptase; RH: RNase H; CHR: Chromodomain, present only in the Chromoviridae genus of RLG. **g-h** Number of TE annotations and relative contribution to total genomic TE load (TE occupancy %) at each size category for RLC (g) and RLG (h). (*) indicates p-value <0.01 (**) indicates p-value <0.0001 (Wilcoxon rank sum test). In all boxplots: median is indicated by solid bar; the boxes extend from the first to the third quartile and whiskers stretch to the furthest value within 1.5 times the interquartile range. At: Arabidopsis, Sp: Spirodela.

achieve levels comparable to those of Arabidopsis (**Fig. 4c, d, Supplemental Fig. S21a, b**). The proportion of TEs displaying high CG methylation rises across TE-size groups. While very few TE-fragments below 500 bp are highly methylated, the majority of long TEs (>4 kb) have mCG above 60% (**Fig. 4c, d**). At the same time, the fraction of TEs with at least one intact cognate protein domain and the number of domains identified per TE annotation increase with each size category, with the majority of long LTR-RTEs containing all domains associated with each superfamily (**Fig. 4e, f**).

Therefore, in Spirodela, most potentially intact TEs display high levels of DNA methylation, which decreased as they degenerated and decayed. Given that long intact LTR-RTEs represent a minority of Spirodela TEs (**Supplemental Fig. S21c, d**), their low abundance helps explain the few TE numbers with high DNA methylation (**Fig. 3c, e**). In addition, the proportion of hypomethylated shorter LTR-RTE fragments relative to methylated intact TEs is higher in Spirodela than in Arabidopsis (**Fig. 4g, h, Supplemental Fig. S21c-f**), resulting in the global reduction of TE methylation (**Fig. 1e, Fig. 3a, b**).

Levels of H3K9me2, but not H3K9me1, are reduced in Spirodela.

Considering the association between DNA and H3K9 methylation in plants, we explored how far DNA methylation in Spirodela would impact the histone methylation landscape. H3K9me2 and H3K9me1 are commonly linked with heterochromatin formation at silenced TEs [6,23,121,122]. Western blot analysis revealed that the H3K9me2 level in Spirodela is lower than in Arabidopsis, whereas H3K9me1 displayed similar levels (**Fig. 5a**). This was unexpected as we anticipated in Spirodela a decrease of both marks, given their connection with DNA methylation and reduction upon loss of non-CG methylation in Arabidopsis [23,123]. To corroborate the observation, we investigated histone post-translational modifications (PTMs) by mass-spectrometry [124,125], which confirmed an approximately 8-10-times lower level of H3K9me2 but similar H3K9me1 levels in Spirodela relative to Arabidopsis (**Supplemental Fig. S22**).

Figure 5: Patterns of H3K9me1/2 distribution in *Spirodela polyrhiza* transposons. a Protein blot analysis of H3K9me1 and H3K9me2 abundance in Spirodela compared to Arabidopsis at different protein amounts (μg). Total H3 protein levels are shown as loading control. **b-c** Heatmap of k-means clustering of transposons in Arabidopsis (b) and Spirodela (c) using H3K9me1, H3K9me2, H3K27me3 and H3K4me3 enrichment over H3, presented as log2[ChIP(X)/ChIP(H3)]. Enrichment is represented by color, while depletion is shown in gray. Each row corresponds to a length-normalized TE annotation ± 1 kb. Transposons are grouped into clusters 1 to 5 based on the abundance of examined marks. **d** Metaplots of average weighted H3K9me1 (solid line) and H3K9me2 (dashed line) enrichments over H3 at cluster 1 and 2 transposons ± 1 kb in Arabidopsis (colored) and Spirodela (gray). **e** Relative genome (%) occupancy of each of the TE clusters defined in b-c in Arabidopsis (grey) and Spirodela (colored). **f** Genome browser capture of the distribution of H3K9me1 and H3K9me2 enrichments, together with DNA methylation, along genes and transposons in Spirodela. **g** Venn diagram of the overlap between H3K9me1 and H3K9me2 peaks along Spirodela genome. **h** Genome occupancy (%) of H3K9me1 and H3K9me2 in Spirodela genome. **i** Scatter plots of DNA methylation in all three contexts against H3K9me1 and H3K9me2 enrichments of TEs in Cluster 1, 2 and 3/4/5 combined in Spirodela. *R* and *p* indicate Pearson correlation coefficients and P-values. **j** Analysis of H3K9me1 and H3K9me2 enrichments in different TE length groups for RLC and RLG combined in Spirodela. In all boxplots: mean is indicated by solid bar; the boxes extend from the first to the third quartile and whiskers stretch to the furthest value within 1.5 times the interquartile range. (*) indicates p-value <0.001. n.s.: not significant.

A low proportion of TEs show high H3K9me2 enrichment in Spirodela.

To further delve into the distribution of H3K9me1/2 in Spirodela and how the general organization of heterochromatic marks is affected by its reduced DNA methylation levels, we performed chromatin immunoprecipitation (ChIP) in Spirodela fronds and Arabidopsis 10 days-old seedlings (**Supplemental Fig. S23**) and focused on the association of these modifications with the TEs. In addition to H3, H3K9me1 and H3K9me2, we generated genomic profiles through ChIP-seq for H3K27me3 and H3K4me3. Although Polycomb-mediated H3K27me3 is generally mediates developmentally regulated gene silencing, its association with TEs was investigated given that it also regulates TEs in the absence or loss of DNA methylation [126–129], and may contribute directly to TE silencing [130]. H3K4me3, was taken as an euchromatic mark control [13,131–134].

As observed for DNA methylation (**Fig. 3a**), H3K9me1/2 is concentrated at TE-dense regions in Spirodela (**Supplemental Fig. S24**), likely centromeric and pericentromeric heterochromatic regions [13,73,121]. The smaller size of such regions, compared to Arabidopsis, and their distribution in the Spirodela 20 chromosomes, fits the speckled pattern of small dense DNA foci in DAPI staining of interphase nuclei of Spirodela (**Supplemental Fig. S25**) [135].

A more detailed K-means clustering analysis of histone mark profiles relative to H3 over TEs, identified 5 clusters with similar features in both Arabidopsis and Spirodela (**Fig. 5b, c**). Clusters 1 and 2 contain TEs enriched for H3K9me1/2 while clusters 3 and 4 consist of those marked by H3K27me3 and H3K4me3 respectively. Lastly, cluster 5 encompasses TEs depleted of any of the profiled marks (**Fig. 5b, c**).

Although cluster 1 and 2 TEs are marked by H3K9me1/2, they display homogeneous H3K9me1 but contrasting H3K9me2 levels (**Fig. 5d, Supplemental Fig. S26**). TEs in cluster 1 are more enriched in H3K9me2 than in H3K9me1 while in cluster 2 TEs H3K9me2 is below H3K9me1 (**Fig. 5d**). Remarkably, the two clusters comprise very different fractions of the two genomes and TE space. In Spirodela,

cluster 1 only comprises about 20% of the TE space, representing less than 6% of the genome, while in Arabidopsis is the most abundant one, encompassing ~35% of the TE space and ~12% of the genome. Cluster 2, on the contrary, occupies 45% of the Spirodela TE space (more than 10% of its genome), but only about 20% in Arabidopsis

(5% of the genome) (**Fig. 5e, Supplemental Fig. S27a**). Given the similar genome size and TE content in both species, the opposed contribution of cluster 1 and cluster 2 to their genomes is in line with the levels of H3K9me2 observed by western blot and mass-spectrometry. Furthermore, similar H3K9me1 levels in both clusters concur with its even abundance in Spirodela and Arabidopsis. Hence, reduced DNA methylation levels in Spirodela correlate with a smaller fraction of the genome displaying a high H3K9me2 enrichment over TEs but it does not impact H3K9me1 distribution, nor it leads to a widespread compensatory regulation of TEs by H3K27me3.

H3K9me1 marks TEs independently of their length and methylation status in Spirodela.

A closer inspection of the distribution of H3K9me1/2 along Spirodela TEs revealed that H3K9me1 uniformly covers TEs, while H3K9me2 peaks are found only at individual TEs (**Fig. 5f**). Furthermore, while more than 95% of H3K9me2 peaks overlapped with H3K9me1 domains, a significant fraction of H3K9me1 peaks (60,5%) did not with H3K9me2-associated regions (**Fig. 5g**). Moreover, most H3K9me2 peaks were embedded within H3K9me1 (**Supplemental Fig. S27b**). Adding to that, H3K9me1-enriched regions occupy 25% of the Spirodela genome whereas H3K9me2 peaks represent less than 12% (**Fig. 5h**). This indicates that, similarly to DNA methylation, high levels of H3K9me2 are only found within few TEs. It prompted us to investigate the association between H3K9me1/2 and DNA methylation in Spirodela and indeed, H3K9me2 levels in Spirodela positively correlate with DNA methylation, including mCHG, like in Arabidopsis (**Fig. 5i, Supplemental Fig. S28**). Parallel to DNA methylation (**Fig. 3**), H3K9me2 increased with TE size (**Fig. 5j**). This further supports that, as previously inferred, the positive feedback loop between mCHG and H3K9me2 operates in Spirodela. However, this did not apply to H3K9me1. While in Arabidopsis TEs enriched in H3K9me1 invariably displayed high levels of DNA methylation (**Supplemental Fig. S28**), they did not so in Spirodela with its broad range of DNA methylation levels at TEs (**Fig. 5i**). Accordingly, H3K9me1 levels vary little across TE sizes (**Fig. 5j**).

These results indicate that, in Spirodela, H3K9me1 serves as broad transposon mark but does not rely on DNA methylation as H3K9me2, suggesting that they might be differentially regulated. Although Arabidopsis H3K9 histone methyl-transferases (HMTs) can catalyze both H3K9me1 and me2 in vitro [121,136], it is unclear how their

mono- and di- methyl-transferase activity is regulated in vivo or if both marks are deposited by the same HMTs. Loss of SUVH4, the main HMT in Arabidopsis, results in a strong reduction of H3K9me2 and loss of epigenetic silencing, only marginally affecting H3K9me1 and global heterochromatin organization [121,137,138]. This originally led to the proposition that H3K9me2 and H3K9me1 play different roles in silencing and heterochromatin maintenance respectively [121]. Indeed, increased H3K9me2 and DNA methylation at long TEs, in contrast to shorter TE fragments, hints that their self-reinforcement is focused at potentially functional or recently active ones to ensure their silencing.

Long, complete TEs show signatures of RdDM activity.

For the reasons explained above, we focused the detailed analysis on the complete LTR-RTEs of the *Copia* (RLC) and *Gypsy* (RLG) superfamilies in which H3K9me2 levels were higher than those of H3K9me1. However, the patterns differed depending on the TE superfamily. While in RLCs had more H3K9me2 towards the center of the TE, the same modification peaked on the edges of RLGs (**Fig. 6a**). Such patterns resembled those found in Arabidopsis intact LTR-RTEs, (**Supplemental Fig. S29a**). DNA methylation at CG and CHG followed a similar trend as H3K9me2 in both cases (**Fig. 6b**), although mCHG was comparatively lower (**Fig. 6b, Supplemental Fig. S29b**). Furthermore, albeit very low, peaks of mCHH were also observable at RLGs edges (**Fig. 6b**). Hence, the layout of epigenetic silencing marks along complete TEs in Spirodela are very similar to those in Arabidopsis.

Given that in Arabidopsis these patterns are partially determined by RdDM [23,68,116], we inspected if small RNA mapped to complete TEs in Spirodela. Indeed, siRNAs derived from regions with DNA methylation and H3K9me2 were present in Spirodela, specially 24-nt siRNAs, despite at lower levels than in Arabidopsis (**Fig. 6c, Supplemental Fig. S29c**). Moreover, 24-nt siRNAs mapping to complete TEs displayed an adenine 5'-nucleotide bias (5'A), characteristic of small RNAs loaded into the RdDM effector AGO4 [29,139] (**Fig. 6d**), although not as preferential as in Arabidopsis (**Supplemental Fig. S29d**). Complementarity of siRNAs with DNA methylation and H3K9me2 within the TE body and/or their LTRs were validated for individual complete TEs (**Fig. 6e-g, Supplemental Fig. S30**).

Figure 6: RdDM signatures in *Spirodela polyrhiza* complete TEs. a Metaplots of average weighted H3K9me1 (solid line) and H3K9me2 (dashed line) enrichments over H3 at complete RLC (left) and RLG (right) transposon annotations ± 2 kb in Spirodela. Complete TEs were defined as those for which all RLC and RLG protein domains where identified. n indicates number of complete elements in each superfamily. **b** Metaplots of averaged weighted DNA methylation in all three contexts over RLC and RLG transposons ± 2 kb flanking regions in Spirodela. mCHH values are scaled to 10% max values instead of 100%. **c** Metaplots of averaged weighted 21-, 22- and 24-nt small RNA abundance over intact RLC and RLG ± 2 kb flanking regions in Spirodela. In all metaplots (a-d), solid lines represent the mean and colored shadows the standard error. **d** 5'-nucleotide (5'nt) composition distribution (as % of reads) of 21-, 22- and 24-nt siRNAs mapping to intact RLC and RLG ± 2 kb flanking regions in Spirodela. **e-g** Genome browser captures of selected complete TEs in Spirodela. From top to bottom tracks display: gene and TE annotations, H3K9me1 and H3K9me2 enrichment profiles; mCG, mCHG and mCHH methylation; 21-, 22- and 24-nt siRNA mapping to Watson (+) and Crick (-) DNA strands; RNA expression by short (illumina) and long reads (IsoSeq). Superfamily, clade and length (in bp) of TEs is indicated above their annotation. Light and dark gray shadow delimit TEs and LTRs respectively across all tracks.

We also detected the presence of very low amounts of 21-22-nt siRNAs at the edges of complete TEs (**Fig. 6c**). In some instances, 24-nt siRNAs overlapped with equally abundant 21-22-nt siRNAs, especially at the flanking LTR sequences (**Fig. 6g, Supplemental Fig. S31**). 21-22-nt siRNAs can arise from TE edges in Arabidopsis (**Supplemental Fig. S29c**) as result of DCL2/4 processing of Pol IV transcripts and direct DNA methylation [57,140–142].

To assess if TE-derived siRNAs might participate in RdDM also in Spirodela, we tested if its AGO4 (SpAGO4) has the same small RNA loading preference as Arabidopsis AGO4 (AtAGO4). To do so, we cloned genomic *AGO4* loci from both species with a Flag-HA tag for heterologous transient expression in *Nicotiana benthamiana* leaves. No Sp*AGO4* protein was detectable due to intron retention events when expressed in *N. benthamiana* (**Supplemental Fig. S332a-c**). This was likely due to the inability of dicots (*N. benthamiana*) to properly process introns of the monocots (Spirodela) [143–145]. To circumvent this limitation, we amplified and cloned full length cDNA from one of the two Spirodela AGO4, *AGO4a*, and expressed it in *N. benthamiana* leaves (**Supplemental Fig. S32d**). To assess their small RNA loading preferences, AtAGO4 and SpAGO4a were immunoprecipitated (IP) from *N. benthamiana* leaf lysates [146] (**Fig. 7a**) and small RNAs extracted and sequenced. Like AtAGO4, SpAGO4a displayed a strong preference for 5'A 24-nt long siRNA and reduced affinity for 21-nt siRNA, compared to the population of siRNAs present in *N. benthamiana* leaves (**Fig. 7b, c**).

Therefore, small RNAs might be indeed involved in maintaining DNA methylation and H3K9me2 in Spirodela, at long complete TEs through canonical RdDM. Our analysis of silencing pathways shows that core components required for RdDM are present in the Spirodela genome and, although poorly expressed, probably active during vegetative clonal propagation.

Post-transcriptional silencing of TEs likely involves DCL2-independent 22-nt-siRNAs in Spirodela.

It is unlikely that the observed TE-derived 21-22-nt siRNAs participate in RdDM, due to the low affinity of SpAGO4 to bind them (**Fig. 7b**), and those mapping to TEs have less 5'A bias but more 5'U (**Fig. 6d, Supplemental Fig. S29c**), a signature of AGO1-loaded small RNAs [29,139]. This hints to a role rather in post-transcriptional

Figure 7: *Spirodela polyrhiza* AGO4a siRNA loading preferences and IR-derived siRNAs in transient expression. a Protein blot analysis of Arabidopsis and Spirodela AGO4 (AtAGO4 and SpAGO4a) input and IP fractions from transient expression in *N. benthamiana* 5 days post infiltration (dpi). n.i.: non-infiltrated. Coomassie (Coom.) staining of blot is shown as control for input and IP fractions. Scheme of constructs infiltrated in *N. benthamiana* leaves for transient expression is shown above. p35S: CaMV 35S promoter; F-HA: in-frame N-term Flag and HA peptide tags. Genomic (gDNA) or cDNA origin of the AGO4 sequences is indicated. **b** Size distribution and abundance in reads per million (rpm) of TraPR purified *N. bethamiana* siRNAs (as control) and siRNAs extracted from AtAGO4 and spAGO4a IPs. **c** 5'-nucleotide (5'nt) composition distribution (as % of reads) of 21-, 22- and 24-nt siRNAs from b. **d** Representative pictures of *N. benthamiana* leaves agroinfitrated for transient expression of RUBY or the scarlet hairpin (*hpScarlet*) IR next to small RNA blot analysis in non-infiltrated and 5 dpi leaves. miRNA159 is shown as loading control. **e** Representative pictures of Spirodela fronds agroinfitrated for transient expression of *RUBY* or *hpScarlet* and small RNA blot analysis in non-infiltrated and infiltrated samples at different timepoints. miRNA159 as loading control. **f** Size distribution and abundance of small RNAs mapping to *hpScarlet* from 5 dpi *N. bethamiana* and 12 dpi Spirodela. **g** Stacked distribution of sense (S) and antisense (AS) small RNA coverage by size along *hpScarlet* hairpin in *N. bethamiana* and Spirodela. Schematic representation of the IR is depicted below. **h** 5'-nucleotide (5'nt) composition distribution of 21-, 22- and 24-nt siRNAs mapping to *hpScarlet* in *N. bethamiana* and Spirodela.

silencing (PTGS). During PTGS, 21-22-nt siRNAs loaded into AGORGONAUTE proteins of the AGO1/5/10/18 and AGO2/7 clades target Pol II transcripts to guide mRNA cleavage or translational repression (reviewed in [147]) Coincidentally, we observed expression of TEs producing 21-22-nt siRNAs in our transcriptome datasets, including full length transcripts by Iso-Seq (**Fig. 6g, Supplemental Fig. S31**). Although PTGS is a well-documented control mechanism against transcriptionally active TEs in Arabidopsis [27,56,58,148–150], we were intrigued by the presence of 22-nt siRNAs, given that any identifiable ortholog of DCL2, their cognate Dicer in plants [151–154], is missing in Spirodela and *Lemnaceae* genomes in general (**Fig. 2b, Supplemental Fig. S8**) [39,73].

To rule out artifacts and to investigate if Spirodela can produce 22-nt siRNAs in the absence of DCL2, we attempted to stimulate small RNA biogenesis through expression of suitable exogenous substrate RNAs. As stable transformation in Spirodela is difficult, we chose Agrobacterium-mediated transient expression. This technique has been widely used to trigger and investigate silencing mechanisms, owing to the quick onset of siRNA production in response to expression of double-stranded RNA (dsRNA) following infiltration of plant tissue with Agrobacterium carrying the corresponding transgenes [155–159]. We developed a modified protocol for frond transformation in *Lemna minor* [160] and applied it successfully to induce transient transgene expression in Spirodela fronds (**Supplemental Fig. S33**). We used the visual reporter *RUBY* [161] and an inverted repeat (IR) derived from the coding sequence of the fluorescent reporter *Scarlet* (*Scarlet* hairpin, *hpScarlet*) [89]. Both constructs triggered siRNAs when infiltrated into *N. benthamiana* as expected (**Fig. 7d**). In Spirodela, however, we could not detect *RUBY*-derived siRNAs even after two weeks post-infiltration. On the other hand, *hpScarlet*-derived siRNAs were observed 12- and 14-days post-infiltration (**Fig. 7e**). Sequencing sRNAs from both *N. benthamiana* and Spirodela infiltrated with *hpScarlet* showed that the IR is processed into 21-22-nt siRNAs in both plants. However, Spirodela produceed relatively equal amounts of 21- and 22-nt siRNAs, as observed for TEs, whereas the profile in *N. benthamiana* is dominated by 21-nt siRNAs (**Fig. 7f**). Both siRNA sizes were distributed along the length of the IR dsRNA (**Fig. 7g**) and displayed the 5'U bias signature of AGO1 loading (**Fig. 7h**). In Arabidopsis, trans-acting sRNAs (tasiRNAs) are processed from transcripts of a specific locus (TAS3) mainly into 21-nt-siRNA by DCL4 [76]. As the TAS3 locus had been previously identified in Spirodela [32], we

scored for matching sRNAs. In Spirodela, abundant 22-nt siRNAs account for about half of TAS3 tasiRNAs (**Supplemental Fig. S34**). Given the similar 21-22-nt siRNA patterns observed between TEs, miRNA-triggered secondary tasiRNA and IR-derived siRNAs, PTGS on TEs is likely the result of of: (i) a shared miRNA targeting event to initiate 2ry siRNAs, shown to regulate some TEs in other plants [58,162], or; (ii) direct processing of RNA hairpins commonly present in LTRs [163].

Hence, Spirodela is capable of processing 22-nt siRNAs in the absence of DCL2, allowing to conclude that 21-22-nt siRNAs mapped to TEs are *bona-fide* siRNAs and that control of TEs in Spirodela combines transcriptional and post-transcriptional silencing mechanisms.

Discussion

Based on high resolution genomic information of a duckweed species during its clonal reproduction, we have provided a detailed analysis of TE content, of the epigenetic configuration at TEs, and several components expected to contribute to the silencing pathways that control TE activity in other angiosperms. Our results reinforce the idea that, with few exceptions, the Spirodela genome encodes only a reduced set of both PTGS and TGS components compared to Arabidopsis, some of which are absent or display a reduced set of paralogues, partially accounting for its reduced levels of DNA methylation and other epigenetic marks generally associated with TEs. The unique epigenome of Spirodela is the result of the loss of DNA methylation, and reduced levels of H3K9me2 associated to it, in highly abundant degenerated TE remnants scattered across the genome which remain nonetheless associated to the heterochromatic mark H3K9me1. In contrast, potentially intact or recently active TEs display all the epigenetic signatures associated with silencing in angiosperms. These changes in epigenetic regulation and silencing might be a consequence of the simplified morphology or reduced tissue and cell types in duckweeds compared to other angiosperms [36,39,45,164,165]. Alternatively, it could be a consequence of their different lifestyle or propagation mode.

Spirodela displays a broad range of adaptations to their free-floating aquatic lifestyle, such as the loss of complete stomatal closure response to prevent water loss [166]. Besides TE control, RdDM plays a role in the regulation of gene expression and physiological adaptation to several abiotic stresses, such as heat, drought, salinity, or

low environmental humidity [20,167,168]. Given that water supply is not limiting in
aquatic environments, and their higher stability to fluctuations in these parameters
compared to terrestrial ones, RdDM activity in adult somatic tissues to respond to
stress might have become less relevant in Spirodela.

Alternatively, but not mutually exclusive, the epigenetic landscape in the
Spirodela genome might be a consequence of their reproduction strategy. Asexual
reproduction has been linked to lower TE activity and content. Transposons require
sexual reproduction to spread horizontally throughout populations, instead of being
locked in vertical clonal lineages where they might become extinct if their proliferation
becomes detrimental for the host without increased purifying selection of new TE
insertions facilitated by sex [169–172]. Furthermore, RdDM activity over short TE
remnants, leading to the production of 24-nt siRNAs, is seen as an immunity memory
against mobilization or invasion by related intact TEs [173], akin to the piRNA pathway
in animals [174]. Thus, TEs potentially being less active during Spirodela clonal
propagation might have led to the loss of DNA methylation, RdDM expression ,and
lack of components required for its activity during vegetative development such as
SHH1 and CLSY1/2 [31] over TE fragments. Maintenance of heterochromatin at such
repetitive TEs remnants through H3K9me1 independently of DNA methylation ensures
proper chromatin organization, preventing their interference with transcriptional and
RNA processes [9,11,12,175] or causing genome instability through illegitimate
recombination [176,177]. Deeper characterization of the chromatin structure (e.g.,
chromatin accessibility) and landscape (additional histone modifications, histone
variants …), together with functional studies, might shed light onto the regulation of
H3K9me1 and its role in shaping heterochromatin and TE regulation in Spirodela and
other plants.

Furthermore, in contrast to vegetative propagation in other plants where clones
originate from differentiated organs such as stems, roots or leaves, with tissue-specific
and developmentally regulated TEs, new individuals in duckweeds originate from a
dedicated group of stem cells [42]. This might have further favored a relaxation of
epigenetic and RdDM-mediated TE control in somatic tissues which do not participate
in vegetative reproduction in Spirodela. Nevertheless, potentially intact TEs are under
several layers of epigenetic control, including RdDM, suggesting the pathway might
be tissue-specific or developmentally regulated.

In contrast to stem cells in other angiosperms, contained in a well-organized shoot apical meristem (SAM) that generates new organs [178], meristems in Spirodela and all duckweeds are formed by unstructured groups of few undifferentiated cells, which generate new meristems that develop into daughter fronds. [42,43,179–181]. In order to maintain genome integrity along the clonal lineage, RdDM might be expressed in the few stem cells within the budding pockets or during the developmental window between the formation of a new meristem and its development into a new frond with its own stem cells.

Furthermore, loss of expression of RdDM and weak CMT3/ZMET activity in somatic tissues, in combination with the absence of CMT2, during development will suffice to explain the global loss of mCHH and low levels of mCHG respectively, which in some instances, might permit transcription of some TEs that then triggers PTGS. Alternatively, transient relaxation of TE control and TE expression has been shown to take place in Arabidopsis meristematic stem cells during developmental transitions [182]. A similar phenomenon taking place during the development of new meristems and fronds might as well explain the paradox of observed expression from TEs covered in silencing marks and their apparent simultaneous regulation by TGS and PTGS. Further tissue- and cell-specific investigation of epigenetic parameters will be able to address these hypotheses.

Nonetheless, although infrequent, Spirodela does flower occasionally [47] and its population genetic structure shows signs of sexual reproduction [183,184]. Hence, RdDM might still be relevant and active during sexual reproduction as suggested by the presence of CLSY3, which recruits RdDM to TEs during flowering in Arabidopsis [31].

Adding to the unique epigenetic regulation of TEs in Spirodela here described, one of the intriguing results of our study is the constatation that, despite lack of DCL2, Spirodela can produce 22-nt siRNAs nearly as abundantly as 21-nt siRNAs, not only from TEs but also from endogenous and exogenous sources of dsRNA. As these sources of siRNAs are generally targets of DCL4 in plants [151,185,186], we consider processing dsRNA into 22-nt siRNAs by Spirodela DCL4 as the most obvious explanation. In Arabidopsis, DCL2 and its capacity to process dsRNA into 22-nt siRNAs have been proposed to play a dual role in antiviral immunity. On one hand its processing of viral dsRNA acts as an innate immunity sensor [187,188]. On the other hand, 22-nt siRNAs in plants guide the recruitment of RDRs to boost siRNAs

production from their targets and amplify silencing [189–192] . Although future genetic
and biochemical characterization of Spirodela DCLs will be required to understand the
factors responsible for the biogenesis of 22-nt siRNAs, their presence of independently
of DCL2 and the lack of several other RNAi factors associated with antiviral defense
in angiosperms raises several intriguing questions about RNA-based defense
mechanisms in duckweeds.

Spirodela might represent a unique model to investigate not only transposon
dynamics and the epigenetic and genetic consequences of long periods of asexual
reproduction but also non-canonical PTGS pathways.

Conclusions

By investigating TE silencing landscape during the clonal reproduction phase
of the angiosperm *Spirodela polyrhiza*, we have shown that despite the lack of
ubiquitous expression of RdDM and loss of DNA maintenance over degenerated TE-
relics, certain heterochromatic marks such as H3K9me1 can be maintained
independently of DNA methylation to ensure proper genome stability and chromatin
organization. Indicating that H3K9me1 and H3K9me2 might be differentially regulated
and play independent roles in heterochromatin biology. Potentially functional TEs are,
however, targets of several silencing mechanisms, including RdDM and PTGS,
probably due to tissue-specific regulation of such pathways during Spirodela clonal
reproduction. Interestingly, despite the absence of a DCL2 orthologue, Spirodela
produced 22-nt siRNAs from TEs and other sources of dsRNA, which has implications
not only for TE silencing but in antiviral immunity as well. Hence, Spirodela represents
a unique model system to gain further insights into PTGS and TGS pathways in plants.

Methods

Spirodela culture and growth conditions:
***Spirodela polyrhiza*.** Spirodela was grown on half-strength Schenk and Hildebrandt
(½ SH, Duchefa Biochemie, #S0225) liquid medium in Magenta™ GA-7 containers
(Sigma Aldrich, #V8505), culture dishes (Greiner Bio-One #664160), 6-wells culture
plates (Greiner Bio-One #657185), or in 1L glass beakers (covered with a sterile
plastic lid), all sealed with Leucopore tape (Duchefa Biochemie #L3301), under long-

day conditions (16h light /8h dark) at 21°C with a light intensity of 85 µM m^{-2} s^{-1}. The

Spirodela polyrhiza #9509 clone was obtained from the Landolt collection (Zürich,

Switzerland), recently transferred to the CNR-IBBA_MIDW collection (Milano, Italy:

https://biomemory.cnr.it/collections/CNR-IBBA-MIDW).

Further information regarding Spirodela sterilization for axenic cultures and long-term

storage, Arabidopsis and *N. benthamiana* material information and culture can be

found within Supplementary Methods in the Supplementary Information files.

Genome sequencing and gene annotation:

Genomic DNA from Spirodela was purified using Dneasy® Plant mini kit (Qiagen

#69104) following the manufacturer's instructions. Library preparation was carried

using NEBNext® Ultra™ II DNA Library Prep Kit for Illumina® (NEB #E7103).

Sequencing was performed on an Illumina HiSeq4A to produce paired-end reds of

125 bp. The Landolt Sp9505 genome was generated by pseudo-assembly with

MaSuRCA [193] (version 3.4.1, default parameters), using as scaffold the

Sp9509_Oxforrd_v3 reference assembly[50]. The pseudo-assembly was polished

using POLCA [194]. Variation, between the two assemblies, was further evaluated

using GAKT[195] (version 4.0.1.2), in particular with HaplotypeCaller [196] (--

standard-min-confidence-threshold-for-calling 30). Variants identified were

summarize into a table using GAKT VariantsToTable (-F CHROM -F POS -F TYPE -

GF AD). Gene annotations were transferred from the latest annotated version of

Sp9509 [39] obtained from (https://www.lemna.org). Annotation transfer was

performed using LiftOff (v1.6.3) [197] using default parameters.

Transposon annotation:

The annotation of transposable elements (TEs) in Spirodela polyrhiza was performed

by combining EDTA [52] and RepeatModeler2 [53] tools. To ensure accuracy, both

tools were run in triplicate and the resulting annotation libraries were aligned and

merged. The classification of TEs was further refined using DeepTE [54] (-sp P). The

presence of the protein domain for LTR-RT was evaluated using TEsorter [198]

(version 1.3, -db rexdb-plant -st nucl).

Transcriptome analysis:

To extract RNA from Spirodela, 100 mg of plant material was flash-frozen upon harvesting and ground. The samples were homogenized in 1 mL of TRIzol™ (ThermoFisher, #15596018) using Silamat S6 (Ivoclar Vivadent). RNA was extracted according to manufacturer's instructions. RNA library preparation was performed using the NEBNext® Ultra™ II RNA Library Prep Kit for Illumina® (NEB #E7775) or the SMRTbell Prep Kit 3.0 for Illumina and Iso-Seq respectively. Illumina sequencing was performed on HiSeq5A to generate 125 bp paired-end reads. Long read sequencing was performed on PacBio® Sequel® II. For Illumina data, base calls were performed using bcl2fastq (v2.17). RNA-seq reads were trimmed using trimgalore (v0.6, --illumina -phred33 -stringency 1 --fastqc --length 20). Trimmed reads were mapped using hiSat2 (v2.1.0). Gene and transposable element expression levels were quantified separately using Kallisto [199] (v0.46) with default parameters. The resulting expression levels (measured in TPM) were then averaged across the three replicates. For Arabidopsis seedlings, data was obtained from (GSM6892967, GSM6892968, GSM6892969).Iso-Seq long reads were processed using nf-core/isoseq (https://github.com/nf-core/isoseq/tree/1.0.0). High quality transcripts were then mapped to Spirodela genome using minimap2 [200] (v2.17, default parameters).

DNA methylation analysis:

For EMseq, DNA extraction was performed as above. Libraries were prepared using the NEBNext® Enzymatic Methyl-seq Kit (NEB, #M7634) following the manufacturer's instructions. Sequencing was performed on an Illumina NovaSeq SP to produce paired-end reads of 150 bp. Briefly, Briefly, basecalls were performed using bcl2fastq (v2.17) and sequenced reads were quality filtered and adaptor trimmed using TrimGalore version 0.6.2 (https://github.com/FelixKrueger/TrimGalore). Enzymatic-converted reads were aligned either to the TAIR10 genome or Spirodela polyrhiza genome using Bismark version 0.22.2 with *bismark* (--non_directional -*q* --score-min *L,0,−0.4*) [201] to generate coverage files per cytosine. Conversion rates were assessed by mapping (samtools v1.9) the reads to the unmethylated lambda phage DNA, with the same parameters used for the genomes BAM files containing clonal deduplicated and uniquely mapped reads were then used to extract weighted methylation rates at each cytosine as described in [202]. Replicates were merged into a single bedFile using the BEDtools [203] merge function and only cytosines with

coverage with ≥4 were considered for further analyses. 5mC was calculated using
ViewBS [204] (v0.1.11) (https://github.com/xie186/ViewBS) using default parameters.

Small RNA extraction:

Total RNA: For small RNA extraction from total RNA and TraPR input fraction, 10µg
of total RNA (as described above) or RNA extracted as described above from TraPR
inputs was separated on a 15% denaturing polyacrylamide-urea gel. For size
reference, 20µl of microRNA marker (New England Biolabs, #N2102S) were loaded in
the same gel. Following gel staining with peqGreen (Peqlab, #732-3196), and using
the marker as reference, small RNAs ranging from 17-to-25 nt were excised and
recovered using the ZR small-RNA PAGE Recovery Kit (Zymo Research, #R1070).

TraPR: Purification of Argonaute-loaded small RNAs was performed according to the
protocol described in [55]. In brief, 100mg of flash-frozen plant tissue was ground to
fine powder using a Silamat S6 (Ivoclar Vivadent) twice for 3s in 1.5 mL
microcentrifuge tubes with ~20 1 mm ø glass beads (Carl Roth, #A554.1). Frozen
powder was incubated with TraPR lysis buffer and the homogenate loaded into the
TraPR columns. TraPR eluates were collected and small RNAs extracted by adding
and mixing 1 volume of acidic Phenol:Chloroform:Isoamyl alcohol (50:49:1, pH 4.5-5)
(Carl Roth, #X985.1). After centrifugation (12000g for 10 min at 4ºC) RNA was
precipitated from the aqueous phase with 1 volume of isopropanol and 10% (v/v) of
sodium actetate 3M pH 5.2, overnight at -20°C. RNA was collected by centrifugation
(12000g for 30 min at 4ºC) and the resulting pellet was washed with 75% ethanol and
resuspended in 10-20 µL RNase free water.

Small RNA library preparation and analysis:

The libraries were prepared according to [205]. In summary, small RNAs purified using
TraPR or total small RNA as described above, were ligated to 3′ barcoded DNA
adapters using truncated T4 RNA ligase 2 (New England Biolabs, #M0373). These
fragments were separated in an 12% denaturing polyacrylamide-urea gel, excised,
and purified using ZR small-RNA PAGE Recovery Kit (Zymo Research, # R1070).
The elution was then ligated to 5′ barcoded RNA adapters using T4 RNA ligase 1 (New
England Biolabs, #M0204). To reduce ligation biases, the sequence of both adaptors
included 4 random nucleotides at the ends. The resulting fragments were reverse
transcribed and amplified by PCR using primers compatible with the Illumina Platform.

The size distribution of the libraries was assessed using the 5200 Fragment Analyser

System (Agilent, #M5310AA). Sequencing was performed on an Illumina NextSeq550

to produce single-end reads of 75 bp. Basecalls were performed using bcl2fastq

(v2.17). Reads were trimmed, quality checked using Trimmomatic [206] (v0.39).

Trimmed reads were mapped to Arabidopsis TAIR10 and Spirodela genomes, and

RUBY or *hpScarlet* sequences using Bowtie2 [207] (v1.2.2; -e 50 -a -v 0 --best --strata

--nomaqround -y --phred33-quals --no-unal --sam). Mapped reads were then

normalized using bamCoverage [208] (Deeptools v,--normalizeUsing CPM --binSize 1

--smoothLength 10 --maxFragmentLength 30). To visualise strand, mapped reads

were split using samtools view [209] (samtools v1.9, using -h -b -F 16 or -h -b -f 16

options).

Protein extraction and western blot:

Protein extraction: For protein extraction from Arabidopsis and Spirodela, plant

material was flash-frozen upon harvesting and ground with a Silamat S6 as described

in the RNA extraction section. For *N. benthamiana*, leaf tissue was grounded in liquid

nitrogen using mortar and pestle. 100mg of ground tissue was mixed and

homogenized for 5 min. with 500µl of extraction buffer (0.7 M sucrose, 0.5 M Tris-HCl,

pH 8, 5 mM EDTA, pH 8, 0.1 M NaCl, 2% b-mercaptoethanol) and cOmplete^{TM} mini

EDTA-free protease inhibitor cocktail (Roche, #04693159001) following

manufacturer's instructions. 1 volume of TRIS-buffered phenol (pH 8) was added and

incubated with agitation for 5 minutes. The upper phenolic phase was recovered after

centrifugation (12000*g* for 10 min at 4ºC) and proteins precipitated with 5 volumes of

0.1 M ammonium acetate dissolved in methanol at -20ºC overnight. Following

centrifugation (5000*g* for 15 min at 4ºC), the resulting protein pellet obtained was

washed twice with 0.1 M ammonium acetate dissolved in methanol and resuspended

in 50-100 µL of resuspension buffer (3% SDS, 62.3 mM Tris-HCl, pH 8, 10% glycerol).

Protein blots: Western blot was carried following standard denaturing SDS-PAGE

and blotting procedures [210]. Unless otherwise specified, 150µg of protein were

resolved on SDS-polyacrylamide gels, transferred onto Inmobilon-P PVDF membrane

(Millipore, # IPVH00010) and incubated with the appropriate antibodies (see below) in

PBS with 0.1% Tween-20 and 5% non-fat dried milk. The primary antibody was

incubated overnight at 4ºC, and the HRP-conjugated secondary antibody for 1 hour at

room temperature. Detection was performed using Clarity Max™ Western ECL

Substrate (Biorad, #1705062), and imaged with the ChemiDoc Imaging system (Bio-

rad). Following detection, loading was verified by coomassie staining of membranes.

Primary antibodies used for western blot: monoclonal (mouse) anti-HA tag [12CA5]

(Abcam, #ab1424), dilution 1:10000. Polyclonal (rabbit) anti-H3 (Abcam #ab1791),

dilution 1:10000. Polyclonal (rabbit) anti-H3K9me1 (Abcam #ab8896), dilution 1:2000.

Monoclonal (mouse) anti-H3K9me2 (Abcam #ab1220), dilution 1:2000.

Secondary antibodies used for western blot: Polyclonal (goat) anti-mouse HRP-

conjugated (ThermoFisher, #62-6520), dilution 1:10000. Polyclonal (goat) anti-rabbit

HRP-conjugated (ThermoFisher, #65-6120), dilution 1:10000.

Chromatin immunoprecipitation (ChIP):

Chromatin immunoprecipitation (ChIP) was conducted on 2 g of Arabidopsis 10-day-

old seedlings (WT Col-0) and *Spirodela polyrhiza* #9509 fronds. Following standard

procedures (see detailed method in Supplementary Methods in the Supplementary

Information files) using the following antibodies: anti-H3 (rabbit polyclonal, Abcam,

#ab1791), anti-H3K27me3 (rabbit polyclonal, Millipore, #07–449), anti-H3K4me3

(rabbit polyclonal, Abcam, #ab8580), anti-H3K9me1 (rabbit polyclonal, Abcam,

3ab8896), or anti-H3K9me2 (mouse monoclonal, Abcam, #ab1220).

ChIP-seq library prep and data processing:

For ChIP-seq, libraries were prepared using the NEBNext® Ultra™ II DNA Library

Prep Kit for Illumina® (NEB, #E7103) according to the manufacturer's instructions.

Sequencing was performed on an Illumina NovaSeq S4 to produce paired-end reads

of 150 bp. The raw BAM files were converted to fastq using BEDTools [203] (v2.27.1,

default parameters) for quality assessment of the sequenced samples. FastQC

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/)   (v0.11.8,   default

parameters) was used to generate quality reports for all sequencing data. Basecalls

were performed using bcl2fastq (v2.17). Trimming of reads was performed using

trim_galore (https://zenodo.org/doi/10.5281/zenodo.5127898) (v0.6.5, --dont_gzip --

stringency 1 --fastqc --length 5'). The aligned reads were then mapped to either the

TAIR10 Arabidopsis genome or Spirodela polyrhiza genome using Bowtie2 [207]

(v2.4.1, default parameters). Duplicate reads were removed using Picard

(http://broadinstitute.github.io/picard/) (v2.22.8, default parameters). Mapped reads

were then normalized using bamCoverage (Deeptools v3.3.1). Correlations between

ChIP samples were evaluated using deepTools [208] (v3.1.2, default parameters). The

bamCompare function from deepTools was used to normalize ChIP samples to Input

or H3 and produce log2 ratio (ChIP/H3) bigwig files.

Agrobacterium-mediated transient expression:

N. benthamiana: Agrobacterium-mediated transient expression in *N. benthamiana*

was performed as in [211]. In brief, *A. tumefaciens* strain GV3101 harbouring the

vectors described below were cultured at 28°C in LB medium with appropriate

antibiotics. After 36 hours, cultures were pelleted at 2000g, 10 minutes at 20°C and

resuspended to an OD_{600} of 0.5 in filtered sterilized agroinfiltration buffer: 2mM MES,

50mM NaH_2PO_4, 0.5% (w/v) glucose, pH 5.6. Acetosyringone was added prior to use

at 100 µM. After 1 hour of incubation at room temperature in darkness, the bacterial

solutions were infiltrated into the abaxial side of young *N. benthamiana* true leaves,

using a 1ml syringe without needle. At 3 or 5 days after infiltration (dpi), leaf discs were

collected, flash-frozen in liquid nitrogen and ground in liquid nitrogen using mortar and

pestle. The ground samples were stored at -80C.

Spirodela polyrhiza: For Agrobacterium-mediated transient expression in Spirodela,

Agrobacterium strain EHA105 was used. Bacteria cultures were prepared as for *N.*

benthamiana infiltrations with the following modifications: Bacteria was resuspended

to OD_{600} 0.8 in Spirodela agroinfiltration media (10 mM MgCl2, 5% sucrose (w/v), pH

5.6). Spirodela fronds were prepared as it follows: Cultures were initiated from axenic

stocks by inoculating fronds into 2L glass beakers containing 0.5-1L of ½ SH media,

covered with the lid of a 145 x 20 mm polystyrene culture dish (Greiner Bio-One

#639102) sealed with 2.5 cm wide Leucopore tape (Duchefa Biochemie #L3301).

Cultures were grown to 50-80% confluence (as surface occupancy). Prior to

agroinfiltration, fronds were pretreated as in [160] by transferring them to an ice-cold

20 mg/L L-Glutamine (PanReac AppliChem #A1420) solution for 20 min. Following

pretreatment, fronds were infiltrated with the *Vir*-induced Agrobacterium solution either

(i) manually, at the budding pouch area with a 1 mL syringe without needle by gently

applying pressure with the plunger on the upper side of fronds or, (ii) trough vacuum

infiltration by transferring the fronds to 100 mL agrobacterium solution complemented

with 0.02% (v/v) Silwet L-77 (Kurt-Obermieier GmbH #7060-10) in a 200 mL beaker in

a vacuum desiccator. -70 kPa vacuum was applied twice for 10 min. Following either

manual or vacuum agroinfiltration, excess of bacteria was removed by rinsing the

fronds in ddH$_2$O and plants transferred to 100 x 20 mm culture dishes (Greiner Bio-

One #664160) containing 30-50 mL ½ SH media + 100 µM acetosyringone for a 3 day

co-culture. 4 days post-infiltration (dpi) transfer fronds to plates containing ½ SH

media. At 7 dpi agrobacterium elimination was initiated by or replacing the media with

½ SH + 250 µM ticarcillin (Duchefa #T0190.0010, lab stock at 250 mg/mL in H2O) +

300 µM cefotaxime (Duchefa #C0111.0005, lab stock at 300 mg/mL in H2O) for a

week. Plants were cultured with regular (1/week) ½ SH media replacements. See

Supplementary Fig. S36 for a graphic depiction of the protocol and workflow.

Small RNA blot analysis:

AGO-loaded small RNAs were purified from 100mg of starting material using TraPR

columns, as described above. The resulting small RNAs were mixed with an equal

volume of 2X Novex^{TM} TBE-Urea sample buffer (ThermoFisher, #LLC6876),

separated on a 17.5% polyacrylamide-urea gel, transferred onto Hybond-NX

membrane (Cytiva, #RPN303T) and chemically crosslinked (12.5M 1-methylimidazol

(Merk, #M50834), 31.25mg/mL N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide

hydrochloride (Merk, #E7750) as in [212]. Probes produced from PCR templates were

radiolabelled with [α-^{32}P]-dCTP (Hartmann Analytic, #SRP-305) using Prime-it II

Random Primer Labelling kit (Agilent, #300385) and oligonucleotides used as a probe

were radiolabelled with [γ-^{32}P]-ATP (Hartmann analytic #SRP-501), using T4

Polynucleotide kinase (Thermofisher, #EK0031) (see Supplementary figures for a list

of oligonucleotides used). The probes were incubated in PerfectHyb™ Plus

Hybridization Buffer (Sigma Aldrich, #H7033) and washed with 2XSSC, 0.1%SDS.

The signal was detected with the Amersham^{TM}Typhoon^{TM} (Cytivia). Probes and

oligonucleotides used are listed in Supplementary figures.

Immunoprecipitation of Flag-tagged AGO4:

100-150mg of *N. benthamiana* ground tissue was homogenized in 1 ml of IP buffer

(50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 10% glycerol, 0.1% NP-40) containing 2 µM

MG-132 and cOmplete^{TM} mini EDTA-free protease inhibitor cocktail (Roche,

#04693159001) for 30 minutes at 4 °C. After lysis, debris were removed by

centrifugation (12000g for 10 min at 4ºC). An aliquot of lysate was kept as input, while

the remainder was incubated with ANTI-FLAG^R M2 Magnetic beads (20µl of

slurry/sample) (Sigma-Aldrich, #M8823) for 3 hours at 4°C, on a orbital shaker. After

incubation, the resin was separated from supernatant in a magnetic rack (DynaMag-

5, Thermofisher, #12321D) and washed with 1 ml of IP buffer for 20 minutes at 4°C,

for 3 times. After the final wash, 100µl of PBS were added and 10µl put aside, used to

check immunoprecipitation quality by western blot by adding Laemli buffer and boiling

the sample for 10 min. at 95ºC. The remaining 90µl was used to analyze the small

RNA composition upon extraction using TRIzol™ (ThermoFisher, #15596018). The

isolated small RNAs were used to generate libraries and sequenced as described

above. TraPR purified small RNAs (described above) from not-infiltrated plants were

used to assess *N. benthamiana* small RNA profile.

Visualization and Statistical analysis:

Data visualization files were generated according to each specific sections. Statistical

tests were performed using R-dependent ggpubr (v0.6.0)

(https://rpkgs.datanovia.com/ggpubr/), statistical test performed are indicated

accordingly on each figure legend figure. Plots under R (v4.2.2) (https://www.r-

project.org) were generated using Circlize [213] (v0.4.15), ggplot2 [214] (v3.4.4),

profileplyr (v1.14.1)

(https://bioconductor.org/packages/release/bioc/html/profileplyr.html), ViewBS [204]

(v0.1.11), or pheatmap (v 1.0.12)

(https://rdocumentation.org/packages/pheatmap/versions/1.0.12).

Supplementary Methods:

Further information regarding plant culture, plant imaging, phylogenetic analysis, mass

spectrometry of histone PTMs, chromatin immunoprecipitation, DAPI staining of nuclei

and molecular cloning procedures can be found within the Supplementary Methods in

the Supplementary Information files.

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

An inventory of all the raw data used to generate each figure panel (as well as
those in Supplementary information) along with all raw image files (including RNA,
protein blots and agarose gels), DNA and protein sequences, alignments, and
phylogenetic tree files can be found in Zenodo (www.zenodo.org) under the doi:
10.5281/zenodo.10911533. NGS data generated for this study are accessible NCBI
SRA under ID number PRJNA1095698, submission ID SUB14339214. Publicly
available Arabidopsis seedling transcriptome from [74] is found under the accession
number GSM6892968. Ready-to-use IGV genome browser tracks for all the NGS
datasets generated here is available in the associated Zenodo dataset. The mass
spectrometry proteomics data have been deposited to the ProteomeXchange
Consortiumvia the PRIDE partner repository with the dataset identifier PXD050443.
Additional Excel files with the MS processed data can be found in the associated
Zenodo dataset. All plasmids generated have been submitted to Addgene and can be
retrieved under the following IDs: p*35S*:*FHA*-At*AGO4*_gDNA (#216838); p*35S*:*FHA*-
Sp*AGO4a*_gDNA (#216841); p*35S*::*FHA*-Sp*AGO4a*_cDNA (#216842). The following
plasmids previously originated in [161] and [215] were obtained from Addgene:
p*35S*:*RUBY* (#160908); p*ZmUbq*:*RUBY* (#160909); p*35S*:*GFP-GUS* (#167122).
p3At*UBQ*:*hpScarlet* was a gift from Marco Incarbone (Max-Planck-Institute of Plant
Physiology, Golm, Germany).

Competing interests

997   The authors declare that they have no competing interests.

Funding

1000

This work was supported by the Gregor Mendel Institute of the Austrian Academy of Sciences core funding attributed to A.M-O.

Author contributions

AM-O and RD conceived and designed the study. RD and VB-B performed most experiments. AM-O and RD performed identification and phylogenetic analysis of silencing components in Spirodela. VB-B cloned and carried out transient AGO4 expression experiments and prepared small RNA libraries. DB-A made nuclei DAPI stainings. JMV-G performed western blot analysis of HPTMs. AM-O, DB-A, and AP-M optimized agroinfiltration-mediated transient expression in duckweeds. AP-M carried out transient expression experiments in Spirodela. VB-B and RE cultured and maintained Spirodela stocks and axenic cultures and provided technical support. RD performed computer and statistical analysis. AM-O and RD analyzed the data and wrote the manuscript.

Acknowledgements

We thank current and past members of the Marí-Ordóñez group and colleagues from the Gregor Mendel Institute and the Vienna BioCenter (VBC) campus for fruitful discussions, ideas, and feedback. We are specially thankful to Ortrun Mitettelsten Scheid for critical feedback and comments on the manuscript. We would also like to thank Klaus Appenroth from the Friedrich Schiller University of Jena for useful advice on duckweed cultivation and storage, Thomas Grentzinger for technical assistance with the TraPR procedure, and Marco Incarbone from the Max-Planck-Institute of Plant Physiology in Golm for kindly providing the pAt*UBQ*:*hpScarlet* plasmid. We are grateful to the Vienna BioCenter Core Facilities (VBCF): NGS for DNA, EM-seq, PacBio, RNA-seq and ChIP-seq library preparation and sequencing, including sequencing of small RNA libraries; the IMP/IMBA/GMI Proteomics Facility, in special Ines Steinmacher for sample preparation, mass-spectrometry using the VBCF instrument pool, and Richard Imre and Elisabeth Rothinger for assistance with data

1035   analysis of histone PTMs; the Plant Science Facility for assistance with plant work; the

1036   IMP Molecular Biology Service for providing reagents and Sanger sequencing. Lastly,

1037   we would like to thank the VBC in-house COVID-19 testing facility for enabling a safe

1038   working environment during the pandemic.

References

1. Wells JN, Feschotte C. A Field Guide to Eukaryotic Transposable Elements. Annu Rev Genet. 2020;54:1–23.

2. Biémont C, Vieira C. Genetics: junk DNA as an evolutionary force. 2006 Oct 5;521–4.

3. Feschotte C. Transposable elements and the evolution of regulatory networks. Nature. 2008;9:397–405.

4. Allshire RC, Madhani HD. Ten principles of heterochromatin formation and function. Nature Reviews Molecular Cell Biology. 2018;19:229–44.

5. Henikoff S, Smith MM. Histone Variants and Epigenetics. Cold Spring Harb Perspect Biol. 2015;7:a019364.

6. Jamge B, Lorković ZJ, Axelsson E, Osakabe A, Shukla V, Yelagandula R, et al. Histone variants shape chromatin states in Arabidopsis. eLife. 2023;12:RP87714.

7. Blumenstiel JP. Birth, School, Work, Death, and Resurrection: The Life Stages and Dynamics of Transposable Element Proliferation. Genes. 2019;10:336.

8. Ito H, Gaubert H, Bucher E, Mirouze M, Vaillant I, Paszkowski J. An siRNA pathway prevents transgenerational retrotransposition in plants subjected to stress. Nature. 2011;472:115–9.

9. Kim MY, Zilberman D. DNA methylation as a system of plant genomic immunity. Trends Plant Sci. 2014;19:320–6.

10. Zervudacki J, Yu A, Amesefe D, Wang J, Drouaud J, Navarro L, et al. Transcriptional control and exploitation of an immune-responsive family of plant retrotransposons. The EMBO journal. 2018;37:e98482.

11. Sammarco I, Pieters J, Salony S, Toman I, Zolotarov G, Placette CL. Epigenetic targeting of transposon relics: beating the dead horses of the genome? Epigenetics. 2022;17:1331–44.

12. Ilık İA, Glažar P, Tse K, Brändl B, Meierhofer D, Müller F-J, et al. Autonomous transposons tune their sequences to ensure somatic suppression. Nature. 2024;1–9.

13. Houben A, Demidov D, Gernand D, Meister A, Leach CR, Schubert I. Methylation of histone H3 in euchromatin of plant chromosomes depends on basic nuclear DNA content. Plant J. 2003;33:967–73.

14. Seymour DK, Koenig D, Hagmann J, Becker C, Weigel D. Evolution of DNA Methylation Patterns in the Brassicaceae is Driven by Differences in Genome Organization. PLoS Genet. 2014;10:e1004785.

15. Sigman MJ, Slotkin RK. The First Rule of Plant Transposable Element Silencing: Location, Location, Location. Plant Cell. 2016;28:304–13.

16. Wyler M, Stritt C, Walser J-C, Baroux C, Roulin AC. Impact of transposable elements on methylation and gene expression across natural accessions of Brachypodium distachyon. Genome Biol Evol. 2020;12:evaa180-.

17. Klein SP, Anderson SN. The evolution and function of transposons in epigenetic regulation in response to the environment. Curr Opin Plant Biol. 2022;69:102277.

18. Moazed D. Small RNAs in transcriptional gene silencing and genome defence. Nature. 2009;457:413–20.

19. Malone CD, Hannon GJ. Small RNAs as Guardians of the Genome. Cell. 2009;136:656–68.

20. Erdmann RM, Picard CL. RNA-directed DNA Methylation. Plos Genet. 2020;16:e1009034.

21. Mirouze M, Reinders J, Bucher E, Nishimura T, Schneeberger K, Ossowski S, et al. Selective epigenetic control of retrotransposition in Arabidopsis. Nature. 2009;461:427–30.

22. Tsukahara S, Kobayashi A, Kawabe A, Mathieu O, Miura A, Kakutani T. Bursts of retrotransposition reproduced in Arabidopsis. Nature. 2010;461:423–6.

23. Stroud H, Do T, Du J, Zhong X, Feng S, Johnson L, et al. Non-CG methylation patterns shape the epigenetic landscape in Arabidopsis. Nat Struct Mol Biol. 2014;21:64–72.

24. He L, Zhao C, Zhang Q, Zinta G, Wang D, Lozano-Durán R, et al. Pathway conversion enables a double-lock mechanism to maintain DNA methylation and genome stability. Proc Natl Acad Sci. 2021;118:e2107320118.

25. Osakabe A, Jamge B, Axelsson E, Montgomery SA, Akimcheva S, Kuehn AL, et al. The chromatin remodeler DDM1 prevents transposon mobility through deposition of histone variant H2A.W. Nat Cell Biol. 2021;23:391–400.

26. Bourc'his D, Voinnet O. A Small-RNA Perspective on Gametogenesis, Fertilization, and Early Zygotic Development. Science. 2010;330:617–22.

27. Marí-Ordóñez A, Marchais A, Etcheverry M, Martin A, Colot V, Voinnet O. Reconstructing de novo silencing of an active plant retrotransposon. Nature Genetics. 2013;45:1029–39.

28. Parhad SS, Theurkauf WE. Rapid evolution and conserved function of the piRNA pathway. R Soc Open Biol. 2019;9:180181.

29. Havecker ER, Wallbridge LM, Hardcastle TJ, Bush MS, Kelly KA, Dunn RM, et al. The Arabidopsis RNA-Directed DNA Methylation Argonautes Functionally Diverge Based on Their Expression and Interaction with Target Loci. THE PLANT CELL ONLINE. 2010;22:321–34.

30. Patel P, Mathioni SM, Hammond R, Harkess AE, Kakrana A, Arikit S, et al. Reproductive phasiRNA loci and DICER-LIKE5, but not microRNA loci, diversified in monocotyledonous plants. Plant Physiol. 2021;185:kiab001.

31. Zhou M, Coruh C, Xu G, Martins LM, Bourbousse C, Lambolez A, et al. The CLASSY family controls tissue-specific DNA methylation patterns in Arabidopsis. Nat Commun. 2022;13:244.

32. Michael TP, Bryant D, Gutierrez R, Borisjuk N, Chu P, Zhang H, et al. Comprehensive definition of genome features in Spirodela polyrhiza by high-depth physical mapping and short-read DNA sequencing strategies. Plant J. 2017;89:617–35.

33. Fourounjian P, Tang J, Tanyolac B, Feng Y, Gelfand B, Kakrana A, et al. Post-transcriptional adaptation of the aquatic plant Spirodela polyrhiza under stress and hormonal stimuli. Plant J. 2019;98:1120–33.

34. Cervantes-Pérez SA, Yong-Villalobos L, Florez-Zapata NMV, Oropeza-Aburto A, Rico-Reséndiz F, Amasende-Morales I, et al. Atypical DNA methylation, sRNA-size distribution, and female gametogenesis in Utricularia gibba. Sci Rep. 2021;11:15725.

35. Ibarra-Laclette E, Lyons E, Hernández-Guzmán G, Pérez-Torres CA, Carretero-Paulet L, Chang T-H, et al. Architecture and evolution of a minute plant genome. Nature. 2013;498:94–8.

36. Wang W, Haberer G, Gundlach H, Gläßer C, Nussbaumer T, Luo MC, et al. The Spirodela polyrhiza genome reveals insights into its neotenous reduction fast growth and aquatic lifestyle. Nat Commun. 2014;5:3311.

37. An D, Zhou Y, Li C, Xiao Q, Wang T, Zhang Y, et al. Plant evolution and environmental adaptation unveiled by long-read whole-genome sequencing of Spirodela. Proc Natl Acad Sci. 2019;116:18893–9.

38. Harkess A, Bewick AJ, Lu Z, Fourounjian P, Messing J, Michael TP, et al. Unusual predominance of maintenance DNA methylation in Spirodela polyrhiza. bioRxiv. 2020;2020.12.03.410332.

39. Ernst E, Abramson B, Acosta K, Hoang PTN, Mateo-Elizalde C, Schubert V, et al. The genomes and epigenomes of aquatic plants (Lemnaceae) promote triploid hybridization and clonal reproduction. bioRxiv. 2023;2023.08.02.551673.

40. Bog M, Appenroth K-J, Sree KS. Duckweed (Lemnaceae): Its Molecular Taxonomy. Front Sustain Food Syst. 2019;3:117.

41. Tippery NP, Les DH, Appenroth KJ, Sree KS, Crawford DJ, Bog M. Lemnaceae and Orontiaceae Are Phylogenetically and Morphologically Distinct from Araceae. Plants. 2021;10:2639.

42. Landolt E. Biosystematic investigations in the family of duckweeds (Lemnaceae) (Vol 2). The family of Lemnaceae a monographic study. ETH V des GI der, editor. Stiftung Ruebel, Switzerland: Veroeffentlichungen des Geobotanischen Instituts der ETH; 1986.

43. Lemon GD, Posluszny U. Comparative Shoot Development and Evolution in the Lemnaceae. Int J Plant Sci. 2000;161:733–48.

44. Lam E, Michael TP. Wolffia, a minimalist plant and synthetic biology chassis. Trends Plant Sci. 2022;27:430–9.

45. Ware A, Jones DH, Flis P, Chrysanthou E, Smith KE, Kümpers BMC, et al. Loss of ancestral function in duckweed roots is accompanied by progressive anatomical reduction and a re-distribution of nutrient transporters. Curr Biol. 2023;33:1795-1802.e4.

46. Pieterse AH. Is flowering in Lemnaceae stress-induced? A review. Aquat Bot. 2013;104:1–4.

47. Fourounjian P, Slovin J, Messing J. Flowering and Seed Production across the Lemnaceae. Int J Mol Sci. 2021;22:2733.

48. Ziegler P, Adelmann K, Zimmer S, Schmidt C, Appenroth K -J. Relative in vitro growth rates of duckweeds (Lemnaceae) – the most rapidly growing higher plants. Plant Biol. 2015;17:33–41.

49. Xu S, Stapley J, Gablenz S, Boyer J, Appenroth KJ, Sree KS, et al. Low genetic variation is associated with low mutation rate in the giant duckweed. Nat Commun. 2019;10:1243.

50. Hoang PNT, Michael TP, Gilbert S, Chu P, Motley ST, Appenroth KJ, et al. Generating a high-confidence reference genome map of the Greater Duckweed by integration of cytogenomic, optical mapping, and Oxford Nanopore technologies. Plant J. 2018;96:670–84.

51. Harkess A, McLoughlin F, Bilkey N, Elliott K, Emenecker R, Mattoon E, et al. Improved Spirodela polyrhiza genome and proteomic analyses reveal a conserved chromosomal structure with high abundance of chloroplastic proteins favoring energy production. J Exp Bot. 2021;72:2491–500.

52. Ou S, Su W, Liao Y, Chougule K, Agda JRA, Hellinga AJ, et al. Benchmarking transposable element annotation methods for creation of a streamlined, comprehensive pipeline. Genome Biol. 2019;20:275.

53. Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, et al. RepeatModeler2 for automated genomic discovery of transposable element families. Proc Natl Acad Sci. 2020;117:9451–7.

54. Yan H, Bombarely A, Li S. DeepTE: a computational method for de novo classification of transposons with convolutional neural network. Bioinformatics. 2020;36:4269–75.

55. Grentzinger T, Oberlin S, Schott G, Handler D, Svozil J, Barragan-Borrero V, et al. A universal method for the rapid isolation of all known classes of functional silencing small RNAs. Nucleic Acids Res. 2020;48:gkaa472-.

56. Teixeira FK, Heredia F, Sarazin A, Roudier F, Boccara M, Ciaudo C, et al. A Role for RNAi in the Selective Correction of DNA Methylation Defects. Science. 2009;323:1600–4.

57. Nuthikattu S, McCue AD, Panda K, Fultz D, DeFraia C, Thomas EN, et al. The Initiation of Epigenetic Silencing of Active Transposable Elements Is Triggered by RDR6 and 21-22 Nucleotide Small Interfering RNAs. PLANT PHYSIOLOGY. 2013;162:116–31.

58. Creasey KM, Zhai J, Borges F, Ex FV, Regulski M, Meyers BC, et al. miRNAs trigger widespread epigenetically activated siRNAs from transposons in Arabidopsis. Nature. 2014;508:411–5.

59. Feng S, Zhong Z, Wang M, Jacobsen SE. Efficient and accurate determination of genome-wide DNA methylation patterns in Arabidopsis thaliana with enzymatic methyl sequencing. Epigenetics Chromatin. 2020;13:42.

60. Niederhuth CE, Bewick AJ, Ji L, Alabady MS, Kim KD, Li Q, et al. Widespread natural variation of DNA methylation within angiosperms. Genome Biol. 2016;17:194.

61. Bewick AJ, Schmitz RJ. Gene body DNA methylation in plants. Curr Opin Plant Biol. 2017;36:103–10.

62. Mathieu O, Reinders J, Čaikovski M, Smathajitt C, Paszkowski J. Transgenerational Stability of the Arabidopsis Epigenome Is Coordinated by CG Methylation. Cell. 2007;130:851–62.

63. Cokus SJ, Feng S, Zhang X, Chen Z, Merriman B, Haudenschild CD, et al. Shotgun bisulphite sequencing of the *Arabidopsis* genome reveals DNA methylation patterning. Nature. 2008;452:215–9.

64. Reinders J, Wulff BBH, Mirouze M, Ordóñez AM, Dapp M, Rozhon W, et al. Compromised stability of DNA methylation and transposon immobilization in mosaic Arabidopsis epigenomes. Genes & development. 2009;23:939–50.

65. Gouil Q, Baulcombe DC. DNA Methylation Signatures of the Plant Chromomethyltransferases. PLoS Genet. 2016;12:e1006526.

66. Li X, Harris CJ, Zhong Z, Chen W, Liu R, Jia B, et al. Mechanistic insights into plant SUVH family H3K9 methyltransferases and their binding to context-biased non-CG DNA methylation. Proc Natl Acad Sci. 2018;115:E8793–802.

67. Fang J, Jiang J, Leichter SM, Liu J, Biswal M, Khudaverdyan N, et al. Mechanistic basis for maintenance of CHG DNA methylation in plants. Nat Commun. 2022;13:3877.

68. Stroud H, Greenberg MVC, Feng S, Bernatavichute YV, Jacobsen SE. Comprehensive Analysis of Silencing Mutants Reveals Complex Regulation of the Arabidopsis Methylome. Cell. 2013;152:352–64.

69. Yaari R, Noy-Malka C, Wiedemann G, Gershovitz NA, Reski R, Katz A, et al. DNA METHYLTRANSFERASE 1 is involved in mCG and mCCG DNA methylation and is essential for sporophyte development in Physcomitrella patens. Plant Mol Biol. 2015;88:387–400.

70. Bewick AJ, Niederhuth CE, Ji L, Rohr NA, Griffin PT, Leebens-Mack J, et al. The evolution of CHROMOMETHYLASES and gene body DNA methylation in plants. Genome Biol. 2017;18:65.

71. Bewick AJ, Ji L, Niederhuth CE, Willing E-M, Hofmeister BT, Shi X, et al. On the origin and evolutionary consequences of gene body DNA methylation. Proc Natl Acad Sci. 2016;113:9111–6.

72. Wendte JM, Zhang Y, Ji L, Shi X, Hazarika RR, Shahryary Y, et al. Epimutations are associated with CHROMOMETHYLASE 3-induced de novo DNA methylation. eLife. 2019;8:e47891.

73. Harkess A, Bewick AJ, Lu Z, Fourounjian P, Michael TP, Schmitz RJ, et al. Unusual predominance of maintenance DNA methylation in Spirodela polyrhiza. G3: Genes, Genomes, Genet. 2024;jkae004

74. Li Z, Wang M, Zhong Z, Gallego-Bartolomé J, Feng S, Jami-Alahmadi Y, et al. The MOM1 complex recruits the RdDM machinery via MORC6 to establish de novo DNA methylation. Nat Commun. 2023;14:4135.

75. Zhan J, Meyers BC. Plant Small RNAs: Their Biogenesis, Regulatory Roles, and Functions. Annu Rev Plant Biol.
2023;74:21–51.

76. Yoshikawa M, Peragine A, Park MY, Poethig RS. A pathway for the biogenesis of trans-acting siRNAs in Arabidopsis.
Genes Development. 2005;19:2164–75.

77. Liu Y, Teng C, Xia R, Meyers BC. PhasiRNAs in Plants: Their Biogenesis, Genic Sources, and Roles in Stress
Responses, Development, and Reproduction. THE PLANT CELL ONLINE. 2020;32:3059–80.

78. Shamandi N, Zytnicki M, Charbonnel C, Elvira-Matelot E, Bochnakian A, Comella P, et al. Plants Encode a General
siRNA Suppressor That Is Induced and Suppressed by Viruses. PLoS Biol. 2015;13:e1002326.

79. Tschopp M-A, Iki T, Brosnan CA, Jullien PE, Pumplin N. A complex of Arabidopsis DRB proteins can impair dsRNA
processing. RNA. 2017;23:782–97.

80. Montavon T, Kwon Y, Zimmermann A, Hammann P, Vincent T, Cognat V, et al. A specific dsRNA-binding protein
complex selectively sequesters endogenous inverted-repeat siRNA precursors and inhibits their processing. Nucleic
Acids Res. 2017;45:1330–44.

81. Zong J, Yao X, Yin J, Zhang D, Ma H. Evolution of the RNA-dependent RNA polymerase (RdRP) genes: Duplications
and possible losses before and after the divergence of major eukaryotic groups. Gene. 2009;447:29–39.

82. Willmann MR, Endres MW, Cook RT, Gregory BD. The Functions of RNA-Dependent RNA Polymerases in
Arabidopsis. Arab Book. 2011;2011:e0146.

83. Deleris A, Gallego-Bartolome J, Bao J, Kasschau KD, Carrington JC, Voinnet O. Hierarchical action and inhibition of
plant Dicer-like proteins in antiviral defense. Science. 2006;313:68–71.

84. Garcia-Ruiz H, Takeda A, Chapman EJ, Sullivan CM, Fahlgren N, Brempelis KJ, et al. Arabidopsis RNA-Dependent
RNA Polymerases and Dicer-Like Proteins in Antiviral Defense and Small Interfering RNA Biogenesis during Turnip
Mosaic Virus Infection. THE PLANT CELL ONLINE. 2010;22:481–96.

85. Harvey JJW, Lewsey MG, Patel K, Westwood J, Heimstädt S, Carr JP, et al. An Antiviral Defense Role of AGO2 in
Plants. Bendahmane M, editor. PLoS ONE. 2011;6:e14639.

86. Carbonell A, Fahlgren N, Garcia-Ruiz H, Gilbert KB, Montgomery TA, Nguyen T, et al. Functional analysis of three
Arabidopsis ARGONAUTES using slicer-defective mutants. THE PLANT CELL ONLINE. 2012;24:3613–29.

87. Yu D, Fan B, MacFarlane SA, Chen Z. Analysis of the Involvement of an Inducible Arabidopsis RNA-Dependent RNA
Polymerase in Antiviral Defense. Mol Plant-Microbe Interact. 2003;16:206–16.

88. Brosseau C, Moffett P. Functional and Genetic Analysis Identify a Role for Arabidopsis ARGONAUTE5 in Antiviral RNA
Silencing. Plant Cell. 2015;27:1742–54.

89. Incarbone M, Bradamante G, Pruckner F, Wegscheider T, Rozhon W, Nguyen V, et al. Salicylic acid and RNA
interference mediate antiviral immunity of plant stem cells. Proc Natl Acad Sci. 2023;120:e2302069120.

90. Bradamante G, Nguyen VH, Incarbone M, Meir Z, Bente H, Donà M, et al. Two ARGONAUTE proteins loaded with
transposon-derived small RNAs are associated with the reproductive cell lineage in Arabidopsis. Plant Cell.
2023;koad295.

91. Law JA, Vashisht AA, Wohlschlegel JA, Jacobsen SE. SHH1, a Homeodomain Protein Required for DNA Methylation,
As Well As RDR2, RDM4, and Chromatin Remodeling Factors, Associate with RNA Polymerase IV. PLoS Genet.
2011;7:e1002195.

92. Law JA, Du J, Hale CJ, Feng S, Krajewski K, Palanca AMS, et al. Polymerase IV occupancy at RNA-directed DNA
methylation sites requires SHH1. Nature. 2013;1–7.

93. Zhou M, Palanca AMS, Law JA. Locus-specific control of the de novo DNA methylation pathway in Arabidopsis by the
CLASSY family. Nat Genet. 2018;50:865–73.

94. Diego-Martin B, Pérez-Alemany J, Candela-Ferre J, Corbalán-Acedo A, Pereyra J, Alabadí D, et al. The TRIPLE PHD
FINGERS proteins are required for SWI/SNF complex-mediated +1 nucleosome positioning and transcription start site
determination in Arabidopsis. Nucleic Acids Res. 2022;50:10399–417.

95. Haag JR, Brower-Toland B, Krieger EK, Sidorenko L, Nicora CD, Norbeck AD, et al. Functional Diversification of Maize
RNA Polymerase IV and V Subtypes via Alternative Catalytic Subunits. Cell Rep. 2014;9:378–90.

96. Wang Y, Zhou X, Luo J, Lv S, Liu R, Du X, et al. Recognition of H3K9me1 by maize RNA-directed DNA methylation
factor SHH2. J Integr Plant Biol. 2021;63:1091–6.

97. Duan C-G, Zhang H, Tang K, Zhu X, Qian W, Hou YJ, et al. Specific but interdependent functions for Arabidopsis
AGO4 and AGO6 in RNA-directed DNA methylation. EMBO. 2014;e201489453.

98. Johnson LM, Law JA, Khattar A, Henderson IR, Jacobsen SE. SRA-Domain Proteins Required for DRM2-Mediated De
Novo DNA Methylation. PLoS Genet. 2008;4:e1000280.

99. Johnson LM, Du J, Hale CJ, Bischof S, Feng S, Chodavarapu RK, et al. SRA- and SET-domain-containing proteins
link RNA polymerase V occupancy to DNA methylation. Nature. 2014;507:124–8.

100. Liu Z-W, Shao C-R, Zhang C-J, Zhou J-X, Zhang S-W, Li L, et al. The SET Domain Proteins SUVH2 and SUVH9 Are
Required for Pol V Occupancy at RNA-Directed DNA Methylation Loci. PLoS Genetics. 2014;10:e1003948.

101. Chan SW-L, Zilberman D, Xie Z, Johansen LK, Carrington JC, Jacobsen SE. RNA Silencing Genes Control de Novo
DNA Methylation. Science. 2004;303:1336–1336.

102. Wierzbicki AT, Ream TS, Haag JR, Pikaard CS. RNA polymerase V transcription guides ARGONAUTE4 to chromatin.
Nature Genetics. 2009;41:630–4.

103. Zhong X, Hale CJ, Nguyen M, Ausin I, Groth M, Hetzel J, et al. DOMAINS REARRANGED METHYLTRANSFERASE3
controls DNA methylation and regulates RNA polymerase V transcript abundance in Arabidopsis. Proc Natl Acad Sci.
2015;112:911–6.

104. Liu W, Duttke SH, Hetzel J, Groth M, Feng S, Gallego-Bartolome J, et al. RNA-directed DNA methylation involves co-
transcriptional small-RNA-guided slicing of polymerase V transcripts in Arabidopsis. Nat Plants. 2018;4:181–8.

105. Fang J, Leichter SM, Jiang J, Biswal M, Lu J, Zhang Z-M, et al. Substrate deformation regulates DRM2-mediated DNA
methylation in plants. Sci Adv. 2021;7:eabd9224.

106. El-Shami M, Pontier D, Lahmy S, Braun L, Picart C, Vega D, et al. Reiterated WG/GW motifs form functionally and
evolutionarily conserved ARGONAUTE-binding platforms in RNAi-related components. Genes Development.
2007;21:2539–44.

107. Bies-Etheve N, Pontier D, Lahmy S, Picart C, Vega D, Cooke R, et al. RNA-directed DNA methylation requires an
AGO4-interacting member of the SPT5 elongation factor family. EMBO Rep. 2009;10:649–54.

108. Rowley MJ, Avrutsky MI, Sifuentes CJ, Pereira L, Wierzbicki AT. Independent Chromatin Binding of ARGONAUTE4
and SPT5L/KTF1 Mediates Transcriptional Gene Silencing. PLoS Genet. 2011;7:e1002120.

109. Huang Y, Kendall T, Forsythe ES, Dorantes-Acosta A, Li S, Caballero-Pérez J, et al. Ancient Origin and Recent
Innovations of RNA Polymerase IV and V. Mol Biol Evol. 2015;32:1788–99.

110. Trujillo JT, Beilstein MA, Mosher RA. The Argonaute-binding platform of NRPE1 evolves through modulation of
intrinsically disordered repeats. N Phytol. 2016;212:1094–105.

111. Zhang X, Yazaki J, Sundaresan A, Cokus S, Chan SWL, Chen H, et al. Genome-wide High-Resolution Mapping and
Functional Analysis of DNA Methylation in Arabidopsis. Cell. 2006;126:1189–201.

112. Regulski M, Lu Z, Kendall J, Donoghue MTA, Reinders J, Llaca V, et al. The maize methylome influences mRNA splice
sites and reveals widespread paramutation-like switches guided by small RNA. Genome Res. 2013;23:1651–62.

113. Wicker T, Sabot F, Hua-Van A, Bennetzen JL, Capy P, Chalhoub B, et al. A unified classification system for eukaryotic
transposable elements. Nature Reviews Genetics. 2007;8:973–82.

114. Zattera ML, Bruschi DP. Transposable Elements as a Source of Novel Repetitive DNA in the Eukaryote Genome. Cells.
2022;11:3373.

115. Zhong X, Hale CJ, Law JA, Johnson LM, Feng S, Tu A, et al. DDR complex facilitates global association of RNA
polymerase V to promoters and evolutionarily young transposons. Nat Struct Mol Biol. 2012;19:870–5.

116. Zemach A, Kim MY, Hsieh P-H, Coleman-Derr D, Eshed-Williams L, Thao K, et al. The Arabidopsis Nucleosome
Remodeler DDM1 Allows DNA Methyltransferases to Access H1-Containing Heterochromatin. Cell. 2013;153:193–
205.

117. Wang Z, Baulcombe DC. Transposon age and non-CG methylation. Nat Commun. 2020;11:1221.

118. Kumar A, Bennetzen JL. Plant retrotransposons. Annual Review of Genetics. 1999;33:479–532.

119. Gorinšek B, Gubenšek F, Kordiš D. Evolutionary Genomics of Chromoviruses in Eukaryotes. Mol Biol Evol.
2004;21:781–98.

120. Neumann P, Novák P, Hoštáková N, Macas J. Systematic survey of plant LTR-retrotransposons elucidates
phylogenetic relationships of their polyprotein domains and provides a reference for element classification. Mob DNA.
2019;10:1.

121. Jackson JP, Johnson L, Jasencakova Z, Zhang X, PerezBurgos L, Singh PB, et al. Dimethylation of histone H3 lysine
9 is a critical mark for DNA methylation and gene silencing in Arabidopsis thaliana. Chromosoma. 2004;112:308–15.

122. Fuchs J, Demidov D, Houben A, Schubert I. Chromosomal histone modification patterns – from conservation to
diversity. Trends Plant Sci. 2006;11:199–208.

123. Choi J, Lyons DB, Zilberman D. Histone H1 prevents non-CG methylation-mediated small RNA biogenesis in
Arabidopsis heterochromatin. eLife. 2021;10:e72676.

124. Johnson L, Mollah S, Garcia BA, Muratore TL, Shabanowitz J, Hunt DF, et al. Mass spectrometry analysis of
Arabidopsis histone H3 reveals distinct combinations of post-translational modifications. Nucleic Acids Res.
2004;32:6511–8.

125. Scheid R, Dowell JA, Sanders D, Jiang J, Denu JM, Zhong X. Histone Acid Extraction and High Throughput Mass
Spectrometry to Profile Histone Modifications in Arabidopsis thaliana. Curr Protoc. 2022;2:e527.

126. Mathieu O, Probst AV, Paszkowski J. Distinct regulation of histone H3 methylation at lysines 27 and 9 by CpG
methylation in Arabidopsis. 2005;24:2783–91.

127. Walter M, Teissandier A, Pérez-Palacios R, Bourc'his D. An epigenetic switch ensures transposon repression upon
dynamic loss of DNA methylation in embryonic stem cells. eLife. 2016;5:e11418.

128. Montgomery SA, Tanizawa Y, Galik B, Wang N, Ito T, Mochizuki T, et al. Chromatin Organization in Early Land Plants
Reveals an Ancestral Association between H3K27me3, Transposons, and Constitutive Heterochromatin. Curr Biol.
2020;30:573-588.e7.

129. Rougée M, Quadrana L, Zervudacki J, Hure V, Colot V, Navarro L, et al. Polycomb mutant partially suppresses DNA
hypomethylation–associated phenotypes in Arabidopsis. Life Sci Alliance. 2021;4:e202000848.

130. Hisanaga T, Romani F, Wu S, Kowar T, Wu Y, Lintermann R, et al. The Polycomb repressive complex 2 deposits
H3K27me3 and represses transposable elements in a broad range of eukaryotes. Curr Biol. 2023;33:4367-4380.e9.

131. Zhang X, Bernatavichute YV, Cokus S, Pellegrini M, Jacobsen SE. Genome-wide analysis of mono-, di- and
trimethylation of histone H3 lysine 4 in Arabidopsis thaliana. Genome Biology. 2009;10:R62.

132. Roudier F, Ahmed I, Bérard C, Sarazin A, Mary-Huard T, Cortijo S, et al. Integrative epigenomic mapping defines four
main chromatin states in Arabidopsis. EMBO J. 2011;30:1928–38.

133. Ding Y, Fromm M, Avramova Z. Multiple exposures to drought "train" transcriptional responses in Arabidopsis. Nat
Commun. 2012;3:740.

134. Pratx L, Wendering P, Kappel C, Nikoloski Z, Bäurle I. Histone retention preserves epigenetic marks during heat stress-
induced transcriptional memory in plants. EMBO J. 2023;42:e113595.

135. Cao HX, Vu GTH, Wang W, Messing J, Schubert I. Chromatin organisation in duckweed interphase nuclei in relation
to the nuclear DNA content. Plant Biol. 2015;17:120–4.

136. Ebbs ML, Bender J. Locus-Specific Control of DNA Methylation by the Arabidopsis SUVH5 Histone Methyltransferase.
Plant Cell. 2006;18:1166–76.

137. Jackson JP, Lindroth AM, Cao X, Jacobsen SE. Control of CpNpG DNA methylation by the KRYPTONITE histone H3
methyltransferase. Nature. 2002;416:556–60.

138. Jasencakova Z, Soppe WJJ, Meister A, Gernand D, Turner BM, Schubert I. Histone modifications in Arabidopsis– high
methylation of H3 lysine 9 is dispensable for constitutive heterochromatin. Plant J. 2003;33:471–80.

139. Mi S, Cai T, Hu Y, Chen Y, Hodges E, Ni F, et al. Sorting of Small RNAs into Arabidopsis Argonaute Complexes Is
Directed by the 5′ Terminal Nucleotide. Cell. 2008;133:116–27.

140. McCue AD, Panda K, Nuthikattu S, Choudury SG, Thomas EN, Slotkin RK. ARGONAUTE 6 bridges transposable
element mRNA-derived siRNAs to the establishment of DNA methylation. The EMBO journal. 2015;34:20–35.

141. Panda K, McCue AD, Slotkin RK. Arabidopsis RNA Polymerase IV generates 21–22 nucleotide small RNAs that can
participate in RNA-directed DNA methylation and may regulate genes. Philos Trans R Soc B. 2020;375:20190417.

142. Sigman MJ, Panda K, Kirchner R, McLain LL, Payne H, Peasari JR, et al. An siRNA-guided ARGONAUTE protein
directs RNA polymerase V to initiate DNA methylation. Nat Plants. 2021;7:1461–74.

143. Keith B, Chua N-H. Monocot and dicot pre-mRNAs are processed with different efficiencies in transgenic tobacco.
EMBO J. 1986;5:2419–25.

144. Goodall GJ, Filipowicz W. Different effects of intron nucleotide composition and secondary structure on pre-mRNA
splicing in monocot and dicot plants. EMBO J. 1991;10:2635–44.

145. Lou H, McCullough AJ, Schuler MA. Expression of maize Adh1 intron mutants in tobacco nuclei. Plant J. 1993;3:393–
403.

146. Jullien PE, Grob S, Marchais A, Pumplin N, Chevalier C, Bonnet DMV, et al. Functional characterization of Arabidopsis ARGONAUTE 3 in reproductive tissues. Plant J. 2020;103:1796–809.

147. Fang X, Qi Y. RNAi in Plants: An Argonaute-Centered View. Plant Cell. 2016;28:272–85.

148. Slotkin RK, Vaughn M, Borges F, TanurdZiC M, Becker JoD, FeijO JA, et al. Epigenetic Reprogramming and Small RNA Silencing of Transposable Elements in Pollen. Cell. 2009;136:461–72.

149. Lee SC, Ernst E, Berube B, Borges F, Parent J-S, Ledon P, et al. Arabidopsis retrotransposon virus-like particles and their regulation by epigenetically activated small RNA. Genome Research. 2020;30:576–88.

150. Oberlin S, Rajeswaran R, Trasser M, Barragán-Borrero V, Schon MA, Plotnikova A, et al. Innate, translation-dependent silencing of an invasive transposon in Arabidopsis. Embo Rep. 2022;23:e53400.

151. Xie Z, Allen E, Wilken A, Carrington JC. DICER-LIKE 4 functions in trans-acting small interfering RNA biogenesis and vegetative phase change in Arabidopsis thaliana. Proceedings of the National Academy of Sciences of the United States of America. 2005;102:12984–9.

152. Henderson IR, Zhang X, Lu C, Johnson L, Meyers BC, Green PJ, et al. Dissecting Arabidopsis thaliana DICER function in small RNA processing, gene silencing and DNA methylation patterning. Nature Genetics. 2006;38:721–5.

153. Wang T, Deng Z, Zhang X, Wang H, Wang Y, Liu X, et al. Tomato DCL2b is required for the biosynthesis of 22-nt small RNAs, the resulting secondary siRNAs, and the host defense against ToMV. Hortic Res. 2018;5:62.

154. Jia J, Ji R, Li Z, Yu Y, Nakano M, Long Y, et al. Soybean DICER-LIKE2 Regulates Seed Coat Color via Production of Primary 22-Nucleotide Small Interfering RNAs from Long Inverted Repeats. Plant Cell. 2020;32:3662–73.

155. Hamilton AJ, Voinnet O, Chappell L, Baulcombe DC. Two classes of short interfering RNA in RNA silencing. Embo Journal. 2002;21:4671–9.

156. Bertazzon N, Raiola A, Castiglioni C, Gardiman M, Angelini E, Borgo M, et al. Transient silencing of the grapevine gene VvPGIP1 by agroinfiltration with a construct for RNA interference. Plant Cell Rep. 2012;31:133–43.

157. Andrieu A, Breitler JC, Siré C, Meynard D, Gantet P, Guiderdoni E. An in planta, Agrobacterium-mediated transient gene expression method for inducing gene silencing in rice (Oryza sativa L.) leaves. Rice. 2012;5:23.

158. Sun S, Kang X-P, Xing X-J, Chen Z-F, Zheng S-W, Xing G-M. Transient expression of siRNA targeted against the TYLCV AV1, AC1 and AC3 genes for high resistance in tomato. Sci Hortic. 2014;179:321–7.

159. Jay F, Brioudes F, Voinnet O. A contemporary reassessment of the enhanced transient expression system based on the tombusviral silencing suppressor protein P19. Plant J. 2023;113:186–204.

160. Yang G-L, Fang Y, Xu Y-L, Tan L, Li Q, Liu Y, et al. Frond transformation system mediated by Agrobacterium tumefaciens for Lemna minor. Plant Mol Biol. 2018;98:319–31.

161. He Y, Zhang T, Sun H, Zhan H, Zhao Y. A reporter for noninvasively monitoring gene expression and plant transformation. Hortic Res. 2020;7:152.

162. Šurbanovski N, Brilli M, Moser M, Si-Ammour A. A highly specific microRNA-mediated mechanism silences LTR retrotransposons of strawberry. The Plant Journal. 2016;85:70–82.

163. Benachenhou F, Sperber GO, Bongcam-Rudloff E, Andersson G, Boeke JD, Blomberg J. Conserved structure and inferred evolutionary history of long terminal repeats (LTRs). Mob DNA. 2013;4:5.

164. Michael TP, Ernst E, Hartwick N, Chu P, Bryant D, Gilbert S, et al. Genome and time-of-day transcriptome of Wolffia australiana link morphological minimization with gene loss and less growth control. Genome Res. 2020;31:gr.266429.120.

165. Abramson BW, Novotny M, Hartwick NT, Colt K, Aevermann BD, Scheuermann RH, et al. The genome and preliminary single-nuclei transcriptome of Lemna minuta reveals mechanisms of invasiveness. Plant Physiol. 2021;188:879–97.

166. Fang Y, Tian X, Jin Y, Du A, Ding Y, Liao Z, et al. Duckweed evolution: from land back to water. bioRxiv. 2023;2023.03.22.533731.

167. Liu J, He Z. Small DNA Methylation, Big Player in Plant Abiotic Stress Responses and Memory. Front Plant Sci. 2020;11:595603.

168. Urquiaga MC de O, Thiebaut F, Hemerly AS, Ferreira PCG. From Trash to Luxury: The Potential Role of Plant LncRNA in DNA Methylation During Abiotic Stress. Front Plant Sci. 2021;11:603246.

169. Arkhipova I, Meselson M. Transposable elements in sexual and ancient asexual taxa. Proc Natl Acad Sci. 2000;97:14473–7.

170. Wright S, Finnegan D. Genome evolution: Sex and the transposable element. Curr Biol. 2001;11:R296–9.

171. Bast J, Jaron KS, Schuseil D, Roze D, Schwander T. Asexual reproduction reduces transposable element load in experimental yeast populations. eLife. 2019;8:e48548.

172. Dechaud C, Volff J-N, Schartl M, Naville M. Sex and the TEs: transposable elements in sexual development and function in animals. Mob DNA. 2019;10:42.

173. Nguyen V, Gutzat R. Epigenetic regulation in the shoot apical meristem. Curr Opin Plant Biol. 2022;69:102267.

174. Czech B, Munafò M, Ciabrelli F, Eastwood EL, Fabry MH, Kneuss E, et al. piRNA-Guided Genome Defense: From Biogenesis to Silencing. Annu Rev Genet. 2018;52:131–57.

175. Huff JT, Zilberman D, Roy SW. Mechanism for DNA transposons to generate introns on genomic scales. Nature. 2016;538:533–6.

176. Chénais B, Caruso A, Hiard S, Casse N. The impact of transposable elements on eukaryotic genomes: From genome size increase to genetic adaptation to stressful environments. Gene. 2012;509:7–15.

177. Bourque G, Burns KH, Gehring M, Gorbunova V, Seluanov A, Hammell M, et al. Ten things you should know about transposable elements. Genome Biol. 2018;19:199.

178. Gaillochet C, Lohmann JU. The never-ending story: from pluripotency to plant developmental plasticity. Development. 2015;142:2237–49.

179. Sree KS, Maheshwari SC, Boka K, Khurana JP, Keresztes Á, Appenroth K-J. The duckweed Wolffia microscopica: A unique aquatic monocot. Flora - Morphol, Distrib, Funct Ecol Plants. 2015;210:31–9.

180. Yang J, Zhao X, Li G, Hu S, Hou H. Frond architecture of the rootless duckweed Wolffia globosa. BMC Plant Biol. 2021;21:387.

181. Li F, Yang J-J, Sun Z-Y, Wang L, Qi L-Y, A S, et al. Plant-on-chip: Core morphogenesis processes in the tiny plant Wolffia australiana. PNAS Nexus. 2023;2:pgad141.

182. Gutzat R, Rembart K, Nussbaumer T, Hofmann F, Pisupati R, Bradamante G, et al. Arabidopsis shoot stem cells display dynamic transcription and DNA methylation patterns. EMBO J. 2020;39:e103667.

183. Ho EKH, Bartkowska M, Wright SI, Agrawal AF. Population genomics of the facultatively asexual duckweed Spirodela polyrhiza. N Phytol. 2019;224:1361–71.

184. Wang Y, Duchen P, Chávez A, Sree SK, Appenroth KJ, Zhao H, et al. Population genomics and epigenomics provide insights into the evolution of facultative asexuality in plants. bioRxiv. 2023;2023.07.18.549461.

185. Fusaro AF, Matthew L, Smith NA, Curtin SJ, Dedic-Hagan J, Ellacott GA, et al. RNA interference-inducing hairpin RNAs in plants act through the viral defence pathway. EMBO Rep. 2006;7:1168–75.

186. Parent J, Bouteiller N, Elmayan T, Vaucheret H. Respective contributions of Arabidopsis DCL2 and DCL4 to RNA silencing. Plant J. 2015;81:223–32.

187. Nielsen CPS, Arribas-Hernández L, Han L, Andersen SU, Pumplin N, Brodersen P. Evidence for an RNAi-independent role of DICER-LIKE2 in conferring growth inhibition and basal antiviral resistance. The Plant Cell. 2024; koae0676.

188. Nielsen CPS, Han L, Arribas-Hernández L, Karelina D, Petersen M, Brodersen P. Sensing of viral RNA in plants via a DICER-LIKE Ribonuclease. bioRxiv. 2023;2023.01.10.523395.

189. Chen H-M, Chen L-T, Patel K, Li Y-H, Baulcombe DC, Wu S-H. 22-Nucleotide RNAs trigger secondary siRNA biogenesis in plants. Proceedings of the National Academy of Sciences. 2010;107:15269–74.

190. Zhang X, Singh J, Li D, Qu F. Temperature-Dependent Survival of Turnip Crinkle Virus-Infected Arabidopsis Plants Relies on an RNA Silencing-Based Defense That Requires DCL2, AGO2, and HEN1. Journal of Virology. 2012;86:6847–54.

191. Taochy C, Gursanscky NR, Cao J, Fletcher SJ, Dressel U, Mitter N, et al. A Genetic Screen for Impaired Systemic RNAi Highlights the Crucial Role of DICER-LIKE 2 . Plant Physiol. 2017;175:1424–37.

192. Chen W, Zhang X, Fan Y, Li B, Ryabov E, Shi N, et al. A Genetic Network for Systemic RNA Silencing in Plants . Plant Physiol. 2018;176:2700–19.

193. Zimin AV, Marçais G, Puiu D, Roberts M, Salzberg SL, Yorke JA. The MaSuRCA genome assembler. Bioinformatics. 2013;29:2669–77.

194. Zimin AV, Salzberg SL. The genome polishing tool POLCA makes fast and accurate corrections in genome assemblies. PLoS Comput Biol. 2020;16:e1007981.

195. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. Genome Res. 2010;20:1297–303.

196. Poplin R, Ruano-Rubio V, DePristo MA, Fennell TJ, Carneiro MO, Auwera GAV der, et al. Scaling accurate genetic
variant discovery to tens of thousands of samples. bioRxiv. 2018;201178.

197. Shumate A, Salzberg SL. Liftoff: accurate mapping of gene annotations. Bioinformatics. 2021;37:btaa1016.

198. Zhang R-G, Li G-Y, Wang X-L, Dainat J, Wang Z-X, Ou S, et al. TEsorter: An accurate and fast method to classify
LTR-retrotransposons in plant genomes. Hortic Res. 2022;9:uhac017.

199. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq quantification. Nat Biotechnol.
2016;34:525–7.

200. Li H. New strategies to improve minimap2 alignment accuracy. Bioinformatics. 2021;37:btab705-.

201. Krueger F, Andrews SR. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq applications. Bioinformatics.
2011;27:1571–2.

202. Schultz MD, He Y, Whitaker JW, Hariharan M, Mukamel EA, Leung D, et al. Human body epigenome maps reveal
noncanonical DNA methylation variation. Nature. 2015;523:212–6.

203. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics.
2010;26:841–2.

204. Huang X, Zhang S, Li K, Thimmapuram J, Xie S, Wren J. ViewBS: a powerful toolkit for visualization of high-throughput
bisulfite sequencing data. Bioinformatics. 2018;34:708–9.

205. Hayashi R, Schnabl J, Handler D, Mohn F, Ameres SL, Brennecke J. Genetic and mechanistic diversity of piRNA 3′-
end formation. Nature. 2016;539:588–92.

206. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics.
2014;30:2114–20.

207. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nature Methods. 2012;9:357–9.

208. Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, et al. deepTools2: a next generation web server
for deep-sequencing data analysis. Nucleic Acids Res. 2016;44:W160–5.

209. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map format and SAMtools.
Bioinformatics. 2009;25:2078–9.

210. Sambrook J, Russell DW. Molecular Cloning. CSHL Press; 2001.

211. Felippes FF de, Weigel D. Transient assays for the analysis of miRNA processing and function. Methods in molecular
biology (Clifton, NJ). 2010;592:255–64.

212. Pall GS, Hamilton AJ. Improved northern blot method for enhanced detection of small RNA. Nature Protocols.
2008;3:1077–84.

213. Gu Z, Gu L, Eils R, Schlesner M, Brors B. circlize implements and enhances circular visualization in R. Bioinformatics.
2014;30:2811–2.

214. Wickham H. Ggplot2: Elegant graphics for data analysis. Springer; 2016.

215. Oberlin S, Sarazin A, Chevalier C, Voinnet O, Ordóñez AM. A genome-wide transcriptome and translatome analysis
of Arabidopsis transposons identifies a unique and conserved genome expression strategy for Ty1/Copia
retroelements. Genome Research. 2017;27:1549–62.

1475

1476