1
2
3
4
5

Endogenous hydrogen peroxide positively regulates secretion of a gut-derived

peptide in neuroendocrine potentiation of the oxidative stress response in *C.*

elegans

Qi Jia[1,3], Drew Young[2,4], Qixin Zhang[3,4] and Derek Sieburth[4,5,*]

[1] Development, Stem Cells and Regenerative Medicine PhD program, Keck School of Medicine, University of Southern California, Los Angeles, CA 90033

[2] Neuroscience Graduate Program, University of Southern California, Los Angeles, CA 90089

[3] Neuromedicine Graduate Program, University of Southern California, Los Angeles, CA 90089

[4] Zilkha Neurogenetic Institute, University of Southern California, Los Angeles, CA 90033

[5] Department of Physiology and Neuroscience, Keck School of Medicine, University of Southern California, Los Angeles, CA 90033

24
25
26
27

* Corresponding author
sieburth@med.usc.edu

Keywords: *C. elegans*, oxidative stress, neuropeptide, hydrogen peroxide, mitochondria, superoxide dismutase, peroxiredoxin, thioredoxin, protein kinase C, SNAP25, SOD-1, SOD-3, PRDX-2, TRX-3, PKC-2, AEX-4.

Abstract

The gut-brain axis mediates bidirectional signaling between the intestine and the nervous system and is critical for organism-wide homeostasis. Here we report the identification of a peptidergic endocrine circuit in which bidirectional signaling between neurons and the intestine potentiates the activation of the antioxidant response in *C. elegans* in the intestine. We identify a FMRF-amide-like peptide, FLP-2, whose release from the intestine is necessary and sufficient to activate the intestinal oxidative stress response by promoting the release of the antioxidant FLP-1 neuropeptide from neurons. FLP-2 secretion from the intestine is positively regulated by endogenous hydrogen peroxide (H_2O_2) produced in the mitochondrial matrix by *sod-3*/superoxide dismutase, and is negatively regulated by *prdx-2*/peroxiredoxin, which depletes H_2O_2 in both the mitochondria and cytosol. H_2O_2 promotes FLP-2 secretion through the DAG and calcium-dependent protein kinase C family member *pkc-2* and by the SNAP25 family member *aex-4* in the intestine. Together, our data demonstrate a role for intestinal H_2O_2 in promoting inter-tissue antioxidant signaling through regulated neuropeptide-like protein exocytosis in a gut-brain axis to activate the oxidative stress response.

Introduction

The gut-brain axis is critical for communication between the intestine and the nervous system to regulate behavior and maintain homeostasis, and altered gut-brain signaling is associated with neurodegeneration, obesity and tumor proliferation (Carabotti et al. 2015; Grenham et al. 2011; Mayer, Nance, and Chen 2022; Mehrian-Shai et al. 2019; Vitali et al. 2022). Over the last decade the importance of peptides that function as signals in gut-brain signaling has gained recognition. Numerous gut peptides are distributed throughout the gastrointestinal (GI) tract with regional specificity (Haber et al. 2017), and gut-secreted peptides can modulate neurocircuits that regulate feeding behavior and glucose metabolism (Batterham and Bloom 2003; Han et al. 2018; Song et al. 2019), inflammatory responses against pathogenic bacteria (Campos-Salinas et al. 2014; Yu et al. 2021), and satiety (Batterham et al. 2002; Chelikani, Haver, and Reidelberger 2005; Gibbs, Young, and Smith 1973; Lutz et al. 1995; Lutz, Del Prete, and Scharrer 1994; West, Fey, and Woods 1984). A gut-released peptide suppresses arousal through dopaminergic neurons during sleep in *Drosophila* (Titos et al. 2023). In *C. elegans*, gut-derived peptides regulate rhythmic behavior and behavioral responses to pathogenic bacteria (Lee and Mylonakis 2017; Singh and Aballay 2019; Wang et al. 2013). Conversely, peptides released from the nervous system regulate many aspects of intestinal function including gut mobility, inflammation and immune defense (Browning and Travagli 2014; Furness et al. 2014; Lai, Mills, and Chiu 2017). In *C. elegans*, the secretion of peptides from various neurons regulates the mitochondrial unfolded protein response (UPR^{mt}), the heat shock response, and the antioxidant response in the intestine (Jia and Sieburth 2021; Maman et al. 2013; Prahlad, Cornelius, and Morimoto 2008; Shao, Niu, and Liu 2016). In spite of the many roles of peptides in the gut-brain axis, the mechanisms underlying the regulation of intestinal peptide secretion and signaling remain to be fully defined.

Hydrogen peroxide (H_2O_2) is emerging as an important signaling molecule that regulates intracellular signaling pathways by modifying specific reactive residues on target proteins. For example, H_2O_2-regulated phosphorylation of inhibitor of nuclear factor κB (NF-κB) kinase, leads to the activation of NF-κB during development, inflammation and immune responses. (Kamata et al. 2002; Oliveira-Marques et al. 2009; Takada et al. 2003). In addition, H_2O_2-induced tyrosine and cysteine modifications contribute to redox regulation of c-Jun N-terminal kinase 2 (JNK2), Src family kinase, extracellular signal-regulated kinases 1 and 2 (ERK1/2), protein kinase C (PKC) and other protein kinases (Kemble and Sun 2009; Konishi et al. 1997; Lee et al. 2003; Nelson et al. 2018). H_2O_2 signaling has been implicated in regulating neurotransmission and transmitter secretion. H_2O_2 at low concentration increases neurotransmission at neuromuscular junctions without influencing lipid oxidation (A R Giniatullin and R A Giniatullin 2003; Giniatullin, Petrov, and Giniatullin 2019; Shakirzyanova et al. 2009), and enhanced endogenous H_2O_2 generation regulates dopamine release (Avshalumov et al. 2005; Avshalumov and Rice 2003; Bao, Avshalumov, and Rice 2005; Chen, Avshalumov, and Rice 2001, 2002). Acute H_2O_2 treatment increases exocytosis of ATP-containing vesicles in astrocytes (Li et al. 2019). Finally, mitochondrially-derived H_2O_2 regulates neuropeptide release from neurons in *C. elegans* (Jia and Sieburth 2021). Cellular H_2O_2 levels are tightly controlled through the regulation of its production from superoxide by superoxide dismutases (SODs) and cytoplasmic oxidases (Fridovich 1995, 1997; Messner and Imlay 2002; Zelko, Mariani,

and Folz 2002), and through its degradation by catalases, peroxidases, and
peroxiredoxins (Chance, Sies, and Boveris 1979; Marinho et al. 2014). In the intestine,
endogenously produced H_2O_2 plays important roles as an antibacterial agent in the lumen,
and in activating the ER unfolded protein response (UPR^{ER}) through protein sulfenylation
(Botteaux et al. 2009; Corcionivoschi et al. 2012; Hourihan et al. 2016; Miller et al. 2020).
Here we demonstrate a role of endogenous H_2O_2 signaling in the intestine in
regulating the release of the intestinal FMRFamide-like peptide, FLP-2, to modulate a
neurocircuit that activates the anti-oxidant response in the intestine in *C. elegans.*
Intestinal FLP-2 signaling functions by potentiating the release of the antioxidant
neuropeptide-like protein FLP-1 from AIY interneurons, which in turn activates the
antioxidant response in the intestine. FLP-2 secretion from the intestine is rapidly and
positively regulated by H_2O_2, whose levels are positively regulated by superoxide
dismutases in the mitochondrial matrix and cytosol, and negatively regulated by the
peroxiredoxin-thioredoxin system in the cytosol. Intestinal FLP-2 release is mediated by
aex-4/SNAP25-dependent exocytosis of dense core vesicles and H_2O_2-induced FLP-2
secretion is dependent upon the production of intestinal diacylglycerol and on *pkc-*
2/PKCα/β kinase activity.
123
124 **Results**
125
126 **Neuronal FLP-1 secretion is regulated by neuropeptide signaling from the intestine.**
127
We previously showed that 10 minute treatment with the mitochondrial toxin
juglone leads to a rapid, reversible and specific increase in FLP-1 secretion from AIY, as
measured by a two-fold increase in coelomocyte fluorescence in animals expressing FLP-
1::Venus fusion proteins in AIY (Fig. 1A and B, (Jia and Sieburth 2021)). Coelomocytes
take up secreted neuropeptides by bulk endocytosis (Fares and Greenwald 2001) and
the fluorescence intensity of Venus in their endocytic vacuoles is used as a measure of
regulated neuropeptide secretion efficacy (Ailion et al. 2014; Ch'ng, Sieburth, and Kaplan
2008; Sieburth, Madison, and Kaplan 2006). To determine the role of the intestine in
regulating FLP-1 secretion, we first examined *aex-5* mutants. *aex-5* encodes an intestinal
subtilisin/kexin type 5 prohormone convertase that functions to proteolytically process
peptide precursors into mature peptides in dense core vesicles (DCVs) (Edwards et al.
2019; Thacker and Rose 2000), and *aex-5* mutants are defective in peptide signaling from
the intestine (Mahoney et al. 2008). We found that *aex-5* mutants expressing FLP-
1::Venus in AIY exhibited no significant difference in coelomocyte fluorescence compared
to wild type controls in the absence of juglone (Fig. 1B). However, coelomocyte
fluorescence did not significantly increase in *aex-5* mutants treated with juglone.
Expression of *aex-5* cDNA selectively in the intestine (under the *ges-1* promoter) fully
restored normal responses to juglone to *aex-5* mutants, whereas *aex-5* cDNA expression
in the nervous system (under the *rab-3* promoter) failed to rescue (Fig. 1B). Thus, peptide
processing in intestinal DCVs is necessary for juglone-induced FLP-1 secretion from AIY.
Next, we examined a number of mutants with impaired SNARE-mediated vesicle
release in the intestine including *aex-1*/UNC13, *aex*-3/MADD, aex-4/SNAP25b, and *aex-*
6/Rab27 (Fig. 2C (Iwasaki et al., 1997; Mahoney et al., 2006; Thacker & Rose, 2000;
Thomas, 1990; Wang et al., 2013)), and they each exhibited no increases in FLP-1

secretion following juglone treatment above levels observed in untreated controls (Fig.
S1A). NLP-40 is a neuropeptide-like protein whose release from the intestine is presumed
to be controlled by *aex-1*, *aex-3*, *aex-4* and *aex-6* (Lin-Moore, Oyeyemi, and Hammarlund
2021; Mahoney et al. 2008; Shi et al. 2022; Wang et al. 2013). Null mutants in *nlp-40* or
its receptor, *aex-2* (Wang et al. 2013), exhibited normal juglone-induced FLP-1 secretion
(Fig. S1B). These results establish a gut-to-neuron signaling pathway that regulates FLP-
1 secretion from AIY that is likely to be controlled by peptidergic signaling distinct from
NLP-40.
160
161 **FLP-2 signaling from the intestine potentiates neuronal FLP-1 secretion and the**
162 **oxidative stress response.**
163
flp-1 protects animals from the toxic effects of juglone (Jia and Sieburth 2021). We
reasoned that the intestinal signal that regulates FLP-1 secretion should also protect
animals from juglone-induced toxicity. We identified the FMRF-amide neuropeptide-like
protein, *flp-2*, in an RNA interference (RNAi) screen for neuropeptides that confer
hypersensitivity to juglone toxicity upon knockdown (Jia and Sieburth 2021). *flp-2*
signaling has been implicated in regulating lifespan, reproductive development,
locomotion during lethargus, and the mitochondrial unfolded protein response (UPR^{mt})
(Chai et al. 2022; Chen et al. 2016; Kageyama et al. 2022; Shao et al. 2016). Putative *flp-*
2(ok3351) null mutants, which eliminate most of the *flp-2* coding region, are superficially
as healthy as wild type animals, but they exhibited significantly reduced survival in the
presence of juglone compared to wild type controls (Fig. 1C). The reduced survival rate
of *flp-2* mutants was similar to that of *flp-1* mutants, and *flp-1; flp-2* double mutants
exhibited survival rates that were not more severe than those of single mutants (Fig. 1C),
suggesting that *flp-1* and *flp-2* may function in a common genetic pathway.
To determine whether *flp-2* signaling regulates FLP-1 secretion from AIY, we
examined FLP-1::Venus secretion. *flp-2* mutants exhibited normal levels of FLP-1
secretion in the absence of stress, but FLP-1 secretion failed to significantly increase
following juglone treatment of *flp-2* mutants (Fig. 1D). *flp-2* is expressed in a subset of
neurons as well as the intestine (Chai et al. 2022), and *flp-2* functions from the nervous
system for its roles in development and the UPR^{mt} (Chai et al. 2022; Chen et al. 2016;
Kageyama et al. 2022; Shao et al. 2016). Expressing a *flp-2* genomic DNA, fragment
(containing both the *flp-2a* and *flp-2b* isoforms that arise by alternative splicing),
specifically in the nervous system failed to rescue the FLP-1::Venus defects of *flp-2*
mutants, whereas expressing *flp-2* selectively in the intestine fully restored juglone-
induced FLP-1::Venus secretion to *flp-2* mutants (Fig. 1D). These results indicate that *flp-*
2 signaling is dispensable for FLP-1 secretion from AIY under normal conditions, but that
flp-2 originating from the intestine is necessary to increase FLP-1 secretion during
oxidative stress.
To address how *flp-2* signaling regulates FLP-1 secretion from AIY, we examined
H$_2$O$_2$ levels in AIY using a mitochondrially targeted pH-stable H$_2$O$_2$ sensor HyPer7 (mito-
HyPer7, Pak et al. 2020). Mito-HyPer7 adopted a punctate pattern of fluorescence in AIY
axons, and the average fluorescence intensity of axonal mito-HyPer7 puncta increased
about two-fold following 10 minute juglone treatment (Fig 1E), in agreement with our

previous studies using HyPer (Jia and Sieburth 2021), confirming that juglone rapidly increases mitochondrial AIY H_2O_2 levels. *flp-2* mutations had no significant effects on the localization or the average intensity of mito-HyPer7 puncta in AIY axons either in the absence of juglone, or in the presence of juglone (Fig 1E), suggesting that *flp-2* signaling promotes FLP-1 secretion by a mechanism that does not increase H_2O_2 levels in AIY. Consistent with this, intestinal overexpression of *flp-2* had no effect on FLP-1::Venus secretion in the absence of juglone, but significantly enhanced the ability of juglone to increase FLP-1 secretion (Fig. 1D). We conclude that both elevated mitochondrial H_2O_2 levels and intact *flp-2* signaling from the intestine are necessary to increase FLP-1 secretion from AIY.

Previously we showed that FLP-1 signaling from AIY positively regulates the activation of the antioxidant transcription factor SKN-1/Nrf2 in the intestine. Specifically, *flp-1* mutations impair the juglone-induced expression of the SKN-1 reporter transgene P*gst-4::gfp* (Fig. 1F and (Jia and Sieburth 2021)). We found that mutations in *flp-2* caused a similar reduction in juglone-induced P*gst-4::gfp* expression as *flp-1* mutants, and that *flp-1; flp-2* double mutants exhibited similar impairments in juglone-induced P*gst-4::gfp* expression as *flp-1* or *flp-2* single mutants (Fig. 1F). Conversely, overexpression of *flp-2* selectively in the intestine elevated juglone-induced P*gst-4::gfp* expression, without altering baseline P*gst-4::gfp* expression, and the elevated P*gst-4::gfp* expression in juglone-treated animals overexpressing *flp-2* was entirely dependent upon *flp-1* (Fig. 1G). It's noteworthy that overexpressing *flp-2* in the intestine did not enhance FLP-1::Venus release or *Pgst-4::gfp* expression in the absence of stress, indicating that the FLP-1 mediated anti-oxidant pathway by *flp-2* is stress-activated. Together this data indicates that *flp-2* signaling originating in the intestine positively regulates the stress-induced secretion of FLP-1 from AIY, as well as the subsequent activation of anti-oxidant response genes in the intestine. We propose that FLP-1 and FLP-2 define a bidirectional gut-neuron signaling axis, whereby during periods of oxidative stress, FLP-2 released from the intestine positively regulates FLP-1 secretion from AIY, and FLP-1, in turn, potentiates the antioxidant response in the intestine (Fig. 1A).

FLP-2 secretion from the intestine is H_2O_2-regulated.

To directly investigate the mechanisms underlying the regulation of FLP-2 secretion, we examined FLP-2::Venus fusion proteins expressed in the intestine under various conditions (Fig. 2A). FLP-2::Venus fusion proteins adopted a punctate pattern of fluorescence throughout the cytoplasm of intestinal cells and at the plasma membrane (Fig. 2A), and FLP-2::Venus puncta co-localized with the DCV cargo protein AEX-5/PCSK5 tagged to mTurquoise2 (AEX-5::mTur2, Fig. 2B). FLP-2::Venus fluorescence was also observed in the coelomocytes (marked by mCherry) (Fig. 2A, D), indicating that FLP-2 is released from the intestine. SNAP25 forms a component of the core SNARE complex, which drives vesicular membrane fusion and transmitter release (Chen and Scheller 2001; Goda 1997; Jahn and Scheller 2006). *aex-4* encodes the *C. elegans* homolog of SNAP25, and mutations in *aex-4* disrupt the secretion of neuropeptides from the intestine (Lin-Moore et al. 2021; Mahoney et al. 2008; Wang et al. 2013) (Fig. 2C). We found that *aex-4* null mutations significantly reduced coelomocyte fluorescence in FLP-2::Venus expressing animals, and expression of *aex-4* cDNA selectively in the

intestine fully restored FLP-2 secretion to *aex-4* mutants (Fig. 2D). Together these results suggest that intestinal FLP-2 can be packaged into DCVs that undergo release via SNARE-dependent exocytosis.

To test whether intestinal FLP-2 secretion is regulated by oxidative stress, we examined coelomocyte fluorescence in FLP-2::Venus-expressing animals that had been exposed to a number of different commonly used oxidative stressors. We found that 10 minute exposure to juglone, thimerosal, or paraquat, which promote mitochondria-targeted toxicity (Castello, Drechsel, and Patel 2007; Elferink 1999; Sharpe, Livingston, and Baskin 2012), each significantly increased Venus fluorescence intensity in the coelomocytes compared to untreated controls (Fig. 2D and E). We conducted four controls for specificity: First, juglone treatment did not significantly alter fluorescence intensity of mCherry expressed in coelomocytes (Fig. S2A). Second, impairing intestinal DCV secretion (by either *aex-4*/SNAP25 or *aex-6*/Rab27 mutations (Lin-Moore et al. 2021; Mahoney et al. 2006, 2008; Thomas 1990), blocked the juglone-induced increase in coelomocyte fluorescence in FLP-2::Venus-expressing animals (Fig. 2C). Third, *nlp-40* and *nlp-27* encode neuropeptide-like proteins that are released from the intestine but are not implicated in stress responses (Liu et al. 2023; Taylor et al. 2021; Wang et al. 2013). Juglone treatment had no detectable effects on coelomocyte fluorescence in animals expressing intestinal NLP-40::Venus or NLP-27::Venus fusion proteins (Fig. S2B and C), and NLP-40::mTur2 puncta did not overlap with FLP-2::Venus puncta in the intestine (Fig. S2D). Finally, *flp-1* mutants exhibited wild type levels of FLP-2 secretion both in the absence and presence of juglone (Fig. 2E). The distribution of FLP-2::Venus puncta in the intestine was not detectably altered by juglone treatment. Together, these results indicate that acute oxidative stress selectively increases the exocytosis of FLP-2-containing DCVs from the intestine, upstream of *flp-1* signaling.

SOD-1 and SOD-3 superoxide dismutases regulate FLP-2 release.

Juglone generates superoxide anion radicals (Ahmad and Suzuki 2019; Paulsen and Ljungman 2005) and juglone treatment of *C. elegans* increases ROS levels (de Castro, Hegi de Castro, and Johnson 2004) likely by promoting the global production of mitochondrial superoxide. Superoxide can then be rapidly converted into H_2O_2 by superoxide dismutase. To determine whether H_2O_2 impacts FLP-2 secretion, we first examined superoxide dismutase mutants. *C. elegans* encodes five superoxide dismutase genes (*sod-1* through *sod-5*). *sod-1* or *sod-3* null mutations blocked juglone-induced FLP-2 secretion without altering baseline FLP-2 secretion, whereas *sod-2*, *sod-4*, or *sod-5* mutations had no effect on FLP-2 secretion in the presence of juglone (Fig. 3A and B, S3A). *sod-1; sod-3* double mutants exhibited juglone-induced FLP-2 secretion defects that was similar to single mutants, without significantly altering FLP-2 secretion in the absence of stress (Fig. 3C). *sod-1* encodes the ortholog of mammalian SOD1, which is a cytoplasmic SOD implicated in the development of amyotrophic lateral sclerosis (ALS) and cancer (Giglio et al. 1994; Papa, Manfredi, and Germain 2014; Wang et al. 2021; Zhang et al. 2007). SOD-1::fusion proteins adopted a diffuse pattern of fluorescence in intestinal cells, consistent with a cytoplasmic localization (Fig. 3D). Transgenes expressing the *sod-1* cDNA selectively in the intestine fully rescued the juglone-induced FLP-2::Venus secretion defects of *sod-1* mutants (Fig. 3A). *sod-3* encodes a homolog of

mammalian SOD2, which is a mitochondrial matrix SOD implicated in protection against oxidative stress-induced neuronal cell death (Fukui and Zhu 2010; Vincent et al. 2007). Intestinal SOD-3::GFP fusion proteins localized to round structures that were surrounded by the outer membrane mitochondrial marker TOMM-20::mCherry (Ahier et al. 2018), consistent with a mitochondrial matrix localization (Fig. 3E). Expression of *sod-3* cDNA in the intestine fully restored juglone-induced FLP-2 release to *sod-3* mutants (Fig. 3B). *sod-3* variants lacking the mitochondrial localization sequence (*sod-3*(ΔMLS)), were no longer localized to mitochondria (Fig. 3F) and failed to restore normal responsiveness to juglone to *sod-3* mutants (Fig. 3B). Thus, the generation of H_2O_2 by either SOD-1 in the cytoplasm or by SOD-3 in the mitochondrial matrix is necessary for juglone to increase FLP-2 secretion.

Next, to determine if H_2O_2 can regulate FLP-2 secretion, we treated animals acutely with exogenous H_2O_2. We found that 10 minute H_2O_2 treatment increased FLP-2::Venus secretion to a similar extent as 10 minute juglone treatment. *aex-4*/SNAP25*, or *aex-6/*Rab27 mutants exhibited no increase in FLP-2 secretion in response to H_2O_2 treatment compared to untreated controls (Fig. 3G). In contrast, *sod-1* or *sod-3* mutants (or *sod-1; sod-3* double mutants) exhibited an increase in FLP-2 secretion in response to H_2O_2 that was similar to that of wild type controls (Fig. 3H), suggesting that exogenous H_2O_2 can bypass the requirement of SODs but not SNAREs to promote FLP-2 secretion. Together these results suggest that H_2O_2 generated by SODs can positively regulate intestinal FLP-2 exocytosis from DCVs (Fig. 3I).

SOD-1 and SOD-3 regulate intestinal mitochondrial H_2O_2 levels.

To directly monitor H_2O_2 levels in the intestine, we generated transgenic animals in which HyPer7 was targeted to either the mitochondrial matrix (matrix-HyPer7) by generating fusion proteins with the cytochrome c MLS, or to the cytosolic face of the outer mitochondrial membrane (OMM-HyPer7) by generating fusion proteins with TOMM-20. When co-expressed in the intestine with the OMM marker TOMM-20::mCherry, matrix-HyPer7 formed round structures throughout the cytoplasm that were surrounded by the OMM, and OMM-HyPer7 formed ring-like structures throughout the cytoplasm that co-localized with the OMM marker (Fig. 3J). Ten minute treatment with H_2O_2 significantly increased the fluorescence intensity by about two-fold of both matrix-HyPer7 and OMM-HyPer7 without altering mitochondrial morphology or abundance (Fig. 3J, S3B and C), validating the utility of HyPer7 as a sensor for acute changes in H_2O_2 levels in and around intestinal mitochondria.

Juglone treatment for 10 minutes led to a similar two-fold increase in matrix-HyPer7 fluorescence as H_2O_2 treatment (Fig. 3J). *sod-3* mutations did not alter baseline H_2O_2 levels in the matrix, but they completely blocked juglone-induced increases H_2O_2 levels, whereas *sod-1* mutations had no effect on either baseline or juglone-induced increases H_2O_2 levels (Fig. 3J). These results indicate that superoxide produced by juglone treatment is likely to be converted into H_2O_2 by SOD-3 in the mitochondrial matrix (Fig. 3I).

Next, we examined H_2O_2 levels on the outer surface of mitochondria using OMM-HyPer7 and we found that juglone treatment led to a two-fold increase in OMM-HyPer7 fluorescence, similar to H_2O_2 treatment (Fig. 3J). *sod-3* or *sod-1* mutations did not alter

baseline H_2O_2 levels on the OMM, but *sod-1* single mutations attenuated, juglone-induced
increases in OMM H_2O_2 levels, while *sod-3* mutations had no effect (Fig. 3J). In *sod-1;*
sod-3 double mutants, the juglone-induced increase in OMM H_2O_2 levels was completely
blocked, whereas baseline H_2O_2 levels in the absence of stress were unchanged (Fig.
3J). These results suggest that *sod-3* and *sod-1* are exclusively required for H_2O_2
production by juglone and that both mitochondrial SOD-3 and cytosolic SOD-1 contribute
to H_2O_2 levels in the cytosol. One model that could explain these results is that juglone-
generated superoxide is converted into H_2O_2 both by SOD-3 in the matrix, and by SOD-
1 in the cytosol, and that the H_2O_2 generated in the matrix can exit the mitochondria to
contribute to cytosolic H_2O_2 levels needed to drive FLP-2 secretion (Fig. 3I).
345
346 **The peroxiredoxin-thioredoxin system regulates endogenous H_2O_2 levels and FLP-**
347 **2 secretion.**
348
To determine whether endogenous H_2O_2 regulates FLP-2 secretion, we examined
mutations in the peroxiredoxin-thioredoxin system. Peroxiredoxins and thioredoxins
detoxify excessive H_2O_2 by converting it into water and they play a critical role in
maintaining cellular redox homeostasis (Netto and Antunes 2016) (Fig. 4A). *C. elegans*
encodes two peroxiredoxin family members, *prdx-2* and *prdx-3*, that are expressed at
high levels in the intestine (Taylor et al. 2021). Null mutations in *prdx-2* significantly
increased FLP-2::Venus secretion compared to wild type animals in the absence of stress
(Fig. 4B), whereas null mutations in *prdx-3* had no effect on FLP-2 secretion (Fig. S4A).
We observed a corresponding increase in both matrix-HyPer7 and OMM-HyPer7
fluorescence intensity in *prdx-2* mutants (Fig. 4C and D), demonstrating that endogenous
H_2O_2 is neutralized by peroxiredoxin and establishing a correlation between endogenous
H_2O_2 levels and FLP-2 secretion. The increase in FLP-2 secretion in *prdx-2* mutants was
not further increased by juglone treatment (Fig. 4B). These results suggest that elevation
in the levels of endogenously produced H_2O_2 in the intestine can positively regulate FLP-
2 secretion.
There are three isoforms of *prdx-2* that arise by the use of alternative
transcriptional start sites (Fig. 4A). Expressing the *prdx-2b* isoform selectively in the
intestine fully rescued the elevated FLP-2::Venus secretion defects of *prdx-2* mutants,
whereas expressing *prdx-2a* or *prdx-2c* isoforms failed to rescue (Fig. 4B, S4B and C).
To independently verify the role of *prdx-2b* function in FLP-2 release, we generated a
prdx-2b-specific knockout mutant by introducing an in frame stop codon within the *prdx-*
2b-specific exon 1 using CRISPR/Cas9 (*prdx-2b(vj380)* Fig. 4A). *prdx-2b(vj380)* mutants
exhibited increased H_2O_2 levels in the mitochondrial matrix and OMM (Fig. 4C and D), as
well as increased FLP-2::Venus secretion compared to wild type controls that were
indistinguishable from *prdx-2* null mutants (Fig. 4B). *prdx-2b* mutations could no longer
increase FLP-2 secretion when either *sod-1* or *sod-3* activity was impaired (Fig. 4E and
Fig. S4D). Thus, the *prdx-2b* isoform normally inhibits FLP-2 secretion likely by promoting
the consumption of H_2O_2 in the mitochondrial matrix and/or cytosol.
Once oxidized, peroxiredoxins are reduced by thioredoxins (TRXs) for reuse (Netto
and Antunes 2016). TRX-3 is an intestine-specific thioredoxin promoting protection
against specific pathogen infections (Jiménez-Hidalgo et al. 2014; Miranda-Vizuete,
Damdimopoulos, and Spyrou 2000; Netto and Antunes 2016). Mutations in *trx-3* elevated

FLP-2::Venus release in the absence of juglone and expressing *trx-3* transgenes in the intestine restored wild type FLP-2 release to *trx-3* mutants (Fig. 4B). Juglone treatment failed to further enhance FLP-2::Venus release in *trx-3* mutants (Fig. 4B). Mutations in cytoplasmic *sod-1* but not in mitochondrial *sod-3* reduced the elevated FLP-2::Venus release in *trx-3* mutants to wild type levels (Fig. 4B). Mutations in *trx-3* increased H_2O_2 levels in the OMM but had no effect on matrix H_2O_2 levels (Fig. 4C and D). Thus TRX-3 likely functions in the cytosol but not in the matrix to neutralize H_2O_2, and elevated H_2O_2 levels in the cytosol are sufficient to drive FLP-2 secretion without SOD-3-mediated H_2O_2 generation in the matrix.

Finally, to investigate the physiological significance of elevated endogenous H_2O_2 levels on the oxidative stress response we examined the effects of *prdx-2b* mutations on expression of *gst-4*. *prdx-2b* mutants had significantly increased P*gst-4::gfp* expression in the intestine compared to wild type controls (Fig. 4F). The increased P*gst-4::gfp* expression in *prdx-2b* mutants was completely dependent upon *flp-2* signaling, since *gst-4* expression was reduced to wild type levels in *prdx-2b; flp-2* double mutants (Fig. 4F). Together our data suggest that *prdx-2b* functions in the intestine to maintain redox homeostasis following SOD-1/SOD-3 mediated H_2O_2 production by regulating the secretion of FLP-2 (Fig. 3I).

PKC-2/PKCα/β mediates H_2O_2 induced FLP-2 secretion from the intestine.

H_2O_2 functions as a cellular signaling molecule by oxidizing reactive cysteines to sulfenic acid, and this modification on target proteins can regulate intracellular signaling pathways (García Santamarina, Boronat i Llop, and Hidalgo Hernando 2014). One of the validated targets of H_2O_2 signaling is the protein kinase C (PKC) family of serine threonine kinases (Jia and Sieburth 2021; Konishi et al. 1997, 2001; Min, Kim, and Exton 1998). *C. elegans* encodes four PKC family members including *pkc-1* and *pkc-2*, which are expressed at highest levels in the intestine (Islas-Trejo et al. 1997; Taylor et al. 2021). *pkc-1* null mutants had no effect on baseline or juglone-induced FLP-2 secretion (Fig. S5A). *pkc-2* null mutations did not alter baseline intestinal FLP-2 secretion, but they eliminated juglone-induced FLP-2 secretion (Fig. 5A). *pkc-2* encodes a calcium and diacylglycerol (DAG) stimulated PKCα/β protein kinase C that regulates thermosensory behavior by promoting transmitter secretion (Edwards et al. 2012; Land and Rubin 2017). Expressing *pkc-2* cDNA selectively in the intestine fully restored juglone-induced FLP-2 secretion to *pkc-2* mutants (Fig. 5A), whereas expressing a catalytically inactive *pkc-2(K375R)* variant (Van et al. 2021) failed to rescue (Fig. 5A). The intestinal site of action of *pkc-2* is in line with prior studies showing that *pkc-2* can function in the intestine to regulate thermosensory behavior (Land and Rubin 2017). *pkc-2* mutants exhibited wild type H_2O_2 levels in the mitochondrial matrix and OMM of the intestine in both the presence and absence of juglone (Fig. 5B and C). Increasing H_2O_2 levels by either acute H_2O_2 treatment or by *prdx-2* mutation failed to increase FLP-2 secretion in *pkc-2* mutants (Fig. 5D and E). To determine whether *pkc-2* can regulate the intestinal secretion of other peptides that are not associated with oxidative stress , we examined expulsion frequency, which is a measure of intestinal NLP-40 secretion (Mahoney et al. 2008; Wang et al. 2013). *pkc-2* mutants showed wild type expulsion frequency (Fig. S5B), indicating that intestinal NLP-40 release is largely unaffected. Together, these results show that *pkc-2*

427 is not a general regulator of intestinal peptide secretion and instead functions downstream
428 or in parallel to H_2O_2 to selectively promote FLP-2 secretion by a mechanism that involves
429 phosphorylation of target proteins.
430
431 **DAG positively regulates FLP-2 secretion.**
432
PKCα/β family members contain two N terminal C1 domains (C1A and C1B) whose
binding to DAG promotes PKC recruitment to the plasma membrane (Burns and Bell 1991;
Darby, Meng, and Fehrenbacher 2017; Johnson, Giorgione, and Newton 2000; Kim et al.
2016; Ono et al. 1989; Yanase et al. 2011). To address the role of DAG in promoting FLP-
2 secretion by PKC-2, we examined mutants that are predicted to have altered DAG levels.
Phosphatidylinositol phospholipase C beta (PLCβ) converts phosphatidyl inositol
phosphate (PIP2) to DAG and inositol triphosphate (IP3, Fig. 6A), and impairing PLC
activity leads to reduced cellular DAG levels (Nebigil 1997). *C. elegans* encodes two PLC
family members whose expression is enriched in the intestine, *plc-2/* PLCβ and *egl-8/*
PLCβ (Taylor et al. 2021). *plc-2* null mutants exhibited baseline and juglone-included FLP-
2 secretion that were similar to wild type controls (Fig. S6). *egl-8* loss-of-function mutants
exhibited wild type baseline FLP-2 secretion, but juglone-induced FLP-2 secretion was
completely blocked (Fig. 6B). H_2O_2 levels in *egl-8* mutants were similar to wild type
controls, both in the presence and absence of juglone (Fig. 6C and D). Thus, *egl-8/*PLCβ
functions downstream of or in parallel to H_2O_2 production to promote FLP-2 secretion.
DAG kinase converts DAG into phosphatidic acid (PA), and is therefore a negative
regulator or DAG levels (Fig. 6A) (van Blitterswijk and Houssa 2000; Topham 2006). In
C. elegans, *dgk-2*/DGKε is the highest expressing DAG kinase in the intestine. Mutations
in *dgk-2* elevated FLP-2::Venus secretion (Fig. 6E) without altering H_2O_2 levels in the
intestinal mitochondrial matrix or OMM (Fig. 6F and G). Expressing *dgk-2* transgenes
selectively in the intestine restored normal FLP-2::Venus secretion to *dgk-2* mutants (Fig.
6E). Finally, the increase in FLP-2 secretion in *dgk-2* mutants was not further increased
by juglone treatment, but it was completely blocked by *pkc-2* mutations or *aex-4*/SNAP25
mutations (Fig. 6E and H). These results show that FLP-2 secretion can be regulated
bidirectionally by DAG, and they suggest that DAG and H_2O_2 function in a common
genetic pathway upstream of *pkc-2* to promote FLP-2 secretion (Fig. 6A).
459
460
461 **Discussion**
462
By screening for intercellular regulators of FLP-1 signaling from the nervous
system in promoting the anti-oxidant response, we have uncovered a function for
peptidergic signaling in mediating gut-to-neuron regulation of the anti-oxidant response
in *C. elegans.* We identified the neuropeptide-like protein FLP-2 as an inter-tissue signal
originating in the intestine to potentiate stress-induced FLP-1 release from AIY neurons
and the subsequent activation of SKN-1 in the intestine. We found that H_2O_2 generated
endogenously in the intestine or exogenously by acute oxidant exposure increases FLP-
2 secretion from intestinal DCVs. H_2O_2 promotes FLP-2 exocytosis through PKC-2, and
AEX-4/SNAP25. The use of oxidant-regulated peptide secretion exemplifies a
mechanism that can allow the gut and the nervous system to efficiently and rapidly

communicate through endocrine signaling to promote organism-wide protection in the
face of intestinal stress (Fig. 6I).
475
476 **A new function for *flp-2* signaling in the antioxidant response.**
477
Previous studies have identified roles for *flp-2* signaling in development and in
stress responses. *flp-2* promotes locomotion during molting (Chen et al. 2016) promotes
entry into reproductive growth (Chai et al. 2022), regulates longevity (Kageyama et al.
2022), and activates the UPR^{mt} cell-non autonomously during mitochondrial stress (Shao
et al. 2016). The function we identified for *flp-2* in the antioxidant response has some
notable similarities with *flp-2*'s other functions. First, *flp-2* mediates its effects at least in
part by regulating signaling by other peptides. *flp-2* signaling increases the secretion of
the neuropeptide like protein PDF-1 during lethargus (Chen et al. 2016) and INS-
35/insulin-like peptide for its roles in reproductive growth choice and longevity (Kageyama
et al. 2022), in addition to regulating AIY FLP-1 secretion (Fig. 5G).. Second, the secretion
of FLP-2 is dynamic. FLP-2 secretion decreases during lethargus (Chen et al. 2016) and
increases under conditions that do not favor reproductive growth (Kageyama et al. 2022),
as well increasing in response to oxidants (Fig. 2C and D). However, in some instances,
the regulation of FLP-2 secretion may occur at the level of flp-2 expression (Kageyama
et al. 2022), rather than at the level of exocytosis (Fig. 2C). Finally, genetic analysis of *flp-
2* has revealed that under normal conditions, *flp-2* signaling may be relatively low, since
flp-2 mutants show no defects in reproductive growth choice when animals are well fed
(Chai et al. 2022), show only mild defects in locomotion during molting in non-sensitized
genetic backgrounds (Chen et al. 2016), and do not have altered baseline FLP-1
secretion or antioxidant gene expression in the absence of exogenous oxidants (Fig. 1D
and E). It is notable that increased ROS levels are associated with molting (Back et al.
2012; Knoefler et al. 2012), ageing (Back et al. 2012; Van Raamsdonk and Hekimi 2010),
starvation (Tao et al. 2017), and mitochondrial dysfunction (Dingley et al. 2010), raising
the possibility that *flp-2* may be used in specific contexts associated with high ROS levels
to affect global changes in physiology, behavior and development.
One major difference we found for *flp-2* signaling in our study is that intestinal, but
not neuronal *flp-2* activates the oxidative stress response, whereas *flp-2* originates from
neurons for its reported roles in development and the UPR^{mt}. The intestine is uniquely
poised to relay information about diet to the rest of the animal, and secretion of a number
of neuropeptide-like proteins from the intestine (e.g, INS-11, PDF-2 and INS-7) is
proposed to regulate responses to different bacterial food sources (Lee and Mylonakis
2017; Murphy, Lee, and Kenyon 2007; O'Donnell et al. 2018). Since bacterial diet can
impact ROS levels in the intestine (Pang and Curran 2014), secretion of FLP-2 from the
intestine could function to relay information about bacterial diet to distal tissues to regulate
redox homeostasis. In addition, the regulation of intestinal FLP-2 release by oxidants may
meet a unique spatial, temporal or concentration requirement for activating the antioxidant
response that cannot be met by its release from the nervous system.
515
516 **AIY as a target for *flp-2* signaling.**
517

AIY interneurons receive sensory information from several neurons primarily as glutamatergic inputs to regulate behavior (Bargmann et al. 2007; Clark et al. 2006; Satoh et al. 2014). Our study reveals a previously undescribed mechanism by which AIY is activated through endocrine signaling originating from FLP-2 secretion from the intestine. FLP-2 could act directly on AIY, or it may function indirectly through upstream neurons that relay FLP-2 signals to AIY. *frpr-18* encodes an orexin-like GPCR that can be activated by FLP-2-derived peptides in transfected mammalian cells (Larsen et al. 2013; Mertens et al. 2005), and *frpr-18* functions downstream of *flp-2* in the locomotion arousal circuit (Chen et al. 2016). *frpr-18* is expressed broadly in the nervous system including in AIY (Chen et al. 2016), and loss-of-function *frpr-18* mutations lead to hypersensitivity to certain oxidants (Ouaakki et al. 2023). FRPR-18 is coupled to the heterotrimeric G protein Gαq (Larsen et al. 2013; Mertens et al. 2005), raising the possibility that FLP-2 may promote FLP-1 secretion from AIY by directly activating FRPR-18 in AIY. However, *flp-2* functions independently of *frpr-18* in the reproductive growth circuit, and instead functions in a genetic pathway with the GPCR *npr-30* (Chai et al. 2022). In addition, FLP-2-derived peptides (of which there are at least three) can bind to the GPCRs DMSR-1, or FRPR-8 in transfected cells (Beets et al. 2023). Identifying the relevant FLP-2 peptide(s), the FLP-2 receptor and its site of action will help to define the circuit used by intestinal *flp-2* to promote FLP-1 release from AIY.

FLP-1 release from AIY is positively regulated by H_2O_2 generated from mitochondria (Jia and Sieburth 2021). Here we showed that H_2O_2-induced FLP-1 release requires intestinal *flp-2* signaling. However, *flp-2* does not appear to promote FLP-1 secretion by increasing H_2O_2 levels in AIY (Fig 1E), and *flp-2* signaling is not sufficient to promote FLP-1 secretion in the absence of H_2O_2 (Fig. 1D). These results point to a model whereby at least two conditions must be met in order for AIY to increase FLP-1 secretion: an increase in H_2O_2 levels in AIY itself, and an increase in *flp-2* signaling from the intestine. Thus AIY integrates stress signals from both the nervous system and the intestine to activate the intestinal antioxidant response through FLP-1 secretion. The requirement of signals from multiple tissues for FLP-1 secretion may function to limit the activation of SKN-1, since unregulated SKN-1 activation can be detrimental to organismal health (Turner, Ramos, and Curran 2024). AIY shows a sporadic Ca^{2+} response regardless of the presence of explicit stimulation (Ashida, Hotta, and Oka 2019; Bargmann et al. 2007; Clark et al. 2006), and FLP-1 secretion from AIY is calcium dependent (Jia and Sieburth 2021)How mitochondrial H_2O_2 levels are established in AIY by intrinsic or extracellular inputs, and how AIY integrates H_2O_2 and *flp-2* signaling to control FLP-1 secretion remain to be defined.

A role for endogenous H_2O_2 in regulated neuropeptide secretion.

Using HyPer7, we showed that acute juglone exposure results in a rapid elevation of endogenous H_2O_2 levels inside and outside intestinal mitochondria and a corresponding increase of FLP-2 release from the intestine that depends on the cytoplasmic superoxide dismutase *sod-1*, and mitochondrial *sod-3*. We favor a model whereby superoxide generated by juglone in the mitochondria is converted to H_2O_2 by SOD-3 in the matrix and by SOD-1 in the cytosol. In this case, both the superoxide generated by juglone and the H_2O_2 generated by SOD-3 would have to be able to exit the

mitochondria and enter the cytosol. Superoxide and H_2O_2 can be transported across
mitochondrial membranes through anion channels and aquaporin channels, respectively
(Bienert and Chaumont 2014; Ferri et al. 2003; Han et al. 2003; Kontos et al. 1985). The
observation that both SOD-1 and SOD-3 activity are necessary to drive FLP-2 release
suggests that H_2O_2 levels much reach a certain threshold in the cytoplasm to promote
FLP-2 release, and this threshold requires the generation of H_2O_2 by both SOD-1 and
SOD-3.
We identified a role for the antioxidant peroxiredoxin-thioredoxin system, encoded
by *prdx-2* and *trx-3*, in maintaining low endogenous H_2O_2 levels in the intestine and in
negatively regulating FLP-2 secretion. We showed that the *prdx-2b* isoform functions to
inhibit FLP-2 secretion and to lower H_2O_2 levels in both the mitochondrial matrix and on
the cytosolic side of mitochondria. These observations are consistent with a subcellular
site of action for PRDX-2B in either the matrix only or in both the matrix and cytosol. In
contrast, *trx-3* mutations do not alter mitochondrial H_2O_2 levels, suggesting that TRX-3
functions exclusively in the cytosol. Thus, the PRDX-2B-TRX-3 combination may function
in the cytosol, and PRDX-2B may function with a different TRX family member in the
matrix. There are several thioredoxin-domain containing proteins in addition to *trx-3* in the
C. elegans genome (including *trx-5*/NXNL2) that could be candidates for this role.
Alternatively, *prdx-2* may function alone or with other redox proteins. PRDX-2 may
function without thioredoxins in its roles in light sensing and stress response in worms (Li
et al. 2016; Oláhová et al. 2008; Oláhová and Veal 2015). PRDX-2B contains a unique N
terminal domain that is distinct from the catalytic domain and is not found on the other
PRDX-2 isoforms. This domain may be important for targeting PRDX-2B to specific
subcellular location(s) where it can regulate FLP-2 secretion.

Regulation of FLP-2 exocytosis by PKC-2/PKCα/β and AEX-4/SNAP25.

We demonstrated that *pkc-2* mediates the effects of H_2O_2 on intestinal FLP-2
secretion, and H_2O_2- and DAG-mediated PKC-2 activation are likely to function in a
common genetic pathway to promote FLP-2 secretion. Our observations that DAG is
required for the effects of juglone (Fig. 6B), are consistent with a two-step activation model
for PKC-2, in which H_2O_2 could first modify PKC-2 in the cytosol, facilitating subsequent
PKC-2 recruitment to the membrane by DAG. Alternatively, DAG could first recruit PKC-
2 to membranes, where it is then modified by H_2O_2. We favor a model whereby H_2O_2
modification occurs in the cytosol, since H_2O_2 produced locally by mitochondria would
have access to cytosolic pools of PKC-2 prior to its membrane translocation.
We defined a role for *aex-4*/SNAP25 in the fusion step of FLP-2 containing DCVs
from the intestine under normal conditions as well as during oxidative stress. In
neuroendocrine cells, phosphorylation of SNAP25 on Ser187 potentiates DCV
recruitment into releasable pools (Nagy et al. 2002; Shu et al. 2008; Yang et al. 2007),
and exocytosis stimulated by the DAG analog phorbol ester (Gao et al. 2016; Shu et al.
2008), without altering baseline SNAP25 function. Interestingly, the residue
corresponding to Ser187 is conserved in AEX-4, raising the possibility that PKC-2
potentiates FLP-2 secretion by phosphorylating AEX-4. Since SNAP25 phosphorylation
on Ser187 has been shown to increase its interaction with syntaxin and promote SNARE
complex assembly in vitro (Gao et al. 2016; Yang et al. 2007), it is possible that elevated

H$_2$O$_2$ levels could promote FLP-2 secretion by positively regulating SNARE-mediated
DCV fusion at intestinal release sites on the basolateral membrane through AEX-
4/SNAP25 phosphorylation by PKC-2. Prior studies have shown that PKC-2
phosphorylates the SNARE-associated protein UNC-18 in neurons to regulate
thermosensory behavior (Edwards et al. 2012; Land and Rubin 2017). Thus, PKC-2 may
have multiple targets in vivo and target selection may be dictated by cell type and/or the
redox status of the cell.
617
618 **Similar molecular mechanisms regulating FLP-1 and FLP-2 release.**
619
The molecular mechanisms we identified that regulate FLP-2 secretion from the
intestine are similar in several respects to those regulating FLP-1 secretion from AIY. First,
the secretion of both peptides is positively regulated by H$_2$O$_2$ originating from
mitochondria. Second, in both cases, H$_2$O$_2$ promotes exocytosis of neuropeptide-
containing DCVs by a mechanism that depends upon the kinase activity of protein kinase
C. Finally, the secretion of both peptides is controlled through the regulation of H$_2$O$_2$ levels
by superoxide dismutases and by the peroxiredoxin-thioredoxin system. H$_2$O$_2$-regulated
FLP-1 and FLP-2 secretion differ in the identity of the family members of some of the
genes involved. *prdx-3-trx-2* and *sod-2* family members regulate H$_2$O$_2$ levels in AIY,
whereas *prdx-2-trx-3* and *sod-1/sod-3* family members regulate H$_2$O$_2$ levels in the
intestine. In addition, *pkc-1* promotes H$_2$O$_2$ induced FLP-1 secretion from AIY whereas
pkc-2 promotes H$_2$O$_2$ -induced FLP-2 secretion from the intestine. Nonetheless, it is
noteworthy that two different cell types utilize largely similar pathways for the H$_2$O$_2$-
mediated regulation of neuropeptide release, raising the possibility that similar
mechanisms may be utilized in other cell types and/or organisms to regulate DCV
secretion in response to oxidative stress.
636
637
638 **Materials and Methods**
639
Strains and transgenic lines
C. elegans strains were maintained at 20°C in the dark on standard nematode
growth medium (NGM) plates seeded with OP50 Escherichia coli as food source, unless
otherwise indicated. All strains were synchronized by picking mid L4 stage animal either
immediately before treatment (for coelomocyte imaging and intestine imaging) or 24h
before treatment (for P*gst-4::gfp* imaging). The wild type strain was Bristol N2. Mutants
used in this study were out-crossed at least four times.
Transgenic lines were generated by microinjecting plasmid mixes into the gonads
of young adult animals following standard techniques (Mello et al. 1991). Microinjection
mixes were prepared by mixing expression constructs with the co-injection markers
pJQ70 (P*ofm-1::rfp*, 25ng/µL), pMH163 (P*odr-1::mCherry*, 40ng/µL), pMH164 (P*odr-
1::gfp*, 40ng/µL) or pDS806 (P*myo-3::mCherry*, 20ng/µL) to a final concentration of
100ng/µL. For tissue-specific expression, a 1.5kb *rab-3* promoter was used for pan-
neuronal expression (Nonet et al. 1997), a 2.0kb *ges-1* or a 3.5kb *nlp-40* promoter was
used for intestinal expression (Egan et al. 1995; Wang et al. 2013). At least three
transgenic lines were examined for each transgene, and one representative line was used

for quantification. Strains and transgenic lines used in this study are listed in the
Supplementary Table.
658

Molecular Biology
All gene expression vectors were constructed with the backbone of pPD49.26.
Promoter fragments including P*rab-3*, and P*ges-1* were amplified from genomic DNA;
genes of interest, including cDNA fragments (*aex-5, snt-5, sod-1b, sod-3, isp-1, prdx-2a,*
prdx-2b, prdx-2c, trx-3, pkc-2b, dgk-2a, aex-4) and genomic fragments (*flp-2, flp-40, nlp-*
36, nlp-27) were amplified from cDNA library and genomic DNA respectively using
standard molecular biology protocols. Expression plasmid of HyPer7 was designed based
on reported mammalian expression plasmid for HyPer7 (Pak et al. 2020) and was
synthesized by Thermo Fisher Scientific with codon optimization for gene expression in
C. elegans. Plasmids and primers used in this study are listed in the Supplementary Table.
669

Toxicity Assay
A stock solution of 50mM juglone in DMSO was freshly made on the same day of
liquid toxicity assay. 120µM working solution of juglone in M9 buffer was prepared using
stock solution before treatment. Around 60-80 synchronized adult animals were
transferred into a 1.5mL Eppendorf tube with fresh M9 buffer and washed three times,
and a final wash was done with either the working solution of juglone with or M9 DMSO
at the concentrations present in juglone-treated animals does not contribute to toxicity
since DMSO treatment alone caused no significant change in survival compared to M9-
treated controls (Fig. S1C). Animals were incubated in the dark for 4h on rotating mixer
before being transferred onto fresh NGM plates seeded with OP50 to recover in the dark
at 20°C for 16 hours. Percentage of survival was assayed by counting the number of alive
and dead animals. Toxicity assays were performed in triplicates.
682

RNAi Interference
Plates for feeding RNAi interference were prepare as described (Kamath 2003).
Around 20-25 gravid adult animals with indicated genotype were transferred onto the
RNAi plates that were seeded with HT115(DE3) bacteria transformed with L4440 vectors
with targeted gene inserts or empty L4440 vectors. Eggs were collected for 4h to obtain
synchronized populations. L4 stage animals were collected for further assays. RNAi
clones were from Ahringer or Vidal RNAi library, or made from genomic DNA. Details were
listed in the Supplementary Table.
691

Behavioral assays
The defecation motor program was assayed as previously described (Liu and
Thomas 1994). Twenty to thirty L4 animals were transferred onto a fresh NGM plate
seeded with OP50 *E. coli* and were stored in a 20°C incubator for 24 hours. After 24 hours,
ten consecutive defecation cycles were observed from three independent animals and
the mean and the standard error was calculated for each genotype. The pBoc and aBoc
steps were recorded using custom Etho software (James Thomas Lab
website:http://depts.washington.edu/jtlab/software/otherSoftware.html)
700

Microscopy and Fluorescence Imaging

Approximately 30-40 age matched animals were paralyzed with 30mg/mL 2,3-
butanedione monoxime (BDM) in M9 buffer and mounted on 2% agarose pads. Images
were captured using the Nikon eclipse 90i microscope equipped with Nikon Plan Apo 20x,
40x, 60x, and 100x oil objective (N.A.=1.40), and a Photometrics Coolsnap ES2 camera
or a Hamamatsu Orca Flash LT+CMOS camera. Metamorph 7.0 software (Universal
Imaging/Molecular Devices) was used to capture serial image stacks and to obtain the
maximum intensity projection image for analysis.
For transcriptional reporter imaging, young adult animals were transferred into a
1.5mL Eppendorf tube with M9 buffer, washed three times and incubated in 50µM working
solution of juglone or M9 buffer control with equivalent DMSO for 1h in the dark on
rotating mixer before recovering on fresh NGM plates with OP50 for 3h in the dark at
20°C. The posterior end of the intestine was imaged with the 60x objective and
quantification for average fluorescence intensity of a 16-pixel diameter circle in the
posterior intestine was calculated using Metamorph.
For coelomocyte imaging, L4 stage animals were transferred in fresh M9 buffer on
a cover slide, washed six times with M9 before being exposed to 300µM juglone in M9
buffer (diluted from freshly made 50mM stock solution), 1mM H_2O_2 in M9 buffer, or M9
buffer. DMSO at the concentrations present in juglone-treated animals does not alter
neuropeptide secretion since DMSO treatment alone caused no significant change in
FLP-1::Venus or FLP-2::Venus coelomocyte fluorescence compared to M9-treated
controls. (Fig. S1D and S2E). . Animals were then paralyzed in BDM and images of
coelomocytes next to the posterior end of intestine were taken using the 100x oil objective.
Average fluorescence intensity of Venus from the endocytic compartments in the posterior
coelomocytes was measured in ImageJ.
For fusion protein fluorescence imaging, L4 stage animals were exposed to M9
buffer or indicated oxidants for 10min before being paralyzed in BDM and images taken
of the posterior end of the intestine using 100x oil objective. For HyPer7 imaging, Z stacks
were obtained using GFP (excitation/emission: 500nm/520nm) and CFP
(excitation/emission: 400nm/520nm) filter sets sequentially, HyPer7 fluorescence signal
was quantified as the ratio of GFP to CFP fluorescence intensity changes with respect to
the baseline $[(F_t − F_0)/F_0]$.
733
CRISPR/Cas9 Editing
prdx-2b(vj380) knock-out mutants were generated using a co-CRISPR protocol
(Arribere et al. 2014). A sgRNA and a repair single stranded oligodeoxynucleotides
(ssODN) targeting *dpy-10* were co-injected with a sgRNA for genes of interest and a
ssODN that induces homology-directed repair (HDR) to introduce Cas9 mediated
mutagenesis. Fifteen young adult animals were injected to produce around thirty singled
F1 animals carrying Dpy or Rol phenotype. F2 animals were genotyped for mutations
based on PCR and enzyme digest. Homozygous mutants were outcrossed with wild type
animals at least four times before being used for assays.
743
Statistics
Statistical analysis was performed on GraphPad Prism 9. Unpaired *t*-test with two
tails was used for two groups and one-way ANOVA with multiple comparison corrections
was used to three or more groups to determine the statistical significance. Statistical

details and n are specified in the figure legends. All comparisons are conducted based on
wild-type controls unless indicated by lines between genotypes. Bar graphs with plots
were generated using GraphPad Prism 9.
751

Acknowledgements

C. elegans strains used in this study were provided by the Caenorhabditis Genetics
Centre (CGC), which is funded by the NIH National Center for Research Resources
(NCRR). We thank members of the Sieburth lab for critical reading and discussion of the
manuscript. This work was supported by grants from National Institute of Health NINDS
R01NS071085 and R01NS110730 to D.S.

Reference

A R Giniatullin and R A Giniatullin. 2003. "Dual Action of Hydrogen Peroxide on Synaptic Transmission at the Frog Neuromuscular Junction." *The Journal of Physiology* 552(1):283–93. doi: 10.1113/jphysiol.2003.050690.

Ahier, Arnaud, Chuan-Yang Dai, Andrea Tweedie, Ayenachew Bezawork-Geleta, Ina Kirmes, and Steven Zuryn. 2018. "Affinity Purification of Cell-Specific Mitochondria from Whole Animals Resolves Patterns of Genetic Mosaicism." *Nature Cell Biology* 20(3):352-. doi: 10.1038/s41556-017-0023-x.

Ahmad, Taseer, and Yuichiro J. Suzuki. 2019. "Juglone in Oxidative Stress and Cell Signaling." *Antioxidants (Basel, Switzerland)* 8(4):91. doi: 10.3390/antiox8040091.

Ailion, Michael, Mandy Hannemann, Susan Dalton, Andrea Pappas, Shigeki Watanabe, Jan Hegermann, Qiang Liu, Hsiao-Fen Han, Mingyu Gu, Morgan Q. Goulding, Nikhil Sasidharan, Kim Schuske, Patrick Hullett, Stefan Eimer, and Erik M. Jorgensen. 2014. "Two Rab2 Interactors Regulate Dense-Core Vesicle Maturation." *Neuron* 82(1):167–80. doi: 10.1016/j.neuron.2014.02.017.

Arribere, Joshua A., Ryan T. Bell, Becky XH Fu, Karen L. Artiles, Phil S. Hartman, and Andrew Z. Fire. 2014. "Efficient Marker-Free Recovery of Custom Genetic Modifications with CRISPR/Cas9 in Caenorhabditis Elegans."

Avshalumov, Marat V., Billy T. Chen, Tibor Koos, James M. Tepper, and Margaret E. Rice. 2005. "Endogenous Hydrogen Peroxide Regulates the Excitability of Midbrain Dopamine Neurons via ATP-Sensitive Potassium Channels." *Journal of Neuroscience* 25(17):4222–31. doi: 10.1523/JNEUROSCI.4701-04.2005.

Avshalumov, Marat V., and Margaret E. Rice. 2003. "Activation of ATP-Sensitive K+ (KATP) Channels by H2O2 Underlies Glutamate-Dependent Inhibition of Striatal Dopamine Release." *Proceedings of the National Academy of Sciences - PNAS* 100(20):11729–34. doi: 10.1073/pnas.1834314100.

Back, Patricia, Winnok H. De Vos, Geert G. Depuydt, Filip Matthijssens, Jacques R. Vanfleteren, and Bart P. Braeckman. 2012. "Exploring Real-Time in Vivo Redox Biology of Developing and Aging Caenorhabditis Elegans." *Free Radical Biology and Medicine* 52(5):850–59. doi: 10.1016/j.freeradbiomed.2011.11.037.

Bao, Li, Marat V. Avshalumov, and Margaret E. Rice. 2005. "Partial Mitochondrial Inhibition Causes Striatal Dopamine Release Suppression and Medium Spiny Neuron Depolarization via H2O2 Elevation, Not ATP Depletion." *The Journal of Neuroscience* 25(43):10029–40. doi: 10.1523/JNEUROSCI.2652-05.2005.

Bargmann, Cornelia I., Sreekanth H. Chalasani, Nikos Chronis, Makoto Tsunozaki, Jesse M. Gray, Daniel Ramot, and Miriam B. Goodman. 2007. "Dissecting a Circuit for Olfactory Behaviour in Caenorhabditis Elegans." *Nature* 450(7166):63–70. doi: 10.1038/nature06292.

Batterham, Rachel L., and Stephen R. Bloom. 2003. "The Gut Hormone Peptide YY Regulates Appetite." *Annals of the New York Academy of Sciences* 994(1):162–68. doi: 10.1111/j.1749-6632.2003.tb03176.x.

Batterham, Rachel L., Michael A. Cowley, Caroline J. Small, Herbert Herzog, Mark A. Cohen, Catherine L. Dakin, Alison M. Wren, Audrey E. Brynes, Malcolm J. Low, Mohammad A. Ghatei, Roger D. Cone, and Stephen R. Bloom. 2002. "Gut Hormone PYY(3-36) Physiologically Inhibits Food Intake." *Nature (London)* 418(6898):650-.

Beets, Isabel, Sven Zels, Elke Vandewyer, Jonas Demeulemeester, Jelle Caers, Esra Baytemur, Amy Courtney, Luca Golinelli, İlayda Hasakioğulları, William R. Schafer, Petra E. Vértes, Olivier Mirabeau, and Liliane Schoofs. 2023. "System-Wide Mapping of Peptide-GPCR Interactions in C. Elegans." *Cell Reports (Cambridge)* 42(9):113058–113058. doi: 10.1016/j.celrep.2023.113058.

Bienert, Gerd P., and François Chaumont. 2014. "Aquaporin-Facilitated Transmembrane Diffusion of Hydrogen Peroxide." *Biochimica et Biophysica Acta* 1840(5):1596–1604. doi: 10.1016/j.bbagen.2013.09.017.

van Blitterswijk, Wim J., and Brahim Houssa. 2000. "Properties and Functions of Diacylglycerol Kinases." *Cellular Signalling* 12(9):595–605. doi: 10.1016/S0898-6568(00)00113-3.

Botteaux, A., C. Hoste, J. E. Dumont, J. Van Sande, and A. Allaoui. 2009. "Potential Role of Noxes in the Protection of Mucosae: H(2)O(2) as a Bacterial Repellent." *Microbes and Infection* 11(5):537–44. doi: 10.1016/j.micinf.2009.02.009.

Browning, Kirsteen N., and R. Alberto Travagli. 2014. "Central Nervous System Control of Gastrointestinal Motility and Secretion and Modulation of Gastrointestinal Functions." *Comprehensive Physiology* 4(4):1339–68. doi: 10.1002/cphy.c130055.

Burns, D. J., and R. M. Bell. 1991. "Protein Kinase C Contains Two Phorbol Ester Binding Domains." *The*
Journal of Biological Chemistry 266(27):18330–38. doi: 10.1016/S0021-9258(18)55274-1.
Campos-Salinas, Jenny, Antonio Cavazzuti, Francisco O'Valle, Irene Forte-Lago, Marta Caro, Stephen M.
Beverley, Mario Delgado, and Elena Gonzalez-Rey. 2014. "Therapeutic Efficacy of Stable Analogues of
Vasoactive Intestinal Peptide against Pathogens." *The Journal of Biological Chemistry* 289(21):14583–99.
doi: 10.1074/jbc.M114.560573.
Carabotti, Marilia, Annunziata Scirocco, Maria Antonietta Maselli, and Carola Severi. 2015. "The Gut-Brain
Axis: Interactions between Enteric Microbiota, Central and Enteric Nervous Systems." *Annals of*
Gastroenterology : Quarterly Publication of the Hellenic Society of Gastroenterology 28(2):203–9.
Castello, Pablo R., Derek A. Drechsel, and Manisha Patel. 2007. "Mitochondria Are a Major Source of
Paraquat-Induced Reactive Oxygen Species Production in the Brain." *The Journal of Biological Chemistry*
282(19):14186–93. doi: 10.1074/jbc.M700827200.
de Castro, Edouard, Sarah Hegi de Castro, and Thomas E. Johnson. 2004. "Isolation of Long-Lived Mutants
in Caenorhabditis Elegans Using Selection for Resistance to Juglone." *Free Radical Biology & Medicine*
37(2):139–45. doi: 10.1016/j.freeradbiomed.2004.04.021.
Chai, Cynthia M., Mahdi Torkashvand, Maedeh Seyedolmohadesin, Heenam Park, Vivek Venkatachalam,
and Paul W. Sternberg. 2022. "Interneuron Control of C. Elegans Developmental Decision-Making." *Current*
Biology 32(10):2316-2324.e4. doi: 10.1016/j.cub.2022.03.077.
Chance, B., H. Sies, and A. Boveris. 1979. "Hydroperoxide Metabolism in Mammalian Organs."
Physiological Reviews 59(3):527–605. doi: 10.1152/physrev.1979.59.3.527.
Chelikani, Prasanth K., Alvin C. Haver, and Roger D. Reidelberger. 2005. "Intravenous Infusion of Peptide
YY(3–36) Potently Inhibits Food Intake in Rats." *Endocrinology (Philadelphia)* 146(2):879–88. doi:
10.1210/en.2004-1138.
Chen, Billy T., Marat V. Avshalumov, and Margaret E. Rice. 2001. "H2O2 Is a Novel, Endogenous Modulator
of Synaptic Dopamine Release." *Journal of Neurophysiology* 85(6):2468–76. doi:
10.1152/jn.2001.85.6.2468.
Chen, Billy T., Marat V. Avshalumov, and Margaret E. Rice. 2002. "Modulation of Somatodendritic Dopamine
Release by Endogenous H2O2: Susceptibility in Substantia Nigra But Resistance in VTA." *Journal of*
Neurophysiology 87(2):1155–58. doi: 10.1152/jn.00629.2001.
Chen, Didi, Kelsey P. Taylor, Qi Hall, and Joshua M. Kaplan. 2016. "The Neuropeptides FLP-2 and PDF-1
Act in Concert To Arouse Caenorhabditis Elegans Locomotion." *Genetics (Austin)* 204(3):1151–59. doi:
10.1534/genetics.116.192898.
Chen, Yu A., and Richard H. Scheller. 2001. "SNARE-Mediated Membrane Fusion." *Nature Reviews.*
Molecular Cell Biology 2(2):98–106. doi: 10.1038/35052017.
Ch'ng, Queelim, Derek Sieburth, and Joshua M. Kaplan. 2008. "Profiling Synaptic Proteins Identifies
Regulators of Insulin Secretion and Lifespan." *PLoS Genetics* 4(11):e1000283. doi:
10.1371/journal.pgen.1000283.
Clark, Damon A., David Biron, Piali Sengupta, and Aravinthan D. T. Samuel. 2006. "The AFD Sensory
Neurons Encode Multiple Functions Underlying Thermotactic Behavior in Caenorhabditis Elegans." *The*
Journal of Neuroscience 26(28):7444–51. doi: 10.1523/JNEUROSCI.1137-06.2006.
Corcionivoschi, Nicolae, Luis AJ Alvarez, Thomas H. Sharp, Monika Strengert, Abofu Alemka, Judith
Mantell, Paul Verkade, Ulla G. Knaus, and Billy Bourke. 2012. "Mucosal Reactive Oxygen Species
Decrease Virulence by Disrupting Campylobacter Jejuni Phosphotyrosine Signaling." *Cell Host & Microbe*
12(1):47–59. doi: 10.1016/j.chom.2012.05.018.
Darby, Lisa M., Hongdi Meng, and Jill C. Fehrenbacher. 2017. "Paclitaxel Inhibits the Activity and Membrane
Localization of PKCα and PKCβI/II to Elicit a Decrease in Stimulated Calcitonin Gene-Related Peptide
Release from Cultured Sensory Neurons." *Molecular and Cellular Neurosciences* 82:105–17. doi:
10.1016/j.mcn.2017.04.001.
Dingley, Stephen, Erzsebet Polyak, Richard Lightfoot, Julian Ostrovsky, Meera Rao, Todd Greco, Harry
Ischiropoulos, and Marni J. Falk. 2010. "Mitochondrial Respiratory Chain Dysfunction Variably Increases
Oxidant Stress in Caenorhabditis Elegans." *Mitochondrion* 10(2):125–36. doi: 10.1016/j.mito.2009.11.003.
Edwards, Mark R., James R. Johnson, Kimberley Rankin, Rosalind E. Jenkins, Carl Maguire, Alan Morgan,
Robert D. Burgoyne, and Jeff W. Barclay. 2012. "PKC-2 Phosphorylation of UNC-18 Ser322 in AFD Neurons
Regulates Temperature Dependency of Locomotion." *The Journal of Neuroscience* 32(20):7042–51. doi:
10.1523/JNEUROSCI.4029-11.2012.

Edwards, Samantha Louise, Lucas Mergan, Bhavesh Parmar, Bram Cockx, Wouter De Haes, Liesbet Temmerman, and Liliane Schoofs. 2019. "Exploring Neuropeptide Signalling through Proteomics and Peptidomics." *Expert Review of Proteomics* 16(2):131–37. doi: 10.1080/14789450.2019.1559733.

Egan, Cay R., May A. Chung, Fran L. Allen, Mark FP Heschl, Cheryl L. Van Buskirk, and James D. McGhee. 1995. "A Gut-to-Pharynx/Tail Switch in Embryonic Expression of the Caenorhabditis Elegans Ges-1 Gene Centers on Two GATA Sequences." *Developmental Biology* 170(2):397–419. doi: 10.1006/dbio.1995.1225.

Elferink, Jan GR. 1999. "Thimerosal A Versatile Sulfhydryl Reagent, Calcium Mobilizer, and Cell Function-Modulating Agent." *General Pharmacology* 33(1):1–6. doi: 10.1016/S0306-3623(98)00258-4.

Fares, Hanna, and Iva Greenwald. 2001. "Genetic Analysis of Endocytosis in Caenorhabditis Elegans: Coelomocyte Uptake Defective Mutants." *Genetics (Austin)* 159(1):133–45. doi: 10.1093/genetics/159.1.133.

Ferri, Domenico, Amelia Mazzone, Giuseppa Esterina Liquori, Grazia Cassano, Maria Svelto, and Giuseppe Calamita. 2003. "Ontogeny, Distribution, and Possible Functional Implications of an Unusual Aquaporin, AQP8, in Mouse Liver." *Hepatology (Baltimore, Md.)* 38(4):947–57. doi: 10.1053/jhep.2003.50397.

Fridovich, Irwin. 1995. "Superoxide Radical and Superoxide Dismutases." *Annual Review of Biochemistry* 64(1):97–112. doi: 10.1146/annurev.bi.64.070195.000525.

Fridovich, Irwin. 1997. "Superoxide Anion Radical (O·2), Superoxide Dismutases, and Related Matters." *The Journal of Biological Chemistry* 272(30):18515–17. doi: 10.1074/jbc.272.30.18515.

Fukui, Masayuki, and Bao Ting Zhu. 2010. "Mitochondrial Superoxide Dismutase SOD2, but Not Cytosolic SOD1, Plays a Critical Role in Protection against Glutamate-Induced Oxidative Stress and Cell Death in HT22 Neuronal Cells." *Free Radical Biology and Medicine* 48(6):821–30. doi: 10.1016/j.freeradbiomed.2009.12.024.

Furness, John B., Brid P. Callaghan, Leni R. Rivera, and Hyun-Jung Cho. 2014. "The Enteric Nervous System and Gastrointestinal Innervation: Integrated Local and Central Control." *Advances in Experimental Medicine and Biology* 817:39–71. doi: 10.1007/978-1-4939-0897-4_3.

Gao, Jing, Makiko Hirata, Akiko Mizokami, Jin Zhao, Ichiro Takahashi, Hiroshi Takeuchi, and Masato Hirata. 2016. "Differential Role of SNAP-25 Phosphorylation by Protein Kinases A and C in the Regulation of SNARE Complex Formation and Exocytosis in PC12 Cells." *Cellular Signalling* 28(5):425–37. doi: 10.1016/j.cellsig.2015.12.014.

García Santamarina, Sarela, Susanna Boronat i Llop, and Elena Hidalgo Hernando. 2014. "Reversible Cysteine Oxidation in Hydrogen Peroxide Sensing and Signal Transduction." doi: 10.1021/bi401700f.

Gibbs, James, Robert C. Young, and Gerade P. Smith. 1973. "Cholecystokinin Elicits Satiety in Rats with Open Gastric Fistulas." *Nature (London)* 245(5424):323–25. doi: 10.1038/245323a0.

Giglio, A. M., T. Hunter, J. V. Bannister, W. H. Bannister, and G. J. Hunter. 1994. "The Copper/Zinc Superoxide Dismutase Gene of Caenorhabditis Elegans." *Biochemistry and Molecular Biology International* 33(1):41–44.

Giniatullin, Arthur, Alexey Petrov, and Rashid Giniatullin. 2019. "Action of Hydrogen Peroxide on Synaptic Transmission at the Mouse Neuromuscular Junction." *Neuroscience* 399:135–45. doi: 10.1016/j.neuroscience.2018.12.027.

Goda, Yukiko. 1997. "SNAREs and Regulated Vesicle Exocytosis." *Proceedings of the National Academy of Sciences - PNAS* 94(3):769–72. doi: 10.1073/pnas.94.3.769.

Grenham, Sue, Gerard Clarke, John F. Cryan, and Timothy G. Dinan. 2011. "Brain-Gut-Microbe Communication in Health and Disease." *Frontiers in Physiology* 2:94–94. doi: 10.3389/fphys.2011.00094.

Haber, Adam L., Moshe Biton, Noga Rogel, Rebecca H. Herbst, Karthik Shekhar, Christopher Smillie, Grace Burgin, Toni M. Delorey, Michael R. Howitt, Yarden Katz, Itay Tirosh, Semir Beyaz, Danielle Dionne, Mei Zhang, Raktima Raychowdhury, Wendy S. Garrett, Orit Rozenblatt-Rosen, Hai Ning Shi, Omer Yilmaz, Ramnik J. Xavier, and Aviv Regev. 2017. "A Single-Cell Survey of the Small Intestinal Epithelium." *Nature (London)* 551(7680):333–39. doi: 10.1038/nature24489.

Han, Derick, Fernando Antunes, Raffaella Canali, Daniel Rettori, and Enrique Cadenas. 2003. "Voltage-Dependent Anion Channels Control the Release of the Superoxide Anion from Mitochondria to Cytosol." *The Journal of Biological Chemistry* 278(8):5557–63. doi: 10.1074/jbc.M210269200.

Han, Wenfei, Luis A. Tellez, Matthew H. Perkins, Isaac O. Perez, Taoran Qu, Jozelia Ferreira, Tatiana L. Ferreira, Daniele Quinn, Zhong-Wu Liu, Xiao-Bing Gao, Melanie M. Kaelberer, Diego V. Bohórquez, Sara J. Shammah-Lagnado, Guillaume de Lartigue, and Ivan E. de Araujo. 2018. "A Neural Circuit for Gut-Induced Reward." *Cell* 175(3):665-678.e23. doi: 10.1016/j.cell.2018.08.049.

Hourihan, John M., Lorenza E. Moronetti Mazzeo, L. Paulette Fernández-Cárdenas, and T. Keith Blackwell. 2016. "Cysteine Sulfenylation Directs IRE-1 to Activate the SKN-1/Nrf2 Antioxidant Response." *Molecular Cell* 63(4):553–66. doi: 10.1016/j.molcel.2016.07.019.

Islas-Trejo, A., M. Land, I. Tcherepanova, J. H. Freedman, and C. S. Rubin. 1997. "Structure and Expression of the Caenorhabditis Elegans Protein Kinase C2 Gene. Origins and Regulated Expression of a Family of Ca2+-Activated Protein Kinase C Isoforms." *The Journal of Biological Chemistry* 272(10):6629–40. doi: 10.1074/jbc.272.10.6629.

Iwasaki, Kouichi, Jane Staunton, Owais Saifee, Michael Nonet, and James H. Thomas. 1997. "Aex-3 Encodes a Novel Regulator of Presynaptic Activity in C. Elegans." *Neuron (Cambridge, Mass.)* 18(4):613–22. doi: 10.1016/S0896-6273(00)80302-5.

Jahn, Reinhard, and Richard H. Scheller. 2006. "SNAREs - Engines for Membrane Fusion." *Nature Reviews. Molecular Cell Biology* 7(9):631–43. doi: 10.1038/nrm2002.

Jia, Qi, and Derek Sieburth. 2021. "Mitochondrial Hydrogen Peroxide Positively Regulates Neuropeptide Secretion during Diet-Induced Activation of the Oxidative Stress Response." *Nature Communications* 12(1):2304–2304. doi: 10.1038/s41467-021-22561-x.

Jiménez-Hidalgo, María, Cyril Léopold Kurz, José Rafael Pedrajas, Francisco José Naranjo-Galindo, María González-Barrios, Juan Cabello, Alberto G. Sáez, Encarnación Lozano, Emma L. Button, Elizabeth A. Veal, Juan Carlos Fierro-González, Peter Swoboda, and Antonio Miranda-Vizuete. 2014. "Functional Characterization of Thioredoxin 3 (TRX-3), a Caenorhabditis Elegans Intestine-Specific Thioredoxin." *Free Radical Biology & Medicine* 68(100):205–19. doi: 10.1016/j.freeradbiomed.2013.11.023.

Johnson, Joanne E., Jennifer Giorgione, and Alexandra C. Newton. 2000. "The C1 and C2 Domains of Protein Kinase C Are Independent Membrane Targeting Modules, with Specificity for Phosphatidylserine Conferred by the C1 Domain." *Biochemistry (Easton)* 39(37):11360–69. doi: 10.1021/bi000902c.

Kageyama, Natsumi, Masayo Nose, Masahiro Ono, Yohei Matsunaga, Takashi Iwasaki, and Tsuyoshi Kawano. 2022. "The FMRFamide-like Peptide FLP-2 Is Involved in the Modulation of Larval Development and Adult Lifespan by Regulating the Secretion of the Insulin-like Peptide INS-35 in Caenorhabditis Elegans." *Bioscience, Biotechnology, and Biochemistry* 86(9):1231–39. doi: 10.1093/bbb/zbac108.

Kamata, Hideaki, Tomoyuki Manabe, Shin-ichi Oka, Keiko Kamata, and Hajime Hirata. 2002. "Hydrogen Peroxide Activates IκB Kinases through Phosphorylation of Serine Residues in the Activation Loops." *FEBS Letters* 519(1):231–37. doi: 10.1016/S0014-5793(02)02712-6.

Kamath, R. 2003. "Genome-Wide RNAi Screening in Caenorhabditis Elegans." *Methods (San Diego, Calif.)* 30(4):313–21. doi: 10.1016/s1046-2023(03)00050-1.

Kemble, David J., and Gongqin Sun. 2009. "Direct and Specific Inactivation of Protein Tyrosine Kinases in the Src and FGFR Families by Reversible Cysteine Oxidation." *Proceedings of the National Academy of Sciences - PNAS* 106(13):5070–75. doi: 10.1073/pnas.0806117106.

Kim, Ki Mo, Deok Rim Heo, Young-A. Kim, Jun Lee, No Soo Kim, and Ok-Sun Bang. 2016. "Coniferaldehyde Inhibits LPS-Induced Apoptosis through the PKC α/β II/Nrf-2/HO-1 Dependent Pathway in RAW264.7 Macrophage Cells." *Environmental Toxicology and Pharmacology* 48:85–93. doi: 10.1016/j.etap.2016.10.016.

Knoefler, Daniela, Maike Thamsen, Martin Koniczek, Nicholas J. Niemuth, Ann-Kristin Diederich, and Ursula Jakob. 2012. "Quantitative In Vivo Redox Sensors Uncover Oxidative Stress as an Early Event in Life." *Molecular Cell* 47(5):767–76. doi: 10.1016/j.molcel.2012.06.016.

Konishi, Hiroaki, Motonari Tanaka, Yukitoshi Takemura, Hidenori Matsuzaki, Yoshitaka Ono, Ushio Kikkawa, and Yasutomi Nishizuka. 1997. "Activation of Protein Kinase C by Tyrosine Phosphorylation in Response to H2O2." *Proceedings of the National Academy of Sciences - PNAS* 94(21):11233–37. doi: 10.1073/pnas.94.21.11233.

Konishi, Hiroaki, Emiko Yamauchi, Hisaaki Taniguchi, Toshiyoshi Yamamoto, Hidenori Matsuzaki, Yukitoshi Takemura, Kyoko Ohmae, Ushio Kikkawa, and Yasutomi Nishizuka. 2001. "Phosphorylation Sites of Protein Kinase C δ in H2O2-Treated Cells and Its Activation by Tyrosine Kinase in Vitro." *Proceedings of the National Academy of Sciences - PNAS* 98(12):6587–92. doi: 10.1073/pnas.111158798.

Kontos, Hermes A., Enoch P. Wei, Earl F. Ellis, Larry W. Jenkins, John T. Povlishock, G. Thomas Rowe, and Michael L. Hess. 1985. "Appearance of Superoxide Anion Radical in Cerebral Extracellular Space during Increased Prostaglandin Synthesis in Cats." *Circulation Research* 57(1):142–51. doi: 10.1161/01.RES.57.1.142.

Lai, N. Y., K. Mills, and I. M. Chiu. 2017. "Sensory Neuron Regulation of Gastrointestinal Inflammation and Bacterial Host Defence." *Journal of Internal Medicine* 282(1):5–23. doi: 10.1111/joim.12591.

Land, Marianne, and Charles S. Rubin. 2017. "A Calcium- and Diacylglycerol-Stimulated Protein Kinase C (PKC), Caenorhabditis Elegans PKC-2, Links Thermal Signals to Learned Behavior by Acting in Sensory Neurons and Intestinal Cells." *Molecular and Cellular Biology* 37(19). doi: 10.1128/MCB.00192-17.

Larsen, Martha J., Elizabeth Ruiz Lancheros, Tracey Williams, David E. Lowery, Timothy G. Geary, and Teresa M. Kubiak. 2013. "Functional Expression and Characterization of the C. Elegans G-Protein-Coupled FLP-2 Receptor (T19F4.1) in Mammalian Cells and Yeast." *International Journal for Parasitology -- Drugs and Drug Resistance* 3:1–7. doi: 10.1016/j.ijpddr.2012.10.002.

Lee, Kiho, and Eleftherios Mylonakis. 2017. "An Intestine-Derived Neuropeptide Controls Avoidance Behavior in Caenorhabditis Elegans." *Cell Reports* 20(10):2501–12. doi: 10.1016/j.celrep.2017.08.053.

Lee, Yoon-Jin, Hae-Nyun Cho, Jae-Won Soh, Gil Ja Jhon, Chul-Koo Cho, Hee-Yong Chung, Sangwoo Bae, Su-Jae Lee, and Yun-Sil Lee. 2003. "Oxidative Stress-Induced Apoptosis Is Mediated by ERK1/2 Phosphorylation." *Experimental Cell Research* 291(1):251–66. doi: 10.1016/S0014-4827(03)00391-4.

Li, Guang, Jianke Gong, Haoyun Lei, Jianfeng Liu, and X. Z. Shawn Xu. 2016. "Promotion of Behavior and Neuronal Function by Reactive Oxygen Species in C. Elegans." *Nature Communications* 7(1):13234. doi: 10.1038/ncomms13234.

Li, Zongwei, Yong Gu, Ruoxue Wen, Fanxia Shen, Heng-Li Tian, Guo-Yuan Yang, and Zhijun Zhang. 2019. "Lysosome Exocytosis Is Involved in Astrocyte ATP Release after Oxidative Stress Induced by H2O2." *Neuroscience Letters* 705:251–58. doi: 10.1016/j.neulet.2019.03.046.

Lin-Moore, Alexander T., Motunrayo J. Oyeyemi, and Marc Hammarlund. 2021. "Rab-27 Acts in an Intestinal Pathway to Inhibit Axon Regeneration in C. Elegans." *PLoS Genetics* 17(11):e1009877–e1009877. doi: 10.1371/journal.pgen.1009877.

Liu, D. W., and J. H. Thomas. 1994. "Regulation of a Periodic Motor Program in C. Elegans." *The Journal of Neuroscience* 14(4):1953–62. doi: 10.1523/jneurosci.14-04-01953.1994.

Liu, Junqiang, Pei Zhang, Zhongfan Zheng, Muhammad Irfan Afridi, Shan Zhang, Zhiqing Wan, Xiumei Zhang, Lukas Stingelin, Yirong Wang, and Haijun Tu. 2023. "GABAergic Signaling between Enteric Neurons and Intestinal Smooth Muscle Promotes Innate Immunity and Gut Defense in Caenorhabditis Elegans." *Immunity* 56(7):1515-1532.e9. doi: 10.1016/j.immuni.2023.06.004.

Lutz, T. A., E. Del Prete, and E. Scharrer. 1994. "Reduction of Food Intake in Rats by Intraperitoneal Injection of Low Doses of Amylin." *Physiology & Behavior* 55(5):891–95. doi: 10.1016/0031-9384(94)90076-0.

Lutz, T. A., N. Geary, M. M. Szabady, E. Del Prete, and E. Scharrer. 1995. "Amylin Decreases Meal Size in Rats." *Physiology & Behavior* 58(6):1197–1202. doi: 10.1016/0031-9384(95)02067-5.

Mahoney, Timothy R., Qiang Liu, Takashi Itoh, Shuo Luo, Gayla Hadwiger, Rose Vincent, Zhao-Wen Wang, Mitsunori Fukuda, and Michael L. Nonet. 2006. "Regulation of Synaptic Transmission by RAB-3 and RAB-27 in Caenorhabditis Elegans." *Molecular Biology of the Cell* 17(6):2617–25. doi: 10.1091/mbc.E05-12-1170.

Mahoney, Timothy R., Shuo Luo, Elaine K. Round, Martin Brauner, Alexander Gottschalk, James H. Thomas, and Michael L. Nonet. 2008. "Intestinal Signaling to GABAergic Neurons Regulates a Rhythmic Behavior in Caenorhabditis Elegans." *Proceedings of the National Academy of Sciences - PNAS* 105(42):16350–55. doi: 10.1073/pnas.0803617105.

Maman, Moria, Filipa Carvalhal Marques, Yuli Volovik, Tatyana Dubnikov, Michal Bejerano-Sagie, and Ehud Cohen. 2013. "A Neuronal GPCR Is Critical for the Induction of the Heat Shock Response in the Nematode C. Elegans." *The Journal of Neuroscience* 33(14):6102–11. doi: 10.1523/JNEUROSCI.4023-12.2013.

Marinho, H. Susana, Carla Real, Luísa Cyrne, Helena Soares, and Fernando Antunes. 2014. "Hydrogen Peroxide Sensing, Signaling and Regulation of Transcription Factors." *Redox Biology* 2(C):535–62. doi: 10.1016/j.redox.2014.02.006.

Mayer, Emeran A., Karina Nance, and Shelley Chen. 2022. "The Gut-Brain Axis." *Annual Review of Medicine* 73(1):439–53. doi: 10.1146/annurev-med-042320-014032.

Mehrian-Shai, Ruty, Juergen K. V. Reichardt, Curtis C. Harris, and Amos Toren. 2019. "The Gut-Brain Axis, Paving the Way to Brain Cancer." *Trends in Cancer* 5(4):200–207. doi: 10.1016/j.trecan.2019.02.008.

Mello, C. C., J. M. Kramer, D. Stinchcomb, and V. Ambros. 1991. "Efficient Gene Transfer in C.Elegans: Extrachromosomal Maintenance and Integration of Transforming Sequences." *The EMBO Journal* 10(12):3959–70. doi: 10.1002/j.1460-2075.1991.tb04966.x.

Mertens, Inge, Tom Meeusen, Tom Janssen, Ronald Nachman, and Liliane Schoofs. 2005. "Molecular Characterization of Two G Protein-Coupled Receptor Splice Variants as FLP2 Receptors in Caenorhabditis

Elegans." *Biochemical and Biophysical Research Communications* 330(3):967–74. doi:
10.1016/j.bbrc.2005.03.071.

Messner, Kevin R., and James A. Imlay. 2002. "Mechanism of Superoxide and Hydrogen Peroxide
Formation by Fumarate Reductase, Succinate Dehydrogenase, and Aspartate Oxidase." *The Journal of*
Biological Chemistry 277(45):42563–71. doi: 10.1074/jbc.M204958200.

Miller, Brittany M., Megan J. Liou, Lillian F. Zhang, Henry Nguyen, Yael Litvak, Eva-Magdalena Schorr,
Kyung Ku Jang, Connor R. Tiffany, Brian P. Butler, and Andreas J. Bäumler. 2020. "Anaerobic Respiration
of NOX1-Derived Hydrogen Peroxide Licenses Bacterial Growth at the Colonic Surface." *Cell Host &*
Microbe 28(6):789-797.e5. doi: 10.1016/j.chom.2020.10.009.

Min, Do Sik, Eung-Gook Kim, and John H. Exton. 1998. "Involvement of Tyrosine Phosphorylation and
Protein Kinase C in the Activation of Phospholipase D by H2O2 in Swiss 3T3 Fibroblasts." *The Journal of*
Biological Chemistry 273(45):29986–94. doi: 10.1074/jbc.273.45.29986.

Miranda-Vizuete, Antonio, Anastasios E. Damdimopoulos, and Giannis Spyrou. 2000. "The Mitochondrial
Thioredoxin System." *Antioxidants & Redox Signaling* 2(4):801–10. doi: 10.1089/ars.2000.2.4-801.

Murphy, Coleen T., Seung-Jae Lee, and Cynthia Kenyon. 2007. "Tissue Entrainment by Feedback
Regulation of Insulin Gene Expression in the Endoderm of Caenorhabditis Elegans." *Proceedings of the*
National Academy of Sciences - PNAS 104(48):19046–50. doi: 10.1073/pnas.0709613104.

Nagy, Gabor, Ulf Matti, Ralf B. Nehring, Thomas Binz, Jens Rettig, Erwin Neher, and Jakob B. Sorensen.
2002. "Protein Kinase C-Dependent Phosphorylation of Synaptosome-Associated Protein of 25 kDa at
Ser187 Potentiates Vesicle Recruitment." *The Journal of Neuroscience* 22(21):9278–86. doi:
10.1523/JNEUROSCI.22-21-09278.2002.

Nebigil, Canan G. 1997. "Suppression of Phospholipase C β, γ, and δ Families Alters Cell Growth and
Phosphatidylinositol 4,5-Bisphosphate Levels." *Biochemistry (Easton)* 36(50):15949–58. doi:
10.1021/bi971721m.

Nelson, Kimberly J., Jesalyn A. Bolduc, Hanzhi Wu, John A. Collins, Elizabeth A. Burke, Julie A. Reisz,
Chananat Klomsiri, Scott T. Wood, Raghunatha R. Yammani, Leslie B. Poole, Cristina M. Furdui, and
Richard F. Loeser. 2018. "H2O2 Oxidation of Cysteine Residues in C-Jun N-Terminal Kinase 2 (JNK2)
Contributes to Redox Regulation in Human Articular Chondrocytes." *The Journal of Biological Chemistry*
293(42):16376–89. doi: 10.1074/jbc.RA118.004613.

Netto, Luis E. S., and Fernando Antunes. 2016. "The Roles of Peroxiredoxin and Thioredoxin in Hydrogen
Peroxide Sensing and in Signal Transduction." *Molecules and Cells* 39(1):65–71. doi:
10.14348/molcells.2016.2349.

Nonet, Michael L., Jane E. Staunton, Michael P. Kilgard, Tim Fergestad, Erika Hartwieg, H. Robert Horvitz,
Erik M. Jorgensen, and Barbara J. Meyer. 1997. "Caenorhabditis Elegans Rab-3 Mutant Synapses Exhibit
Impaired Function and Are Partially Depleted of Vesicles." *The Journal of Neuroscience* 17(21):8061–73.
doi: 10.1523/jneurosci.17-21-08061.1997.

O'Donnell, Michael P., Pin-Hao Chao, Jan E. Kammenga, and Piali Sengupta. 2018. "Rictor/TORC2
Mediates Gut-to-Brain Signaling in the Regulation of Phenotypic Plasticity in C. Elegans." *PLoS Genetics*
14(2):e1007213–e1007213. doi: 10.1371/journal.pgen.1007213.

Oláhová, Monika, Sarah R. Taylor, Siavash Khazaipoul, Jinling Wang, Brian A. Morgan, Kunihiro Matsumoto,
T. Keith Blackwell, and Elizabeth A. Veal. 2008. "Redox-Sensitive Peroxiredoxin That Is Important for
Longevity Has Tissue- and Stress-Specific Roles in Stress Resistance." *Proceedings of the National*
Academy of Sciences - PNAS 105(50):19839–44. doi: 10.1073/pnas.0805507105.

Oláhová, Monika, and Elizabeth A. Veal. 2015. "A Peroxiredoxin, PRDX‐2, Is Required for Insulin Secretion
and Insulin/IIS‐dependent Regulation of Stress Resistance and Longevity." *Aging Cell* 14(4):558–68. doi:
10.1111/acel.12321.

Oliveira-Marques, Virgínia, H. Susana Marinho, Luísa Cyrne, and Fernando Antunes. 2009. "Role of
Hydrogen Peroxide in NF-κB Activation: From Inducer to Modulator." *Antioxidants & Redox Signaling*
11(9):2223–43. doi: 10.1089/ars.2009.2601.

Ono, Yoshitaka, Tomoko Fujii, Koichi Igarashi, Takayoshi Kuno, Chikako Tanaka, Ushio Kikkawa, and
Yasutomi Nishizuka. 1989. "Phorbol Ester Binding to Protein Kinase C Requires a Cysteine-Rich Zinc-
Finger-Like Sequence." *Proceedings of the National Academy of Sciences - PNAS* 86(13):4868–71. doi:
10.1073/pnas.86.13.4868.

Ouaakki, Hafsa, Heetanshi Joshi, Laxmi Rathor, and Sung Min Han. 2023. "Frpr-18, a Neuropeptide
Receptor, Regulates Organismal Lifespan and Stress Tolerance in C. Elegans." *microPublication Biology*
2023. doi: 10.17912/micropub.biology.000840.
Pak, Valeriy V., Daria Ezeriņa, Olga G. Lyublinskaya, Brandán Pedre, Pyotr A. Tyurin-Kuzmin, Natalie M.
Mishina, Marion Thauvin, David Young, Khadija Wahni, Santiago Agustín Martínez Gache, Alexandra D.
Demidovich, Yulia G. Ermakova, Yulia D. Maslova, Arina G. Shokhina, Emrah Eroglu, Dmitry S. Bilan, Ivan
Bogeski, Thomas Michel, Sophie Vriz, Joris Messens, and Vsevolod V. Belousov. 2020. "Ultrasensitive
Genetically Encoded Indicator for Hydrogen Peroxide Identifies Roles for the Oxidant in Cell Migration and
Mitochondrial Function." *Cell Metabolism* 31(3):642-653.e6. doi: 10.1016/j.cmet.2020.02.003.
Pang, Shanshan, and Sean P. Curran. 2014. "Adaptive Capacity to Bacterial Diet Modulates Aging in
C. Elegans." *Cell Metabolism* 19(2):221–31. doi: 10.1016/j.cmet.2013.12.005.
Papa, Luena, Giovanni Manfredi, and Doris Germain. 2014. "SOD1, an Unexpected Novel Target for
Cancer Therapy." *Genes & Cancer* 5(1–2):15–21. doi: 10.18632/genesandcancer.4.
Paulsen, Michelle T., and Mats Ljungman. 2005. "The Natural Toxin Juglone Causes Degradation of P53
and Induces Rapid H2AX Phosphorylation and Cell Death in Human Fibroblasts." *Toxicology and Applied*
Pharmacology 209(1):1–9. doi: 10.1016/j.taap.2005.03.005.
Prahlad, Veena, Tyler Cornelius, and Richard I. Morimoto. 2008. "Regulation of the Cellular Heat Shock
Response in Caenorhabditis Elegans by Thermosensory Neurons." *Science (American Association for the*
Advancement of Science) 320(5877):811–14. doi: 10.1126/science.1156093.
Satoh, Yohsuke, Hirofumi Sato, Hirofumi Kunitomo, Xianfeng Fei, Koichi Hashimoto, and Yuichi Iino. 2014.
"Regulation of Experience-Dependent Bidirectional Chemotaxis by a Neural Circuit Switch in
Caenorhabditis Elegans." *The Journal of Neuroscience* 34(47):15631–37. doi: 10.1523/JNEUROSCI.1757-
14.2014.
Shakirzyanova, A. V., A. I. Malomouzh, N. V. Naumenko, and E. E. Nikolsky. 2009. "The Effect of Hydrogen
Peroxide on Spontaneous Quantal and Nonquantal Acetylcholine Release from Rat Motor Nerve Endings."
Doklady. Biological Sciences 424(1):18–20. doi: 10.1134/S0012496609010062.
Shao, Li-Wa, Rong Niu, and Ying Liu. 2016. "Neuropeptide Signals Cell Non-Autonomous Mitochondrial
Unfolded Protein Response." *Cell Research* 26(11):1182–96. doi: 10.1038/cr.2016.118.
Sharpe, Martyn A., Andrew D. Livingston, and David S. Baskin. 2012. "Thimerosal-Derived Ethylmercury Is
a Mitochondrial Toxin in Human Astrocytes: Possible Role of Fenton Chemistry in the Oxidation and
Breakage of mtDNA." *Journal of Toxicology* 2012:373678–373678. doi: 10.1155/2012/373678.
Shi, Yanjun, Lu Qin, Mengting Wu, Junyu Zheng, Tao Xie, and Zhiyong Shao. 2022. "Gut Neuroendocrine
Signaling Regulates Synaptic Assembly in C. Elegans." *EMBO Reports* 23(8):e53267-n/a. doi:
10.15252/embr.202153267.
Shu, Yilong, Xin Liu, Yan Yang, Masami Takahashi, and Kevin D. Gillis. 2008. "Phosphorylation of SNAP-
25 at Ser187 Mediates Enhancement of Exocytosis by a Phorbol Ester in INS-1 Cells." *The Journal of*
Neuroscience 28(1):21–30. doi: 10.1523/JNEUROSCI.2352-07.2008.
Sieburth, Derek, Jon M. Madison, and Joshua M. Kaplan. 2006. "PKC-1 Regulates Secretion of
Neuropeptides." *Nature Neuroscience* 10(1):49–57. doi: 10.1038/nn1810.
Singh, Jogender, and Alejandro Aballay. 2019. "Intestinal Infection Regulates Behavior and Learning via
Neuroendocrine Signaling." *eLife* 8. doi: 10.7554/eLife.50033.
Song, Youngmi, Jacqueline A. Koehler, Laurie L. Baggio, Alvin C. Powers, Darleen A. Sandoval, and Daniel
J. Drucker. 2019. "Gut-Proglucagon-Derived Peptides Are Essential for Regulating Glucose Homeostasis
in Mice." *Cell Metabolism* 30(5):976-986.e3. doi: 10.1016/j.cmet.2019.08.009.
Takada, Yasunari, Asok Mukhopadhyay, Gopal C. Kundu, Ganapati H. Mahabeleshwar, Sujay Singh, and
Bharat B. Aggarwal. 2003. "Hydrogen Peroxide Activates NF-Kappa B through Tyrosine Phosphorylation of
I Kappa B Alpha and Serine Phosphorylation of P65: Evidence for the Involvement of I Kappa B Alpha
Kinase and Syk Protein-Tyrosine Kinase." *The Journal of Biological Chemistry* 278(26):24233–41. doi:
10.1074/jbc.M212389200.
Tao, Jun, Qin-Yi Wu, Yi-Cheng Ma, Yuan-Li Chen, and Cheng-Gang Zou. 2017. "Antioxidant Response Is
a Protective Mechanism against Nutrient Deprivation in C. Elegans." *Scientific Reports* 7(1):43547–43547.
doi: 10.1038/srep43547.
Taylor, Seth R., Gabriel Santpere, Alexis Weinreb, Alec Barrett, Molly B. Reilly, Chuan Xu, Erdem Varol,
Panos Oikonomou, Lori Glenwinkel, Rebecca McWhirter, Abigail Poff, Manasa Basavaraju, Ibnul Rafi,
Eviatar Yemini, Steven J. Cook, Alexander Abrams, Berta Vidal, Cyril Cros, Saeed Tavazoie, Nenad Sestan,

Marc Hammarlund, Oliver Hobert, and David M. Miller. 2021. "Molecular Topography of an Entire Nervous
System." *Cell* 184(16):4329-4347.e23. doi: 10.1016/j.cell.2021.06.023.

Thacker, Colin, and Ann M. Rose. 2000. "A Look at the Caenorhabditis Elegans Kex2/Subtilisin-like
Proprotein Convertase Family." *BioEssays* 22(6):545–53. doi: 10.1002/(SICI)1521-
1878(200006)22:6<545::AID-BIES7>3.0.CO;2-F.

Thomas, J. H. 1990. "Genetic Analysis of Defecation in Caenorhabditis Elegans." *Genetics (Austin)*
124(4):855–72. doi: 10.1093/genetics/124.4.855.

Titos, Iris, Alen Juginović, Alexandra Vaccaro, Keishi Nambara, Pavel Gorelik, Ofer Mazor, and Dragana
Rogulja. 2023. "A Gut-Secreted Peptide Suppresses Arousability from Sleep." *Cell* 186(10):2273–74. doi:
10.1016/j.cell.2023.04.005.

Topham, Matthew K. 2006. "Signaling Roles of Diacylglycerol Kinases." *Journal of Cellular Biochemistry*
97(3):474–84. doi: 10.1002/jcb.20704.

Turner, Chris D., Carmen M. Ramos, and Sean P. Curran. 2024. "Disrupting the SKN-1 Homeostat:
Mechanistic Insights and Phenotypic Outcomes." *Frontiers in Aging* 5.

Van, An-Angela N., Maya T. Kunkel, Timothy R. Baffi, Gema Lordén, Corina E. Antal, Sourav Banerjee, and
Alexandra C. Newton. 2021. "Protein Kinase C Fusion Proteins Are Paradoxically Loss of Function in
Cancer." *The Journal of Biological Chemistry* 296:100445–100445. doi: 10.1016/j.jbc.2021.100445.

Van Raamsdonk, Jeremy Michael, and Siegfried Hekimi. 2010. "Reactive Oxygen Species and Aging in
Caenorhabditis Elegans: Causal or Casual Relationship?" *Antioxidants & Redox Signaling* 13(12):1911–53.
doi: 10.1089/ars.2010.3215.

Vincent, Andrea M., James W. Russell, Kelli A. Sullivan, Carey Backus, John M. Hayes, Lisa L. McLean,
and Eva L. Feldman. 2007. "SOD2 Protects Neurons from Injury in Cell Culture and Animal Models of
Diabetic Neuropathy." *Experimental Neurology* 208(2):216–27. doi: 10.1016/j.expneurol.2007.07.017.

Vitali, Roberta, Clara Prioreschi, Laura Lorenzo Rebenaque, Eleonora Colantoni, Daniela Giovannini,
Sarah Frusciante, Gianfranco Diretto, Francisco Marco-Jiménez, Mariateresa Mancuso, Arianna Casciati,
and Simonetta Pazzaglia. 2022. "Gut–Brain Axis: Insights from Hippocampal Neurogenesis and Brain
Tumor Development in a Mouse Model of Experimental Colitis Induced by Dextran Sodium Sulfate."
International Journal of Molecular Sciences 23(19):11495-. doi: 10.3390/ijms231911495.

Wang, Han, Kelly Girskis, Tom Janssen, Jason P. Chan, Krishnakali Dasgupta, James A. Knowles, Liliane
Schoofs, and Derek Sieburth. 2013. "Neuropeptide Secreted from a Pacemaker Activates Neurons to
Control a Rhythmic Behavior." *Current Biology* 23(9):746–54. doi: 10.1016/j.cub.2013.03.049.

Wang, Xiaowen, Hong Zhang, Russell Sapio, Jun Yang, Justin Wong, Xin Zhang, Jessie Y. Guo, Sharon
Pine, Holly Van Remmen, Hong Li, Eileen White, Chen Liu, Megerditch Kiledjian, Dimitri G. Pestov, and X.
F. Steven Zheng. 2021. "SOD1 Regulates Ribosome Biogenesis in KRAS Mutant Non-Small Cell Lung
Cancer." *Nature Communications* 12(1):2259–2259. doi: 10.1038/s41467-021-22480-x.

West, D. B., D. Fey, and S. C. Woods. 1984. "Cholecystokinin Persistently Suppresses Meal Size but Not
Food Intake in Free-Feeding Rats." *American Journal of Physiology. Regulatory, Integrative and
Comparative Physiology* 246(5):776-R787. doi: 10.1152/ajpregu.1984.246.5.r776.

Yanase, Yuhki, Izumi Hide, Shoji Mihara, Yasuhito Shirai, Naoaki Saito, Yoshihiro Nakata, Michihiro Hide,
and Norio Sakai. 2011. "A Critical Role of Conventional Protein Kinase C in Morphological Changes of
Rodent Mast Cells." *Immunology and Cell Biology* 89(1):149–59. doi: 10.1038/icb.2010.67.

Yang, Yan, Tim J. Craig, Xiaohui Chen, Leonora F. Ciufo, Masami Takahashi, Alan Morgan, and Kevin D.
Gillis. 2007. "Phosphomimetic Mutation of Ser-187 of SNAP-25 Increases Both Syntaxin Binding and Highly
Ca2+-Sensitive Exocytosis." *The Journal of General Physiology* 129(3):233–44. doi:
10.1085/jgp.200609685.

Yu, Hong Bing, Hyungjun Yang, Joannie M. Allaire, Caixia Ma, Franziska A. Graef, Arthur Mortha, Qiaochu
Liang, Else S. Bosman, Gregor S. Reid, James A. Waschek, Lisa C. Osborne, Harry Sokol, Bruce A.
Vallance, and Kevan Jacobson. 2021. "Vasoactive Intestinal Peptide Promotes Host Defense against
Enteric Pathogens by Modulating the Recruitment of Group 3 Innate Lymphoid Cells." *Proceedings of the
National Academy of Sciences - PNAS* 118(41):1-. doi: 10.1073/pnas.2106634118.

Zelko, Igor N., Thomas J. Mariani, and Rodney J. Folz. 2002. "Superoxide Dismutase Multigene Family: A
Comparison of the CuZn-SOD (SOD1), Mn-SOD (SOD2), and EC-SOD (SOD3) Gene Structures, Evolution,
and Expression." *Free Radical Biology & Medicine* 33(3):337–49. doi: 10.1016/S0891-5849(02)00905-X.

Zhang, Fujian, Anna-Lena Ström, Kei Fukada, Sangmook Lee, Lawrence J. Hayward, and Haining Zhu.
2007. "Interaction between Familial Amyotrophic Lateral Sclerosis (ALS)-Linked SOD1 Mutants and the
Dynein Complex." *The Journal of Biological Chemistry* 282(22):16691–99. doi: 10.1074/jbc.M609743200.

Figures

1203 Figure 1. Peptidergic gut-to-neuron FLP-2 signaling potentiates the oxidative stress response.
A (Top) Schematic showing the positions of AIY, intestine and coelomocytes of transgenic animals co-
expressing FLP-1::Venus in the intestine and mCherry in coelomocytes. Representative image of the
posterior coelomocyte that has taken up Venus into the endocytic compartment. Scale bar: 5µM. (Bottom)
Schematic showing FLP-1 and FLP-2 peptides as inter-tissue signals in gut-intestine regulation of the
antioxidant response.
B Representative images and quantification of average coelomocyte fluorescence of the indicated mutants
expressing FLP-1::Venus fusion proteins in AIY following M9 or 300µM juglone treatment for 10min.
Neuronal *aex-5* denotes expression of *aex-5* cDNA under the *rab-3* promoter; intestinal *aex-5* denotes

expression of *aex-5* cDNA under the *ges-1* promoter. Unlined *** denotes statistical significance compared
to "wild type". n = 30, 30, 24, 30, 26, 30, 30 independent animals. Scale bar: 5µM.
C Average percentage of surviving young adult animals of the indicated genotypes after 16h recovery
following 4h juglone treatment. Unlined ** denotes statistical significance compared to "wild type". n = 213,
156, 189, 195 independent biological samples over three independent experiments.
D Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-1::Venus
fusion proteins in AIY following M9 or 300µM juglone treatment for 10min. Neuronal *flp-2* denotes
expression of *flp-2* gDNA under the *rab-3* promoter; intestinal *flp-2* denotes expression of *flp-2* gDNA under
the *ges-1* promoter; intestinal *flp-2(OE)* denotes expression of *flp-2* gDNA under the *ges-1* promoter in wild
type animals. Unlined *** and ns denote statistical significance compared to "wild type". n = 20, 20, 25, 20,
20, 20, 25, 22 independent animals.
E Representative images and quantification of fluorescence of mitochondrial matrix-targeted HyPer7 in the
axon of AIY following M9 or 300µM juglone treatment for 10min. Arrowheads denote puncta marked by
mito::HyPer7 fusion proteins (Excitation: 500 and 400nm; emission: 520nm). Ratio of images taken with
500nM (GFP) and 400nM (CFP) for excitation was used to measure H_2O_2 levels. Unlined *** and ns denote
statistical significance compared to "wild type". n = 24, 22, 25, 24 independent animals. Scale bar: 10µM.
F Representative images and quantification of average fluorescence in the posterior intestine of transgenic
animals expressing P*gst-4::gfp* after 1h M9 or juglone exposure and 3h recovery. Asterisks mark the
intestinal region used for quantification. P*gst-4::gfp* expression in the body wall muscles, which appears as
fluorescence on the edge animals in some images, was not quantified. Unlined *** and ns denote statistical
significance compared to "wild type"; unlined ## and ### denote statistical significance compared to "wild
type+juglone". n = 25, 26, 25, 25, 25, 25, 25, 25 independent animals. Scale bar: 10µM.
G Representative images and quantification of average fluorescence in the posterior region of transgenic
animals expressing P*gst-4::gfp* after 1h M9 or juglone exposure and 3h recovery. Asterisks mark the
intestinal region for quantification. P*gst-4::gfp* expression in the body wall muscles, which appears as
fluorescence on the edge animals in some images, was not quantified. Unlined *** denotes statistical
significance compared to "wild type"; unlined ### denotes statistical significance compared to "wild
type+juglone". n = 23, 25, 25, 26, 24, 25 independent animals. Scale bar: 10µM.
B-G Data are mean values ± s.e.m normalized to wild type controls. ns. not significant, ** and ## $P < 0.01$,
*** and ### $P < 0.001$ by Brown-Forsythe and Welch ANOVA with Dunnett's T3 multiple comparisons test.

1242
Figure 2. FLP-2 secretion from the intestine is stress regulated.
A Schematic showing the positions of intestine and coelomocytes of transgenic animals co-expressing FLP-
2::Venus in the intestine and mCherry in coelomocytes. Representative images of the posterior
coelomocyte that have taken up Venus into the endocytic compartment (Scale bar: 5μM) and the posterior
intestinal region showing the distribution of FLP-2::Venus in puncta in the intestine are shown (Scale bar:
15μM).
B Representative images of fluorescence distribution in the posterior intestinal region of transgenic animals
co-expressing FLP-2::Venus and AEX-5::mTur2 fusion proteins. Arrowheads denote puncta marked by both
fusion proteins. Scale bar: 5μM.
C Schematic showing the locations of AEX-1/UNC13, AEX-3/MADD, AEX-4/SNAP25 and AEX-6/Rab27
relative to a DCV.
D Representative images and quantification of average coelomocyte fluorescence of the indicated mutants
expressing FLP-2::Venus fusion proteins in the intestine following M9 or 300μM juglone for 10min. Unlined
*** and ns denote statistical significance compared to "wild type". n = 29, 25, 24, 30, 23, 30, 25, 25, 25
independent animals. Scale bar: 5μM.
E Quantification of average coelomocyte fluorescence of transgenic animals expressing FLP-2::Venus
fusion proteins in the intestine following treatment with M9 buffer or the indicated stressors for 10min.
Unlined *** denotes statistical significane compared to "M9". n = 23, 25, 25 independent animals.
F Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-2::Venus
fusion proteins in the intestine following M9 or 300μM juglone treatment for 10min. Unlined ** denotes
statistical significance compared to "wild type"; unlined ## denotes statistical significance compared to "*flp-
1*"; a denotes statistical significance compared to "wild type+juglone". n = 30, 30, 30, 30 independent
animals.
D-F Data are mean values ± s.e.m normalized to wild type controls. ns. not significant, ** and ## $P < 0.01$,
*** and ### $P < 0.001$ by Brown-Forsythe and Welch ANOVA with Dunnett's T3 multiple comparisons test.

1268
1269 **Figure 3. SOD-1/SOD-3 mediates endogenous H₂O₂ regulates FLP-2 release from the intestine.**

A Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-2::Venus fusion proteins in the intestine following M9 or 300μM juglone treatment for 10min. Intestinal *sod*-1 denotes expression of *sod-1b* cDNA under the *ges-1* promoter. Unlined *** denotes statistical significance compared to "wild type". n = 25, 22, 24, 24, 25 independent animals.

B Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-2::Venus fusion proteins in the intestine following M9 or 300μM juglone treatment for 10min. Intestinal *sod-3* and *sod-3(ΔMLS)* denote intestinal expression of *sod-3* cDNA and *sod-3(ΔMLS)* variants, which lacks the mitochondrial localization sequence, under the *ges-1* promoter. Unlined *** denotes statistical significance compared to "wild type". n = 25, 25, 25, 25, 25, 25 independent animals.

C Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-2::Venus fusion proteins in the intestine following M9 or 300μM juglone treatment for 10min. Unlined *** denotes statistical significance compared to "wild type". n = 25, 25, 22, 25 independent animals.

D Representative images of fluorescence distribution in the posterior intestinal region of transgenic animals expressing SOD-1b::GFP fusion proteins in contrast against auto-fluorescence of gut granules. Scale bar: 10μM.

E Representative images of fluorescence distribution in the posterior intestinal region of transgenic animals co-expressing SOD-3::GFP and TOMM-20::mCherry (to target mitochondria) fusion proteins. Scale bar: 15μM.

F Representative images of fluorescence distribution in the posterior intestinal region of transgenic animals co-expressing SOD-3(ΔMLS)::GFP and TOMM-20::mCherry fusion proteins. Scale bar: 15μM.

G Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-2::Venus fusion proteins in the intestine following M9, 300μM juglone or 1mM H_2O_2 treatment for 10min. Unlined *** and ns denote statistical significance compared to "wild type". n = 29, 30, 25, 25, 25, 24, 25 independent animals.

H Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-2::Venus fusion proteins in the intestine following M9 or 1mM H_2O_2 treatment for 10min. n = independent animals.

I Schematic showing that SOD-1 and SOD-3 mediate juglone-induced H_2O_2 production in promoting FLP-2 release, and the PRDX-2/TRX-3 system detoxifies excessive H_2O_2.

J Schematic, representative images and quantification of fluorescence in the posterior region of the indicated transgenic animals co-expressing mitochondrial matrix targeted HyPer7 (matrix-HyPer7) or mitochondrial outer membrane targeted HyPer7 (OMM-HyPer7) with TOMM-20::mCherry following M9 or 300μM juglone treatment. Ratio of images taken with 500nM (GFP) and 400nM (CFP) for excitation and 520nm for emission was used to measure H_2O_2 levels. Unlined *** and ns denote statistical significance compared to "wild type". Unlined ## and ### denote statistical significance compared to "wild type+juglone". (top) n = 20, 20, 18, 20, 19, 19, 20, 20 independent animals. (bottom) n = 20, 20, 19, 20, 20, 20, 20, 20 independent animals. Scale bar: 5μM.

A-C, G-H and J Data are mean values ± s.e.m normalized to wild type controls. ns. not significant, * *P* < 0.05, ## *P* < 0.01, *** and ### *P* < 0.001 by Brown-Forsythe and Welch ANOVA with Dunnett's T3 multiple comparisons test.

1309
Figure 4. PRDX-2/PRDX and TRX-3/TRX regulate endogenous H_2O_2 and FLP-2 secretion.
A (Top) Schematic showing the PRDX/TRX system in H_2O_2 detoxification. (Bottom) Schematic showing the
three isoforms of *prdx-2* transcripts and *vj380* allele of *prdx-2b* knockout.
B Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-2::Venus
fusion proteins in the intestine following M9 or 300μM juglone treatment for 10min. Intestinal *prdx-2b*
denotes expression of *prdx-2b* cDNA under the *ges-1* promoter. Intestinal *trx-3* denotes expression of *trx-3*
cDNA under the *ges-1* promoter. Unlined *** denotes statistical significance compared to "wild type";
unlined ## and ### denote statistical significance compared to "*trx-3*". n= 25, 23, 25, 25, 25, 25, 25, 25, 25,
25, 25 independent animals.
C and D Quantification of fluorescence in the posterior region of the indicated transgenic animals co-
expressing matrix-HyPer7 (C) or OMM-HyPer7 (D) with TOMM-20::mCherry following M9 or 300μM juglone
treatment. Ratio of images taken with 500nM (GFP) and 400nM (CFP) for excitation and 520nm for
emission was used to measure H_2O_2 levels. Unlined *** and ns denote statistical significance compared to
"wild type. (C) n = 20, 20, 20, 20, 20 independent animals. (D) n = 20, 20, 20, 20, 20 independent animals.
E Quantification of average coelomocyte FLP-2::Venus fluorescence of transgenic animals fed with RNAi
bacteria targeting the indicated genes following M9 treatment for 10min. Unlined *** denotes statistical
significance compared to "empty vector". n = 25, 23, 24 independent animals.
F Representative images and quantification of average fluorescence in the posterior region of transgenic
animals expressing P*gst-4::gfp* after 1h M9 or juglone exposure and 3h recovery. Asterisks mark the
intestinal region for quantification. P*gst-4::gfp* expression in the body wall muscles, which appears as
fluorescence on the edge animals in some images, was not quantified. Unlined ** denotes statistical
significance compared to "wild type", unlined ## denotes statistical analysis compared to "*prdx-2b*". n = 25,
25, 25 independent animals. Scale bar: 10μM.
B-F Data are mean values ± s.e.m normalized to wild type controls. n.s. not significant, ** and ## $P < 0.01$,
*** and ### $P < 0.001$ by Brown-Forsythe and Welch ANOVA with Dunnett's T3 multiple comparisons test.

1335
Figure 5. PKC-2/PKCα/β activation by H₂O₂ promotes FLP-2 secretion from the intestine.
A Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-2::Venus
fusion proteins in the intestine following M9 or 300μM juglone treatment for 10min. Intestinal *pkc-2* denotes
expression of *pkc-2b cDNA* under the ges-1 promoter. Intestinal *pkc-2b(K375R)* denotes expression of *pkc-
2b(K375R)* variants under the *ges-1* promoter. Unlined *** and ns denote statistical significance compared
to "wild type"; ### denote statistical significance compared to "*pkc*-2+juglone". n = 24, 24, 25, 25, 25, 25
independent animals.
B and C Quantification of fluorescence in the posterior region of the indicated transgenic animals co-
expressing matrix-HyPer7 (B) or OMM-HyPer7 (C) with TOMM-20::mCherry following M9 or 300μM juglone
treatment. Ratio of images taken with 500nM (GFP) and 400nM (CFP) for excitation and 520nm for
emission was used to measure H₂O₂ levels. Unlined *** denotes statistical significance compared to "wild
type"; unlined ### denotes statistical analysis compared to "*pkc-2*". (B) n = 20, 20, 19, 20 independent
animals, (C) n = 20, 20, 20, 20 independent animals.
D Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-2::Venus
fusion proteins in the intestine following M9 or 1mM H₂O₂ treatment for 10min. n = 23, 25, 25 independent
animals.
E Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-2::Venus
fusion proteins in the intestine following M9 treatment for 10min. Unlined *** denotes statistical significance
compared to "wild type"; unlined ### denotes statistical significance compared to "*prdx-2*". n = 25, 25, 25,
25 independent animals.
A-E Data are mean values ± s.e.m normalized to wild type controls. ns. not significant, *** and ### $P <$
0.001 by Brown-Forsythe and Welch ANOVA with Dunnett's T3 multiple comparisons test.

1358
Figure 6. DAG promotes PKC-2 mediated FLP-2 secretion from the intestine.
A Schematic showing PLC and DGK mediates DAG metabolism and DAG functions in H_2O_2-mediated FLP-
2 signaling.
B Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-2::Venus
fusion proteins in the intestine following M9 or juglone treatment for 10min. n = 25, 25, 25, 25 independent
animals.
C and D Quantification of fluorescence in the posterior region of the indicated transgenic animals co-
expressing matrix-HyPer7 (C) or OMM-HyPer7 (D) with TOMM-20::mCherry following M9 or 300µM juglone
treatment. Ratio of images taken with 500nM (GFP) and 400nM (CFP) for excitation and 520nm for
emission was used to measure H_2O_2 levels. Unlined *** denotes statistical significance compared to "wild
type"; unlined ### denotes statistical significance compared to "*egl-8*". (C) n = 22, 20, 20, 21 independent
animals, (D) n = 20, 20, 20, 20 independent animals.
E Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-2::Venus
fusion proteins in the intestine following M9 or 300µM juglone treatment for 10min. Intestinal *dgk-2* denotes
expression of *dgk-2a* cDNA under the *ges-1* promoter. Unlined *** denotes statistical significance
compared to "wild type"; unlined ### denotes statistical significance compared to "*dgk-2*/DGKε". n = 25,
25, 25, 25, 24 independent animals.

F and G Quantification of fluorescence in the posterior region of the indicated transgenic animals co-
expressing matrix-HyPer7 (F) or OMM-HyPer7 (G) with TOMM-20::mCherry following M9 treatment. Ratio
of images taken with 500nM (GFP) and 400nM (CFP) for excitation and 520nm for emission was used to
measure H_2O_2 levels. (F) n = 20, 20 independent animals, (G) n = 20, 20 independent animals.
H Quantification of average coelomocyte fluorescence of the indicated transgenic animals fed with RNAi
bacteria targeting the indicated genes in the intestine following M9 treatment for 10min. n = 25, 24, 25, 30
independent animals.
I (Top) Schematic showing the position of intestine and AIY neurons in FLP-1-FLP-2 mediated axis. (Bottom)
Schematic showing endogenous H_2O_2 promotes PKC-2/AEX-4 mediated FLP-2 release from the intestine
in FLP-1-FLP-2 regulated inter-tissue axis.
B-H Data are mean values ± s.e.m normalized to wild type controls. **B-E and H** ns. not significant, ** $P <$
0.01, *** and ### $P < 0.001$ by Brown-Forsythe and Welch ANOVA with Dunnett's T3 multiple comparisons
test. **F and G** ns. not significant by unpaired t test with Welch's correction.

Supplementary Figures

1391
1392 Supplementary Figure 1. The effect of intestinal DCV secretion mutations on FLP-1 release from AIY.
A Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-1::Venus
fusion proteins in AIY following M9 or 300μM juglone treatment for 10min. Unlined *** and ### denotes
statistical significance compared to "wild type". n = 30, 30, 29, 30, 30, 30 independent animals.
B Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-1::Venus
fusion proteins in AIY following M9 or 300μM juglone treatment for 10min. Unlined *** and ns denote
statistical analysis compared to "wild type". n = 24, 24, 25, 25, 30, 30 independent animals.
C Average percentage of surviving young adult animals of the indicated genotypes after 16h recovery
following 4h DMSO treatment. n = 203, 174 independent biological samples over three independent
experiments.
D Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-1::Venus
fusion proteins in AIY following M9, DMSO or juglone treatment for 10min. Unlined ns and *** denote
statistical significance compared to "M9". n = 20, 20, 19 independent animals.
A-D Data are mean values ± s.e.m normalized to wild type controls. **A, B and D** ns. not significant, *** and
P < 0.001 by Brown-Forsythe and Welch ANOVA with Dunnett's T3 multiple comparisons test. **C** ns.
not significant by unpaired t test with Welch's correction.

1408
1409 **Supplementary Figure 2. Specificity of juglone on intestinal peptide secretion, and FLP-2 and NLP-**
1410 **40 localization in the intestine.**

1411
A Quantification of average coelomocyte fluorescence of the indicated mutants co-expressing FLP-
2::Venus in the intestine (under the *ges-1* promoter) and mCherry in the coelomocytes (under the *ofm-1*
promoter) following M9 or 300µM juglone treatment for 10min. n = 23, 19 independent animals.
B Quantification of average coelomocyte fluorescence of transgenic animals expressing NLP-40::Venus
fusion proteins in the intestine following M9 or 300µM juglone exposure for 10min. n = 25, 24 independent
animals.
C Quantification of average coelomocyte fluorescence of transgenic animals expressing NLP-27::Venus
fusion proteins in the intestine following M9 or 300µM juglone exposure for 10min. n = 23, 25 independent
animals.
D Representative images of fluorescence distribution in the posterior intestinal region of transgenic animals
co-expressing FLP-2::Venus fusion proteins (marked by arrowheads) and NLP-40::mTur2 fusion proteins
(marked by arrows). Scale bar: 5µM.
E Quantification of average coelomocyte fluorescence of transgenic animals expressing FLP-2::Venus
fusion proteins in the intestine following M9, DMSO or 300µM juglone exposure for 10min. Unlined ns and
*** denote statistical significance compared to "M9". n = 20, 20, 20 independent animals.
A-C and E Data are mean values ± s.e.m normalized to wild type controls. **A-C** ns. not significant by
unpaired t test with Welch's correction. **E** ns. not significant by Brown-Forsythe and Welch ANOVA with
Dunnett's T3 multiple comparisons test.

1430
Supplementary Figure 3. SODs function in juglone induced FLP-2 release from the intestine and
mitochondrial mCherry control.
A Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-2::Venus
fusion proteins in the intestine following M9 or juglone treatment for 10min. Unlined *** denotes statistical
significance compared to "wild type"; unlined ### and ns denote statistical significance compared to "wild
type + juglone" n = 29, 27, 29, 27, 25, 26, 24 independent animals.
B and C Representative images and quantification of average fluorescence intensity of TOMM-20::mCherry
proteins in transgenic animals co-expressing matrix-HyPer7 (B) or OMM-HyPer7 (C) following M9 or H_2O_2
treatment for 10min. (B) Scale bar: Scale bar: 5µM. n = 20, 20 independent animals. (C) Scale bar: Scale
bar: 5µM. n = 20, 22 independent animals.
A-C Data are mean values ± s.e.m normalized to wild type controls. **A** ns. not significant, *** and ### $P <$
0.001 by Brown-Forsythe and Welch ANOVA with Dunnett's T3 multiple comparisons test. **B and C** ns. not
significant by unpaired t test with Welch's correction.

1444
Supplementary Figure 4. PRDX-2 intestinal rescue and mediates SOD-3 dependent regulation of
FLP-2 release.
A Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-2::Venus
fusion proteins in the intestine following M9 treatment for 10min. n = 30, 29 independent animals.
B Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-2::Venus
fusion proteins in the intestine following M9 treatment for 10min. Intestinal *prdx-2a* denotes expression of
prdx-2a cDNA under the *ges-1* promoter. n = 30, 30, 25 independent animals.
C Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-2::Venus
fusion proteins in the intestine following M9 treatment for 10min. Intestinal *prdx-2c* denotes expression of
prdx-2c cDNA under the *ges-1* promoter. n = 25, 25, 25 independent animals.
D Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-2::Venus
fusion proteins in the intestine following M9 treatment for 10min. n = 25, 23, 22 independent animals.
A-D Data are mean values ± s.e.m normalized to wild type controls. **A** ns. not significant by unpaired t test
with Welch's correction. **B-D** ns. not significant, *** and ### *P* < 0.001 by Brown-Forsythe and Welch
ANOVA with Dunnett's T3 multiple comparisons test.

1460
Supplementary Figure 5. Juglone promotes FLP-2 release in *pkc-1* mutants and expulsion analysis.
A Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-2::Venus
fusion proteins in the intestine following M9 or juglone treatment for 10min. Unlined ns and ** denote
statistical significance compared to "wild type". n = 24, 25, 20, 25 independent animals.
B Quantification of the number of expulsions (Exp) per defecation cycle in adult animals of the indicated
genotypes. Unlined *** and ns denote statistical significance compared to "wild type". n = 30, 30, 30, 30 in
three independent animals.
A-B Data are mean values ± s.e.m normalized to wild type controls. ns. not significant, ** and ## P < 0.01,
*** P < 0.001 by Brown-Forsythe and Welch ANOVA with Dunnett's T3 multiple comparisons test.

intestinal FLP-2::Venus secretion

1470
Supplementary Figure 6. Juglone promotes FLP-2 release in *plc-2* mutants.
Quantification of average coelomocyte fluorescence of the indicated mutants expressing FLP-2::Venus
fusion proteins in the intestine following M9 or juglone treatment for 10min. Unlined *** denotes statistical
significance compared to "wild type"; unlined ### denotes statistical significance compared to "*plc-2*/PLCβ".
n = 25, 25, 23, 28 independent animals.
Data are mean values ± s.e.m normalized to wild type controls. ns. not significant, *** and ### *P* < 0.001
by Brown-Forsythe and Welch ANOVA with Dunnett's T3 multiple comparisons test.

Table S1. Strains, transgenic lines and plasmids used in this study

Strain	Genotype	Figures
N2	wild type Bristol strain	Fig. 1C, S1C, S5B
OJ6555	*flp-1(ok2811) IV*	Fig. 1C
OJ5490	*flp-2(ok3351) X*	Fig. 1C
OJ10228	*flp-1(ok2811);flp-2(ok3351)*	Fig. 1C
OJ7466	*aex-4(sa22) X*	Fig. S5B
VC127	*pkc-2(ok328) X*	Fig. S5B
OJ3614	*vjIs150[pJQ60(Pttx-3::flp-1::Venus)]*	Fig. 1A, 1B, 1D, S1B, S1D
OJ5616	*aex-5(sa23);vjIs150*	Fig. 1B
OJ5780	*vjEx1748[pJQ298(Prab-3::aex-5 cDNA)];aex-5(sa23);vjIs150*	Fig. 1B
OJ5785	*vjEx1753[pJQ299(Pges-1::aex-5 cDNA)];aex-5(sa23);vjIs150*	Fig. 1B
OJ6334	*vjEx1753;aex-5(sa23);flp-2(ok3351);vjIs150*	Fig. 1B
OJ5264	*flp-2(ok3351);vjIs150*	Fig. 1D
OJ8818	*vjEx2882[pJQ366(Prab-3::flp-2 gDNA)];flp-2(ok3351);vjIs150*	Fig. 1D
OJ8813	*vjEx2877[pJQ305(ges-1::flp-2 gDNA)];flp-2(ok3511);vjIs150*	Fig. 1D
OJ10229	*vjEx2877;vjIs150*	Fig. 1D
CL2166	*dvIs19[pAF15(Pgst-4::GFP::NLS)]*	Fig. 1F, 1G, 4F
OJ2547	*flp-1(ok2811);dvIs19*	Fig. 1F, 1G
OJ10230	*flp-2(ok3351);dvIs19*	Fig. 1F
OJ6544	*flp-1(ok2811);flp-2(ok3511);dvIs19*	Fig. 1F
OJ10231	*vjEx2877;dvIs19*	Fig. 1G
OJ10232	*vjEx2877;flp-1(ok2281);dvIs19*	Fig. 1G
OJ5888	*aex-1(sa9);vjIs150*	Fig. S1B
OJ5890	*aex-3(js815);vjIs150*	Fig. S1B
OJ5891	*aex-4(sa22);vIs150*	Fig. S1B
OJ5892	*aex-6(sa24);vjIs150*	Fig. S1B
OJ5615	*nlp-40(tm4085);vjIs150*	Fig. S1C
OJ5889	*aex-2(sa3);vjIs150*	Fig. S1C
OJ6405	*vjEx2035[pJQ305(Pges-1::flp-2 gDNA::Venus)]*	Fig. 2A, 2C, 2D, 2F, S2A, S2E, 3A, 3B, 3C, 3G, 3H, S3A, 4B, 4E, S4A, S4B, S4C, S4D, 5A, 5D, 5E, S5A, 6B, 6E, 6H, S6
OJ9469	*vjEx3069[pDY10(Pges-1::aex-5::mTur2)];vjEx2035*	Fig. 2B
OJ6409	*aex-4(sa22);vjEx2035*	Fig. 2C, 3G
OJ8345	*aex-6(sa24);vjEx2035*	Fig. 2C, 3G
OJ6641	*flp-1(ok2811);vjEx2035*	Fig. 2F
OJ1002	*vjIs40[pDS292(Pnlp-40::nlp-40::Venus)]*	Fig. S2B
OJ10237	*vjEx3263[pJQ370(Pges-1::nlp-27 gDNA::Venus)]*	Fig. S2C
OJ9567	*vjEx3062[pDY(Pges-1::nlp-40::mTur2)];vjEx2035*	Fig. S2D
OJ9797	*sod-1(tm783);vjEx2035*	Fig. 3A, 3H, S3A
OJ8588	*vjEx2814[pJQ419(Pges-1::sod-1b cDNA)];sod-1(tm783);vjEx2035*	Fig. 3A
OJ8341	*sod-3(tm760);vjEx0235*	Fig. 3B, 3H, S3A
OJ8933	*vjEx2910[pJQ389(Pges-1::sod-3 cDNA)];sod-3(tm760);vjEx2035*	Fig. 3B
OJ9106	*vjEx2973[pJQ408(Pges-1::sod-3(ΔMLS) cDNA)];sod-3(tm760);vjEx2035*	Fig. 3B
OJ10234	*sod-1(tm783);sod-3(tm760);vjEx2035*	Fig. 3C, 3H
OJ10243	*vjEx3266[pJQ420(Pges-1::sod-1b cDNA::GFP)]*	Fig. 3D
OJ9141	*vjEx2993[pJQ407(Pges-1::sod-3 cDNA::GFP)]*	Fig. 3E
OJ9144	*vjEx2996[pJQ409(Pges-1::sod-3(ΔMLS) cDNA::GFP)]*	Fig. 3F

OJ9230	vjEx3020[pJQ383(Pges-1::MLS::HyPer7)]	Fig. 3I, S3B, 5B, 6C, 6F
OJ9196	vjEx3014[pJQ411(Pges-1::tomm-20::HyPer7)]	Fig. 3I, S3C, 5C ,6D, 6G
OJ9281	sod-1(tm783);vjEx3020	Fig. 3I
OJ9259	sod-3(tm760);vjEx3020	Fig. 3I
OJ10244	sod-1(tm783);sod-3(tm760);vjEx3020	Fig. 3I
OJ9795	sod-1(tm783);vjEx3014	Fig. 3I
OJ9280	sod-3(tm760);vjEx3014	Fig. 3I
OJ10245	sod-1(tm783);sod-3(tm760);vjEx3014	Fig. 3I
OJ10238	sod-2(ok1030);vjEx2035	Fig. S3A
OJ10239	sod-4(gk101);vjEx2035	Fig. S3A
OJ10240	sod-5(tm1146);vjEx2035	Fig. S3A
OJ8991	prdx-2(gk169);vjEx2035	Fig. 4B, S3E, S3F ,4C, 5E
OJ10251	prdx-2b(vj380);vjEx2035	Fig. 4B, 4E, S4B
OJ8996	vjEx2926[pJQ381(Pges-1::prdx-2b cDNA)];prdx-2(gk169);vjEx2035	Fig. 4B
OJ9249	trx-3(tm2820);vjEx2035	Fig. 4B
OJ9496	vjEx3091[pJQ422(Pges-1::trx-3 cDNA)];trx-3(tm2820);vjEx2035	Fig. 4B
OJ10252	trx-3(tm2820);sod-1(tm783);vjEx2035	Fig. 4B
OJ10253	trx-3(tm2820);sod-3(tm760);vjEx2035	Fig. 4B
OJ9237	prdx-2(gk169);vjEx3020	Fig. 4C
OJ10247	prdx-2b(vj380);vjEx3020	Fig. 4C
OJ10249	trx-3(tm2820);vjEx3020	Fig. 4C
OJ10246	prdx-2(gk169);vjEx3014	Fig. 4D
OJ10248	prdx-2b(vj380);vjEx3014	Fig. 4D
OJ10250	trx-3(tm2820);vjEx3014	Fig. 4D
OJ10254	prdx-2b(vj380);dvIs19	Fig. 4F
OJ10255	prdx-2b(vj380);flp-2(tm3351);dvIs19	Fig. 4F
OJ10256	prdx-3(gk529);vjEx2035	Fig. S4A
OJ10258	vjEx3268[pJQ380(Pges-1::prdx-2a cDNA)];prdx-2(gk169);vjEx2035	Fig. S4B
OJ10260	vjEx3270[pJQ399(Pges-1::prdx-2c cDNA)];prdx-2(gk169);vjEx2035	Fig. S4C
OJ9250	prdx-2b(vj380);sod-3(tm760);vjEx2035	Fig. S4D
OJ9682	pkc-2(ok328);vjEx2035	Fig. 5A, 5D, 5E
OJ8682	vjEx2828[pJQ376(Pges-1::pkc-2b cDNA)];pkc-2(ok328);vjEx2035	Fig. 5A
OJ9657	vjEx3131[pJQ446(Pges-1::pkc-2(K375R) cDNA)];pkc-2(ok328);vjEx2035	Fig. 5A
OJ10279	pkc-2(ok328);vjEx3020	Fig. 5B
OJ10280	pkc-2(ok328);vjEx3014	Fig. 5B
OJ8939	prdx-2(gk169);pkc-2(ok328);vjEx2035	Fig. 5E
OJ10278	pkc-1(nj3);vjEx2035	Fig. S5A
OJ9863	egl-8(sa47);vjEx2035	Fig. 6B
OJ10281	egl-8(sa47);vjEx3020	Fig. 6C
OJ10282	egl-8(sa47);vjEx3014	Fig. 6D
OJ10263	dgk-2(gk124);vjEx2035	Fig. 6E, 6H
OJ10264	vjEx327[pJQ460(Pges-1::dgk-2a cDNA)];dgk-2(gk124);vjEx2035	Fig. 6E
OJ10266	dgk-2(gk124);pkc-2(ok328);vjEx2035	Fig. 6E
OJ10283	dgk-2(gk124);vjEx3020	Fig. 6F
OJ10284	dgk-2(gk124);vjEx3014	Fig. 6G
OJ9809	plc-2(ok1761);vjEx2035	Fig. S6
OJ9028	vjEx2936[pJQ382(Pttx-3::MLS::HyPer7)]	Fig. 1E

| OJ10595 | *vjEx2936;flp-2(ok3351)* | Fig. 1E |

1479