Lower baseline immunity in invasive Egyptian goose compared to sympatric native waterfowls

Simone Messina[1,2], Hanna Prüter[3], Gábor Árpád Czirják[4], David Costantini[1]

[1] Department of Ecological and Biological Sciences, University of Tuscia, Largo dell'Università s.n.c., 01100 Viterbo, Italy

[2] Behavioural Ecology & Ecophysiology Group, Department of Biology, University of Antwerp, Universiteitsplein 1, 2610 Wilrijk, Belgium

[3] Department of Ornithology, Max Planck Institute for Biological Intelligence, Eberhard-Gwinner-Strasse, 82319, Seewiesen, Germany

[4] Department of Wildlife Diseases, Leibniz Institute for Zoo and Wildlife Research, Berlin, Germany

Corresponding author: Simone Messina
Email: simone.messina@unitus.it
ORCID: https://orcid.org/0000-0001-6034-7450

Abstract

Successful invasive species out compete native species for vital resources. To increase their spreading success, invasive species need to trade-off nutritional and metabolic resources allocated to reproduction and range expansion with other costly body functions. One proposed mechanism for the reallocation of resources is a trade-off with the immune function. According to the evolution of increased competitive ability hypothesis, a reduced investment in immunity would be favoured by parasite loss in the colonised habitats. It is also suggested that invaders would reallocate resources among different immune effectors depending on the new pathogens they encounter in the new habitat. Reallocation of resources may also involve the regulation of oxidative status given its fundamental link with the immune system. Relying on a panel of blood-based markers of immune function and oxidative status quantified in an invasive species (Egyptian goose) and two native competing species (mallard and mute swan) in Germany, we tested the hypothesis that the invasive species would have a lower investment in immune function compared to the native species. We predicted lower levels of baseline immune markers associated with systemic inflammatory response, and higher levels of certain low-cost immunological effectors (e.g. humoral effectors) in the invasive species compared to the two native species. If Egyptian geese reduced their investment in immune function, we would also expect that geese generated less oxidative damage and had a lower expression of antioxidant defences than native species. We found lower levels of several immune markers associated either with inflammatory response or humoral innate and adaptive immunity in the invasive species compared to the two native species. Bacteria killing ability of plasma was the only immune marker that was upregulated in Egyptian geese. Markers of oxidative status were higher in mallards compared to the other species. The results of our study point to an overall reduced investment in immune function in the invasive species as a possible energy-saving immunological strategy due to the loss of parasites in the new colonised habitats. Thus, a lower investment in immune function may benefit other energy-demanding activities, such as reproduction, dispersal, and territoriality.

Introduction

The occurrence of biological invasions is increasing worldwide as a consequence of ongoing globalization. Successful invasive species engage in competition for vital resources with native species, sometimes even replacing native species in their natural habitat, and thus concurs with the ongoing biodiversity crisis (Mollot et al., 2017). Moreover, invaders can disrupt ecosystem function and facilitate the distribution of novel pathogens, including some with zoonotic potential, generating massive economic costs and challenges for public and animal health (Henry et al., 2023; Turbelin et al., 2023; Zhang et al., 2022).

Non-native introduced species need to establish and spread over the new region to become invasive. High population growth rate and superior dispersal ability are two main life-history traits that promote successful invasions (Sakai et al., 2001). However, increased need for nutritional and metabolic resources for particularly high reproductive investment and dispersion means that invaders need to trade-off those resources with other important body functions. Since invasive plants and animals may experience a loss of parasites during the invasion process (Dunn et al., 2012; Torchin et al., 2003), a reduced investment in the immune function might concur with the reallocation of resources (White and Perkins, 2012).

The immune system of vertebrates involves different pathways that require relatively higher or lower energetic costs. The evolution of increased competitive ability (EICA) hypothesis predicts that a loss of natural parasites in a habitat enables invasive species to colonise with reduced investment in defence functions and thus allocate more resources towards fitness-related traits (Blossey and Notzold, 1995). However, as an overall reduction of the immune response of invading hosts might increase their susceptibility towards pathogens specific to the novel habitat, it has been proposed that the reallocation of resources between different immune effectors would reduce the immune costs while keeping the system functioning at the same time (Lee and Klasing, 2004; White and Perkins, 2012).

The EICA hypothesis was refined by Lee and Klasing (2004), who presented an analysis focused on the trade-offs between immune effectors in invasive vertebrates. The revised EICA hypothesis proposed that successful invasive species minimize systemic inflammatory responses that are triggered by strong innate or cell-mediated immune response to infectious agents, as they come at elevated energetic and immunopathological costs. Rather, a successful strategy for invaders to fight novel pathogens might be the

increase of humoral defences, which includes both constitutive levels of natural antibodies and adaptive

antibody responses (Lee and Klasing, 2004).

Studies on invasive populations of different vertebrate species show mixed evidence for trade-offs

between immune effectors and host traits. For example, invasive cane toads (*Rhinella marina*) moving further

distances showed decreased bacteria-killing ability and phagocytic activity, but higher swelling response

compared to their more sedentary conspecifics, suggesting trade-offs between dispersal distance and

immunity (Brown and Shine, 2014). An immune challenge experiment found minor metabolic changes and

reduced movement in American invasive house sparrows (*Passer domesticus*) compared to less-invasive tree

sparrows (*Passer montanus*). Additionally, immune-challenged tree sparrows had lower reproductive success

compared to house sparrows indicating greater immune and fitness costs in the tree sparrows (Lee et al.,

2005). Recently, a study on Egyptian geese (*Alopochen aegyptiaca*) found higher baseline values of

haemolysis, heterophil granulocytes and lymphocytes in the German invasive population compared to a

native population (Prüter et al., 2020). Although we still have a poor understanding of how variation in

immunological traits affects the success of biological invasions, multi-trait comparative studies might provide

valuable information on the mechanisms involved.

Energy-based trade-off models provide valuable insight into the regulation of immune function in

invaders. There are, however, other relevant physiological costs that can underlie immune investment. One

important physiological mechanism linking immune response to individual fitness is the regulation of cellular

oxidative status (Costantini and Møller, 2009; Hasselquist and Nilsson, 2012). Oxidative status is determined

by levels of reactive oxygen species (ROS) generation, oxidative damage caused by ROS to biomolecules, and

antioxidant molecules. During the phagocytosis of microbes, there is a rapid increase of ROS (respiratory

burst) that enables phagocytic cells to degrade internalized particles and bacteria. Such an increase of ROS

can induce oxidative stress if not sufficiently counterbalanced by antioxidants, with potential negative effects

for the host fitness (Costantini, 2022).

Antioxidants act as free radicals scavengers by preventing and repairing damages caused by ROS, and

can be upregulated to prevent disease or to protect the organism from immunopathology induced by the

inflammatory response (Jena et al., 2023; Pham-Huy et al., 2008). Thus, upregulated baseline levels of endogenous antioxidants may help hosts to prevent a strong immune response and consequent oxidative stress. For example, white-browed sparrow weavers (*Plocepasser mahali*) with stronger superoxide dismutase (SOD) activity produced a smaller swelling response after an immune challenge (Cram et al., 2015). We are not aware of any studies comparing baseline levels of oxidative status markers between invasive and native species.

Important behavioural and life-history traits of successful invasive species, such as increased dispersal, endurance, and high reproductive rate may require particularly high levels of circulating oxygen. Haematocrit is an indirect marker of oxygen carrying capacity of blood (Ritchison, 2023) and thus, it might be upregulated in invasive species. This hypothesis was not supported by a study that compared cane toads (*Rhinella marina*) from the invasion front (spreaders) with cane toads from the core range (resident; Tracy et al., 2012). To our knowledge there is no study that compares levels of oxygen carrying capacity of blood between invasive and non-invasive species.

In this study, we compared the immune-physiological profile of the invasive Egyptian goose to that of two sympatric European waterfowl species. Specifically, we quantified a number of blood-based markers of immune function, oxidative status, and oxygen carrying capacity. The Egyptian goose is a successful invasive species which has been established in Europe since the 1980s and is currently expanding its range (Gyimesi and Lensink, 2012). The other two waterfowl species included in this study are native to the study area and share the same habitat with the Egyptian goose: the mallard (*Anas platyrhynchos*) and the mute swan (*Cygnus olor*).

In accordance with Lee and Klasing (2004), we predicted lower levels of baseline immune markers associated with systemic inflammatory response (i.e., granulocytes, macrophages, and lymphocytes) and higher levels of humoral and low-cost immunological effectors (i.e., lysozymes, immunoglobulins, bacteria killing ability, haemolysis and haemagglutination) in the invasive species compared to the native species. In addition, if Egyptian geese have reduced inflammatory responses, we might expect either (i) lower investment in endogenous antioxidant defences to save energy to allocate in other fitness traits or (ii) higher investment

in antioxidant defences for a better control of inflammatory responses compared to the other two species.

Baseline levels of oxidative damage are expected to show an opposite pattern to that of antioxidant defences.

Finally, we expect higher levels of oxygen carrying capacity in Egyptian geese compared to mallards and mute

swans.

Materials and methods

Study species and area

Mallard ducks and mute swans are waterfowl species native to Europe, while the Egyptian goose is native to

sub-Saharan Africa and was introduced to Europe in the twentieth century. The invasion of the Egyptian

goose started from the Netherlands, where they were introduced as ornamental species in urban parks. From

the 1980s onwards, Egyptian geese also invade Germany where its population size increased rapidly.

Nowadays, the Egyptian goose is the most common and widespread African waterfowl species in Europe

(Gyimesi and Lensink, 2012). Mallard ducks, mute swans and Egyptian geese are resident, territorial and

socially monogamous species. While mute swans and Egyptian geese show biparental care, mallards perform

female-only parental care. Additionally, mallard populations show adult sex-ratios which are usually male-

biased and forced extra-pair copulations frequently occur. Anseriformes are known for their aggressive

behaviour towards conspecifics and other species, particularly during the breeding season, to protect their

territory, nest and brood (Hohmann and Woog, 2021). All three species live in a variety of water habitats,

with a clear preference for freshwater ponds, rivers and lakes (both natural and artificial) sometimes in close

proximity to humans.

Adult individuals of the study species were live trapped in June and July 2016 in the Rhine and Mosel

area in Western Germany (50.4° N, 7.6° E; Fig. S1). All individuals of this study were locally breeding and

mostly in the phase of guiding goslings/ducklings at the time of sampling. Birds were caught using either

loops or landing nets. Each bird was individually ringed, and body mass and sex were recorded. A blood

sample from each bird was collected, either from the vena metatarsalia plantaris superficialis or the vena

ulnaris, using a needle of 0.06 mm diameter. Blood samples were kept cool during field procedures, then

centrifuged for 15 minutes to separate plasma from red blood cells which were stored separately in liquid

nitrogen within 8 hours from blood collection. At the end of fieldwork, samples were stored at -80 °C until

laboratory analyses.

Sample sizes for each assay were dependent on the total amount of plasma and red blood cells

available from each individual, and therefore differ between the tests that were carried out (Table 1).

Sampling was authorized in terms of animal welfare by the Landesuntersuchungsamt Rheinland-Pfalz (G 15-

20-005) and Landesamt für Natur, Umwelt und Verbraucherschutz Nordrhein-Westfalen (LANUV) (84-

08.04.2015.A266). The experimental procedures were approved by the Internal Committee for Ethics and

Animal Welfare of the Leibniz Institute for Zoo and Wildlife Research (permit #2014-11-03).

Laboratory analyses

Several immunological assays were carried out to characterize both the cellular and humoral components of

the adaptive and innate immune system of the study species. We used non-specific assays that had previously

been applied to samples from a variety of free-living avian species, including different waterfowl (Bourgeon

et al., 2010; Matson et al., 2006). Specifically, we counted basophils, eosinophils, heterophils and monocytes

as effectors of cellular innate immunity. We quantified the amounts of lysozyme, bacteria killing ability,

haemagglutination and haemolysis as markers for humoral innate immunity. We measured the total

immunoglobulin Y (IgY) concentration and the number of lymphocytes as markers of, respectively, humoral

and cellular adaptive immunity (Table 1; (Prüter et al., 2020)).

In addition to the analyses of immune markers, we measured the haematocrit as a proxy for the

oxygen carrying capacity of the blood. Further, we measured the total non-enzymatic antioxidant capacity

(OXY-adsorbent assay), and the activity of antioxidant enzymes superoxide dismutase (SOD) and glutathione

peroxidase, to estimate the antioxidant capacity of the blood. We measured the concentrations of protein

carbonyls and reactive oxygen metabolites (d-ROMS test) to evaluate blood levels of oxidative damage. More

details on the assays used for this study can be found in the Supplementary Material.

Statistical analyses

The dataset included 74 actively reproducing individuals of which 28 were Egyptian geese, 10 mallards, and

36 mute swans. We collected blood samples from both female and male Egyptian geese (females = 12; males

= 16) and mute swans (females = 21; males = 15), and only female mallards. Sample sizes varied slightly for

different immunological markers due to limited amounts of blood for certain individuals (Table 1).

We performed a principal component analysis (PCA) on the correlation matrix between all markers

of the immune system and of oxidative status to reduce the number of variables by identifying the main

sources of variation. Because PCAs do not accept missing data, we imputed missing values through the R

package 'missMDA' (Husson 2016) and performed a PCA on the completed dataset using the *prcomp* function

(R Core Team 2023). Then, we extracted the first three main axes of the PCA (PC1, PC2, and PC3) and tested

their correlations with values of each immune and oxidative status marker to assess how markers loaded on

each of the three axes. Correlations above the threshold value of 0.5 were considered as main correlations

(Table S1; Cohen, 2008).

To test for species differences in immune-physiological traits, we ran three linear regression models

(LMs) including each principal component (PC) as a response variable and species as the predictor variable.

Initially, we also included the interaction between sex and species to test for intraspecific and interspecific

differences between sexes. However, the interaction showed no significant effects and was removed from the

models.

Results

The first three PCs of the PCA explained 47.2% of the total variance in markers of the immune system and

oxidative status (Fig. S2). In particular, PC1 captured 22.85% of the total variance and reflected both cellular

innate immunity and adaptive immunity: the immune markers that strongly correlated with PC1 were

heterophil granulocytes (corr. = 0.63), lysozymes (corr. = 0.68), haemagglutination (corr. = 0.66), haemolysis

(corr. = 0.72), bacteria killing ability (corr. = -0.61), lymphocytes (corr. = 0.58) and IgY (corr. = 0.67). PC2

captured 12.62% of the total variance and was mainly associated with the basophil granulocytes (corr. = -

0.77), eosinophil granulocytes (corr. = -0.58), and two markers of the oxidative status, superoxide dismutase

(corr. = -0.64) and reactive oxygen metabolites (corr. = -0.54; Fig. 1). PC3 captured 9.56% of the total variance

and was associated with haemagglutination (corr. = -0.60) and haemolysis (corr. = -0.55; Table S1, Fig. S3).

ANOVA tests of the LMs showed significant differences among species in their immune characteristics

(Table 2). Pairwise comparisons of estimated marginal means showed that Egyptian geese had lower levels

of both cellular innate and adaptive immune markers compared to mallards and mute swans. Specifically,

mute swans showed the highest levels of immune function among the three species, while mallards showed

intermediate value. The bacteria killing ability was the only marker negatively correlated with the Principal

Component and is thus higher in Egyptian geese compared to mallards and mute swans. We also found

significantly higher levels of eosinophil and basophil granulocytes, superoxide dismutase and oxidative

damage (ROMs) in mallards compared to both Egyptian geese and mute swans. Those markers correlate

negatively with PC2, which means their values were higher for negative values of the PC2. No other significant

differences between species were found (Fig. 2; Table 2).

Discussion

In this study we compared the baseline immune-physiological profile of an invasive species, the Egyptian

goose, to that of two sympatric European waterfowl species, the mallard and the mute swan. As expected,

we found lower levels of cell-mediated immune markers in the blood of the invasive species compared to the

two native species. Specifically, the Egyptian goose showed lower levels of heterophil granulocytes and

lymphocytes. Surprisingly, we also found reduced levels of humoral defences, represented by non-specific

natural antibodies (haemagglutination), complement activity (haemolysis) and immunoglobulins-Y (IgY), but

higher level of plasma bacteria killing ability (BKA), in Egyptian geese compared to mute swans and mallards.

In addition, we found higher levels of basophil and eosinophil granulocytes, superoxide dismutase and

reactive oxygen metabolites in mallards compared to Egyptian geese and mute swans.

Immunological effectors

Heterophils and lymphocytes are the most abundant leucocytes in birds, comprising up to 95% of all white

blood cells (Davis et al., 2008; Ritchison, 2023). Heterophils are the major phagocytic leucocytes and

important mediators of acute inflammatory responses (Scanes et al., 2022). Lymphocytes play important roles in both cell- and antibody-mediated adaptive immunity. Specifically, T-lymphocytes attack infected or abnormal cells inducing inflammation, and B-lymphocytes produce immunoglobulins (Guriec et al., 2018). According to White and Perkins (2012), lower levels of leucocytes might be due to lower parasitic prevalence and/or reduced immune resistance to infections in the invasive species compared to the native species.

Recent studies found that German populations of Egyptian geese show lower prevalence of infections for a variety of blood parasites and macro-parasites compared to conspecifics from their native habitats and show lower seroprevalences of some common waterfowl viruses (Prüter et al., 2018, 2020). Such findings support the hypothesis that lower levels of leucocytes in Egyptian geese in their invasive range may be due to reduced prevalences of infectious agents, which would be compatible with the predictions of the evolution of increased competitive ability hypothesis. In addition, invasive hosts may tolerate certain levels of parasitic infections because it reduces the overall cost associated with the active removal of parasites (i.e., resistance) and associated sickness behaviours (Lennon et al., 2023; White and Perkins, 2012). Thus, Egyptian geese might benefit from reduced levels of heterophil granulocytes and lymphocytes compared to sympatric species, because it might prevent strong inflammatory responses in case of infection. Reduced cellular and inflammatory response were also found in other successful invasive species, such as the European paper wasp (*Polistes dominula*) and the house sparrow (*Passer domesticus*), compared to sympatric congeneric species (Lee et al., 2005; Wilson-Rich and Starks, 2010).

A previous study on this invasive population of Egyptian geese found higher levels of heterophils and leucocytes than Egyptian geese that were sampled in their native habitat, but higher variance of immune markers (Prüter et al., 2020). This might indicate that immune plasticity may be a key factor for successful invasions (Poirier, 2019; Pyšek and Richardson, 2010; White and Perkins, 2012). Invasive Egyptian geese might defend efficiently against novel pathogens by upregulating plasma BKA while simultaneously investing less energy in other costly immune functions. Our results point to reduced overall costs of immunity in the invasive population of Egyptian geese compared to mallards and mute swans in the same habitat. Therefore, invasive

Egyptian geese might have more energy to allocate towards reproduction, dispersal, territoriality, and survival than their competitors.

Contrary to our predictions, we found lower levels of humoral immune markers, such as haemagglutination, haemolysis, lysozymes and IgY, in Egyptian geese compared to mallards and mute swans. The only exception was represented by the plasma BKA, that was effectively higher in the invasive species. The mechanisms that allow plasma to kill bacteria are not well understood, but might be strongly dependent on antibacterial plasma proteins and peptides (Dugovich et al., 2019; Gao et al., 2015). Accordingly, Egyptian geese might invest relatively more energy in antimicrobial proteins and peptides, which come to lower (i) energetic costs compared to immune cells and (ii) fitness costs compared to inflammatory effectors.

Haemagglutination and haemolysis reflect responses to foreign antigens driven by natural antibodies and the complement system, respectively (Matson et al., 2006). Natural antibodies are produced in absence of antigenic stimulation, independently from previous infections, whereas the complement system can be activated directly by pathogens or indirectly by immunoglobulins (Ochsenbein and Zinkernagel, 2000). Natural antibodies and the complement system are part of the innate immune system and act as a first line of defence against infections. Therefore, lower levels of haemagglutination and haemolysis are in accordance with previous studies (Prüter et al., 2020, 2018) that Egyptian geese have a reduced prevalence of infections compared to mallards and mute swans.

IgY is the most abundant circulating immunoglobulin in birds (Ritchison, 2023). As every other immunoglobulin, IgY are produced by B-lymphocytes. Thus, the lower levels of constitutive IgY in Egyptian geese are in agreement with the reduced number of lymphocytes. Furthermore, native species often show higher levels of IgY and immune resistance due to the co-evolution between hosts and parasites (Koskella, 2018). For example, an experimental study on house finches (*Haemorhous mexicanus*) found higher levels of IgY in populations that co-evolved with the inoculated virus (*Mycoplasma gallisepticum*) compared to unexposed populations (Gates et al., 2021). Since higher resistance to infections leads to higher costs to host fitness (Rigby et al., 2002), invasive species may benefit from minor resistance to infections for reducing fitness costs (Medzhitov et al., 2012). This might explain why invasive species, which are mainly exposed to

novel pathogens, produce less immunoglobulins. Further studies are, however, needed to explore this

hypothesis in more detail.

Oxygen carrying capacity and oxidative status

We found no evidence for higher baseline oxygen carrying capacity in Egyptian geese compared to the other

species. It might be that differences in oxygen carrying capacity emerge only during the phase of range

expansion of invasive species as the haematocrit is rather sensitive to current metabolic requests (Ritchison

2023). All Egyptian geese included in our study were reproductively active individuals holding a territory, so

their immune-physiological function might differ from that of geese at the invasion front (Gyimesi and

Lensink, 2012). While there are studies showing different regulations of the immune system between

individuals living at the population core versus those at the front of invasion (Brown and Shine, 2014; Romeo

et al., 2023), further studies are required to assess the role of oxygen carrying capacity in individuals that are

at the invasion front.

Finally, we found that mallards had higher levels of eosinophils, basophils, oxidative damage (ROMs),

and activity of the antioxidant enzyme superoxide dismutase compared to Egyptian geese and mute swans.

In our study region, mallards are subject to increased competition and aggressive behaviour due to the

presence of Egyptian geese (Hohmann and Woog, 2021), which might result in increased metabolic activity

and ROS generation (Quque et al., 2022). It might also be that the higher pace of life of mallards compared

to Egyptian geese and mute swans determines higher baseline generation of ROS.

Conclusions

Our comparative analysis of multiple immune-physiological traits suggests that invasive Egyptians geese

might strategically reduce their investment in immune function compared to sympatric related native species

because of their lower parasite exposure. Native species that co-evolved with parasites in their natural habitat

might exhibit a stronger immune response aiming to clear the infection (i.e., resistance). By contrast, invasive

species are likely no longer exposed to the pathogens they encountered in their natural habitats. Moreover,

if they are exposed to novel pathogens in the invaded habitat, they might benefit from a less strong immune

response (i.e., more tolerant) to avoid immunopathological effects. Thus, invasive species might invest less energy to sustain immune function, which might (i) come at the benefit for other energy-demanding host traits, such as reproduction, dispersal, and territoriality, and (ii) bear minor risks of metabolic and behavioural impairments caused by acute inflammatory responses.

The results of our work also indicate that constitutive immune cells involved in acute inflammatory response might be the main downregulated effectors of immune function. This finding supports the hypothesis developed by Lee and Klasing (2004) that invasive species reduce the risk of systemic inflammatory responses as this is energetically and pathologically expensive and therefore ecologically maladaptive. Future work is needed to understand how baseline differences in immunological parameters translate into trade-offs with fitness traits, such as reproduction or survival. We also recognize that some variation in markers among species might also reflect species-specific differences that are not due to their history of colonisation. This is particularly true given that mallards show important behavioural, morphological, and pace-of-life differences with the other two species, which might contribute to among species variation in the regulation of oxidative status and immune function (i.e., relatively higher importance of eosinophil and basophil granulocytes as immune effectors, and upregulated oxidative status). On the other hand, physiological plasticity is an important driver of local acclimation and success of invasion (McCann et al., 2018). In accordance with Prüter et al. (2020), high levels of physiological plasticity in Egyptian geese might be one main mechanism underlying their successful invasion to Europe. Determining immune-physiological traits underlying the phenotype of successful invaders will foster our understanding of the proximate mechanisms through which species can successfully colonise novel environments and help to better predict novel alien species and potential mitigation strategies (Lennox et al., 2015).

.

Acknowledgements

S.M. was funded by the National Recovery and Resilience Plan (NRRP), Mission 4, Component 2, Investment 1.2 of the Italian Ministry for University and Research, funded by the European Union – NextGenerationEU, through the Young Researchers program (ID: SOE_0000189). The sample collection for this study was financially supported by the Ministry of Rhineland-Palatinate (Ministerium für Umwelt, Energie, Ernährung

und Forsten, Project Nr: Gz. 105-63 313/2015-40). We thank Sönke Twietmeyer and Niklas Böhm for their

great help with organizing and carrying out the field work. Also, we thank Lorena Derezanin, Sophie Ewert,

Lea Jäger, Oliver Krone, Manuela Merling de Chapa, Katja Pohle, Felix Prüter, and Jannis Twietmeyer for their

assistance during field or laboratory work. At last, we thank Rebecca Nagel for revising language and grammar

of the manuscript. The laboratory analysis was supported by funds from the Leibniz Institute for Zoo and

Wildlife Research, Berlin.

Bibliography

Blossey, B., Notzold, R., 1995. Evolution of Increased Competitive Ability in Invasive Nonindigenous Plants: A
Hypothesis. J. Ecol. 83, 887. https://doi.org/10.2307/2261425

Bourgeon, S., Kauffmann, M., Geiger, S., Raclot, T., Robin, J.-P., 2010. Relationships between metabolic
status, corticosterone secretion and maintenance of innate and adaptive humoral immunities in fasted re-
fed mallards. J. Exp. Biol. 213, 3810–3818. https://doi.org/10.1242/jeb.045484

Brown, G.P., Shine, R., 2014. Immune response varies with rate of dispersal in invasive cane toads (Rhinella
marina). PLoS ONE 9, e99734. https://doi.org/10.1371/journal.pone.0099734

Cohen, J., 2008. Statistical power analysis for the behavioral sciences, 2nd ed. Routledge.
https://doi.org/10.4324/9780203771587

Costantini, D., 2022. A meta-analysis of impacts of immune response and infection on oxidative status in
vertebrates. Conserv. Physiol. 10, coac018. https://doi.org/10.1093/conphys/coac018

Costantini, D., Møller, A.P., 2009. Does immune response cause oxidative stress in birds? A meta-analysis.
Comp. Biochem. Physiol. A. Mol. Integr. Physiol. 153, 339–344. https://doi.org/10.1016/j.cbpa.2009.03.010

Cram, D.L., Blount, J.D., York, J.E., Young, A.J., 2015. Immune response in a wild bird is predicted by oxidative
status, but does not cause oxidative stress. PLOS ONE 10, e0122421.
https://doi.org/10.1371/journal.pone.0122421

Davis, A.K., Maney, D.L., Maerz, J.C., 2008. The use of leukocyte profiles to measure stress in vertebrates: a
review for ecologists. Funct. Ecol. 22, 760–772. https://doi.org/10.1111/j.1365-2435.2008.01467.x

Dugovich, B.S., Crane, L.L., Alcantar, B.B., Beechler, B.R., Dolan, B.P., Jolles, A.E., 2019. Multiple innate
antibacterial immune defense elements are correlated in diverse ungulate species. PLOS ONE 14, e0225579.
https://doi.org/10.1371/journal.pone.0225579

Dunn, A.M., Torchin, M.E., Hatcher, M.J., Kotanen, P.M., Blumenthal, D.M., Byers, J.E., Coon, C.A.C., Frankel,
V.M., Holt, R.D., Hufbauer, R.A., Kanarek, A.R., Schierenbeck, K.A., Wolfe, L.M., Perkins, S.E., 2012. Indirect
effects of parasites in invasions. Funct. Ecol. 26, 1262–1274. https://doi.org/10.1111/j.1365-
2435.2012.02041.x

Gao, W., Xing, L., Qu, P., Tan, T., Yang, N., Li, D., Chen, H., Feng, X., 2015. Identification of a novel cathelicidin antimicrobial peptide from ducks and determination of its functional activity and antibacterial mechanism. Sci. Rep. 5, 17260. https://doi.org/10.1038/srep17260

Gates, D.E., Staley, M., Tardy, L., Giraudeau, M., Hill, G.E., McGraw, K.J., Bonneaud, C., 2021. Levels of pathogen virulence and host resistance both shape the antibody response to an emerging bacterial disease. Sci. Rep. 11, 8209. https://doi.org/10.1038/s41598-021-87464-9

Guriec, N., Bussy, F., Gouin, C., Mathiaud, O., Quero, B., Le Goff, M., Collén, P.N., 2018. Ulvan Activates Chicken Heterophils and Monocytes Through Toll-Like Receptor 2 and Toll-Like Receptor 4. Front. Immunol. 9, 2725. https://doi.org/10.3389/fimmu.2018.02725

Gyimesi, A., Lensink, R., 2012. Egyptian Goose Alopochen aegyptiaca: An introduced species spreading in and from the Netherlands. Wildfowl 62, 128–145.

Hasselquist, D., Nilsson, J.-Å., 2012. Physiological mechanisms mediating costs of immune responses: what can we learn from studies of birds? Anim. Behav. 83, 1303–1312. https://doi.org/10.1016/j.anbehav.2012.03.025

Henry, M., Leung, B., Cuthbert, R.N., Bodey, T.W., Ahmed, D.A., Angulo, E., Balzani, P., Briski, E., Courchamp, F., Hulme, P.E., Kouba, A., Kourantidou, M., Liu, C., Macêdo, R.L., Oficialdegui, F.J., Renault, D., Soto, I., Tarkan, A.S., Turbelin, A.J., Bradshaw, C.J.A., Haubrock, P.J., 2023. Unveiling the hidden economic toll of biological invasions in the European Union. Environ. Sci. Eur. 35, 43. https://doi.org/10.1186/s12302-023-00750-3

Hohmann, R., Woog, F., 2021. How aggressive are Egyptian Geese Alopochen aegyptiaca? Interactions with Greylag Geese Anser anser and other birds in an urban environment, Wildfowl.

Jena, A.B., Samal, R.R., Bhol, N.K., Duttaroy, A.K., 2023. Cellular Red-Ox system in health and disease: The latest update. Biomed. Pharmacother. 162, 114606. https://doi.org/10.1016/j.biopha.2023.114606

Koskella, B., 2018. Resistance gained, resistance lost: An explanation for host–parasite coexistence. PLOS Biol. 16, e3000013. https://doi.org/10.1371/journal.pbio.3000013

Lee, K.A., Klasing, K.C., 2004. A role for immunology in invasion biology. Trends Ecol. Evol. 19, 523–529. https://doi.org/10.1016/j.tree.2004.07.012

Lee, K.A., Martin, L.B., Wikelski, M.C., 2005. Responding to inflammatory challenges is less costly for a successful avian invader, the house sparrow (Passer domesticus), than its less-invasive congener. Oecologia 145, 243–250. https://doi.org/10.1007/s00442-005-0113-5

Lennon, R.J., Ronanki, S., Hegemann, A., 2023. Immune challenge reduces daily activity period in free-living birds for three weeks. Proc. R. Soc. B Biol. Sci. 290, 20230794. https://doi.org/10.1098/rspb.2023.0794

Lennox, R., Choi, K., Harrison, P.M., Paterson, J.E., Peat, T.B., Ward, T.D., Cooke, S.J., 2015. Improving science-based invasive species management with physiological knowledge, concepts, and tools. Biol. Invasions 17, 2213–2227. https://doi.org/10.1007/s10530-015-0884-5

Matson, K.D., Cohen, A.A., Klasing, K.C., Ricklefs, R.E., Scheuerlein, A., 2006. No simple answers for ecological immunology: relationships among immune indices at the individual level break down at the species level in waterfowl. Proc. R. Soc. B Biol. Sci. 273, 815–822. https://doi.org/10.1098/rspb.2005.3376

McCann, S.M., Kosmala, G.K., Greenlees, M.J., Shine, R., 2018. Physiological plasticity in a successful invader: rapid acclimation to cold occurs only in cool-climate populations of cane toads (Rhinella marina). Conserv. Physiol. 6. https://doi.org/10.1093/conphys/cox072

Medzhitov, R., Schneider, D.S., Soares, M.P., 2012. Disease Tolerance as a Defense Strategy. Science 335,
936–941. https://doi.org/10.1126/science.1214935

Mollot, G., Pantel, J.H., Romanuk, T.N., 2017. The Effects of Invasive Species on the Decline in Species
Richness, in: Advances in Ecological Research. Elsevier, pp. 61–83.
https://doi.org/10.1016/bs.aecr.2016.10.002

Ochsenbein, A.F., Zinkernagel, R.M., 2000. Natural antibodies and complement link innate and acquired
immunity. Immunol. Today 21, 624–630. https://doi.org/10.1016/S0167-5699(00)01754-0

Pham-Huy, L.A., He, H., Pham-Huy, C., 2008. Free radicals, antioxidants in disease and health. Int. J. Biomed.
Sci. IJBS 4, 89–96.

Poirier, M., 2019. A trade-off model for immunocompetence: The potential contribution of immunological
regulation in invasive vertebrate success. J. Exp. Zool. Part Ecol. Integr. Physiol. 331, 478–484.
https://doi.org/10.1002/jez.2314

Prüter, H., Czirják, G.Á., Twietmeyer, S., Harder, T., Grund, C., Mühldorfer, K., Lüschow, D., 2018. Sane and
sound: a serologic and molecular survey for selected infectious agents in neozootic Egyptian geese
(Alopochen aegyptiacus) in Germany. Eur. J. Wildl. Res. 64, 71. https://doi.org/10.1007/s10344-018-1231-9

Prüter, H., Franz, M., Twietmeyer, S., Böhm, N., Middendorff, G., Portas, R., Melzheimer, J., Kolberg, H., Von
Samson-Himmelstjerna, G., Greenwood, A.D., Lüschow, D., Mühldorfer, K., Czirják, G.Á., 2020. Increased
immune marker variance in a population of invasive birds. Sci. Rep. 10, 21764.
https://doi.org/10.1038/s41598-020-78427-7

Pyšek, P., Richardson, D.M., 2010. Invasive Species, Environmental Change and Management, and Health.
Annu. Rev. Environ. Resour. 35, 25–55. https://doi.org/10.1146/annurev-environ-033009-095548

Quque, M., Ferreira, C., Sosa, S., Schull, Q., Zahn, S., Criscuolo, F., Bleu, J., Viblanc, V.A., 2022. Cascading
Effects of Conspecific Aggression on Oxidative Status and Telomere Length in Zebra Finches. Physiol.
Biochem. Zool. 95, 416–429. https://doi.org/10.1086/721252

Rigby, M.C., Hechinger, R.F., Stevens, L., 2002. Why should parasite resistance be costly? Trends Parasitol.
18, 116–120. https://doi.org/10.1016/S1471-4922(01)02203-6

Ritchison, G., 2023. In a Class of Their Own: A Detailed Examination of Avian Forms and Functions,
Fascinating Life Sciences. Springer International Publishing, Cham. https://doi.org/10.1007/978-3-031-
14852-1

Romeo, C., Filipe, J., Wauters, L.A., Comazzi, S., Riva, F., Ferrari, N., 2023. Shifts in immune responses of an
invasive alien species: A test of the evolution of increased competitive ability hypothesis using American
Eastern gray squirrels in Italy. Sci. Total Environ. 900, 165747.
https://doi.org/10.1016/j.scitotenv.2023.165747

Sakai, A.K., Allendorf, F.W., Holt, J.S., Lodge, D.M., Molofsky, J., With, K.A., Baughman, S., Cabin, R.J., Cohen,
J.E., Ellstrand, N.C., McCauley, D.E., O'Neil, P., Parker, I.M., Thompson, J.N., Weller, S.G., 2001. The
population biology of invasive species. Annu. Rev. Ecol. Syst. 32, 305–332.
https://doi.org/10.1146/annurev.ecolsys.32.081501.114037

Scanes, C.G., Dridi, S., Sturkie, P.D. (Eds.), 2022. Sturkie's avian physiology, Seventh edition. ed. Academic
Press, an imprint of Elsevier, London, United Kingdom.

Torchin, M.E., Lafferty, K.D., Dobson, A.P., McKenzie, V.J., Kuris, A.M., 2003. Introduced species and their
missing parasites. Nature 421, 628–630. https://doi.org/10.1038/nature01346

Tracy, C.R., Christian, K.A., Baldwin, J., Phillips, B.L., 2012. Cane toads lack physiological enhancements for
dispersal at the invasive front in Northern Australia. Biol. Open 1, 37–42.
https://doi.org/10.1242/bio.2011024

Turbelin, A.J., Cuthbert, R.N., Essl, F., Haubrock, P.J., Ricciardi, A., Courchamp, F., 2023. Biological invasions
are as costly as natural hazards. Perspect. Ecol. Conserv. 21, 143–150.
https://doi.org/10.1016/j.pecon.2023.03.002

White, T.A., Perkins, S.E., 2012. The ecoimmunology of invasive species. Funct. Ecol. 26, 1313–1323.
https://doi.org/10.1111/1365-2435.12012

Wilson-Rich, N., Starks, P.T., 2010. The Polistes war: weak immune function in the invasive P. dominulus
relative to the native P. fuscatus. Insectes Sociaux 57, 47–52. https://doi.org/10.1007/s00040-009-0049-6

Zhang, L., Rohr, J., Cui, R., Xin, Y., Han, L., Yang, X., Gu, S., Du, Y., Liang, J., Wang, X., Wu, Z., Hao, Q., Liu, X.,
2022. Biological invasions facilitate zoonotic disease emergences. Nat. Commun. 13, 1762.
https://doi.org/10.1038/s41467-022-29378-2

471
472

Tables and figures

Table 1 – Sample size (n), mean and standard deviation (SD) for immunological effectors, and for haematocrit and markers of oxidative status.

Immunological effectors	Egyptian goose		Mallard		Mute swan	
	n	mean±SD	n	mean±SD	n	mean±SD
Lysozymes	27	3.07±2	10	3.89±1.85	34	7.06±3.17
Haemagglutination	21	6.66±1.42	10	7.1±0.99	34	7.67±1.49
Haemolysis	21	5.76±1.13	10	6±0.94	34	6.94±1.59
IgY	23	0.55±0.19	10	0.81±0.22	36	0.86±0.2
Bacteria Killing Ability	27	83.61±24.77	10	84.01±25.77	27	47.61±23.04
Basophils	28	3.29±2.72	10	10.53±6.33	36	3.98±2.44
Eosinophils	28	26.99±13.97	10	33.14±19.78	36	25.21±15.48
Heterophils	28	6.97±4.25	10	11.17±11.26	36	16.69±9.83
Lymphocytes	28	35.18±14.61	10	51.38±19.6	36	49.87±18.99
Monocytes	28	15.1±6.2	10	15.57±10.01	36	10.89±7.24
Haematocrit and oxidative status						
Haematocrit	24	44.79±8.5	7	40.57±8.81	21	39.9±4.51
OXY	27	197.47±46.99	10	180.61±46.22	32	207.69±43.73
ROMS	23	0.33±0.15	8	0.68±0.26	32	0.44±0.22
Protein Carbonyls	18	43.23±17.97	8	40.27±14.61	30	35.31±18.05
GPx	28	0.29±0.11	10	0.27±0.16	36	0.38±0.21
SOD	28	2.05±0.61	10	2.4±0.53	36	2.11±0.68

Table 2 – Between-species comparisons of estimated marginal means obtained from three ANOVA models applied on the Principal Components (PC). Significant results are shown in bold. ALEG = Egyptian geese; CYOL = mute swans; ANPL = mallards; f = females; m = males.

	Contrast	Estimate	SE	df	t.ratio	P-value
PC1	ALEG – ANPL	-1.64	0.50	71	-3.25	**<0.01**
	ALEG – CYOL	-2.92	0.35	71	-8.46	**<0.01**
	ANPL – CYOL	-1.28	0.49	71	-2.61	**0.03**
PC2	ALEG – ANPL	2.05	0.45	71	4.52	**<0.01**
	ALEG – CYOL	-0.19	0.31	71	-0.61	0.82
	ANPL – CYOL	-2.23	0.44	71	-5.09	**<0.01**
PC3	ALEG – ANPL	0.28	0.46	71	0.61	0.81
	ALEG – CYOL	0.26	0.31	71	0.82	0.69
	ANPL – CYOL	-0.02	0.45	71	-0.05	1.00

Fig. 1 – **Immune-physiological differences between the study species.** The position of each species is projected into the space defined by the first two Principal Components (PC) of a Principal Components Analysis performed on 16 parameters of immunity and oxidative status. Arrows represent the contribution of each parameter to PC1 and PC2. Species clusters are based on 95% confidence ellipses. Silhouettes of mallard and mute swan were sourced from PhyloPic. Each silhouette is available under CC BY-NC-SA 4.0 Deed licence, with credits to Maija Karala and Andy Wilson.

493　Fig. 2 – Estimated marginal means ± standard error (SE) of the ANOVA models based on Principal Components
494　(PC). Note that immune markers that are negatively correlated with the PC must be interpreted as higher
495　mean values (e.g., bacteria killing ability is negatively correlated with PC1, thus is higher in the Egyptian goose
496　compared to the other species). See Table S1 for results of correlations between the scores of PCs and the
497　immune markers. Significant results are indicated by different superscripts letters.