1 Weighting of sensory cues reflect changing patterns of visual investment
2 during ecological divergence in *Heliconius* butterflies.
3

4 José Borrero[1], Elisa Mogollon Perez[1], Daniel Shane Wright[1], Daniela Lozano[1,2], Geraldine

5 Rueda-Muñoz[1,2], Carolina Pardo-Diaz[2], Camilo Salazar[2], Stephen H. Montgomery[3]* &

6 Richard M. Merrill[1]*

7 Corresponding authors: jose.borrero@lmu.de (J.B.) & merrill@bio.lmu.de (R.M.M)

8
9   *[1] Division of Evolutionary Biology, Faculty of Biology, LMU Munich, Germany*
10   *[2] Faculty of Natural Sciences, Universidad del Rosario; Bogotá, Colombia*
11   *[3] School of Biological Science, University of Bristol, Bristol, United Kingdom*

Abstract

Integrating information across sensory modalities enables animals to orchestrate a wide

range of complex behaviours. The relative importance placed on one sensory modality over

another reflects the reliability of cues in a particular environment and corresponding

differences in neural investment. As populations diverge across environmental gradients, the

reliability of sensory cues may shift, favouring divergence in neural investment and the

weight given to different sensory modalities. During their divergence across closed-forest

and forest-edge habitats, closely related butterflies *Heliconius cydno* and *H. melpomene*

evolved distinct brain morphologies, with the former investing more in vision. Quantitative

genetic analyses suggest selection drove these changes, but their behavioural effects

remain uncertain. We hypothesised that divergent neural investment may alter sensory

weighting. We trained individuals in an associative learning experiment using multimodal

colour and odour cues. When positively rewarded stimuli were presented in conflict pairing

positively trained colour with negatively trained odour, and *vice-versa*, *H. cydno* favoured

visual cues more strongly than *H. melpomene*. Hence, differences in sensory weighting may

evolve early during divergence and are predicted by patterns of neural investment. These

findings, alongside other examples, imply that differences in sensory weighting stem from

divergent investment as adaptations to local sensory environments.

Introduction

Animals rely on multiple sensory systems to perceive and navigate their environments. The integration of environmental information across different sensory modalities (e.g., vision and olfaction) facilitates complex behaviours such as foraging, predator avoidance, and mate finding [1,2]. Cues perceived by these different sensory modalities can work in concert to enhance signal detection [3], but individuals may also prioritise one sensory modality over another [1,4]. This prioritisation of different sensory modalities is likely shaped by divergent selection acting on heritable variation across different habitats [5,6], leading to evolutionary shifts in their relative importance between populations that exploit different habitats.

Evidence for divergence in sensory weighting, defined here as the relative emphasis given to different modalities during behavioural decision-making [4,7,8], has been observed across a handful of taxa where populations or closely related species exploit different sensory environments. For instance, stingless bees rely less on visual cues, such as colour, than honeybees, possibly due to reduced investment in eye size [9]. In butterflies and moths, brain areas processing sensory information are among the most variable structure between species, and studies investigating multisensory integration emphasise associations between interspecific differences in neuroanatomy and behaviour [10,11]. For example, while hawkmoths broadly use both visual and olfactory cues during foraging [4,12], nocturnal species invest relatively more in brain regions associated with olfactory processing, whereas diurnal species are characterised by larger investments in visual neuropils. This shift in neural investment is mirrored in the relative importance given to olfactory and visual cues during foraging [7]. How such specialisations in the weighting of sensory cues evolve over smaller evolutionary time scales and across more subtle ecological gradients at the early stages of divergence is less clear.

The Neotropical *Heliconius* butterflies provide an excellent opportunity to study how sensory environments can shape multisensory integration during the early stages of divergence. Although olfaction also plays a role [8,13–17], these butterflies rely heavily on vision during foraging [18,19], mate recognition [20–22] and oviposition [23,24]. In *Heliconius*, speciation is often associated with ecological transitions across forest types [25–28], such that closely related species experience different sensory conditions [26,29,30]. Sister taxa are also known to differ in multiple aspects of both the peripheral and central sensory pathways, likely reflecting adaptation to local sensory conditions [31–33].

If these shifts in neural investment and sensory weighting are the result of divergent selection imposed by differences in sensory ecology, we expect populations diverging across similar environmental gradients to show associated changes in behaviour [34–36]. The *Heliconius* species pair, *H. melpomene* and *H. cydno,* are a well-studied case of ecological speciation which provide an opportunity to test this hypothesis. These species exist in "mosaic sympatry", where *H. melpomene* inhabits open forest edges and *H. cydno* occupies more closed canopy, inner forest habitats (Figure 1) [26,37,38]. *H. cydno* has larger eyes and invests more heavily in brain structures associated with vision when compared to *H. melpomene* . Multiple sources of data strongly suggesting these differences have likely been shaped by adaptive processes [32,33,39]. In *Heliconius*, visual information is relayed from the optic lobes to the mushroom bodies (the major sensory integration centres), which also receives input from the olfactory pathway [11]. Changes in investment in specific sensory pathways may therefore affect the strength of input received, consequently influencing the weight given to different sensory cues during behavioural decision-making processes.

Here, we test how *H. cydno* and *H. melpomene* prioritise visual and olfactory information during foraging, predicting that increased investment in the visual pathway may lead to *H. cydno* prioritising visual cues. To this end, we trained butterflies to a combination of visual and olfactory stimuli using flower models, and then quantified the relative weights given to visual and olfactory cues by presenting the rewarded stimuli in conflict.

Materials & Methods

Butterfly rearing and maintenance

We maintained outbred stock populations of *H. cydno* and *H. melpomene* at the Experimental Station José Celestino Mutis – Universidad del Rosario, near La Vega, Colombia (5.0005° N, 74.3394° W), established from wild individuals caught in the local area. Outdoor cages (1x3x2 m) housed butterfly groups with access to a ~20% sugar solution*, Gurania, Lantana* and *Psiguria spp.* flowers as pollen sources, and *Passiflora* vines for oviposition. All individuals used in the experiments were reared under common garden conditions.

Experimental set-up and stimuli

To test for visual and olfactory preferences, learning and sensory weighting, we presented male and female butterflies with artificial feeders with both visual (red or blue) and olfactory (lavender or rose essential oils) stimuli on two 25x25 cm hard PVC sheet 'arrays' separated by 200 cm in an 1x3x2m insectary (Figure 2A). Each array had four feeders made from a 5

ml Eppendorf tube surrounded by a 3 cm radius circle printed on waterproof paper. One array all red circles, and on the other all blue. Olfactory stimuli were presented in a 200µl PCR-tube below the paper circles, containing either lavender (*Lavandula angustifolia*) or rose *(Rosa damascena)* essential oils diluted with paraffin oil to 1% concentration (Figure 2B). By keeping the olfactory cues separate underneath the waterproof paper and not mixing with the feeding solution, we ensured that olfactory associations were not confounded by gustatory reception. All four feeders on the same array had the same olfactory stimuli to avoid odour mixing.

We trained and tested butterflies in same sex groups of 20-24 uniquely marked individuals (10-12 per species). Butterflies were introduced to the experimental cages at least 12 hours prior to the initial testing period to allow for acclimatisation and were food deprived overnight. Butterflies ware trained to associate the combination of blue colour and rose odour with a positive reward [blue/rose] (20% sugar-water solution) and the combination of [red/lavender] with a negative stimulus (~18% Vitamin C solution; [8,40]). Butterflies were tested with the visual and olfactory in the *trained* test (i.e. [blue/rose] vs [red/lavender]: days 1 and 5), in '*conflict'* (i.e. [blue/lavender] vs [red/rose]: day 6), or with *colour alone* (i.e. [blue] vs [red]: day 7) (Figure 3A). In contrast to the training phases, during testing, clean Eppendorf tubes were only filled with water (i.e., with no positive or negative 'reward') and fresh visual stimuli were used.

Training and testing

i) Naive test (Day 1): We recorded the naive preferences toward combinations of the visual and olfactory stimuli [blue/rose] vs [red/lavender] for one hour (10:00-11:00) using only water feeders. Each butterfly was then hand fed once on a positive [blue/rose (+)], and once on a negative rewarded feeder [red/lavender (-)], and then allowed to feed freely for 2 hours from the positive and negative rewarded feeders (11:00-13:00).

ii)Training (Days 2-4): Training was reinforced on experimental days 2-4. First, each individual was hand fed once on a positive [blue/rose (+)], and once on a negative rewarded feeder [red/lavender (-)]. Afterwards, butterflies were allowed to feed freely for 3 hours (10:00-13:00) from the positive or negative rewarded feeders. The feeders' position was randomised (coin toss) every day to prevent positional learning.

iii) Trained test (Day 5*)*: On day 5, we recorded the trained preferences for one hour (between 10:00-11:00) using only water in the feeders with the stimuli in *trained* conditions

(i.e. [blue/rose] vs [red/lavender]). This was immediately followed by two hours of

reinforcement training with rewarded feeders, i.e. [blue/rose (+)] and [red/lavender (-)].

iv) Conflict test (Day 6): To test for differences in sensory weighting, we presented the visual

and olfactory in conflict (i.e. [blue/lavender] vs [red/rose]; Figure 3A). As for the previous

tests, feeding behaviour was recorded for one hour (between 10:00-11:00) using only water

in the feeders. This was then followed by reinforcement training for two hours, i.e. [blue/rose

(+)] and [red/lavender (-)].

v) Colour test (Day 7): We tested visual preferences in isolation for one hour (between

10:00-11:00) (i.e., blue vs red, without olfactory stimuli, and using water only).

151   Behavioural tests

All feeding test assays [(i) Naive, (ii) Trained, (iii) Conflict and (iv) Colour] were filmed from

above with an AKASO Brave 4 Pro action camera. We scored the videos in BORIS [41],

recording the total number of feeding attempts directed toward each stimulus per individual.

Feeding attempts were defined as a proboscis extension within the coloured circle feeders.

We scored instances where an individual moved outside the circle and subsequently

returned as a new feeding attempt. In contrast, we scored instances where an individual

remained within a circle and probed continuously as a single attempt, unless the proboscis

was retracted for 3 seconds or longer.

161   Training criterion

We established a stringent training criterion to incorporate only individuals that had

successfully learned the correct stimulus combination, following similar learning studies in

lizards [42,43] and butterflies [19]. Out of 260 individuals, 73 were excluded from the analysis

for not attempting to feed on the trained test (day 5) (Table S2). To investigate species

differences in sensory weighting, we included only those individuals that demonstrated

robust associative learning of the trained cue combinations (i.e. >=75% of feeding choices

towards the positive multimodal stimuli [blue/rose] on the trained test (day 5)), resulting in a

reduced sample size from 187 to 73 individuals. Although biased towards *H. melpomene* (52

individuals vs 21 *H. cydno* individuals), this rigorous criterion ensured that our analysis

focused solely on individuals who had successfully learned. However, our results did not

quantitatively change when including all 117 individuals with a lower threshold of learning

(i.e. >50% correct; see supplementary results Table S1 and Tables S5-S6).

<u>Statistical analysis</u>

We used general linear mixed models (GLMM) with a binomial distribution in the *lme4* [44] in

R [45] to assess if species differed in their feeding attempts when the positively rewarded

stimuli agreed or were in conflict. The dependent variable was the proportion of choices

toward the blue feeders and we included species, and treatment [Trained test (day 5),

Conflict test (day 6) or Colour test (day 7)], and their interaction as fixed factors. To account

for the varying number of choices made by each butterfly, individual observations are

weighted by their total number of feeding attempts [46]. Individual ID nested within group ID

was included as random effects to account for repeated measures (Table S1). We applied a

stepwise model reduction approach to identify the minimum adequate statistical models,

using likelihood ratio tests (LRT) via the *drop1* function to determine significance. When fixed

effects or their interactions were significant, we conducted pairwise comparisons tests using

post hoc Bonferroni corrections with the *emmeans* package [47]. Plots were made using

ggplot2 [48].

Results
189

Among the 73 individuals (52 *H. melpomene* and 21 *H. cydno*) that passed our stringent

training criterion (i.e., >75% correct choices on day 5), we conducted tests to quantify the

species' reliance on visual and olfactory cues. Importantly, during these tests, all feeders

were unrewarded. *H. cydno* did not differ from *H. melpomene* in the trained test (Table 1

post-hoc: *H. cydno* Trained vs. *H. melpomene* Trained, Z= -1.375, P= 1.0). In contrast, *H.*

melpomene and *H. cydno* responded differently when presented with the olfactory and visual

stimuli in conflict, (Figure 3B). Importantly, the interaction between species and trial type was

retained in our model, confirming that this was an effect of how the stimuli were presented

(species*treatment LRT: 2ΔlnL = 22.876, df=2, p < 0.001. Table S3). In the conflict test, *H.*

cydno fed mostly on the blue feeder, while *H. melpomene* primarily chose the feeder with the

rose scent, suggesting a difference between these species in how olfactory and visual cues

are weighed during behavioural decisions (Table 1, post-hoc: *H. cydno* Conflict vs. *H.*

melpomene Conflict, Z= 3.210, P= 0.019). We found no effect of sex, which was excluded

from our model (sex LRT: 2ΔlnL = 3.067, df=1, p = 0.079. Table S3).

Our post-hoc analysis revealed that individuals of both species made fewer feeding attempts

on the blue feeder when the stimuli were presented in conflict, suggesting that both species

use both visual and olfactory stimuli when making foraging decisions (Table 1, post-hoc: *H.*

cydno Trained vs. *H. cydno* Conflict, Z= 6.932, P= <.0001; *H. melpomene* Trained vs. *H.*

melpomene Conflict, Z=20.614, P= <.0001). *H. cydno* made fewer feeding attempts on the

blue feeder when the positively trained olfactory and visual stimuli were in conflict, than when they were in the trained combinations confirming the use of both cues when present. However, there was no distinction between the trained test and the colour-only test (Table 1, post-hoc: *H. cydno* Trained vs. *H. cydno* Colour, Z= 2.321, P= 0.304), further emphasising that *H. cydno* was more attentive to the visual cue in our assay. Conversely, *H. melpomene* differed between the trained test and the colour-only test, indicating greater use of the olfactory stimulus (Table 1, post-hoc: *H. melpomene* Trained vs. *H. melpomene* Colour, Z= 10.920, P= <.0001)

Discussion

We explored how two closely related butterfly species, *H. cydno* and *H. melpomene*, weigh visual and olfactory information in a foraging context. Our findings indicate a significant divergence in behaviour between these two species, with *H. cydno* favouring visual cues over olfactory cues compared to *H. melpomene.* Importantly, our results mirror differences in brain morphology and suggest that divergence in the sensory environment may have led to changes in brain composition and how these species process, integrate, and prioritise information across sensory modalities.

To assess how *H. cydno* and *H. melpomene* prioritise visual and olfactory information, we presented trained butterflies with conflicting stimuli - pairing positively trained colours with negatively trained odours, and *vice versa*. Both species chose the 'correct' colour less frequently in these conflict scenarios compared to their trained preferences or when presented with colours alone, indicating that both visual and olfactory cues influence foraging decisions. However, when presented with conflicting cues, the species differed in their reliance on each sensory modality. *H. cydno* predominantly fed on the positively rewarded visual cue (the blue feeder). In contrast, *H. melpomene* showed a stronger preference for the feeder containing the positive trained odour stimulus (rose), indicating a difference between two species in the weight given to the two cues during behavioural decision-making. Additionally, *H. melpomene* differed between the trained and colour only test, making fewer correct choices when only colours were presented, again suggesting a larger reliance on the olfaction stimulus compared to *H. cydno*. As the visual and olfactory cues employed in our study are not assumed to be ecologically relevant, we suggest they reflect the baseline relative preferences of the species for each sensory modality. Given that the behavioural variation we observe is from common-garden-reared individuals, these interspecific shifts are likely to have a genetic basis. In addition, our results align with divergent patterns of investment in sensory pathways between the two species, whereby *H.*

cydno exhibits greater relative investment in the visual system [32,33,39]. Therefore, heritable shifts in neural investment and sensory weighting may result from divergent selection imposed by differences in sensory ecology.

One potential limitation of our study is that we managed to successfully train a larger number of *H. melpomene* than *H. cydno* individuals. This discrepancy in training could be due to *H. melpomene* being more active during the trials, possibly due to the outdoor insectaries being found in the forest edge, which closely resembles the *H. melpomene* habitat [26,30]. Indeed, in contrast to *H. cydno, H. melpomene* is frequently observed flying next to the insectaries (pers. obs. J. Borrero). Despite this difference, individuals of both species successfully learned the rewarded combination under our rigorous training criteria (>= 75% correct choices) and our main conclusions hold even under less rigorous training criteria (see supplementary results).

Adaptation to the local sensory environment may play an important role during the early stages of divergence. The environmental gradient observed in our studied species, *H. cydno* and *H. melpomene*, from closed forest canopy to open forest edge [26,30], resembles the transition across an altitudinal gradient, from complex broad-leaf forest environments to dry Andean forests seen in another distantly related *Heliconius* species-pair, *H. erato* and *H. himera* [49]. *H. erato cyrbia* and *H. himera* exhibit similar patterns of neural divergence, with *H. erato cyrbia* showing increased, heritable investment in the visual pathway, but with *H. himera* also showing some increased investment in the olfactory pathway [8,31]. Despite methodological differences (group vs individual testing and the exact visual and olfactory stimuli used), our results qualitatively reassemble a previous study in *H. erato* and *H. himera* [8]. Together, these findings demonstrate similar shifts in sensory weighting among *Heliconius* butterflies, with species inhabiting more closed habitats investing more in perceiving and processing visual information and favouring visual over olfactory cues. This suggests changes in the light environment is a common source of divergent selection across the two species pairs.

More broadly, changes in sensory weighting have been observed across different taxa and across different evolutionary timescales. For instance, urban Great tits prefer olfactory cues, while forest birds rely more on visual cues [50]. Nine-spined sticklebacks show plasticity in their sensory reliance, for instance compared to fish reared in visually unrestricted habitats, individuals raised in visually restricted environments display a larger reliance on chemical over visual cues. This suggests neural plasticity may underpin initial behavioural responses

to novel environments that are later consolidated by selection [51]. In threespine stickleback

fish, populations in different lakes segregate into limnetic and benthic ecomorphs along an

environmental gradient [52]. These stickleback populations show repeated evolutionary

shifts in brain regions that are correlated with differences in their sensory environments;

limnetic species have larger structures for visual processing and benthic ectomorphs

possess larger olfactory bulbs. Comparing across larger evolutionary timescales, studies

have found that nocturnal hawkmoths prioritise olfactory cues while foraging, as evidenced

by their increased investment in olfactory processing. Conversely, diurnal hawkmoths show

larger investment in visual neuropils and place a greater emphasis on visual cues. These

changes in the relative weight given to different sensory modalities, and the accompanying

changes in brain investment, are likely driven by the transition from a nocturnal to a diurnal

habitat and the associated alterations in their sensory environment [7]. Combined with our

findings, these studies underscore the role of environmental factors in driving changes in

neuroanatomy, sensory integration, and behaviour across species.

In conclusion, our results highlight shifts in sensory weighting between species associated

with neural investment and ecological divergence, consistent with patterns across *Heliconius*

butterflies. We demonstrated that *H. cydno*, from a lower light habitat [30] which is

characterised with increased visual investment [32,33], favours visual cues more strongly

than *H. melpomene*. Alongside previous results in *Heliconius* butterflies and other taxa

making [7,8], our research reveals repeated shifts in sensory weighting associated with

changes in neural investment and habitat use. This highlights the potential role for the

sensory environment to shape differences in brain structure, which subsequently may

influence key sensory behaviours crucial during the initial stages of species divergence.

Figure 1: Habitat preference and the sensory environment are associated with differences in brain investment in *Heliconius* butterflies. *H. cydno* invests more in visual neuropils [32] visual periphery [33, 39] and inhabits more-closed canopy forest habitats, than its close relative, *H. melpomene,* which is found in the forest edge [21, 30].

Figure 2: Experimental set-up A) Groups of *H. cydno* and *H. melpomene* were presented with artificial flower arrays at the end of the cages and were trained to a certain colour/odour combination. **B)** Artificial flower array containing the multimodal visual cues (coloured circles) and olfactory cue (PCR tube containing essential oil).

Figure 3: *H. cydno* and *H. melpomene* **weight** visual and olfactory cues **differently while** foraging. **A)** Butterflies were trained to feed on blue feeders with a rose oil odour and negatively reinforced red feeders with a lavender smell. In the conflict test, the positively trained feeders were presented in combination with the negatively trained odours and vice versa. During training positive stimuli were rewarded with sugar water and negative stimuli with (~18% Vitamin C), but during all tests (naive, trained, conflict and colour) feeders were only filled with water (i.e., with no positive or negative reward). Training and reinforcement training periods are delineated by white rectangles, while behavioural tests are indicated by grey rectangles (water only). **B)** Proportion of feeding attempts towards blue feeder for *H. cydno* and *H. melpomene*. Each point represents the feeding choices of an individual; solid dots and error bars represent the mean choices per species ±95% confidence intervals. Blue circles represent *H. cydno* and orange triangles represent *H. melpomene*, shapes in grey represent naive individuals that did not train. On naive and trained tests, feeder combinations were blue/rose and red/lavender; in the conflict test, the positively trained colour (blue) was presented in conflict with the negatively rewarded odour (lavender) and *vice versa*. In the colour tests, individuals were presented only with red or blue feeders without an odour. Only individuals that passed our training criteria (>=75% on the trained day) are shown (see methods).

309
310 **Table 1:** Pairwise p-value comparison matrix of post-hoc test with Bonferroni correction for the minimal adequate
311 model for the effects of the interaction between species and treatment (Trained, Conflict, and Colour) on choices
312 towards blue feeders. Only individuals that passed our training criteria (>=75% on the trained day) are included
313 Results are from a binomial GLMM: *see methods for statistical details. Proportion blue ~ Species * Level +*
314 *(1|Group/ID)*
315

	H. cydno Trained	*H. melpomene* Trained	*H. cydno* Conflict	*H. melpomene* Conflict	*H. cydno* Colour	*H. melpomene* Colour
H. cydno Trained	[2.329]	1	<.0001	<.0001	0.3041	0.008
H. melpomene Trained		[2.874]	<.0001	<.0001	0.0116	<.0001
H. cydno Conflict			[0.465]	0.0199	<.0001	1
H. melpomene Conflict				[-0.601]	<.0001	<.0001
H. cydno Colour					[1.639]	1
H. melpomene Colour						[0.989]

Acknowledgements

319 We thank Universidad del Rosario for insectary access at JCM and Isabel León for

320 assistance, as well as Amulya Hosur and Alejandro Mancino for video scoring help.

Funding information

323 Research Funded by ERC Starter Grant 851040 to R.M.M.

Conflict of interest

326 The authors have no conflict of interest to declare.

References

1. Talsma D, Senkowski D, Soto-Faraco S, Woldorff MG. 2010 The multifaceted interplay between attention and multisensory integration. *Trends in Cognitive Sciences* **14**, 400–410. (doi:10.1016/j.tics.2010.06.008)

2. Stevens M. 2013 *Sensory ecology, behaviour, and evolution*. First edition. Oxford, United Kingdom: Oxford University Press.

3. Leonard AS, Dornhaus A, Papaj DR. 2011 Flowers help bees cope with uncertainty: signal detection and the function of floral complexity. *Journal of Experimental Biology* **214**, 113–121. (doi:10.1242/jeb.047407)

4. Balkenius A, Rosén W, Kelber A. 2006 The relative importance of olfaction and vision in a diurnal and a nocturnal hawkmoth. *J Comp Physiol A* **192**, 431–437. (doi:10.1007/s00359-005-0081-6)

5. Dell'Aglio DD, Rivas-Sánchez DF, Wright DS, Merrill RM, Montgomery SH. 2023 The Sensory Ecology of Speciation. *Cold Spring Harb Perspect Biol* , a041428. (doi:10.1101/cshperspect.a041428)

6. Raine NE, Chittka L. 2007 The Adaptive Significance of Sensory Bias in a Foraging Context: Floral Colour Preferences in the Bumblebee Bombus terrestris. *PLOS ONE* **2**, e556. (doi:10.1371/journal.pone.0000556)

7. Stöckl A, Heinze S, Charalabidis A, El Jundi B, Warrant E, Kelber A. 2016 Differential investment in visual and olfactory brain areas reflects behavioural choices in hawk moths. *Scientific Reports* **6**, 1–10. (doi:10.1038/srep26041)

8. Dell'Aglio DD, McMillan WO, Montgomery SH. 2022 Shifting balances in the weighting of sensory modalities are predicted by divergence in brain morphology in incipient species of Heliconius butterflies. *Animal Behaviour* **185**, 83–90. (doi:10.1016/j.anbehav.2022.01.003)

9. Koethe S, Fischbach V, Banysch S, Reinartz L, Hrncir M, Lunau K. 2020 A Comparative Study of Food Source Selection in Stingless Bees and Honeybees: Scent Marks, Location, or Color. *Frontiers in Plant Science* **11**. (doi:10.3389/fpls.2020.00516)

10. Couto A, Wainwright JB, Morris BJ, Montgomery SH. 2020 Linking ecological specialisation to adaptations in butterfly brains and sensory systems. *Current Opinion in Insect Science* **42**, 55–60. (doi:10.1016/j.cois.2020.09.002)

11. Couto A *et al.* 2023 Rapid expansion and visual specialisation of learning and memory centres in the brains of Heliconiini butterflies. *Nat Commun* **14**, 4024. (doi:10.1038/s41467-023-39618-8)

12. Balkenius A, Kelber A. 2006 Colour preferences influences odour learning in the hawkmoth, Macroglossum stellatarum. *Naturwissenschaften* **93**, 255–258. (doi:10.1007/s00114-006-0099-9)

13. Andersson S, Dobson HEM. 2003 Behavioral foraging responses by the butterfly Heliconius melpomene to Lantana camara floral scent. *Journal of Chemical Ecology* **29**, 2303–2318. (doi:10.1023/A:1026226514968)

14. Thiele SC, Rodrigues D, Moreira GRP. 2016 Oviposition in Heliconius erato (Lepidoptera, Nymphalidae): how Essential Is Drumming Behavior for Host-Plant Selection? *J Insect Behav* **29**, 283–300. (doi:10.1007/s10905-016-9559-z)

15. Darragh K *et al.* 2017 Male sex pheromone components in Heliconius butterflies released by the
androconia affect female choice. *PeerJ* **5**, e3953. (doi:10.7717/peerj.3953)

16. Silva DS, Barp EA, Kucharski LCR, Moreira GRP. 2018 Sensing the Plant Surface Prior to Feeding
and Oviposition: Differences in External Ultrastructure and Function Among Tarsi of Heliconius
erato. *Neotrop Entomol* **47**, 85–95. (doi:10.1007/s13744-017-0508-0)

17. van Schooten B, Meléndez-Rosa J, van Belleghem SM, Jiggins CD, Tan JD, Owen McMillan W, Papa
R. 2020 Divergence of chemosensing during the early stages of speciation. *Proceedings of the
National Academy of Sciences of the United States of America* **117**, 16438–16447.
(doi:10.1073/pnas.1921318117)

18. Zaccardi G, Kelber A, Sison-Mangus MP, Briscoe AD. 2006 Color discrimination in the red range
with only one long-wavelength sensitive opsin. *Journal of Experimental Biology* **209**, 1944–1955.
(doi:10.1242/jeb.02207)

19. Toure MW, Young FJ, McMillan WO, Montgomery SH. 2020 Heliconiini butterflies can learn time-
dependent reward associations. *Biology Letters* **16**, 20200424. (doi:10.1098/rsbl.2020.0424)

20. Jiggins CD, Naisbit RE, Coe RL, Mallet J. 2001 Reproductive isolation caused by colour pattern
mimicry. *Nature* **411**, 302–305. (doi:10.1038/35077075)

21. Estrada C, Jiggins CD. 2008 Interspecific sexual attraction because of convergence in warning
colouration: Is there a conflict between natural and sexual selection in mimetic species? *Journal
of Evolutionary Biology* **21**, 749–760. (doi:10.1111/j.1420-9101.2008.01517.x)

22. Merrill RM, Chia A, Nadeau NJ. 2014 Divergent warning patterns contribute to assortative mating
between incipient Heliconius species. *Ecology and Evolution* **4**, 911–917. (doi:10.1002/ece3.996)

23. Gilbert LE. 1982 The Coevolution of a Butterfly and a Vine. *Scientific American* **247**, 110–121.
(doi:10.1038/scientificamerican0882-110)

24. Dell'Aglio DD, Losada ME, Jiggins CD. 2016 Butterfly learning and the diversification of plant leaf
shape. *Frontiers in Ecology and Evolution* **4**, 1–7. (doi:10.3389/fevo.2016.00081)

25. Jiggins CD, Mallet J. 2000 Bimodal hybrid zones and speciation. *Trends in Ecology & Evolution* **15**,
250–255. (doi:10.1016/S0169-5347(00)01873-5)

26. Estrada C, Jiggins CD. 2002 Patterns of pollen feeding and habitat preference among Heliconius
species. *Ecological Entomology* **27**, 448–456. (doi:10.1046/j.1365-2311.2002.00434.x)

27. Jiggins CD. 2008 Ecological speciation in mimetic butterflies. *BioScience*

28. Merrill RM *et al.* 2015 The diversification of Heliconius butterflies: What have we learned in 150
years? *Journal of Evolutionary Biology* **28**, 1417–1438. (doi:10.1111/jeb.12672)

29. Endler JA. 1993 The Color of Light in Forests and Its Implications. *Ecological Society of America* **63**,
1–27.

30. Seymoure BM. 2016 Heliconius In A New Light: The Effects of Light Environments on Mimetic
Coloration, Behavior, and Visual Systems. Doctoral Dissertation, Arizona State University, Tempe,
AZ. See https://hdl.handle.net/2286/R.I.39452.

31. Montgomery SH, Merrill RM. 2017 Divergence in brain composition during the early stages of
ecological specialization in Heliconius butterflies. *Journal of Evolutionary Biology* **30**, 571–582.
(doi:10.1111/jeb.13027)

32. Montgomery SH, Rossi M, McMillan WO, Merrill RM. 2021 Neural divergence and hybrid
disruption between ecologically isolated *Heliconius* butterflies. *Proc. Natl. Acad. Sci. U.S.A.* **118**,
e2015102118. (doi:10.1073/pnas.2015102118)

33. Wright DS, Rodriguez-Fuentes J, Ammer L, Darragh K, Kuo C-Y, McMillan WO, Jiggins CD,
Montgomery SH, Merrill RM. 2023 Adaptive divergence in the eyes of *Heliconius* butterflies likely
contributes to pre- and post-mating isolation. (doi:10.1101/2023.10.26.564160)

34. Johnson MA, Revell LJ, Losos JB. 2009 BEHAVIORAL CONVERGENCE AND ADAPTIVE RADIATION:
EFFECTS OF HABITAT USE ON TERRITORIAL BEHAVIOR IN ANOLIS LIZARDS. *Evolution* **64**, 1151–
1159. (doi:10.1111/j.1558-5646.2009.00881.x)

35. Henry CS, Brooks SJ, Duelli P, Johnson JB, Wells MM, Mochizuki A. 2012 Parallel evolution in
courtship songs of North American and European green lacewings (Neuroptera: Chrysopidae):
PARALLEL EVOLUTION IN LACEWING SONGS. *Biological Journal of the Linnean Society* **105**, 776–
796. (doi:10.1111/j.1095-8312.2011.01845.x)

36. Bolnick DI, Barrett RDH, Oke KB, Rennison DJ, Stuart YE. 2018 (Non)Parallel Evolution. *Annu. Rev.*
Ecol. Evol. Syst. **49**, 303–330. (doi:10.1146/annurev-ecolsys-110617-062240)

37. Smiley J. 1978 The Host Plant Ecology of Heliconius Butterflies in Northeastern Costa Rica.
Universtiy of Texas at Austin.

38. Mérot C, Salazar C, Merrill RM, Jiggins CD, Joron M. 2017 What shapes the continuum of
reproductive isolation? Lessons from Heliconius butterflies. *Proceedings of the Royal Society B:*
Biological Sciences **284**, 20170335. (doi:10.1098/rspb.2017.0335)

39. Seymoure BM, Mcmillan WO, Rutowski R. 2015 Peripheral eye dimensions in Longwing
(Heliconius) butterflies vary with body size and sex but not light environment nor mimicry ring.
The Journal of Research on the Lepidoptera **48**, 83–92. (doi:10.5962/p.266475)

40. Bernays EA, Glendinning JI, Chapman RF. 1998 Plant acids modulate chemosensory responses in
Manduca sexta larvae. *Physiological Entomology* **23**, 193–201. (doi:10.1046/j.1365-
3032.1998.233079.x)

41. Friard O, Gamba M. 2016 BORIS: a free, versatile open-source event-logging software for
video/audio coding and live observations. *Methods in Ecology and Evolution* **7**, 1325–1330.
(doi:10.1111/2041-210X.12584)

42. Damas-Moreira I, Oliveira D, Santos JL, Riley JL, Harris DJ, Whiting MJ. 2018 Learning from others:
an invasive lizard uses social information from both conspecifics and heterospecifics. *Biology*
Letters **14**, 20180532. (doi:10.1098/rsbl.2018.0532)

43. Batabyal A, Thaker M. 2019 Lizards from suburban areas learn faster to stay safe. *Biology Letters*
(doi:10.1098/rsbl.2019.0009)

44. Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting Linear Mixed-Effects Models Using lme4.
Journal of Statistical Software **67**, 1–48. (doi:10.18637/jss.v067.i01)

45. R Core Team. 2023 R: a Language and Environment for Statistical Computing.

46. Crawley MJ. 2013 *The R book*. Second edition. Chichester, West Sussex, UK: Wiley.

47. Lenth R, Singmann H, Love J, Buerkner P, Herve M. 2019 Package 'emmeans'.

48. Wickham H. 2011 ggplot2. *Wiley interdisciplinary reviews: computational statistics* **3**, 180–185.
(doi:10.1002/wics.147)

49. Jiggins CD, McMillan WO, Neukirchen W, Mallet J. 1996 What can hybrid zones tell us about
speciation? The case of Heliconius erato and H. himera (Lepidoptera: Nymphalidae). *Biological*
Journal of the Linnean Society **59**, 221–242. (doi:10.1111/j.1095-8312.1996.tb01464.x)

50. Rubene D, Low M, Brodin A. 2023 Birds differentially prioritize visual and olfactory foraging cues
depending on habitat of origin and sex. *Royal Society Open Science* **10**, 221336.
(doi:10.1098/rsos.221336)

51. Pike TW, Ramsey M, Wilkinson A. 2018 Environmentally induced changes to brain morphology
predict cognitive performance. *Philosophical Transactions of the Royal Society B: Biological*
Sciences **373**, 20170287. (doi:10.1098/rstb.2017.0287)

52. Keagy J, Braithwaite VA, Boughman JW. 2018 Brain differences in ecologically differentiated
sticklebacks. *Current Zoology* **64**, 243–250. (doi:10.1093/cz/zox074)

458
459

Supplementary material:

Supplementary Results

Among the 117 individuals (82 *H. melpomene* and 35 *H. cydno*) that passed our less stringent training criterion (i.e., >50% correct choices on day 5), *H. cydno* did not differ from *H. melpomene* in the trained test, indicating that both species were able to train to the multimodal stimulus (Table S1 post-hoc: *H. cydno* Trained vs. *H. melpomene* Trained, Z= -1.333, P= 1.0). However, *H. melpomene* and *H. cydno* responded differently when presented with the olfactory and visual stimuli in conflict (Figure S1). Importantly, the interaction between species and trial type was retained in our model, confirming that this was not due to existing differences amongst "trained" individuals, but rather the effect of how the stimuli were presented (species*treatment LRT: 2ΔlnL = 44.537, df=2, p < 0.001). Table S5). In the conflict test, *H. cydno* fed mostly on the blue feeder, while *H. melpomene* primarily chose the feeder with the rose scent, suggesting a difference between these species in how olfactory and visual cues are weighed during behavioural decisions (Table S1, post-hoc: *H. cydno* Conflict vs. *H. melpomene* Conflict, Z= 4.402, P< 0.001). We found no effect of sex, which was excluded from our model (sex LRT: 2ΔlnL = 0.297, df=1, p = 0.586. Table S5).

Post-hoc analysis revealed that individuals of both species made fewer feeding attempts on the blue feeder when the stimuli were presented in conflict, indicating that both species use both visual and olfactory stimuli when making foraging decisions (Table S1, post-hoc: *H. cydno* Trained vs. *H. cydno* Conflict, Z= 5.760, P= <.0001) (Table S1, post-hoc: *H. melpomene* Trained vs. *H. melpomene* Conflict, Z=23.222, P= <.0001). Notably, *H. cydno* did not differ from the trained test and the colour-only test indicating that this species largely relies on visual cues (Table S1, post-hoc: *H. cydno* Trained vs. *H. cydno* Conflict, Z= 0.337, P= 1.0). In contrast, *H. melpomene* differed between the trained test and the colour-only test further suggesting a difference in sensory weighting (Table 1, post-hoc: *H. melpomene* Trained vs. *H. melpomene* Conflict, Z= 8.433, P= <.0001).

Figure S 1: Our result that *H. cydno* and *H. melpomene* weight visual and olfactory cues differently when foraging is also supported when adopting a less stringent learning criteria. Proportion of feeding attempts towards blue feeder for *H. cydno* and *H. melpomene*. Each point represents the feeding choices of and individual; solid dots and error bars represent the mean choices per species ±95% confidence intervals. Blue circles represent *H. cydno* and orange triangles represent *H. melpomene*, shapes in grey represent naive individuals that did not train. On naive and trained tests, feeder combinations were blue/rose and red/lavender; In the conflict test, the positively trained colour (blue) was presented in conflict with the negatively rewarded odour (lavender) and *vice versa*. In the colour tests, individuals were presented only with red or blue feeders without an odour. Only individuals that passed our training criteria (>=50% on the trained day) are show (see methods).

490 **Table S 1:** Pairwise p-value comparison matrix of post-hoc test with Bonferroni correction for the minimal
491 adequate model for the effects of the interaction between species, treatment (Trained, Conflict, and Colour) and
492 their interaction on choices towards blue feeders. Only individuals that passed our training criteria (>=50% on the
493 trained day) are on this dataset. Results are from binomial GLMM model, see methods for statistical details.

	H. cydno Trained	*H. melpomene* Trained	*H. cydno* Conflict	*H. melpomene* Conflict	*H. cydno* Colour	*H. melpomene* Colour
***H. cydno* Trained**	[1.331]	1	<.0001	<.0001	1	0.338
***H. melpomene* Trained**		[1.656]	<.0001	<.0001	1	<.0001
***H. cydno* Conflict**			[0.360]	0.0002	<.0001	1
***H. melpomene* Conflict**				[-0.649]	<.0001	<.0001
***H. cydno* Colour**					[1.265]	0.7534
***H. melpomene* Colour**						[0.771]

Figure S2: Behavioural choices including individuals that did not reach our training criterion (>=75%). Proportion of feeding attempts towards blue feeder for *H. cydno* and *H. melpomene*. Each point represents the feeding choices of and individual; solid dots and error bars represent the mean choices per species ±95% confidence intervals. Blue circles represent *H. cydno* and orange triangles represent *H. melpomene*, shapes in grey represent individuals that did not train. Gray error bars represent the mean choices per species ±95% confidence intervals including all individuals.

496 **Table S 2:** Number of individuals by sex that participated in each treatment (Naïve, Trained, Conflict, Colour)
497 separated by number of individuals that passed the training criterion (>=50% or >=75%) of correct choices on the
498 trained day (day 5).

Species	Level	Sex	All individuals	>=50% training	>=75% training
H. cydno	Naive	Female	23	5	2
H. cydno	Naive	Male	38	8	6
H. cydno	Trained	Female	30	13	6
H. cydno	Trained	Male	49	22	15
H. cydno	Conflict	Female	35	11	5
H. cydno	Conflict	Male	56	20	14
H. cydno	Colour	Female	29	9	6
H. cydno	Colour	Male	48	18	12
H. melpomene	Naive	Female	23	14	10
H. melpomene	Naive	Male	51	34	15
H. melpomene	Trained	Female	40	30	24
H. melpomene	Trained	Male	68	52	28
H. melpomene	Conflict	Female	41	26	20
H. melpomene	Conflict	Male	68	48	28
H. melpomene	Colour	Female	35	21	16
H. melpomene	Colour	Male	61	45	26

Figure S3: Estimated marginal means (e.m.m.) of the minimum adequate statistical models for the proportion of feeding attempts towards blue feeder demonstrating the significant effects of the interaction between species (*H. cydno* and *H. melpomene*) and treatment [Trained test (day 5), Conflict test (day 6) or Colour test (day 7)]. Only individuals that passed our training criteria (>=75% on the trained day) are on this dataset. (see methods).

502   **Table S 3:** Model 1 minimal adequate model for the effects of species, treatment (Trained, Conflict, Colour) and
503   their interaction on choices towards blue feeders. Only individuals that passed our training criteria (>=75% on the
504   trained day) are on this dataset. Results are from a binomial GLMM model, see methods for statistical details.
505   Proportion blue ~ Species * Level + (1|Group/ID)

	2ΔlnL	df	Pr
Species	27.846	3	3.912e-06 ***
Treatment	681.740	4	< 2.2e-16 ***
Treatment:Species	22.876	2	1.078e-05 ***
Sex	3.067	1	0.07988

508 **Table S 4:** Pairwise comparison post-hoc test with Bonferroni correction for the minimal adequate model for the
509 effects of the interaction between species and treatment (Trained, Conflict, and Colour) on choices towards blue
510 feeders. Only individuals that passed our training criteria (>=75% on the trained day) are on this dataset. Results
511 are from binomial GLMM model, see methods for statistical details. Proportion blue ~ Species * Level +
512 (1|Group/ID)

	estimate	SE	z-ratio	p-value (Adjusted)
H. cydno Trained - *H. melpomene* Trained	-0.545	0.396	-1.375	1.000
H. cydno Trained - *H. cydno* Conflict	1.864	0.269	6.932	<.0001
H. cydno Trained - *H. melpomene* Conflict	2.930	0.381	7.696	<.0001
H. cydno Trained - *H. cydno* Colour	0.690	0.297	2.321	0.304
H. cydno Trained - *H. melpomene* Colour	1.340	0.387	3.464	0.008
H. melpomene Trained - *H. cydno* Conflict	2.409	0.351	6.860	<.0001
H. melpomene Trained - *H. melpomene* Conflict	3.475	0.169	20.614	<.0001
H. melpomene Trained - *H. cydno* Colour	1.235	0.367	3.361	0.012
H. melpomene Trained - *H. melpomene* Colour	1.884	0.173	10.920	<.0001
H. cydno Conflict - *H. melpomene* Conflict	1.066	0.332	3.210	0.020
H. cydno Conflict - *H. cydno* Colour	-1.174	0.237	-4.956	<.0001
H. cydno Conflict - *H. melpomene* Colour	-0.524	0.340	-1.544	1.000
H. melpomene Conflict - *H. cydno* Colour	-2.240	0.351	-6.385	<.0001
H. melpomene Conflict - *H. melpomene* Colour	-1.590	0.141	-11.244	<.0001
H. cydno Colour - *H. melpomene* Colour	0.650	0.357	1.819	1.000

514 **Table S 5:** Model 2 minimal adequate model for the effects of species, treatment (Trained, Conflict, Colour) and
515 their interaction on choices towards blue feeders. (>=50% on the trained day) are on this dataset. Results are
516 from binomial GLMM model, see methods for statistical details.
517 Proportion blue ~ Species * Level + (1|Group/ID)

	$2\Delta\ln L$	df	Pr
Species	50.004	3	7.972e-11 ***
Treatment	690.730	4	< 2.2e-16 ***
Treatment:Species	44.537	2	2.132e-10 ***
Sex	0.297	1	0.58550

519
520

Figure S4: Estimated marginal means (e.m.m.) of the minimum adequate statistical models for the proportion of feeding attempts towards blue feeder demonstrating the significant effects of the interaction between species (*H. cydno* and *H. melpomene)* and treatment [*Trained test* (day 5), *Conflict test* (day 6) or *Colour test* (day 7)]. Only individuals that passed our training criteria (>=50% on the trained day) are on this dataset. (see methods).

522 **Table S 6**: Pairwise comparison post-hoc test with Bonferroni correction for the minimal adequate model for the

523 effects of the interaction between species and treatment (Trained, Conflict, and Colour) on choices towards blue

524 feeders. Only individuals that passed our training criteria (>=50% on the trained day) are on this dataset. Results

525 are from binomial GLMM model, see methods for statistical details.

	estimate	SE	z-ratio	p-value
H. cydno Trained - *H. melpomene* Trained	-0.326	0.244	-1.333	1.000
H. cydno Trained - *H. cydno* Conflict	0.971	0.169	5.760	<.0001
H. cydno Trained - *H. melpomene* Conflict	1.980	0.241	8.199	<.0001
H. cydno Trained - *H. cydno* Colour	0.066	0.195	0.337	1.000
H. cydno Trained - *H. melpomene* Colour	0.560	0.245	2.281	0.338
H. melpomene Trained - *H. cydno* Conflict	1.296	0.233	5.571	<.0001
H. melpomene Trained - *H. melpomene* Conflict	2.305	0.099	23.222	<.0001
H. melpomene Trained - *H. cydno* Colour	0.391	0.251	1.556	1.000
H. melpomene Trained - *H. melpomene* Colour	0.885	0.105	8.433	<.0001
H. cydno Conflict - *H. melpomene* Conflict	1.009	0.229	4.402	0.000
H. cydno Conflict - *H. cydno* Colour	-0.905	0.183	-4.942	<.0001
H. cydno Conflict - *H. melpomene* Colour	-0.411	0.234	-1.760	1.000
H. melpomene Conflict - *H. cydno* Colour	-1.914	0.249	-7.697	<.0001
H. melpomene Conflict - *H. melpomene* Colour	-1.420	0.100	-14.263	<.0001
H. cydno Colour - *H. melpomene* Colour	0.494	0.252	1.958	0.753

528 **Table S7**: Average proportion of feeding attempts separated by species and treatment (Naïve, Trained, Conflict,
529 Colour) that passed the training criterion (>=50% or >=75%) of correct choices on the trained day (day 5).

Species	Level	Mean ≥50% training	95CI ≥50% training	Mean ≥75% training	95CI ≥75% training
H. cydno	Naive	0.252	(-0.001, 0.506)	0.159	(-0.111, 0.43)
H. cydno	Trained	0.806	(0.748, 0.864)	0.925	(0.885, 0.966)
H. cydno	Conflict	0.603	(0.49, 0.716)	0.638	(0.495, 0.782)
H. cydno	Colour	0.771	(0.65, 0.892)	0.858	(0.753, 0.964)
H. melpomene	Naive	0.227	(0.114, 0.339)	0.202	(0.049, 0.355)
H. melpomene	Trained	0.823	(0.785, 0.861)	0.938	(0.915, 0.961)
H. melpomene	Conflict	0.331	(0.267, 0.395)	0.359	(0.27, 0.448)
H. melpomene	Colour	0.728	(0.648, 0.809)	0.7845	(0.684, 0.885)

530 **Table S 8:** Model X minimal adequate model for the effects of species, treatment (Trained, Conflict, Colour) and
531 their interaction on the total number of feeding attempts. Only individuals that passed our training criteria (>=75%
532 on the trained day 5) are on this dataset. Results are from a binomial GLMM model, see methods for statistical
533 details. Total_feeding_attempts ~ Species * Level + (1|Group/ID)

	$2\Delta \ln L$	df	Pr
Species	6.9359	3	0.0738 n.s.
Treatment	8.3859	4	0.0784 n.s.
Treatment:Species	5.3948	2	0.06738 n.s.

Table S9: Total number of feeding attempts and proportion of feeding attempts towards blue feeders grouped by individual. Only individuals that passed our training criteria (>=75% on the trained day5) are on this dataset.

ID	Species	Sex	Level	Day	Red	Blue	Total	Prop_blue	Group
C22_004	MMC	M	Naive	1	0	3	3	1	G14
C22_107	CC	F	Naive	1	1	0	1	0	G15
C22_091	MMC	M	Naive	1	2	0	2	0	G16
C22_173	MMC	F	Naive	1	0	1	1	1	G17
C22_226	MMC	F	Naive	1	3	0	3	0	G17
C22_273	MMC	F	Naive	1	2	0	2	0	G17
C22_028	MMC	M	Naive	1	9	0	9	0	G18
C22_433	MMC	F	Naive	1	1	0	1	0	G20
C22_590	MMC	F	Naive	1	1	0	1	0	G21
C22_831	MMC	F	Naive	1	1	0	1	0	G21
C22_898	MMC	F	Naive	1	12	0	12	0	G21
C22_918	MMC	F	Naive	1	8	0	8	0	G21
C22_555	MMC	M	Naive	1	7	0	7	0	G22
C22_699	CC	M	Naive	1	4	31	35	0	G22
A23_055	MMC	M	Naive	1	23	18	41	0	G22
A23_094	MMC	M	Naive	1	1	0	1	0	G23
C22_704	MMC	M	Naive	1	14	19	33	0	G23
A23_280	CC	F	Naive	1	1	0	1	0	G25
A23_054	CC	M	Naive	1	14	9	23	0	G25
A23_242	MMC	M	Naive	1	5	1	6	0	G25
A23_306	MMC	M	Naive	1	1	11	12	0	G25
C22_911	MMC	M	Naive	1	7	0	7	0	G26
C22_935	CC	M	Naive	1	1	0	1	0	G26
C22_936	CC	M	Naive	1	6	0	6	0	G26
A23_098	MMC	M	Naive	1	1	0	1	0	G27
A23_148	MMC	M	Naive	1	1	0	1	0	G27
A23_182	MMC	M	Naive	1	1	0	1	0	G27
A23_258	MMC	M	Naive	1	1	24	25	0	G25
A23_259	MMC	M	Naive	1	3	0	3	0	G27
A23_303	CC	M	Naive	1	5	0	5	0	G27
A23_428	CC	M	Naive	1	6	0	6	0	G27
A23_224	MMC	F	Naive	1	2	0	2	0	G28
A23_283	MMC	F	Naive	1	1	0	1	0	G28
C22_004	MMC	M	Trained	5	0	16	16	1	G14
C22_063	MMC	F	Trained	5	13	52	65	0	G25
C22_107	CC	F	Trained	5	0	2	2	1	G15
C22_134	MMC	F	Trained	5	0	1	1	1	G15
C22_025	CC	M	Trained	5	0	1	1	1	G16
C22_032	MMC	M	Trained	5	0	6	6	1	G16

C22_091	MMC	M	Trained	5	0	26	26	1	G16
C22_173	MMC	F	Trained	5	0	1	1	1	G17
C22_198	CC	F	Trained	5	3	17	20	0	G17
C22_226	MMC	F	Trained	5	2	22	24	0	G17
C22_265	MMC	F	Trained	5	0	20	20	1	G17
C22_273	MMC	F	Trained	5	0	9	9	1	G17
C22_285	MMC	F	Trained	5	0	49	49	1	G17
C22_028	MMC	M	Trained	5	0	1	1	1	G18
C22_242	CC	M	Trained	5	2	7	9	0	G25
C22_227	MMC	M	Trained	5	3	11	14	0	G25
C22_290	CC	M	Trained	5	0	6	6	1	G19
C22_341	MMC	M	Trained	5	12	36	48	0	G19
C22_367	MMC	M	Trained	5	2	12	14	0	G19
C22_448	MMC	M	Trained	5	4	39	43	0	G19
C22_186	MMC	F	Trained	5	1	4	5	0	G19
C22_230	MMC	F	Trained	5	2	14	16	0	G19
C22_422	MMC	F	Trained	5	2	24	26	0	G19
C22_433	MMC	F	Trained	5	0	46	46	1	G20
C22_590	MMC	F	Trained	5	0	11	11	1	G21
C22_831	MMC	F	Trained	5	0	8	8	1	G21
C22_898	MMC	F	Trained	5	2	16	18	0	G21
C22_918	MMC	F	Trained	5	0	26	26	1	G21
C22_555	MMC	M	Trained	5	0	4	4	1	G22
C22_699	CC	M	Trained	5	0	5	5	1	G22
C22_756	MMC	M	Trained	5	0	1	1	1	G22
A23_055	MMC	M	Trained	5	0	4	4	1	G23
A23_094	MMC	M	Trained	5	2	7	9	0	G23
C22_704	MMC	M	Trained	5	0	7	7	1	G23
C22_848	MMC	M	Trained	5	3	20	23	0	G23
A23_264	MMC	F	Trained	5	0	3	3	1	G25
A23_280	CC	F	Trained	5	2	6	8	0	G25
A23_030	MMC	M	Trained	5	0	1	1	1	G26
A23_054	CC	M	Trained	5	2	7	9	0	G26
A23_185	MMC	M	Trained	5	0	4	4	1	G26
A23_242	MMC	M	Trained	5	1	13	14	0	G26
A23_301	CC	M	Trained	5	0	7	7	1	G26
A23_306	MMC	M	Trained	5	0	3	3	1	G26
A23_319	CC	M	Trained	5	0	5	5	1	G26
C22_843	CC	M	Trained	5	0	4	4	1	G26
C22_911	MMC	M	Trained	5	5	16	21	0	G26
C22_935	CC	M	Trained	5	0	1	1	1	G26
C22_936	CC	M	Trained	5	3	23	26	0	G26
C22_949	CC	M	Trained	5	0	2	2	1	G26

A23_056	MMC	M	Trained	5	5	17	22	0	G26
A23_083	CC	M	Trained	5	0	18	18	1	G27
A23_098	MMC	M	Trained	5	1	19	20	0	G27
A23_148	MMC	M	Trained	5	1	21	22	0	G27
A23_182	MMC	M	Trained	5	4	33	37	0	G27
A23_258	MMC	M	Trained	5	0	2	2	1	G27
A23_259	MMC	M	Trained	5	1	5	6	0	G27
A23_268	MMC	F	Trained	5	0	11	11	1	G27
A23_289	MMC	M	Trained	5	0	19	19	1	G27
A23_303	CC	M	Trained	5	1	6	7	0	G27
A23_345	MMC	M	Trained	5	0	8	8	1	G27
A23_347	MMC	M	Trained	5	1	10	11	0	G27
A23_421	CC	M	Trained	5	2	17	19	0	G27
A23_428	CC	M	Trained	5	2	16	18	0	G27
A23_009	CC	F	Trained	5	0	5	5	1	G28
A23_027	CC	F	Trained	5	1	5	6	0	G28
A23_209	CC	F	Trained	5	5	55	60	0	G28
A23_221	MMC	F	Trained	5	0	10	10	1	G28
A23_224	MMC	F	Trained	5	0	33	33	1	G28
A23_283	MMC	F	Trained	5	0	4	4	1	G28
C22_809	MMC	F	Trained	5	5	21	26	0	G28
C22_891	MMC	F	Trained	5	0	8	8	1	G28
C22_900	MMC	F	Trained	5	3	28	31	0	G28
C22_920	MMC	F	Trained	5	3	28	31	0	G28
C22_004	MMC	M	Conflict	6	1	20	21	0	G28
C22_063	MMC	F	Conflict	6	22	11	33	0	G28
C22_134	MMC	F	Conflict	6	5	0	5	0	G15
C22_025	CC	M	Conflict	6	5	0	5	0	G16
C22_032	MMC	M	Conflict	6	2	22	24	0	G16
C22_091	MMC	M	Conflict	6	5	8	13	0	G16
C22_198	CC	F	Conflict	6	14	8	22	0	G16
C22_226	MMC	F	Conflict	6	15	3	18	0	G16
C22_265	MMC	F	Conflict	6	39	15	54	0	G16
C22_273	MMC	F	Conflict	6	8	0	8	0	G17
C22_285	MMC	F	Conflict	6	21	9	30	0	G17
C22_028	MMC	M	Conflict	6	30	4	34	0	G17
C22_242	CC	M	Conflict	6	0	21	21	1	G18
C22_227	MMC	M	Conflict	6	5	0	5	0	G19
C22_341	MMC	M	Conflict	6	4	2	6	0	G19
C22_367	MMC	M	Conflict	6	19	1	20	0	G19
C22_448	MMC	M	Conflict	6	5	1	6	0	G19
C22_422	MMC	F	Conflict	6	10	1	11	0	G19
C22_433	MMC	F	Conflict	6	10	2	12	0	G19

C22_590	MMC	F	Conflict	6	17	12	29	0	G19
C22_831	MMC	F	Conflict	6	0	3	3	1	G21
C22_898	MMC	F	Conflict	6	34	3	37	0	G21
C22_555	MMC	M	Conflict	6	3	11	14	0	G21
C22_699	CC	M	Conflict	6	1	3	4	0	G21
C22_756	MMC	M	Conflict	6	0	13	13	1	G22
A23_055	MMC	M	Conflict	6	5	4	9	0	G22
A23_094	MMC	M	Conflict	6	20	0	20	0	G23
C22_704	MMC	M	Conflict	6	10	28	38	0	G23
C22_848	MMC	M	Conflict	6	11	3	14	0	G23
A23_264	MMC	F	Conflict	6	1	0	1	0	G25
A23_280	CC	F	Conflict	6	7	12	19	0	G25
A23_030	MMC	M	Conflict	6	10	2	12	0	G25
A23_054	CC	M	Conflict	6	2	25	27	0	G25
A23_185	MMC	M	Conflict	6	5	10	15	0	G25
A23_242	MMC	M	Conflict	6	4	27	31	0	G25
A23_301	CC	M	Conflict	6	9	18	27	0	G25
A23_306	MMC	M	Conflict	6	19	18	37	0	G25
A23_319	CC	M	Conflict	6	0	8	8	1	G26
C22_843	CC	M	Conflict	6	3	8	11	0	G26
C22_911	MMC	M	Conflict	6	31	20	51	0	G26
C22_935	CC	M	Conflict	6	2	5	7	0	G26
C22_936	CC	M	Conflict	6	7	25	32	0	G26
C22_949	CC	M	Conflict	6	4	5	9	0	G26
A23_056	MMC	M	Conflict	6	17	8	25	0	G26
A23_083	CC	M	Conflict	6	14	4	18	0	G26
A23_098	MMC	M	Conflict	6	22	3	25	0	G26
A23_148	MMC	M	Conflict	6	9	2	11	0	G26
A23_182	MMC	M	Conflict	6	28	7	35	0	G26
A23_258	MMC	M	Conflict	6	8	6	14	0	G26
A23_259	MMC	M	Conflict	6	2	2	4	0	G26
A23_268	MMC	F	Conflict	6	7	0	7	0	G27
A23_289	MMC	M	Conflict	6	10	8	18	0	G27
A23_303	CC	M	Conflict	6	6	7	13	0	G27
A23_345	MMC	M	Conflict	6	2	1	3	0	G27
A23_347	MMC	M	Conflict	6	5	4	9	0	G27
A23_421	CC	M	Conflict	6	15	2	17	0	G27
A23_428	CC	M	Conflict	6	6	5	11	0	G27
A23_009	CC	F	Conflict	6	0	2	2	1	G28
A23_027	CC	F	Conflict	6	1	3	4	0	G28
A23_209	CC	F	Conflict	6	2	29	31	0	G28
A23_221	MMC	F	Conflict	6	12	0	12	0	G28
A23_224	MMC	F	Conflict	6	7	6	13	0	G28

A23_283	MMC	F	Conflict	6	19	11	30	0	G28
C22_809	MMC	F	Conflict	6	1	1	2	0	G28
C22_891	MMC	F	Conflict	6	0	2	2	1	G28
C22_900	MMC	F	Conflict	6	6	0	6	0	G28
C22_920	MMC	F	Conflict	6	9	2	11	0	G28
C22_004	MMC	M	Colour	Day7	10	18	28	0	G28
C22_063	MMC	F	Colour	Day7	22	16	38	0	G28
C22_107	CC	F	Colour	Day7	13	3	16	0	G28
C22_134	MMC	F	Colour	Day7	0	14	14	1	G15
C22_032	MMC	M	Colour	Day7	0	5	5	1	G16
C22_091	MMC	M	Colour	Day7	0	14	14	1	G16
C22_173	MMC	F	Colour	Day7	8	0	8	0	G17
C22_198	CC	F	Colour	Day7	0	17	17	1	G17
C22_226	MMC	F	Colour	Day7	0	1	1	1	G17
C22_265	MMC	F	Colour	Day7	2	11	13	0	G17
C22_285	MMC	F	Colour	Day7	9	7	16	0	G17
C22_227	MMC	M	Colour	Day7	7	5	12	0	G17
C22_290	CC	M	Colour	Day7	4	15	19	0	G17
C22_341	MMC	M	Colour	Day7	1	5	6	0	G17
C22_367	MMC	M	Colour	Day7	24	1	25	0	G17
C22_448	MMC	M	Colour	Day7	7	14	21	0	G17
C22_590	MMC	F	Colour	Day7	0	6	6	1	G21
C22_831	MMC	F	Colour	Day7	0	4	4	1	G21
C22_898	MMC	F	Colour	Day7	25	1	26	0	G21
C22_555	MMC	M	Colour	Day7	3	19	22	0	G21
C22_699	CC	M	Colour	Day7	2	3	5	0	G21
C22_756	MMC	M	Colour	Day7	3	0	3	0	G22
A23_055	MMC	M	Colour	Day7	0	7	7	1	G23
A23_094	MMC	M	Colour	Day7	0	7	7	1	G23
C22_704	MMC	M	Colour	Day7	0	9	9	1	G23
C22_848	MMC	M	Colour	Day7	0	7	7	1	G23
A23_280	CC	F	Colour	Day7	4	14	18	0	G23
A23_030	MMC	M	Colour	Day7	10	10	20	0	G23
A23_054	CC	M	Colour	Day7	0	29	29	1	G26
A23_185	MMC	M	Colour	Day7	4	12	16	0	G26
A23_242	MMC	M	Colour	Day7	0	47	47	1	G26
A23_301	CC	M	Colour	Day7	0	14	14	1	G26
A23_306	MMC	M	Colour	Day7	21	18	39	0	G26
A23_319	CC	M	Colour	Day7	0	10	10	1	G26
C22_843	CC	M	Colour	Day7	0	9	9	1	G26
C22_911	MMC	M	Colour	Day7	27	13	40	0	G26
C22_935	CC	M	Colour	Day7	0	6	6	1	G26
C22_936	CC	M	Colour	Day7	9	21	30	0	G26

C22_949	CC	M	Colour	Day7	0	7	7	1	G26
A23_056	MMC	M	Colour	Day7	0	10	10	1	G27
A23_083	CC	M	Colour	Day7	0	10	10	1	G27
A23_098	MMC	M	Colour	Day7	0	20	20	1	G27
A23_148	MMC	M	Colour	Day7	0	3	3	1	G27
A23_182	MMC	M	Colour	Day7	0	10	10	1	G27
A23_258	MMC	M	Colour	Day7	0	5	5	1	G27
A23_259	MMC	M	Colour	Day7	0	8	8	1	G27
A23_268	MMC	F	Colour	Day7	0	2	2	1	G27
A23_289	MMC	M	Colour	Day7	0	14	14	1	G27
A23_303	CC	M	Colour	Day7	0	4	4	1	G27
A23_345	MMC	M	Colour	Day7	0	3	3	1	G27
A23_421	CC	M	Colour	Day7	0	6	6	1	G27
A23_009	CC	F	Colour	Day7	1	5	6	0	G27
A23_027	CC	F	Colour	Day7	4	14	18	0	G27
A23_209	CC	F	Colour	Day7	9	34	43	0	G27
A23_221	MMC	F	Colour	Day7	0	3	3	1	G28
A23_224	MMC	F	Colour	Day7	2	10	12	0	G28
A23_283	MMC	F	Colour	Day7	0	5	5	1	G28
C22_809	MMC	F	Colour	Day7	1	7	8	0	G28
C22_900	MMC	F	Colour	Day7	0	5	5	1	G28
C22_920	MMC	F	Colour	Day7	0	5	5	1	G28

539