1   # Sensory weighting reflects changing patterns of visual investment during
2   # ecological divergence in *Heliconius* butterflies.
3

4   José Borrero[1], Elisa Mogollon Perez[1], Daniel Shane Wright[1], Daniela Lozano[1,2], Geraldine

5   Rueda-Muñoz[1,2], Carolina Pardo-Diaz[2], Camilo Salazar[2], Stephen H. Montgomery[3]* &

6   Richard M. Merrill[1]*

7   Corresponding authors: jose.borrero@lmu.de (J.B.) & merrill@bio.lmu.de (R.M.M)

8
9   [1] *Division of Evolutionary Biology, Faculty of Biology, LMU Munich, Germany*
10   [2] *Faculty of Natural Sciences, Universidad del Rosario; Bogotá, Colombia*
11   [3] *School of Biological Science, University of Bristol, Bristol, United Kingdom*

12   ## Abstract
13

Integrating information across sensory modalities enables animals to orchestrate a wide

range of complex behaviours. The relative importance placed on one sensory modality over

another reflects the reliability of cues in a particular environment and corresponding

differences in neural investment. As populations diverge across environmental gradients, the

reliability of sensory cues may shift, favouring divergence in neural investment and sensory

weighting. During their divergence across closed forest and forest-edge habitats, *Heliconius*

butterflies *H. cydno* and *H. melpomene* evolved distinct brain morphologies, with the former

investing more in vision. Molecular and anatomical data suggest selection drove these

changes, but their behavioural effects remain uncertain. We hypothesised that divergent

investment in neuropils may alter sensory weighting during behavioural tasks. To address

this, we trained individuals in an associative learning experiment using multimodal colour

and odour cues. When positively rewarded stimuli were presented in conflict pairing

positively trained colour with negatively trained odour, and *vice-versa*, *H. cydno* prioritised

visual cues more strongly than *H. melpomene*. Hence, differences in sensory weighting may

evolve early during divergence and are potentially predicted by patterns of neural

investment. These findings, alongside other ecological divergence examples, imply that

differences in sensory weighting stem from sensory investment as adaptations to local

sensory environments.

Introduction

Animals rely on multiple sensory systems to perceive and navigate their environments. The integration of environmental information across different sensory modalities (e.g., vision and olfaction) facilitates complex behaviours such as foraging, predator avoidance, and mate finding [1,2]. Cues from different sensory modalities can work in concert to enhance signal detection [3], but individuals may also prioritise one sensory modality over another [1,4]. The relative importance animals place on one sensory modality over another is likely shaped by natural selection [5], leading to evolutionary shifts in how different sensory modalities are weighted between populations that exploit different habitats.

Evidence for divergence in sensory weighting within and between species exploiting different sensory environments has been observed across a handful of taxa. For instance, nine-spined sticklebacks show plasticity in their sensory reliance, with individuals raised in visually restricted environments displaying a larger reliance on chemical cues over visual cues, compared to fish raised in visually unrestricted habitats [6]. Stingless bees rely less on visual cues, such as colour, than honeybees, possibly due to reduced investment in eye size [7]. In butterflies and moths, brain areas processing sensory information are among the most variable structure between species, and studies investigating multisensory integration emphasise associations between interspecific differences in neuroanatomy and behaviour [8,9]. For example, while hawkmoths broadly use both visual and olfactory cues during foraging [4,10], nocturnal species invest relatively more in brain regions associated with olfactory processing, whereas diurnal species are characterised by larger investments in visual neuropils. This shift in neural investment is mirrored in the relative importance given to olfactory and visual cues during foraging [11]. However, how sensory weighting evolves in smaller time scales and across more subtle ecological gradients at the early stages of divergence is less clear.

The Neotropical *Heliconius* butterflies provide an excellent opportunity to study how sensory environments can shape multisensory integration during the early stages of divergence. Although olfaction also plays a role [12–17], these butterflies rely heavily on vision, including during foraging [18,19], mate recognition [20–22] and oviposition [23,24]. In *Heliconius*, speciation is often associated with ecological transitions across forest types [25–28], such that closely related species experience different sensory conditions [26,29,30].

67 Sister taxa are also known to differ in multiple aspects of both the peripheral and central
68 sensory pathways, likely reflecting adaptation to local sensory conditions [31–33].
69
If these shifts in neural investment and sensory weighting are the result of divergent
selection imposed by differences in sensory ecology, we expect populations diverging
across similar environmental gradients to show associated changes in behaviour [34–36].
The *Heliconius* species pair, *H. melpomene* and *H. cydno,* are a well-studied case of
ecological speciation which provide an opportunity to test this hypothesis. These species
exist in "mosaic sympatry", where *H. melpomene* inhabits open forest edges and *H. cydno*
occupies more closed canopy, inner forest habitats (Figure 1) [26,37,38]. *H. cydno* has larger
eyes and invests more heavily in brain structures associated with vision when compared to
H. melpomene, with multiple sources of data strongly suggesting these differences have
been shaped by adaptive processes [32,33,39]. In *Heliconius*, visual information is relayed
from the optic lobes to the mushroom bodies, major sensory integration centres, which also
receives input from the olfactory pathway [9]. Changes in investment in specific sensory
pathways may therefore affect the strength of input received, consequently influencing the
weight given to different sensory cues during behavioural decision-making processes.
84
85 Here, we test how *H. cydno* and *H. melpomene* prioritise visual and olfactory information
86 during foraging, predicting that increased investment in visual pathways may lead to *H.*
87 *cydno* prioritising these cues. To this end, we trained butterflies to a particular combination of
88 visual and olfactory stimuli using flower models, and then quantified the relative weights of
89 visual and olfactory cues by presenting the rewarded stimuli in conflict.

Materials & Methods
<u>Butterfly rearing and maintenance</u>
We maintained outbred stock populations of *H. cydno* and *H. melpomene* at the
Experimental Station José Celestino Mutis – Universidad del Rosario, near La Vega,
Colombia (5.0005° N, 74.3394° W), established from wild individuals caught in the local
area. Butterflies were kept in groups in outdoor 1x3x2 m insectary cages with access to
~20% sugar solution*, Gurania, Lantana* and *Psiguria spp.* flowers as pollen sources, and
Passiflora vines for oviposition. All individuals used in the experiments were reared under
common garden conditions.
99
<u>Experimental set-up and stimuli</u>
To test for visual and olfactory preferences, learning and sensory weighting, we presented
male and female butterflies with artificial feeders with both visual (red or blue) and olfactory

(lavender or rose essential oils) stimuli, on two 25x25 cm hard PVC sheet 'arrays' separated

by 200 cm in an 1x3x2m insectary (Figure S1A). Each array had four feeders made from a 5

ml Eppendorf tube surrounded by a 3 cm radius circle printed on waterproof paper. On one

array all the paper circles were red, and on the other all paper circles were blue. Olfactory

stimuli were presented in a 200µl PCR-tube below the paper circles, containing either

lavender (*Lavandula angustifolia*) or rose *(Rosa damascene)* essential oils diluted with

paraffin oil to 1% concentration (Figure S1B). By keeping the olfactory cues separate

underneath waterproof paper and avoiding their mixing with the feeding solution, this

ensured that olfactory associations were not confounded by gustatory reception. All four

feeders on the same array had the same olfactory stimuli to avoid odour mixing.

We trained and tested butterflies in same sex groups of 20-24 uniquely marked individuals

(10-12 per species). Butterflies were introduced to the experimental cages at least 12 hours

prior to the initial testing period to allow for acclimatisation and were food deprived overnight.

We trained butterflies to associate the combination of blue colour and rose odour with a

positive reward [blue/rose] (20% sugar-water solution) and the combination of [red/lavender]

with a negative stimulus (~18% Vitamin C solution; [17,40]). Butterflies were tested with the

visual and olfactory in the *trained* test (i.e. [blue/rose] vs [red/lavender]: days 1 and 5), in

'*conflict'* (i.e. [blue/lavender] vs [red/rose]: day 6), or with *colour alone* (i.e. [blue] vs [red]:

day 7) (Figure 2A). During testing, clean Eppendorf tubes were filled solely with water (i.e.,

no positive or negative 'reward') and fresh visual stimuli were used.

125   <u>Training and testing</u>

i) Naive test (Day 1): We recorded the naive preferences toward combinations of the visual

and olfactory stimuli [blue/rose] vs [red/lavender] for one hour (between 10:00-11:00). Each

butterfly was then hand fed once on a positive [blue/rose (+)], and once on a negative

rewarded feeder [red/lavender] (-)], and then allowed to feed freely for 2 hours from the

positive and negative rewarded feeders (11:00-13:00).

ii)Training (Days 2-4): Training was reinforced on experimental days 2-4. First, each

individual was hand fed once on a positive [blue/rose (+)], and once on a negative rewarded

feeder [red/lavender] (-)]. Afterwards, butterflies were allowed to feed freely for 3 hours

(10:00-13:00) from the positive or negative rewarded feeders. The feeders' position was

randomised (coin toss) every day to prevent positional learning.

iii) Trained test (Day 5*)*: On day 5, we recorded the trained preferences for one hour

(between 10:00-11:00) using only water in the feeders with the stimuli in *trained* conditions

(i.e. [blue/rose] vs [red/lavender]). This was immediately followed by two hours of

reinforcement training with rewarded feeders, i.e. [blue/rose (+)] and [red/lavender (-)].

iv) Conflict test (Day 6*)*: To test for differences in sensory weighting, we presented the visual

and olfactory in conflict (i.e. [blue/lavender] vs [red/rose]; Figure 2A). As for the previous

tests, feeding behaviour was recorded for one hour (between 10:00-11:00) using only water

in the feeders. This was then followed by reinforcement training for two hours, i.e. [blue/rose

(+)] and [red/lavender (-)].

v) Colour test (Day 7*):* We tested visual preferences in isolation for one hour (between

10:00-11:00) (i.e., blue vs red, without olfactory stimuli, and using water only).

152   <u>Behavioural tests</u>

All feeding test assays [(i) Naive, (ii) Trained, (iii) Conflict and (iv) Colour] were filmed from

above with an AKASO Brave 4 Pro action camera. We scored the videos in BORIS [41],

recording the total number of feeding attempts directed toward each stimulus per individual.

Feeding attempts were defined as a proboscis extension within the coloured circle feeders.

We scored instances where an individual moved outside the circle and subsequently

returned as a new feeding attempt. In contrast, we scored instances where an individual

remained within a circle and probed continuously as a single attempt, unless the proboscis

was retracted for 3 seconds or longer.

162   <u>Training criterion</u>

We established a stringent training criterion to incorporate only individuals that had

successfully learned the correct stimulus combination, following similar learning studies in

lizards [42,43] and butterflies [19]. Out of 260 individuals, 73 did not attempt to feed on the

trained test (day 5) and were subsequently excluded from further analyses (See Table S2 for

sample sizes by different training criterion). To investigate species differences in sensory

weighting, we included only those individuals that demonstrated robust associative learning

of the trained cue combinations (i.e. >=75% of feeding choices towards the positive

multimodal stimuli [blue/rose] on the trained test (day 5)), resulting in a reduced sample size

from 187 to 73 individuals. Although biased towards *H. melpomene* (52 individuals vs 21 *H.*

cydno individuals), this rigorous criterion ensured that our analysis focused solely on

individuals who had successfully learned. However, our results did not quantitatively change

when including all 117 individuals with a lower threshold of learning (i.e. >50% correct; see

supplementary results Table S1 and Tables S5-S6).

<u>Statistical analysis</u>

We used general linear mixed models (GLMM) with a binomial distribution in the *lme4* [44] in

R [45] to assess if species differed in their feeding attempts when the positively rewarded

stimuli agreed or were in conflict. The dependent variable was the proportion of choices

toward the blue feeders, species, and treatment [*Trained test* (day 5), *Conflict test* (day 6) or

Colour test (day 7)], and their interaction, were included as fixed factors. Individual ID nested

within group ID were included as random effects to account for repeated measures (Table

S1). We applied a stepwise model reduction approach to identify the minimum adequate

statistical models, using likelihood ratio tests (LRT) via the *drop1* function to determine

significance. When fixed effects or their interactions were significant, we conducted pairwise

comparisons tests using post hoc Bonferroni corrections with the *emmeans* package [46].

Plots were made using *ggplot2* [47].

Results

Among the 73 individuals (52 *H. melpomene* and 21 *H. cydno*) that passed our training criterion (i.e., >75% correct choices on day 5), *H. cydno* did not differ from *H. melpomene* in the trained test (Table 1 post-hoc: *H. cydno* Trained vs. *H. melpomene* Trained, Z= -1.375, P= 1.0). In contrast, *H. melpomene* and *H. cydno* respond differently when presented with the olfactory and visual stimuli in conflict (Figure 2B). Importantly, the interaction between species and trial type was retained in our model, confirming that this was an effect of how the stimuli were presented (species*treatment LRT: $2\Delta lnL = 22.876$, df=2, p < 0.001. Table S3). In the conflict test, *H. cydno* fed mostly on the blue feeder, while *H. melpomene* primarily chose the feeder with the rose scent, suggesting a difference in sensory weighting between these species (Table 1, post-hoc: *H. cydno* Conflict vs. *H. melpomene* Conflict, Z= 3.210, P= 0.019). We found no effect of sex, which was excluded from our model (species*trial type LRT: $2\Delta lnL = 3.067$, df=1, p = 0.079. Table S3).

Our post-hoc analysis revealed that individuals of both species made fewer feeding attempts on the blue feeder when the stimuli were presented in conflict, suggesting that both species integrate visual and olfactory stimuli when making foraging decisions (Table 1, post-hoc: *H. cydno* Trained vs. *H. cydno* Conflict, Z= 6.932, P= <.0001) (Table 1, post-hoc: *H. melpomene* Trained vs. *H. melpomene* Conflict, Z=20.614, P= <.0001). *H. cydno* made fewer feeding attempts on the blue feeder when the positively rewarded olfactory and visual stimuli were in conflict than when they were in the trained combinations confirming the use of both cues when present. However, there was no distinction between the trained test and the colour-only test (Table 1, post-hoc: *H. cydno* Trained vs. *H. cydno* Colour, Z= 2.321, P= 0.304), further emphasising that *H. cydno* has heavier reliance on the visual cue in our assay. Conversely, *H. melpomene* did display differences between the trained test and the colour-only test, further indicating a difference in sensory weighting and greater attention to the olfactory stimulus (Table 1, post-hoc: *H. melpomene* Trained vs. *H. melpomene* Colour, Z= 10.920, P= <.0001).

Discussion

We explored how two closely related butterfly species, *H. cydno* and *H. melpomene*, weigh visual and olfactory information in a foraging context. Our findings indicate a significant divergence in behaviour between these two species, with *H. cydno* placing a greater emphasis on visual cues over olfactory cues compared to *H. melpomene.* Importantly, our results mirror differences in brain morphology and suggest that divergence in the sensory

environment may have led to changes in brain composition and how these species process,

integrate, and prioritise information across sensory modalities.

To assess how *H. cydno* and *H. melpomene* prioritise visual and olfactory information, we

presented trained butterflies with conflicting stimuli - pairing positively trained colours with

negatively trained odours, and *vice versa*. Both species chose the 'correct' colour less

frequently in these conflict scenarios compared to their trained preferences or when

presented with colours alone, indicating that both visual and olfactory cues influence

foraging decisions . However, when forced to choose between conflicting cues, the species

differed in their reliance on each sensory modality. *H. cydno* predominantly fed on the

positively rewarded visual cue (the blue feeder), suggesting they prioritise vision. In contrast,

H. melpomene, showed a stronger preference for the feeder containing the positive trained

odour stimulus (rose), indicating a difference in sensory weighting between the two species.

Additionally, *H. melpomene* differed between the trained and colour only test, making fewer

correct choices when only colours were presented, again suggesting a larger reliance on the

olfaction stimulus compared to *H. cydno.* As the visual and olfactory cues employed in our

study are assumed to not be ecologically relevant, we suggest they reflect the baseline

relative preferences of the species for each sensory modality. Given that the behavioural

variation we observed was among common-garden-reared individuals, these interspecific

shifts are likely to have a genetic basis. In addition, our results align with divergent patterns

of investment in sensory pathways between the two species, whereby *H. cydno* exhibits

greater relative investment in the visual components of the sensory systems [32,33,39].

Therefore, heritable shifts in neural investment and sensory weighting may result from

divergent selection imposed by differences in sensory ecology.

One potential limitation of our study is that we managed to successfully train a larger number

of *H. melpomene* than *H. cydno* individuals. This discrepancy in training could be due to *H.*

melpomene being more active during the trials, possibly due to the outdoor insectaries being

found in the forest edge, which closely resembles *H. melpomene's* habitat [26,30]. Indeed, in

contrast to *H. cydno, H. melpomene* is frequently observed flying next to the insectaries

(pers. obs. J. Borrero). Despite this difference, individuals of both species successfully

learned the rewarded combination under our rigorous training criteria (>= 75% correct

choices) and our main conclusions hold even under less rigorous training criteria (See

supplementary results).

Adaptation to the local sensory environment may play an important role during the early

stages of divergence. The environmental gradient observed in our studied species, *H. cydno*

and *H. melpomene*, from closed forest canopy to open forest edge [26,30], resembles the

transition across an altitudinal gradient, from complex broad-leaf forest environments to dry

Andean forests seen in another distantly related *Heliconius* species-pair, *H. erato* and *H.*

himera [48]. *H. erato cyrbia* and *H. himera* exhibit similar patterns of neural divergence,with

H. erato cyrbia showing increased, heritable investment in the visual pathway, but with *H.*

himera also showing some increased investment in the olfactory pathway [17,31]. Despite

methodological differences (group vs individual testing and the exact visual and olfactory

stimuli used), our results qualitatively reassemble this previous study on sensory weighting

in *H. erato* and *H. himera.* Together these findings demonstrate repeated shifts in sensory

weighting among *Heliconius* butterflies, with species inhabiting relatively more closed

habitats investing more in perceiving and processing visual information, and preferentially

weighting visual over olfactory cues. This suggests changes in the light environment is a

common source of divergent selection across the two species pairs.

More broadly, changes in sensory weighting have been observed across different taxa and

across different evolutionary timescales. For instance, urban Great tits prefer olfactory cues,

while forest birds rely more on visual cues [49]. In threespine stickleback fish, populations in

different lakes segregate into limnetic and benthic ecomorphs along an environmental

gradient [50]. These groups show repeated evolutionary shifts in brain regions, with limnetic

species having larger structures for visual processing and benthic ectomorphs possessing

larger olfactory bulbs, and are correlated with differences in their sensory environments [50].

By altering the visual environment of nine-spined sticklebacks, researchers demonstrated

plastic changes in the size of brain regions involved in processing visual and chemical

information, which altered reliance on chemical over visual cues than fish housed in visually

unrestricted conditions [6]. This potentially suggests neural plasticity may underpin initial

behavioural responses to novel environments that are later consolidated by selection.

Comparing across larger evolutionary timescales, studies have found that nocturnal

hawkmoths prioritise olfactory cues while foraging, as evidenced by their correlated

increased investment in olfactory processing. Conversely, diurnal hawkmoths show larger

investments in visual neuropils and place a greater emphasis on visual cues. These changes

in sensory weighting and brain investment are likely driven by the transition from a nocturnal

to a diurnal habitat and the associated alterations in their sensory environment [11].

Combined with our findings, these studies underscore the role of environmental factors in

driving changes in neuroanatomy, sensory integration, and behaviour across species.

In conclusion, our results highlight repeated shifts in sensory weighting reflecting patterns of neural investment and ecological divergence in *Heliconius* butterflies. We demonstrated that *H. cydno*, which is characterised by a larger visual investment and more challenging light habitat, prioritises vision more strongly than *H. melpomene*. Alongside previous results in *Heliconius* butterflies and other taxa, our research reveals repeated shifts in sensory weighting associated with changes in neural investment and habitat use. This highlights the potential role for the sensory environment to shape differences in brain structure, which subsequently may influence key sensory behaviours crucial during the initial stages of species divergence.

Figure 1: Habitat preference and the sensory environment are associated with differences in brain investment in *Heliconius* butterflies. *H. cydno* invests more in visual neuropils [31] visual periphery [33, 40] and inhabits more-closed canopy forest habitats, than its close relative, *H. melpomene*, which is found in the forest edge [26, 30].

Figure 2: *H. cydno* and *H. melpomene* **weigh the importance of visual and olfactory cues differently when foraging. A)** Butterflies were trained to feed on blue feeders with a rose oil odour and negatively reinforced red feeders with a lavender smell. In the conflict test, the positively trained feeders were presented in combination with the negatively trained odours and vice versa. Training and reinforcement training periods are delineated by white rectangles, while behavioural tests are indicated by grey rectangles. **B)** Proportion of feeding attempts towards blue feeder for *H. cydno* and *H. melpomene*. Each point represents the feeding choices of an individual; solid dots and error bars represent the mean choices per species ±95% confidence intervals. Blue circles represent *H. cydno* and orange triangles represent *H. melpomene*, shapes in grey represent naive individuals that did not train. On naive and trained tests, feeder combinations were blue/rose and red/lavender; In the conflict test, the positively trained colour (blue) was presented in conflict with the negatively rewarded odour (lavender) and *vice versa*. In the colour tests, individuals were presented only with red or blue feeders without an odour. Only individuals that passed our training criteria (>=75% on the trained day) are shown (see methods).

307
308
309 **Table 1:** Pairwise p-value comparison matrix of post-hoc test with Bonferroni correction for the minimal adequate
310 model for the effects of the interaction between species and treatment (Trained, Conflict, and Colour) on choices
311 towards blue feeders. Only individuals that passed our training criteria (>=75% on the trained day) are on this
312 dataset. Results are from a binomial GLMM model, see methods for statistical details. Proportion blue ~ Species
313 * Level + (1|Group/ID)

	H. cydno Trained	*H. melpomene* Trained	*H. cydno* Conflict	*H. melpomene* Conflict	*H. cydno* Colour	*H. melpomene* Colour
H. cydno **Trained**	[2.329]	1	<.0001	<.0001	0.3041	0.008
H. melpomene **Trained**		[2.874]	<.0001	<.0001	0.0116	<.0001
H. cydno **Conflict**			[0.465]	0.0199	<.0001	1
H. melpomene **Conflict**				[-0.601]	<.0001	<.0001
H. cydno **Colour**					[1.639]	1
H. melpomene **Colour**						[0.989]

Acknowledgements

We thank Universidad del Rosario for insectary access at JCM and Isabel León for assistance, as well as Amulya Hosur and Alejandro Mancino for video scoring help.

Funding information

Research Funded by ERC Starter Grant 851040 to R.M.M.

Conflict of interest

The authors have no conflict of interest to declare.

References

1. Talsma D, Senkowski D, Soto-Faraco S, Woldorff MG. 2010 The multifaceted interplay between attention and multisensory integration. *Trends in Cognitive Sciences* **14**, 400–410. (doi:10.1016/j.tics.2010.06.008)

2. Stevens M. 2013 *Sensory ecology, behaviour, and evolution*. First edition. Oxford, United Kingdom: Oxford University Press.

3. Leonard AS, Dornhaus A, Papaj DR. 2011 Flowers help bees cope with uncertainty: signal detection and the function of floral complexity. *Journal of Experimental Biology* **214**, 113–121. (doi:10.1242/jeb.047407)

4. Balkenius A, Rosén W, Kelber A. 2006 The relative importance of olfaction and vision in a diurnal and a nocturnal hawkmoth. *J Comp Physiol A* **192**, 431–437. (doi:10.1007/s00359-005-0081-6)

5. Dell'Aglio DD, Rivas-Sánchez DF, Wright DS, Merrill RM, Montgomery SH. 2023 The Sensory Ecology of Speciation. *Cold Spring Harb Perspect Biol* , a041428. (doi:10.1101/cshperspect.a041428)

6. Pike TW, Ramsey M, Wilkinson A. 2018 Environmentally induced changes to brain morphology predict cognitive performance. *Philosophical Transactions of the Royal Society B: Biological Sciences* **373**, 20170287. (doi:10.1098/rstb.2017.0287)

7. Koethe S, Fischbach V, Banysch S, Reinartz L, Hrncir M, Lunau K. 2020 A Comparative Study of Food Source Selection in Stingless Bees and Honeybees: Scent Marks, Location, or Color. *Frontiers in Plant Science* **11**. (doi:10.3389/fpls.2020.00516)

8. Couto A, Wainwright JB, Morris BJ, Montgomery SH. 2020 Linking ecological specialisation to adaptations in butterfly brains and sensory systems. *Current Opinion in Insect Science* **42**, 55–60. (doi:10.1016/j.cois.2020.09.002)

9. Couto A *et al.* 2023 Rapid expansion and visual specialisation of learning and memory centres in the brains of Heliconiini butterflies. *Nat Commun* **14**, 4024. (doi:10.1038/s41467-023-39618-8)

10. Balkenius A, Kelber A. 2006 Colour preferences influences odour learning in the hawkmoth, Macroglossum stellatarum. *Naturwissenschaften* **93**, 255–258. (doi:10.1007/s00114-006-0099-9)

11. Stöckl A, Heinze S, Charalabidis A, El Jundi B, Warrant E, Kelber A. 2016 Differential investment in visual and olfactory brain areas reflects behavioural choices in hawk moths. *Scientific Reports* **6**, 1–10. (doi:10.1038/srep26041)

12. Andersson S, Dobson HEM. 2003 Behavioral foraging responses by the butterfly Heliconius melpomene to Lantana camara floral scent. *Journal of Chemical Ecology* **29**, 2303–2318. (doi:10.1023/A:1026226514968)

13. Thiele SC, Rodrigues D, Moreira GRP. 2016 Oviposition in Heliconius erato (Lepidoptera, Nymphalidae): how Essential Is Drumming Behavior for Host-Plant Selection? *J Insect Behav* **29**, 283–300. (doi:10.1007/s10905-016-9559-z)

14. Darragh K *et al.* 2017 Male sex pheromone components in Heliconius butterflies released by the androconia affect female choice. *PeerJ* **5**, e3953. (doi:10.7717/peerj.3953)

15. Silva DS, Barp EA, Kucharski LCR, Moreira GRP. 2018 Sensing the Plant Surface Prior to Feeding
and Oviposition: Differences in External Ultrastructure and Function Among Tarsi of Heliconius
erato. *Neotrop Entomol* **47**, 85–95. (doi:10.1007/s13744-017-0508-0)

16. van Schooten B, Meléndez-Rosa J, van Belleghem SM, Jiggins CD, Tan JD, Owen McMillan W,
Papa R. 2020 Divergence of chemosensing during the early stages of speciation. *Proceedings of*
the National Academy of Sciences of the United States of America **117**, 16438–16447.
(doi:10.1073/pnas.1921318117)

17. Dell'Aglio DD, McMillan WO, Montgomery SH. 2022 Shifting balances in the weighting of
sensory modalities are predicted by divergence in brain morphology in incipient species of
Heliconius butterflies. *Animal Behaviour* **185**, 83–90. (doi:10.1016/j.anbehav.2022.01.003)

18. Zaccardi G, Kelber A, Sison-Mangus MP, Briscoe AD. 2006 Color discrimination in the red range
with only one long-wavelength sensitive opsin. *Journal of Experimental Biology* **209**, 1944–1955.
(doi:10.1242/jeb.02207)

19. Toure MW, Young FJ, McMillan WO, Montgomery SH. 2020 Heliconiini butterflies can learn
time-dependent reward associations. *Biology Letters* **16**, 20200424.
(doi:10.1098/rsbl.2020.0424)

20. Jiggins CD, Naisbit RE, Coe RL, Mallet J. 2001 Reproductive isolation caused by colour pattern
mimicry. *Nature* **411**, 302–305. (doi:10.1038/35077075)

21. Estrada C, Jiggins CD. 2008 Interspecific sexual attraction because of convergence in warning
colouration: Is there a conflict between natural and sexual selection in mimetic species? *Journal*
of Evolutionary Biology **21**, 749–760. (doi:10.1111/j.1420-9101.2008.01517.x)

22. Merrill RM, Chia A, Nadeau NJ. 2014 Divergent warning patterns contribute to assortative
mating between incipient Heliconius species. *Ecology and Evolution* **4**, 911–917.
(doi:10.1002/ece3.996)

23. Gilbert LE. 1982 The Coevolution of a Butterfly and a Vine. *Scientific American* **247**, 110–121.
(doi:10.1038/scientificamerican0882-110)

24. Dell'Aglio DD, Losada ME, Jiggins CD. 2016 Butterfly learning and the diversification of plant leaf
shape. *Frontiers in Ecology and Evolution* **4**, 1–7. (doi:10.3389/fevo.2016.00081)

25. Jiggins CD, Mallet J. 2000 Bimodal hybrid zones and speciation. *Trends in Ecology & Evolution* **15**,
250–255. (doi:10.1016/S0169-5347(00)01873-5)

26. Estrada C, Jiggins CD. 2002 Patterns of pollen feeding and habitat preference among Heliconius
species. *Ecological Entomology* **27**, 448–456. (doi:10.1046/j.1365-2311.2002.00434.x)

27. Jiggins CD. 2008 Ecological speciation in mimetic butterflies. *BioScience*

28. Merrill RM *et al.* 2015 The diversification of Heliconius butterflies: What have we learned in 150
years? *Journal of Evolutionary Biology* **28**, 1417–1438. (doi:10.1111/jeb.12672)

29. Endler JA. 1993 The Color of Light in Forests and Its Implications. *Ecological Society of America*
63, 1–27.

30. Seymoure BM. 2016 Heliconius In A New Light: The Effects of Light Environments on Mimetic
Coloration, Behavior, and Visual Systems.

31. Montgomery SH, Merrill RM. 2017 Divergence in brain composition during the early stages of
ecological specialization in Heliconius butterflies. *Journal of Evolutionary Biology* **30**, 571–582.
(doi:10.1111/jeb.13027)

32. Montgomery SH, Rossi M, McMillan WO, Merrill RM. 2021 Neural divergence and hybrid
disruption between ecologically isolated *Heliconius* butterflies. *Proc. Natl. Acad. Sci. U.S.A.* **118**,
e2015102118. (doi:10.1073/pnas.2015102118)

33. Wright DS, Rodriguez-Fuentes J, Ammer L, Darragh K, Kuo C-Y, McMillan WO, Jiggins CD,
Montgomery SH, Merrill RM. 2023 Adaptive divergence in the eyes of *Heliconius* butterflies
likely contributes to pre- and post-mating isolation. (doi:10.1101/2023.10.26.564160)

34. Johnson MA, Revell LJ, Losos JB. 2009 BEHAVIORAL CONVERGENCE AND ADAPTIVE RADIATION:
EFFECTS OF HABITAT USE ON TERRITORIAL BEHAVIOR IN ANOLIS LIZARDS. *Evolution* **64**, 1151–
1159. (doi:10.1111/j.1558-5646.2009.00881.x)

35. Henry CS, Brooks SJ, Duelli P, Johnson JB, Wells MM, Mochizuki A. 2012 Parallel evolution in
courtship songs of North American and European green lacewings (Neuroptera: Chrysopidae):
PARALLEL EVOLUTION IN LACEWING SONGS. *Biological Journal of the Linnean Society* **105**, 776–
796. (doi:10.1111/j.1095-8312.2011.01845.x)

36. Bolnick DI, Barrett RDH, Oke KB, Rennison DJ, Stuart YE. 2018 (Non)Parallel Evolution. *Annu. Rev.
Ecol. Evol. Syst.* **49**, 303–330. (doi:10.1146/annurev-ecolsys-110617-062240)

37. Smiley J. 1978 The Host Plant Ecology of Heliconius Butterflies in Northeastern Costa Rica.
Universtiy of Texas at Austin.

38. Mérot C, Salazar C, Merrill RM, Jiggins CD, Joron M. 2017 What shapes the continuum of
reproductive isolation? Lessons from Heliconius butterflies. *Proceedings of the Royal Society B:*
Biological Sciences **284**, 20170335. (doi:10.1098/rspb.2017.0335)

39. Seymoure BM, Mcmillan WO, Rutowski R. 2015 Peripheral eye dimensions in Longwing
(Heliconius) butterflies vary with body size and sex but not light environment nor mimicry ring.
The Journal of Research on the Lepidoptera **48**, 83–92. (doi:10.5962/p.266475)

40. Bernays EA, Glendinning JI, Chapman RF. 1998 Plant acids modulate chemosensory responses in
Manduca sexta larvae. *Physiological Entomology* **23**, 193–201. (doi:10.1046/j.1365-
3032.1998.233079.x)

41. Friard O, Gamba M. 2016 BORIS: a free, versatile open-source event-logging software for
video/audio coding and live observations. *Methods in Ecology and Evolution* **7**, 1325–1330.
(doi:10.1111/2041-210X.12584)

42. Damas-Moreira I, Oliveira D, Santos JL, Riley JL, Harris DJ, Whiting MJ. 2018 Learning from
others: an invasive lizard uses social information from both conspecifics and heterospecifics.
Biology Letters **14**, 20180532. (doi:10.1098/rsbl.2018.0532)

43. Batabyal A, Thaker M. 2019 Lizards from suburban areas learn faster to stay safe. *Biology Letters*
(doi:10.1098/rsbl.2019.0009)

44. Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting Linear Mixed-Effects Models Using lme4.
Journal of Statistical Software **67**, 1–48. (doi:10.18637/jss.v067.i01)

45. R Core Team. 2023 R: a Language and Environment for Statistical Computing.

46. Lenth R, Singmann H, Love J, Buerkner P, Herve M. 2019 Package 'emmeans'.

47. Wickham H. 2011 ggplot2. *Wiley interdisciplinary reviews: computational statistics* **3**, 180–185.
(doi:10.1002/wics.147)

48. Jiggins CD, McMillan WO, Neukirchen W, Mallet J. 1996 What can hybrid zones tell us about
speciation? The case of Heliconius erato and H. himera (Lepidoptera: Nymphalidae). *Biological*
Journal of the Linnean Society **59**, 221–242. (doi:10.1111/j.1095-8312.1996.tb01464.x)

49. Rubene D, Low M, Brodin A. 2023 Birds differentially prioritize visual and olfactory foraging cues
depending on habitat of origin and sex. *Royal Society Open Science* **10**, 221336.
(doi:10.1098/rsos.221336)

50. Keagy J, Braithwaite VA, Boughman JW. 2018 Brain differences in ecologically differentiated
sticklebacks. *Current Zoology* **64**, 243–250. (doi:10.1093/cz/zox074)

453
454

455   Supplementary material:

Figure S 1: **Experimental set-up. A**) Groups of *H. cydno* and *H. melpomene* were presented with artificial flower arrays at the end of the cages and were trained to a certain colour/odour combination. **B)** Artificial flower array containing the multimodal visual cues (coloured circles) and olfactory cue (PCR tube containing essential
oil).

Supplementary Results

Among the 117 individuals (82 *H. melpomene* and 35 *H. cydno*) that passed our less

stringent training criterion (i.e., >50% correct choices on day 5), *H. cydno* did not differ from

H. melpomene in the trained test, indicating that both species were able to train to the

multimodal stimulus (Table S1 post-hoc: *H. cydno* Trained vs. *H. melpomene* Trained, Z= -

1.333, P= 1.0). However, *H. melpomene* and *H. cydno* did respond differently when

presented with the olfactory and visual stimuli in conflict (Figure S2). Importantly, the

interaction between species and trial type was retained in our model, confirming that this

was not due to existing differences amongst "trained" individuals, but rather the effect of how

the stimuli were presented (species*treatment LRT: 2ΔlnL = 44.537, df=2, p < 0.001). Table

S5). In the conflict test, *H. cydno* fed mostly on the blue feeder, while *H. melpomene*

primarily chose the feeder with the rose scent, suggesting a difference in sensory weighting

between these species (Table S1, post-hoc: *H. cydno* Conflict vs. *H. melpomene* Conflict,

Z= 4.402, P< 0.001). We found no effect of sex, which was excluded from our model

(species*trial type LRT: 2ΔlnL = 0.297, df=1, p = 0.586. Table S5).

Our post-hoc analysis revealed that individuals of both species made fewer feeding attempts

on the blue feeder when the stimuli were presented in conflict, indicating that both species

integrate visual and olfactory stimuli when making foraging decisions (Table S1, post-hoc: *H.*

cydno Trained vs. *H. cydno* Conflict, Z= 5.760, P= <.0001) (Table S1, post-hoc: *H.*

melpomene Trained vs. *H. melpomene* Conflict, Z=23.222, P= <.0001). Notably, *H. cydno*

did not differ from the trained test and the colour-only test indicating that this species largely

relies on visual cues (Table S1, post-hoc: *H. cydno* Trained vs. *H. cydno* Conflict, Z= 0.337,

P= 1.0). In contrast, *H. melpomene* differed between the trained test and the colour-only test

further suggesting a difference in sensory weighting (Table 1, post-hoc: *H. melpomene*

Figure S 2: *H. cydno* and *melpomene* weigh the importance of visual and olfactory cues differently when foraging. Proportion of feeding attempts towards blue feeder for *H. cydno* and *H. melpomene*. Each point represents the feeding choices of and individual; solid dots and error bars represent the mean choices per species ±95% confidence intervals. Blue circles represent *H. cydno* and orange triangles represent *H. melpomene*, shapes in grey represent naive individuals that did not train. On naive and trained tests, feeder combinations were blue/rose and red/lavender; In the conflict test, the positively trained colour (blue) was presented in conflict with the negatively rewarded odour (lavender) and *vice versa*. In the colour tests, individuals were presented only with red or blue feeders without an odour. Only individuals that passed our training criteria (>=50% on the trained day) are show (see methods).

482 Trained vs. *H. melpomene* Conflict, Z= 8.433, P= <.0001).

483
484 **Table S 1:** Pairwise p-value comparison matrix of post-hoc test with Bonferroni correction for the minimal
485 adequate model for the effects of the interaction between species, treatment (Trained, Conflict, and Colour) and
486 their interaction on choices towards blue feeders. Only individuals that passed our training criteria (>=50% on the
487 trained day) are on this dataset. Results are from binomial GLMM model, see methods for statistical details.

	H. cydno Trained	*H. melpomene* Trained	*H. cydno* Conflict	*H. melpomene* Conflict	*H. cydno* Colour	*H. melpomene* Colour
H. cydno **Trained**	[1.331]	1	<.0001	<.0001	1	0.338
H. melpomene **Trained**		[1.656]	<.0001	<.0001	1	<.0001
H. cydno **Conflict**			[0.360]	0.0002	<.0001	1
H. melpomene **Conflict**				[-0.649]	<.0001	<.0001
H. cydno **Colour**					[1.265]	0.7534
H. melpomene **Colour**						[0.771]

Figure S 3: Behavioural choices including individuals that did not reach our training criterion (>=75%). Proportion of feeding attempts towards blue feeder for *H. cydno* and *H. melpomene*. Each point represents the feeding choices of and individual; solid dots and error bars represent the mean choices per species ±95% confidence intervals. Blue circles represent *H. cydno* and orange triangles represent *H. melpomene*, shapes in grey represent individuals that did not train. Gray error bars represent the mean choices per species ±95% confidence intervals including all individuals.

491 **Table S 2:** Number of individuals that participated in each treatment (Naïve, Trained, Conflict, Colour) separated
492 by number of individuals that passed the training criterion (>=50% or >=75%) of correct choices on the trained
493 day (day 5).

	H. cydno	*H. melpomene*	*H. cydno* (>=50%) training	*H. melpomene* (>=50%) training	*H. cydno* (>=75%) training	*H. melpomene* (>=75%) training
Naive	61	74	13	48	8	25
Trained	79	108	35	82	21	52
Conflict	91	109	31	74	19	48
Colour	71	96	27	66	18	42

Figure S 4: Estimated marginal means (e.m.m.) of the minimum adequate statistical models for the proportion of feeding attempts towards blue feeder demonstrating the significant effects of the interaction between species (H. cydno and H. melpomene) and treatment [Trained test (day 5), Conflict test (day 6) or Colour test (day 7)]. Only individuals that passed our training criteria (>=75% on the trained day) are on this dataset. (see methods).

496
497 **Table S 3:** Model 1 minimal adequate model for the effects of species, treatment (Trained, Conflict, Colour) and
498 their interaction on choices towards blue feeders. Only individuals that passed our training criteria (>=75% on the
499 trained day) are on this dataset. Results are from a binomial GLMM model, see methods for statistical details.
500 Proportion blue ~ Species * Level + (1|Group/ID)
501

	2ΔlnL	df	Pr
Species	27.846	3	3.912e-06 ***
Treatment	681.740	4	< 2.2e-16 ***
Treatment:Species	22.876	2	1.078e-05 ***
Sex	3.067	1	0.07988

503
504 **Table S 4:** Pairwise comparison post-hoc test with Bonferroni correction for the minimal adequate model for the
505 effects of the interaction between species and treatment (Trained, Conflict, and Colour) on choices towards blue
506 feeders. Only individuals that passed our training criteria (>=75% on the trained day) are on this dataset. Results
507 are from binomial GLMM model, see methods for statistical details. Proportion blue ~ Species * Level +
508 (1|Group/ID)
509

	estimate	SE	z-ratio	p-value (Adjusted)
H. cydno Trained - *H. melpomene* Trained	-0.545	0.396	-1.375	1.000
H. cydno Trained - *H. cydno* Conflict	1.864	0.269	6.932	<.0001
H. cydno Trained - *H. melpomene* Conflict	2.930	0.381	7.696	<.0001
H. cydno Trained - *H. cydno* Colour	0.690	0.297	2.321	0.304
H. cydno Trained - *H. melpomene* Colour	1.340	0.387	3.464	0.008
H. melpomene Trained - *H. cydno* Conflict	2.409	0.351	6.860	<.0001
H. melpomene Trained - *H. melpomene* Conflict	3.475	0.169	20.614	<.0001
H. melpomene Trained - *H. cydno* Colour	1.235	0.367	3.361	0.012
H. melpomene Trained - *H. melpomene* Colour	1.884	0.173	10.920	<.0001
H. cydno Conflict - H. melpomene Conflict	1.066	0.332	3.210	0.020
H. cydno Conflict - *H. cydno* Colour	-1.174	0.237	-4.956	<.0001
H. cydno Conflict - H. melpomene Colour	-0.524	0.340	-1.544	1.000
H. melpomene Conflict - *H. cydno* Colour	-2.240	0.351	-6.385	<.0001
H. melpomene Conflict - *H. melpomene* Colour	-1.590	0.141	-11.244	<.0001
H. cydno Colour - *H. melpomene* Colour	0.650	0.357	1.819	1.000

Figure S 5: Estimated marginal means (e.m.m.) of the minimum adequate statistical models for the proportion of feeding attempts towards blue feeder demonstrating the significant effects of the interaction between species (*H. cydno* and *H. melpomene)* and treatment [*Trained test* (day 5), *Conflict test* (day 6) or *Colour test* (day 7)]. Only individuals that passed our training criteria (>=50% on the trained day) are on this dataset. (see methods).

511
512
513 **Table S 5:** Model 2 minimal adequate model for the effects of species, treatment (Trained, Conflict, Colour) and
514 their interaction on choices towards blue feeders. (>=50% on the trained day) are on this dataset. Results are
515 from binomial GLMM model, see methods for statistical details.
516 Proportion blue ~ Species * Level + (1|Group/ID)

	2ΔlnL	df	Pr
Species	50.004	3	7.972e-11 ***
Treatment	690.730	4	< 2.2e-16 ***
Treatment:Species	44.537	2	2.132e-10 ***
Sex	0.297	1	0.58550

519 **Table S 6**: Pairwise comparison post-hoc test with Bonferroni correction for the minimal adequate model for the

520 effects of the interaction between species and treatment (Trained, Conflict, and Colour) on choices towards blue

521 feeders. Only individuals that passed our training criteria (>=50% on the trained day) are on this dataset. Results

522 are from binomial GLMM model, see methods for statistical details.

	estimate	SE	z-ratio	p-value (Adjusted)
H. cydno Trained - *H. melpomene* Trained	-0.326	0.244	-1.333	1.000
H. cydno Trained - *H. cydno* Conflict	0.971	0.169	5.760	<.0001
H. cydno Trained - *H. melpomene* Conflict	1.980	0.241	8.199	<.0001
H. cydno Trained - *H. cydno* Colour	0.066	0.195	0.337	1.000
H. cydno Trained - *H. melpomene* Colour	0.560	0.245	2.281	0.338
H. melpomene Trained - *H. cydno* Conflict	1.296	0.233	5.571	<.0001
H. melpomene Trained - *H. melpomene* Conflict	2.305	0.099	23.222	<.0001
H. melpomene Trained - *H. cydno* Colour	0.391	0.251	1.556	1.000
H. melpomene Trained - *H. melpomene* Colour	0.885	0.105	8.433	<.0001
H. cydno Conflict - *H. melpomene* Conflict	1.009	0.229	4.402	0.000
H. cydno Conflict - *H. cydno* Colour	-0.905	0.183	-4.942	<.0001
H. cydno Conflict - *H. melpomene* Colour	-0.411	0.234	-1.760	1.000
H. melpomene Conflict - *H. cydno* Colour	-1.914	0.249	-7.697	<.0001
H. melpomene Conflict - *H. melpomene* Colour	-1.420	0.100	-14.263	<.0001
H. cydno Colour - *H. melpomene* Colour	0.494	0.252	1.958	0.753

H. cydno

H. melpomene

Optic lobe
(+ eye size + facet number)

light intensity

Closed tropical forest

Forest edge/open

A

2 m

B

PCR-tube containing odour cue

Visual cue printed on waterproof paper