PDGFRα signaling regulates Srsf3 transcript binding to affect PI3K signaling and endosomal

trafficking

Thomas E. Forman[1,2], Marcin P. Sajek[3,4,5], Eric D. Larson[6,7], Neelanjan Mukherjee[3,4], Katherine

A. Fantauzzo[1,4,*]

[1]Department of Craniofacial Biology, School of Dental Medicine, University of Colorado

Anschutz Medical Campus, Aurora, CO, USA

[2]Medical Scientist Training Program, University of Colorado Anschutz Medical Campus, Aurora,

CO, USA

[3]Department of Biochemistry and Molecular Genetics, School of Medicine, University of

Colorado Anschutz Medical Campus, Aurora, CO, USA

[4]RNA Bioscience Initiative, University of Colorado Anschutz Medical Campus, Aurora, CO, USA

[5]Institute of Human Genetics, Polish Academy of Sciences, Poznan, Poland

[6]Department of Otolaryngology – Head and Neck Surgery, University of Colorado Anschutz

Medical Campus, Aurora, CO, USA

[7]Basic and Translational Sciences, Penn Dental Medicine, University of Pennsylvania,

Philadelphia, PA, USA

*Corresponding author. Email: katherine.fantauzzo@cuanschutz.edu

Abstract

Signaling through the platelet-derived growth factor receptor alpha (PDGFRα) plays a

critical role in craniofacial development, as mutations in *PDGFRA* are associated with cleft

lip/palate in humans and *Pdgfra* mutant mouse models display varying degrees of facial clefting.

Phosphatidylinositol 3-kinase (PI3K)/Akt is the primary effector of PDGFRα signaling during

skeletal development in the mouse. We previously demonstrated that Akt phosphorylates the

RNA-binding protein serine/arginine-rich splicing factor 3 (Srsf3) downstream of PI3K-mediated

PDGFRα signaling in mouse embryonic palatal mesenchyme (MEPM) cells, leading to its

nuclear translocation. We further showed that ablation of *Srsf3* in the murine neural crest

lineage results in severe midline facial clefting, due to defects in proliferation and survival of

cranial neural crest cells, and widespread alternative RNA splicing (AS) changes. Here, we

sought to determine the molecular mechanisms by which Srsf3 activity is regulated downstream

of PDGFRα signaling to control AS of transcripts necessary for craniofacial development. We

demonstrated via enhanced UV-crosslinking and immunoprecipitation (eCLIP) of MEPM cells

that PDGF-AA stimulation leads to preferential binding of Srsf3 to exons and loss of binding to

canonical Srsf3 CA-rich motifs. Through the analysis of complementary RNA-seq data, we

showed that Srsf3 activity results in the preferential inclusion of exons with increased GC

content and lower intron to exon length ratio. Moreover, we found that the subset of transcripts

that are bound by Srsf3 and undergo AS upon PDGFRα signaling commonly encode regulators

of PI3K signaling and early endosomal trafficking. Functional validation studies further

confirmed that Srsf3 activity downstream of PDGFRα signaling leads to retention of the receptor

in early endosomes and increases in downstream PI3K-mediated Akt signaling. Taken together,

our findings reveal that growth factor-mediated phosphorylation of an RNA-binding protein

underlies gene expression regulation necessary for mammalian craniofacial development.

Introduction

Craniofacial development is a complex morphogenetic process that requires a precise

interplay of multiple cell and tissue types to generate the frontonasal skeleton. Disruption of this

process can result in some of the most common birth defects in humans, such as cleft lip and

palate (Mai et al., 2019). Signaling through the platelet derived growth factor receptor alpha

(PDGFRα) receptor tyrosine kinase (RTK) is essential for human craniofacial development.

Heterozygous missense mutations in the coding region of *PDGFRA* that alter amino acids within

the extracellular, transmembrane or cytoplasmic domains of the receptor, in addition to single

base-pair substitutions in the 3' untranslated region (3' UTR), are associated with nonsyndromic

cleft palate (Rattanasopha et al., 2012). Further, single-nucleotide polymorphisms that repress

transcriptional activity of the promoter upstream of *PDGFC*, which encodes one of two PDGFRα

ligands, are associated with cleft lip and palate (Choi et al., 2009). This role of PDGFRα

signaling in craniofacial development is conserved in mice, as *Pdgfra* mutant mouse models

display a variety of defects that range from cleft palate to complete facial clefting (Klinghoffer et

al., 2002; Soriano, 1997; Tallquist & Soriano, 2003; Fantauzzo & Soriano, 2014; He & Soriano,

2013). These phenotypes are recapitulated in embryos lacking both *Pdgfa* and *Pdgfc* (Ding et

al., 2004). Phosphatidylinositol 3-kinase (PI3K) is the primary effector of PDGFRα signaling

during skeletal development in the mouse (Klinghoffer et al., 2002). Following activation, PI3K

increases phosphatidylinositol-3,4,5-trisphosphate (PIP$_3$) levels at the cell membrane, leading to

the recruitment and subsequent phosphorylation of the serine/threonine kinase Akt. Akt

subsequently dissociates from the membrane to phosphorylate an array of target proteins that

are involved in wide-ranging cellular processes (Manning & Cantley, 2007). We previously

identified proteins phosphorylated by Akt downstream of PI3K-mediated PDGFRα signaling in

primary mouse embryonic mesenchyme (MEPM) cells (Fantauzzo & Soriano, 2014). Gene

ontology analysis revealed that 25% of the 56 proteins were involved in RNA processing, with a

particular enrichment for RNA splicing (Fantauzzo & Soriano, 2014).

Alternative RNA splicing (AS) is a process by which different combinations of exons from

the same gene are incorporated into mature RNA transcripts, thereby contributing to gene

expression regulation and enhancing the diversity of protein isoforms (Licatalosi & Darnell,

2010). AS occurs in approximately 95% of multi-exon human genes, frequently in a tissue-

specific manner (Pan et al., 2008; Wang et al., 2008). Dysregulation of AS causes a number of

diseases, due to mutations in precursor RNA sequences, mutations in core components of the

spliceosome complex and/or mutations in auxiliary RNA-binding proteins (RBPs) (Scotti &

Swanson, 2016). These *trans*-acting auxiliary RBPs bind to specific sequence and/or structural

motifs in a target RNA via one or more RNA-binding domains to promote or inhibit exon

inclusion (Fu & Ares, 2014; Licatalosi & Darnell, 2010). The phenotypes resulting from global

and/or tissue-specific knockout of multiple RBPs have established that RBP-mediated AS is an

essential process during mouse craniofacial development (Bebee et al., 2015; Cibi et al., 2019;

Dennison et al., 2021; Forman et al., 2021; Lee et al., 2020). We previously demonstrated that

ablation of *Srsf3* in the murine neural crest lineage results in severe midline facial clefting and

facial bone hypoplasia, due to defects in proliferation and survival of cranial neural crest cells,

and widespread AS changes (Dennison et al., 2021).

Srsf3 belongs to the serine/arginine-rich (SR) protein family of splicing factors that

generally promote exon inclusion by binding to exonic and intronic splicing enhancers and by

recruiting spliceosome components to the 5' and 3' splice sites (Fu & Ares, 2014; Licatalosi &

Darnell, 2010). Srsf3 specifically was shown to bind pyrimidine-rich motifs, with a preference for

exonic regions (Akerman et al., 2009; Änkö et al., 2012). Srsf3 is phosphorylated downstream of

PDGF and EGF stimulation and all-*trans* retinoic acid treatment, and by the kinases Akt and

SRPK2 (Bavelloni et al., 2014; Dennison et al., 2021; Fantauzzo & Soriano, 2014; Long et al.,

2019; Zhou et al., 2012). Phosphorylation of the Akt consensus sites within the C-terminal

arginine/serine-rich (RS) domain of Srsf3 drives its translocation to the nucleus, where AS takes

place (Bavelloni et al., 2014; Dennison et al., 2021; Long et al., 2019). Moreover,

phosphorylation of the RS domain can alter the ability of SR proteins to interact with other

proteins, such as the U1 small nuclear ribonucleoprotein (snRNP) component of the

spliceosome and additional RBPs, and affect the ability of SR proteins to bind RNA (Huang et

al., 2004; Shen & Green, 2006; Shin et al., 2004; Xiao & Manley, 1997). However, the molecular

mechanisms by which Srsf3 activity is regulated downstream of specific signaling inputs in a

context-specific manner to regulate RNA binding and/or sequence specificity remain

undetermined.

Here, we identified changes in Srsf3-dependent AS and Srsf3 transcript binding in the

absence or presence of PDGF-AA ligand in MEPM cells. RNA-sequencing (RNA-seq) analysis

revealed that Srsf3 activity and PDGFRα signaling have more pronounced effects on AS than

gene expression, as well as a significant dependence in regulating AS. Using enhanced UV-

crosslinking and immunoprecipitation (eCLIP), we found a shift from intronic to exonic Srsf3

binding and loss of CA-rich sites upon PDGF-AA stimulation. Further, we demonstrated that the

subset of transcripts that are bound by Srsf3 and undergo AS upon PDGFRα signaling

commonly encode regulators of PI3K signaling and early endosomal trafficking, ultimately

serving as a feedback mechanism to affect trafficking of the receptors. Combined, our findings

provide significant insight into the mechanisms underlying RBP-mediated gene expression

regulation in response to growth factor stimulation within the embryonic mesenchyme.

Results

PDGFRα signaling for one hour minimally affects gene expression

To determine the Srsf3-dependent changes in gene expression and AS downstream of

PDGFRα signaling, we stably integrated a scramble shRNA (scramble) or an shRNA targeting

the 3' UTR of *Srsf3* (shSrsf3) into immortalized MEPM (iMEPM) cells via lentiviral transduction

(Fig. 1A). Western blotting revealed a 66% decrease in Srsf3 protein levels in the shSrsf3 cell

line (Fig. 1B). We previously demonstrated that phosphorylated Srsf3 levels peaked in the

nucleus of iMEPM cells following 1 hour of PDGF-AA ligand treatment (Dennison et al., 2021).

As such, cells were left unstimulated (-PDGF-AA) or treated with 10 ng/mL PDGF-AA for 1 hour

(+PDGF-AA). RNA was isolated and sequenced from three biological replicates across each of

the four conditions (Fig. 1A; Table S1). We first examined Srsf3-dependent differentially-

expressed (DE) genes by comparing scramble versus shSrsf3 samples across the same PDGF-

AA stimulation condition (-PDGF-AA or +PDGF-AA). We detected 827 DE genes in the absence

of ligand treatment and 802 DE genes upon ligand stimulation (Fig. 1C, Table S2). There was

high overlap (1,042 out of 1,629; 64.0%) of DE genes across ligand treatment conditions (Fig.

1C,D; Fig. S1A). Of these 521 shared DE genes, 514 (98.7%) had the same directionality,

including 273 (52%) with shared increases in expression in the shSrsf3 samples and 241 (46%)

with shared decreases in expression in the shSrsf3 samples (Fig. 1C,D). These findings

demonstrate that expression of a set of genes (521) depends on Srsf3 activity independent of

PDGFRα signaling, while a similarly sized set of genes (587) is differentially expressed in

response to both Srsf3 activity and PDGFRα signaling (Fig. 1C,D). Gene ontology (GO)

analysis of the Srsf3-dependent DE genes revealed that the most significant terms for biological

process commonly involved regulation of osteoblast differentiation, calcium-dependent cell-cell

adhesion, regulation of cell migration and canonical Wnt signaling, while only a handful of

significant terms for molecular function were detected in unstimulated cells, relating to cation

channel activity and phosphatase activity (Fig. S2A,B).

We next examined PDGF-AA-dependent DE genes by comparing -PDGF-AA versus

+PDGF-AA across the same Srsf3 condition (scramble or shSrsf3). We detected only 37 DE

genes in scramble cells and 14 DE genes in shSrsf3 cells (Fig. 1E, Table S2). There was low

overlap (8 out of 51; 15.7%) of DE genes across Srsf3 conditions (Fig. 1E,F; Fig. S1B). Of these

4 shared DE genes, 3 (75%) had shared increases in expression upon PDGF-AA stimulation

(Fig. 1E,F). These findings demonstrate that, unlike Srsf3 activity, PDGFRα signaling minimally

affects gene expression at one hour of ligand stimulation, consistent with our previous findings

in mouse embryonic facial mesenchyme (Dennison et al., 2021). Further, these results show

that expression of a set of genes (4) depends on PDGFRα signaling independent of Srsf3

activity, while a larger set of genes (43) is differentially expressed in response to both PDGFRα

signaling and Srsf3 activity (Fig. 1E,F). GO analysis of the PDGF-AA-dependent DE genes

revealed significant terms for biological process in the scramble cells relating to cell migration,

response to growth factor stimulation and regulation of transcription (Fig. S2A). The most

significant terms for molecular function commonly involved DNA binding (Fig. S2B).

PDGFRα signaling for one hour has a more pronounced effect on alternative RNA splicing

158   We previously demonstrated that AS is an important mechanism of gene expression

regulation downstream of PI3K/Akt-mediated PDGFRα signaling in the murine mid-gestation

palatal shelves (Dennison et al., 2021). Accordingly, we next assessed AS in our same RNA-

seq dataset. In examining Srsf3-dependent alternatively-spliced transcripts, we detected 1,354

differential AS events in the absence of ligand treatment and 1,071 differential AS events upon

ligand stimulation (Fig. 2A). When filtered to include events detected in at least 10 reads in

either condition, we obtained a list of 1,113 differential AS events in the absence of ligand

treatment and 795 differential AS events upon ligand stimulation (Fig. 2B, Tables S3 and S4).

There was low overlap (406 out of 1,908; 21.3%) of alternatively-spliced transcripts across

ligand treatment conditions. Of these 203 shared alternatively-spliced transcripts, 100% had the

same directionality, including 81 (40%) with shared negative changes in percent spliced in (PSI)

(exon included more often in shSrsf3 samples) and 122 (60%) with shared positive changes in

PSI (exon skipped more often in shSrsf3 samples) (Fig. 2A,B). We confirmed the differential AS

of two transcripts, *Arhgap12* and *Cep55*, between scramble and shSrsf3 samples by qPCR

using primers within constitutively-expressed exons flanking the alternatively-spliced exon (Fig.

S3A,B). These findings demonstrate that AS of a set of transcripts (203) depends on Srsf3

activity independent of PDGFRα signaling, while a much larger set of transcripts (1,502) is

alternatively spliced in response to both Srsf3 activity and PDGFRα signaling (Fig. 2A,B). GO

analysis of these Srsf3-dependent alternatively-spliced transcripts revealed that the most

significant terms for biological process and molecular function commonly involved regulation of

RNA splicing, and RNA binding and cadherin binding, respectively (Fig. S4A,B).

In examining PDGF-AA-dependent alternatively-spliced transcripts, we detected 595

differential AS events in scramble cells and 398 differential AS events in shSrsf3 cells (Fig. 2C).

When filtered to include events detected in at least 10 reads in either condition, we obtained a

list of 375 differential AS events in scramble cells and 256 differential AS events in shSrsf3 cells

(Fig. 2D, Tables S5 and S6). There was extremely low overlap (18 out of 631; 2.85%) of

alternatively-spliced transcripts across Srsf3 conditions (Fig. 2C,D). Of these 9 shared

alternatively-spliced transcripts, 100% had the same directionality, including 5 (56%) with

shared negative changes in PSI (exon included more often in +PDGF-AA samples) and 4 (44%)

with shared positive changes in PSI (exon skipped more often in +PDGF-AA samples) (Fig.

2C,D). Taken together, these findings demonstrate that both Srsf3 activity and PDGFRα

signaling have more pronounced effects on AS than gene expression. While more transcripts

and genes are subject to these regulatory mechanisms upon loss of Srsf3 activity, the

magnitude of events is more greatly skewed towards AS upon PDGF-AA stimulation. Further,

these results show that AS of a set of transcripts (9) depends on PDGFRα signaling

independent of Srsf3 activity, while a much larger set of transcripts (613) is alternatively spliced

in response to both PDGFRα signaling and Srsf3 activity (Fig. 2C,D). When combined with the

data above (Fig. 2B), this points to a profound dependence between PDGFRα signaling and

Srsf3 in regulating AS of transcripts in the facial mesenchyme. GO analysis of these PDGF-AA-

dependent alternatively-spliced transcripts demonstrated only a handful of significant terms for

molecular function in the scramble cells, all relating to protein kinase activity (Fig. S4B).

The vast majority of Srsf3-dependent (70.2-73.7%) and PDGF-AA dependent (62.1-

65.1%) AS events involved skipped exons, with minimal contribution from retained introns,

mutually exclusive exons, alternative 5' splice sites, or alternative 3' splice sites (Fig. 2E). For

the Srsf3-dependent skipped exon events, there were more transcripts with +ΔPSI (exon

skipped more often in shSrsf3 samples) (44.8%) as opposed to -ΔPSI (exon included more

often in shSrsf3 samples) (25.4%) in the absence of PDGF-AA stimulation (Fig. 2E), consistent

with previous results that SR proteins tend to promote exon inclusion (Fu & Ares, 2014;

Licatalosi & Darnell, 2010). However, PDGF-AA ligand treatment led to an increase in the

percentage of transcripts with -ΔPSI (36.5%) (Fig. 2E), indicating that PDGFRα signaling

promotes exon skipping in the presence of Srsf3. Among the PDGF-AA-dependent skipped

exon events, there was a significant shift in transcripts with -ΔPSI (exon included more often in

+PDGF-AA samples) in the absence (41.0%) versus presence (19.5%) of Srsf3, and a

corresponding shift in transcripts with +ΔPSI (exon skipped more often in +PDGF-AA samples)

(21.1% and 45.6%, respectively) (Fig. 2E), again suggesting that PDGF-AA stimulation causes

increased exon skipping when Srsf3 is present.

Srsf3 exhibits differential transcript binding upon PDGFRα signaling

To determine direct binding targets of Srsf3 downstream of PDGFRα signaling, we

conducted eCLIP (Van Nostrand et al., 2016, 2017) of iMEPM cells that were left unstimulated (-

PDGF-AA) or treated with 10 ng/mL PDGF-AA for 1 hour (+PDGF-AA) as above (Fig. 3A; Table

S7). Immunoprecipitation with a previously validated Srsf3 antibody (Dennison et al., 2021)

successfully enriched for Srsf3 in UV-crosslinked cells (Fig. 3B). We detected 6,555 total eCLIP

peaks in protein-coding genes in the -PDGF-AA samples and 8,584 total peaks in the +PDGF-

AA samples (Table S8). Among the -PDGF-AA peaks, 3,727 (56.9%) were located in exons

(CDS) and 1,690 (25.8%) were located within introns, with the rest binding within 5' UTRs (768,

11.7%) and 3' UTRs (367, 5.60%) (Fig. 3C,D; Table S9). We observed substantial shifts in

Srsf3-bound regions upon PDGF-AA stimulation. Many more +PDGF-AA peaks were located

within exons (7,139, 83.2%), and less were located within introns (765, 8.91%), 5' UTRs (389,

4.53%) and 3' UTRs (287, 3.34%) (Fig. 3C,D; Table S9). Given that SR proteins are crucial for

exon definition and bind to exonic splicing enhancers to recruit and stabilize core splicing

machinery (Fu & Maniatis, 1990; Krainer et al., 1991; Zahler et al., 1993), we investigated Srsf3

binding around 5' and 3' splice sites in response to PDGFRα signaling. Consistent with the

results above and previous findings (Änkö et al., 2012), Srsf3 binding was enriched in exonic

regions, as opposed to intronic regions, surrounding the splice sites (Fig. 3E). There was

greater mean coverage of Srsf3 peaks in the +PDGF-AA condition versus the -PDGF-AA

condition within 100 nucleotides upstream of the 5' splice site and within 100 nucleotides

downstream of the 3' splice site (Fig. 3E). Additionally, we detected decreased mean coverage

in the +PDGF-AA condition within 25 nucleotides downstream of the 5' splice site boundary

(Fig. 3E). Taken together, these data show that PDGFRα signaling leads to increased binding of

Srsf3 to exons. Finally, we performed unbiased motif enrichment analysis of Srsf3 peaks in

unstimulated and PDGF-AA-treated samples, revealing that the mostly highly enriched motifs in

-PDGF-AA samples were CACACA and AAGAAG (Fig. 3F). Of note, these CA-rich motifs have

previously been identified as canonical Srsf3 motifs in a CLIP study utilizing a stably integrated

SRSF3-EGFP transgene (Änkö et al., 2012). However, in PDGF-AA-stimulated samples, the

most highly enriched motifs were GAAGCG, GAAGAA, and AGAAGA (Fig. 3G), suggesting that

PDGFRα signaling influences Srsf3 binding specificity.

Srsf3 and PDGFRα signaling are associated with differential GC content and length of

alternatively-spliced exons

We next probed our RNA-seq dataset to determine whether specific transcript features

were associated with Srsf3-dependent AS. When comparing significant AS events between

scramble and shSrsf3 samples in the absence of PDGF-AA stimulation, we found that included

exons had a significantly higher GC content (median value of 53.4%) than both skipped exons

(50.0%) and exons that were not differentially alternatively spliced (51.2%) when Srsf3 is

present (Fig. 4A, Table S10). Additionally, we observed that included exons had a significantly

lower ratio of downstream intron to exon GC content (0.856) than both skipped exons (0.901)

and exons that were not differentially alternatively spliced (0.888) in the presence of Srsf3 (Fig.

4B, Table S10). The same comparisons revealed that the ratio of upstream and downstream

intron to exon length was significantly decreased in included exons (median values of 12.4 and

10.6, respectively) as compared to both skipped exons (19.5 and 20.0) and exons that were not

differentially alternatively spliced (14.2 and 13.3) in the presence of Srsf3 (Fig. 4C,D, Table

S10). Taken together, our data demonstrate that exons which are included in the presence of

Srsf3 tend to have a higher GC content and lower intron to exon length ratio.

To determine whether PDGFRα signaling had an effect on the transcript features to

which Srsf3 bound, we subsequently examined our eCLIP dataset. We found that the exon GC

content was significantly increased in exons bound by Srsf3 in the absence of ligand treatment

(median value of 57.9%) as compared to unbound exons (51.5%) (Fig. 4E, Table S10).

However, exon GC content was similar between unbound exons and those bound by Srsf3

upon PDGF-AA stimulation (53.8%) (Fig. 4E, Table S10). These findings indicate that

PDGFRα signaling mediates binding of Srsf3 to exons with a lower GC content.

Transcripts bound by Srsf3 that undergo alternative splicing upon PDGFRα signaling encode

regulators of PI3K signaling

To determine which transcripts are directly bound by Srsf3 and subject to DE and/or AS,

we cross-referenced the eCLIP and RNA-seq datasets. We collated transcripts with an Srsf3

eCLIP peak that were uniquely detected in the -PDGF-AA or +PDGF-AA samples (2,660

transcripts across 3,388 peaks) (Tables S8 and S11). Similarly, we gathered differentially-

expressed genes (596) or differentially alternatively-spliced transcripts (985) uniquely found in

one of the four treatment comparisons (Fig. 5A, Table S11). Only 32 (5.4%) of the DE genes

were directly bound by Srsf3, while 233 (23.7%) of the alternatively-spliced transcripts had an

Srsf3 eCLIP peak, with very little overlap (1 transcript) between all three categories (Fig. 5A).

We next correlated the eCLIP peaks with AS events across all four treatment

comparisons by identifying transcripts in which Srsf3 bound within an alternatively-spliced exon

or within 250 bp of the neighboring introns (Tables S12-S15). In agreement with our findings

above for the entire eCLIP dataset, Srsf3 exhibited differential binding in exons and surrounding

the 5' and 3' splice sites upon PDGF-AA stimulation (Fig. S5A,B). We performed an unbiased

motif enrichment analysis of Srsf3 peaks within the overlapping datasets, again revealing

different motifs between ligand treatment conditions and an enrichment of GA-rich motifs in the

+PDGF-AA samples (Fig. S5C,D).

To determine whether transcripts that are differentially bound by Srsf3 and undergo

differential AS downstream of PDGFRα signaling contribute to shared biological outputs, we

conducted GO analysis of the 149 unique transcripts from the overlapping datasets. The most

significant terms for biological process involved protein phosphorylation and deacetylation, and

RNA metabolism (Fig. 5B). Given that PI3K-mediated PDGFRα signaling is critical for

craniofacial development and regulates AS in this context (Dennison et al., 2021; Fantauzzo &

Soriano, 2014; Klinghoffer et al., 2002), we turned our focus toward GO terms associated with

this signaling pathway. We noted enrichment of PI3K-related GO terms (Fig. 5C), which

encompassed the genes *Becn1, Wdr81* and *Acap3* (Fig. 5D). Related to their roles in PI3K

signaling, each of these gene products has been shown to regulate membrane and/or endocytic

trafficking. Phosphatidylinositol 3-phosphate (PI(3)P) is a critical component of early endosomes

and is mainly generated by conversion of phosphatidylinositol (PI) by the class III PI3K complex,

which includes Beclin-1 (encoded by *Becn1*) (Wallroth & Haucke, 2018). WDR81 and Beclin-1

have been shown to interact, resulting in decreased endosomal PI(3)P synthesis, PI(3)P

turnover and early endosome conversion to late endosomes (Liu et al., 2016). Importantly, this

role of WDR81 contributes to RTK degradation (Rapiteanu et al., 2016).

Within our data, Srsf3 binding was increased in *Becn1* exon 7 upon PDGF-AA

stimulation, at an enriched motif within the overlapping datasets, and we observed a

corresponding increase in retention of adjacent intron 7 (Fig. 5D,E). As *Becn1* intron 7 contains

a premature termination codon (PTC), this event is predicted to result in nonsense-mediated

decay (NMD) in the absence of Srsf3 (Fig. 5E). Srsf3 binding was also increased in *Wdr81* exon

8 in response to PDGF-AA treatment, and our analyses revealed a corresponding increase in

excision of adjacent exon 9 (Fig. 5D,E). Because *Wdr81* exon 9 encodes two WD-repeat

domains, which are generally believed to form a β propellor structure required for protein

interactions (Li & Roberts, 2001), this AS event is predicted to result in a protein missing a

functional domain (Fig. 5E). These splicing patterns predict increased levels of Beclin-1 and

decreased levels of functional Wdr81 in the presence of Srsf3 and PDGF-AA stimulation,

resulting in augmented early endosome formation.

Srsf3 regulates early endosome size and phosphorylation of Akt downstream of PDGFRα

signaling

Given that Srsf3 differentially binds to transcripts that encode proteins involved in early

endosomal trafficking downstream of PDGFRα signaling, we first examined the formation of

Rab5-positive early endosomes (Zerial & McBride, 2001) in response to a time course of PDGF-

AA ligand stimulation in scramble versus shSrsf3 cells. As expected, Rab5 puncta size

increased from 0 min ($9.79 \times 10^{-4} \pm 1.13 \times 10^{-4}$ arbitrary units; mean $\pm$ s.e.m.) to 15 min of ligand

stimulation ($2.01 \times 10^{-3} \pm 7.20 \times 10^{-4}$ arbitrary units) in scramble cells, and significantly so by 60

min ($2.58 \times 10^{-3} \pm 9.20 \times 10^{-4}$ arbitrary units) (Fig. 6A, C-C", E-E", G-G"). However, this increase

was not observed in the absence of Srsf3 ($9.21 \times 10^{-4} \pm 1.61 \times 10^{-4}$ arbitrary units at 60 min)

(Fig. 6A, D-D", F-F", H-H"), demonstrating that Srsf3-mediated PDGFRα signaling leads to

enlarged early endosomes.

We next examined colocalization of PDGFRα with Rab5, as an estimate of receptor

levels in the early endosome. Colocalization levels increased from 0 min (0.332 ± 0.0832

Pearson's correlation coefficient (PCC); mean $\pm$ s.e.m.) to 15 min (0.429 ± 0.108 PCC) of

PDGF-AA treatment in scramble cells, then decreased to near baseline levels by 60 min (0.348

± 0.0885 PCC) (Fig. 6B, C-C", E-E", G-G") as a subset of PDGFRα homodimers are likely

trafficked to late endosomes (Rogers et al., 2022). Strikingly, shSrsf3 cells exhibited a

significant decrease in colocalization between PDGFRα and Rab5 by 60 min of ligand treatment

(0.186 ± 0.0102 PCC) (Fig. 6B, D-D", F-F", H-H"), indicating that Srsf3 activity downstream of

PDGFRα signaling results in retention of PDGFRα in early endosomes.

Finally, we previously demonstrated that rapid internalization of PDGFRα homodimers

following PDGF-AA ligand stimulation is critical for downstream AKT phosphorylation (Rogers et

al., 2022). As such, we examined phospho-Akt levels as a readout of PDGFRα activation in

early endosomes. While PDGF-AA treatment for 15 min induced a robust phospho-Akt

response in scramble cells (13.5 ± 7.39 relative induction, mean $\pm$ s.e.m.) this response was

muted in shSrsf3 cells at the same timepoint (5.73 ± 1.91) (Fig. 6I). These findings further

suggest that retention of PDGFRα in early endosomes leads to increases in downstream PI3K-

mediated Akt signaling. Collectively, our data point to a feedback loop in which PI3K/Akt-

mediated PDGFRα signaling results in the nuclear translocation of Srsf3 and the subsequent

AS of transcripts to decrease levels of proteins that promote PDGFRα trafficking out of early

endosomes (Fig. 6J).

Discussion

In this study, we confirmed our prior *in vivo* results in the mouse embryonic facial

mesenchyme (Dennison et al., 2021) that PDGFRα signaling primary regulates gene expression

via AS. PDGF-AA-dependent differential gene expression was minimal following one hour of

ligand treatment and led to increased expression of immediate early genes *Klf10, Egf3* and

Egr1, consistent with previous findings (Vasudevan et al., 2015) and in line with the enriched

GO terms of regulation of transcription and DNA binding. Alternatively, PDGF-AA-dependent

alternatively-spliced transcripts were enriched for protein kinase activity, consistent with our

prior AS findings upon disruption of PI3K-mediated PDGFRα signaling in the palatal shelf

mesenchyme (Dennison et al., 2021). Importantly, our results demonstrated a significant

dependence on the RBP Srsf3 for AS downstream of PDGFRα activation. In fact, we found that

79% of Srsf3-dependent and 97% of PDGF-AA-dependent alternatively-spliced transcripts were

responsive to both Srsf3 activity and PDGFRα signaling. As discussed above, Srsf3 is

phosphorylated downstream of multiple stimuli (Bavelloni et al., 2014; Dennison et al., 2021;

Fantauzzo & Soriano, 2014; Long et al., 2019; Zhou et al., 2012), and it is likely that these

additional inputs contributed to Srsf3-dependent AS that was non-responsive to PDGF-AA

treatment. Further, our previous mass spectrometry screen identified 11 additional RBPs

involved in AS that are phosphorylated by Akt downstream of PI3K-mediated PDGFRα signaling

in primary MEPM cells (Fantauzzo & Soriano, 2014), which may account for the small fraction of

PDGF-AA-dependent AS that was non-responsive to Srsf3 knockdown. However, our RNA-seq

results together with the phenotypic overlap of embryos with neural crest-specific ablation of

Srsf3 and mutant mouse models of *Pdgfra* and/or its ligands (Andrae et al., 2016; Dennison et

al., 2021; Ding et al., 2004; Fantauzzo & Soriano, 2014; Fredriksson et al., 2012; Klinghoffer et

al., 2002; Soriano, 1997) make a compelling case for Srsf3 serving as a critical effector of

PDGFRα signaling in the facial mesenchyme.

Here, we observed that Srsf3-dependent skipped exon events were enriched for

transcripts with a +ΔPSI (exon skipped more often in shSrsf3 samples) in the absence of PDGF-

AA stimulation, consistent with Srsf3 promoting exon inclusion. Alternatively, PDGF-AA ligand

treatment led to an increase in the percentage of Srsf3-dependent transcripts with a -ΔPSI

(exon included more often in shSrsf3 samples), suggesting that PDGFRα signaling causes

decreased exon inclusion in the presence of Srsf3. Interestingly, a recent paper found that

tethering Srsf3 downstream of the 5' splice site or upstream of the 3' splice site using MS2 stem

loops did not lead to AS of a splicing reporter (Schmok et al., 2024). However, the assay did not

test tethering within the exon and used a single minigene sequence context, and thus had the

potential to lead to false negative results (Schmok et al., 2024). Whether phosphorylation of

Srsf3 directly influences its binding to target RNAs or acts to modulate Srsf3 protein-protein

interactions which then contribute to differential RNA binding remains to be determined, though

findings from Schmok et al., 2024 may argue for the latter mechanism. Studies identifying

proteins that differentially interact with Srsf3 in response to PDGF-AA ligand stimulation are

ongoing and will shed light on these mechanisms.

This study represents the first endogenous CLIP analysis of Srsf3 in the absence of

protein tagging, and thus circumvents potential limitations with prior approaches in which

assayed RBPs were overexpressed and/or fused to another protein. Our eCLIP analyses

revealed several changes in Srsf3 transcript binding downstream of PDGF-AA stimulation,

including increased Srsf3 binding to exons and loss of Srsf3 binding to canonical CA-rich motifs.

A previous CLIP study using a stably integrated SRSF3-EGFP transgene in mouse P19 cells

determined that SRSF3 binding was enriched in exons, particularly within 100 nucleotides of the

5' and, to a lesser extent, 3' splice sites (Änkö et al., 2012), consistent with our results. This

same study identified a CA-rich canonical SRSF3 motif (Änkö et al., 2012). While such motifs

were identified here in the absence of PDGF-AA treatment, they were lost upon ligand

stimulation. Again, this shift could be due to loss of RNA binding owing to electrostatic repulsion

and/or changes in ribonucleoprotein composition and will be the subject of future studies.

Our findings additionally pointed to novel properties of exons whose inclusion is

dependent on Srsf3 in the absence of PDGFRα signaling. We demonstrated that these included

exons had a higher GC content, a lower ratio of downstream intron to exon GC content and a

decreased ratio of upstream and downstream intron to exon length. These findings are

consistent with previous results demonstrating that included exons tend to have higher GC

content than the flanking introns (Amit et al., 2012). Of note, PDGF-AA ligand stimulation

resulted in binding of Srsf3 to exons with a lower GC content, again suggesting that

phosphorylation of the RBP downstream of this signaling axis promotes exon skipping.

By cross-referencing our RNA-seq and eCLIP datasets, we showed that 24% of the

alternatively-spliced transcripts across our four treatment comparisons had an Srsf3 eCLIP

peak. As Srsf3 also has functions in the cytoplasm, such as RNA trafficking, translation and

degradation (Howard & Sanford, 2015), the additional eCLIP peaks may reflect alternate roles

for Srsf3 in RNA metabolism. Conversely, Srsf3-mediated AS may be delayed following

transcript binding in the short timeframe of our experimental design. However, the extent of

overlap that we observed is in line with previous studies correlating alternatively-spliced

transcripts upon knockdown of an RBP with endogenous eCLIP results for that same RBP,

including Rbfox2 (10%) (Moss et al., 2023) and LUC7L2 (18-26%) (Jourdain et al., 2021). The

degree to which our RNA-seq and eCLIP datasets overlapped here points to the robustness and

biological significance of our findings.

Our combined analyses demonstrated that Srsf3 binds and mediates the AS of

transcripts that encode proteins that regulate PI3K signaling and early endosomal trafficking

downstream of PDGFRα activation, including *Becn1, Wdr81* and *Acap3.* Homozygous missense

mutations in *WDR81* cause Cerebellar ataxia, impaired intellectual development, and

dysequilibrium syndrome 2 (OMIM 610185) in humans (Gulsuner et al., 2011), with some

patients exhibiting coarse facial features and strabismus (Garcias & Roth, 2007), pointing to a

critical role for this gene product in craniofacial development. Consistently, we found that related

GO terms, such as phosphatidylinositol phosphate binding and endosome to lysosome

transport, were significantly enriched among alternatively-spliced transcripts in murine

embryonic facial mesenchyme upon loss of PI3K binding to PDGFRα and/or knockdown of

Srsf3 (Dennison et al., 2021). These data further confirm that our iMEPM model system served

as a powerful platform to uncover mechanisms that are utilized during craniofacial development

in vivo.

Our subsequent *in vitro* validation studies showed that Srsf3-mediated PDGFRα

signaling results in enlarged early endosomes, retention of the receptor in these early

endosomes and increased downstream PI3K-mediated Akt signaling. Relatedly, we and others

have linked spatial organization with the propagation of PDGFRα signaling, such that rapid

internalization of the receptors into early endosomes or autophagy of the receptors are required

for maximal phosphorylation of AKT downstream of PDGFRα activation (Rogers et al., 2022;

Simpson et al., 2024). Together, these results indicate a feedback loop at play in the

craniofacial mesenchyme in which stimulation of PDGFRα homodimer signaling leads to Srsf3-

dependent AS of transcripts, a subsequent increase in the levels of proteins that maintain the

receptor in early endosomes and a corresponding decrease in the levels of proteins that

promote trafficking of the receptor to late endosomes for eventual degradation. These findings

thus represent a novel mechanism by which PDGFRα activity is maintained and propagated

within the cell. Whether similar mechanisms exist downstream of alternate RTKs or contribute to

increases in the phosphorylation of effector proteins other than Akt remain to be determined.

Taken together, our findings significantly enhance our understanding of the molecular

mechanisms by which Srsf3 activity is regulated downstream of growth factor stimulation.

Interestingly, a recent study demonstrated that retention of another shuttling SR protein, Srsf1,

exclusively in the nucleus resulted in widespread ciliary defects in mice (Haward et al., 2021).

This finding indicates that dissecting nuclear from cytoplasmic functions of SR proteins can

provide powerful insight into the physiological relevance of each. Going forward, it will be critical

to explore the *in vivo* consequences of abrogating Akt-mediated phosphorylation of Srsf3 and

comparing the resulting phenotype to those of embryos with constitutive or conditional ablation

of *Srsf3* in the neural crest lineage (Dennison et al., 2021; Jumaa et al., 1999). These

experiments are ongoing and should shed considerable light on the roles of RBP post-

translational modifications during development.

Materials and Methods

Generation of scramble and Srsf3 *shRNA iMEPM cell lines*

Immortalized mouse embryonic palatal mesenchyme (iMEPM) cells were derived from a

male $Cdkn2a^{-/-}$ embryo as previously described (Fantauzzo & Soriano, 2017). iMEPM cells were

cultured in growth medium [Dulbecco's modified Eagle's medium (Gibco, Thermo Fisher

Scientific, Waltham, MA, USA) supplemented with 50 U/mL penicillin (Gibco), 50 µg/mL

streptomycin (Gibco) and 2 mM L-glutamine (Gibco) containing 10% fetal bovine serum (FBS)

(Hyclone Laboratories Inc., Logan, UT, USA)] and grown at 37°C in 5% carbon dioxide. iMEPM

cells were tested for mycoplasma contamination using the MycoAlert Mycoplasma Detection Kit

(Lonza Group Ltd, Basel, Switzerland). Packaged lentiviruses containing pLV[shRNA]-

EGFP:T2A:Puro-U6>Scramble_shRNA (vectorID: VB010000-0009mxc) with sequence

CCTAAGGTTAAGTCGCCCTCGCTCGAGCGAGGGCGACTTAACCTTAGG or pLV[shRNA]-

EGFP:T2A:Puro-U6>mSrsf3[shRNA#1] (vectorID: VB900060-7699yyh) with sequence

GAATGATAAAGCGGTGTTTACTCGAGTAAACACCGCTTTATCATTCC were purchased from

VectorBuilder (Chicago, IL, USA). Medium containing lentivirus for a multiplicity of infection of

10 for 200,000 cells with the addition of 10 ug/mL polybrene was added to iMEPM cells for 16 h,

and cells were subsequently grown in the presence of 4 ug/mL puromycin for 10 days. Cells

with the highest GFP expression (top 20%) were isolated on a Moflo XDP 100 cell sorter

(Beckman Coulter Inc., Brea, CA, USA) and expanded. Srsf3 expression in scramble and *Srsf3*

shRNA cell lines was confirmed by western blotting. Once the stable cell lines were established,

they were split at a ratio of 1:4 for maintenance. Scramble and *Srsf3* shRNA cells were used for

experiments at passages 9-20.

Immunoprecipitation and western blotting

To induce PDGFRα signaling, cells at ~70% confluence were serum starved for 24 h in

starvation medium [Dulbecco's modified Eagle's medium (Gibco) supplemented with 50 U/mL

penicillin (Gibco), 50 μg/mL streptomycin (Gibco) and 2 mM L-glutamine (Gibco) containing

0.1% FBS (Hyclone Laboratories Inc.)] and stimulated with 10 ng/mL rat PDGF-AA ligand (R&D

Systems, Minneapolis, MN, USA) diluted from a 1.5 μg/mL working solution in 40 nM HCl

containing 0.1% BSA for the indicated length of time. When applicable, UV-crosslinking was

performed at 254 nm and 400 mJ/cm^2 using a Vari-X-Link system (UVO3 Ltd, Cambridgeshire,

UK). For immunoprecipitation of Srsf3, cells were resuspended in ice-cold CLIP lysis buffer [50

mM Tris-HCl pH 7.4, 100 mM NaCl, 1% NP-40, 0.1% SDS, 0.5% sodium deoxycholate, 1x

complete Mini protease inhibitor cocktail (Roche, MilliporeSigma, Burlington, MA, USA), 1 mM

PMSF, 10 mM NaF, 1 mM Na$_3$VO$_4$, 25 mM β-glycerophosphate]. Cleared lysates were collected

by centrifugation at 18,000 *g* for 20 min at 4°C. Anti-Srsf3 antibody (10 μg/sample) (ab73891,

Abcam, Waltham, MA, USA) was added to protein A Dynabeads (125 μL/sample) (Thermo

Fisher Scientific) washed twice in ice-cold CLIP lysis buffer and incubated for 45 min at room

temperature. Cells lysates were incubated with antibody-conjugated Dynabeads or Dynabeads

M-280 sheep anti-rabbit IgG (Thermo Fisher Scientific) washed twice in ice-cold CLIP lysis

buffer overnight at 4°C. The following day, Dynabeads were washed twice each with ice-cold

high salt wash buffer [50 mM Tris-HCl pH 7.4, 1M NaCl, 1 mM EDTA, 1% NP-40, 0.1% SDS,

0.5% sodium deoxycholate] followed by ice-cold wash buffer [20 mM Tris-HCl pH 7.4, 10 mM

$MgCl_2$, 0.2% Tween-20]. The precipitated proteins were eluted with 1x NuPAGE LDS buffer

(Thermo Fisher Scientific) containing 100 mM dithiothreitol, heated for 10 minutes at 70°C, and

separated by SDS-PAGE. For western blotting of whole-cell lysates, protein lysates were

generated by resuspending cells in ice-cold NP-40 lysis buffer (20 mM Tris-HCl pH 8, 150 mM

NaCl, 10% glycerol, 1% Nonidet P-40, 2 mM EDTA, 1x complete Mini protease inhibitor cocktail

(Roche), 1 mM PMSM, 10 mM NaF, 1 mM Na_3VO_4, 25 mM β-glycerophosphate) and collecting

cleared lysates by centrifugation at 13,400 g at 4°C for 20 min. Laemmli buffer containing 10%

β-mercaptoethanol was added to the lysates, which were heated for 5 min at 100°C. Proteins

were subsequently separated by SDS-PAGE. Western blot analysis was performed according to

standard protocols using horseradish peroxidase-conjugated secondary antibodies. Blots were

imaged using a ChemiDoc XRS+ (Bio-Rad Laboratories, Inc., Hercules, CA, USA) or a

ChemiDoc (Bio-Rad Laboratories, Inc.). The following primary antibodies were used for western

blotting: Srsf3 (1:1,000, ab73891, Abcam), Gapdh (1:50,000, 60004, Proteintech Group, Inc.,

Rosemont, IL, USA), phospho-Akt (Ser473) (1:1,000, 9271, Cell Signaling Technology, Inc.,

Danvers, MA, USA), Akt (1:1,000, 9272, Cell Signaling Technology, Inc.), horseradish

peroxidase-conjugated goat anti-mouse IgG (1:20,000, 115035003, Jackson ImmunoResearch

Inc., West Grove, PA, USA), horseradish peroxidase-conjugated goat anti-rabbit IgG (1:20,000,

111035003;, Jackson ImmunoResearch Inc.). Quantifications of signal intensity were performed

with ImageJ software (version 1.53t, National Institutes of Health, Bethesda, MD, USA). Relative

phospho-Akt levels were determined by normalizing to total Akt levels. When applicable,

statistical analyses were performed with Prism 10 (GraphPad Software Inc., San Diego, CA,

USA) using a two-tailed, ratio paired *t*-test within each cell line and a two-tailed, unpaired *t*-test

with Welch's correction between each cell line. Immunoprecipitation and western blotting

experiments were performed across three independent experiments.

RNA-sequencing and related bioinformatics analyses

8×10^5 cells obtained from each of three independent biological replicates per treatment

were frozen on liquid nitrogen and stored at -80°C. Following thawing, total RNA was

simultaneously isolated from all samples using the RNeasy Mini Kit (Qiagen, Inc., Germantown,

MD, USA) according to the manufacturer's instructions. RNA was forwarded to the University of

Colorado Cancer Center Genomics Shared Resource for quality control, library preparation, and

sequencing. RNA purity, quantity and integrity were assessed with a NanoDrop (Thermo Fisher

Scientific) and a 4200 TapeStation System (Agilent Technologies, Inc., Santa Clara, CA, USA)

prior to library preparation. Total RNA (200 ng) was used for input into the Universal Plus

mRNA-Seq kit with NuQuant (Tecan Group Ltd., Männedorf, Switzerland). Dual index, stranded

libraries were prepared and sequenced on a NovaSeq 6000 Sequencing System (Illumina, San

Diego, CA, USA) to an average depth of ~54 million read pairs (2x150 bp reads).

Raw sequencing reads were de-multiplexed using bcl2fastq (Illumina). Trimming, filtering

and adapter contamination removal was performed using BBDuk (from the BBmap v35.85 tool

suite) (Bushnell, 2015). For differential expression analysis, transcript abundance was quantified

using Salmon (v1.4.0) (Patro et al., 2017) and a decoy-aware transcriptome index prepared

using GENCODE (Frankish et al., 2019) GRCm39 M26. Gene level summaries were calculated

using tximport (Soneson et al., 2016) in R and differential expression was measured using

DESeq2 (v.1.32.0) (Love et al., 2014). Significant changes in gene-level expression are

reported for cases with adjusted $P \leq 0.05$ and fold change $|FC| \geq 2$. Spearman correlation was

computed between conditions for differentially-expressed genes. For alternative splicing

analysis, raw FASTQ were trimmed to a uniform length of 125 bp. Reads were aligned to the

mouse genome (GRCm39 Gencode M26) using STAR (v.2.7.9a) (Dobin et al., 2013). Additional

parameters for STAR: --outFilterType BySJout --outFilterMismatchNmax 10 --

outFilterMismatchNoverLmax 0.04 --alignEndsType EndToEnd --runThreadN 16 --

alignSJDBoverhangMin 4 --alignIntronMax 300000 --alignSJoverhangMin 8 --alignIntronMin 20.

All splice junctions detected in at least 1 read from the first pass alignment were used in a

second pass alignment, per software documentation. Alternative splicing events were detected

using rMATS (v4.0.2, default parameters plus '—cstat 0.0001') (Shen et al. 2014). Reads

mapping to the splice junction as well as those mapping to the exon body were used in

downstream analyses. Detected events were compared between treatment groups and

considered significant with false discovery rate (FDR) $\leq$ 0.05, a difference in percent spliced in

(|ΔPSI|) $\geq$ 0.05 and event detection in at least 10 reads in either condition. Raw read pairs,

trimmed read pairs for Salmon input, Salmon mapping rate per sample, trimmed read pairs (125

bp) for STAR input and STAR unique mapping rate can be found in Table S1. Gene ontology

analysis was performed with various libraries from the Enrichr gene list enrichment analysis tool

(Chen et al., 2013; Kuleshov et al., 2016) and terms with P < 0.05 were considered significant.

qPCR

Total RNA was isolated using the RNeasy mini kit (Qiagen, Germantown, MD, USA)

according to the manufacturer's instructions. First-strand cDNA was synthesized using a ratio of

2:1 random primers:oligo (dT) primer and SuperScript II RT (Invitrogen, Thermo Fisher

Scientific) according to the manufacturer's instructions. All reactions were performed with 1×

ThermoPol buffer [0.02 M Tris (pH 8.8), 0.01 M KCl, 0.01 M $(NH_4)_2SO_4$, 2 mM $MgSO_4$ and 0.1%

Triton X-100], 200 μM dNTPs, 200 nM primers (Integrated DNA Technologies, Inc., Coralville,

IA, USA), 0.6 U Taq polymerase and 1 μg cDNA in a 25 μL reaction volume. The primers used

can be found in Table S16. The following PCR protocol was used for *Arhgap12*: step1, 3 min at

94°C; step 2, 30 s at 94°C; step 3, 30 s at 47°C; step 4, 30 s at 72°C; repeat steps 2-4 for 34

cycles; and step 5, 5 min at 72°C. The following PCR protocol was used for *Cep55*: step1, 3 min

at 94°C; step 2, 30 s at 94°C; step 3, 30 s at 48°C; step 4, 30 s at 72°C; repeat steps 2-4 for 34

cycles; and step 5, 5 min at 72°C. Two-thirds of total PCR products were electrophoresed on a

2% agarose/TBE gel containing ethidium bromide and photographed on an Aplegen Omega

Fluor Gel Documentation System (Aplegen Inc., Pleasanton, CA, USA). Quantifications of band

intensity were performed with ImageJ software (version 1.53t, National Institutes of Health). The

PSI was calculated independently for each sample as the percentage of the larger isoform

divided by the total abundance of all isoforms within the given gel lane. Statistical analyses were

performed with Prism 10 (GraphPad Software) using a two-tailed, unpaired *t*-test with Welch's

correction. qPCR reactions were performed using three biological replicates.

Enhanced UV-crosslinking and immunoprecipitation and related bioinformatics analyses

Experiments were performed as previously described in biological duplicates (Van

Nostrand et al., 2016, 2017). Briefly, 2 million cells per treatment were serum starved and

treated with 10 ng/mL PDGF-AA as described above. Cells were subsequently UV-crosslinked

at 254 nm and 400 mJ/cm^2, scraped in 1x phosphate buffered saline (PBS) and transferred to

1.5 mL Eppendorf tubes, at which point excess PBS was removed and cells were frozen on

liquid nitrogen and stored at -80°C. Following thawing, cells were lysed in ice-cold CLIP lysis

buffer, sonicated by BioRuptor (Diagenode, Denville, NJ, USA) and treated with RNase I

(Thermo Fisher Scientific). 2% of lysates were set aside as size-matched input samples. Srsf3-

RNA complexes were immunoprecipitated with anti-Srsf3 antibody (10 μg per sample)

(ab73891, Abcam) conjugated to protein A Dynabeads (Thermo Fisher Scientific). IP samples

were washed and dephosphorylated with FastAP (New England Biolabs, Ipswich, MA, USA)

and T4 PNK (New England Biolabs). IP samples underwent on-bead ligation of barcoded RNA

adapters (/5phos/rArGrArUrCrGrGrArArGrArGrCrGrUrCrGrUrG/3SpC3/) to the 3' end using T4

RNA ligase (New England Biolabs). Following elution, protein-RNA complexes were run on 4-

12% Bis-Tris 1.5 mm gels (Thermo Fisher Scientific) and transferred onto nitrocellulose

membranes. The 20-75 kDa region was excised and digested with proteinase K (New England

Biolabs). RNA was isolated with acid phenol/chloroform/isoamyl alcohol (pH 6.5) (Thermo

Fisher Scientific), reverse transcribed with Superscript III (Thermo Fisher Scientific) and treated

with ExoSAP-IT (Affymetrix, Thermo Fisher Scientific) to remove excess primers and

unincorporated nucleotides. Samples underwent 3' ligation of barcoded DNA adapters

(/5Phos/NNNNNNNNNNAGATCGGAAGAGCACACGTCTG/3SpC3/), clean-up with Dynabeads

MyOne Silane (Thermo Fisher Scientific) and qPCR to determine the appropriate number of

PCR cycles. Libraries were then amplified with Q5 PCR mix (New England Biolabs) for a total of

16-25 cycles. Libraries were forwarded to the University of Colorado Cancer Center Genomics

Shared Resource for quality control and sequencing. Sample integrity was assessed with a

D1000 ScreenTape System (Agilent Technologies, Inc.) and sequenced on a NovaSeq 6000

Sequencing System (Illumina) to an average depth of ~20 million read pairs (2x150 bp reads).

Raw sequencing reads were de-multiplexed using bcl2fastq (Illumina). Adapters were

trimmed using cutadapt (v.1.18) (Martin, 2011). Trimmed reads were quality filtered and

collapsed using a combination of FASTX-Toolkit (v.0.0.14)

(http://hannonlab.cshl.edu/fastx_toolkit), seqtk (v.1.3-r106) (https://github.com/lh3/seqtk) and

custom scripts. After collapsing the reads, unique molecular identifiers were removed using

seqtk. STAR index for repetitive elements was created using repetitive sequences from

msRepDB (Liao et al., 2022). Reads ≥ 18 nt were mapped to the repetitive elements using

STAR (v.2.7.9a) (Dobin et al. 2013). Reads unmapped to the repetitive elements were mapped

to the mouse genome (GRCm39 Gencode M26) using STAR (v.2.7.9a) with parameters

alignEndsType: EndtoEnd and outFilterMismatchNoverReadLmax: 0.04. Peaks were called

using omniCLIP (v.0.20) (Drewe-Boss et al., 2018) with the foreground penalty (--fg_pen)

parameter set to 5. Peaks were annotated and motif analyses performed using RCAS (v.1.19.0)

(Uyar et al., 2017) and custom R script. For visualization purposes, bigWig files were created

from bam files using deepTools (v.3.5.5) (Ramírez et al., 2016). Peaks were visualized in

Integrative Genomics Viewer (v.2.13.0) (Robinson et al., 2011). Intron and exon features were

calculated using Matt (v.1.3.1) (Gohr & Irimia, 2019), and statistical analyses were performed

using a Mann-Whitney U test. For overlap of eCLIP peaks and alternative splicing events, peak

coordinates were taken from omniCLIP bed files and alternative splicing coordinates were taken

from rMATS output. Overlapping coordinates from alternative splicing events were defined

following the rMAPS default values (Park et al., 2016). Overlap was calculated using valr

(v.0.6.4) (Riemondy et al., 2017) and custom R scripts. Raw read pairs, trimmed read pairs,

collapsed reads, reads after removing repetitive elements, mapped reads, peaks and annotated

peaks can be found in Table S7. Gene ontology analysis was performed with various libraries

from the Enrichr gene list enrichment analysis tool (Chen et al., 2013; Kuleshov et al., 2016) and

terms with $P < 0.05$ were considered significant.

Immunofluorescence analysis

Cells were seeded onto glass coverslips at ~40% confluency per 24-well plate well in

iMEPM growth medium. After 24 h, cells were serum starved and treated with 10 ng/mL PDGF-

AA as described above. Cells were fixed in 4% paraformaldehyde (PFA) in PBS with 0.1%

Triton X-100 for 10 min and washed in PBS. Cells were blocked for 1 h in 5% normal donkey

serum (Jackson ImmunoResearch Inc.) in PBS and incubated overnight at 4°C in primary

antibody diluted in 1% normal donkey serum in PBS. After washing in PBS, cells were

incubated in Alexa Fluor 488-conjugated donkey anti-rabbit secondary antibody (1:1,000;

A21206; Invitrogen) or Alexa Fluor 546-conjugated donkey anti-mouse secondary antibody

(1:1,000; A10036; Invitrogen) diluted in 1% normal donkey serum in PBS with 2 µg/ml DAPI

(Sigma-Aldrich, St. Louis, MO, USA) for 1 h. Cells were mounted in VECTASHIELD HardSet

Antifade Mounting Medium (Vector Laboratories, Inc., Burlingame, CA, USA) and photographed

using an Axiocam 506 mono digital camera (Carl Zeiss Microscopy LLC, White Plains, NY,

USA) fitted onto an Axio Observer 7 fluorescence microscope (Carl Zeiss Microscopy LLC) with

the 63x oil objective with a numerical aperture of 1.4 at room temperature. The following

antibodies were used for immunofluorescence analysis: Rab5 (1:200, C8B1, 3547, Cell

Signaling Technology Inc.), PDGFRα (1:20, AF1062, R&D Systems). For assessment of Rab5

puncta size and colocalization experiments, three independent trials, or biological replicates,

were performed. For each biological replicate, 20 technical replicates consisting of individual

cells were imaged with Z-stacks (0.24 µm between Z-stacks with a range of 1–6 Z-stacks) per

timepoint. Images were deconvoluted using ZEN Blue software (Carl Zeiss Microscopy LLC)

using the 'Better, fast (Regularized Inverse Filter)' setting. Extended depth of focus was applied

to Z-stacks using ZEN Blue software (Carl Zeiss Microscopy LLC) to generate images with the

maximum depth of field. For assessment of Rab5 puncta size, images were converted to 8-bit

using Fiji software (version 2.14.0/1.54f). Images were subsequently converted to a mask and

watershed separation was applied. A region of interest (ROI) was drawn around each Rab5-

positive cell and particles were analyzed per cell using the "analyze particles" function. For

colocalization measurements, an ROI was drawn around each PDGFRα-positive cell in the

corresponding Cy3 (marker) channel using Fiji. For each image with a given ROI, the Cy3

channel and the EGFP channel were converted to 8-bit images. Colocalization was measured

using the Colocalization Threshold function, where the rcoloc value [Pearson's correlation

coefficient (PCC)] was used in statistical analysis. Statistical analyses were performed on the

average values from each biological replicate with Prism 10 (GraphPad Software Inc.) using a

two-way ANOVA followed by uncorrected Fisher's LSD test.

Acknowledgements

We are grateful to Jessica Johnston and Erin Binne for technical assistance, and Drs. Allison

Swain, Justin Roberts and Aaron Johnson at the University of Colorado Anschutz Medical

Campus for advice on eCLIP experiments. Cell sorting was performed at the University of

Colorado Cancer Center Flow Cytometry Shared Resource with assistance from Dr. Dmitry

Baturin. RNA-seq and eCLIP sequencing experiments were performed at University of Colorado

Cancer Center Genomics Shared Resource. We thank members of the Fantauzzo laboratory for

their critical comments on the manuscript.

Competing Interests

The authors declare no competing or financial interests.

Author contributions

Conceptualization: TEF, MS, NM, KAF; Methodology: TEF, MS, NM, KAF; Formal analysis:

TEF, MS, EDL, KAF; Investigation: TEF, MS, EDL; Writing – Original Draft: TEF, KAF; Writing –

Review & Editing: MS, EDL, NM; Visualization: TEF, MS, KAF; Supervision: NM, KAF; Project

administration: KAF; Funding acquisition: TEF, KAF.

Funding

This work was supported by National Institutes of Health grants R01DE030864 (to K.A.F.),

R35GM147025 (to N.M.), the University of Colorado Anschutz Medical Campus RNA

Bioscience Initiative (to N.M. and M.P.S.) and F31DE032252 (to T.E.F.). The Flow Cytometry

Shared Resource and Genomics Shared Resource are supported by National Institutes of

Health grant P30CA046934.

Data Availability

The eCLIP and RNA-sequencing datasets generated during this study have been deposited in

GEO under SuperSeries accession number GSE263170. Custom analysis scripts will be

provided by Dr. Larson through GitHub.

References

Akerman, M., David-Eden, H., Pinter, R. Y., & Mandel-Gutfreund, Y. (2009). A computational

approach for genome-wide mapping of splicing factor binding sites. *Genome Biology*,

10(3), R30. https://doi.org/10.1186/gb-2009-10-3-r30

Amit, M., Donyo, M., Hollander, D., Goren, A., Kim, E., Gelfman, S., Lev-Maor, G., Burstein, D.,

Schwartz, S., Postolsky, B., Pupko, T., & Ast, G. (2012). Differential GC Content

between Exons and Introns Establishes Distinct Strategies of Splice-Site Recognition.

Cell Reports, *1*(5), 543–556. https://doi.org/10.1016/j.celrep.2012.03.013

Andrae, J., Gouveia, L., Gallini, R., He, L., Fredriksson, L., Nilsson, I., Johansson, B. R.,

Eriksson, U., & Betsholtz, C. (2016). A role for PDGF-C/PDGFRα signaling in the

formation of the meningeal basement membranes surrounding the cerebral cortex.

Biology Open, *5*(4), 461–474. https://doi.org/10.1242/bio.017368

Änkö, M.-L., Müller-McNicoll, M., Brandl, H., Curk, T., Gorup, C., Henry, I., Ule, J., &

Neugebauer, K. M. (2012). The RNA-binding landscapes of two SR proteins reveal

unique functions and binding to diverse RNA classes. *Genome Biology*, *13*(3), R17.

https://doi.org/10.1186/gb-2012-13-3-r17

Bavelloni, A., Piazzi, M., Faenza, I., Raffini, M., D'Angelo, A., Cattini, L., Cocco, L., & Blalock,

W. L. (2014). Prohibitin 2 represents a novel nuclear AKT substrate during all- *trans*

retinoic acid–induced differentiation of acute promyelocytic leukemia cells. *The FASEB*

Journal, *28*(5), 2009–2019. https://doi.org/10.1096/fj.13-244368

Bebee, T. W., Park, J. W., Sheridan, K. I., Warzecha, C. C., Cieply, B. W., Rohacek, A. M.,

Xing, Y., & Carstens, R. P. (2015). The splicing regulators Esrp1 and Esrp2 direct an

epithelial splicing program essential for mammalian development. *eLife*, *4*, e08954.

https://doi.org/10.7554/eLife.08954

Bushnell, B. (2015). *BBMap*. Https://Sourceforge.Net/Projects/Bbmap/.

https://sourceforge.net/projects/bbmap/

Chen, E. Y., Tan, C. M., Kou, Y., Duan, Q., Wang, Z., Meirelles, G. V., Clark, N. R., & Ma'ayan,

A. (2013). Enrichr: Interactive and collaborative HTML5 gene list enrichment analysis

tool. *BMC Bioinformatics*, *14*, 128. https://doi.org/10.1186/1471-2105-14-128

Choi, S. J., Marazita, M. L., Hart, P. S., Sulima, P. P., Field, L. L., McHenry, T. G., Govil, M.,

Cooper, M. E., Letra, A., Menezes, R., Narayanan, S., Mansilla, M. A., Granjeiro, J. M.,

Vieira, A. R., Lidral, A. C., Murray, J. C., & Hart, T. C. (2009). The PDGF-C regulatory

region SNP rs28999109 decreases promoter transcriptional activity and is associated

with CL/P. *European Journal of Human Genetics*, *17*(6), 774–784.

https://doi.org/10.1038/ejhg.2008.245

Cibi, D. M., Mia, M. M., Guna Shekeran, S., Yun, L. S., Sandireddy, R., Gupta, P., Hota, M.,

Sun, L., Ghosh, S., & Singh, M. K. (2019). Neural crest-specific deletion of Rbfox2 in

mice leads to craniofacial abnormalities including cleft palate. *eLife*, *8*, e45418.

https://doi.org/10.7554/eLife.45418

Dennison, B. J. C., Larson, E. D., Fu, R., Mo, J., & Fantauzzo, K. A. (2021). Srsf3 mediates

alternative RNA splicing downstream of PDGFRα signaling in the facial mesenchyme.

Development, *148*(14), dev199448. https://doi.org/10.1242/dev.199448

Ding, H., Wu, X., Boström, H., Kim, I., Wong, N., Tsoi, B., O'Rourke, M., Koh, G. Y., Soriano, P.,

Betsholtz, C., Hart, T. C., Marazita, M. L., Field, L. L., Tam, P. P. L., & Nagy, A. (2004).

A specific requirement for PDGF-C in palate formation and PDGFR-α signaling. *Nature*

Genetics, *36*(10), 1111–1116. https://doi.org/10.1038/ng1415

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,

M., & Gingeras, T. R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics,*

29(1), 15-21. https://doi.org/10.1093/bioinformatics/bts635

Drewe-Boss, P., Wessels, H.-H., & Ohler, U. (2018). omniCLIP: Probabilistic identification of

protein-RNA interactions from CLIP-seq data. *Genome Biology*, *19*(1), 183.

https://doi.org/10.1186/s13059-018-1521-2

Fantauzzo, K. A., & Soriano, P. (2014). PI3K-mediated PDGFR signaling regulates survival and

proliferation in skeletal development through p53-dependent intracellular pathways.

Genes & Development, *28*(9), 1005–1017. https://doi.org/10.1101/gad.238709.114

Fantauzzo, K. A., & Soriano, P. (2017). Generation of an immortalized mouse embryonic palatal

mesenchyme cell line. *PLOS ONE*, *12*(6), e0179078.

https://doi.org/10.1371/journal.pone.0179078

Forman, T. E., Dennison, B. J. C., & Fantauzzo, K. A. (2021). The Role of RNA-Binding Proteins

in Vertebrate Neural Crest and Craniofacial Development. *Journal of Developmental*

Biology, 19.

Frankish, A., Diekhans, M., Ferreira, A. M., Johnson, R., Jungreis, I., Loveland, J., Mudge, J.

M., Sisu, C., Wright, J., Armstrong, J., Barnes, I., Berry, A., Bignell, A., Carbonell Sala,

S., Chrast, J., Cunningham, F., Di Domenico, T., Donaldson, S., Fiddes, I. T., … Flicek,

P. (2019). GENCODE reference annotation for the human and mouse genomes. *Nucleic*

Acids Research, *47*(D1), D766-763. https://doi.org/10.1093/nar/gky955

Fredriksson, L., Nilsson, I., Su, E. J., Andrae, J., Ding, H., Betsholtz, C., Eriksson, U., &

Lawrence, D. A. (2012). Platelet-Derived Growth Factor C Deficiency in C57BL/6 Mice

Leads to Abnormal Cerebral Vascularization, Loss of Neuroependymal Integrity, and

Ventricular Abnormalities. *The American Journal of Pathology*, *180*(3), 1136–1144.

https://doi.org/10.1016/j.ajpath.2011.12.006

Fu, X.-D., & Ares, M. (2014). Context-dependent control of alternative splicing by RNA-binding

proteins. *Nature Reviews Genetics*, *15*(10), 689–701. https://doi.org/10.1038/nrg3778

Fu, X.-D., & Maniatis, T. (1990). Factor required for mammalian spliceosome assembly is

localized to discrete regions in the nucleus. *Nature, 343*(6257), 437–441.

https://doi.org/10.1038/343437a0

Garcias, G. D. L., & Roth, M. D. G. M. (2007). A Brazilian family with quadrupedal gait, severe

mental retardation, coarse facial characteristics, and hirsutism. *International Journal of*

Neuroscience, 117(7), 927–933. https://doi.org/10.1080/00207450600910721

Gohr, A., & Irimia, M. (2019). *Matt*: Unix tools for alternative splicing analysis. *Bioinformatics,*

35(1), 130–132. https://doi.org/10.1093/bioinformatics/bty606

Gulsuner, S., Tekinay, A. B., Doerschner, K., Boyaci, H., Bilguvar, K., Unal, H., Ors, A., Onat, O.

E., Atalar, E., Basak, A. N., Topaloglu, H., Kansu, T., Tan, M., Tan, U., Gunel, M., &

Ozcelik, T. (2011). Homozygosity mapping and targeted genomic sequencing reveal the

gene responsible for cerebellar hypoplasia and quadrupedal locomotion in a

consanguineous kindred. *Genome Research, 21*(12), 1995-2003.

https://doi.org/10.1101/gr.126110.111.

Haward, F., Maslon, M. M., Yeyati, P. L., Bellora, N., Hansen, J. N., Aitken, S., Lawson, J., von

Kriegsheim, A., Wachten, D., Mill, P., Adams, I. R., & Caceres, J. F. (2021). Nucleo-

cytoplasmic shuttling of splicing factor SRSF1 is required for development and cilia

function. *eLife, 10*, e65104. https://doi.org/10.7554/eLife.65104

He, F., & Soriano, P. (2013). A critical role for PDGFRalpha signaling in medial nasal process

development. *PLoS Genetics, 9*(9), e1003851.

https://doi.org/10.1371/journal.pgen.1003851

Howard, J. M., & Sanford, J. R. (2015). The RNAissance family: SR proteins as multifaceted

regulators of gene expression: *Wiley Interdisciplinary Reviews: RNA, 6*(1), 93–110.

https://doi.org/10.1002/wrna.1260

Huang, Y., Yario, T. A., & Steitz, J. A. (2004). A molecular link between SR protein

dephosphorylation and mRNA export. *Proceedings of the National Academy of*

Sciences, *101*(26), 9666–9670. https://doi.org/10.1073/pnas.0403533101

Jourdain, A. A., Begg, B. E., Mick, E., Shah, H., Calvo, S. E., Skinner, O. S., Sharma, R., Blue,

S. M., Yeo, G. W., Burge, C. B., & Mootha, V. K. (2021). Loss of LUC7L2 and U1 snRNP

subunits shifts energy metabolism from glycolysis to OXPHOS. *Molecular Cell*, *81*(9),

1905-1919.e12. https://doi.org/10.1016/j.molcel.2021.02.033

Jumaa, H., Wei, G., & Nielsen, P. J. (1999). Blastocyst formation is blocked in mouse embryos

lacking the splicing factor SRp20. *Current Biology*, *9*(16), 899–902.

https://doi.org/10.1016/S0960-9822(99)80394-7

Klinghoffer, R. A., Hamilton, T. G., Hoch, R., & Soriano, P. (2002). An Allelic Series at the

PDGFaR Locus Indicates Unequal Contributions of Distinct Signaling Pathways During

Development. *Developmental Cell*, *2*, 103–113.

Krainer, R., Mayeda, A., & Kozak, D. (1991). Functional Expression of Cloned Human Splicing

Factor SF2: Homology to RNA-Binding Proteins, UI 70K, and Drosophila Splicing

Regulators. *Cell*, *66*, 383–394.

Kuleshov, M. V., Jones, M. R., Rouillard, A. D., Fernandez, N. F., Duan, Q., Wang, Z., Koplev,

S., Jenkins, S. L., Jagodnik, K. M., Lachmann, A., McDermott, M. G., Monteiro, C. D.,

Gundersen, G. W., & Ma'ayan, A. (2016). Enrichr: A comprehensive gene set

enrichment analysis web server 2016 update. *Nucleic Acids Research*, *44*(W1), W90–

W97. https://doi.org/10.1093/nar/gkw377

Lee, S., Sears, M. J., Zhang, Z., Li, H., Salhab, I., Krebs, P., Xing, Y., Nah, H.-D., Williams, T., &

Carstens, R. P. (2020). Cleft lip and cleft palate in *Esrp1* knockout mice is associated

with alterations in epithelial-mesenchymal crosstalk. *Development*, *147*(21), dev187369.

https://doi.org/10.1242/dev.187369

Li, D., & Roberts, R. (2001). WD-repeat proteins: Structure characteristics, biological function,

and their involvement in human diseases. *Cellular and Molecular Life Sciences*, *58*.

Liao, X., Hu, K., Salhi, A., Zou, Y., Wang, J., & Gao, X. (2022). msRepDB: A comprehensive

repetitive sequence database of over 80 000 species. *Nucleic Acids Research*, *50*(D1),

D236–D245. https://doi.org/10.1093/nar/gkab1089

Licatalosi, D. D., & Darnell, R. B. (2010). RNA processing and its regulation: Global insights into

biological networks. *Nature Reviews Genetics*, *11*(1), 75–87.

https://doi.org/10.1038/nrg2673

Liu, K., Jian, Y., Sun, X., Yang, C., Gao, Z., Zhang, Z., Liu, X., Li, Y., Xu, J., Jing, Y., Mitani, S.,

He, S., & Yang, C. (2016). Negative regulation of phosphatidylinositol 3-phosphate

levels in early-to-late endosome conversion. *Journal of Cell Biology, 212*(2), 181-198.

https://doi.org/10.1083/jcb.201506081.

Long, Y., Sou, W. H., Yung, K. W. Y., Liu, H., Wan, S. W. C., Li, Q., Zeng, C., Law, C. O. K.,

Chan, G. H. C., Lau, T. C. K., & Ngo, J. C. K. (2019). Distinct mechanisms govern the

phosphorylation of different SR protein splicing factors. *Journal of Biological Chemistry*,

294(4), 1312–1327. https://doi.org/10.1074/jbc.RA118.003392

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550.

https://doi.org/10.1186/s13059-014-0550-8

Mai, C. T., Isenburg, J. L., Canfield, M. A., Meyer, R. E., Correa, A., Alverson, C. J., Lupo, P. J.,

Riehle-Colarusso, T., Cho, S. J., Aggarwal, D., Kirby, R. S., & National Birth Defects

Prevention Network. (2019). National population-based estimates for major birth defects,

2010–2014. *Birth Defects Research*, *111*(18), 1420–1435.

https://doi.org/10.1002/bdr2.1589

Manning, B. D., & Cantley, L. C. (2007). AKT/PKB Signaling: Navigating Downstream. *Cell*,

129(7), 1261–1274. https://doi.org/10.1016/j.cell.2007.06.009

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing

reads. *EMBnet.Journal*, *17*(1), 10. https://doi.org/10.14806/ej.17.1.200

Moss, N. D., Wells, K. L., Theis, A., Kim, Y.-K., Spigelman, A. F., Liu, X., MacDonald, P. E., &

Sussel, L. (2023). Modulation of insulin secretion by RBFOX2-mediated alternative

splicing. *Nature Communications*, *14*(1), 7732. https://doi.org/10.1038/s41467-023-

43605-4

Pan, Q., Shai, O., Lee, L. J., Frey, B. J., & Blencowe, B. J. (2008). Deep surveying of alternative

splicing complexity in the human transcriptome by high-throughput sequencing. *Nature*

Genetics, *40*(12), 1413–1415. https://doi.org/10.1038/ng.259

Park, J. W., Jung, S., Rouchka, E. C., Tseng, Y-T., & Xing, Y. (2016). rMAPS: RNA map

analysis and plotting server for alternative exon regulation. *Nucleic Acids Research*

44(W1), 333-338. https://doi.org/10.1093/nar/gkw410

Patro, R., Duggal, G., Love, M. I., Irizarry, R. A., & Kingsford, C. (2017). Salmon provides fast

and bias-aware quantification of transcript expression. *Nature Methods*, *14*(4), 417–419.

https://doi.org/10.1038/nmeth.4197

Ramírez, F., Ryan, D. P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A. S., Heyne, S.,

Dündar, F., & Manke, T. (2016). deepTools2: A next generation web server for deep-

sequencing data analysis. *Nucleic Acids Research*, *44*(W1), W160–W165.

https://doi.org/10.1093/nar/gkw257

Rapiteanu, R., Davis, L. J., Williamson, J. C., Timms, R. T., Paul Luzio, J., & Lehner, P. J.

(2016). A Genetic Screen Identifies a Critical Role for the WDR81-WDR91 Complex in

the Trafficking and Degradation of Tetherin. *Traffic*, *17*(8), 940–958.

https://doi.org/10.1111/tra.12409

Rattanasopha, S., Tongkobpetch, S., Srichomthong, C., Siriwan, P., Suphapeetiporn, K., &

Shotelersuk, V. (2012). PDGFRa mutations in humans with isolated cleft palate.

879   *European Journal of Human Genetics*, *20*(10), 1058–1062.

880   https://doi.org/10.1038/ejhg.2012.55

881 Riemondy, K. A., Sheridan, R. M., Gillen, A., Yu, Y., Bennett, C. G., & Hesselberth, J. R. (2017).

882   valr: Reproducible genome interval analysis in R. *F1000Research*, *6*, 1025.

883   https://doi.org/10.12688/f1000research.11997.1

884 Robinson, J. T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E. S., Getz, G., &

885   Mesirov, J. P. (2011). Integrative genomics viewer. *Nature Biotechnology*, *29*(1), 24–26.

886   https://doi.org/10.1038/nbt.1754

887 Rogers, M. A., Campaña, M. B., Long, R., & Fantauzzo, K. A. (2022). PDGFR dimer-specific

888   activation, trafficking and downstream signaling dynamics. *Journal of Cell Science*,

889   *135*(17), jcs259686. https://doi.org/10.1242/jcs.259686

890 Schmok, J. C., Jain, M., Street, L. A., Tankka, A. T., Schafer, D., Her, H.-L., Elmsaouri, S.,

891   Gosztyla, M. L., Boyle, E. A., Jagannatha, P., Luo, E.-C., Kwon, E. J., Jovanovic, M., &

892   Yeo, G. W. (2024). Large-scale evaluation of the ability of RNA-binding proteins to

893   activate exon inclusion. *Nature Biotechnology*. https://doi.org/10.1038/s41587-023-

894   02014-0

895 Scotti, M. M., & Swanson, M. S. (2016). RNA mis-splicing in disease. *Nature Reviews Genetics*,

896   *17*(1), 19–32. https://doi.org/10.1038/nrg.2015.3

897 Shen, H., & Green, M. R. (2006). RS domains contact splicing signals and promote splicing by a

898   common mechanism in yeast through humans. *Genes & Development*, *20*(13), 1755–

899   1765. https://doi.org/10.1101/gad.1422106

900 Shen, S., Park, J. W., Lu, Z., Lin, L., Henry, M. D., Wu, Y. N., Zhou, Q., & Xing, Y. (2014).

901   rMATS: Robust and flexible detection of differential alternative splicing from replicate

902   RNA-Seq data. *Proceedings of the National Academy of Sciences*, *111*(51), E5593–

903   E5601. https://doi.org/10.1073/pnas.1419161111

Shin, C., Feng, Y., & Manley, J. L. (2004). Dephosphorylated SRp38 acts as a splicing

repressor in response to heat shock. *Nature*, *427*(6974), 553–558.

https://doi.org/10.1038/nature02288

Simpson, J. E., Muir, M. T., Lee, M., Naughton, C., Gilbert, N., Pollard, S. M., & Gammoh, N.

(2024). Autophagy supports PDGFRA-dependent brain tumor development by

enhancing oncogenic signaling. *Developmental Cell*, S1534580723006214.

https://doi.org/10.1016/j.devcel.2023.11.023

Soneson, C., Love, M. I., & Robinson, M. D. (2016). Differential analyses for RNA-seq:

Transcript-level estimates improve gene-level inferences [version 2; referees: 2

approved]. *F1000Research*, *4*, 1521. https://doi.org/10.12688/F1000RESEARCH.7563.2

Soriano, P. (1997). The PDGFα receptor is required for neural crest cell development and for

normal patterning of the somites. *Development*, *124*, 2691–2700.

Tallquist, M. D., & Soriano, P. (2003). Cell autonomous requirement for PDGFRα in populations

of cranial and cardiac neural crest cells. *Development*, *130*(3), 507–518.

https://doi.org/10.1242/dev.00241

Uyar, B., Yusuf, D., Wurmus, R., Rajewsky, N., Ohler, U., & Akalin, A. (2017). RCAS: An RNA

centric annotation system for transcriptome-wide regions of interest. *Nucleic Acids*

Research, *45*(10), e91–e91. https://doi.org/10.1093/nar/gkx120

Van Nostrand, E. L., Nguyen, T. B., Gelboin-Burkhart, C., Wang, R., Blue, S. M., Pratt, G. A.,

Louie, A. L., & Yeo, G. W. (2017). Robust, Cost-Effective Profiling of RNA Binding

Protein Targets with Single-end Enhanced Crosslinking and Immunoprecipitation

(seCLIP). In Y. Shi (Ed.), *mRNA Processing* (Vol. 1648, pp. 177–200). Springer New

York. https://doi.org/10.1007/978-1-4939-7204-3_14

Van Nostrand, E. L., Pratt, G. A., Shishkin, A. A., Gelboin-Burkhart, C., Fang, M. Y.,

Sundararaman, B., Blue, S. M., Nguyen, T. B., Surka, C., Elkins, K., Stanton, R., Rigo,

F., Guttman, M., & Yeo, G. W. (2016). Robust transcriptome-wide discovery of RNA-

930   binding protein binding sites with enhanced CLIP (eCLIP). *Nature Methods*, *13*(6), 508–

931   514. https://doi.org/10.1038/nmeth.3810

932 Vasudevan, H. N., Mazot, P., He, F., & Soriano, P. (2015). Receptor tyrosine kinases modulate

933   distinct transcriptional programs by differential usage of intracellular pathways. *eLife*, *4*,

934   e07186. https://doi.org/10.7554/eLife.07186

935 Wallroth, A., & Haucke, V. (2018). Phosphoinositide conversion in endocytosis and the

936   endolysosomal system. *Journal of Biological Chemistry*, *293*(5), 1526–1535.

937   https://doi.org/10.1074/jbc.R117.000629

938 Wang, E. T., Sandberg, R., Luo, S., Khrebtukova, I., Zhang, L., Mayr, C., Kingsmore, S. F.,

939   Schroth, G. P., & Burge, C. B. (2008). Alternative isoform regulation in human tissue

940   transcriptomes. *Nature*, *456*(7221), 470–476. https://doi.org/10.1038/nature07509

941 Xiao, S.-H., & Manley, J. (1997). Phosphorylation of the ASF/SF2 RS domain affects both

942   protein-protein and protein-RNA interactions and is necessary for splicing. *Genes &*

943   *Development*, 12.

944 Zahler, A. M., Neugebauer, K. M., Lane, W. S., & Roth, M. B. (1993). Distinct Functions of SR

945   Proteins in Alternative pre-mRNA Splicing. *Science*, *260*(5105), 219–222.

946   https://doi.org/10.1126/science.8385799

947 Zerial, M., & McBride, H. (2001). Rab proteins as membrane organizers. *Nature Reviews*

948   *Molecular Cell Biology*, *2*(2), 107–117. https://doi.org/10.1038/35052055

949 Zhou, Z., & Fu, X.-D. (2013). Regulation of splicing by SR proteins and SR protein-specific

950   kinases. *Chromosoma*, *122*(3), 191–207. https://doi.org/10.1007/s00412-013-0407-z

951 Zhou, Z., Qiu, J., Liu, W., Zhou, Y., Plocinik, R. M., Li, H., Hu, Q., Ghosh, G., Adams, J. A.,

952   Rosenfeld, M. G., & Fu, X.-D. (2012). The Akt-SRPK-SR Axis Constitutes a Major

953   Pathway in Transducing EGF Signaling to Regulate Alternative Splicing in the Nucleus.

954   *Molecular Cell*, *47*(3), 422–433. https://doi.org/10.1016/j.molcel.2012.05.014

Figure 1: PDGFRα signaling for one hour minimally affects gene expression. (A)

Schematic of RNA-seq experimental design. iMEPM cells were transduced to stably express a

scramble shRNA (scramble) or shRNA targeting the 3' UTR of *Srsf3* (shSrsf3). iMEPM cells

expressing either scramble or shSrsf3 were left unstimulated or stimulated with 10 ng/mL

PDGF-AA for 1 hour and RNA was isolated for RNA-seq analysis. **(B)** Western blot (WB)

analysis of whole-cell lysates from scramble and shSrsf3 cell lines with anti-Srsf3 and anti-

Gapdh antibodies. The percentage of Srsf3 expression normalized to Gapdh expression is

indicated below. **(C)** Volcano plots depicting differentially-expressed genes in scramble versus

shSrsf3 cell lines in the absence (left) or presence (right) of PDGF-AA stimulation. Log_2(fold

change) (FC) values represent log_2(shSrsf3 normalized counts/scramble normalized counts).

Significant changes in gene-level expression are reported for genes with adjusted *P* (padj) <

0.05 and fold change |FC| $\geq$ 2. **(D)** Venn diagram of significant genes from C. **(E)** Volcano plots

depicting differentially-expressed genes in the absence versus presence of PDGF-AA ligand in

scramble (left) or shSrsf3 (right) cell lines. Log_2(FC) values represent log_2(+PDGF-AA

normalized counts/-PDGF-AA normalized counts). **(F)** Venn diagram of significant genes from

E.

Figure 2: PDGFRα signaling for one hour has a more pronounced effect on alternative

RNA splicing. (A) Volcano plots depicting alternatively-spliced transcripts in scramble versus

shSrsf3 cell lines in the absence (left) or presence (right) of PDGF-AA stimulation. Difference in

percent spliced in (ΔPSI) values represent scramble PSI – shSrsf3 PSI. Significant changes in

alternative RNA splicing are reported for events with a false discovery rate (FDR) $\leq$ 0.05 and a

difference in percent spliced in (|ΔPSI|) $\geq$ 0.05. **(B)** Venn diagram of significant transcripts from

A, filtered to include events detected in at least 10 reads in either condition. **(C)** Volcano plots

depicting alternatively-spliced transcripts in the absence versus presence of PDGF-AA ligand in

scramble (left) or shSrsf3 (right) cell lines. Difference in percent spliced in (ΔPSI) values

represent -PDGF-AA PSI – +PDGF-AA PSI. **(D)** Venn diagram of significant transcripts from C,

filtered to include events detected in at least 10 reads in either condition. **(E)** Bar graph

depicting alternative RNA splicing events in scramble versus shSrsf3 cell lines in the absence or

presence of PDGF-AA stimulation (left) or in the absence versus presence of PDGF-AA ligand

in scramble or shSrsf3 cell lines (right).

Figure 3: Srsf3 exhibits differential transcript binding upon PDGFRα signaling. (A)

Schematic of eCLIP experimental design. iMEPM cells were left unstimulated or stimulated with

10 ng/mL PDGF-AA for 1 hour and processed for eCLIP analysis. **(B)** Immunoprecipitation (IP)

of Srsf3 from cells that were UV-crosslinked or not UV-crosslinked with IgG or an anti-Srsf3

antibody followed by western blotting (WB) of input, supernatant (Sup), and IP samples with an

anti-Srsf3 antibody. **(C)** Mapping of eCLIP peaks to various transcript locations in the absence

or presence of PDGF-AA stimulation. 5' UTR, 5' untranslated region; CDS, coding sequence; 3'

998 UTR, 3' untranslated region. **(D,E)** Mean coverage of eCLIP peaks across various transcript

999 locations (D) and surrounding the 5' and 3' splice sites (E) in the absence or presence of PDGF-

1000 AA stimulation. **(F,G)** Top three motifs enriched in eCLIP peaks in the absence (F) or presence

1001 (G) of PDGF-AA stimulation.

1002

1003

Figure 4: Srsf3 and PDGFRα signaling are associated with differential GC content and

length of alternatively-spliced exons. (A) Box and whisker plot depicting the percentage of

exon GC content in exons that are not differentially alternatively spliced, and exons that are

included or skipped when Srsf3 is present from the rMATS analysis. **(B)** Box and whisker plot

depicting the ratio of downstream intron to exon GC content in exons that are not differentially

alternatively spliced, and exons that are included or skipped when Srsf3 is present from the

rMATS analysis. **(C,D)** Box and whisker plots depicting the ratio of upstream intron to exon

length (C) and downstream intron to exon length (D) in exons that are not differentially

alternatively spliced, and exons that are included or skipped when Srsf3 is present from the

rMATS analysis. **(E)** Violin and box and whisker (inset) plots depicting the percentage of exon

GC content in exons that are not bound by Srsf3, and exons that are bound in the absence

and/or presence of PDGF-AA stimulation from the eCLIP analysis. *, $P < 0.05$; **, $P < 0.01$; ***,

$P < 0.001$.

1017

1018

Figure 5: Transcripts bound by Srsf3 that undergo alternative splicing upon PDGFRα

signaling encode regulators of PI3K signaling. (A) Venn diagram of genes with differential

expression (DE) or transcripts subject to alternative RNA splicing (AS) across the four treatment

comparisons that overlap with transcripts with Srsf3 eCLIP peaks in the absence or presence of

PDGF-AA stimulation. **(B,C)** Top ten (B) and PI3K-related (C) biological process gene ontology

(GO) terms for transcripts from the overlapping datasets. p.val, *P*. **(D)** Difference in percent

spliced in (ΔPSI) values for PI3K/endosome-related transcripts of interest. FDR, false detection

rate. **(E)** Peak visualization for input and eCLIP samples in the absence or presence of PDGF-

AA stimulation from Integrative Genomics Viewer (left) with location of motifs from Figure S5

indicated below for PI3K/endosome-related transcripts of interest. Predicted alternative splicing

outcomes for PI3K/endosome-related transcripts of interest (right).

1030

Figure 6: Srsf3 regulates early endosome size and phosphorylation of Akt downstream of PDGFRα signaling. (A,B) Scatter dot plots depicting average size of Rab5 puncta per cell (A) and Pearson's correlation coefficient of signals from anti-Rab5 and anti-PDGFRα antibodies (B) in scramble and shSrsf3 cell lines in the absence or presence (15-60 min) of PDGF-AA stimulation. Data are mean $\pm$ s.e.m. *, $P < 0.05$. Shaded shapes correspond to independent experiments. Summary statistics from biological replicates consisting of independent experiments (large shapes) are superimposed on top of data from all cells; n = 20 technical replicates across each of three biological replicates. **(C-H")** PDGFRα antibody signal (white or magenta) and Rab5 antibody signal (white or green) as assessed by immunofluorescence analysis of scramble and shSrsf3 cells in the absence or presence (15-60 min) of PDGF-AA stimulation. Nuclei were stained with DAPI (blue). White arrows denote colocalization. Scale bars: 20 μm (main images), 3 μm (insets). **(I)** Western blot (WB) analysis of whole-cell lysates (WCL) from scramble (left) and shSrsf3 (right) cell lines following a time course of PDGF-AA stimulation from 15 min to 4 h, with anti-phospho-(p)-Akt and anti-Akt antibodies. Line graphs depicting quantification of band intensities from n = 3 biological replicates as above. Data are mean $\pm$ s.e.m. *, $P < 0.05$; **, $P < 0.01$. **(J)** Model of experimental results in which PI3K/Akt-mediated PDGFRα signaling results in the nuclear translocation of Srsf3 and the subsequent AS of transcripts to decrease levels of proteins that promote PDGFRα trafficking out of early endosomes.