Human V4 size predicts crowding distance

Jan W. Kurzawski*[1,2], Brenda S. Qiu[3], Najib J. Majaj[4], Noah Benson[5], Denis G. Pelli[1,4], Jonathan Winawer[1,4]

[1]Department of Psychology, New York University, New York, NY, USA
[2]Faculty of Psychology and Neuroscience, Maastricht University, Maastricht, Netherlands
[3]Department of Psychology, University of Washington, Seattle, WA, USA
[4]Center for Neural Science, New York University, New York, NY, USA
[5]eScience Institute, University of Washington, Seattle, WA, USA

*Corresponding author

Abstract

Visual recognition is limited by both object size (acuity) and spacing. The spacing limit, called "crowding", is the failure to recognize an object in the presence of other objects. Here, we take advantage of individual differences in crowding to investigate its biological basis. Crowding distance, the minimum object spacing needed for recognition, varies 2-fold among healthy adults. We test the conjecture that this variation in psychophysical crowding distance is due to variation in cortical map size. To test this, we made paired measurements of brain and behavior in 50 observers. We used psychophysics to measure crowding distance and calculate λ, the number of letters that fit into each observer's visual field without crowding. In the same observers, we used fMRI to measure the surface area A (mm²) of retinotopic maps V1, V2, V3, and V4. Across observers, λ is proportional to the surface area of V4 but is uncorrelated with the surface area of V1 to V3. The proportional relationship of λ to area of V4 indicates conservation of *cortical crowding distance* across individuals: letters can be recognized if they are spaced by at least 1.4 mm on the V4 map, irrespective of map size and psychophysical crowding distance. We conclude that the size of V4 predicts the spacing limit of visual perception.

Introduction

Visual recognition is limited by two independent scale factors: the size and spacing of objects[1]. The size limit is understood but the spacing limit is not. The size limit (acuity) is set by properties of the eye, including optical quality and the sampling density of the retinal cell mosaics[2]. However, even when object size is well above the acuity limit, recognition fails if objects are insufficiently spaced (crowding; **Fig. 1**).

Figure 1. Example of crowding. Top: While fixating the cross, the letter V is unrecognizable ("crowded") because the center-to-center letter spacing is less than the crowding distance. Bottom: The letter V is recognizable when the letter spacing is greater than the crowding distance.

Crowding limits performance of important everyday activities, including reading and search[3,4]. The severity of this limitation varies among healthy individuals and between groups: Crowding thresholds are elevated in several disorders, including dyslexia[5], dyscalculia[6], strabismic amblyopia[1] and apperceptive agnosia[7]. Although studied for over a century[8], the biological basis of crowding is largely unknown. Physiological studies have yielded conflicting claims about the anatomical site of crowding (**Table 1**). Perhaps the only certain conclusion is that crowding has a cortical basis, because it persists when target and clutter are presented separately to the two eyes[9,10].

Brain area	Species	Reference
V1	human	Fang and He 2008[11]
V1-V3	human	Kwon et al. 2014[9]
V1-V4	human	Millin et al. 2014[12]
V1-V4	human	Anderson et al. 2012[13]
beyond V1	human	He et al. 1996[14]
V2	human	He, Wang, and Fang 2019[15]
V2	macaque, human	Freeman and Simoncelli 2011; Freeman et al. 2013[16,17]
V2-V3	human	Bi et al. 2009[18]
V2-LOC	human	Jastrzębowska et al. 2021[19]
V3	human	Tyler et al. 2007[20]
beyond V3	human	Liu et al. 2009[21]
V4	macaque	Henry and Kohn 2022[22]
V4	macaque	Motter 2018[23]
V4	macaque	Kim et al. 2023[24]

Table 1. Physiology-based links between crowding and visual brain areas.

Here, we take an anatomical rather than physiological approach to study the basis of crowding. We do so by taking advantage of variation in crowding distance and cortical map size across observers. Individual differences in crowding[25-28] and map size[29] are large even among young healthy observers. We conjecture that crowding results from limited neural resources at some processing stage, resembling the way acuity is limited by retinal cell density. Retinotopic maps in the human cortex are obvious candidates to mediate crowding. We ask whether the surface area of one or more retinotopic

maps is associated with crowding distance variability across observers. Further, we test the more
specific hypothesis that the *cortical crowding distance* (threshold object spacing on the cortical surface)
is a constant for one or more retinotopic maps, despite large individual differences in psychophysical
crowding distance and map size.

Some quantities are known to be conserved in cortical architecture (for review see ref [30]). For
example, in visual cortex, V1 cortical magnification and population receptive field size both vary
substantially across eccentricities and individuals, yet the product of these two measures, "the
population point image", is approximately constant[31]. This indicates that a point in the visual field is
represented by the same amount of V1 cortex, irrespective of the observer and eccentricity. A second
example is found in the mapping between visual areas: Receptive fields in macaque V4 grow with
eccentricity when measured in degrees of visual field; however, within an individual animal, they
maintain a constant size, independent of eccentricity, when projected onto the map of V1, showing
constancy in intracortical sampling[32]. This implies that the way higher level brain areas sample V1 is
relatively constant within an individual. The quantities that are conserved are likely to be important for
explaining the functional organization of the nervous system.

Here we use the concept of cortical conservation to test the possibility that variation in psychophysical
crowding distance is predicted by the surface area of retinotopic maps.

Results

Cortical conservation hypothesis

In each of 50 observers, we psychophysically measure the crowding distance with a letter identification
task, and use fMRI to measure the surface area of retinotopic maps V1 to V4. Inspired by the letter
acuity charts introduced by Anstis in 1974[33], we calculate λ, the maximum number of letters that fit into
an observer's visual field without crowding (**Figure 2a**). The smaller the crowding distance the more
letters fit in their visual field. Conservation of cortical crowding distance implies that variation in λ is
entirely due to variation in surface area A of one or more maps, such that observers with larger maps
can recognize proportionately more letters in their visual field (**Figure 2b**). We express this relationship
as,

$$\lambda \ = \ kA, \tag{1}$$

80 where λ (letters) and A (mm^2) are vectors with one element per observer, and k is a constant scalar
81 (letters per mm^2 of cortex). Writing k as a constant expresses the hypothesis that variation in λ across
82 observers is due only to variation in A.

Figure 2. Conservation of cortical crowding distance. (a) Two hypothetical examples of uncrowded visual fields. In each case, the letters are spaced at the minimum distance necessary for successful recognition. In both examples, the spacing increases with eccentricity. The denser array in Example 1 indicates that this hypothetical observer is more tolerant to clutter (smaller crowding distance), and hence can fit more letters into the uncrowded visual field than observer 2 ($\lambda = 25$ vs $\lambda = 9$). (b) The projection of the letter arrays onto hypothetical retinotopic maps on the cortical surface. The greater λ for the first observer is explained by a larger cortical map, but with equal center-to-center letter spacing on the cortex. Hence the conservation conjecture predicts that the number of letters in the uncrowded field is proportional to surface area. The projections assume cortical conservation of crowding with a constant $k = 1$ letter / mm^2.

Prior work observed that radial crowding distance, in degrees of visual angle, increases linearly with eccentricity (the Bouma law), and that the radial cortical magnification, measured in mm/deg, decreases inversely with eccentricity, such that the product of the two functions is roughly constant[34] (but see [35]). This amounts to conservation across eccentricity of the threshold center-to-center letter spacing on the cortical surface. Critically, this conservation applies equally to any retinotopic map where cortical magnification decreases inversely with eccentricity, without favoring any map for association with psychophysical crowding[34]. In contrast, conservation across individuals might favor a single map, enabling us to link crowding to a brain locus. Indeed, across individuals, the sizes of retinotopic maps vary largely independently, meaning that the size of one map (e.g., V1) only weakly predicts the size of others (e.g., V4), with the exception of adjacent maps such as V1 and V2[29,36]. Here we test each retinotopic map for conservation of crowding distance across individuals, with reason to expect that it cannot be conserved in multiple maps.

Individual differences in crowding and map size

Consider two actual observers, one with a large crowding distance and one with a small crowding distance (**Figure 3a**). As expected, for both observers crowding distance grows with eccentricity. Dividing the crowding distance by eccentricity gives the Bouma factor, which is 0.22±0.03 for observer 1 and 0.34±0.05 for observer 2 [mean±SD$_{\text{test-retest}}$]. Using the Bouma factors, we then compute , the number of letters that can fit into the visual field spaced at their psychophysically measured crowding distances (**Figure 3b**). This value is much larger for Observer 1 than 2 (748 vs. 302). We next estimate map surface area for each observer using retinotopy (**Figure 3c**). Observer 1's V4 surface area is twice Observer 2's, whereas V1 to V3 surface areas are about the same for the two observers (**Table 2**).

Figure 3. Individual differences in crowding and **map size**. (a) Crowding distances for Observers 1 and 2. The crowding distance was measured radially at two eccentricities, 5º and 10º, in the four cardinal directions, twice at each location. The eight target locations are indicated by black dots. Crowding thresholds are displayed as horizontal and vertical lines. The endpoints of the lines indicate the neighboring letter locations at threshold spacing. Observer 2's lines are slightly offset to avoid overlap. Dotted sectors represent the part of the visual field shown in Panel b. (b) For each observer, we show a portion of the visual field with letters spaced according to their Bouma factor. The portion of the visual field shown is 3º to 8.5º eccentricity and ±25º polar angle. The values count letters for the whole visual field up to 10º eccentricity, the extent of our psychophysics and retinotopy. (c) Retinotopic map boundaries (V1 to V4) from both hemispheres of the same two observers. Visual maps were independently drawn by two researchers (dashed and solid lines). Polar angle and eccentricity maps for the two observers are shown in **Supplementary Figure 1**. Maps for all 49 participants are included as Supplementary Data 2.

	A (mm^2)			
	V1	V2	V3	V4
Observer 1	748	3052	2767	2053	1483
Observer 2	302	2580	2815	1907	618
Ratio	2.5	1.2	1.0	1.1	2.4

Table 2. Number of letters in uncrowded visual field and surface area A in two observers. The numbers in the table
indicate each observer's and each observer's surface area A, in mm^2 of the bilateral retinotopic maps and the ratios between
observers. Surface areas are averaged across the two researchers (**Figure 3C**). The two observers are the same as the two in
Figure 3. Data for all observers are provided in Supplementary Data 1.

Test-retest reliability of and A

Testing the conservation conjecture requires reliable measurement. To assess reliability, we estimated
twice per observer and found a high correlation between sessions ($r = 0.96$; **Figure 4A**). The data lie
slightly above the identity line due to a small improvement across sessions (average of 10%). Similarly,
we measured A twice by having two researchers independently draw the retinotopic map boundaries
(from the same fMRI data) and found high agreement in surface area between the two researchers ($r = $
0.94, 0.88, 0.74, 0.73 for V1 to V4, respectively; **Figure 4B**). Although we did not have two separate
scan sessions per observer, retinotopic mapping parameters in the early visual cortex are stable across
scans[37,38], and the uncertainty in A from scan-to-scan variability is negligible compared to differences in
how researchers use heuristics to draw map boundaries[39].

138
139 **Figure 4. Measurement reliability of crowding distance and map size**. (a) Test-retest correlation of . Each dot is one
140 observer, with one value of measured from each of two test sessions on different days. (b) Correlation in map surface area
141 across two researchers. Two researchers independently delineated the boundaries of the four retinotopic maps in each
142 hemisphere of each observer. The surface areas plotted are bilateral (summed across L and R), and are limited to 10 deg of
143 eccentricity.

Cortical crowding distance is conserved in V4 but not V1, V2, or V3

Relating surface area to crowding, we plot each observer's A vs. their ⬚, separately for V1 to V4

(**Figure 5a**). Only V4 shows high correlation between ⬚ and A (r = 0.65; **Figure 5b**). In V1 to V3, there

is no relationship between ⬚ and A, as indicated by the practically zero correlation coefficients and the

circular covariance ellipses in the scatterplots. The correlation of ⬚ with the size of V4 but not V1 to V3

suggests that V4 size is independent of the sizes of the V1 to V3 maps. This is indeed the case in our

dataset, with correlations between V4 size and V1, V2 and V3 of $r \leq 0.11$ (**Supplementary figure 6**).

Previous reports also found no correlation between the size of V1 and V4[39], and little correlation

between the size of V1 and V3[36].

If a map conserves cortical crowding distance across observers, not only will ⬚ and A be correlated, but

the two measures will be related by a scale factor with no offset (**Eq. 1**). The V4 data are close to

scaling, indicated by the orientation of the covariance ellipse being close to the conservation line in

Figure 5a. We quantified how well the conservation hypothesis explains variation in ⬚ by fitting the

scale factor k in equation 1. We find that the conservation conjecture for V4 surface area accounts for

43% of the ⬚ variance, whereas for V1 to V3 it accounts for none (**Figure 5c**).

Figure 5. A test of cortical crowding distance conservation in V1 to V4 in 49 participants. (a) Each panel plots one point per observer as ⬚ vs A. Both ⬚ and A are computed for the bilateral visual field up to 10 deg eccentricity. Black outlines around two dots in each panel show data for the two observers from Figure 2. The colored ellipses are the 1-SD and 2-SD contours from the covariance ellipses. The black dashed lines are the fits by regression with one parameter, slope. Hence these lines are the predictions assuming cortical conservation of crowding. The dotted lines show 68th percentile of the conservation fit. (b) The correlation coefficient r between ⬚ and A for V1 to V4. (c) The coefficient of determination R^2 by assuming conservation (dashed lines in panel a). For panels b and c, we bootstrapped the data 10,000 times across observers; the bars are the medians and the error bars the 68% (thick) and 95% (thin) confidence intervals across bootstraps. See supplementary Figure 2 for results split by which of the two researchers defined the cortical maps, and supplementary Figure 3 for results split by hemisphere.

159 The slope of the conservation line (dashed line in **Figure 5a**) allows us to estimate the required spacing
160 between letters on the V4 map. We do so by re-expressing **Equation 1** in terms of the cortical crowding
161 distance *c*, which measures center-to-center threshold spacing of letters in mm on the cortex:

$$A = \lambda c^2, \tag{2}$$

162 where

$$c = \frac{1}{\sqrt{k}}. \tag{3}$$

For V4, the slope k is 0.54 letters per mm^2 [CI 0.52 - 0.57] which implies that the cortical crowding
distance *c* is 1.36 mm [CI 1.33 - 1.39]. According to the conservation model, this single number, 1.4
mm, is the required letter-to-letter spacing, despite very different map sizes and crowding distances
across observers. This estimate of *c*, 1.4 mm, is robust to variations in the assumed parameter values
used to compute λ (**Supplementary Analyses 1 and 2**). Moreover, the value of *c* obtained here is
close to the value that we obtained in a pilot experiment (1.25 vs. 1.36; **Supplementary Figure 5).**
There are only three common participants between the pilot and the main experiment.

Discussion

Our main finding is that despite large individual differences in psychophysical crowding distance and
map surface area, crowding distance is conserved on the V4 map. We concluded this by comparing the
number of letters that fit into the uncrowded visual field λ and the surface area of retinotopic maps *A*,
finding that λ scales with *A* for V4 but not for V1 to V3. A model assuming scaling explains over 40% of
the variance in λ for V4, and none for V1 to V3. The scaling implies a cortical crowding distance in V4 of
1.4 mm.

Why is cortical crowding distance conserved?

We speculate that this 1.4-mm crowding distance is conserved across observers because there is a
computational unit composed of a fixed number of V4 neurons needed to isolate an object. A fixed
neural count implies a fixed area, which we estimate to be $A/\lambda = 2$ mm^2 of cortex. This area contains
about a quarter million neurons[40]. This proposed V4 computational unit isolating an object is analogous
to a V1 hypercolumn analyzing local scale and orientation[41], about 1.5 mm^2 in human (and 0.6 mm^2 in
macaque)[42].

The V1 hypercolumn is a well-defined concept. Functional domains have been identified in V4
(reviewed by Pasupathy et al[43]), but are not sufficiently well defined to measure their size. Nonetheless,
an indirect measure of cortical organization, the spatial extent of direct lateral connectivity, has been
quantified in macaque: Tracer studies estimate that 95% of intrinsic horizontal connections in V4 are
within 1.9 mm, and 80% within 1.4 mm[44], close to our estimated V4 cortical crowding distance.

A related interpretation is that the spacing limit for recognition relates to receptive field size rather than
cortical distance. In this interpretation, if multiple letters are within the same receptive field, the
representations of the separate objects would be degraded, as suggested in biased competition
theory[45]. The two possibilities are not mutually exclusive. Receptive field size tends to be inversely
correlated with map size, at least in human V1[31] and so spacing and receptive field size are likely to
make similar predictions. Here we focused on area rather than receptive field size because map size is

a robust, reliable measurement, whereas receptive field size estimates have considerable uncertainty,
both in human neuroimaging[46] and single unit electrophysiology[47].

Measuring anatomy vs. function

Physiological methods have linked crowding to a variety of visual maps, with no clear consensus that a
single visual area is closely linked to crowding distance (**Table 1**). Physiological studies of crowding,
and computational models based on physiology, tend to compare neural responses to a target with
different crowding conditions, such as small vs large spacing[12,22,23]. When the stimulus changes, neural
responses also change. As a result, the particular metric calculated from the neural responses by the
experimenter, such as target decodability from a set of sampled neurons[22] or BOLD response
magnitude, is likely to be affected in most visual areas. This creates ambiguity in identifying a single
brain area as the crowding bottleneck. This is unsurprising since object recognition relies on
computations spanning much of the visual system[48].

In contrast, our search for a neural basis of crowding measured anatomy rather than physiology and
exploited individual differences, measuring both crowding and map size in the same 49 individuals. An
advantage of anatomical measures such as map size is that they are robust and do not depend on
task, stimulus, and attentional state (for example, see Extended Figure 9 in ref [[49]]). Unlike the
physiological approach, we find a sharp divide: no relationship between crowding psychophysics and
map size in V1 to V3, and a strong relationship in V4.

V4 as a gateway to the ventral stream

The sharp divide between the results in V4 and those in V1 to V3 surprised us. Why should the neural
resources in a single brain area be linked to crowding thresholds, when image recognition involves
many stages of processing, from optics to retina to the brain? We interpret this sharp divide as
evidence for a spacing bottleneck in recognition. Suppose that each visual area has its own spacing
threshold, such that neural responses to a target object are suppressed whenever there are similar
objects closer to the target than its threshold separation. Our interpretation is that the *psychophysical*
crowding distance is determined by the visual area in which the separation threshold is largest,
meaning the area most sensitive to stimulus spacing. Our study implies that among posterior maps V4
is the most sensitive, i.e., has the largest spacing threshold. This result is consistent with known
properties of visual maps. For example, from V1 to V4, the size of receptive fields increases[50-52] and the
surface area decreases[29,39,53], so that there are decreasing numbers of neurons to analyze a given part
of the visual field in higher visual areas. Hence if objects are sufficiently spaced for V4 (greater than
about 1.4 mm), they will also be sufficiently spaced for earlier areas such as V1 to V3. The opposite,
however, is not true: if stimuli are sufficiently spaced for V1, they can still be insufficiently spaced for
V4. This is why visual areas prior to V4 are not spacing bottlenecks and do not correlate with crowding
measures.

This argument still leaves open the question of why V4 size should be a spacing bottleneck rather than
an even later stage of cortical processing. Indeed, there is spatial sensitivity in ventral temporal brain
areas anterior to V4[39,54-56]. Nonetheless, V4 is likely one of the final processing stages prior to visual
information being analyzed by specialized regions such as those involved in recognizing words[57],
faces[58], and places[59]. Hence if object spacing is sufficient for processing in V4, then recognition by
downstream areas is possible. This conclusion is consistent with connectivity analyses suggesting that
V4 acts as a gateway to the ventral stream recognition areas[43,60]. It is possible that the size of other,
downstream ventral temporal regions would also correlate with crowding behavior, perhaps as strongly

as, or more strongly than, V4. The relationship between crowding and cortical size is more difficult to
study in these areas, however, as retinotopic mapping is less precise in anterior maps, and the criteria
for defining boundaries, and even the number of discrete areas, remains uncertain for category-
selective regions [61]. Thus, we can conclude that V4 is the *first* visual area to show a strong relationship
to crowding behavior, but we cannot draw conclusions about whether it is the only such area.

Our proposal that V4 acts as a spacing bottleneck for recognition is consistent with several properties
of crowding. First, crowding distance is similar for many types of stimuli (letters, faces, other objects)[62],
which seems inconsistent with crowding arising independently in areas that are selective to different
kinds of object. Second, the nonlinearity in the growth of crowding distance with eccentricity[63] matches
the nonlinear decline of cortical magnification with eccentricity in human V4 map[64]. This nonlinearity is
not present in V1, V2 and V3[36].

Homology between human and macaque V4

Our results reveal a tight link between the size of the human V4 map and psychophysical crowding
distance. In arguing that V4 mediates crowding, we referred to literature on both macaque and human
V4. However, the extent of homology between human and macaque V4 remains uncertain[65]. In both
species, V4 has been identified as part of the ventral visual processing stream, with an important role in
recognition[53,66]. On the other hand, human V4 is located on the ventral surface of the occipital
lobe[53,67,68] (but see Refs [69,70]), whereas macaque V4 is split into a ventral and dorsal part[52].
Nonetheless, macaque recognition, like human recognition, is limited by crowding[23,71], and it is
plausible that V4 acts as a bottleneck in both species. Whether or not the surface area of macaque V4
predicts individual differences in macaque crowding thresholds requires independent validation.
259

The signal and the noise

The reader may share our curiosity about the relationship between λ and A at a finer-grained scale than
we reported. Unfortunately, splitting the data into small parcels decreases the signal-to-noise ratio. For
example, the test-retest correlation of λ, pooled across the four meridians, was 0.96, whereas the test-
retest correlations for the left and right meridians alone were 0.84 and 0.71, respectively
(Supplementary Figure 4). The lower reliability of the single meridian behavioral estimates reduces the
power to detect a relationship with cortical measures. Nonetheless, the main pattern of results holds for
various subsets of our data (Supplementary figures 2 and 3), albeit with weaker relationships. For
example, the correlation coefficients between λ and *A* for only one V4 hemisphere and one meridian
are *r*=0.58 and *r*=0.30 (right hemisphere/left meridian, and left hemisphere/right meridian), compared to
r=0.65 bilaterally.
271

What limits the precision our measurements? Retinotopy with fMRI is effective but imperfect[38,72,73].
There are many non-neural factors contributing to the fMRI signal[74]. For example, vascular artifacts
could cause a portion of a map to have poor signal and be excluded from our estimate of *A*, particularly
in the left hemisphere of V4[75,76]. Also, some participants have unusual V2 or V3 maps[77], which might
not be well captured by our manual delineation procedures. Such artifacts can introduce systematic
bias at the level of the individual subject, meaning both researchers underestimate or both overestimate
A. However, as fMRI artifacts are unlikely to be related to behavior, individual biases translate to
random (unbiased) noise at the level of correlation across the group. Random noise puts an upper limit
on goodness of fit but has little effect on estimate of slope.
281

Conclusion

We exploited individual differences to uncover a strong connection between visual crowding and V4 size. Individual differences can help characterize visual mechanisms[78], and they are important in their own right. Extreme individual variation becomes a disorder, as in dyslexia, dyscalculia and apperceptive agnosia, which are associated with unusually large crowding distances[5-7]. Our discovery that the size of the V4 map predicts crowding might help understand these disorders and is a step in settling the long-standing dispute about the cortical locus of crowding.

Methods

Participants

Fifty participants (32 females, 18 males, mean age 27 years) were recruited from New York University. All participants had normal or corrected-to-normal vision. The participants completed one fMRI session to measure retinotopic maps (45 min) and two behavioral sessions to measure crowding distance (30 min each). One participant was dropped from the analysis due to incomplete fMRI data which did not allow for reliable estimation of retinotopic maps. All participants provided written informed consent and approved the public release of anonymized data. The experiment was conducted under the Declaration of Helsinki and was approved by the NYU ethics committee on activities involving human participants. Data from one observer was excluded because the fMRI session was not completed.

Pilot study

We conducted a pilot study with 26 participants and reported the results in a conference abstract[79]. The new experiments reported here were conducted to include test-retest measures, obtain a larger number of participants, and assess replicability. Eight of the pilot subjects were re-tested in the main experiment including both new crowding data and new retinotopic mapping. We performed the same analysis of the pilot data as presented in the main text. The results reported in the pilot study are largely independent and highly consistent with those reported in the main text (**Supplementary Figure 5**).

fMRI stimulus display

Participants viewed a retinotopic mapping stimulus in the MRI scanner using a ProPixx DLP LED projector (VPixx Technologies Inc., Saint-Bruno-de-Montarville, QC, Canada). The stimulus image was projected onto an acrylic back-projection screen (60 cm $\times$ 36.2 cm) in the scanner bore. The projected image had a resolution of 1920 $\times$ 1080 and a refresh rate of 60 Hz. The display luminance was 500 cd/m^2. Participants viewed the screen at 83.5 cm (from eyes to the screen) using an angled mirror that was mounted on the head coil.

fMRI Stimuli

The mapping stimulus was generated in MATLAB 2017a and was presented using the Psychophysics Toolbox v3[80] and custom vistadisp software (https://github.com/WinawerLab/vistadisp) on an iMac computer. For 27 participants, stimulus images were shown within a sweeping bar aperture, identical to the stimulus in Himmelberg et al[37]. For 33 participants, the stimulus image patterns were shown within two types of apertures in alternative scans, a sweeping bar and a wedge+ring. These stimulus apertures were similar to the ones used in the NSD dataset[49]. Unlike NSD, our stimuli changed location in discrete steps and had a larger maximum eccentricity. In short, there were six runs of retinotopic mapping, each lasting 5 minutes (300 TRs of 1 sec). In recent work we found that the inclusion of the wedge+ring aperture was important for getting reliable estimates of pRFs centered near the fovea[37].

Each aperture type was shown three times in an interleaved order. The carrier image pattern was
presented with a 3 Hz temporal frequency and was composed of colorful objects, faces, and scenes at
multiple scales. The faces, objects and scenes were made for a study of object recognition[81], and then
later incorporated into retinotopic mapping, with the addition of an achromatic pink-noise (1/f)
background[82]. The motivation for including faces, objects and pink noise was that a more varied carrier
pattern might effectively drive responses across visual cortex, especially higher-level areas The
stimulus image pattern was windowed within a circular aperture (12.4° of maximum radius). The
apertures were superimposed on a polar fixation grid placed upon a uniform gray background, with a
red or green dot at the center (3 pixels, or 0.07°). Participants completed a fixation task to ensure that
they were maintaining central fixation and remained alert throughout the scan. Participants were
required to respond, via button press, when the central fixation dot changed from green to red, or vice
versa.

Data acquisition and preprocessing

Structural and functional data were acquired on a 3T Siemens MAGNETOM Prisma MRI scanner
(Siemens Medical Solutions, Erlangen, Germany) at the Center for Brain Imaging at NYU. Both the
structural and functional images were acquired using a Siemens 64-channel head coil. MPRAGE
anatomical images were acquired for each participant (TR, 2400 ms; TE, 2.4 ms; voxel size, 0.8mm^3
isotropic; flip angle, 8°) and were auto-aligned to a template to ensure a similar slice prescription for all
participants. Six functional echo-planar images (EPIs) were collected from each participant using a T2*-
weighted multiband EPI sequence. The parameters were as follows: repetition time (TR) of 1000 ms,
echo time (TE) of 37 ms, voxel size of 2 mm³, flip angle of 68°, multiband acceleration factor of 6, and
posterior-to-anterior phase encoding. Additionally, two distortion maps were acquired to correct
susceptibility distortions in the functional images: one spin-echo image with anterior-posterior (AP) and
one with posterior-anterior (PA) phase encoding. Anatomical and functional preprocessing was
performed using fMRIPrep v.23.1.2[83]. For each participant and each run, fMRIprep produced a BOLD
time series on the participant's native FreeSurfer surface.

Population receptive field model

We first averaged the time-series data from the three repetitions of bar aperture and three repetitions of
wedge and ring apertures resulting in two functional time series. For the 27 participants who saw
images only through a bar aperture, we averaged all 6 runs. We used these average data to fit the pRF
model. The analysis was conducted using Vistasoft (https://github.com/WinawerLab/prfVista;). A pRF
is modeled as a circular 2D-Gaussian, as described in Dumoulin and Wandell[84]. The Gaussian is
parameterized by values at each vertex for x, y, and σ. The x and y parameters specify the center
position of the 2D-Gaussian in the visual field. The σ parameter, the standard deviation of the 2D-
Gaussian, specifies the size of the receptive field. The 2D-Gaussian is multiplied pointwise by the
stimulus contrast aperture (apertures were prepared at a resolution of 101 pixels $\times$ 101 pixels), and
then convolved with a hemodynamic response function (HRF) to predict the BOLD percent signal
change. The HRF is parameterized by 5 values, describing a difference of two gamma functions, as
used previously[84]. The HRF was assumed to be the same across vertices within a participant but
differed among participants. We use a two-stage coarse-to-fine approach described in detail by
Dumoulin and Wandell[84], and the addition of the HRF fit is described in detail by Harvey and
Dumoulin[31]

Defining the size of visual areas (V1-V4)

Two authors, JWK and BSQ, independently defined ROIs (V1-V4) by hand using neuropythy (https://github.com/noahbenson/neuropythy)[85]. To define visual maps, researchers followed the heuristics of Wandell[86] for V1-V3, and of Winawer and Witthoft for V4[87]. Note that while we have developed automated tools for defining visual maps, our tools have only been validated for V1 to V3[73]. To match the maximum eccentricity of the psychophysical stimuli we restricted all maps to a maximum eccentricity of 10 deg. The size of each map was calculated as the sum of the surface areas of all vertices included in the map on the mid-gray cortical meshes, defined by FreeSurfer as the surfaces hallway between the meshes at the gray/white interface and the pial/gray interface. Finally, each map surface area was summed across hemispheres.

Measuring crowding distance

Data were acquired with the CriticalSpacing software[88], allowing for reliable and relatively fast measurement of crowding distances. Our crowding database consists of measurements of crowding distance with the Sloan font with radial flankers. We measured crowding at 8 different peripheral locations in the visual field: 2 eccentricities (5, 10 deg) along the four cardinal meridians (upper, lower, left and right). Each observer participated in two crowding sessions (each lasting about 40 min). Sessions were separated at least by a day with a maximum of five days. In session one first block showed stimuli at 5 deg and the second showed stimuli at 10 deg. In the second session the order was reversed.

Crowding stimulus display

Each testing session was completed on an Apple iMac 27" with an external monitor for stimulus presentation. Screen resolution was 5120×2880. Apple iMac has AMD graphics for optimal compatibility with the Psychtoolbox imaging software. The Systems Preference: Displays: Brightness slider was set (by calling MacDisplaySettings.m in the Psychtoolbox) to 0.86 (range 0 to 1) to achieve a white background luminance of about 250 cd/m^2. The observer viewed the screen binocularly at 40 cm distance from the screen. Stimuli were rendered in MATLAB 2021 using the Psychtoolbox[80]

Crowding stimuli

To measure crowding we show a three-letter trigram. For each trial, the three letters are drawn randomly, without replacement, from the 9 possible letters. Letters are rendered as black in the Sloan font, presented on a uniform white background of about 250 cd/m2. For the Sloan font we use the letters "DHKNORSVZ" as possible targets and flankers. For crowding, each trigram is always arranged radially. Each testing session is about 30 minutes, measuring crowding for targets at two eccentricities, 5º and 10º. The eccentricities are measured in separate blocks of about 15 min each. Each block measures crowding thresholds in the four cardinal directions, with four interleaved staircases. Fixation is displayed until the observer presses a key to initiate the trial. Then the letter trigram appears on the screen for 150 ms and the computer waits for the observer to identify the middle letter by choosing the letter using a mouse click. Letter choices appear on the screen after stimulus offset. Observers are instructed to return their eyes to fixation before clicking their response. The computer waits indefinitely for each response. 300 ms after the observer's response, the next trial begins as soon as the observer fixates the center of the crosshair for 250 ms.

Eye tracking

For all thresholds, we controlled for eye movements. To measure fixation, we used an Eyelink 1000 eye tracker (SR Research, Ottawa, Ontario, Canada) with a sampling rate of 1000 Hz. To allow for short

viewing distance (40 cm) we used their Tower mount setup with a 25 mm lens mounted on the Eyelink
camera. We allowed participants to move their eyes within a 1.5-degree radius from fixation. We
checked eye position during the stimulus presentation. If during the stimulus presentation, observers
moved their eyes more than 1.5 deg away from the fixation, the fixation cross turned red, and observers
were asked to repeat the trial. Each threshold was estimated based on 35 trials with correct fixation. On
average, 10% of trials had to be repeated.

Measuring crowding distance

Pelli's[88] procedure estimates the crowding distance. Letter spacing is controlled by QUEST. Spacing
scales with letter size, maintaining a fixed ratio of 1.4:1. We set the guessing rate parameter gamma to
the reciprocal of the characters in the test alphabet for that font, usually 9. We set the "finger error"
probability delta to 1%, so that QUEST can better cope with an occasional reporting mistake. At the end
of each block, QUEST estimates the threshold (crowding distance). We devote 35 trials to measure the
threshold for a condition. In each block we randomly interleaved the four meridian locations, to get a
threshold for each condition, while keeping the observer uncertain as to which condition is present on
each trial.

Calculating the Bouma factor *b*

For each participant we measured 16 crowding distances (2 eccentricities $\times$ 4 cardinal meridians $\times$ 2
sessions). For each session, crowding distances were converted to Bouma factors by dividing them by
the target eccentricity. (This calculation omits   because it's negligible, 0.24 deg << 2.5 deg.) Next, we
estimated one Bouma factor for each session by taking the geometric mean across all 8 visual field
locations. The final estimate for each observer was calculated as the geometric mean of Bouma factors
from sessions 1 and 2.

Calculating the number of uncrowded letters

431 We used the Bouma law to calculate the number   of letters that fit into the visual field without
432 crowding. The Bouma law states that the radial crowding distance s_r, in deg, grows linearly with
433 eccentricity   , in deg,

$$(4)$$

434 where   and   are fitted constants. Tangential crowding distance s_t is generally found to be
435 proportional to radial crowding distance,

$$s_t = \frac{s_r}{\alpha} \tag{5}$$

436 where   (typically 2) is the ratio of radial-to-tangential crowding distances[26].

437 When letters are packed as tightly as possible without crowding, the local density, in letters per deg2, is
438   the reciprocal of the product of the radial and tangential crowding distances. The number   of
439 uncrowded letters in the visual field is the integral of that density over the visual field,

$$\int_{\varphi=0\ \text{deg}}^{\varphi_{max}} \int_{\theta=0}^{2\pi} \frac{\varphi}{s_r s_t}\, d\theta\, d\varphi \tag{6}$$

440 where is the eccentricity (in deg), is the maximum extent of the visual field (in deg), is the
441 polar angle, and and are the radial and tangential crowding distances (in deg). Integrating yields a
442 formula,

$$\lambda = 2\pi\alpha b^{-2} \left(\ln\left(1 + \frac{\varphi_{max}}{\varphi_0}\right) - \frac{1}{1 + \frac{\varphi_0}{\varphi_{max}}} \right)$$

(7)

443 The last term is an approximately logarithmic function of the ratio $_{max}/_0$ (**Supplementary Analysis 2**).
444 We set $_{max} = 10$ deg, corresponding to the maximum eccentricity of both the psychophysical and
445 retinotopic data. We took measured values, $_0 = 0.24$ deg and $= 2$, from our prior study[63]. With these
446 values, the formula reduces to

$$\lambda = \frac{34.9}{b^2}.$$

(8)

447 Our measure of interest, *c*, the cortical letter spacing at threshold separation, depends on . We
448 analyzed the sensitivity of to the values of and (**Supplementary Analysis 1**).

Relationship between *A and*
We quantified the relationship between *A*, cortical map surface area, and , number of letters in the
uncrowded visual field, in two ways. First, we computed the correlation (Pearson's *r*) and covariance
between the two variables. These calculations treat the two variables symmetrically. We visualize the
covariance ellipses at 1-SD and 2-SD, scaling the x and y axes so that one standard deviation of each
variable is the same length (**Figure 3a**). This ensures that the covariance ellipses are circular when the
correlation is 0. Second, to explicitly test for conservation, we fit a linear model to *A* (independent
variable) and (dependent variable), with the intercept set to 0, and asked how much of the variance in
is explained (**Figure 3c**). Note that the variance explained can be negative, because a line through
the origin may be a worse predictor (larger residuals in) than just assuming the mean (a horizontal
line). For both the correlation coefficient and the variance explained, 68% confidence intervals were
calculated by bootstrapping participants with replacement ($n = 10000$).

Bootstrapping and confidence intervals
We quantified variability in the data with confidence intervals derived from bootstrapping over
participants. Bootstrapping has the advantage over parametric statistical measures in that it does not
make assumptions about the shape of the distributions. In a normal distribution, 68% of the data fall
within one standard deviation of the mean, and 95% within two standard deviations. Hence by analogy,
we report 68% and 95% CIs from bootstrapping. While we do not report formal null hypothesis
significance tests (their scientific value is controversial[89,90]), the provided CIs allow for such inferences:
when the 68% CIs of two groups do not overlap, this is similar to rejecting a null hypothesis that the two
groups do not differ with an alpha parameter of 0.05. When the 95% CIs do not include 0, this is similar
to rejecting the null hypothesis that the measure does not differ from 0 with an alpha of 0.05.
471

Code and data availability
473 Code to produce figures and raw crowding data together with MRI surface files are deposited on
474 GitHub. fMRI dataset in BIDS format is available on OpenNeuro
475 (https://openneuro.org/datasets/ds005639/versions/1.0.0).

Acknowledgments

Funding support from NIH NEI Grant R01-EY033628, NIH NEI Grant R01-EY027401, NIH NEI Grant R01-EY027964, NIH NEI Grant R01-EY031446, and NIH NEI core grant for vision science P30EY013079. The authors thank Professor Michael Landy for comments and discussion.

References

Song, S., Levi, D. M. & Pelli, D. G. A double dissociation of the acuity and crowding limits to letter identification, and the promise of improved visual screening. *Journal of Vision* **14**, 3 (2014). https://doi.org/10.1167/14.5.3

Geisler, W. S. Physical limits of acuity and hyperacuity. *J Opt Soc Am A* **1**, 775-782 (1984). https://doi.org/10.1364/josaa.1.000775

Pelli, D. G. *et al.* Crowding and eccentricity determine reading rate. *Journal of Vision* **7**, 20 (2007). https://doi.org/10.1167/7.2.20

Rosenholtz, R., Huang, J., Raj, A., Balas, B. J. & Ilie, L. A summary statistic representation in peripheral vision explains visual search. *Journal of Vision* **12** (2012). https://doi.org/10.1167/12.4.14

Bertoni, S., Franceschini, S., Ronconi, L., Gori, S. & Facoetti, A. Is excessive visual crowding causally linked to developmental dyslexia? *Neuropsychologia* **130**, 107-117 (2019). https://doi.org/10.1016/j.neuropsychologia.2019.04.018

Castaldi, E., Turi, M., Gassama, S., Piazza, M. & Eger, E. Excessive visual crowding effects in developmental dyscalculia. *Journal of Vision* **20**, 7 (2020). https://doi.org/10.1167/jov.20.8.7

Strappini, F., Pelli, D. G., Di Pace, E. & Martelli, M. Agnosic vision is like peripheral vision, which is limited by crowding. *Cortex* **89**, 135-155 (2017). https://doi.org/https://doi.org/10.1016/j.cortex.2017.01.012

Korte, W. Über die Gestaltauffassung im indirekten Sehen. *Zeitschrift für Psychologie* **93**, 17-82 (1923).

Kwon, M., Bao, P., Millin, R. & Tjan, B. S. Radial-tangential anisotropy of crowding in the early visual areas. *Journal of Neurophysiology* **112**, 2413-2422 (2014). https://doi.org/10.1152/jn.00476.2014

Flom, M. C., Heath, G. G. & Takahashi, E. Contour Interaction and Visual Resolution: Contralateral Effects. *Science* **142**, 979-980 (1963). https://doi.org/10.1126/science.142.3594.979

Fang, F. & He, S. Crowding alters the spatial distribution of attention modulation in human primary visual cortex. *Journal of Vision* **8**, 6 (2008). https://doi.org/10.1167/8.9.6

Millin, R., Arman, A. C., Chung, S. T. L. & Tjan, B. S. Visual Crowding in V1. *Cereb Cortex* **24**, 3107-3115 (2014). https://doi.org/10.1093/cercor/bht159

Anderson, E. J., Dakin, S. C., Schwarzkopf, D. S., Rees, G. & Greenwood, J. A. The neural correlates of crowding-induced changes in appearance. *Curr Biol* **22**, 1199-1206 (2012). https://doi.org/10.1016/j.cub.2012.04.063

He, S., Cavanagh, P. & Intriligator, J. Attentional resolution and the locus of visual awareness. *Nature* **383**, 334-337 (1996). https://doi.org/10.1038/383334a0

He, D., Wang, Y. & Fang, F. The Critical Role of V2 Population Receptive Fields in Visual Orientation Crowding. *Curr Biol* **29**, 2229-2236.e2223 (2019). https://doi.org/10.1016/j.cub.2019.05.068

Freeman, J. & Simoncelli, E. P. Metamers of the ventral stream. *Nat. Neurosci.* **14**, 1195-1201 (2011). https://doi.org/10.1038/nn.2889

Freeman, J., Ziemba, C. M., Heeger, D. J., Simoncelli, E. P. & Movshon, J. A. A functional and perceptual signature of the second visual area in primates. *Nat Neurosci* **16**, 974-981 (2013). https://doi.org/10.1038/nn.3402

18   Bi, T., Cai, P., Zhou, T. & Fang, F. The effect of crowding on orientation-selective adaptation in
human early visual cortex. *Journal of Vision* **9**, 13.11-10 (2009). https://doi.org/10.1167/9.11.13
19   Jastrzębowska, M. A., Chicherov, V., Draganski, B. & Herzog, M. H. Unraveling brain
interactions in vision: The example of crowding. *NeuroImage* **240**, 118390 (2021).
https://doi.org/10.1016/j.neuroimage.2021.118390
20   Tyler, C. W. & Likova, L. T. Crowding: a neuroanalytic approach. *J Vis* **7**, 16 11-19 (2007).
https://doi.org/10.1167/7.2.16
21   Liu, T., Jiang, Y., Sun, X. & He, S. Reduction of the crowding effect in spatially adjacent but
cortically remote visual stimuli. *Curr Biol* **19**, 127-132 (2009).
https://doi.org/10.1016/j.cub.2008.11.065
22   Henry, C. A. & Kohn, A. Feature representation under crowding in macaque V1 and V4
neuronal populations. *Curr Biol* **32**, 5126-5137.e5123 (2022).
https://doi.org/10.1016/j.cub.2022.10.049
23   Motter, B. C. Stimulus conflation and tuning selectivity in V4 neurons: a model of visual
crowding. *Journal of Vision* **18**, 15 (2018). https://doi.org/10.1167/18.1.15
24   Kim, T. & Pasupathy, A. Neural Correlates of Crowding in Macaque Area V4. *J Neurosci* **44**
(2024). https://doi.org/10.1523/JNEUROSCI.2260-23.2024
25   Kurzawski, J. W. *et al.* EasyEyes — A new method for accurate fixation in online vision testing.
Front. Hum. Neurosci. **17** (2023). https://doi.org/10.3389/fnhum.2023.1255465
26   Toet, A. & Levi, D. M. The two-dimensional shape of spatial interaction zones in the parafovea.
Vision Res **32**, 1349-1357 (1992). https://doi.org/10.1016/0042-6989(92)90227-a
27   Veríssimo, I. S., Hölsken, S. & Olivers, C. N. L. Individual differences in crowding predict visual
search performance. *Journal of Vision* **21**, 29 (2021). https://doi.org/10.1167/jov.21.5.29
28   Greenwood, J. A., Szinte, M., Sayim, B. & Cavanagh, P. Variations in crowding, saccadic
precision, and spatial localization reveal the shared topology of spatial vision. *Proc Natl Acad*
Sci U S A **114**, E3573-E3582 (2017). https://doi.org/10.1073/pnas.1615504114
29   Benson, N. C. *et al.* Variability of the Surface Area of the V1, V2, and V3 Maps in a Large
Sample of Human Observers. *J. Neurosci.* **42**, 8629-8646 (2022).
https://doi.org/10.1523/JNEUROSCI.0690-21.2022
30   Karbowski, J. Constancy and trade-offs in the neuroanatomical and metabolic design of the
cerebral cortex. *Front Neural Circuits* **8**, 9 (2014). https://doi.org/10.3389/fncir.2014.00009
31   Harvey, B. M. & Dumoulin, S. O. The Relationship between Cortical Magnification Factor and
Population Receptive Field Size in Human Visual Cortex: Constancies in Cortical Architecture.
J. Neurosci. **31**, 13604-13612 (2011). https://doi.org/10.1523/JNEUROSCI.2572-11.2011
32   Motter, B. C. Central V4 Receptive Fields Are Scaled by the V1 Cortical Magnification and
Correspond to a Constant-Sized Sampling of the V1 Surface. *J. Neurosci.* **29**, 5749-5757
(2009). https://doi.org/10.1523/JNEUROSCI.4496-08.2009
33   Anstis, S. M. Letter: A chart demonstrating variations in acuity with retinal position. *Vision Res*
14, 589-592 (1974). https://doi.org/10.1016/0042-6989(74)90049-2
34   Pelli, D. G. Crowding: a cortical constraint on object recognition. *Curr Opin Neurobiol* **18**, 445-
451 (2008). https://doi.org/10.1016/j.conb.2008.09.008
35   Strasburger, H. On the cortical mapping function - Visual space, cortical space, and crowding.
Vision Res **194**, 107972 (2022). https://doi.org/10.1016/j.visres.2021.107972
36   Dougherty, R. F. *et al.* Visual field representations and locations of visual areas V1/2/3 in human
visual cortex. *J. Vis.* **3**, 586-598 (2003). https://doi.org/10:1167/3.10.1
37   Himmelberg, M. M. *et al.* Cross-dataset reproducibility of human retinotopic maps. *NeuroImage*
244, 118609 (2021). https://doi.org/10.1016/j.neuroimage.2021.118609
38   van Dijk, J. A., de Haas, B., Moutsiana, C. & Schwarzkopf, D. S. Intersession reliability of
population receptive field estimates. *Neuroimage* **143**, 293-303 (2016).
https://doi.org/10.1016/j.neuroimage.2016.09.013

39 Brewer, A. A., Liu, J., Wade, A. R. & Wandell, B. A. Visual field maps and stimulus selectivity in
human ventral occipital cortex. *Nat Neurosci* **8**, 1102-1109 (2005).
https://doi.org/10.1038/nn1507
40 Ribeiro, P. F. *et al.* The human cerebral cortex is neither one nor many: neuronal distribution
reveals two quantitatively different zones in the gray matter, three in the white matter, and
explains local variations in cortical folding. *Front Neuroanat* **7**, 28 (2013).
https://doi.org/10.3389/fnana.2013.00028
41 Hubel, D. H. & Wiesel, T. N. Ferrier lecture. Functional architecture of macaque monkey visual
cortex. *Proc R Soc Lond B Biol Sci* **198**, 1-59 (1977). https://doi.org/10.1098/rspb.1977.0085
42 Garcia-Marin, V., Kelly, J. G. & Hawken, M. J. Neuronal composition of processing modules in
human V1: laminar density for neuronal and non-neuronal populations and a comparison with
macaque. *Cereb Cortex* **34**, bhad512 (2024). https://doi.org/10.1093/cercor/bhad512
43 Pasupathy, A., Popovkina, D. V. & Kim, T. Visual Functions of Primate Area V4. *Annu Rev Vis*
Sci **6**, 363-385 (2020). https://doi.org/10.1146/annurev-vision-030320-041306
44 Markov, N. T. *et al.* Weight Consistency Specifies Regularities of Macaque Cortical Networks.
Cerebral Cortex **21**, 1254-1272 (2011). https://doi.org/10.1093/cercor/bhq201
45 Desimone, R. & Duncan, J. Neural mechanisms of selective visual attention. *Annu Rev*
Neurosci **18**, 193-222 (1995). https://doi.org/10.1146/annurev.ne.18.030195.001205
46 Lerma-Usabiaga, G., Benson, N., Winawer, J. & Wandell, B. A. A validation framework for
neuroimaging software: The case of population receptive fields. *PLOS Computational Biology*
16, e1007924 (2020). https://doi.org/10.1371/journal.pcbi.1007924
47 Cavanaugh, J. R., Bair, W. & Movshon, J. A. Nature and interaction of signals from the
receptive field center and surround in macaque V1 neurons. *J Neurophysiol* **88**, 2530-2546
(2002). https://doi.org/10.1152/jn.00692.2001
48 DiCarlo, J. J., Zoccolan, D. & Rust, N. C. How does the brain solve visual object recognition?
Neuron **73**, 415-434 (2012). https://doi.org/10.1016/j.neuron.2012.01.010
49 Allen, E. J. *et al.* A massive 7T fMRI dataset to bridge cognitive neuroscience and artificial
intelligence. *Nat Neurosci* **25**, 116-126 (2022). https://doi.org/10.1038/s41593-021-00962-x
50 Wandell, B. A. & Winawer, J. Computational neuroimaging and population receptive fields.
Trends Cogn Sci **19**, 349-357 (2015). https://doi.org/10.1016/j.tics.2015.03.009
51 Gattass, R., Gross, C. G. & Sandell, J. H. Visual topography of V2 in the macaque. *J Comp*
Neurol **201**, 519-539 (1981). https://doi.org/10.1002/cne.902010405
52 Gattass, R., Sousa, A. P. & Gross, C. G. Visuotopic organization and extent of V3 and V4 of the
macaque. *J Neurosci* **8**, 1831-1845 (1988). https://doi.org/10.1523/jneurosci.08-06-01831.1988
53 Wade, A. R., Brewer, A. A., Rieger, J. W. & Wandell, B. A. Functional measurements of human
ventral occipital cortex: retinotopy and colour. *Philos Trans R Soc Lond B Biol Sci* **357**, 963-973
(2002). https://doi.org/10.1098/rstb.2002.1108
54 Arcaro, M. J., McMains, S. A., Singer, B. D. & Kastner, S. Retinotopic organization of human
ventral visual cortex. *J Neurosci* **29**, 10638-10652 (2009).
https://doi.org/10.1523/jneurosci.2807-09.2009
55 DeWitt, I. & Rauschecker, J. P. Phoneme and word recognition in the auditory ventral stream.
Proc Natl Acad Sci U S A **109**, E505-514 (2012). https://doi.org/10.1073/pnas.1113427109
56 Kay, K. N., Weiner, K. S. & Grill-Spector, K. Attention reduces spatial uncertainty in human
ventral temporal cortex. *Curr Biol* **25**, 595-600 (2015). https://doi.org/10.1016/j.cub.2014.12.050
57 Cohen, L. *et al.* The visual word form area: spatial and temporal characterization of an initial
stage of reading in normal subjects and posterior split-brain patients. *Brain* **123 (Pt 2)**, 291-307
(2000). https://doi.org/10.1093/brain/123.2.291
58 Kanwisher, N., McDermott, J. & Chun, M. M. The fusiform face area: a module in human
extrastriate cortex specialized for face perception. *J Neurosci* **17**, 4302-4311 (1997).
https://doi.org/10.1523/JNEUROSCI.17-11-04302.1997
59 Epstein, R. & Kanwisher, N. A cortical representation of the local visual environment. *Nature*
392, 598-601 (1998). https://doi.org/10.1038/33402

60   Ungerleider, L. G., Galkin, T. W., Desimone, R. & Gattass, R. Cortical connections of area V4 in
the macaque. *Cereb Cortex* **18**, 477-499 (2008). https://doi.org/10.1093/cercor/bhm061

61   Weiner, K. S. & Grill-Spector, K. The improbable simplicity of the fusiform face area. *Trends*
Cogn Sci **16**, 251-254 (2012). https://doi.org/10.1016/j.tics.2012.03.003

62   Pelli, D. G. & Tillman, K. A. The uncrowded window of object recognition. *Nat Neurosci* **11**,
1129-1135 (2008). https://doi.org/10.1038/nn.2187

63   Kurzawski, J. W. *et al.* The Bouma law accounts for crowding in 50 observers. *Journal of Vision*
23, 6 (2023). https://doi.org/10.1167/jov.23.8.6

64   Winawer, J. & Witthoft, N. Human V4 and ventral occipital retinotopic maps. *Vis. Neurosci.* **32**,
E020 (2015). https://doi.org/10.1017/S0952523815000176

65   Vanduffel, W., Zhu, Q. & Orban, G. A. Monkey cortex through fMRI glasses. *Neuron* **83**, 533-
550 (2014). https://doi.org/10.1016/j.neuron.2014.07.015

66   Desimone, R. & Schein, S. J. Visual properties of neurons in area V4 of the macaque: sensitivity
to stimulus form. *J Neurophysiol* **57**, 835-868 (1987). https://doi.org/10.1152/jn.1987.57.3.835

67   Winawer, J. & Witthoft, N. Human V4 and ventral occipital retinotopic maps. *Vis Neurosci* **32**,
E020 (2015). https://doi.org/10.1017/S0952523815000176

68   McKeefry, D. J. & Zeki, S. The position and topography of the human colour centre as revealed
by functional magnetic resonance imaging. *Brain* **120 (Pt 12)**, 2229-2242 (1997).

69   Hansen, K. A., Kay, K. N. & Gallant, J. L. Topographic organization in and near human visual
area V4. *J. Neurosci.* **27**, 11896-11911 (2007).

70   Glasser, M. F. *et al.* A multi-modal parcellation of human cerebral cortex. *Nature* **536**, 171-178
(2016). https://doi.org/10.1038/nature18933

71   Crowder, E. A. & Olson, C. R. Macaque monkeys experience visual crowding. *J Vis* **15**, 14
(2015). https://doi.org/10.1167/15.5.14

72   Bordier, C., Hupe, J. M. & Dojat, M. Quantitative evaluation of fMRI retinotopic maps, from V1 to
V4, for cognitive experiments. *Front Hum Neurosci* **9**, 277 (2015).
https://doi.org/10.3389/fnhum.2015.00277

73   Benson, N. C. & Winawer, J. Bayesian analysis of retinotopic maps. *Elife* **7** (2018).
https://doi.org/10.7554/eLife.40224

74   Kurzawski, J. W., Gulban, O. F., Jamison, K., Winawer, J. & Kay, K. Non-Neural Factors
Influencing BOLD Response Magnitudes within Individual Subjects. *J Neurosci* **42**, 7256-7266
(2022). https://doi.org/10.1523/JNEUROSCI.2532-21.2022

75   Winawer, J., Horiguchi, H., Sayres, R. A., Amano, K. & Wandell, B. A. Mapping hV4 and ventral
occipital cortex: the venous eclipse. *Journal of Vision* **10**, 1 (2010).
https://doi.org/10.1167/10.5.1

76   Boyd Taylor, H. G., Puckett, A. M., Isherwood, Z. J. & Schira, M. M. Vascular effects on the
BOLD response and the retinotopic mapping of hV4. *PLoS One* **14**, e0204388 (2019).
https://doi.org/10.1371/journal.pone.0204388

77   Ribeiro, F. L. *et al.* Variability of visual field maps in human early extrastriate cortex challenges
the canonical model of organization of V2 and V3. *Elife* **12** (2023).
https://doi.org/10.7554/eLife.86439

78   Wilmer, J. B. How to use individual differences to isolate functional organization, biology, and
utility of visual functions; with illustrative proposals for stereopsis. *Spat Vis* **21**, 561-579 (2008).
https://doi.org/10.1163/156856808786451408

79   Kurzawski, J. W., Pelli, D. G. & Winawer, J. A. Conservation across individuals of cortical
crowding distance in human V4. *Journal of Vision* **21**, 2675 (2021).
https://doi.org/10.1167/jov.21.9.2675

80   Kleiner, M. *et al.* What's new in psychtoolbox-3. *Perception* **36**, 1-16 (2007).

81   Kiani, R., Esteky, H., Mirpour, K. & Tanaka, K. Object category structure in response patterns of
neuronal population in monkey inferior temporal cortex. *J Neurophysiol* **97**, 4296-4309 (2007).
https://doi.org/10.1152/jn.00024.2007

82   Benson, N. C. *et al.* The Human Connectome Project 7 Tesla retinotopy dataset: Description
and population receptive field analysis. *Journal of Vision* **18**, 23-23 (2018).
https://doi.org/10.1167/18.13.23
83   Esteban, O. *et al.* fMRIPrep: a robust preprocessing pipeline for functional MRI. *Nat Methods*
16, 111-116 (2019). https://doi.org/10.1038/s41592-018-0235-4
84   Dumoulin, S. O. & Wandell, B. A. Population receptive field estimates in human visual cortex.
NeuroImage **39**, 647-660 (2008). https://doi.org/10.1016/j.neuroimage.2007.09.034
85   Benson, N. C. & Winawer, J. Bayesian analysis of retinotopic maps. *eLife* **7**, e40224 (2018).
https://doi.org/10.7554/eLife.40224
86   Wandell, B. A., Dumoulin, S. O. & Brewer, A. A. Visual field maps in human cortex. *Neuron* **56**,
366-383 (2007). https://doi.org/10.1016/j.neuron.2007.10.012
87   Winawer, J. & Witthoft, N. Identification of the ventral occipital visual field maps in the human
brain. *F1000Res* **6**, 1526 (2017). https://doi.org/10.12688/f1000research.12364.1
88   Pelli, D. G. *et al.* A clinical test for visual crowding. Report No. 5:81, (F1000Research, 2016).
89   Loftus, G. R. *A picture is worth a thousand p values: On the irrelevance of hypothesis testing in*
the microcomputer age 25 thesis, Psychonomic Society, (1993).
90   Vol. 46   806-834 (American Psychological Association, US, 1978).
695

Supplementary materials

Supplementary Figure 1. Retinotopic maps for Observers 1 and 2

Supplementary Figure 1. Retinotopic maps for Observers 1 and 2. For each observer we show flattened representations of their left and right hemispheres. The overlays show eccentricity (top), polar angle (middle) and underlying thresholded cortical curvature (bottom; dark regions are sulci, light regions are gyri). The two sets of outlines (blue and white) are the map boundaries drawn independently by two researchers. In rows 1 and 2, the color overlays are thresholded by pRF variance explained > 10%. Maps for all participants are available as images in Supplementary Data 1 (retinotopicMaps.zip).

Supplementary Figure 2. Model fits of λ vs *A* for V1 to V4, based on map boundaries from single researchers.

Supplementary Figure 2. Model fits of λ vs *A* for V1 to V4, based on map boundaries from single researchers. Panel a shows the correlation coefficient (left) and variance explained from linear regression with 0 intercept (right) when using cortical map definitions from Researcher 1. Panel b is the same but for Researcher 2. The thick and thin error bars are 68% and 95% confidence intervals. The results can be compared to Figure 5b and 5c in the main text, which compute the same statistics for map surface area averaged across the two researchers.

Supplementary Figure 3.

Supplementary Figure 3. Model fits of   vs *A* for V1 to V4, separated by cortical hemisphere. Both panels are as in Supplementary Figure 2, except that: (1) the map surface area is computed only for the right hemisphere (panel a) or left hemisphere (panel b) and   is derived from crowding data only on the left horizontal meridian (Panel a) or right horizontal meridian (panel b). Surface area measures are averaged across the two researchers.

Supplementary Figure 4.

Supplementary Figure 4. Measurement reliability of crowding distance and map size separated by hemisphere and visual field location. Both panels are the same as in Figure 4, except that panel a plots from the left meridian and right hemisphere, while panel b plots the data from right meridian and left hemisphere.

Supplementary Figure 5.

		Pilot	Main
Methods	Number of participants	26	49 (50 planned)
	Nature of research	Exploratory	Confirmatory
	Crowding sessions per subject	1	2
	Number of researchers defining map boundaries	1	2
	pRF stimulus design for better mapping of the fovea[1]	no	yes
	Gaze contingent stimulus display	no	yes
	Removing trials with fixation errors	no	yes
Results	k (letters / mm^2) in V4	0.64 [CI 0.60 - 0.69]	0.54 [CI 0.52 - 0.57]
	c (mm per letter) in V4	1.25 [CI 1.20 - 1.30]	1.36 [CI 1.33 - 1.39]
	r (crowding distance, surface area) V4	0.46	0.66
	R^2 (conservation prediction), V4	0.22 [CI 0.09 - 0.33]	0.43 [CI 0.33 - 0.50]

Supplementary Figure 5. A comparison of pilot and main experiments a, Number of letters λ vs cortical surface area A for the 26 observers in the pilot dataset. Otherwise plotted as in Figure 3a. b, The correlation between λ and A for V1 to V4 in the pilot dataset. Otherwise plotted as in **Figure 3bc**, Variance explained by conservation prediction. Plotted as in **Figure 3c**. d, A comparison of methods and results between the pilot and main experiments.

Supplementary Figure 6.

Supplementary Figure 6. Correlations between surface areas of different maps. The left panel shows correlations for bilateral maps. Note that V4 has little correlation with the surface area of V1 to V3. The panel on the right shows the same but separated by hemisphere. The values in the cells are Pearson correlations (n=49). The asterisks indicate statistical significance of the null hypothesis test that the correlation between surface areas of two maps is exactly 0. One asterisk is p < 0.05, two asterisks is p < 0.01, and three asterisks mean p < 0.001.

Supplementary Analysis 1: Sensitivity analysis of the estimated cortical crowding distance *c*

Effects of α and φ_0 on c.

We concluded that cortical crowding distance is conserved on the V4 map, meaning that there is a single value for *c* (cortical crowding distance) across participants, estimated to be 1.4 mm. We estimated this value by regressing λ, number of uncrowded letters, on *A*, the surface area of the V4 map, assuming an intercept of 0 (eq 1 in main text), and then solving for *c* (equation 3 in main text).

Further, we reported uncertainty in *c* from bootstrapping across participants (**Figure 3**). Here we assess how c is affected by variation of λ across participants. The main-text calculation of λ assumed fixed values from our previous work[2] for two parameters: the tangential-to-radial ratio α of crowding distances and the Bouma law intercept φ_0. Here we calculate how the known parameter variations would affect the *c* estimate. We computed *c* 20,000 times, each time assuming different α values with φ_0 fixed (10,000 calculations), or different φ_0 values with α fixed (10,000 calculations). The values for the two parameters were based on the measured distributions, rather than assuming the mean as in the main text. From our prior study, the mean value of α was 2.10 (sd = 0.38), and the mean value of φ_0 was 0.24 (sd 0.05). Supplementary Table 1 below reports the *c* median and 95% confidence intervals from the calculations.

Median [95% CI] estimate of *c*, assuming: α ($\square$ = 2.10, $\square$ = 0.38), φ_0 = 0.24	Median [95% CI] estimate of c, assuming: α = 2.10, φ_0 ($\square$ = 0.24, $\square$ = 0.05)
1.32 [1.14,1.64]	1.36 [1.25,1.45]

Supplementary Table 1. Sensitivity of cortical crowding distance to variation in α and φ_0.

Supplementary Analysis 2. *Effect of* ϕ_{max} *on* λ.

The formula (Eq. 7) for number λ of uncrowded letters grows approximately logarithmically with ϕ_{max}/ϕ_0 (**Supplementary Figure 7,** below). Our decision to measure (in psychophysics and fMRI) out to 10 deg eccentricity was somewhat arbitrary, but we can use equation 7 to consider how many more uncrowded letters might fit in the far periphery. As expected of logarithmic dependence, large changes in the maximum eccentricity produce only modest changes in the number of letters. For instance, doubling the maximum eccentricity ϕ_{max} from 10 to 20 deg would increase the number λ of uncrowded letters by only 13%, and increasing it tenfold, from 10 to 100 deg, would increase λ by 47%.

Supplementary Figure 7. Sensitivity of number λ of uncrowded letters to maximum eccentricity ϕ_{max}. Plots the formula (Eq. 7) for λ vs. ϕ_{max}. This paper mostly sets ϕ_{max} to 10 deg, shown by the circle. The nearly straight curve shows that the formula's dependence on ϕ_{max} is practically logarithmic.

Supplementary Data 1

Please see attached zip file which includes images of visual maps in all 49 subjects.

Supplementary Data 2

Please see attached csv file which includes a table of visual map sizes and crowding measures for all subjects.

Supplementary References

1   Himmelberg, M. M. *et al.* Cross-dataset reproducibility of human retinotopic maps. *NeuroImage*
         244, 118609 (2021). https://doi.org/10.1016/j.neuroimage.2021.118609
2   Kurzawski, J. W. *et al.* The Bouma law accounts for crowding in 50 observers. *Journal of Vision*
         23, 6 (2023). https://doi.org/10.1167/jov.23.8.6