Masayoshi Teraishi

Title

Identification of novel candidate genes associated symbiotic compatibility of soybean

with rhizobia under natural conditions.

Masayoshi Teraishi, Kosuke Sakaguchi, Takahiro Tsuchimoto, Takanori Yoshikawa

MT,KS,TT

Graduate School of Agriculture, Kyoto University, Kyoto 606-8502, Japan

TY

National Institute of Genetics, Mishima 411-8540, Japan

Keyword: Soybean, root nodule, QTL analysis, RNA-seq

Corresponding author: Masayoshi Teraishi (teraishi.masayoshi.5e@kyoto-u.ac.jp)

Masayoshi Teraishi

Abstruct

A robust symbiotic relationship between soybean and rhizobia can enhance soybean yield

and quality with reduced nitrogen fertilizer inputs, contributing to sustainable agriculture.

The genetic interplay between soybean cultivars and the rhizobial species that colonize

their root under natural conditions remains underexplored. This study builds on the

observation that the prevalence of rhizobial species associated with the soybean cultivars

Peking and Tamahomare varies significantly. We performed a QTL analysis on the

proportion of rhizobial species associated with these cultivars using recombinant inbred

lines derived from a cross between Peking and Tamahomare. A major QTL on

chromosome 18 was identified, accounting for 42% of the phenotypic variation and was

subsequently localized to a 240kb region. RNA-seq analysis indicated that a single gene,

featuring nucleotide-binding and leucine-rich repeat (NBS-LRR) domains, exhibited

markedly different expression levels between the parent cultivars within the QTL region.

As this locus is distinct from the chromosomal regions containing known nodule-related

genes such as *Rj* or *rj*, it likely represents a novel gene involved in the symbiosis between

rhizobia and soybeans.

1 Introduction

Soybean (*Glycine max*) is one of the most economical important crops in the world. It is highly suitable for human and animal diets and is the source of 30% of the world's oil derived from processed crops. Soybean is originated from north China, and now expanding to be worldwide crop, especially north and south America. The use in biofuel production will further increase the economic impact of soybean. Its rising economic importance has led to increased efforts over the past several years to improve soybean productivity.

Legume plants, including soybean, have the important characteristic of symbiosis with rhizobia for nitrogen fixation (Doyle et al., 2003). Nitrogen fixation by legumes is a major source of nitrogen to soil (Schaedel et al., 2021). Legume crops occupy 12-15% of the world's arable land （Graham et al., 2003） and nitrogen fixation by legume crops provides 40million tons of nitrogen annually to agricultural land （Galloway et al., 2008）. The nitrogen-fixing capacity of legumes contributes to the supply of nitrogen to natural and arable ecosystems. Effective utilization of nitrogen-fixing capacity is effective for the development of low-input and sustainable agriculture.

Several genetic factors, called *Rj* or *rj* genes, regulating nodulation traits, were found in soybean. They were identified under inoculated condition of specific rhizobia strains.

Genetic factors under non-inoculated condition or natural condition are poorly

investigated.

Our previous research (Ramongolalaina et al, 2018), performed in natural environment

under non-inoculated condition, showed that the novel genes, located on chromosome 18

in soybean, controls compatibility of rhizobia strains with host plants. No genes

responsible for nodulation have been found in this chromosome region. Here, we report

narrowing down the QTL region, and picking the candidate genes with RNA-seq analysis.

2. Results

2.1 QTL analysis

Ramongolalaina et al. (2018) found a major QTL associated with compatibility of

indigenous rhizobia on Chromosome 18, using 93 recombinant inbred lines (F14) (RILs)

crossed between Peking and Tamahomare.

We redeveloped another set of RILs, which consist of 113 F6 between Peking and

Tamahomare, and applied them for high resolution linkage mapping with GRAS-Di

technology. Linkage map was made of total 860 markers, total length 2846.3cM and

average distance between markers was 3.3cM (Table S1).

Rhizobia strains attached on root were divided into two groups, including *B. japonicum*

and *B. elkanii*, with amplicon sequence analysis of partial ITS region between tRNA-Ala

and end of 23S rRNA. Result of QTL analysis for ratio of rhizobia strains showed that a

single major QTL associated with compatibility of indigenous rhizobia in soil was located

between A08389 (52,593kb) and A35793(57,317kb) on chromosome 18, responsible for

42% of the phenotypic variation (Table 1, Fig. S1). Frequency distribution of the RILs

for ratio of rhizobia strains between two homozygous genotype group on the QTL region

was shown clear bias (Fig.S2). The Peking homozygous RILs had tendency toward to

high frequency of *B.elkanii*, and the Tamahomare homozygous RILs to high frequency of

B. Japonicum. Top LOD value was located on between A01978(56,531kb) and

A35793(57,317kb) (Fig.S3). The previous report (Ramongolalaina et al, 2018) was also

detected QTLs in the same region at the vicinity of the SSR marker Sat_064 (56,334kb)

on chromosome 18.

2.2 Fine mapping of the QTL

To narrow down the QTL, RILs harboring recombination among the QTL region were

selected, and additional PCR markers for detailed genotyping were designed (Table S2).

Finaly, location of the responsible genes for compatibility with rhizobia was successfully

narrowed down from 56,433kbp to 56,675kbp on chromosome 18 (Fig.1). There are 22

predicted gene models, named Glyma.18g280800-Glyma.18g283100, according to

genome assembly sequence data of Williams 82 (Glyma.Wm82.a2).(Table2)

2.3・RNA-seq analysis and candidate genes

To identify potential candidate genes responsible for the QTL, RNA-seq analysis was

performed using RNA derived from roots of Peking and Tamahomare at one-month after

germination. RNA-seq analysis revealed only a single gene, named as Glyma.18g281700

was differentially expressed between the parents within the QTL region with padj<0.05

and fold change >2 (Fig.S4, TableS3). Glyma.18g281700 in Peking was significantly

higher than that in Tamahomare. Glyma.18g281700 encode a NBS-ARC domain and a

leucine-rich repeat (LRR) domain, which of them often found in disease resistance genes.

3 Discussion

In soybean, several genes regulating nodulation were reported. They were called *Rj*(s) or

rj(s) genes. *Rj2*, *Rj3*, *Rj4* and *Rfg1* are known as dominant alleles that restrict nodulation

with specific strains (Hayashi et al., 2012). *Rj2* and *Rfg1* are allelic genes located on

chromosome 16 and encode a member of the TIR-NBS-LRR (Toll-interleukin

receptor/nucleotide-binding site/leucine-rich repeat) resistance protein (Fan et al., 2017).

Rj4 are located on chromosome 1 and encoded a thaumatin-like protein (Hayashi et al.,

2014; Tang et al., 2014; Tang et al., 2016). *Rj3* is restricted nodulation by *B. elkanii*

USDA33 (Vest 1970) but not yet identified. Other recessive genes for regulating

nodulation induce non-nodulation phenotypes (*rj1*, *rj5* and *rj6*) (Pracht et al., 1993; Lee

et al., 2011; Indrasumunar et al., 2011) and hyper-nodulation phenotypes(*rj7*) (Nishimura

et al., 2002a; Nishimura et al., 2002b).

Ramongolalaina et al (2018) demonstrated for the first time to reveal a chromosome

location of genes responsible for proportion of nodulated bacteria strains on root under

natural non-inoculated conditions. In this paper, we have successfully narrow down the

location of the gene underlying the QTL into 240kb interval, in which 22 predicted gene

models are expected. The following RNA-seq revealed that there was single significantly

differential expressed gene between Peking and Tamahomare in the QTL interval.

Three consecutive genes, including Glyma.18g281500, Glyma.18g281600 and

Glyma.18g281700, have ULP1 domain, NBS-ARC domain and LRR domain

(Jebanathirajah et al., 2002; Steele et al., 2019; Wei et al., 2023). The three tandemly

arranged genes showed sequence similarities higher than 90%. Glyma.18g281500 was

just outside of the QTL region and Glyma.18g281600 was within the QTL region but

these two genes were not different at expression level among the parents. Only

Glyma.18g281700 differed substantially among the parents.

Two genes, *Rpp1* and *Rpp1b*, which confers resistance to *Phakopsora pachyrhizi*, the

causal agent of rust, were located at the same region on chromosome 18. (Pedley et al.,

2019; Barros et al., 2023; Wei et al., 2023; Yamanaka et al.,2023). Pedley et al. (2019)

reported that eight genes had homology to the NBS-LRR family of disease R genes on

the four bacterial artificial chromosome (BAC) contigs on chromosome18 including the

Rpp1 locus. The three genes, namely Glyma.18g281500, 281600 and 281700 were

considered as the candidate genes for *Rpp1* and Glyma.18g281600 was the most highly

expressed candidate gene in PI 200492. But Barros et al. (2023) showed that *Rpp1* gene

in PI594756 was located slightly upstream of *Rpp1* in PI200492. Barros et al. (2023)

indicated that genomic variation such as presence/absence and copy number variation

causes the diversification of disease resistance genes.

In conclusion, a single major QTL, located in chromosome 18, was associated with

symbiotic compatibility with nodule. No genes, related with symbiosis between rhizobia

and soybean, have been found in chromosome 18. Single candidate gene was identified

with differential transcription levels between Peking and Tamahomare. This candidate

gene bears NBS-LRR domain, and already reported as one of candidate genes for soybean

rust resistance, *Rpp1*.

4 **Materials and Methods**

4.1 Plant materials

113 F_6-derived recombinant inbred lines (RILs) were provided with their parents, cultivar

Tamahomare and Peking. Major nodulating rhizobia on root of Tamahomare belong to

B.japonicum, while that of Peking to *B. elkanii.*

Soybean seeds of each line, sterilized by being soaked in 70% ethanol for 30 second and

2.5% sodium hypochlorite solution for 3 min, were sowed in a pot (10cm x 20cm x 15cm),

which filled with soil derived from soybean planted field in a Kyoto University

experimental farm and grown in an incubator set at 28°C with 18/6 h photoperiod. Plants

were pulled out from soil at the fourth trifoliolate stage, on one month later after sowing.

Twenty-four nodules per line were harvested randomly from roots.

4.2 DNA extraction

The nodules were washed to remove soil, and surface sterilized with 70% ethanol for

1min and 2.5% sodium hypochlorite solution for 3min and rinsed with sterilized water.

They were put into tubes one by one, added 300ul TPS (100mM Tris-HCl, 1M KCl,

10mM EDTA) buffer and then crashed with multi-beads shocker (Yasui Kikai co, Japan).

After centrifuged at 2500rpm for 1min, the supernatants were mixed with isopropanol

and washed twice with 70% ethanol and dissolved with TE buffer.

4.3 PCR amplification and classification of rhizobia species

DNA sequence analysis of internal transcribed spacer (ITS) region between the 16S and

23S ribosomal RNA of nodule bacteria was used for classification of rhizobia species. In

previous report (Ramongolalaina et al, 2018), classification of rhizobia species was based

on restriction fragment length polymorphism of the PCR products (PCR-RFLP) of the

ITS region among the 16S-23S rRNA gene. In this paper, amplicon sequence of a part of

the ITS region by illumina Miseq was adopted for classification of nodule bacteria

because of high-through put analysis.

Forward primer for first PCR was designed on the tRNA-Ala sequence conserved within

ITS region of the *Bradyrhizobium* spp. Reverse primer for first PCR was used the ITS-R

sequence at the end of 23S rRNA (Akao et al., 2004). Eight-base index, called i5/i7 index,

was added between the first primers and P5/P7 sequence (Fig.S5). Primers used for

amplicon sequence were listed in Table S4. The primers were pre-mixed to be each 10uM

forward and reverse primer, finally to create 576 primer sets mixed from 24 different

forward primers and 24 different reverse primers.

0.1uL DNA templates were mixed with 10uL Emerald Amp Max PCR Master (Takara

Bío. co., Japan) and 0.2uL of 10uM primer set up to 20ul with distilled water for 1st PCR

analysis. PCR cycle consisted of a pre run at 96℃ for 2min, denaturation at 98℃ for

20sec., annealing at 62℃ for 15sec. and extension at 72℃ for 1min. This was repeated

for a total 25 cycles and was followed by a final post-run at 72℃ for 1 min. The PCR

products were purified using the LaboPass Gel Extraction kit (Hokkaido system science

co., Japan) after electrophoresis and mixed with totally 576 PCR products into one tube,

followed by amplicon sequencing using a MiSeq platform (2x300bp) (Illimina, USA)

conducted by Bioengineering Lab.Co., Ltd (Japan).

4.4 Estimation of rhizobial species

Primer and index sequences were trimmed using cutadapt (Martin et al., 2011) from fastq

files and quality filtering was performed using the fastp (Chen et al., 2018), subsequently

fastq files were divided into each RILs. Manifest file was also made for import to QIIME2

(Bolyen et al., 2019). Classification of nodule bacteria was used QIIME2 based on known

bradyrhizobium ITS sequences.

4.5Construction of linkage map

Linkage map of the RILs was constructed with GRAS-Di (genotyping by random

amplicon sequencing, direct) technology, developed by TOYOTA (Enoki et al., 2018).

DNA samples of RILs were sent to Eurofins Genomics, which performed genotyping by

GRAS-Di technology. Codominant DNA markers, accrued by GRAS-Di, were used for

linkage map construction by MapMaker 3.0 (Lander et al., 1987).

QTL analysis was performed using the composite interval mapping (CIM) function of the

R/qtl package with the Haley-Knott regression method. The LOD significance threshold

for detecting QTLs was calculated by performing 1000 iterations using the R/qtl

permutation test.

4.6 RNA-seq analysis

For RNA-seq, total RNA was extracted with RNAiso Plus (TakaraBio, Japan) from roots

of soybean which cultivated on soil for one month, and purified with Qiagen RNeasy

Mini Spin Columns, according to manufacturer protocols. Purified RNAs were sent to the

Bioengineering Lab. Co., Ltd. (Japan) for performing RNA-seq. Analysis of RNA-seq

data was performed by RaNA-Seq program

(https://doi.org/10.1093/bioinformatics/btz854).

REFERENCE

Akao S (2004) Grouping of Bradyrhizobium USDA strains by sequence analysis of 168 rDNA and 168-238 rDNA internal transcribed spacer region. Soil Sci. Plant Nutr.**50:** 517-525

Barros LG, Avelino BB, da Silva DCG, Ferreira EGC, Castanho FM, Ferreira ME, Lopes-Caitar VS, Marin SRR, Arias CAA, de O N Lopes I, Abdelnoor RV, Marcelino-Guimarães FC (2023) Mapping of a soybean rust resistance in PI 594756 at the *Rpp1* locus. Mol. Breed. **43:** 12

Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, Alexander H, Alm EJ, Arumugam M, Asnicar F, Bai Y, Bisanz JE, Bittinger K, Brejnrod A, Brislawn CJ, Brown CT, Callahan BJ, Caraballo-Rodríguez AM, Chase J, Cope EK, Da Silva R, Diener C, Dorrestein PC, Douglas GM, Durall DM, Duvallet C, Edwardson CF, Ernst M, Estaki M, Fouquier J, Gauglitz JM, Gibbons SM, Gibson DL, Gonzalez A, Gorlick K, Guo J, Hillmann B, Holmes S, Holste H, Huttenhower C, Huttley GA, Janssen S, Jarmusch AK, Jiang L, Kaehler BD, Kang KB, Keefe CR, Keim P, Kelley ST, Knights D, Koester I, Kosciolek T, Kreps J, Langille MGI, Lee J, Ley R, Liu Y-X, Loftfield E, Lozupone C, Maher M, Marotz C, Martin BD, McDonald D, McIver LJ, Melnik AV, Metcalf JL, Morgan SC, Morton JT, Naimey AT, Navas-Molina JA, Nothias LF, Orchanian SB, Pearson T, Peoples SL, Petras D, Preuss ML, Pruesse E, Rasmussen LB, Rivers A, Robeson MS, Rosenthal P, Segata N, Shaffer M, Shiffer A, Sinha R, Song SJ, Spear JR, Swafford AD, Thompson LR, Torres PJ, Trinh P, Tripathi A, Turnbaugh PJ, Ul-Hasan S, van der Hooft JJJ, Vargas F, Vázquez-Baeza Y, Vogtmann E, von Hippel M, Walters W, Wan Y, Wang M, Warren J, Weber KC, Williamson CHD, Willis AD, Xu ZZ, Zaneveld JR, Zhang Y, Zhu Q, Knight R, Caporaso JG (2019) Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. Nat. Biotechnol. **37:** 852–857

Chen S, Zhou Y, Chen Y, Gu J (2018) fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics (Oxford, England) **34:** i884–i890

Doyle JJ, Luckow MA (2003) The rest of the iceberg. Legume diversity and evolution in a phylogenetic context. Plant physiol. **131:** 900–910

Enoki H, Takeuchi Y (2018) New genotyping technology, GRAS-Di, using next generation sequencer. Proceedings of the Plant and Animal genome conference

XXVI. San Diego, CA.

Fan Y, Liu J, Lyu S, Wang Q, Yang S, Zhu H (2017) The Soybean *Rfg1* Gene Restricts
Nodulation by *Sinorhizobium fredii* USDA193. Front. Plant Sci. **8:** 1548-1548

Galloway JN, Townsend AR, Erisman JW, Bekunda M, Cai Z, Freney JR,
Martinelli LA, Seitzinger SP, Sutton MA (2008) Transformation of the nitrogen
cycle: recent trends, questions, and potential solutions. Science **320:** 889–892

Graham PH, Vance CP (2003) Legumes: importance and constraints to greater use.
Plant Physiol. **131:** 872–877

Hayashi M, Saeki Y, Haga M, Harada K, Kouchi H, Umehara Y (2012) *Rj* (*rj*) genes
involved in nitrogen-fixing root nodule formation in soybean. Breed. Sci. **61:**
544–553

Hayashi M, Shiro S, Kanamori H, Mori-Hosokawa S, Sasaki-Yamagata H, Sayama
T, Nishioka M, Takahashi M, Ishimoto M, Katayose Y, Kaga A, Harada K,
Kouchi H, Saeki Y, Umehara Y (2014) A thaumatin-like protein, *Rj4*, controls
nodule symbiotic specificity in soybean. Plant Cell Physiol. **55:** 1679–1689

Indrasumunar A, Searle I, Lin M-H, Kereszt A, Men A, Carroll BJ, Gresshoff PM
(2011) Nodulation factor receptor kinase 1α controls nodule organ number in
soybean (*Glycine max* L. Merr). Plant J. **65:** 39–50

Jebanathirajah JA, Peri S, Pandey A (2002) Toll and interleukin-1 receptor (TIR)
domain-containing proteins in plants: a genomic perspective. Trends in Plant Sci.
7: 388–391

Lander ES, Green P, Abrahamson J, Barlow A, Daly MJ, Lincoln SE, Newberg LA
(1987) MAPMAKER: an interactive computer package for constructing primary
genetic linkage maps of experimental and natural populations. Genomics **1:** 174–
181

Lee WK, Jeong N, Indrasumunar A, Gresshoff PM, Jeong SC (2011) *Glycine max*
non-nodulation locus *rj1*: a recombinogenic region encompassing a SNP in a
lysine motif receptor-like kinase (GmNFR1α). Theor. Appl. Genet. **122:** 875–884

Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing
reads. EMBnet J. **17:** 10

Nishimura R, Hayashi M, Wu G-J, Kouchi H, Imaizumi-Anraku H, Murakami Y,
Kawasaki S, Akao S, Ohmori M, Nagasawa M, Harada K, Kawaguchi M
(2002a) *HAR1* mediates systemic regulation of symbiotic organ development.
Nature **420:** 426-429

Nishimura R, Ohmori M, Fujita H, Kawaguchi M (2002b) A Lotus basic leucine
zipper protein with a RING-finger motif negatively regulates the developmental
program of nodulation. Proc. Natl. Acad. Sci. U. S. A. **99:** 15206–15210

Pedley KF, Pandey AK, Ruck A, Lincoln LM, Whitham SA, Graham MA (2019)
Rpp1 Encodes a ULP1-NBS-LRR Protein That Controls Immunity to *Phakopsora*
pachyrhizi in Soybean. Mol Plant Microbe Interact. **32:** 120–133

Pracht JE, Nickell CD, Harper JE (1993) Genes Controlling Nodulation in Soybean:
Rj5 and Rj6. Crop Sci. **33:** 711-713

Ramongolalaina C, Teraishi M, Okumoto Y (2018) QTLs underlying the genetic
interrelationship between efficient compatibility of Bradyrhizobium strains with
soybean and genistein secretion by soybean roots. PloS one **13:** e0194671

Schaedel M, Hidrobo G, Grossman J (2021) From microns to meters: exploring
advances in legume microbiome diversity for agroecosystem benefits. Front.
Sustain. Food Syst. **5:** 668195

Steele JFC, Hughes RK, Banfield MJ (2019) Structural and biochemical studies of an
NB-ARC domain from a plant NLR immune receptor. PloS one **14:** e0221226

Tang F, Yang S, Liu J, Gao M, Zhu H (2014) Fine mapping of the *Rj4* locus, a gene
controlling nodulation specificity in soybean. Mol. Breed. **33:** 691-700

Tang F, Yang S, Liu J, Zhu H (2016) *Rj4*, a Gene controlling nodulation specificity in
soybeans, encodes a Thaumatin-like protein but not the one previously reported.
Plant Physiol. **170:** 26–32

Vest G (1970) *Rj3* A gene conditioning ineffective nodulation in soybean. Crop Sci. **10:**
34-35

Wei W, Wu X, Garcia A, McCoppin N, Viana JPG, Murad PS, Walker DR,
Hartman GL, Domier LL, Hudson ME, Clough SJ (2023) An NBS-LRR
protein in the *Rpp1* locus negates the dominance of *Rpp1*-mediated resistance
against *Phakopsora pachyrhizi* in soybean. Plant J. **113:** 915–933

Yamanaka N, Aoyagi LN, Hossain MM, Aoyagi MBF, Muraki Y (2023) Genetic
mapping of seven kinds of locus for resistance to Asian soybean rust. Plants
(Basel) **12:** 2263

Masayoshi Teraishi

Data availability statement

RNA-seq raw data used in this study was downloaded from the DRA database in DDBJ

with the Bioproject accession numbers of PRJDB17790.

Author contributions

MT conceived and designed the experiments. TY and TT developed the mapping

population and constructed the genetic linkage map and performed fine mapping. TT and

KS collected phenotypic data. MT performed RNA-seq analysis and wrote the manuscript.

All authors contributed to the article and approved the submitted version.

Funding

This work was supported by Grant in Aid for Scientific Research (C) (General) (for MT)

(JSPS KAKENHI Grant numbers 22K05571) and by a The Kyoto University Foundation

(for MT)

Conflict of interest

The authors declare no conflict of interest.

ORCID

Masayoshi Teraishi : 0000-0002-6768-8052

Takanori Yoshikawa : 0000-0002-8631-4216

Kosuke Sakaguchi: 0009-0008-0943-7862

Table 1. QTLs detected for nodule number ratios of *B. japonicum* and *B. elkanii*

QTL name	Trait	Chromosome	Peak (cM)[1]	LOD[2]	Variance (%)
qBJ_18	Ratio of *B. japonicum*	18	77.6	15.13	42.33
qBE_18	Ratio of *B. elkanii*	18	77.6	15.39	42.92

(1) QTL position from the first marker.
(2) Maximum likelihood LOD score for the individual QTL.
(3) Phenotypic variation explained by each QTL.

Table 2　Candidate genes within the narrowed QTL region.

marker	Gene Name (Wm82.a4.v1)	Gene Name (Wm82.a2.v1)	Annotations
Mk1-4			
	Glyma18g51591	Glyma.18G280800	ROTUNDIFOLIA like 21
	Glyma18g51600	Glyma.18G280900	allene-oxide cyclase
	Glyma18g51626	Glyma.18G281000	uncharacterized protein
	Glyma18g51643	Glyma.18G281100	DNA-directed RNA polymerase I and III subunit Rpc19
	Glyma18g51660	Glyma.18G281200	60S ribosomal protein L30-like
	Glyma18g51670	Glyma.18G281300	Potential peptide N-acetyl transferase
	Glyma18g51680	Glyma.18G281400	ethylene-responsive transcription factor
Mk1-5			
		Glyma.18G281451	MEI2-like protein 5
	Glyma18g51715	Glyma.18G281500	Disease resistance protein
	Glyma18g51741	Glyma.18G281600	NB-ARC domain disease resistance protein
A01978			
	Glyma18g51765	Glyma.18G281700	NB-ARC domain disease resistance
	Glyma18g51790	Glyma.18G281800	myb/SANT-like DNA-binding domain protein
	Glyma18g51810	Glyma.18G282000	ADP-ribosylation factor GTPase-activating
	Glyma18g51820	Glyma.18G282100	Protein kinase superfamily protein
	Glyma18g51830	Glyma.18G282200	serine carboxypeptidase-like
	Glyma18g51840	Glyma.18G282300	C-terminal domain phosphatase-like
	Glyma18g51850	Glyma.18G282400	protein bicaudal C homolog 1-B-like
	Glyma18g51871	Glyma.18G282500	E3 ubiquitin protein ligase DRIP2-like
	Glyma18g51880	Glyma.18G282600	Disease resistance-responsive
	Glyma18g51890	Glyma.18G282700	histone-lysine N-methyltransferase SUVR5-like isoform X2
	Glyma18g51900	Glyma.18G282800	Ran binding protein
		Glyma.18G282900	Unknown protein
	Glyma18g51911	Glyma.18G283000	RNA recognition motif
	Glyma18g51920	Glyma.18G283100	Eukaryotic aspartyl protease family
Mk2-1			

Fig.1 Fine mapping of the QTL locus on Chr.18

Marker name	A04404*	Mk1-1	Mk1-2	Mk1-3	Mk1-4	Mk1-5	A01978*	Mk2-1	Mk2-2	Mk2-3	Mk2-4	A35793*	B. japonicum (%)	B.elkanii (%)
Position	56,118kb	56,218kb	56,296kb	56,399kb	56,433kb	56,493kb	56,531kb	56,674kb	56,852kb	56,982kb	57,189kb	57,317kb		
Peking	A	A	A	A	A	A	A	A	A	A	A	A	27.8	72.2
Tamahomare	B	B	B	B	B	B	B	B	B	B	B	B	75.0	25.0
PT109	B	B	B	B	B	B	B	B	B	B	B	A	58.3	41.6
PT72	B	B	B	B	B	B	B	B	B	B	B	A	83.3	8.3
PT84	B	B	B	B	B	B	B	B	B	B	B	A	62.5	33.3
PT166	B	B	B	B	B	B	B	B	B	B	A	A	83.3	16.7
PT32	B	B	B	B	B	B	B	B	A	A	A	A	50.0	45.8
PT90	B	B	B	B	B	B	B	B	A	A	A	A	50.0	50.0
PT94	B	B	B	B	B	B	B	A	A	A	A	A	79.1	20.8
PT77	A	A	A	A	A	A	A	A	A	A	A	B	33.3	66.7
PT158	A	A	A	A	A	A	A	A	B	B	B	B	15.4	84.6
PT36	A	A	A	A	A	B	B	B	B	B	B	B	87.5	12.5
PT10	A	B	B	B	B	B	B	B	B	B	B	B	54.2	37.5
PT37	A	B	B	B	B	B	B	B	B	B	B	B	75.0	25.0
PT197	A	B	B	B	B	B	B	B	B	B	B	B	95.8	4.2

*: GrasDi marker
A and B means Peking and Tamahomare genotypes, respectively.
PT number indicates line number of progenies crossed between Peking and Tamahomare.