Hexokinase 2 expression in apical enterocytes correlates with inflammation severity in

patients with inflammatory bowel disease

Saskia Weber-Stiehl[1*], Jan Taubenheim[2*], Lea Järke[1*], Christoph Röcken[3], Stefan Schreiber[4],

Konrad Aden[1,4], Christoph Kaleta[2], Philip Rosenstiel[1], Felix Sommer[1]

[1] Institute of Clinical Molecular Biology, University of Kiel, Rosalind-Franklin-Straße 12,

24105 Kiel, Germany

[2] Institute of Experimental Medicine, University of Kiel, Michaelisstr. 5, 24105 Kiel, Germany

[3] Department of Pathology, University Hospital Schleswig-Holstein, Campus Kiel, Arnold-

Heller-Straße 3/House U33, 24105 Kiel, Germany

[4] Department of Internal Medicine I, University Hospital Schleswig-Holstein, Campus Kiel,

24105 Kiel, Germany

[*] These authors contributed equally with shared first authorship.

Correspondence: Felix Sommer, Institute of Clinical Molecular Biology (IKMB), Christian-

Albrechts-University (CAU) Kiel, Rosalind-Franklin-Str. 12, Kiel D-24105, Germany; tel: +49(0)

431/500-15146; email: f.sommer@ikmb.uni-kiel.de

Keywords: inflammation / hexokinase / HK2 / human biopsies

Abstract

Background: Inflammation is characterized by a metabolic switch promoting glycolysis and

lactate production. Hexokinases (HK) catalyze the first reaction of glycolysis and inhibition of

epithelial HK2 protected from colitis in mice. HK2 expression has been described as elevated

in patients with intestinal inflammation, however there is conflicting data from few cohorts

especially with severely inflamed individuals, thus systematic studies linking disease activity

with HK2 levels are needed.

Methods: We examined the relationship between HK2 expression and inflammation severity

using bulk transcriptome data derived from the mucosa of thoroughly phenotyped patients

suffering from intestinal inflammation of two independent cohorts. Analyzing publicly

available single cell RNA sequencing data and performing immunofluorescence on colonic

biopsies of unrelated patients with intestinal inflammation confirmed the RNA-based findings

on cellular and protein level.

Results: HK2 expression gradually increased from mild to intermediate inflammation, yet

strongly declined at high inflammation scores. Expression of epithelial marker genes also

declined at high inflammation scores, whereas that of candidate immune marker genes

increased, indicating a cellular remodeling of the mucosa during inflammation with an

infiltration of HK2-negative immune cells and a loss of the apical epithelium – the main site

of HK2 expression. Normalizing for the enterocyte loss clearly identified epithelial HK2

expression as gradually increasing with disease activity and remaining elevated at high

inflammation scores. HK2 protein expression was mostly restricted to brush border

enterocytes and these cells along with HK2 levels vanished with increasing disease severity.

Conclusions: Our findings clearly define dysregulated epithelial HK2 expression as an

indicator of disease activity in intestinal inflammation and suggest targeted HK2-inhibition as

a potential therapeutic avenue.

Background

Prevalence of inflammatory bowel disease (IBD) is rising globally causing severe health issues and drastically reducing quality of life. These diseases are multifactorial with complex interactions of multiple genetic and environmental factors[1]. Despite decades of intense research, the exact etiology of IBD remains mostly unknown limiting treatment options[2]. In recent years it became evident that ongoing inflammation is characterized by changes in metabolic activity, in particular glucose metabolism[3–5]. Hexokinases (HK) catalyze the first and irreversible step of glycolysis – the phosphorylation of glucose to glucose-6-phosphate – and are therefore crucial for glucose metabolism and maintenance of homeostatic body functions. In mammals, five HK isoenzymes have been identified: HK1, HK2, HK3, GCK (Glucokinase) and HKDC1 (HK domain containing 1). Each of these HK isoenzymes display specific tissue expression patterns and glucose affinities. HK2 is of particular interest as it is the most abundant HK member within the intestine, exerts high HK activity, responds to various internal as well as external stimuli and shows elevated levels during inflammation[6]. Recently, we further revealed an upregulated expression of HK2 specifically in inflamed compared to non-inflamed tissue of the same patient, irrespective of the type of intestinal inflammation and demonstrated that ablation of HK2 in intestinal epithelial cells (IEC) protects from acute intestinal inflammation, suppresses cell death and alters mitochondrial function[6]. These findings placed HK2 as a molecular target to treat intestinal inflammation. However, when expanding our research about the role of HK2 in intestinal inflammation, we encountered transcriptome datasets with contradictory findings of either an unaltered HK2 expression or even downregulation, especially in severely inflamed patients undergoing bowel resection[7]. We therefore aimed to investigate the cause of this apparent contradiction with the goal to clarify the relation between HK2 expression and intestinal inflammation. Here, we show that with increasing inflammation severity the intestinal mucosa is gradually remodeled, which comprises a partial loss of the apical epithelium – the primary source of HK2 expression – and a simultaneous infiltration of immune cells. Normalizing for this cellular remodeling clearly demonstrates a gradual upregulation of HK2 expression with severity of intestinal inflammation.

Methods

Human studies

We used mucosal transcriptomic data and fixed intestinal specimen of two independent large longitudinal clinical studies, namely the EMED[9] (191 samples) and FUTURE[8] (87 samples) cohorts (trial IDs: EudraCT number 2016-000205-36 and ClinicalTrials.gov NCT02694588), for which accompanying disease activity scores (Harvey-Bradshaw Index (HBI)/Mayo Score) were available (see Table S1 for cohort clinical characteristics). Note that multiple samples were collected from each patient. Sample origin was incorporated in all following analyses. The HBI[26] is used to quantify disease activity in patients suffering from Crohn's disease (CD), one of the two IBD subtypes. Here, variables such as general well-being, abdominal pain and abdominal mass with each having scores of 0-3, as well as the number of liquid stools per day (1 point per stool) and other complications (1 point per complication) yield the "open-ended" total score (Table S3). The Mayo score is used for ulcerative colitis (UC), the other clinical IBD subtype. This scoring system accounts for general well-being, rectal bleeding, endoscopic results and stool frequency with scores of 0-3 per category for an overall score ranging from 0 to 12 (Table S3). To facilitate a direct comparison of both scores despite them being "open-ended" and "discrete", we calculated a general inflammation score by setting the highest score in the dataset for each HBI and Mayo score to 1 and scaled the score of each patient accordingly (see Table S3).

Transcriptome data analysis

RNA sequencing data derived from mucosal biopsies of patients with various grades of intestinal inflammation were retrieved from two previously published IBD cohorts[8,9]. Read counts were transformed to Transcripts Per Million (TPM) values to normalize for differential sequencing depths among samples. TPM data were then plotted per sample against the inflammation score using R (version 4.2.2) and ggplot2 (3.4.3). Trendlines were calculated including all data points and using the distance to the actual location as a weight to enable a robust calculation and avoiding overfitting especially in areas of scarce sampling. To test for an association of gene expression to changes in the inflammation score we used variance-stabilization transformation of raw read counts and fitted linear mixed models with following form: vst(geneExpression) ~ Age + Sex + Diagnosis + InflammationScore. Statistical analyses

106   were performed using R (version 4.2.2[27] with the following packages: lm4 (version 1.1-29)[28],

107   lmerTest (version 3.1-3)[29], car (version 3.0-13)[30], ggplot2 (version 3.3.6)[31] and DESeq2

108   (version 1.38.3)[32].

Cellular deconvolution

We deconvoluted the bulk RNA data using the MuSiC package (version 1.0.0)[10] with default

values. As reference, we used a single cell data set from UC patients described in a study by

Smillie et. al.[11]. The original single cell data set was split in "epithelial", "immunogenic" and

"stromal" subsets, which we rejoined in our analysis. Due to limited resolution in

deconvolution approaches and to reduce cell diversity, we pooled ontogenetically closely

related cell types (Table S4). We restricted the analysis to the eight most abundant cell types.

Single cell RNA sequencing

Single cell data was obtained from the Single Cell Portal (accession SCP259) initially described

by Smillie and colleagues[11]. The different cell populations were rejoined and general quality

controls were performed. In short, we filtered cells with read counts below 1000 (low quality).

Further, we removed cells with less than 200 or with more than 2720 (two-fold of the

standard deviation of expressed genes across all cells) differently expressed genes to account

for spurious sequencing depth and removal of duplicates. All cells with a mitochondrial RNA

content above 5% were also removed. For plotting, gene expression values where log-

transformed. Plotting and quality filtering was performed in R (version 4.2.2) using ggplot2

(version 3.4.3) and Seurat (version 4.4.0)[33].

Immunofluorescence

5 μm sections of paraffin-embedded colonic biopsies were deparaffinized with Xylol

substitute (Roth), incubated in citrate buffer for 3 minutes and subsequently blocked in 5 %

BSA-PBS and 0,2 % TritonX for 30 minutes. Primary anti-E-cadherin (1:400 in 1 % BSA, #3195,

Cell Signaling Technology) and anti-Hexokinase 2 (1:500 in 1% BSA, Cat# NBP2-02272, Novus

Biologicals, Colorado, US) antibodies were incubated over-night. Sections were washed,

incubated with secondary antibodies (Alexa Fluor 488 goat anti mouse, Invitrogen, A32731

and Alexa Fluor 555 goat anti rabbit, Invitrogen, A21430) for 45 minutes at room

temperature. DAPI (1:40000 in PBS, D9542, Sigma Aldrich, St. Louis, US) was used for DNA

counterstaining. Slides were mounted using antifade mounting media (DAKO, Hovedstaden,

Denmark). The quantitative analysis was performed using fluorescence microscopy for

stained samples using the imager Z1 microscope (ZEISS, Jena, Germany) and ZEN software

(version 3.0). Images were taken by a digital camera system (AxioCam HrC/HrM, Zeiss, Jena,

Germany) and ApoTome (ZEISS, Jena, Germany). Fluorescence signal intensity was measured

using Fiji/ImageJ software.

Statistics

General statistical analyses were performed using the GraphPad Prism 9 (GraphPad Software

Inc., La Jolla, USA). For pairwise comparisons the Mann-Whitney-U test was used whereas for

multiple comparisons one-way ANOVA with false discovery rate (FDR) correction were

performed. Data are shown as mean ± standard error of the mean (SEM). A p-value of ≤ 0.05

was considered as significant (*). A p-value of ≤ 0.01 was considered as strongly significant

(**) and p-value of ≤ 0.001 as highly significant (***). For longitudinal data, linear mixed effect

models were fitted where gene expression was used as dependent variable, while

inflammation score and patient ID were fixed and random factors. The models were fitted

using lme4 (version 1.1-31) and statistical testing for coefficients was performed using

lmerTest (version 3.1-3). Model validity was tested against a null-model using log-likelihood

ratio tests and model assumption were evaluated by diagnostic plotting of model residuals,

data point influence strength and random factor fitting.

Results

Epithelial HK2 expression increases with high inflammation scores

To determine a possible correlation of HK2 expression and disease activity, we analyzed RNA sequencing (RNA-seq) data derived from intestinal biopsies of patients suffering from IBD of two independent clinical studies[8,9], for which accompanying disease activity scores for each sample were available (Table S1). This analysis revealed that HK2 expression initially increased gradually reaching peak expression at mid disease activity (inflammation score ~ 0.4) and then declined with high inflammation scores (Figure 1A). As within the intestine HK2 is mainly expressed by epithelial cells of the apical mucosa (Figure S1), we hypothesized that with very high levels of inflammation these apical epithelial cells and therefore also the main site of HK2 expression might be lost due to shedding and apoptosis or transdifferentiation resulting in decreased HK2 expression levels in bulk RNA-seq data. To investigate this hypothesis, we assessed the expression levels of three different epithelial marker genes, namely E-cadherin (*ECAD*), epithelial cell adhesion molecule (*EPCAM*) and Villin1 (*VIL1*). Supporting our hypothesis, the expression levels of these epithelial markers were all declining with increasing inflammation severity (Figure 1B-D), which indicates progressive epithelial damage caused by the inflammation. To account for this epithelial loss, we then normalized the expression of HK2 to that of these epithelial markers, thereby yielding an epithelial HK2 expression. Importantly, epithelial HK2 expression gradually increased in correlation to disease activity and then remained elevated at high inflammation scores (Figure 1E). To test whether the observed expression changes were unique to the combined clinical cohorts, we also performed these analyses separately for each clinical cohort (Figure S2). Importantly, we observed for both clinical cohorts the same changes in *HK2*, *ECAD*, *EPCAM*, *ECAD* and epithelial HK2 expression. We also tested for potential differences between the two main types of IBD, Crohn´s disease (CD) and ulcerative colitis (UC) and found similar responses both for CD and UC (Figure S3). Finally, correlation of HK2 expression with disease severity was confirmed using linear regression models (Figure S4 and Table S2).

Immune cell infiltration during intestinal inflammation

Another source for cellular remodeling during inflammation is tissue infiltration by immune

cells. Using the bulk RNA-seq data we also investigated the proportions of various immune

cell types in relation to disease severity. First, we performed a candidate gene-based analysis

and chose several immune marker genes to assess infiltration of T cells, T helper 1 (Th1) cells,

T helper 2 (Th2) cells, B cells, basophils, neutrophils, eosinophils, and macrophages (Figure

2A). In contrast to *HK2* and the epithelial marker genes, the expression of these individual

immune cell genes increased with the inflammation score. However, slight alterations in the

expression patterns could be observed, mostly depending on the immune cell type.

Expression of the T cell marker genes *TRAC*, *CD3D* and *CD3E,* the B cell marker gene *CD19,* the

eosinophil marker gene *CCR2*, the neutrophil marker genes *CD14* and *CXCR4* as well as the

basophil marker genes *ITGA2B* and *ITGA4* all gradually increased in a linear fashion with the

inflammation score (Figure 2). In contrast, the marker genes for Th2 (*IL4*, *IL5*, *IL13*), Th1 (*IFNG*,

IL12) cells and macrophages (*ITGAM*, *IL1B*) all only displayed biphasic expression patterns

with a first phase characterized by slight increases in expression until an inflammation score

of approximately 0.5 and a second phase at higher disease severity characterized by larger

changes in their expression. The greatest expression changes were detected for the

macrophage marker genes *ITGAM* and *IL1B* suggesting the greatest relative increase of these

cells with increasing disease activity. Normalizing *HK2* expression to the candidate immune

cell genes revealed gradual declines with increasing inflammation scores for all tested

immune cell types indicating that the immune cell infiltration leads to more cells present in

the mucosa, which do not or only express very little *HK2* (Figure 2B).

Next, we moved from the candidate gene-based to a systematic cellular deconvolution of the

bulk RNA-seq data using MuSiC[10]. This program uses a reference single cell data set and cell

type-specific expression profiles to derive the abundance of the individual cell types from bulk

RNA-seq data. We restricted this analysis to the eight most abundant cell types. The

proportion of epithelial cells gradually declined with increasing inflammation scores whereas

the proportion of macrophages, B cells and regulatory T cells increased (Figure 3A). The

proportion of mast, cytotoxic T cells and dendritic cells remained mostly unchanged.

Therefore, this data indicates a massive gradual cellular remodeling of the intestinal mucosa

during inflammation and the findings of the systematic cellular deconvolution supported

those of the candidate gene-based analyses. Furthermore, the drastic decrease in the

proportion of epithelial cells with increasing inflammation scores demonstrates again that the
main source of *HK2* expression is lost during the disease course. This is therefore also
reflected by the outcome that after accounting for the abundance of epithelial cells the
normalized epithelial *HK2* expression gradually increased with disease severity (Figure 3B).

Apical HK2 expression in mature enterocytes correlates with inflammation severity

To validate the transcriptome data and to elucidate the biogeography of the changes in tissue
architecture in relation to HK2 expression, we performed immunostainings on colonic
biopsies of an independent set of UC patients that had been thoroughly scored for histological
disease severity (Nancy score). HK2 protein levels increased non-significantly in mildly
inflamed (score 1) compared to non-inflamed specimen (score 0) and remained unaltered at
medium or strong inflammation (scores 2 and 4) (Figure 4A,C). In contrast, protein levels of
the epithelial marker E-cadherin (ECAD) were unchanged in mild (score 1) but non-
significantly reduced at medium or strong inflammation (scores 2 and 4) compared to non-
inflamed control samples (Figure 4A,C). Notably, structural epithelial damage of inflamed
biopsies was detectable with increasing inflammatory scores (Figure 4C) with loss of epithelial
cells and an infiltration of other cells, probably immune cells, into the inflamed mucosa. In
addition, many epithelial cells that were not shed but still present at the inflamed site had
reduced ECAD expression, which could indicate transdifferention of these inflamed apical
enterocytes. Normalizing the HK2 protein levels to those of ECAD revealed that epithelial HK2
protein levels (Figure 4B) significantly increased with inflammation regardless of disease
severity (scores 1-4). This epithelial HK2 protein expression in inflamed intestinal mucosal
biopsies mirrored and confirmed the transcript patterns of the RNA sequencing analysis.

Furthermore, we used recently-published data from single cell RNA sequencing of mucosal
biopsies from IBD patients[11] to investigate the *HK2* expression on a cellular level in relation to
intestinal inflammation. This analysis of single cell RNA sequencing data confirmed our
findings from the bulk RNA sequencing and immunofluorescence, as *HK2* was mainly
expressed by mature enterocytes and then followed by goblet cells and immature
enterocytes (Figure 5A,B). Next, we checked for *HK2* expression pattern in enterocytes at
different inflammatory stages (healthy vs. inflamed vs. non-inflamed) and found that the
lowest *HK2* expression can be observed in healthy controls, while cells from inflamed samples

showed an increased HK2 expression (Figure 5C). Non-inflamed samples had intermediate

HK2 expression levels (Figure 5C), which indicates that HK2 expression is dysregulated in

enterocytes of IBD patients also in the absence of an overt inflammation. This pattern of *HK2*

expression was also present although less predominant in immature enterocytes and goblet

cells. Finally, we looked into the number of these three cell types that are detected in the

three health states and found that mature and immature enterocytes are lost during

inflammation, while incomplete recovery can be observed in non-inflamed tissue of IBD

patients (Figure 5D). Altogether, the single cell data therefore supports a cellular

reprogramming during inflammation with loss of mature enterocytes (the main cell type of

the apical epithelium) and a dysregulated *HK2* expression in inflamed apical epithelial cells.

In summary, both the RNA and protein data clearly demonstrate that epithelial HK2 was

overexpressed during intestinal inflammation, which highlights HK2 as an indicator of active

disease. Building upon previous findings from murine models that inhibition of HK2

expression either genetically or via the microbial metabolite butyrate protect from

experimental colitis[6], these new data now show that indeed HK2 expression is dysregulated

in the mucosa of patients with active inflammation and therefore suggest that targeting HK2

may represent a promising approach to suppress intestinal inflammation in humans.

Discussion

There is a growing need for valid disease biomarkers and molecular targets for intestinal inflammation as prevalence is increasing and despite significant efforts the cause of these diseases remains unknown. In a recent study, HK2 was identified as upregulated under inflammatory conditions in mice and humans irrespective of disease subtype (CD, UC or non-IBD colitis) and suppressing HK2 expression even protected from experimental colitis[6] placing HK2 as a potential disease marker. However, we found other transcriptome datasets generated from severely inflamed patients in which HK2 expression was unaltered or even reduced[7]. Here, we therefore thoroughly investigated the relationship between HK2 expression and disease severity by analyzing transcriptome data derived from intestinal biopsies of patients with intestinal inflammation for which corresponding data on disease severity (HBI, Mayo score, Nancy score) were available. In addition, we performed cellular deconvolution analyses of the RNA sequencing data to infer the changes in cell proportions during inflammation. Finally, we used immunofluorescence staining to clarify the changes in HK2 protein biogeography in the mucosa during inflammation. Using these approaches, we demonstrated that raw HK2 RNA and protein expression at first gradually increased with the inflammation scores, yet after reaching a critical inflammation score HK2 expression declined again at very severe inflammation. These findings integrate the seemingly conflicting data from previous studies[7,12–14] by linking HK2 expression to disease severity. We further were able to demonstrate that loss of HK2 expression at high inflammation levels was dependent on the disruption of brush border enterocytes as both the expression of epithelial marker genes were decreasing with the inflammation score and immunofluorescence analyses clearly pointed to a destruction of the apical epithelium – the main site of HK2 expression[15,16]. These findings are in line with clinical practice, in which epithelial damage is a key criterion to classify an increasing inflammatory state during IBD[17], for example with epithelial erosion being a feature of histological inflammation in the Nancy score. By analyzing single cell RNA sequencing data derived from an independent set of UC patients, we confirmed that in the gut epithelial cells are the main source of *HK2* expression, both in terms of per cell expression as well as the number of cells contributing to overall expression (Figure 5). These cells are consequently lost due to increasing inflammation in the single cell data set, which would explain the decrease of *HK2* expression at high levels of inflammation[11]. Finally, we were able

to deconstruct the simultaneous epithelial erosion and immune cell infiltration into the

submucosa during intestinal inflammation from the RNA sequencing data. Based on the

overall abundance and their induction (fold change) during inflammation, infiltrating

macrophages seemed most relevant, but also other cell types such as B cells increased in

proportion. Especially the expansion of macrophages could be important and contributing to

the loss of epithelial cells and therefore HK2 expression during inflammation as IL-1β

interferes with the tight junction complexes between IECs[11–13] and thereby increases

intestinal permeability[21]. Similar to epithelial erosion, immune cell infiltration is a classical

feature of inflammation and used in clinical practice to histologically evaluate disease

severity, in particular as a feature of the Nancy score[22]. In addition, transdifferentiation of

HK2-positive mature enterocytes into other cell types such as HK2-negative immature

enterocytes and stem cells[23,24] or maybe even into mesenchymal cells as during epithelial–

mesenchymal transition [25] could also contribute to a reduction of HK2 levels in the inflamed

mucosa. In summary, our findings imply that during the course of inflammation, changes in

the cellular composition of the mucosa, i.e. loss of brush border epithelial cells expressing

high levels of HK2 and a corresponding massive recruitment of immune cells expressing little

to no HK2 to the site of inflammation, affect the overall bulk HK2 expression leading to an

overall reduction despite higher expression levels in fewer and fewer HK2-positive cells

(Figure 6). Overall, we want to highlight the importance of taking into consideration the

biogeography and changes in cell type expression ratios, when trying to identify disease

biomarkers, especially regarding complex diseases such as intestinal inflammation involving

various internal and external factors. Most previous studies used bulk expression data in the

search for disease biomarkers or therapeutic targets and therefore may have missed other

locally restricted but disease relevant genes like *HK2*. In our analyses HK2 expression was only

significantly associated with disease severity after considering epithelial cell abundance

(Figure 1E and Figure 3B) and we speculate that this pattern will also hold true for other genes

with an apical or otherwise restricted expression. For example, other hexokinases (HK1,

HKDC1), proteins responsible for the uptake of dietary nutrients (e.g. SGLT1, SLC15A1) or

those contributing to the intestinal epithelial barrier (e.g. CLDN15, ZO-1) also show a

predominant or even restricted apical expression and therefore possibly have not yet been

identified as disease biomarker during severe intestinal inflammation due to the loss of their

primary expression site. Notably, epithelial remodeling not only includes the partial loss of

apical enterocytes and infiltration of lymphocytes, but also includes hyperregeneration,

metaplasia and a loss of goblet cells. The advent of single cell technologies such as single cell

RNA sequencing promise to pave the way for more refined analyses that will enable the

discovery of more suitable disease markers and to enhance our understanding of the

molecular and cellular mechanisms during disease progression.

Conclusions

HK2 has not yet been described as IBD risk gene, most probably due to its site-restricted expression, but here we were able to show an association of HK2 and epithelial status during intestinal inflammation. Therefore, epithelial HK2 expression correlates with disease severity making it a useful indicator of intestinal inflammation and highlighting the therapeutic potential of targeting HK2. However ultimately, clinical studies are required to indeed demonstrate the feasibility and efficacy of a HK2-targeted intervention in IBD patients.

References

1. Kaplan, G. G. & Windsor, J. W. The four epidemiological stages in the global evolution of

inflammatory bowel disease. *Nat Rev Gastroenterol Hepatol* **18**, 56–66 (2021).

2. Kaplan, G. G. The global burden of IBD: from 2015 to 2025. *Nat Rev Gastroenterol*

Hepatol **12**, 720–727 (2015).

3. Rath, E. & Haller, D. Intestinal epithelial cell metabolism at the interface of microbial

dysbiosis and tissue injury. *Mucosal Immunology* **15**, 595–604 (2022).

4. Gaber, T., Strehl, C. & Buttgereit, F. Metabolic regulation of inflammation. *Nature*

Reviews Rheumatology **13**, 267–279 (2017).

5. Soto-Heredero, G., Gómez de las Heras, M. M., Gabandé-Rodríguez, E., Oller, J. &

Mittelbrunn, M. Glycolysis – a key player in the inflammatory response. *FEBS J* **287**,

3350–3369 (2020).

6. Hinrichsen, F. *et al.* Microbial regulation of hexokinase 2 links mitochondrial metabolism

and cell death in colitis. *Cell Metab* **33**, 2355-2366.e8 (2021).

7. Peters, L. A. *et al.* A functional genomics predictive network model identifies regulators of

inflammatory bowel disease. *Nat Genet* **49**, 1437–1449 (2017).

8. Schreiber, S. *et al.* Therapeutic Interleukin-6 Trans-signaling Inhibition by Olamkicept

(sgp130Fc) in Patients With Active Inflammatory Bowel Disease. *Gastroenterology* **160**,

2354-2366.e11 (2021).

9. Zeissig, S. *et al.* Vedolizumab is associated with changes in innate rather than adaptive

immunity in patients with inflammatory bowel disease. *Gut* **68**, 25–39 (2019).

10. Wang, X., Park, J., Susztak, K., Zhang, N. R. & Li, M. Bulk tissue cell type deconvolution

with multi-subject single-cell expression reference. *Nat Commun* **10**, 380 (2019).

11. Smillie, C. S. *et al.* Intra- and Inter-cellular Rewiring of the Human Colon during

Ulcerative Colitis. *Cell* **178**, 714-730.e22 (2019).

12. Marigorta, U. M. *et al.* Transcriptional risk scores link GWAS to eQTLs and predict

complications in Crohn's disease. *Nat Genet* **49**, 1517–1521 (2017).

13. Haberman, Y. *et al.* Pediatric Crohn disease patients exhibit specific ileal transcriptome

and microbiome signature. *J. Clin. Invest.* **124**, 3617–3633 (2014).

14. Quraishi, M. N. *et al.* A Pilot Integrative Analysis of Colonic Gene Expression, Gut

Microbiota, and Immune Infiltration in Primary Sclerosing Cholangitis-Inflammatory

Bowel Disease: Association of Disease With Bile Acid Pathways. *Journal of Crohn's and*

Colitis **14**, 935–947 (2020).

15. Sommer, F., Nookaew, I., Sommer, N., Fogelstrand, P. & Bäckhed, F. Site-specific

programming of the host epithelial transcriptome by the gut microbiota. *Genome Biol*

16, 62 (2015).

16. The Tabula Muris Consortium *et al.* Single-cell transcriptomics of 20 mouse organs

creates a Tabula Muris. *Nature* **562**, 367–372 (2018).

17. Guan, Q. A Comprehensive Review and Update on the Pathogenesis of Inflammatory

Bowel Disease. *Journal of Immunology Research* **2019**, 1–16 (2019).

18. Coccia, M. *et al.* IL-1β mediates chronic intestinal inflammation by promoting the

accumulation of IL-17A secreting innate lymphoid cells and CD4+ Th17 cells. *Journal of*

Experimental Medicine **209**, 1595–1609 (2012).

19. Chotikatum, S., Naim, H. Y. & El-Najjar, N. Inflammation induced ER stress affects

absorptive intestinal epithelial cells function and integrity. *International*

Immunopharmacology **55**, 336–344 (2018).

20. Lin, Y. *et al.* Non-hematopoietic STAT6 induces epithelial tight junction dysfunction and

promotes intestinal inflammation and tumorigenesis. *Mucosal Immunology* **12**, 1304–

1315 (2019).

21. He, L. *et al.* Gut Epithelial Vitamin D Receptor Regulates Microbiota-Dependent Mucosal

Inflammation by Suppressing Intestinal Epithelial Cell Apoptosis. *Endocrinology* **159**,

967–979 (2018).

22. Marchal-Bressenot, A. *et al.* Development and validation of the Nancy histological index

for UC. *Gut* **66**, 43–49 (2017).

23. Moor, A. E. *et al.* Spatial Reconstruction of Single Enterocytes Uncovers Broad Zonation

along the Intestinal Villus Axis. *Cell* **175**, 1156-1167.e15 (2018).

24. Wei, X. *et al.* Extensive jejunal injury is repaired by migration and transdifferentiation of

ileal enterocytes in zebrafish. *Cell Reports* **42**, 112660 (2023).

25. López-Novoa, J. M. & Nieto, M. A. Inflammation and EMT: an alliance towards organ

fibrosis and cancer progression. *EMBO Mol Med* **1**, 303–314 (2009).

26. Harvey, R. F. & Bradshaw, J. M. A Simple Index Of Crohn's-Disease Activity. *The Lancet*

315, 514 (1980).

27. R Core Team. *R: A Language and Environment for Statistical Computing*. (R Foundation

for Statistical Computing, Vienna, 2022).

28. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using

lme4. *Journal of Statistical Software* **67**, 1–48 (2015).

29. Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. lmerTest Package: Tests in Linear

Mixed Effects Models. *Journal of Statistical Software* **82**, 1–26 (2017).

30. Fox, J. & Weisberg, S. *An R Companion to Applied Regression*. (Sage, Thousand Oaks CA,

2019).

31. Wickham, H. *Ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag, New York,

2016).

32. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

33. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29

(2021).

Declarations

Ethics approval and consent to participate

All research complied with relevant ethical regulations, and usage of human tissue material and transcriptome data was approved by the ethics committee of the Medical Faculty at Kiel University (A156/03, A124/14, A102/16).

Consent for publication

Not applicable

Availability of data and material

Not applicable. All data was retrieved (see method section).

Competing interests

SS reports indirect stock ownership in Gerion Biotech GmbH as well as consulting and personal fees from AbbVie, Allergosan, Amgen, Arena, BMS, Biogen, Celltrion, Celgene, Falk, Ferring, Fresenius, Galapagos/Gilead, HIKMA, I-Mab, Janssen, Lilly, Morphic, MSD, Mylan, Pfizer, Prometheus, Protagonist, Provention Bio, Sandoz/Hexal, Takeda, and Theravance. PR reports stock ownership in Gerion Biotech GmbH and consulting fees from Takeda.

All other authors declare no competing interests.

Funding

This work was supported by the German Research Foundation (DFG) through the individual grant SO1141/10-1, the Research Unit FOR5042 "miTarget - The Microbiome as a Target in Inflammatory Bowel Diseases" (project P5), the Excellence Cluster EXS2167 "Precision Medicine in Chronic Inflammation" and an intramural grant of the medical faculty of Kiel University (grant no K126408) to FS. The funding bodies had no part or influence on the design of the study and data collection, analysis, or interpretation.

Authors' contributions

447 SW, LJ and FS designed research. SW, JT, LJ and FS performed experiments and analyzed the
448 data. SS, CR, KA and PR provided samples and RNAseq data. SW, JT, LJ and FS prepared the
449 figures. KA, PR, CK and FS obtained funding. SW, JT, LJ and FS co-wrote the manuscript with
450 critical input from all authors. All authors read and approved the final manuscript.

Acknowledgements

453 The authors thank Sabine Kock, Stefanie Baumgarten, Maren Reffelmann, Vivian Wegner,
454 Tanja Klostermeier, Dorina Ölsner, Meike Hansen, Ronja Möhring and Sophie Reiher for
455 excellent technical assistance. We further thank Florian Tran and Finn Hinrichsen for valuable
456 input regarding the description of the clinical cohort characteristics.

Authors' information (optional)

Not applicable

Figures and tables

Figure 1: Epithelial HK2 expression correlates with inflammation severity. A-D) Expression
(TPM, transcript per million) of *HK2* **(A)** and the epithelial marker genes *ECAD* **(B)**, *VIL1* **(C)** and
EPCAM **(D)** in the intestinal mucosa of patients with various degrees of gut inflammation.
Note that at high inflammation scores (>0.4) expression of *HK2* and the epithelial marker
genes all decrease indicating epithelial erosion. The inflammation score was calculated as a
scaled Harvey-Bradshaw index (Crohn's disease) or Mayo score (ulcerative colitis) to
accommodate both disease types. **E)** *HK2* expression increases with inflammation scores after
normalization to epithelial marker gene expression. The red lines represent the mean
expression trendline. The number in the upper right corner represents the p value for the
correlation between gene expression and inflammation score as determined by linear mixed
model.

Figure 2: Expression of candidate immune marker genes increases with inflammation severity. Expression of selected marker genes, which are characteristic for individual immune cell types, were evaluated in patient biopsies. Expression of virtually all candidate immune marker genes increase with inflammation severity indicating an expansion of immune cells – a known feature of intestinal inflammation.

Figure 3: Deconvolution of mucosal transcriptomes demonstrates cellular remodeling during inflammation. A) MuSiC cell deconvolution package was used to estimate ratios of the most abundant cell types based on the mucosal patient transcriptomes. Notably, epithelial cell abundance is decreasing with inflammation, whereas abundances of macrophages and B cells, the two most frequent immune cell types that combine for approximately 35 % proportion, are increasing with inflammation severity. Therefore, cell deconvolution supports the candidate gene analyses indicating mucosal cellular remodeling characterized by a reduction of epithelial cells and an expansion of immune cells with increasing inflammation severity. **B)** *HK2* expression increases with inflammation scores after normalization to epithelial cell abundance.

Figure 4: Epithelial HK2 protein expression increases with inflammation severity. Colonic mucosa biopsies of unrelated UC patients were stained for HK2, epithelial cells (ECAD) and nuclei (DAPI). All biopsies were analyzed for classical histological signs of inflammation using the Nancy Score (see method section). **A)** Quantification of HK2 and ECAD signal intensity in apical intestinal epithelium. Note that HK2 and ECAD level decrease with higher Nancy Scores. **B)** HK2 protein expression normalized to ECAD levels. Note that after epithelial normalization HK2 expression increases with inflammation severity, i.e. Nancy Score. Significance testing was performed using one-way ANOVA compared to Score = 0. **C)** Representative images of the multiplex immunofluorescence stainings (HK2, ECAD, DAPI). Scale bar indicates 50 μm. n = 7 – 10 per group. Note that microscopic images were taken from areas with rather intact epithelium for a better visualization of the mucosal structure.

Figure 5: *HK2* is mainly expressed by mature enterocytes and increased during
inflammation. *HK2* expression was analyzed in published single cell RNA sequencing data
derived from mucosal biopsies of UC patients[11]. **A)** *HK2* expression per cell type. **B)**
Abundances of cell types expressing *HK2*. **C)** *HK2* expression in mature & immature
enterocytes and goblet cells divided into samples from healthy controls, inflamed and non-
inflamed mucosa. **D)** Abundances of cell types expressing *HK2* after stratification into disease
group.

Figure 6: Model of HK2 expression changes in relation to intestinal inflammation severity.
HK2 is predominantly expressed by apical epithelial cells and HK2 expression in these cells increases with disease severity. However, during the course of inflammation the cellular composition of the mucosa changes. In severely inflamed tissue, the epithelium is disrupted, brush border epithelial cells with high levels of HK2 are shed and lost, whereas immune cells with little to no HK2 expression are recruited. Therefore, overall HK2 expression is reduced in bulk tissue samples under severe inflammation despite local high expression in remaining HK2-positive epithelial cells. Created with BioRender.com

522 **Figure S1: HK2 expression in the intestinal mucosa.** Biopsies from small and large intestine

523 were immuno-stained to localize the HK2 protein in the mucosa. Note that HK2 expression is

524 mainly confined to epithelial cells of the apical mucosa both in the small and large intestine.

Figure S2: Epithelial *HK2* expression correlates with disease severity in both clinical cohorts.

Expression data (TPM, transcript per million) of *HK2* and the epithelial marker genes *ECAD,*

VIL1 and *EPCAM* in the intestinal mucosa of patients with various degrees of gut inflammation

split per clinical cohort FUTURE and EMED.

Figure S3: Epithelial *HK2* expression correlates with inflammation severity regardless of disease subtype. Expression data (TPM, transcript per million) of *HK2* and the epithelial marker genes *ECAD, VIL1* and *EPCAM* in the intestinal mucosa of patients with various degrees of gut inflammation split per disease subtype CD and UC.

Figure S4: Correlation analysis for HK2 expression and inflammation score. Three different

linear mixed effect models (PatientID as random factor) were fitted to the data. First, all data

points were used to estimate the overall effect of *HK2* expression and inflammation score.

Afterwards the data set was split into "low scores" (inflammation score < 0.5), "high scores"

(inflammation score >0.5) and fitted models for these data subsets. While the first model

showed no significant association between *HK2* expression and inflammation, there is a

positive association at "low scores" and a negative association at "high scores" (see also table

S2) indicating that *HK2* expression increases with inflammation during lower disease scores

and then decreases at higher disease scores.

 Table S1: Cohort clinical characteristics

Cohort	EMED	FUTURE	Combined	CD	UC
Patients	41	16	57	28	29
Samples	191	87	278	141	137
Age in years, min - max (median)	21 - 65 (26)	16 -74 (36)	16 - 74 (35)	16 - 74 (35)	18 - 65 (35)
Sex assigned at birth	57 female 30 male	119 female 72 male	176 female 102 male	95 female 56 male	81 female 56 male
Smoking	5 smoker 17 previous 65 non-smoker	53 smoker 30 previous 108 non-smoker	58 smoker 47 previous 173 non-smoker	40 smoker 25 previous 76 non-smoker	18 smoker 22 previous 97 non-smoker
Age at first diagnosis in years, min - max (median)	6 - 60 (22)	3 -61 (25)	3 - 61 (24)	3 - 61 (22)	13 - 60 (26)
Time since first diagnosis in years, min - max (median)	0 - 20 (4)	0 - 29 (9)	0 - 29 (6)	0 - 27 (8)	0 - 29 (5)
Inflammation score, min - max (median)	0 - 1 (0.375)	0 - 1 (0.625)	0 - 1 (0.4375)	0 - 1 (0.5)	0 - 1 (0.375)

547 **Table S2: Comparison of linear mixed model statistics for the coefficients explaining HK2**

548 **expression with inflammation scores for models created with all data, low (score <0.5) or**

549 **high (score >0.05) inflammation scores only.**

	Estimate	Std. Error	DF	t-value	p-value	
All data	44.73	32.07	179.94	1.395	0.165	
Low score	195.86	59.48	154.44	3.293	0.0123	**
High score	-178.6	78	114.87	-2.29	0.0239	*

551 **Table S3: Translation of the HBI and Mayo Score into a general Inflammation Score.** The

552 definitions of the HBI and Mayo scores are listed. The Inflammation Score was calculated by

553 scaling each the HBI and Mayo Score from 0-1 and then merging the scores.

HBI		
Criterium	**Score**	
General well-being	Very well	0
	Slightly poor	1
	Poor	2
	Very poor	3
	Terrible	4
Abdominal pain	None	0
	Mild	1
	Moderate	2
	Severe	3
Abdominal mass	None	0
	Dubious	1
	Definite	2
	Definite with tenderness	3
Stool frequency	Number of liquid stools	1 for each lquid stool per day
Complications	Arthralgia, Uveitis, Erythema nodosum, Apthous ulcers, Pyoderma gangrenosum, Anal fissure, New fistula, Abscess	1 point each

Mayo Score		
Criterium	**Score**	
Number of stools per day	Normal for the patient	0
	1-2 stools more than normal	1
	3-4 stools more than normal	2
	5+ stools more than normal	3
Rectal blood	No blood	0
	Blood streaks in less the half of evacuations	1
	Evidence of fresh blood in most of the evacuations	2
	Bowel movements with fresh blood	3
Endoscopy	Normal or inactive disease	0
	Mild disease (erythema, decreased vascular pattern, mild friability)	1
	Moderate disease (marked erythema, lack of vascular pattern, friability, erosions)	2
	Severe disease (spontaneous bleeding, ulceration)	3
Global medical asessment	Normal	0
	Mild disease	1
	Moderate disease	2
	Severe disease	3

555
556

Calculation of inflammation score								
HBI	InfScore	Mayo	InfScore	>>>	InfScore	HBI	Mayo
0	0.00	0	0.00	merged	0.00	0	0
1	0.06	1	0.08		0.06	1	
2	0.12	2	0.17		0.08		1
3	0.18	3	0.25		0.12	2	
4	0.24	4	0.33		0.17		2
5	0.29	5	0.42		0.18	3	
6	0.35	6	0.50		0.24	4	
7	0.41	7	0.58		0.25		3
8	0.47	8	0.67		0.29	5	
9	0.53	9	0.75		0.33		4
10	0.59	10	0.83		0.35	6	
11	0.65	11	0.92		0.41	7	
12	0.71	12	1.00		0.42		5
13	0.76				0.47	8	
14	0.82				0.50		6
15	0.88				0.53	9	
16	0.94				0.58		7
17	1.00				0.59	10	
					0.65	11	
					0.67		8
					0.71	12	
					0.75		9
					0.76	13	
					0.82	14	
					0.83		10
					0.88	15	
					0.92		11
					0.94	16	
					1.00	17	12

559 **Table S4: Original and pooled deconvolution cell types.** Originally classified cell types that

560 are ontogenetically related, e.g. Paneth cells, enterocytes, goblet cells and stem cells, were

561 pooled and classified into "epithelial cells" to reduce cell type diversity.

Original cell type description	Grouped into cell type
TA 1	epis
TA 2	epis
Enterocyte Progenitors	epis
Immature Goblet	epis
Immature Enterocytes 1	epis
Immature Enterocytes 2	epis
Cycling TA	epis