Protoxylem microtubule patterning requires ROP pattern co-alignment, realistic microtubule-based nucleation, and sufficient microtubule flexibility.

Bas Jacobs[1], Marco Saltini[1], Jaap Molenaar[1], Laura Filion[2], Eva E. Deinum[1*]

[1*]Mathematical and Statistical Methods (Biometris), Plant Science Group, Wageningen University & Research, Wageningen, 6708 PB, The Netherlands.
[2]Soft Condensed Matter group, Debye Institute for Nanomaterials Science, Utrecht University, Utrecht, 3584 CC, The Netherlands.

*Corresponding author(s). E-mail(s): eva.deinum@wur.nl;

Abstract

The development of the water transporting xylem tissue in plants involves an intricate interplay of Rho of Plant proteins (ROPs) and cortical microtubules to generate highly functional secondary cell wall patterns, such as the ringed or spiral patterns in early-developing protoxylem. Simulation models have been crucial in understanding the self-organisation of the highly dynamic cortical microtubule array, suggesting important roles for microtubule-dependent nucleation and co-alignment of the microtubule array and ROP pattern in speeding up the band formation process. So far, however, no one has been able to reproduce a banded pattern for realistic microtubule behaviour, as past implementations of microtubule-based nucleation created an inhomogeneity problem that has only recently been solved. Here, we study the requirements of protoxylem band formation with computer simulations in corticalSim, extended to include finite microtubule persistence length and a novel algorithm for microtubule-based nucleation that allows for local feedback on microtubule density but avoids global competition for nucleation events. We find that microtubule-flexibility is required to increase the mismatch tolerance of the ROP and microtubule pattern to realistic levels. At the same time, flexibility leads to more catastrophes, both induced catastrophes and catastrophes from the microtubule-hostile gap regions. This effect results in a loss of microtubule density, making it harder to maintain microtubule bands. Microtubule-dependent nucleation helps to counteract this effect

by moving some nucleations from the gap regions to the bands once microtubules disappear from the gaps. Our results reveal the main mechanisms required for efficient protoxylem band formation.

Keywords: Plant cortical microtubules, Microtubule persistence length, Microtubule nucleation, Protoxylem, Stochastic simulation, corticalSim

1 Background

To meet the water demand of the above ground organs, plants depend on their water transporting tissue, the xylem. Xylem consists of an interconnected tubular network that stretches from young root tips to the above ground water sinks. Inside the xylem, the water pressure is often negative [1]. This means that the constituting elements have to be sufficiently strong to prevent collapse [2]. This is extra challenging close to the root tips. The earliest maturing xylem vessels, called protoxylem, mature while the surrounding tissue is still elongating [3]. This elongation precludes homogeneously thick cell walls. Therefore, protoxylem elements show banded or spiralled secondary cell wall reinforcements. These bands provide strength [4], but also allow for elongation. In contrast, the later maturing metaxylem has much more solid secondary cell wall reinforcements, with a number of ellipsoid gaps [5]. More general, these two xylem types are studied as a model system for pattern formation in cell wall structure [6].

The patterned secondary cell wall reinforcements initially consist mostly of cellulose microfibrils deposited by cellulose synthase (CESA) complexes [5, 7]. Upon maturation, they are additionally lignified [8]. CESA delivery to the membrane and their subsequent movement through it is guided by the cortical microtubule array [9–12], with the cellulose microfibrils in the cell wall themselves as a secondary directing mechanism in absence of microtubule contact [13]. Understanding how the characteristic protoxylem cell wall patterns form, therefore, requires understanding how the corresponding patterns arise in the cortical microtubule array.

Microtubule dynamics during xylem patterning can be studied by ectopic expression of the transcriptional master regulators VASCULAR NAC DOMAIN (VND) 6 and 7, which induce metaxylem and protoxylem like patterns, respectively [14, 15]. In these systems, the cortical microtubule array adopts the pattern that the subsequent secondary cell wall reinforcements will follow [16, 17]. This pattern is formed in interaction with Rho-of-Plants (ROP) proteins and their downstream effectors. In metaxylem, active AtROP11 accumulates in future gap regions, where it recruits MICROTUBULE DEPLETION DOMAIN 1 (MIDD1) and Kinesin-13A, leading to local microtubule depolymerization [18–20]. In protoxylem, a similar interaction between ROPs and microtubules is highly likely, and striated ROP and MIDD1 patterns have been observed [17, 21]. Many ROPs are expressed in the zone of protoxylem patterning [22], however, and the ROPs responsible for the protoxylem pattern remain elusive as even *AtROP7/8/11* triple knockouts still form a banded protoxylem pattern [17], possibly due to a high degree of redundancy. Microtubules themselves also influence the shape of the ROP pattern by anisotropically restricting active ROP diffusion [19, 23], which

can orient the ROP pattern along the microtubules [24]. This coupling results in a degree of co-alignment between the orientation of the original microtubule array and the banded ROP pattern, particularly in highly aligned arrays. Previous simulations suggest that this co-alignment increases the speed of pattern formation [16], but it has not been thoroughly quantified to what extent this effect is required. Consequently, we do not know the likelihood that individual arrays meet this requirement in practice.

Individual microtubules are highly dynamic, particularly at their plus-end (figure 1A). They display phases of growth and rapid shrinkage, with stochastic switches between the two so-called catastrophe and rescue, respectively [25–27]. As the cortical microtubules are attached to the inside of the cell membrane, they are bound to interact via frequent collisions. The outcome of these collisions depends on the relative angle of the colliding and obstructing microtubule [28]. For small angles, the colliding microtubule bundles with the obstructing one, while for large angles it either crosses over or undergoes and induced catastrophe [28]. After a crossover (figure 1B), the latest arriving microtubule, i.e., on the cytoplasmic side, is likely to be severed by katanin at a later time (figure 1B) [29, 30]. Computer simulations and theoretical models have been indispensable in understanding how the above interactions can lead to the spontaneous self-organization of the cortical array into highly aligned patterns [31, 32]. As computer simulations are more easily amended to complex situations, they have been the approach of choice in studying protoxylem development [16, 33]. Full reproduction of the patterning process with interacting microtubules turned out to be a hard problem and has not been achieved yet [16].

One major reason is related to microtubule nucleation. Most microtubules in the cortical array are nucleated from existing microtubules with a specific distribution of relative nucleation angles [34]. The most common implementation of microtubule based nucleation [35, 36], however, introduces a global competition for nucleation, leading to highly inhomogeneous arrays [33] and aggregation of microtubule density in one or few bands of the protoxylem pattern only [16, 33]. Recently, it has been shown that this so-called "inhomogeneity problem" is a consequence of an incomplete understanding of the nucleation process [33]. A more realistic nucleation algorithm based on detailed experimental observations was indeed sufficient to produce regular banded protoxylem arrays in a simplified context of transverse, non-interacting microtubules [33]. The question is whether a similar algorithm [37] will be sufficient to produce timely band formation in the full interacting microtubule array.

Another potentially important aspect of microtubules that is often disregarded in modelling studies is their flexibility (figure 1G). Most cortical array studies model microtubule segments in between collision points as perfectly straight line [16, 32, 38–40], justified by a millimetre range persistence length of isolated microtubules [41]. In microscopic images of plant cells, however, microtubules appear less straight, e.g. [16, 42–45], suggesting a lower effective persistence length in the cell context. Unfortunately, the actual persistence length in plant cells has never been quantified yet, and the recent studies that do model microtubules as semiflexible polymers [46, 47], base themselves on measurements from animal cells [48, 49]. The values they use are similar to the values found *in vitro* for microtubules with reduced persistence length due to the addition MAP70-5 [41], suggesting that persistence length values from this

context are too low for transverse arrays and protoxylem bands. MAP70-5 is thought to be essential for the formation of microtubule rings around the ellipsoid gaps in metaxylem and induces spontaneous formation of microtubule rings *in vitro* [41]. The effect of flexibility on protoxylem patterning remains an open question. Flexibility may both help and hinder the patterning process, as incorrectly oriented microtubules may find their way back to a band, while correctly oriented ones may curve out of it.

Here, we study the effect of three different concepts on protoxylem microtubule patterning: (1) a co-alignment between microtubules and the microtubule-hostile future band regions likely specified by ROPs, (2) realistic microtubule-based nucleation through a recently developed computationally efficient nucleation algorithm that captures the critical aspect of localized positive feedback [37], and (3) a realistic degree of microtubule flexibility.

2 Methods

2.1 Microtubule simulations

We performed our simulations using an extended version of the cortical microtubule simulation software 'corticalSim' [50], fast, event-driven software for simulating cortical microtubule dynamics and interactions on the cell surface (cortex) [16, 32, 36, 38, 50]. Unless stated otherwise, we used a cylindrical geometry with dimensions representative of a developing protoxylem cell (a height of 60 μm and a radius of 7.5 μm). The edges of the cylinder induced catastrophes to favour a transverse array orientation [51]. Simulated microtubules showed dynamic instability by switching between growing and shrinking states (Fig. 1A). Microtubule collisions could result in bundling, crossovers, and induced catastrophes depending on relative collision angles (Fig. 1B). A constant rate of severing the overlaying microtubules at intersections represented katanin activity (Fig. 1C). For a detailed overview of all parameter values, see Appendix A.

2.2 Microtubule nucleation

In many microtubule simulation studies [38, 46, 52] new microtubules appear by isotropic nucleation, i.e, with random, uniformly distributed locations and orientations (Fig. 1D). Because of this frequent use, we use isotropic nucleation for comparison. Otherwise, we use that most microtubules are nucleate from existing microtubules [34] with the specific angle distribution as modelled in [36]. For this, we use a recently developed nucleation algorithm dubbed "local density-dependent nucleation" [37] that effectively approximates the diffusion of nucleation complexes from their insertion into the plasma membrane to the point where they either dissociate or nucleate. Specifically, the algorithm inserts nucleation complexes at random, uniformly distributed positions within the simulation domain at a rate of r_{ins}. Then, three outcomes are possible: the complex dissociates, it results in a microtubule-bound nucleation, or it results in an unbound nucleation. Based on empirical observation [33, 53], a complex reaching the lattice of a microtubule dissociates with probability 76%, while a freely diffusing complex dissociates with probability 98% based on a constant dissociation rate and a increased nucleation rate on microtubules (for calculations, see [33]). From

Fig. 1 Microtubule simulation details. (A) Microtubule plus ends grow or shrink at constant rates v^+ or v^-, respectively, and minus ends retract at constant rate v^{tm}. Spontaneous catastrophes (switch from growing to shrinking) occur at a rate r_c and rescues (switch from shrinking to growing) at a rate r_r. (B) Microtubule-microtubule collision outcomes depend on the collision angle. At angles below θ^* the impinging microtubule bundles with, i.e., continues growing along, the obstructing one. At greater collision angles, the incoming microtubule undergoes an induced catastrophe with a probability P_{cat} and crosses over the other microtubule otherwise [38, 50]. (C) Katanin severs the overlying microtubule at crossovers at a constant rate r_x per crossover. (D) With isotropic nucleation new microtubules appear at a constant rate r_n, in a uniformly random location and direction. (E) With local density-dependent nucleation, new microtubules appear in a way that more realistically mimics the behaviour of nucleation complexes (NCs), taking into account their insertion, movement and potential microtubule binding. (F) In addition, nucleation complexes stay occupied for a duration $t_{occupied}$, temporarily reducing their global availability. (G) For simulations with semiflexible microtubules, a finite persistence length is simulated with deflections in the microtubule growth direction. In addition, microtubules in bundles follow their bundle around bends below an angle θ_b ('bundle tracking'). Deflection angles in cartoons are exaggerated for visibility. (H) In band formation simulations, an initial transverse array is artificially enforced by drawing nucleation angles in the first half hour (α_{init}) from a normal distribution with an average of α_{bias} and a variance of α_{noise}. (I) Protoxylem band formation is simulated with predefined band and gap regions, where the catastrophe rate in the gap regions is increased by a factor f_{cat} after a 2h initiation phase, following [16], except that f_{cat} is reduced to 3.

each insertion point, $n = 6$ possible linear paths (or metatrajectories) are generated (Fig. 1E). These paths have a maximum distance of $d_{\max} = 1.5\mu$m and are oriented in random, uniformly distributed directions with equal angular separation. A path that intersects a microtubule lattice at a distance d identifies a potential nucleation site. The probability that a nucleation complex reaches a site this far from its insertion point without nucleating before is:

$$p_{\text{bound}}(d_i) = e^{\dfrac{-r_u d_i^2}{4D}} ,$$ (1)

where r_u represents the rate of unbound nucleation, and D is the nucleation complex diffusion coefficient at the membrane. Therefore, the probability of a potential isotropic nucleation event follows:

$$p_{\text{iso}} = 1 - \frac{1}{n} \sum_j p_{\text{bound}}(d_j),$$ (2)

with the summation over j including only those paths that intersect a lattice of a microtubule. After this partitioning, the above rejection probabilities are applied. Isotropic nucleations are all executed at the insertion point. In case of a microtubule-based nucleation, new microtubules are nucleated parallel, antiparallel, or branched to either side with an angular distribution with the mode at $35°$ with respect to the parent microtubule, exactly as in [36], based on the experimental data by [34].

To accommodate the observation that the total number of nucleation complexes is relatively constant, which leads to an increase of the local rate per surface area in bands when the pattern starts emerging [16], we modify the above algorithm by introducing nucleation complex occupancy. Upon successful nucleation, a nucleation complex becomes occupied for a duration $t_{occupied} = 60s$ (Fig. 1F), which is the average time until a nucleation complex is released from a new microtubule by katanin [53]. The instantaneous insertion rate is reduced proportional to the fraction of available complexes by rejection of a fraction of attempted nucleations.

2.3 Microtubule flexibility

We extended our model with semiflexible microtubules in a way similar to [46], i.e., by abrupt changes of the microtubule growth direction, but with larger and random intervals between these "deflections". Deflection step sizes were drawn from an exponential distribution with mean $\bar{l}$ and deflection angles were drawn uniformly from $[-m, m]$, where m is the maximum deflection angle. Deflection angles with absolute values lower than a minimal deflection angle q were set to zero to prevent numerical artefacts. With these parameters we could control persistence length p in a computationally tractable way within the simulation environment. To prevent microtubules from leaving bundles at every deflection point, we made microtubules follow their bundle along bends smaller than $10°$ (Fig. 1G). For further implementation details, see Appendix B.

2.4 Protoxylem simulations

Experimental observations of protoxylem development show that microtubule patterning starts from a well-established transversely oriented array [16, 45]. We, therefore, started our simulations with a two hour initiation phase without band-gap differences, with a biased orientation of microtubule nucleations during the first half an hour similar to [16]. In this biased phase, all microtubules were nucleated at a variable angle α_{init}, drawn from a normal distribution with a mean of α_{bias} and a variance of α_{noise} (Fig. 1H). After this first half an hour, the nucleation mode was switched to the one that would be used in the rest of the simulation. After the two hour initiation phase, we simulated protoxylem band formation by imposing predefined gap regions, where the catastrophe rate was increased by a factor f_{cat}, and band regions where it was not (Fig. 1I), similar to simulations by Schneider et al. [16]. These predefined gap regions represented the local activity of proteins specifying the banded pattern, most likely ROPs and their downstream effectors [17, 19, 21].

2.5 Expected mismatch angles

Although a ROP pattern is expected to follow the general orientation of the initial microtubule array, this match may not be exact, as the ROP pattern needs to wrap smoothly around the geometry, while maintaining an intrinsic band spacing [24]. We have previously shown [24] that the orientation of a spiral ROP pattern that maintains the distance between bands follows:

$$\vartheta = \arcsin\left(\frac{H \cdot n}{W \cdot n_{bands}}\right), \tag{3}$$

where H is the domain length, W the domain circumference, n the spiral number (1 for a single spiral, 2 for a double spiral, etc.), and n_{bands} is the number of bands in a banded array. For our 10 bands, this equation gives a set of discreet angles that a ROP pattern is likely to follow. Assuming that the ROP pattern will always adopt the orientation closest to that of the microtubule array (reasonable at least for low spiral numbers; see [24]), microtubule arrays with orientations in between these discreet spiral angles will have the mismatch shown in Fig. 2E.

3 Results

3.1 Strong co-alignment of the microtubule array with a pre-existing band pattern facilitates rapid microtubule band formation

When starting with a microtubule array that is strongly co-aligned with the orientation of the band regions, we found that microtubule bands could form rapidly, both for isotropic nucleation (Fig. S.3 and S.4) and for local density-dependent nucleation (Fig. 2A–D). A slight mismatch of 3.6° (0.02π rad) in this co-alignment, however, already resulted in an extremely slow band formation process for both nucleation modes, similar to simulations in [16] with isotropic nucleations.

Fig. 2 Fast protoxylem patterning is sensitively dependent on co-alignment between microtubules and the underlying pattern. (A) Snapshots from protoxylem simulations using starting arrays with bias angles α_{bias} of 0.5π (90°), 0.51π (91.8°), 0.52π (93.6°), and 0.53π (95.4°) in the first half hour with only minor deviations ($\alpha_{noise} = 0.01\ rad^2$). Histograms below showing local microtubule density ρ share the same axis within a time series, but not between time series. (B) Average microtubule density in the band regions. (C) Number of populated bands, defined as bands with a microtubule density greater than three times the average density in the gaps. (D) Ratio of density in gaps and bands. Quantities in (B–D) were calculated from 100 simulations. The band formation phase starts at $t = 2h$, i.e., at the end of the grey area. Lines indicate the average and shaded areas the standard deviation. (E) Minimal expected mismatch angle of a spiral (or banded) ROP pattern following a microtubule array of different orientations based on geometrical constraints for a distance of 6 μm between the centres of bands and a cylindrical domain with a radius of 7.5 μm. Dotted line indicates the largest mismatch angle for which a microtubule can still span the circumference within a band. (F) Largest mismatch angle α^* at which a microtubule (bundle) can fit within a band along the entire circumference of the cell. Black dot indicates default simulation values. (G) Largest length that a microtubule (bundle) can have while still fitting entirely within a band region at varying mismatch angles. Band width in simulations is 1 μm (orange line). Dashed line indicates cell circumference for comparison.

We observed that with a mismatch up to 1.8° (0.01π rad), the microtubule density in bands was maintained, whereas with mismatches of at least 3.6°, this density was lost at the beginning of the patterning process after which bands were "rediscovered" at a slow rate (Fig. 2B). These results suggest that timely microtubule band formation requires co-alignment between the initial microtubule array and the underlying pattern proposed to be formed by ROPs.

Some degree of co-alignment between the initial microtubule array and this underlying ROP pattern is biologically realistic, since the orientation of the microtubule array helps shape the orientation of the ROP pattern [19, 24]. However, the orientation of a ROP pattern is also influenced by geometrical constraints, as it has to form either rings or spirals that wrap around the circumference [24]. This requirement on the ROP pattern creates a minimal expected mismatch with the initial array, which, unlike the ROP pattern, is not limited to a discrete set of possible orientations. From this consideration, we calculated a minimal expected mismatch that can be as high as 3.6° (Fig. 2E). Our simulations suggested that this mismatch was too large for band formation in without full breakdown and rediscovery (Fig. 2A–D).

As a proxy for calculating the maximum tolarable mismatch angle, we calculated the largest mismatch angle for which a straight microtubule could still span the circumference once while staying within the band (Fig. 2F) as well as the largest stretch of microtubule that could fit in a band given the mismatch angle (Fig. 2G). The point at which band formation began to suffer from the mismatch was similar to the point at which a single microtubule (bundle) could no longer stay in a band region along the entire circumference of the cell (Fig. 2F). For greater mismatch angles the largest length of microtubule (bundle) that could fit within a band without bending rapidly decreased (Fig. 2G). Variation of cell circumference and band width within the biologically relevant range suggests that the co-alignment requirement for straight microtubules is too strict to guarantee timely band formation.

The effect of co-alignment alone, therefore, could not explain timely band formation, and the use of more realistic microtubule nucleations could not improve upon this effect.

3.2 Microtubule flexibility can lead to density loss from bands under isotropic nucleation

Previous models with microtubule flexibility [46, 47] used values of $p = 20$–30 μm measured *in vivo* in animal cells [48, 49]. For microtubule dynamics parameters based largely on measurements in developing protoxylem [16], these persistence length values caused so many extra collisions that it resulted in a loss of density and alignment even in simulations without band formation (Fig. S.5A–C). Possibly, microtubule persistence lengths may have lower values in (often smaller) animal cells. Evidence from *in vitro* measurements in gliding assays suggests persistence lengths of 60 μm and 100 μm for microtubules with many and few interactions, respectively, while measurements of microtubules suspended in flow cells gave persistence lengths of around 2 mm for non-interacting microtubules [41]. Since the deflections due to microtubule interactions are modelled directly in our simulations, it is not just the 'intrinsic' persistence length that contributes to the measured persistence length and so the measured persistence

length will be smaller than the value we use to determine deflections. Therefore, we investigated multiple larger persistence lengths, up to the millimetre range consistently measured in *in vitro* experiments with individual microtubules [54]. For persistence lengths of hundreds of micrometers, aligned arrays did form (Fig. S.5A–C). Curiously, for these larger persistence lengths, the edge-induced catastrophes were not always sufficient to give the array a transverse orientation (Fig. S.5D). For our study of protoxylem band formation we therefore continued using the biased initiation phase.

Band formation was actually hindered by semiflexible microtubules when using isotropic nucleation. For $p = 100$ or $200 \, \mu m$, stable starting arrays could be formed, but density was lost when band formation started (Fig. 3A and S.6). It would seem, therefore, that the extra flexibility, rather than helping microtubules find bands, actually makes microtubules already in bands bend out and suffer from the increased catastrophe rate in the gap regions. Proper bands only formed for more rigid microtubules with $p = 500$ or $1000 \, \mu m$.

The loss of density in the band regions at the lower persistence lengths could at least partially be counteracted when a significant portion of isotropic nucleations was moved to the gap regions (Fig. S.7). Such a shift could be expected to occur dynamically in cells as gap regions start to empty, when taking into account microtubule-bound nucleations, suggesting that a more realistic implementation of nucleations could be essential to band formation when taking microtubule flexibility into account.

3.3 Local density-dependent microtubule-based nucleation helps to keep bands populated with semiflexible microtubules, even for misaligned starting arrays

When combining semiflexible microtubules with local density-dependent nucleation, band formation improved for $p = 100 \, \mu m$, but most microtubule density was still initially lost, and took a long time to recover. However, band formation was now possible at $p = 200 \, \mu m$, lower than for isotropic nucleation, as nucleations were automatically allocated to the denser band regions (Fig. 3B–E).

Furthermore, the combination of semiflexible microtubules and local density-dependent nucleation also greatly improved timely band formation for a significant mismatch between the orientations of the starting array and the band pattern. A mismatch as high as 18° (0.1π rad) in the angle of the nucleations in the initiation phase still yielded a partially banded pattern after five hours of band formation (Fig. 4A–D). The actual mismatch of the starting array was, in practice, somewhat reduced, as we used boundary conditions promoting transversely oriented arrays. Consequently, the initial mismatch of $\alpha_{bias} = 18°$ 0.1π rad in the initiation phase nucleation angle yielded a measured mismatch of about 10.8° (0.06π rad) in the orientation of the starting array (Fig. 4F), substantially more than the worst mismatches we expected theoretically (Fig. 2E).

A reason for the increased tolerance for mismatches might be that the semiflexible microtubules inherently cover more different angles, of which a substantial portion could align with the band regions, even though the average orientation does not. This would give the array more opportunities to correct its course, by microtubules bending back into the array, or by 'wrong' orientations getting catastrophes in gap regions and

Fig. 3 At lower persistence lengths too much density is lost to maintain bands. This problem is less severe local density-dependent nucleation than with isotropic nucleation. (A,B) Snapshots from protoxylem simulations with isotropic and local density-dependent nucleation for different microtubule persistence lengths. Starting arrays were obtained with transverse nucleations in the first half hour ($\alpha_{bias} = 0.5\pi$, $\alpha_{noise} = 0.01$ rad^2). Histograms below showing local microtubule density ρ are plotted on the same scale within a time series, but not between time series. (C) Average microtubule density in the band regions. (D) Number of populated bands, defined as bands with a microtubule density greater than three times the average density in the gaps. (E) Ratio of density in gaps and bands. Quantities in (C–E) were calculated from 100 simulations. The band formation phase starts at $t = 2h$, i.e., at the end of the grey area. Lines indicate the average and shaded areas the standard deviation.

rescues at a point where the orientation better matches the underlying ROP pattern. An investigation of the angles at the start of the band formation process in simulations with microtubule flexibility showed that there is indeed a broad distribution of microtubule segment angles (Fig. 4E).

Fig. 4 With local density-dependent nucleation (Fig. 1), greater mismatches in the initial orientation still allow fast band formation. (A) Snapshots from protoxylem simulations for $p = 200 \ \mu m$ with local density-dependent nucleation using starting arrays with different bias angles α_{bias} in the first half hour with only minor deviations ($\alpha_{noise} = 0.01 \ \text{rad}^2$). Histograms below showing local microtubule density ρ are plotted on the same scale within a time series, but not between time series. (B) Average microtubule density in the band regions. (C) Number of populated bands, defined as bands with a microtubule density greater than three times the average density in the gaps. (D) Ratio of density in gaps and bands. (E) Distribution of microtubule segment angles, weighted by segment length, at $t = 2h$ from the individual example simulations shown in (A). Dashed lines indicate the the overall array orientation. (F) Average array orientation at $t = 2h$ as a function of the bias angle in the initiation phase. Quantities in (B–D), and (J) were calculated from 100 simulations. The band formation phase starts at $t = 2h$, i.e., at the end of the grey area. Lines indicate the average and shaded areas the standard deviation.

4 Discussion

We have identified three important requirements for microtubule patterning in developing protoxylem by doing extensive simulations: (1) co-alignment between the microtubule array and the underlying ROP pattern, (2) microtubule flexibility, and (3) realistic microtubule-based nucleation. Together, these aspects allow microtubule bands to form on a realistic time scale in fully interacting microtubule array simulations with idealised microtubule dynamics parameter values based on measured data [16].

One important aspect that we did not model explicitly is the interplay between microtubule patterning and ROP patterning. Here, we modelled the ROP pattern as a static, banded prepattern, whereas in reality, it is most likely shaped by a reaction diffusion process in which a combination of positive feedback activation and a difference in diffusion between active and inactive ROPs leads to spontaneous pattern formation (we extensively reviewed the details of such processes in plants elsewhere [55]). This ROP patterning has been modelled for both metaxylem [56] and protoxylem [24]. For protoxylem, the formation of the banded ROP pattern involves microtubules acting as ROP diffusion barriers that orient the pattern [24]. We partly modelled this orienting effect by generating starting arrays with various degrees of co-alignment. However, while real microtubule arrays generally align in a transverse orientation before the start of protoxylem patterning, they rarely have exactly the same orientation along the entire length [16, 45]. The resulting bands also tend to have a variation in their orientation and they are not all perfectly straight or equally spaced in ways that reflect initial densities and orientations in the starting arrays [16, 45]. These observations suggest that the underlying ROP pattern also provides co-alignment at a local level, which may help microtubule band formation, given moderate flexibility to follow the curved features. At the same time, the possibility of curved bands results in an increased rate of straight-growing microtubules leaving band regions, which we found could be detrimental to band formation. To help growing microtubules follow these curved bands, microtubule associated proteins (MAPs) involved in microtubule bundling may be important, e.g., MAP65-8, which is expressed in developing xylem [57]. In addition, microtubules must be sufficiently flexible to follow these curves.

In our simulations, we showed the importance of microtubule flexibility even when assuming straight band and gap regions. Flexibility improved the ability of microtubules to follow predefined band regions in spite of small mismatches in the orientation of the microtubules and the band regions. However, our simulations also pointed at a trade-off, where increased flexibility means more microtubules curve into gap regions where they are more likely to suffer catastrophes. Here, too, microtubule bundling proteins may be important to help prevent this problem.

The need for microtubule flexibility becomes even more obvious when we consider metaxylem patterning, where arrays need to form roughly circular gaps. In microscopic pictures, microtubules appear to curve around these gaps [58]. The gapped structure also means that microtubule patterning cannot rely on co-alignment with the ROP pattern as in protoxylem patterning. Therefore, metaxylem patterning may require additional proteins to help the microtubules form this structure that are absent or less important in protoxylem patterning. Recent evidence suggests involvement of

the microtubule-associated protein Map70-5, which lowers microtubule persistence lengths, enabling the formation of microtubule loops [41]. This is consistent with earlier observations that MAP70-5 lines the borders of the pits and its overexpression increases the ratio of pitted to spiral cell wall patterns [59]. CORD proteins may also be involved. These proteins disorder microtubules by partially detaching them from the membrane [58], possibly facilitating corrections in the microtubule orientation. Other potential factors include BDR1, Wallin, and actin networks that form in the pits, though these may only act in a later stage, during formation of the pit borders [60].

In addition to the importance of microtubule flexibility, our simulations also showed the importance of realistic microtubule-based nucleation. We have previously shown the importance of locally saturating microtubule-based nucleation for array homogeneity in [33]. There, we assumed a constant, uniform supply of nucleation complexes as an important source of local saturation. Here, we relaxed this assumption by allowing microtubules to draw from a global pool of nucleation complexes, while maintaining a local density-dependence of the nucleation rate. Consequently, the reduction of microtubule density in the gap regions would free up nucleation complexes that could then be used to boost the nucleation rate in the bands. This effect helped compensate for microtubule loss from catastrophes suffered by microtubules leaving the band region. This partial shift in the location of nucleation complexes is in line with microscopic observations of nucleation complexes during protoxylem development [16] and upon oryzalin treatment [33].

Another interesting finding was that the introduction of a finite persistence length made it harder to obtain global alignment from edge-induced catastrophes alone. This global alignment problem already occurred with rigid microtubules on large domains [61], but microtubule flexibility reduced the domain size at which locally aligned patches formed. Microtubule flexibility, therefore, increases the importance of local orienting cues, such as the observed sensitivity of microtubules to mechanical stress [62–66]. Alternatively, for alignment in a longitudinal orientation, a reduction in the number of membrane linkers may contribute, allowing microtubules to minimise their bending energy along longer stretches [67, 68].

There is a large gap between the microtubule persistence lengths of 30-100 μm measured in gliding assays with and without Map70-5 [41], which are comparable to the 20-30 μm measured in animal cells [48, 49] and those in the millimetre range measured in flow cells [41]. This difference suggests a large uncertainty in the value of the microtubule persistence length in plant cells, as well as a possibility for cells to modulate it. In absence of sufficient membrane attachment, strong microtubule bending can be induced by forces generated by active cellulose synthase complexes [69] and cytoplasmic streaming [70, 71]. Therefore, good candidates for persistence length modulation are proteins involved in microtubule-membrane linkage, such as CMU (cellulose-microtubule uncoupling) proteins [69], and certain IQD proteins [72], of which IQD13 functions in metaxylem development [23]. IQD proteins can also be regulated dynamically, in particular through calcium signalling, as they have calmodulin-binding domains [73, 74]. This kind of regulation may dynamically influence microtubule persistence length. Therefore, persistence lengths measured in one

situation may not necessarily apply to the next, making it necessary to obtain *in vivo* persistence length measurements for plant cells of different types.

5 Conclusion

In conclusion, we have shown that co-alignment, microtubule flexibility, and local density-dependent nucleation are important aspects of protoxylem patterning. Our findings lay the groundwork for future studies on patterns generated by microtubule-ROP interactions. These studies may include simulations of other systems, such as metaxylem, as well as simulations that combine existing ROP models (e.g., [24]) with microtubule simulations.

Appendix A Simulation details and parameter values

Simulations were performed with an extended version of 'corticalSim', a well established and fast two-dimensional microtubule simulation platform [16, 32, 36, 38, 50]. To account for the katanin severings at crossover intersections that may have been interpreted as induced catastrophes [32] in the original experiments by Dixit and Cyr [28], the probability P_{cat} that collisions at large angles result in catastrophes was lowered compared to previous studies [36, 38]. Contrary to those studies, we included katanin severing by default, with a rate of 0.023 s^{-1} per crossover, similar to values found in experiments [29, 44, 75].

Protoxylem simulations were performed with 1 μm wide band regions separated by 5 μm wide gap regions as in Schneider et al. [16]. Simulations started with a 2h initiation phase without bands followed by a 5h or 33h band formation phase in which the catastrophe rate in the gap regions was increased by a factor $f_{cat} = 3$, which in experimental observations tends to be achieved or exceeded during a substantial part of the patterning process [16]. In the first 0.5h of the initiation phase, nucleations were distributed uniformly and given an angle α_{bias}, with a small amount of normally distributed noise, with a variance α_{noise} of 0.01 rad^2. For simulations with rigid microtubules, this biased phase has almost no collisions that could lead to induced catastrophes or crossovers. To compensate the nucleation rate for this phase was reduced by a factor 4 for simulations with rigid microtubules. The remainder of the initiation phase used the same nucleation mode as the band formation phase. Simulations without bands were run for 7h starting from an empty array with additional isotropic nucleations added at the beginning speed up the population of the array. These 'seeded' nucleations were added at a density of 0.1 $\mu m^{-2}s^{-1}$ and a rate of 0.003 s^{-1} as in Lindeboom et al. [40]. See Table A1 for default parameter values.

For calculating nucleation parameters, we aimed at an overall target nucleation rate $r_{n,target}$ for homogeneous arrays of 0.001 nucleations per μm^2 per s, consistent with previous work [16, 33, 36]. As a first estimate, we assumed that the fraction of microtubule-bound nucleations in a fully populated array would be negligible.

From there, we calculate back to a parameter $r_{n,max}$ (the nucleation rate when all nucleation complexes are free and all nucleations microtubule-bound). For this,

Table A1 Default parameter values used in the simulations.

Parameter	Value	Unit	Description	Source
Simulation domain				
H	60	μm	Domain length	[16]
W	$7.5 \cdot 2\pi$	μm	Domain circumference	[16]
Dynamic instability				
v^+	0.05	$\mu m/s$	Growth speed	[16]
v^-	0.08	$\mu m/s$	Shrinkage speed	[16]
v^{tm}	0.01	$\mu m/s$	Treadmilling speed	[16]
r_c	0.0016	s^{-1}	Catastrophe rate (bands)	[16]
r_r	0.001	s^{-1}	Rescue rate	[16]
Microtubule interaction				
θ_c	40°		Collision angle where outcome switches from bundling to crossover or induced catastrophe	[28]
P_{cat}	0.09		Fraction of induced catastrophes	[32]
r_x	0.023	s^{-1}	Crossover severing rate	
Nucleation				
r_n	0.001	$\mu m^{-2}s^{-1}$	Nucleation rate for isotropic nucleation	[16]
$r_{n,target}$	0.001	$\mu m^{-2}s^{-1}$	Nucleation rate for a populated homogeneous array with local density-dependent nucleation	[16]
r_{ins}	0.018	$\mu m^{-2}s^{-1}$	Nucleation complex insertion rate when all nucleation complexes are free.	
$t_{occupied}$	60	s	Time a nucleation complex stays occupied after a nucleation	[53]
Band formation				
f_{cat}	3		Factor increased catastrophe rate in gaps	
Band width	1	μm	Width of band regions	[16]
Gap width	5	μm	Width of gap regions	[16]
Semiflexible microtubules				
p	200	μm	Persistence length	
$\bar{l}$	$0.01p$	μm	Average deflection step size	
q	0.1°		Minimum deflection angle	
θ_b	10°		Maximum bundle tracking angle	

we use the factor difference f_{rn} between the nucleation rate when all complexes are available and the target nucleation rate. We estimate this rate from the ratio between microtubule-associated insertions in nearly empty oryzalin-treated arrays (0.013 insertions per μm^2 per s) and total, mostly microtubule-associated, insertions in established arrays (0.0037 insertions per μm^2 per s) that were measured in [33]. This approach gives:

$$r_{n,max} = f_{rn} \cdot r_{n,target} = \frac{0.013}{0.0037} \cdot 0.001 = 0.0035 \mu m^{-2}s^{-1}. \tag{A1}$$

With a rejection probablity of 0.76 for microtubule-bound nucleations, we compute a required maximum insertion rate of:

$$r_{ins} = 0.0035/(1 - 0.76) = 0.015 \mu m^{-2}s^{-1} \tag{A2}$$

We further use a fixed duration $t_{occupied}$ of 60s during which a nucleation complex remains occupied upon nucleation, based on an average of 58.9s observed in experimental work [53]. Using our estimates for parameters $r_{n,target}$, $r_{n,max}$, and $t_{occupied}$, we can now calculate the remaining parameter N_{tot}. At nucleation rate $r_{n,target}$, a number of $N_{occ,target}$ nucleation complexes are expected to be occupied, following:

$$r_{n,target} = r_{n,max} \frac{N_{tot} - N_{occ,target}}{N_{tot}}, \tag{A3}$$

where N_{tot} is the total number of nucleation complexes. Using this expression and the fact that the number of occupied complexes depends on the duration of occupancy $t_{occupied}$ and the rate at which they become occupied (i.e., the global nucleation rate), we find the following expression for N_{tot}:

$$N_{tot} = \frac{r_{n,max} \cdot N_{occ,target}}{r_{n,max} - r_{n,target}} = \frac{r_{n,max} \cdot r_{n,target} \cdot t_{occupied} \cdot A}{r_{n,max} - r_{n,target}}, \tag{A4}$$

where A is the total domain area.

From initial simulations of homogeneous arrays, we found that this approach resulted in a realised nucleation rate of about 80% of $r_{n,target}$ (Fig. S.1). To keep the realised nucleation rate close to the target, we increased the insertion rate by 20% to 0.018 $\mu m^{-2} s^{-1}$.

Appendix B Semiflexible microtubules

To implement semiflexible microtubules, we adapted an approach from Mirabet et al. [46]. We gave microtubules deflections in their growth direction every step of variable size l that they grow, with l drawn from an exponential distribution with mean $\bar{l}$. Deflection angles were drawn from a uniform interval $[-m, m]$. Angles of which the absolute value was smaller than the minimum deflection angle $q = 0.1°$ were set to zero to avoid numerical problems. For $\bar{l}$ we chose a value of 1% of the desired persistence length p by default. This value prevents large numbers of very small deflections from resulting in many very similar trajectories that give technical difficulties with the simulation software, while still keeping the step size small relative to the persistence length. For details on how the maximum deflection angle m was chosen to give microtubules a given persistence length p, see the next section.

In our current simulations, microtubule bundles consist of multiple microtubules on the same trajectory, without any space in between. Therefore, there is no distinction between microtubules on the sides, which could still deflect outwards, and microtubules on the inside, that don't have any room to deflect at all. To partially compensate for the lack of deflection restriction of microtubules in bundles, we rejected a portion of deflections in bundles, keeping only $1/(n + 1)$ deflections, where n is the number of other microtubules in the bundle at the point of the deflection.

Another bundling-related issue is that microtubules in bundles now regularly encounter bend points (deflection points or bundling points) from the microtubules to which they are bundled. Without any adjustment, bundles would regularly separate at

these points. Therefore, we made microtubules track their bundles along bend points. Since bundling events also create bend points and these may have considerably larger angles (up to 40°), we implemented a maximum bundle tracking angle of $\theta_b = 10°$. If a bundle splits with an angle below this value, an incoming microtubule randomly follows one of the bundles, proportional to their occupancy.

For individual microtubules averaged over sufficient length, the length of the average deflection step size $\bar{l}$ does not matter for the persistence length of the total microtubule as long as the appropriate deflection angle is used. However, microtubules in a populated array interact with each other, and so there may well be a difference between many small deflections and fewer larger deflections. Therefore, we tested the effect of different step sizes for the same persistence length, for a range of step sizes feasible in the current simulation setup (Fig. S.2). For the same persistence length, the deflection step size seemed to have little effect on the alignment and orientation of arrays without bands (Fig. S.2D,E). For lower persistence lengths in particular, there seems to be a small effect on the overall density, likely resulting from differences in the rates of encounters that could lead to crossover-severing [32] or induced catastrophes (Fig. S.2A–C). Therefore, the precise way in which microtubules are flexible, may also have some impact on the array as a whole, but the magnitude of this impact on array alignment, orientation, and density is limited.

Appendix C Persistence length calculations

Persistence length p measures how fast the correlation between the orientation of two different points on a microtubule decays with the microtubule length between these points. We use the following definition of persistence length p:

$$\langle r_n \cdot r_{n+k} \rangle = e^{-\frac{L}{2p}}, \tag{C5}$$

where $\langle r_n \cdot r_{n+k} \rangle$ is the average inner product of unit vectors r_i in the direction of the microtubule at points n and $n+k$ and L is the length along the microtubule between these two points.

Since Eq. C5 holds for any two points n and $n+k$, and we are using independent deflections, it is sufficient to look at a single deflection after length l between points n and $n+1$:

$$\langle r_n \cdot r_{n+1} \rangle = e^{-\frac{l}{2p}}. \tag{C6}$$

For a given deflection angle ϑ (Fig. C1A), setting the arbitrary initial angle to 0, r_{n+1} is given by:

$$r_{n+1} = \begin{bmatrix} \cos(\vartheta) \\ \sin(\vartheta) \end{bmatrix}. \tag{C7}$$

For the inner product in Eq. C6, we then get:

$$\langle r_n \cdot r_{n+1} \rangle = \left\langle \begin{bmatrix} 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} \cos(\vartheta) \\ \sin(\vartheta) \end{bmatrix} \right\rangle = \langle \cos(\vartheta) \rangle. \tag{C8}$$

Fig. C1 Semiflexible microtubule implementation details. (A) Simulated microtubules get deflections of angle ϑ every deflection step size l. Angles in cartoon have been exaggerated for visibility. (B) Deflection angles are drawn from a uniform interval $[-m, m]$, with all angles in $[-q, q]$ set to zero. Value of q in graph is exaggerated for visibility. (C) Deflection step sizes are drawn from an exponential distribution with average $\bar{l}$.

We take deflection angles ϑ, drawn from a uniform interval $[-m, m]$, with the smallest angles ($|\vartheta| < q$) set to zero for numerical reasons (Fig. C1B). In that case, we get:

$$\langle\cos(\vartheta)\rangle = \frac{q}{m}\cos(0) + \frac{m-q}{m}\frac{1}{m-q}\int_q^m \cos(\vartheta)d\vartheta = \frac{\sin(m) - \sin(q) + q}{m}. \quad \text{(C9)}$$

Taking deflection step lengths l drawn from an exponential distribution with mean $\bar{l}$ (Fig. C1C), we can now determine the persistence length p for a given value of m:

$$p = \langle p\rangle = \left\langle -\frac{l}{2\ln(\langle r_n \cdot r_{n+1}\rangle)}\right\rangle = \left\langle -\frac{l}{2\ln\left(\frac{\sin(m)-\sin(q)+q}{m}\right)}\right\rangle = -\frac{\bar{l}}{2\ln\left(\frac{\sin(m)-\sin(q)+q}{m}\right)}.$$
$$\text{(C10)}$$

Therefore, if we draw our deflection step sizes from a distribution with average $\bar{l}$, we can solve the boundary value m of the interval from which we draw the deflection angles from the above equation for the desired persistence length p. This means that we can control the microtubule persistence length in our simulations using $\bar{l}$, m, and q as input parameters, with one of m and $\bar{l}$ calculated to obtain the desired p.

Appendix D   Summary statistics

Array alignment was quantified using the two-dimensional nematic order parameter S_2 as commonly used in polymer physics and often used for quantifying cortical microtubule alignment [36, 38, 50]:

$$S_2 = \sqrt{\langle\langle\cos(2\theta)\rangle\rangle^2 + \langle\langle\sin(2\theta)\rangle\rangle^2}, \quad \text{(D11)}$$

where θ is the angle of individual microtubule segments with the x-axis of the simulation domain, and double angular brackets indicate a length-weighted average over

all microtubule segments. Alignment parameter S_2 is zero for a completely isotropic array and one for a perfectly aligned array (ignoring microtubule polarity).

Overall array orientation Θ was also computed as commonly used in simulation studies [36, 50]:

$$\Theta = \arctan\left(\frac{\langle\langle\sin(2\theta)\rangle\rangle}{\langle\langle\cos(2\theta)\rangle\rangle + S_2}\right). \tag{D12}$$

Band counts were calculated as the number of bands with a microtubule density at least three times higher than the average density in the gaps. This measure was selected after testing multiple options for best representing visual inspections of array snapshots and histograms.

Averages and standard deviations of orientations from multiple simulation runs were calculated using the average and standard deviation for angular quantities. The average orientation $\bar{\phi}$ over N simulations is:

$$\bar{\phi} = \frac{1}{2}\arg\left(\frac{1}{N}\sum_{n=1}^{N}e^{i2\phi_n}\right) \tag{D13}$$

where arg is the argument of a complex number, and ϕ_n is the orientation of the n^{th} simulation. The factor 2 in the exponent and factor $1/2$ before the argument correct the fact that we consider orientations without direction (so in $[0, \pi)$ rather than $[0, 2\pi)$). The circular standard deviation sd_ϕ is:

$$sd_\phi = -2\ln\left(\left|\frac{1}{N}\sum_{n=1}^{N}e^{i2\phi_n}\right|\right), \tag{D14}$$

where ln is the natural logarithm and the vertical bars indicate the absolute value of a complex number.

References

[1] Brown, H.R.: The theory of the rise of sap in trees: Some historical and conceptual remarks. Physics in Perspective **15**(3), 320–358 (2013) https://doi.org/10.1007/s00016-013-0117-1

[2] Venturas, M.D., Sperry, J.S., Hacke, U.G.: Plant xylem hydraulics: What we understand, current research, and future challenges. Journal of Integrative Plant Biology **59**(6), 356–389 (2017) https://doi.org/10.1111/jipb.12534 https://onlinelibrary.wiley.com/doi/pdf/10.1111/jipb.12534

[3] Růžička, K., Ursache, R., Hejátko, J., Helariutta, Y.: Xylem development – from the cradle to the grave. New Phytologist **207**(3), 519–535 (2015) https://doi.org/10.1111/nph.13383 https://nph.onlinelibrary.wiley.com/doi/pdf/10.1111/nph.13383

[4] Roumeli, E., Ginsberg, L., McDonald, R., Spigolon, G., Hendrickx, R., Ohtani, M., Demura, T., Ravichandran, G., Daraio, C.: Structure and biomechanics during xylem vessel transdifferentiation in Arabidopsis thaliana. Plants **9**(12) (2020) https://doi.org/10.3390/plants9121715

[5] Turner, S., Gallois, P., Brown, D.: Tracheary element differentiation. Annual Review of Plant Biology **58**(1), 407–433 (2007) https://doi.org/10.1146/annurev.arplant.57.032905.105236 https://doi.org/10.1146/annurev.arplant.57.032905.105236

[6] Xu, H., Giannetti, A., Sugiyama, Y., Zheng, W., Schneider, R., Watanabe, Y., Oda, Y., Persson, S.: Secondary cell wall patterning—connecting the dots, pits and helices. Open Biology **12**(5), 210208 (2022)

[7] Kamon, E., Ohtani, M.: Xylem vessel cell differentiation: A best model for new integrative cell biology? Current Opinion in Plant Biology **64**, 102135 (2021) https://doi.org/10.1016/j.pbi.2021.102135

[8] Barros, J., Serk, H., Granlund, I., Pesquet, E.: The cell biology of lignification in higher plants. Annals of Botany **115**(7), 1053–1074 (2015) https://doi.org/10.1093/aob/mcv046 https://academic.oup.com/aob/article-pdf/115/7/1053/17635489/mcv046.pdf

[9] Paredez, A.R., Somerville, C.R., Ehrhardt, D.W.: Visualization of cellulose synthase demonstrates functional association with microtubules. Science **312**(5779), 1491–1495 (2006) https://doi.org/10.1126/science.1126551 http://science.sciencemag.org/content/312/5779/1491.full.pdf

[10] Gutierrez, R., Lindeboom, J.J., Paredez, A.R., Emons, A.M.C., Ehrhardt, D.W.: Arabidopsis cortical microtubules position cellulose synthase delivery to the plasma membrane and interact with cellulose synthase trafficking compartments. Nature Cell Biology **11**(7), 797–806 (2009) https://doi.org/10.1038/ncb1886

[11] Crowell, E.F., Bischoff, V., Desprez, T., Rolland, A., Stierhof, Y.-D., Schumacher, K., Gonneau, M., Höfte, H., Vernhettes, S.: Pausing of Golgi Bodies on Microtubules Regulates Secretion of Cellulose Synthase Complexes in Arabidopsis. The Plant Cell **21**(4), 1141–1154 (2009) https://doi.org/10.1105/tpc.108.065334 https://academic.oup.com/plcell/article-pdf/21/4/1141/37012423/plcell_v21_4_1141.pdf

[12] Watanabe, Y., Meents, M.J., McDonnell, L.M., Barkwill, S., Sampathkumar, A., Cartwright, H.N., Demura, T., Ehrhardt, D.W., Samuels, A.L., Mansfield, S.D.: Visualization of cellulose synthases in Arabidopsis secondary cell walls. Science **350**(6257), 198–203 (2015) https://doi.org/10.1126/science.aac7446 http://science.sciencemag.org/content/350/6257/198.full.pdf

[13] Chan, J., Coen, E.: Interaction between autonomous and microtubule guidance

systems controls cellulose synthase trajectories. Current Biology **30**(5), 941–947 (2020)

[14] Yamaguchi, M., Goué, N., Igarashi, H., Ohtani, M., Nakano, Y., Mortimer, J.C., Nishikubo, N., Kubo, M., Katayama, Y., Kakegawa, K., Dupree, P., Demura, T.: VASCULAR-RELATED NAC-DOMAIN6 and VASCULAR-RELATED NAC-DOMAIN7 effectively induce transdifferentiation into xylem vessel elements under control of an induction system. Plant Physiology **153**(3), 906–914 (2010) https://doi.org/10.1104/pp.110.154013 http://www.plantphysiol.org/content/153/3/906.full.pdf

[15] Yamaguchi, M., Mitsuda, N., Ohtani, M., Ohme-Takagi, M., Kato, K., Demura, T.: VASCULAR-RELATED NAC-DOMAIN 7 directly regulates the expression of a broad range of genes for xylem vessel formation. The Plant Journal **66**(4), 579–590 (2011) https://doi.org/10.1111/j.1365-313X.2011.04514.x

[16] Schneider, R., Klooster, K.v., Picard, K.L., Gucht, J., Demura, T., Janson, M., Sampathkumar, A., Deinum, E.E., Ketelaar, T., Persson, S.: Long-term single-cell imaging and simulations of microtubules reveal principles behind wall patterning during proto-xylem development. Nature Communications **12**(1), 669 (2021)

[17] Higa, T., Kijima, S.T., Sasaki, T., Takatani, S., Asano, R., Kondo, Y., Wakazaki, M., Sato, M., Toyooka, K., Demura, T., Fukuda, H., Oda, Y.: Microtubule-associated phase separation of MIDD1 tunes cell wall spacing in xylem vessels in Arabidopsis thaliana. Nature Plants **10**(1), 100–117 (2024)

[18] Oda, Y., Iida, Y., Kondo, Y., Fukuda, H.: Wood cell-wall structure requires local 2D-microtubule disassembly by a novel plasma membrane-anchored protein. Current Biology **20**(13), 1197–1202 (2010) https://doi.org/10.1016/j.cub.2010.05.038

[19] Oda, Y., Fukuda, H.: Initiation of cell wall pattern by a rho- and microtubule-driven symmetry breaking. Science **337**(6100), 1333–1336 (2012) https://doi.org/10.1126/science.1222597 http://science.sciencemag.org/content/337/6100/1333.full.pdf

[20] Oda, Y., Fukuda, H.: Rho of plant GTPase signaling regulates the behavior of Arabidopsis kinesin-13A to establish secondary cell wall patterns. The Plant Cell Online **25**(11), 4439–4450 (2013) https://doi.org/10.1105/tpc.113.117853 http://www.plantcell.org/content/25/11/4439.full.pdf

[21] Brembu, T., Winge, P., Bones, A.M.: The small GTPase AtRAC2/ROP7 is specifically expressed during late stages of xylem differentiation in Arabidopsis. Journal of Experimental Botany **56**(419), 2465–2476 (2005) https://doi.org/10.1093/jxb/eri239

[22] Brady, S.M., Orlando, D.A., Lee, J.-Y., Wang, J.Y., Koch, J., Din-neny, J.R., Mace, D., Ohler, U., Benfey, P.N.: A high-resolution root spatiotemporal map reveals dominant expression patterns. Science **318**(5851), 801–806 (2007) https://doi.org/10.1126/science.1146265 http://science.sciencemag.org/content/318/5851/801.full.pdf

[23] Sugiyama, Y., Wakazaki, M., Toyooka, K., Fukuda, H., Oda, Y.: A novel plasma membrane-anchored protein regulates xylem cell-wall deposition through microtubule-dependent lateral inhibition of rho GTPase domains. Current Biology **27**(16), 2522–25284 (2017) https://doi.org/10.1016/j.cub.2017.06.059

[24] Jacobs, B., Molenaar, J., Deinum, E.E.: Robust banded protoxylem pattern formation through microtubule-based directional ROP diffusion restriction. Journal of Theoretical Biology **502**, 110351 (2020) https://doi.org/10.1016/j.jtbi.2020.110351

[25] Desai, A., Mitchison, T.J.: Microtubule polymerization dynamics. Annual Review of Cell and Developmental Biology **13**(1), 83–117 (1997) https://doi.org/10.1146/annurev.cellbio.13.1.83 https://doi.org/10.1146/annurev.cellbio.13.1.83

[26] Aher, A., Akhmanova, A.: Tipping microtubule dynamics, one protofilament at a time. Current Opinion in Cell Biology **50**, 86–93 (2018) https://doi.org/10.1016/j.ceb.2018.02.015

[27] Gudimchuk, N.B., McIntosh, J.R.: Regulation of microtubule dynamics, mechanics and function through the growing tip. Nature Reviews Molecular Cell Biology **22**(12), 777–795 (2021)

[28] Dixit, R., Cyr, R.: Encounters between dynamic cortical microtubules promote ordering of the cortical array through angle-dependent modifications of microtubule behavior. The Plant Cell Online **16**(12), 3274–3284 (2004) https://doi.org/10.1105/tpc.104.026930 http://www.plantcell.org/content/16/12/3274.full.pdf

[29] Zhang, Q., Fishel, E., Bertroche, T., Dixit, R.: Microtubule severing at crossover sites by katanin generates ordered cortical microtubule arrays in Arabidopsis. Current Biology **23**(21), 2191–2195 (2013) https://doi.org/10.1016/j.cub.2013.09.018

[30] Lindeboom, J.J., Nakamura, M., Hibbel, A., Shundyak, K., Gutierrez, R., Ketelaar, T., Emons, A.M.C., Mulder, B.M., Kirik, V., Ehrhardt, D.W.: A mechanism for reorientation of cortical microtubule arrays driven by microtubule severing. Science **342**(6163), 1245533 (2013) https://doi.org/10.1126/science.1245533 http://science.sciencemag.org/content/342/6163/1245533.full.pdf

[31] Deinum, E.E., Mulder, B.M.: Modelling the role of microtubules in plant cell morphology. Current Opinion in Plant Biology **16**(6), 688–692 (2013) https://doi.org/10.1016/j.pbi.2013.10.001

[32] Deinum, E.E., Tindemans, S.H., Lindeboom, J.J., Mulder, B.M.: How selective severing by katanin promotes order in the plant cortical microtubule array. Proceedings of the National Academy of Sciences **114**(27), 6942–6947 (2017) https://doi.org/10.1073/pnas.1702650114

[33] Jacobs, B., Schneider, R., Molenaar, J., Filion, L., Deinum, E.E.: Microtubule nucleation complex behavior is critical for cortical array homogeneity *and* xylem wall patterning. Proceedings of the National Academy of Sciences **119**(50), 2203900119 (2022) https://doi.org/10.1073/pnas.2203900119 https://www.pnas.org/doi/pdf/10.1073/pnas.2203900119

[34] Chan, J., Sambade, A., Calder, G., Lloyd, C.: Arabidopsis cortical microtubules are initiated along, as well as branching from, existing microtubules. The Plant Cell Online **21**(8), 2298–2306 (2009) https://doi.org/10.1105/tpc.109.069716 http://www.plantcell.org/content/21/8/2298.full.pdf

[35] Allard, J.F., Wasteneys, G.O., Cytrynbaum, E.N.: Mechanisms of self-organization of cortical microtubules in plants revealed by computational simulations. Molecular Biology of the Cell **21**(2), 278–286 (2010) https://doi.org/10.1091/mbc.E09-07-0579 http://www.molbiolcell.org/content/21/2/278.full.pdf+html

[36] Deinum, E.E., Tindemans, S.H., Mulder, B.M.: Taking directions: the role of microtubule-bound nucleation in the self-organization of the plant cortical array. Physical Biology **8**(5), 056002 (2011) https://doi.org/10.1088/1478-3975/8/5/056002

[37] Saltini, M., Deinum, E.E.: A realistic algorithm for microtubule-based nucleation reveals large sensitivity to cell geometry of the plant cortical array. bioRxiv (2024) https://doi.org/10.1101/2024.03.25.586463

[38] Tindemans, S.H., Hawkins, R.J., Mulder, B.M.: Survival of the aligned: Ordering of the plant cortical microtubule array. Physical Review Letters **104**, 058103 (2010) https://doi.org/10.1103/PhysRevLett.104.058103

[39] Eren, E.C., Dixit, R., Gautam, N.: A three-dimensional computer simulation model reveals the mechanisms for self-organization of plant cortical microtubules into oblique arrays. Molecular Biology of the Cell **21**(15), 2674–2684 (2010) https://doi.org/10.1091/mbc.e10-02-0136 https://doi.org/10.1091/mbc.e10-02-0136

[40] Lindeboom, J.J., Lioutas, A., Deinum, E.E., Tindemans, S.H., Ehrhardt, D.W., Emons, A.M.C., Vos, J.W., Mulder, B.M.: Cortical microtubule arrays are initiated from a nonrandom prepattern driven by atypical microtubule initiation. Plant Physiology **161**(3), 1189–1201 (2013) https://doi.org/10.1104/pp.112.204057 http://www.plantphysiol.org/content/161/3/1189.full.pdf

[41] Sasaki, T., Saito, K., Inoue, D., Serk, H., Sugiyama, Y., Pesquet, E., Shimamoto,

Y., Oda, Y.: Confined-microtubule assembly shapes three-dimensional cell wall structures in xylem vessels. Nature Communications **14**(6987) (2023)

[42] Fu, Y., Gu, Y., Zheng, Z., Wasteneys, G., Yang, Z.: Arabidopsis interdigitating cell growth requires two antagonistic pathways with opposing action on cell morphogenesis. Cell **120**(5), 687–700 (2005) https://doi.org/10.1016/j.cell.2004.12.026

[43] Chomicki, G., Wightman, R., Turner, S.R.: A specific class of short treadmilling microtubules enhances cortical microtubule alignment. Molecular Plant **9**(8), 1214–1216 (2016) https://doi.org/10.1016/j.molp.2016.05.008

[44] Nakamura, M., Lindeboom, J.J., Saltini, M., Mulder, B.M., Ehrhardt, D.W.: Spr2 protects minus ends to promote severing and reorientation of plant cortical microtubule arrays. J Cell Biol **217**(3), 915–927 (2018)

[45] Schneider, R., Tang, L., Lampugnani, E.R., Barkwill, S., Lathe, R., Zhang, Y., McFarlane, H.E., Pesquet, E., Niittyla, T., Mansfield, S.D., Zhou, Y., Persson, S.: Two complementary mechanisms underpin cell wall patterning during xylem vessel development. The Plant Cell **29**(10), 2433–2449 (2017) https://doi.org/10.1105/tpc.17.00309 http://www.plantcell.org/content/29/10/2433.full.pdf

[46] Mirabet, V., Krupinski, P., Hamant, O., Meyerowitz, E.M., Jönsson, H., Boudaoud, A.: The self-organization of plant microtubules inside the cell volume yields their cortical localization, stable alignment, and sensitivity to external cues. PLOS Computational Biology **14**(2), 1–23 (2018) https://doi.org/10.1371/journal.pcbi.1006011

[47] Durand-Smet, P., Spelman, T.A., Meyerowitz, E.M., Jönsson, H.: Cytoskeletal organization in isolated plant cells under geometry control. Proceedings of the National Academy of Sciences **117**(29), 17399–17408 (2020) https://doi.org/10.1073/pnas.2003184117 https://www.pnas.org/content/117/29/17399.full.pdf

[48] Brangwynne, C.P., MacKintosh, F.C., Weitz, D.A.: Force fluctuations and polymerization dynamics of intracellular microtubules. Proceedings of the National Academy of Sciences **104**(41), 16128–16133 (2007) https://doi.org/10.1073/pnas.0703094104 https://www.pnas.org/content/104/41/16128.full.pdf

[49] Pallavicini, C., Levi, V., Wetzler, D., Angiolini, J., Benseñor, L., Despósito, M., Bruno, L.: Lateral motion and bending of microtubules studied with a new single-filament tracking routine in living cells. Biophysical Journal **106**(12), 2625–2635 (2014) https://doi.org/10.1016/j.bpj.2014.04.046

[50] Tindemans, S., Deinum, E., Lindeboom, J., Mulder, B.: Efficient event-driven simulations shed new light on microtubule organization in the plant cortical array. Frontiers in Physics **2**, 19 (2014) https://doi.org/10.3389/fphy.2014.00019

[51] Ambrose, C., Allard, J.F., Cytrynbaum, E.N., Wasteneys, G.O.: A CLASP-modulated cell edge barrier mechanism drives cell-wide cortical microtubule organization in Arabidopsis. Nature Communications **2**, 430 (2011) https://doi.org/10.1038/ncomms1444

[52] Deinum, E.E., Mulder, B.M.: Modelling the plant microtubule cytoskeleton. In: Morris, R.J. (ed.) Mathematical Modelling in Plant Biology, pp. 53–67. Springer, Cham (2018). https://doi.org/10.1007/978-3-319-99070-5_4

[53] Nakamura, M., Ehrhardt, D.W., Hashimoto, T.: Microtubule and katanin-dependent dynamics of microtubule nucleation complexes in the acentrosomal Arabidopsis cortical array. Nature Cell Biology **12**(11), 1064–1070 (2010) https://doi.org/10.1038/ncb2110

[54] Hawkins, T., Mirigian, M., Selcuk Yasar, M., Ross, J.L.: Mechanics of microtubules. Journal of Biomechanics **43**(1), 23–30 (2010) https://doi.org/10.1016/j.jbiomech.2009.09.005

[55] Deinum, E.E., Jacobs, B.: Rho of Plants patterning: linking mathematical models and molecular diversity. Journal of Experimental Botany **75**(5), 1274–1288 (2023) https://doi.org/10.1093/jxb/erad447 https://academic.oup.com/jxb/article-pdf/75/5/1274/56751586/erad447.pdf

[56] Nagashima, Y., Tsugawa, S., Mochizuki, A., Sasaki, T., Fukuda, H., Oda, Y.: A Rho-based reaction-diffusion system governs cell wall patterning in metaxylem vessels. Scientific Reports **8**(1), 11542 (2018) https://doi.org/10.1038/s41598-018-29543-y

[57] Kubo, M., Udagawa, M., Nishikubo, N., Horiguchi, G., Yamaguchi, M., Ito, J., Mimura, T., Fukuda, H., Demura, T.: Transcription switches for protoxylem and metaxylem vessel formation. Genes & Development **19**(16), 1855–1860 (2005) https://doi.org/10.1101/gad.1331305 http://genesdev.cshlp.org/content/19/16/1855.full.pdf+html

[58] Sasaki, T., Fukuda, H., Oda, Y.: CORTICAL MICROTUBULE DIS-ORDERING1 is required for secondary cell wall patterning in xylem vessels. The Plant Cell (2017) https://doi.org/10.1105/tpc.17.00663 http://www.plantcell.org/content/early/2017/11/13/tpc.17.00663.full.pdf

[59] Pesquet, E., Korolev, A.V., Calder, G., Lloyd, C.W.: The microtubule-associated protein AtMAP70-5 regulates secondary wall patterning in arabidopsis wood cells. Current Biology **20**(8), 744–749 (2010) https://doi.org/10.1016/j.cub.2010.02.057

[60] Sugiyama, Y., Nagashima, Y., Wakazaki, M., Sato, M., Toyooka, K., Fukuda, H., Oda, Y.: A Rho-actin signaling pathway shapes cell wall boundaries in Arabidopsis xylem vessels. Nature Communications **10**(1), 468 (2019)

[61] Deinum, E.E.: Simple models for complex questions on plant development. PhD thesis, Wageningen University, S.l. (2013). https://edepot.wur.nl/259414

[62] Hamant, O., Heisler, M.G., Jönsson, H., Krupinski, P., Uyttewaal, M., Bokov, P., Corson, F., Sahlin, P., Boudaoud, A., Meyerowitz, E.M., Couder, Y., Traas, J.: Developmental patterning by mechanical signals in arabidopsis. Science **322**(5908), 1650–1655 (2008) https://doi.org/10.1126/science.1165594 https://science.sciencemag.org/content/322/5908/1650.full.pdf

[63] Sampathkumar, A., Krupinski, P., Wightman, R., Milani, P., Berquand, A., Boudaoud, A., Hamant, O., Jönsson, H., Meyerowitz, E.M.: Subcellular and supracellular mechanical stress prescribes cytoskeleton behavior in *Arabidopsis* cotyledon pavement cells. eLife **3**, 01967 (2014) https://doi.org/10.7554/eLife.01967

[64] Verger, S., Long, Y., Boudaoud, A., Hamant, O.: A tension-adhesion feedback loop in plant epidermis. eLife **7**, 34460 (2018) https://doi.org/10.7554/eLife.34460

[65] Takatani, S., Verger, S., Okamoto, T., Takahashi, T., Hamant, O., Motose, H.: Microtubule response to tensile stress is curbed by NEK6 to buffer growth variation in the arabidopsis hypocotyl. Current Biology **30**(8), 1491–15032 (2020) https://doi.org/10.1016/j.cub.2020.02.024

[66] Hamant, O., Inoue, D., Bouchez, D., Dumais, J., Mjolsness, E.: Are microtubules tension sensors? Nature Communications **10**(1), 2360 (2019) https://doi.org/10.1038/s41467-019-10207-y

[67] Bachmann, S., Froese, R., Cytrynbaum, E.N.: A buckling instability and its influence on microtubule orientation in plant cells. SIAM Journal on Applied Mathematics **79**(5), 2132–2149 (2019) https://doi.org/10.1137/19M1264588 https://doi.org/10.1137/19M1264588

[68] Tian, T.Y.Y., Macdonald, C.B., Cytrynbaum, E.N.: A stochastic model of cortical microtubule anchoring and mechanics provides regulatory control of microtubule shape. Bulletin of Mathematical Biology **85**(11), 103 (2023)

[69] Liu, Z., Schneider, R., Kesten, C., Zhang, Y., Somssich, M., Zhang, Y., Fernie, A.R., Persson, S.: Cellulose-microtubule uncoupling proteins prevent lateral displacement of microtubules during cellulose synthesis in Arabidopsis. Developmental Cell **38**(3), 305–315 (2016) https://doi.org/10.1016/j.devcel.2016.06.032

[70] Shaw, S.L., Kamyar, R., Ehrhardt, D.W.: Sustained microtubule treadmilling in Arabidopsis cortical arrays. Science **300**(5626), 1715–1718 (2003) https://doi.org/10.1126/science.1083529 http://science.sciencemag.org/content/300/5626/1715.full.pdf

[71] Sainsbury, F., Collings, D.A., Mackun, K., Gardiner, J., Harper, J.D.I., Marc, J.: Developmental reorientation of transverse cortical microtubules to longitudinal directions: a role for actomyosin-based streaming and partial microtubule-membrane detachment. The Plant Journal **56**(1), 116–131 (2008) https://doi.org/10.1111/j.1365-313X.2008.03574.x https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1365-313X.2008.03574.x

[72] Bürstenbinder, K., Möller, B., Plötner, R., Stamm, G., Hause, G., Mitra, D., Abel, S.: The IQD family of calmodulin-binding proteins links calcium signaling to microtubules, membrane subdomains, and the nucleus. Plant Physiology **173**(3), 1692–1708 (2017) https://doi.org/10.1104/pp.16.01743 https://academic.oup.com/plphys/article-pdf/173/3/1692/37240133/plphys_v173_3_1692.pdf

[73] Bürstenbinder, K., Mitra, D., Quegwer, J.: Functions of IQD proteins as hubs in cellular calcium and auxin signaling: A toolbox for shape formation and tissue-specification in plants? Plant Signaling & Behavior **12**(6), 1331198 (2017) https://doi.org/10.1080/15592324.2017.1331198 https://doi.org/10.1080/15592324.2017.1331198

[74] Kölling, M., Kumari, P., Bürstenbinder, K.: Calcium- and calmodulin-regulated microtubule-associated proteins as signal-integration hubs at the plasma membrane–cytoskeleton nexus. Journal of Experimental Botany **70**(2), 387–396 (2018) https://doi.org/10.1093/jxb/ery397 https://academic.oup.com/jxb/article-pdf/70/2/387/27435272/ery397.pdf

[75] Lindeboom, J.J., Nakamura, M., Saltini, M., Hibbel, A., Walia, A., Ketelaar, T., Emons, A.M.C., Sedbrook, J.C., Kirik, V., Mulder, B.M., Ehrhardt, D.W.: CLASP stabilization of plus ends created by severing promotes microtubule creation and reorientation. Journal of Cell Biology **218**(1), 190–205 (2018) https://doi.org/10.1083/jcb.201805047 https://rupress.org/jcb/article-pdf/218/1/190/1379512/jcb_201805047.pdf