1

Originator: Computational Framework Separating Single-Cell RNA-Seq by Genetic and Contextual Information

Thatchayut Unjitwattana[1], Qianhui Huang[2], Yuheng Du[2], Youqi Yang[3], Mengtian Zhou[4], Lana X. Garmire[2]

[1]Department of Biomedical Engineering, University of Michigan, Ann Arbor, MI, USA

[2]Department of Computation Medicine and Bioinformatics, University of Michigan, Ann Arbor, MI, USA

[3]Department of Biostatistics, University of Michigan, Ann Arbor, MI, USA

[4]Department of Statistics, University of Michigan, Ann Arbor, MI, USA

Abbreviations:
CCC: Cell-cell communication
DE: Differential expression
GSEA: Gene set enrichment analysis
PBMC: Peripheral blood mononuclear cells
PDAC: Pancreatic ductal adenocarcinoma
QC: Quality control
scRNA-Seq: Single-cell RNA sequencing

Keywords: scRNA-Seq, blood cell, genetic origin, tissue heterogeneity, placenta, cancer

Abstract

Single-cell RNA sequencing data from complex human tissues are prone to blood contamination in sample preparation, and some comprise cells of different genetic makeups, necessitating rigorous preprocessing and cell filtering prior to the downstream functional analysis. Our proposed new computational framework, Originator, deciphers single cells by genetic origin and separates blood cells from tissue-resident cells. It improves the quality of data analysis, exemplified by pancreatic ductal adenocarcinoma and placenta tissues.

Main

Single-cell RNA sequencing (scRNA-Seq) data analysis of complex (*e.g.* placenta) or inherently mixed (*e.g.* tumor) tissues pose significant challenges in computational biology. For example, in tumor studies, precisely separating blood and tissue-resident immune cells in tumor tissues is crucial for understanding the tumor microenvironment [1]. In placenta tissues interfacing with the mother and fetus, isolating the maternal and fetal cells is pivotal to reveal cellular and immunological differences between them [2]. Moreover, many tissues contain blood vessels, and the resulting scRNA-seq data often include blood cell types (e.g., T cells) having highly similar expression profiles with the same cell types in the resident tissues. Therefore, a rigorous preprocessing step is required to enhance the quality of the data of interest, subsequently improving the downstream analysis. Here, we propose a novel computational pipeline called Originator. It deconvolutes barcoded cells into different origins using inferred genotype information from scRNA-Seq data, as well as separating cells in the blood from those in solid tissues, an issue often encountered in scRNA-Seq experimentation.

Fig. 1. Illustration of Originator framework. The input data are the scRNA-Seq experiment on tissue

sections. (1) data preprocessing and cell type annotation. (2) separating barcoded cells into different

origins by blood vs tissue residents context and optionally by inferred genotype information. (3) using the

results in steps (1) and (2) to dissect tissue heterogeneity. (4) the functional downstream analyses with

respect to cells' origins.

The proposed Originator framework is illustrated in **Fig 1**. Cells undergo standard scRNA-Seq

preprocessing steps, including quality control (QC), normalization, data integration, clustering,

and cell-type annotation. Then, Originator utilizes the differences in the genotype information

from the scRNA-Seq reads to separate the barcoded cells into different (default N=2) genetic

origins, for example, maternal vs. fetal origins. Next, given that many tissues have blood

contamination, the subsequent step separates immune cells by blood vs. tissue-resident context

using the publicly available whole blood scRNA-Seq data as the reference [3]. We validated the

pipeline on artificially mixed data from a PBMCs dataset and a cell mixture dataset containing

three breast cancer lines (T47D, BT474, MCF7), monocytes, lymphocytes, and stem cells

(Supplementary Notes) [4, 5, 6]. As shown by the UMAP plot in **Supplementary Figure 1,**

Originator separates the two compartments highly accurately, with AUC of 0.97, F-1 score of

0.97 and Area under the Precision-Recall Curve (AUCPR) of 0.96 (**Supplementary Table 1)**.

After dissecting the heterogeneity of the scRNA-seq data, various downstream functional

analysis can be performed.

We first applied Originator to a pancreatic ductal adenocarcinoma (PDAC) scRNA-seq dataset

[7]**,** to demonstrate its utility in removing cells that originate from blood. Originator successfully

separates immune cells in the blood from those in the tumor tissue **(Fig 2a),** and mast and

Myeloid-derived suppressor cells (MDSCs) are exclusive in tissues as expected. The immune

cell type proportions of the two compartments are shown in **Fig 2b.** The blood has larger

proportions of macrophages and regulatory T-cells, but fewer T-cells. We then performed the

differential expression (DE) analysis comparing the common immune cell types between the

blood and tumor tissues, and detected significant differences in gene expression in each

immune cell type, especially for macrophage and T-cells **(Fig. 2c, Supplementary Table 2-5)**.

For example, CCL4 and CCL5 expression is higher in T-cells in the tumor tissue compared to

those in the blood (**Supplementary Figure 2a**), consistent with the previous study that these

genes are directly associated with T-cell-inflamed phenotype and antigen-presenting cell

mediated processes in PDAC [8]. We subsequently performed Gene Set Enrichment Analysis

(GSEA) comparing the immune cells' gene expression difference between the blood vs tissue

compartments **(Fig. 2d)**. Toll-like receptor (TLR) signaling pathway and NOD-like receptor

signaling pathway are more active in tissue-resident T-cells and macrophages compared to

those in blood. However, natural killer cell-mediated cytotoxicity signaling pathway is

downregulated in NK cells from the tumor compared to the blood compartment, consistent with

observed NK cell dysfunction in PDAC via the reduced cytotoxic granule components,

granzyme B, and perforin [9]. To demonstrate the impact of blood cell contamination in

functional interpretation, we inferred cell-cell communications CCC in tumor tissues before and

after removing blood cells **(Fig 2e,f).** While the interactions among the cell types are globally

less frequent/noisy in the tumor compartment, those between fibroblast/epithelium and other

cell types are strengthened in blood cells, as reflected by the increased node size of these two

cell types **(Fig 2e)**. Such a trend of change is most drastic for the MIF signaling pathway (**Fig**

2f), which was previously reported to drive the malignant character of pancreatic cancer [10].

Particularly, MIF signaling between T-cells/regulatory T-cells and other cell types (eg. epithelial

cells) is absent after removing blood immune cells, indicating a compromised anti-tumor

immune response in PDAC [11].

Some tissues may have different genetic lineages. For example, the placenta has both maternal

and fetal tissues. We next applied Originator to the human placenta, and separated the single

cells into fetal and maternal origins as well as blood vs. tissue-resident cells. The cells from the

fetal origin account for the majority of the cell populations, and the tissue-resident cells

significantly over-weigh blood cells, as expected **(Fig 2g)**. The framework correctly and

exclusively assigns trophoblast cells, including cytotrophoblasts, syncytiotrophoblasts, and

extravillous trophoblasts, to the fetal tissue (**Fig 2g,2h**) [12]. On the contrary, immune cells,

vascular endothelial cells, and fibroblasts (type 1 and type 2) appear in both maternal and fetal

tissues, also as expected **(Fig 2h)**. We further performed DE analysis on the common cell types

between fetal and maternal tissues and discovered drastic differences in gene expression

related to the local tissue environment **(Fig 2i, Supplementary Table 6-8).** For example, The

top-ranked DE gene EGFL6 between maternal vs. fetal fibroblast cells is highly expressed in

fibroblasts subtype 1 in fetal tissues as expected (**Supplementary Figure 3a**) [13]. However, it

is much lower in fibroblast subtype 1 or mostly absent in fibroblast subtype 2 in maternal

tissues, similar to a previous report [14]. Among the DE genes from macrophage cells

(**Supplementary Table 8**), SEPP1 is expressed at much higher levels in Hofbauer cells

originating from the fetal tissue compared to the maternal tissue (**Supplementary Figure 3b)**, in

accordance with a previous study [15]. We additionally performed GSEA on the DE genes of the

common cell types between fetal and maternal cells in the placenta tissue **(Fig 2j)**. The MAPK

signaling pathway is enriched in vascular endothelial cells from the fetal tissues, which agrees

with its role in growth factor-induced fetoplacental angiogenesis [16]. On the other hand,

fibroblast type 2 from the fetal origin is enriched with several other signaling pathways, such as

calcium, hippo, and Ras signaling pathways as well as ECM-receptor interaction **(Fig 2j),** which

indicates extracellular matrix (ECM) remodeling during trophoblast differentiation [17]. We also

compared the CCC among the common cell types between the fetal and maternal tissues **(Fig**

2k, 2l). Type 1 and type 2 fibroblasts both show different degrees of CCC with some other cell

types when comparing the maternal and fetal tissue contexts. A higher interaction between

fibroblast cells (type 1 and type 2) and vascular endothelial cells in fetal tissues reflects the

active formation of the villous stroma underneath the syncytiotrophoblast and surrounding fetal

capillaries [18]. On the contrary, higher interactions between type 2 fibroblast and macrophages

in maternal tissue may assist trophoblast invasion through growth factors and cytokines [19].

Thus teasing apart the cell types by their genetic and local tissue context helps to refine the

molecular analysis in placenta scRNA-seq data.

In conclusion, we demonstrate that Originator can systematically decipher scRNA-Seq data by

genetic origin and blood/tissue contexts in heterogeneous tissues, exemplified by PDAC and

placenta. It can be used as an effective tool to remove the undesirable blood cells in scRNA-

Seq data. It can also provide improved cell type annotations and other downstream functional

analyses, based on the genetic background.

Fig. 2. Applications of Originator to pancreatic ductal adenocarcinoma (PDAC) and placenta

scRNA-seq data. a, UMAP plot of cells in PDAC tissues (top-left) separated into blood and tissue-

resident immune cells (bottom-left). UMAP plot of cells in PDAC tissues after blood immune cell removal

is shown on the top-right. **b,** The cell type proportion barplot shows the immune cell types in blood and

PDAC tumor tissue, as well as all cell types in the tumor tissue before and after blood immune cell

removal. **c,** Venn diagram of the significant DE genes identified when comparing immune cells in blood

and tumor tissue. **d,** GSEA results comparing blood and tissue-resident immune cell types. **e,** Overall

CCC in the tumor tissues before and after blood immune cell removal. **f,** CCC of MIF signaling pathway

in tumor tissues before and after blood immune cell removal. **g,** UMAP plot of cells in placenta tissues

deciphered into fetal vs. maternal origins and blood vs. placenta-resident cells. **h,** Cell type proportions in

the different compartments related to the placenta. **i,** Upset plot of significant DE genes identified in

common cell types between fetal and maternal placenta tissues. T-cell is excluded due to the near 0

counts in the fetal tissue. **j,** GSEA results comparing fetal and maternal tissues among common cell

types. **k-l,** CCC of common cell types in fetal (**k**) and maternal (**l**) tissues.

Methods
169

scRNA-Seq case study data sets

Two datasets are used in this report. One is a PDAC scRNA-Seq dataset, in which two tumor

tissues are used in this study (GSE212966) [7]. The other is the scRNA-Seq data from placenta

tissues,comprising　eight　placenta　samples　from　the　previous　study　(EGA;

https://www.ebi.ac.uk/ega/) hosted by the European Bioinformatics Institute (EBI; accession no.

EGAS00001002449) [20]. Cell types in the placenta and PDAC tissues are annotated using the

cell-type-specific marker genes (**Supplementary Table 9-10**).

Description of Originator for scRNA-Seq data analysis

Originator is a multi-module framework that can be used to preprocess scRNA-Seq data from

heterogeneous tissues. It consists of one mandatory step based on the tissue vs blood

compartments and also an optional step based on the genetic origins. Originator takes the gene

expression matrix processed by Cell Ranger (version 7.1.0) of 10x genomics as the input [21].

For genetic background based deciphering, it also uses the BAM file. It processes and outputs

the R data serialization file (RDS). This output file contains gene expression and related

information, including the cell type, blood and tissue immune cell annotation, and annotation by

genetic information if this function is needed.

First, Originator separates immune cells in blood vs. those in the tissue using the blood immune

cell scRNA-Seq reference constructed from the publicly available scRNA-seq data from the

whole blood cells [3]. Annotated immune cells of interest (monocytes, macrophages, T-cells,

regulatory T-cells, plasma cells, NK cells, and B-cells) from the datasets of this study are

aligned to the whole blood scRNA-Seq reference data using package Seurat (4.3.0) [22]. For

each immune cell type, the top 10 embeddings from the UMAP analysis of each scRNA-Seq

sample are used to compute a pairwise Euclidean distance matrix between each query immune

cell and the reference whole blood. To separate each query of immune cells into blood or

tissue-resident cells, k-means clustering (k = 2) is applied using Euclidean distances relative to

the whole blood immune cell reference. The cluster more similar to the whole blood immune cell

references is annotated as the blood immune cells, and the other cluster is determined as the

tissue-resident immune cells.

If the tissue contains the mixture of cells of different genetic background, such as maternal and

fetal origins, Originator provides an optional step to decipher the barcoded cells by the genetic

origin relies on the overlapping genetic variants extracted from the scRNA-Seq data. Genetic

variants are extracted by *bcftools* (version 1.17) [23]. Variants informative of genetic origins are

determined by excluding single nucleotide polymorphisms (SNPs) with minor allele frequency

(MAF) greater than x percentages (default x=10), using the 1000 Genomes Phase 3 reference

panel (release date: 2013/05/02) [24]. The *freemuxlet* package is used to separate the single

cells with the parameter '--nsample N' (default N=2), as implemented in popscle

(https://github.com/statgen/popscle) [25]. N denotes the number of genetic origins of the single

cells (e.g., N = 2 for the maternal and fetal origins in a placenta scRNA-Seq data). Mosaic

doublets (a single barcoded cell with a 50% genetic mixture from two individuals) are also

identified in this step as part of the quality control. As trophoblast cells in the placenta are of

fetal origin, the cluster that contains trophoblast cells is identified with the fetal origin, while the

other cluster is marked as the maternal origin.

Simulation

The artificially mixed tissue-blood-resident data was generated to assess the ability of Originator

to separate blood cells and tissue immune residents. We included only mixture 1

(GSM6807287) in the simulated data to reduce the computational cost during the origins and

tissue-blood deconvolution process (**Supplementary Notes**) [6].

Downstream analysis

DE analysis was performed using *FindMarkers()* in the package Seurat (4.3.0) [22]. GSEA is

done using *gseKEGG()* in the package ClusterProfiler (4.8.2) with the gene set information from

the Kyoto Encyclopedia of Genes and Genomes (KEGG) database [26, 27]. CCC inference was

performed using the package CellChat [28].

Code availability

Originator is freely available to use at URL: https://github.com/lanagarmire/Originator

Acknowledgments

TU is supported by the Royal Thai Government fellowship from the Thai Government. LXG is

supported by grants by NIH/NIGMS, R01 LM012373 and R01 LM012907 awarded by NLM, and

R01 HD084633 awarded by NICHD. The authors acknowledge all lab members of Garmire

Group for helpful discussions.

Author Contributions

L.X.G. and Q.H. conceived this project and supervised the study. T.U. modified the pipeline,

carried the analysis, and wrote the manuscript. Q.H. wrote the Originator pipeline. M.Z., Y.Y.,

and Y.D. assisted the analysis and tested the package.

Competing Interests

LXG is a member of the Scientific Advisory Boards of Simulations Plus.

Reference

1. Del Prete, A., & Wu, Q. (2022). Editorial: Tissue-resident immune cells in tumor immunity and

immunotherapy. *Frontiers in Cell and Developmental Biology*, *10*, 1068720.

https://doi.org/10.3389/fcell.2022.1068720

2. Hussain, T., Murtaza, G., Kalhoro, D. H., Kalhoro, M. S., Yin, Y., Chughtai, M. I., Tan, B.,

Yaseen, A., & Rehman, Z. U. (2022). Understanding the Immune System in Fetal Protection

and Maternal Infections during Pregnancy. *Journal of Immunology Research*, *2022*, 1–12.

https://doi.org/10.1155/2022/7567708

3. The Tabula Sapiens Consortium*, Jones, R. C., Karkanias, J., Krasnow, M. A., Pisco, A. O.,

Quake, S. R., Salzman, J., Yosef, N., Bulthaup, B., Brown, P., Harper, W., Hemenez, M.,

Ponnusamy, R., Salehi, A., Sanagavarapu, B. A., Spallino, E., Aaron, K. A., Concepcion, W.,

Gardner, J. M., … Wyss-Coray, T. (2022). The Tabula Sapiens: A multiple-organ, single-cell

transcriptomic atlas of humans. *Science*, *376*(6594), eabl4896.

https://doi.org/10.1126/science.abl4896

4. Hansen, K. D., Risso, D., & Hicks, S. (2023). *TENxPBMCData: PBMC data from 10X*

Genomics [Computer software]. doi:10.18129/

5. 10x Genomics. (2017). *8k PBMCs from a Healthy Donor, Single Cell Gene Expression*

Dataset by Cell Ranger 1.3.0 [dataset]. https://support.10xgenomics.com/single-cell-gene-

expression/datasets/1.3.0/pbmc8k

6. Sumazin, P. (2023). *GSE220606, Effective methods for bulk RNA-seq deconvolution using*

scnRNA-seq transcriptomes [cell mixtures scRNA-seq] [dataset].

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE220606

7. Chen, K., Wang, Q., Liu, X., Tian, X., Dong, A., & Yang, Y. (2023). Immune profiling and

prognostic model of pancreatic cancer using quantitative pathology and single-cell RNA

sequencing. *Journal of Translational Medicine*, *21*(1), 210. https://doi.org/10.1186/s12967-023-

04051-4

8. Romero, J. M., Grünwald, B., Jang, G.-H., Bavi, P. P., Jhaveri, A., Masoomian, M., Fischer,

S. E., Zhang, A., Denroche, R. E., Lungu, I. M., De Luca, A., Bartlett, J. M. S., Xu, J., Li, N.,

Dhaliwal, S., Liang, S.-B., Chadwick, D., Vyas, F., Bronsert, P., … Gallinger, S. (2020). A Four-

Chemokine Signature Is Associated with a T-cell–Inflamed Phenotype in Primary and Metastatic

Pancreatic Cancer. *Clinical Cancer Research*, *26*(8), 1997–2010. https://doi.org/10.1158/1078-

0432.CCR-19-2803

9. Kim, H.-A., Kim, H., Nam, M.-K., Park, J. K., Lee, M.-Y., Chung, S., Lee, K.-M., & Kuh, H.-J.

(2023). Suppression of the antitumoral activity of natural killer cells under indirect coculture with

cancer-associated fibroblasts in a pancreatic TIME-on-chip model. *Cancer Cell International*,

23(1), 219. https://doi.org/10.1186/s12935-023-03064-9

10. Yang, S., He, P., Wang, J., Schetter, A., Tang, W., Funamizu, N., Yanaga, K., Uwagawa, T.,

Satoskar, A. R., Gaedcke, J., Bernhardt, M., Ghadimi, B. M., Gaida, M. M., Bergmann, F.,

Werner, J., Ried, T., Hanna, N., Alexander, H. R., & Hussain, S. P. (2016). A Novel MIF

Signaling Pathway Drives the Malignant Character of Pancreatic Cancer by Targeting NR3C2.

Cancer Research, *76*(13), 3838–3850. https://doi.org/10.1158/0008-5472.CAN-15-2841

11. Bacher, M., Metz, C. N., Calandra, T., Mayer, K., Chesney, J., Lohoff, M., Gemsa, D.,

Donnelly, T., & Bucala, R. (1996). An essential regulatory role for macrophage migration

inhibitory factor in T-cell activation. *Proceedings of the National Academy of Sciences*, *93*(15),

7849–7854. https://doi.org/10.1073/pnas.93.15.7849

12. Red-Horse, K., Zhou, Y., Genbacev, O., Prakobphol, A., Foulk, R., McMaster, M., & Fisher,

S. J. (2004). Trophoblast differentiation during embryo implantation and formation of the

maternal-fetal interface. *Journal of Clinical Investigation*, *114*(6), 744–754.

https://doi.org/10.1172/JCI200422991

13. Suryawanshi, H., Morozov, P., Straus, A., Sahasrabudhe, N., Max, K. E. A., Garzia, A.,

Kustagi, M., Tuschl, T., & Williams, Z. (2018). A single-cell survey of the human first-trimester

placenta and decidua. *Science Advances*, *4*(10), eaau4788.

https://doi.org/10.1126/sciadv.aau4788

14. Tang, C.-T., Zhang, Q.-W., Wu, S., Tang, M.-Y., Liang, Q., Lin, X.-L., Gao, Y.-J., & Ge, Z.-Z.

(2020). Thalidomide targets EGFL6 to inhibit EGFL6/PAX6 axis-driven angiogenesis in small

bowel vascular malformation. *Cellular and Molecular Life Sciences*, *77*(24), 5207–5221.

https://doi.org/10.1007/s00018-020-03465-3

15. Liang, G., Zhou, C., Jiang, X., Zhang, Y., Huang, B., Gao, S., Kang, Z., Ma, D., Wang, F.,

Gottgens, B., Wang, H., Han, J.-D. J., & Liu, F. (2021). De novo generation of macrophage from

placenta-derived hemogenic endothelium. *Developmental Cell*, *56*(14), 2121-2133.e6.

https://doi.org/10.1016/j.devcel.2021.06.005

16. Wang, K., & Zheng, J. (2012). Signaling regulation of fetoplacental angiogenesis. *Journal of*

Endocrinology, *212*(3), 243–255. https://doi.org/10.1530/JOE-11-0296

17. MacPhee, D. J., Mostachfi, H., Han, R., Lye, S. J., Post, M., & Caniggia, I. (2001). Focal

Adhesion Kinase Is a Key Mediator of Human Trophoblast Development. *Laboratory*

Investigation, *81*(11), 1469–1483. https://doi.org/10.1038/labinvest.3780362

18. Palaiologou, E., Etter, O., Goggin, P., Chatelet, D. S., Johnston, D. A., Lofthouse, E. M.,

Doherty, R., Pearson☐Farr, J., Sengers, B. G., Torrens, C., Cleal, J. K., Page, A. M., & Lewis,

R. M. (2020). Human placental villi contain stromal macrovesicles associated with networks of

stellate cells. *Journal of Anatomy*, *236*(1), 132–141. https://doi.org/10.1111/joa.13082

19. Ueshima, C., Kataoka, T. R., Osakabe, M., Sugimoto, A., Ushirokawa, A., Shibata, Y.,

Nakamura, H., Shibuya, R., Minamiguchi, S., Sugai, T., & Haga, H. (2022). Decidualization of

Stromal Cells Promotes Involvement of Mast Cells in Successful Human Pregnancy by

Increasing Stem Cell Factor Expression. *Frontiers in Immunology*, *13*, 779574.

https://doi.org/10.3389/fimmu.2022.779574

20. Tsang, J. C. H., Vong, J. S. L., Ji, L., Poon, L. C. Y., Jiang, P., Lui, K. O., Ni, Y.-B., To, K. F.,

Cheng, Y. K. Y., Chiu, R. W. K., & Lo, Y. M. D. (2017). Integrative single-cell and cell-free

plasma RNA transcriptomics elucidates placental cellular dynamics. *Proceedings of the National*

Academy of Sciences, *114*(37). https://doi.org/10.1073/pnas.1710470114

21. Zheng, G. X. Y., Terry, J. M., Belgrader, P., Ryvkin, P., Bent, Z. W., Wilson, R., Ziraldo, S.

B., Wheeler, T. D., McDermott, G. P., Zhu, J., Gregory, M. T., Shuga, J., Montesclaros, L.,

Underwood, J. G., Masquelier, D. A., Nishimura, S. Y., Schnall-Levin, M., Wyatt, P. W.,

Hindson, C. M., … Bielas, J. H. (2017). Massively parallel digital transcriptional profiling of

single cells. *Nature Communications*, *8*(1), 14049. https://doi.org/10.1038/ncomms14049

22. Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W. M., Zheng, S., Butler, A., Lee, M. J., Wilk,

A. J., Darby, C., Zager, M., Hoffman, P., Stoeckius, M., Papalexi, E., Mimitou, E. P., Jain, J.,

Srivastava, A., Stuart, T., Fleming, L. M., Yeung, B., … Satija, R. (2021). Integrated analysis of

multimodal single-cell data. *Cell*, *184*(13), 3573-3587.e29.

https://doi.org/10.1016/j.cell.2021.04.048

23. Li, H. (2011). A statistical framework for SNP calling, mutation discovery, association

mapping and population genetical parameter estimation from sequencing data. *Bioinformatics*,

27(21), 2987–2993. https://doi.org/10.1093/bioinformatics/btr509

24. The 1000 Genomes Project Consortium, Corresponding authors, Auton, A., Abecasis, G. R.,

Steering committee, Altshuler, D. M., Durbin, R. M., Abecasis, G. R., Bentley, D. R.,

Chakravarti, A., Clark, A. G., Donnelly, P., Eichler, E. E., Flicek, P., Gabriel, S. B., Gibbs, R. A.,

Green, E. D., Hurles, M. E., Knoppers, B. M., … Abecasis, G. R. (2015). A global reference for

human genetic variation. *Nature*, *526*(7571), 68–74. https://doi.org/10.1038/nature15393

25. Zhang, F., & Min Kang, H. (2021). *popscle: A suite of population scale analysis tools for*

single-cell genomics data (freemuxlet) [Computer software]. https://github.com/statgen/popscle

26. Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L.,

Fu, X., Liu, S., Bo, X., & Yu, G. (2021). clusterProfiler 4.0: A universal enrichment tool for

interpreting omics data. *The Innovation*, *2*(3), 100141.

https://doi.org/10.1016/j.xinn.2021.100141

27. Kanehisa, M., & Goto, S. (n.d.). *KEGG: Kyoto Encyclopedia of Genes and Genomes*.

28. Jin, S., Guerrero-Juarez, C. F., Zhang, L., Chang, I., Ramos, R., Kuan, C.-H., Myung, P.,

Plikus, M. V., & Nie, Q. (2021). Inference and analysis of cell-cell communication using

CellChat. *Nature Communications*, *12*(1), 1088. https://doi.org/10.1038/s41467-021-21246-9

Fig. 1. Illustration of Originator framework.

1 Perform scRNA-seq preprocessing including QC, normalization, data integration, clustering, and cell-type annotation

2 Reference

Blood residents   Tissue residents

Separate immune cells into tissue and blood residents using references

optional

Separate barcoded cells into different origins using inferred genotypes

3 Dissect heterogeneity of the tissues in combinations of results from the previous two separation methods

4 Perform downstream analysis

Fig. 2. Applications of Originator to pancreatic ductal adenocarcinoma (PDAC) and placenta scRNA-seq data.