1

Single-cell RNA-Seq data have prevalent blood contamination but can be rescued by Originator, a computational tool separating single-cell RNA-Seq by genetic and contextual information

Thatchayut Unjitwattana[1], Qianhui Huang[2], Yiwen Yang[2], Leyang Tao[1], Youqi Yang[3], Mengtian Zhou[4], Yuheng Du[2], Lana X. Garmire[1,2]*

[1]Department of Biomedical Engineering, University of Michigan, 2200 Bonisteel, Ann Arbor, MI, 48109, USA

[2]Department of Computation Medicine and Bioinformatics, University of Michigan, 100 Washtenaw Ave Ann Arbor, MI, 48109, USA

[3]Department of Biostatistics, University of Michigan, 1415 Washington Heights, Ann Arbor, MI, 48109, USA

[4]Department of Statistics, University of Michigan,1085 S University Ave, Ann Arbor, MI, 48109, USA

*:Corresponding author. Email: lgarmire@umich.edu

Abstract

Single-cell RNA sequencing (scRNA-Seq) data from complex human tissues have prevalent blood cell contamination during the sample preparation process. They may also comprise cells of different genetic makeups. These issues demand rigorous preprocessing and filtering prior to the downstream functional analysis, to avoid biased conclusions due to cell types not of interest. Towards this, we propose a new computational framework, Originator, which deciphers single cells by the genetic origin and separates immune cells of blood contamination from those of expected tissue-resident cells. We demonstrate the accuracy of Originator at separating immune cells from the blood and tissue as well as cells of different genetic origins, using a variety of artificially mixed and real datasets. We show the prevalence of blood contamination in scRNA-Seq data of many tissue types. We alert the significant consequences if failing to adjust for these confounders, using scRNA-Seq data of pancreatic cancer and placentas as examples.

Keywords: scRNA-Seq, blood cell, contamination, genetic origin, tissue heterogeneity, placenta, cancer

Background

Single-cell RNA sequencing (scRNA-Seq) data analysis of complex (*e.g.* placenta) or inherently

mixed (*e.g.* tumor) tissues pose significant challenges in computational biology. Many tissues

contain blood vessels, and the resulting scRNA-seq data often include blood cell types (e.g., T

cells) having highly similar expression profiles with the same cell types in the resident tissues.

For example, in tumor studies precisely separating blood and tissue-resident immune cells in

tumor tissues is crucial for understanding the tumor microenvironment [1]. In placenta tissues

interfacing with the mother and fetus, isolating the maternal and fetal cells is pivotal to reveal

cellular and immunological differences between them [2]. Therefore, a rigorous preprocessing

step is required to enhance the quality of the data of interest, subsequently improving the

downstream analysis. However, currently there is no dedicated tool to perform such a function.

Here we fill in the void and propose a novel computational pipeline called Originator. It

deconvolutes barcoded cells into different origins using inferred genotype information from

scRNA-Seq data, as well as separating cells in the blood from those in solid tissues, an issue

often encountered in scRNA-Seq experimentation.

Fig. 1. Illustration of Originator framework and the prevalence of blood immune cells in tissue samples. a, The input data are the scRNA-Seq experiment on tissue sections. (1) data preprocessing and cell type annotation. (2) separating barcoded cells into different origins by blood vs tissue residents context and optionally by inferred genotype information. (3) using the results in steps (1) and (2) to dissect tissue heterogeneity. (4) the functional downstream analyses with respect to cells' origins. **b,** Heatmap plots show inferred immune cell type proportions originated from the blood (left) vs. the tissue (right), from scRNA-Seq data of a wide-variety of organs including liver, lung, spleen, kidney, pancreas and in the normal or cancer tissues. The cell types were annotated by the original publications and blood vs tissue identification were done by Originator. NA: the cell type does not exist in the original publication.

Results and Discussion

The proposed Originator framework is illustrated in **Fig 1a**. Cells undergo standard scRNA-Seq

preprocessing steps, including quality control (QC), normalization, data integration, clustering,

and cell-type annotation [3]. Then, Originator takes advantages of the differences in the

genotype information from the scRNA-Seq reads, and utilize freemuxlet, a reference-free

version of the popsicle suite [4] to separate the barcoded cells into different (default N=2)

genetic origins, for example, maternal vs. fetal origins. Freemuxlet is chosen, as it showed

better accuracies at recovering the genetic origins in comparison to scSplit [5], in the scRNA-

Seq data of two placentas and the paired clear cell renal cell carcinoma (ccRCC) and PBMC

cells from the same patient (**Supplementary Table 1**). Next, given that many tissues have

blood contamination, the subsequent step separates immune cells by blood vs. expected tissue-

resident context using the publicly available whole blood scRNA-Seq data as the reference [6].

After dissecting the heterogeneity of the scRNA-seq data, various downstream functional

analysis can be performed.

Current public scRNA-Seq data lack paired PBMC (or blood) and complex primary tissue

completely free of blood contamination from the same person. We therefore first validated the

pipeline using the artificially mixed data from a PBMCs dataset and a cell mixture dataset

containing three breast cancer lines (T47D, BT474, MCF7), monocytes, lymphocytes, and stem

cells **(Supplementary Notes)** [7–9]. Before creating the artificial mixture scRNA-Seq data, we

removed the batch effect between the PBMC and the breast cancer cell lines (**Supplementary**

Figure 1), using Harmony [10] which showed good performance for this task previously [11]. As

shown by the UMAP plot in **Supplementary Figure 1,** Originator separates the two

compartments highly accurately, with AUC of 0.96, F-1 score of 0.97 and Area under the

Precision-Recall Curve (AUCPR) of 0.96 (**Supplementary Table 2)**. Next, we tested Originator

on the dataset of Krishna et al. 2021 [12], which contains paired ccRCC tissue and PBMC from

the same patients. To generate the "ground truth" cell types for blood-eliminated ccRCC tissue,

we applied Originator (or Seurat, for comparison) to the ccRCC to remove the potential blood

immune cells from those residing in the tissue. We then integrated the cleaned ccRCC tissues

and PBMC to generate an artificial mixture. Next, we ran Originator on this pre-cleaned mixture

dataset for five iterations, and obtained averaged F-scores of 0.98, 0.93, 0.91, and 0.99 for B

cell, CD4 T-cell, CD8 T-cell and NK cell respectively on the pre-cleaned, mixed dataset

(**Supplementary Figure 2**). We thus conclude that Originator is effective at removing blood

contamination from the results of both datasets above.

To signify the prevalence of blood immune cells in tissue samples, we applied Originator to

eight publicly available scRNA-Seq data from a variety of organs including lung, liver, spleen,

kidney, and pancreas, with normal (or adjacent normal) and/or tumor tissue samples [13–20].

As summarized in **Fig.1b**, in most datasets the majority of the immune cells are from the blood,

rather than the tissue. The detailed side-by-side comparison of UMAP plots of blood vs tissue

immune cells is in **Supplementary Figure 3-4.** These results unambiguously demonstrate the

vast amount of immune cells from blood, rather than the tissue microenvironment. Failure to

remove these immune cells may yield significantly biased results.

To directly demonstrate the potential biases in the downstream analyses due to contamination

of the immune cells from the blood, we first applied Originator to a pancreatic ductal

adenocarcinoma (PDAC) scRNA-seq dataset [20]**,** to demonstrate its utility in removing cells

that originate from blood. Originator successfully separates immune cells in the blood from

those in the tumor tissue **(Fig 2a),** and mast and Myeloid-derived suppressor cells (MDSCs) are

exclusive in tissues as expected. The immune cell type proportions of the two compartments are

shown in **Fig 2b.** The blood has larger proportions of macrophages and regulatory T-cells, but

fewer T-cells. We then performed the differential expression (DE) analysis comparing the

common immune cell types between the blood and those expected from the tumor tissues, and

detected significant differences in gene expression in each immune cell type, especially for

macrophage and T-cells **(Fig. 2c, Supplementary Table 3-6)** [21,22]. For example, CCL4 and

CCL5 expression is higher in T-cells in the tumor tissue compared to those in the blood

(**Supplementary Figure 5a**), consistent with the previous study that these genes are directly

associated with T-cell-inflamed phenotype and antigen-presenting cell mediated processes in

PDAC [23]. We also examined the genes common in some immune cell types, but show

significant differences between the blood vs. tumor compartments (**Supplementary Notes**). We

subsequently performed Gene Set Enrichment Analysis (GSEA) comparing the immune cells'

gene expression difference between the blood vs tissue compartments **(Fig. 2d)**. Toll-like

receptor (TLR) signaling pathway and NOD-like receptor signaling pathway are more active in

expected tissue-resident T-cells and macrophages compared to those in blood. However,

natural killer cell-mediated cytotoxicity signaling pathway is downregulated in NK cells from the

tumor compared to the blood compartment, consistent with observed NK cell dysfunction in

PDAC via the reduced cytotoxic granule components, granzyme B, and perforin [24]. To

demonstrate the impact of blood cell contamination in functional interpretation, we inferred cell-

cell communications CCC in tumor tissues before and after removing blood cells **(Fig 2e,f).**

While the interactions among the cell types are globally less frequent/noisy in the tumor

compartment, those between fibroblast/epithelium and other cell types are strengthened in

blood cells, as reflected by the increased node size of these two cell types **(Fig 2e)**. Such a

trend of change is most drastic for the MIF signaling pathway (**Fig 2f**), which was previously

reported to drive the malignant character of pancreatic cancer [25]. Particularly, MIF signaling

between T-cells/regulatory T-cells and other cell types (eg. epithelial cells) are absent after

removing blood immune cells, indicating a compromised anti-tumor immune response in PDAC

[26].

153
Some tissues may have different genetic lineages that also need to be addressed. For example,

the placenta has both maternal and fetal tissues. We then applied Originator to the human

placenta [27], and separated the single cells into fetal and maternal origins as well as blood vs.

expected tissue-resident cells. The cells from the fetal origin account for the majority of the cell

populations, and the expected tissue-resident cells significantly over-weigh blood cells, as

expected **(Fig 2g)**. The framework correctly and exclusively assigns trophoblast cells, including

cytotrophoblasts, syncytiotrophoblasts, and extravillous trophoblasts, to the fetal tissue (**Fig**

2g,2h) [28]. On the contrary, immune cells, vascular endothelial cells, and fibroblasts (type 1

and type 2) appear in both maternal and fetal tissues, also as expected **(Fig 2h)**. We further

performed DE analysis on the common cell types between fetal and maternal tissues and

discovered drastic differences in gene expression related to the local tissue environment **(Fig 2i,**

Supplementary Table 7-9). For example, The top-ranked DE gene EGFL6 between maternal

vs. fetal fibroblast cells is highly expressed in fibroblasts subtype 1 in fetal tissues as expected

(**Supplementary Figure 6a**) [29]. However, it is much lower in fibroblast subtype 1 or mostly

absent in fibroblast subtype 2 in maternal tissues, similar to a previous report [30]. Among the

DE genes from macrophage cells (**Supplementary Table 9**), SEPP1 is expressed at much

higher levels in Hofbauer cells originating from the fetal tissue compared to the maternal tissue

(**Supplementary Figure 6b)**, in accordance with a previous study [31]. We additionally

performed GSEA on the DE genes of the common cell types between fetal and maternal cells in

the placenta tissue **(Fig 2j)**. The MAPK signaling pathway is enriched in vascular endothelial

cells from the fetal tissues, which agrees with its role in growth factor-induced fetoplacental

angiogenesis [32]. On the other hand, fibroblast type 2 from the fetal origin is enriched with

several other signaling pathways, such as calcium, hippo, and Ras signaling pathways as well

as ECM-receptor interaction **(Fig 2j),** which indicates extracellular matrix (ECM) remodeling

during trophoblast differentiation [33]. We also compared the CCC among the common cell

types between the fetal and maternal tissues **(Fig 2k, 2l)**. Type 1 and type 2 fibroblasts both

show different degrees of CCC with some other cell types when comparing the maternal and

fetal tissue contexts. A higher interaction between fibroblast cells (type 1 and type 2) and

vascular endothelial cells in fetal tissues reflects the active formation of the villous stroma

underneath the syncytiotrophoblast and surrounding fetal capillaries [34]. On the contrary,

higher interactions between type 2 fibroblast and macrophages in maternal tissue may assist

trophoblast invasion through growth factors and cytokines [35]. Thus, teasing apart the cell

types by their genetic and local tissue context helps to refine the molecular analysis in placenta

scRNA-seq data.

Fig. 2. Applications of Originator to pancreatic ductal adenocarcinoma (PDAC) and placenta

scRNA-seq data. a, UMAP plot of cells in PDAC tissues (top-left) separated into blood and expected

tissue-resident immune cells (bottom-left). UMAP plot of cells in PDAC tissues after blood immune cell

removal is shown on the top-right. **b,** The cell type proportion barplot shows the immune cell types in

blood and PDAC tumor tissue, as well as all cell types in the tumor tissue before and after blood immune

cell removal. **c,** Venn diagram of the significant DE genes identified when comparing immune cells in

blood and expected tumor resident tissue. **d,** GSEA results comparing blood and expected tissue-resident

immune cell types. **e,** Overall CCC in the tumor tissues before and after blood immune cell removal. **f,**

CCC of MIF signaling pathway in tumor tissues before and after blood immune cell removal. **g,** UMAP plot

of cells in placenta tissues deciphered into fetal vs. maternal origins and blood vs. placenta-resident cells.

h, Cell type proportions in the different compartments related to the placenta. **i,** Upset plot of significant

DE genes identified in common cell types between fetal and maternal placenta tissues. T-cell is excluded

due to the near 0 counts in the fetal tissue. **j,** GSEA results comparing fetal and maternal tissues among

common cell types. **k-l,** CCC of common cell types in fetal (**k**) and maternal (**l**) tissues.

204   Discussion and conclusions

Originator is the first dedicated systematic tool to decipher scRNA-Seq data by genetic origin

and blood/tissue contexts in heterogeneous tissues. It can be used as an effective tool to

remove the undesirable blood cells in scRNA-Seq data. It can also provide improved cell type

annotations and other downstream functional analyses, based on the genetic background.

Future work will be focused on generating pan-tissue immune cell atlas which are free of

immune cells originating from blood contamination, which will better annotate the expected

tissue-resident immune cells truthfully from the tissues of interest.

Methods

scRNA-Seq case study data sets

scRNA-Seq data from liver, lung, pancreas and kidney are used in this study. For lung, we used

a large scRNA-Seq dataset from the integrated human lung cell atlas (v1.0), containing samples

from 107 individuals [14], as well as lung cancer datasets from Xing et al. and Bischoff et al.

[18,19]. For liver tissues, we used the single-cell liver landscape dataset from five individuals

[16], and another healthy liver dataset from Guilliams et al. [17]. Additionally, we used lung,

spleen, and liver scRNA-Seq datasets from Domínguez Conde et al. [15]. We obtained the

paired blood and kidney cancer dataset from Krishna et al. [13]. For pancreatic ductal

adenocarcinoma (PDAC), we use a scRNA-Seq dataset from GSE212966, which includes two

tumor tissues [20]. The last dataset is the scRNA-Seq data from placenta tissues, comprising

eight placenta samples from the previous study (EGA; https://www.ebi.ac.uk/ega/) hosted by the

European Bioinformatics Institute (EBI; accession no. EGAS00001002449) [27]. Cell types in

the placenta and PDAC tissues are annotated using the cell-type-specific marker genes

[36](**Supplementary Table 10-11**).

Description of Originator for scRNA-Seq data analysis

Originator is a multi-module framework that can be used to preprocess scRNA-Seq data from

heterogeneous tissues. It consists of one mandatory step based on the tissue vs blood

compartments and also an optional step based on the genetic origins. Originator takes the gene

expression matrix processed by Cell Ranger (version 7.1.0) of 10x genomics as the input [37].

For genetic background based deciphering, it also uses the BAM file. It processes and outputs

the R data serialization file (RDS). This output file contains gene expression and related

information, including the cell type, blood and tissue immune cell annotation, and annotation by

genetic information if this function is needed.

First, Originator separates immune cells in blood vs. those in the tissue using the blood immune

cell scRNA-Seq reference constructed from the publicly available scRNA-seq data from the

whole blood cells [6]. Annotated immune cells of interest (monocytes, macrophages, T-cells,

regulatory T-cells, plasma cells, NK cells, and B-cells) from the datasets of this study are

aligned to the whole blood scRNA-Seq reference data using package Seurat (4.3.0) [38]. For

each immune cell type, the top 10 latent variables from each scRNA-Seq sample are used to

compute a pairwise Euclidean distance matrix between each query immune cell and the

reference whole blood. The latent variables can be obtained by UMAP or PCA based dimension

reduction. We compared UMAP-based and PCA-based dimension reduction on the data of

Krishna et al. [13], and found that the former yielded higher accuracies in detecting the immune

cell types (**Supplementary Figure 7**). Thus we used UMAP-based dimension reduction in this

report. To separate each query of immune cells into blood or expected tissue-resident cells, k-

means clustering (default k = 2) is applied using Euclidean distances relative to the whole blood

immune cell reference. The cluster more similar to the whole blood immune cell references is

annotated as the blood immune cells, and the other cluster is determined as the expected

tissue-resident immune cells.

If the tissue contains the mixture of cells of different genetic background, such as maternal and

fetal origins, Originator provides an optional step to decipher the barcoded cells by the genetic

origin relies on the overlapping genetic variants extracted from the scRNA-Seq data. Genetic

variants are extracted by *bcftools* (version 1.17) [39]. Variants informative of genetic origins are

determined by excluding single nucleotide polymorphisms (SNPs) with minor allele frequency

(MAF) greater than x percentages (default x=10), using the 1000 Genomes Phase 3 reference

panel (release date: 2013/05/02) [40]. The *freemuxlet* package is used to separate the single

cells with the parameter '--nsample N' (default N=2), as implemented in popscle

(https://github.com/statgen/popscle) [4]. N denotes the number of genetic origins of the single

cells (e.g., N = 2 for the maternal and fetal origins in a placenta scRNA-Seq data). Mosaic

doublets (a single barcoded cell with a 50% genetic mixture from two individuals) are also

identified in this step as part of the quality control. As trophoblast cells in the placenta are of

fetal origin, the cluster that contains trophoblast cells is identified with the fetal origin, while the

other cluster is marked as the maternal origin.

Testing the performance of Originator

The artificially mixed tissue-blood-resident data were generated to assess the ability of

Originator to separate blood cells and tissue immune residents. We included two datasets: (1)

an artificially mixed data from a PBMCs dataset and a cell mixture dataset containing three

breast cancer lines (T47D, BT474, MCF7), monocytes, lymphocytes, and stem cells [7–9]. (2) A

paired pre-cleaned ccRCC tissues and PBMC from the same patient provided by Krishna et al.

[13]. The details of the evaluation are in **Supplementary Notes.** Additionally, we also

benchmarked freemuxlet [4](used in Originator) against scSplit [5] in identifying cells from

different genetic origins using two different datasets, including (1) scRNA-Seq data of two

placenta samples and (2) mixed PBMC scRNA-Seq data from two ccRCC patients provided by

Krishna et al. [13,27] **(Supplementary Notes)**.

Downstream analysis

DE analysis was performed using *FindMarkers()* in the package Seurat (4.3.0) [38]. GSEA is

done using *gseKEGG()* in the package ClusterProfiler (4.8.2) with the gene set information from

the Kyoto Encyclopedia of Genes and Genomes (KEGG) database [41,42]. CCC inference was

performed using the package CellChat [43].

Code availability

Originator is freely available to use at URL: https://github.com/lanagarmire/Originator

Abbreviations:

CCC: Cell-cell communication

DE: Differential expression

GSEA: Gene set enrichment analysis

PBMC: Peripheral blood mononuclear cells

PDAC: Pancreatic ductal adenocarcinoma

QC: Quality control

scRNA-Seq: Single-cell RNA sequencing

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

We use a healthy lung scRNA-Seq dataset from the integrated human lung cell atlas (v1.0),

containing samples from 107 individuals [14]. Additionally, we use lung, spleen, and liver

scRNA-Seq datasets from Domínguez Conde et al. [15]. For liver tissues, we used the single-

cell liver landscape dataset from five individuals [16], as well as another healthy liver dataset

from Guilliams et al. [17]. For cancer tissues, we incorporated lung cancer datasets from Xing et

al. and Bischoff et al. [18,19], and a kidney cancer dataset from Krishna et al. [13]. For

pancreatic ductal adenocarcinoma (PDAC), we use a scRNA-Seq dataset from GSE212966,

which includes two tumor tissues [20]. The last dataset is the scRNA-Seq data from placenta

tissues, comprising eight placenta samples from the previous study (EGA;

https://www.ebi.ac.uk/ega/) hosted by the European Bioinformatics Institute (EBI; accession no.

EGAS00001002449) [27]. For simulation, we generated the data using an 8k PBMCs dataset

from a healthy donor (blood-resident cells) and an *in vitro* cell mixture (expected tissue-resident

cells) containing three breast cancer lines (T47D, BT474, MCF7), monocytes (Thp1),

lymphocytes (Jukrat), and stem cells (hMSC) [8,9].

Competing interests

LXG is a member of the Scientific Advisory Boards of Simulations Plus.

Funding

TU is supported by the Royal Thai Government fellowship from the Thai Government. LXG is

supported by grants by NIH/NIGMS, R01 LM012373 and R01 LM012907 awarded by NLM, and

R01 HD084633 awarded by NICHD.

Authors' contributions

L.X.G. and Q.H. conceived this project and supervised the study. T.U. modified the pipeline,

carried the analysis, and wrote the manuscript. Q.H. wrote the Originator pipeline. M.Z., Y.Y.,

Y.Y., Y.D. and L.T. assisted the analysis and tested the package.

Acknowledgements

The authors acknowledge all lab members of Garmire Group for helpful discussions.

331 References

1. Del Prete A, Wu Q. Editorial: Tissue-resident immune cells in tumor immunity and

immunotherapy. Front Cell Dev Biol. 2022 Nov 22;10:1068720.

2. Hussain T, Murtaza G, Kalhoro DH, Kalhoro MS, Yin Y, Chughtai MI, et al. Understanding

the Immune System in Fetal Protection and Maternal Infections during Pregnancy. Xu H,

editor. J Immunol Res. 2022 Jun 24;2022:1–12.

3.   Stanojevic S, Li Y, Ristivojevic A, Garmire LX. Computational Methods for Single-Cell Multi-

Omics Integration and Alignment. Genomics Proteomics Bioinformatics. 2022 Oct

1;20(5):836–49.

4.   Kang HM, Zhang F. popscle: A suite of population scale analysis tools for single-cell

genomics data (freemuxlet) [Internet]. 2021. Available from:

https://github.com/statgen/popscle

5.   Xu J, Falconer C, Nguyen Q, Crawford J, McKinnon BD, Mortlock S, et al. Genotype-free

demultiplexing of pooled single-cell RNA-seq. Genome Biol. 2019 Dec;20(1):290.

6.   The Tabula Sapiens Consortium*, Jones RC, Karkanias J, Krasnow MA, Pisco AO, Quake

SR, et al. The Tabula Sapiens: A multiple-organ, single-cell transcriptomic atlas of humans.

Science. 2022 May 13;376(6594):eabl4896.

7.   Hansen K, Risso D, Hicks S. TENxPBMCData: PBMC data from 10X Genomics [Internet].

2023. Available from: https://bioconductor.org/packages/TENxPBMCData

8.   10x Genomics. 8k PBMCs from a Healthy Donor, Single Cell Gene Expression Dataset by

Cell Ranger 1.3.0 [Internet]. 2017. Available from: https://support.10xgenomics.com/single-

cell-gene-expression/datasets/1.3.0/pbmc8k

9.   Sumazin P. GSE220606, Effective methods for bulk RNA-seq deconvolution using scnRNA-

seq transcriptomes [cell mixtures scRNA-seq] [Internet]. 2023. Available from:

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE220606

10.   Korsunsky I, Millard N, Fan J, Slowikowski K, Zhang F, Wei K, et al. Fast, sensitive and

accurate integration of single-cell data with Harmony. Nat Methods. 2019 Dec;16(12):1289–

96.

11.   Luecken MD, Büttner M, Chaichoompu K, Danese A, Interlandi M, Mueller MF, et al.

Benchmarking atlas-level data integration in single-cell genomics. Nat Methods. 2022

Jan;19(1):41–50.

12. Krishna C, DiNatale RG, Kuo F, Srivastava RM, Vuong L, Chowell D, et al. Single-cell

sequencing links multiregional immune landscapes and tissue-resident T cells in ccRCC to

tumor topology and therapy efficacy. Cancer Cell. 2021 May;39(5):662-677.e6.

13. Krishna C, DiNatale RG, Kuo F, Srivastava RM, Vuong L, Chowell D, et al. Single-cell

sequencing links multiregional immune landscapes and tissue-resident T cells in ccRCC to

tumor topology and therapy efficacy. Cancer Cell. 2021 May;39(5):662-677.e6.

14. Sikkema L, Ramírez-Suástegui C, Strobl DC, Gillett TE, Zappia L, Madissoon E, et al. An

integrated cell atlas of the lung in health and disease. Nat Med. 2023 Jun;29(6):1563–77.

15. Domínguez Conde C, Xu C, Jarvis LB, Rainbow DB, Wells SB, Gomes T, et al. Cross-tissue

immune cell analysis reveals tissue-specific features in humans. Science. 2022 May

13;376(6594):eabl5197.

16. MacParland SA, Liu JC, Ma XZ, Innes BT, Bartczak AM, Gage BK, et al. Single cell RNA

sequencing of human liver reveals distinct intrahepatic macrophage populations. Nat

Commun. 2018 Oct 22;9(1):4383.

17. Guilliams M, Bonnardel J, Haest B, Vanderborght B, Wagner C, Remmerie A, et al. Spatial

proteogenomics reveals distinct and evolutionarily conserved hepatic macrophage niches.

Cell. 2022 Jan;185(2):379-396.e38.

18. Xing X, Yang F, Huang Q, Guo H, Li J, Qiu M, et al. Decoding the multicellular ecosystem of

lung adenocarcinoma manifested as pulmonary subsolid nodules by single-cell RNA

sequencing. Sci Adv. 2021 Jan 29;7(5):eabd9738.

19. Bischoff P, Trinks A, Obermayer B, Pett JP, Wiederspahn J, Uhlitz F, et al. Single-cell RNA

sequencing reveals distinct tumor microenvironmental patterns in lung adenocarcinoma.

Oncogene. 2021 Dec 16;40(50):6748–58.

20. Chen K, Wang Q, Liu X, Tian X, Dong A, Yang Y. Immune profiling and prognostic model of

pancreatic cancer using quantitative pathology and single-cell RNA sequencing. J Transl

Med. 2023 Mar 21;21(1):210.

21. Xu J, He B, Carver K, Vanheyningen D, Parkin B, Garmire LX, et al. Heterogeneity of

neutrophils and inflammatory responses in patients with COVID-19 and healthy controls.

Front Immunol. 2022 Nov 16;13:970287.

22. He B, Xiao Y, Liang H, Huang Q, Du Y, Li Y, et al. ASGARD is A Single-cell Guided Pipeline

to Aid Repurposing of Drugs. Nat Commun. 2023 Feb 22;14(1):993.

23. Romero JM, Grünwald B, Jang GH, Bavi PP, Jhaveri A, Masoomian M, et al. A Four-

Chemokine Signature Is Associated with a T-cell–Inflamed Phenotype in Primary and

Metastatic Pancreatic Cancer. Clin Cancer Res. 2020 Apr 15;26(8):1997–2010.

24. Kim HA, Kim H, Nam MK, Park JK, Lee MY, Chung S, et al. Suppression of the antitumoral

activity of natural killer cells under indirect coculture with cancer-associated fibroblasts in a

pancreatic TIME-on-chip model. Cancer Cell Int. 2023 Sep 27;23(1):219.

25. Yang S, He P, Wang J, Schetter A, Tang W, Funamizu N, et al. A Novel MIF Signaling

Pathway Drives the Malignant Character of Pancreatic Cancer by Targeting NR3C2. Cancer

Res. 2016 Jul 1;76(13):3838–50.

26. Bacher M, Metz CN, Calandra T, Mayer K, Chesney J, Lohoff M, et al. An essential

regulatory role for macrophage migration inhibitory factor in T-cell activation. Proc Natl Acad

Sci. 1996 Jul 23;93(15):7849–54.

27. Tsang JCH, Vong JSL, Ji L, Poon LCY, Jiang P, Lui KO, et al. Integrative single-cell and

cell-free plasma RNA transcriptomics elucidates placental cellular dynamics. Proc Natl Acad

Sci [Internet]. 2017 Sep 12 [cited 2024 May 31];114(37). Available from:

https://pnas.org/doi/full/10.1073/pnas.1710470114

28. Red-Horse K, Zhou Y, Genbacev O, Prakobphol A, Foulk R, McMaster M, et al. Trophoblast

differentiation during embryo implantation and formation of the maternal-fetal interface. J

Clin Invest. 2004 Sep 15;114(6):744–54.

29. Suryawanshi H, Morozov P, Straus A, Sahasrabudhe N, Max KEA, Garzia A, et al. A single-

cell survey of the human first-trimester placenta and decidua. Sci Adv. 2018 Oct

5;4(10):eaau4788.

30. Tang CT, Zhang QW, Wu S, Tang MY, Liang Q, Lin XL, et al. Thalidomide targets EGFL6 to

inhibit EGFL6/PAX6 axis-driven angiogenesis in small bowel vascular malformation. Cell

Mol Life Sci. 2020 Dec;77(24):5207–21.

31. Liang G, Zhou C, Jiang X, Zhang Y, Huang B, Gao S, et al. De novo generation of

macrophage from placenta-derived hemogenic endothelium. Dev Cell. 2021

Jul;56(14):2121-2133.e6.

32. Wang K, Zheng J. Signaling regulation of fetoplacental angiogenesis. J Endocrinol. 2012

Mar;212(3):243–55.

33. MacPhee DJ, Mostachfi H, Han R, Lye SJ, Post M, Caniggia I. Focal Adhesion Kinase Is a

Key Mediator of Human Trophoblast Development. Lab Invest. 2001 Nov;81(11):1469–83.

34. Palaiologou E, Etter O, Goggin P, Chatelet DS, Johnston DA, Lofthouse EM, et al. Human

placental villi contain stromal macrovesicles associated with networks of stellate cells. J

Anat. 2020 Jan;236(1):132–41.

35. Ueshima C, Kataoka TR, Osakabe M, Sugimoto A, Ushirokawa A, Shibata Y, et al.

Decidualization of Stromal Cells Promotes Involvement of Mast Cells in Successful Human

Pregnancy by Increasing Stem Cell Factor Expression. Front Immunol. 2022 Jan

31;13:779574.

36. Huang Q, Liu Y, Du Y, Garmire LX. Evaluation of Cell Type Annotation R Packages on

Single-Cell RNA-Seq Data. Genomics Proteomics Bioinformatics. 2021 Apr 1;19(2):267–81.

37. Zheng GXY, Terry JM, Belgrader P, Ryvkin P, Bent ZW, Wilson R, et al. Massively parallel

digital transcriptional profiling of single cells. Nat Commun. 2017 Jan 16;8(1):14049.

38. Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A, et al. Integrated analysis

of multimodal single-cell data. Cell. 2021 Jun;184(13):3573-3587.e29.

39. Li H. A statistical framework for SNP calling, mutation discovery, association mapping and

population genetical parameter estimation from sequencing data. Bioinformatics. 2011 Nov

1;27(21):2987–93.

40. The 1000 Genomes Project Consortium, Corresponding authors, Auton A, Abecasis GR,

Steering committee, Altshuler DM, et al. A global reference for human genetic variation.

Nature. 2015 Oct 1;526(7571):68–74.

41. Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal enrichment

tool for interpreting omics data. The Innovation. 2021 Aug;2(3):100141.

42. Kanehisa M, Goto S. KEGG: Kyoto Encyclopedia of Genes and Genomes.

43. Jin S, Guerrero-Juarez CF, Zhang L, Chang I, Ramos R, Kuan CH, et al. Inference and

analysis of cell-cell communication using CellChat. Nat Commun. 2021 Feb 17;12(1):1088.

Fig. 1. Illustration of Originator framework and the prevalence of blood immune cells in tissue samples.

a

1 Perform scRNA-seq preprocessing including QC, normalization, data integration, clustering, and cell-type annotation

2 Separate immune cells into tissue and blood residents using references

Separate barcoded cells into different origins using inferred genotypes

optional

Reference

Blood residents   Tissue residents

3 Dissect heterogeneity of the tissues in combinations of results from the previous two separation methods

4 Perform downstream analysis

b

Fig. 2. Applications of Originator to pancreatic ductal adenocarcinoma (PDAC) and placenta scRNA-seq data.