Enhanced behavioural and neural sensitivity to punishments in chronic pain and fatigue

Flavia Mancini *[1], **Pranav Mahajan**[2], **Anna á V. Guttesen**[1,2], **Jakub Onysk**[1,3], **Ingrid Scholtes**[4,5], **Nicholas Shenker**[4], **Michael Lee**[5], and **Ben Seymour**[2,6]

[1]**Computational and Biological Learning Unit, Department of Engineering, University of Cambridge, Cambridge CB2 1PZ, UK**
[2]**Wellcome Centre for Integrative Neuroimaging, FMRIB, Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford OX3 9DU, UK**
[3]**Applied Computational Psychiatry Lab, Max Planck Centre for Computational Psychiatry and Ageing Research, Queen Square Institute of Neurology and Mental Health Neuroscience Department, Division of Psychiatry, University College London, UK**
[4]**Cambridge University Hospital, Cambridge, UK**
[5]**Department of Medicine, University of Cambridge**
[6]**Institute of Biomedical Engineering, University of Oxford**

ABSTRACT

Chronic pain and fatigue in musculoskeletal disease contribute significantly to disability, and recent studies suggest an association with reduced motivation and excessive fear avoidance. In this behavioural neuroimaging study in chronic inflammatory arthritis participants and healthy controls, we aimed to identify the specific behavioral and neural changes associated pain and fatigue during reward and loss decision-making. Computational modeling of behaviour identified a parametric signature, characterized most notably by increased punishment sensitivity. This signature is distinct from patterns previously reported in psychiatric conditions and it aligns with predictions of mechanistic models of chronic pain such as the fear avoidance model. Neural activity associated with the punishment prediction error was enhanced in the right posterior insular cortex, putamen, pallidum, and dorsolateral prefrontal cortex. Functional network connectivity analysis showed that insula centrality correlated with subjective reports of fatigue and pain. Overall, the findings show that pain and fatigue in chronic pain relate to objective behavioural changes, and can be mapped to a specific pattern of activity in brain circuits of motivation and decision-making.

1 INTRODUCTION

Chronic pain and fatigue are the cardinal symptoms of musculoskeletal disease and lead to a level of disability that has significant socioeconomic cost (Woolf and Pfleger, 2003). Although multiple factors may contribute to the risk of developing chronic pain, a key hypothesis is that reduced central motivational drive to engage in physical activity leads to physical deconditioning that in turn exacerbates pain, a corollary of the fact that physiotherapy is clearly beneficial in most subacute and chronic musculoskeletal conditions (Meulders, 2019; Vlaeyen et al., 2016). This 'fear avoidance' hypothesis links with other neurobiological and cognitive factors known to contribute to risk for pain chronification, embedded within the broader biopsychosocial model that captures the complexity of chronic pain in everyday clinical contexts (Gatchel et al., 2007). The mechanism by which pain reduces central motivation therefore represents a critical link in the pathway to chronic pain and fatigue development (Seymour et al., 2023), but its nature and neurobiology are not known.

Injuries are common across animal species, for example after accidents and within/cross species contests, and lead to recognisable protective and recuperative behaviours (Walters and Williams, 2019). These are

---

*Corresponding author: flavia.mancini@eng.cam.ac.uk

presumed to be evolutionarily adaptive because the injured state makes animals more vulnerable to further harm, and so being extra careful whilst recovering benefits survival (Walters, 1994). Peripheral and spinal sensitization to pain provide one route to hyper-protective behaviour, and excessive pain hypersensitivity is a key focus of many current models of chronic pain. But recent ecological studies indicate a broader hyper-sensitivity to threat that goes beyond pain, suggesting a central, supramodal enhanced sensitivity to punishments as a mechanism of behavioural homeostasis after an injury, and a potential contributing factor to chronic pain and disability (Lister et al., 2020). Whilst enhanced punishment sensitivity would be adaptive for simple musculoskeletal injuries encountered in ecological contexts, it would ultimately be maladaptive in the context of clinical musculoskeletal conditions where inactivity is detrimental (Seymour et al., 2023).

Computational models of the neural circuits of learning and motivation allow us to dissect its core information processing steps, thus helping identify the underlying components of post-injury homeostasis such as modulation of punishment sensitivity (Seymour and Mancini, 2020). A similar approach has been applied to human behavioural experiments to identify, for example, reduced reward sensitivity in major depressive disorder, and aversive generalisation in anxiety disorder (Aylward et al., 2019; Huys et al., 2013; Dunsmoor and Paz, 2015). In this study, we aimed to identify i) whether there is a specific behavioural signature of pain and fatigue manifest during reward and loss decision making, and ii) if there is, does it relate to specific, localisable activity in the brain, in particular in circuits associated with motivation and learning.

Reward and loss decision-making captures a fundamental process that underlies much of our everyday behaviour - doing things that benefit us, and avoiding things that don't. We studied this using a '4-armed bandit' task (figure 1; (Daw et al., 2006; Seymour et al., 2016; Aylward et al., 2019)). Computational modelling of behavioural data can then be used to interrogate brain activity to find corresponding neural processing steps, and their relation to pain and associated fatigue. In the human brain, the mesolimbic and insula regions are thought to play a key role in pain-related motivation and avoidance learning, alongside a broader role in behavioural homeostasis (e.g. in illness and sickness states (Uddin et al., 2017)). We reasoned that the insula in particular might link the physiological identification of injury with enhanced responsiveness to punishments, which ultimate contribute to the pain and fatigue phenotype.

Figure 1. Task design. At the beginning of each trial, four cues were presented and participants chose one of them. Each cue was associated with an independent probability of winning and losing a token. After each choice, the outcome was displayed (green coin = win £1).

2 RESULTS

2.1 Choice behaviour

We recruited 60 volunteers to a behavioural neuroimaging experiment: 30 patients with painful inflammatory arthritis, and 30 age and sex matched controls (see Methods). In the 4-armed bandit task, participants were presented with four cues, and they need to choose one of them on each trial. Each cue was associated with an independent, non-stationary probability of winning or losing a coin worth £1. The mean reward and punishment rate were independent of each other, and noisy - slowly changing across trials (as a Gaussian random walk). After selecting one of the cues, their choice yielded one of four possible outcomes: one green token (win £1), one red token (lose £1), one green and one red token (win £1 but lose £1, so effectively £0 when added at the end of the game), and no outcome. This nature of the task means that participants need to constantly learn about the outcomes, with each option being fully independent of others, and importantly to learn independently about rewards and losses (i.e. so we can dissociate differences in reward and loss learning).

Basic performance measures (net gains, reaction times, frequency of choice switches) were comparable between patients and controls (see Supplementary Behavioural Results). This similar level of net gains is as expected: as the task is effectively a dynamic learning environment for probabilistic outcomes, different levels of performance do not yield big differences in outcomes, which protects against the potentially confounding effects of accruing different winnings. In contrast, any differences between groups is embedded in the trial-by-trial dynamics as a result of exploration and learning.

To capture these learning effects we used a hierarchical Bayesian inference approach to fit multiple variants of Reinforcement Learning (RL) models to the choice data, first to all pooled participants and then separately for the two groups (see Methods). These are biological informed models, arising from basic animal learning theory, that are known to capture behaviour and brain activity during this general class of task (Seymour and Mancini, 2020). The hierarchical Bayesian approach allows for individual differences while aggregating information across participants of the same group and make meaningful group comparisons (Ahn et al., 2017). The choice data of the pooled participants and of the patient group (Table 1) were best fitted by two models (1 and 3) that included two sensitivity parameters for rewards and punishments, which were used to scale the prediction error, a lapse and a decay parameter capturing forgetting in choice behaviour. In absolute terms, model 1 fitted best the data, but it was not significantly different than model 3. The only difference between models 1 and 3 lies in their learning rates: model 1 has two learning rates ($alpha$ and $beta$) that were added for updating reward values (reward LR $= \alpha + \beta$) and subtracted for updating punishment values (punishment LR $= \alpha - \beta$), whereas model 3 has separate, fixed learning rates for rewards and punishments. In the control group, model 3 showed the best absolute fit, but it was not significantly different than the other models we tested.

We compared the group estimates of the parameters of model 1 between the arthritis patient group and the control group, to investigate any core underlying differences in choice behaviour.

The group-level estimates of each free parameter of model 1 are reported in Table 2, and the individual participant's estimates are shown in Figure 2. In keeping with our core hypothesis, participants with chronic arthritis were more sensitive to punishments when compared to controls. In addition, we also found that they had faster memory decay and greater lapses in the choice behaviour. Finally, the differential component of the learning rate (β) was also higher in patients.

2.2 Neural correlates of punishment PE, PPE

Participants performed the bandit task whilst undergoing functional brain MRI. We aimed to identify brain changes associated with the behaviourally manifest difference in punishment sensitivity. Since punishment sensitivity scales the magnitude of the value of the punishment as used for learning avoidance actions, this is reflected in brain responses correlated with the punishment prediction error (PPE), with the individually estimated punishment sensitivity as a covariate. Thus, we evaluated whether the neural activity associated with the PPE, and modulated by punishment sensitivity, differed across groups. The arthritis group showed increased activity in the right posterior insular cortex, putamen, pallidum, and dorsolateral prefrontal cortex (figure 3; table 3. There was no significant cluster of decreased activity, at the cluster threshold used (z>2.6,

Table 1. Model comparison for the 4-armed bandit task. Five models were fitted to all participants, as well as the arthritis and control groups separately: (1) differential learning rates (LR), lapse and decay parameters; (2) differential LR with lapse parameter, no decay; (3) separate LRs for rewards and punishments, lapse and decay parameters; (4) separate LRs for rewards and punishments, lapses but no decay; (5) separate LRs for rewards and punishments, without lapse or decay parameters. We report the expected log predictive density (ELPD) difference between the best performing model (lowest LOOIC (leave-one-out cross-validation information criterion)) and each model are displayed, alongside the standard error (SE) of the difference and the sigma effect (the ratio between the ELPD and SE difference, which is a proxy for significance).

Group	Model	ELPD difference	SE difference	Sigma effect	LOOIC
All	1	0	0		24386.3539
	3	-53.422223	37.8139003	-1.4127668	24493.1983
	2	-276.00621	90.2032862	-3.0598243	24938.3663
	4	-280.33323	91.9252409	-3.0495785	24947.0203
	5	-336.24668	97.0027097	-3.4663638	25058.8472
Arthritis	1	0	0		12754.9737
	3	-46.954621	37.2017901	-1.2621603	12848.8829
	4	-221.65393	80.4995507	-2.7534803	13198.2815
	2	-226.78714	77.0709632	-2.9425757	13208.5479
	5	-264.26755	83.3235728	-3.171582	13283.5088
Controls	3	0	0		11667.014
	1	-2.6956325	9.89283891	-0.2724832	11672.4052
	4	-46.800276	48.2061096	-0.970837	11760.6145
	2	-48.534824	49.9225293	-0.9722028	11764.0836
	5	-56.156688	49.6040047	-1.1320999	11779.3273

Table 2. Group-level parameter estimates for the arthritis and control groups (mean and SD) in model 3 (separately fitted for each group) and HDI group differences (lower and upper 95% HDI). Significant differences do not include 0 and are highlighted with *.

Parameter	Arthritis	Controls	Lower 95% HDI	Upper 95% HDI
reward sensitivity	12.750 (5.911)	9.417 (5.031)	-1.004	7.695
punishment sensitivity	9.953 (2.577)	5.778 (3.715)	1.388*	8.016*
α	0.585 (0.300)	0.569 (0.178)	-0.133	0.183
β	0.106 (0.100)	0.016 (0.018)	0.0130*	0.101*
reward learning rate ($\alpha+\beta$)	0.679 (0.069)	0.590 (0.043)	-0.067	0.25
punishment learning rate ($\alpha-\beta$)	0.547 (0.074)	0.570 (0.047)	-0.199	0.142
lapse ξ	0.105 (0.043)	0.026 (0.003)	0.036*	0.105*
decay	0.603 (0.311)	0.469 (0.261)	0.008*	0.403*

Figure 2. Participant's estimates of each parameter. For each parameter (a.u.), we show the raincloud plots which include the (half) probability distribution function, scatterplots and boxplots of the data of arthritis patients (red) and controls (green).

Figure 3. Increased neural processing of the Punishment Prediction Error in the arthritis group vs. controls, modulated by punishment sensitivity. Increased BOLD responses associated with PPE in the right posterior insula, putamen, pallidum and the dorsolateral prefrontal cortex. The colorscale represents z scores ranging from 0 to 5, thresholded at z>2.6, p<0.05. The statistical contrast was overlaid over a structural brain image, obtained by averaging the high-resolution structural images of the study participants.

p<0.05). For completeness, common activity across the two groups is reported in Supplementary Figure S3 and Supplementary Table S6.

Cluster Id	Voxels	P	Z peak	X	Y	Z
2a	808	0.0015	4.28	36	-6	-6
2b			3.78	28	-4	-4
2c			3.72	16	4	2
2d			3.51	52	10	6
2e			3.47	24	12	-8
2f			3.23	50	-8	10
1a	468	0.0282	4.1	28	46	32
1b			3.51	14	18	34
1c			3.49	16	22	30
1d			3.39	16	32	32

Table 3. Activation clusters of increased activity in arthritis participants vs controls, associated with PPE and modulated by the punishment sensitivity parameter. Coordinates in MNI space.

2.3 Neural correlates of reward PE, RPE

Rewards prediction error (RPE) activity modulated by reward sensitivity was decreased in arthritis participants vs. controls in the a rostral region of the paracingulate gyrus and neighbouring white matter, and increased in the left occipital fusiform gyrus, area V4 (table 4). There was no significant cluster of increased activity, at the cluster threshold used (z>2.6, p<0.05). Common activity across groups is reported in Supplementary Figure S4 and Table S7.

2.4 Connectivity analyses

The correlation of punishment sensitivity with evoked BOLD responses in the insula is in keeping with the core hypothesis that the insula is a central part of a behavioural homeostatic network that mediates motivational changes in the pain state. To look at this further, we aimed to identify whether the insula connectivity during the task correlates with pain behaviour and task parameters. To do this, we calculated the network centrality (here, degree centrality), which captures the importance or influence of the insula as a network hub during the task. We parcelled the brain into 180 symmetrical ROIs using a functional (Glasser) brain atlas, and constructed a binarised connectivity network using thresholded pairwise Pearson correlations (see Methods). This analysis ignores the task events, and simply considers overall connectivity during the task as a whole. We found that degree centrality in posterior insula subregions correlated with subjective reports of daily average pain and fatigue scores (see figure 5A). There was evidence for a

Figure 4. Increased (red, A) and decreased (blue, B) activity associated with the reward prediction error in the arthritis group vs. controls. Colorscale represents z scores ranging from 0 to 5, thresholded at z>2.6, p<0.05. The statistical contrast was overlaid over a structural brain image, obtained by averaging the high-resolution structural images of the study participants.

Cluster Id	Voxels	P	Z peak	X	Y	Z
Arthritis > Controls						
1a	662	1.31e-06	3.76	-24	-82	-10
1b			3.75	-16	-82	-12
1c			3.21	-14	-88	-6
1d			3.11	-28	-74	-8
1e			2.89	-20	-64	-8
1f			2.81	-34	-78	-12
Arthritis < Controls						
1a	218	1.65	3.59	16	44	22
1b			2.72	8	50	14
1c			2.48	6	52	20

Table 4. Activation clusters of increased and decreased activity in arthritis patients vs controls, associated with RPE and modulated by the reward sensitivity parameter. Coordinates in MNI space.

correlation between Posterior Insular Area 2 and measures of pain (average daily pain rating from weekly diaries: Pearson's r= 0.566, LogBF10= 3.413; BPI severity score: Pearson's r= 0.544, LogBF10= 2.958), as well as fatigue (average daily fatigue rating from weekly diaries: Pearson's r= 0.517, LogBF10= 2.468). These correlations were less apparent when performing the same analysis using the resting state fMRI data, which has also been observed in other connectivity analyses (Zhao et al., 2023) (figure 5B).

This provides evidence that posterior insula activity not only correlates with pain related changes in objective task behaviour, but that it sits within a network of regions whose activity correlates with clinical subjective ratings of fatigue and pain.

Figure 5. (A) Bayesian Pearson correlation of degree centrality of 10 insular nodes with clinical outcomes and model parameters for the patient group in task fMRI. (B) Bayesian Pearson correlation of degree centrality of 10 insular nodes with clinical outcomes and model parameters for the patient group in resting-state fMRI Significant correlations: Area Post Insular 1 with ESR (Pearson's r= 0.515, LogBF10= 2.43) (C) A colour-coded map of insular nodes derived from the Glasser atlas (number denotes the parcel number in the Glasser atlas). Results for other network centralities are provided in Fig.S5.

3 DISCUSSION

The results reveal specific computational signature of chronic pain and fatigue, reflected in the pattern of changes of the underlying parameters that govern motivated behaviour. At the heart of this signature is an increase in punishment sensitivity, which demonstrates a generalised form of hyper-protective behaviour that enhances learning of aversive outcomes and may lead to excessive avoidance. This allows the organism to enhance defence beyond transient pain, which can mediated by central and peripheral hypersensitivity, to other aversive outcomes.

The signature also includes other components of behaviour, including the learning rate differential between rewards and punishments, alongside increased forgetting and choice lapses. This reveals a specific pattern that is distinct from that seen in other clinical conditions, using the same or similar task. For instance, depression is associated to the opposite tendency, a reduced sensitivity to rewards (Huys et al., 2013)). Anxiety is associated with increased punishment learning rates (Aylward et al., 2019), opposite to relative increase in reward learning rate seen here, suggesting more dynamic updating of rewards compared punishments. This is equivalent to a more persistent punishment memory, as has been demonstrated in previous studies of fibromyalgia - a non-inflammatory pain condition with dominant pain and fatigue (Meulders et al., 2017). The increased sensitivity to punishments in inflammatory arthritis is also different from the reduced general outcome sensitivity observed in Parkinson's disease as the result of deep brain stimulation (Seymour et al., 2016) and the reduced reward outcome value reported following tryptophan depletion (Seymour et al., 2012). An increase in the lapse and decay rates was also found in Parkinson's

patients, as in our study. This adds randomness to the decisions and probably reflects a non-specific, disruptive effect of pain and fatigue on attention. Together, this illustrates how the chronic pain and fatigue state seen in inflammatory arthritis is clearly computationally distinct from depression, anxiety and Parkison's disease.

The behavioural manifestation of computational components of learning and decision making maps well to the predictions of chronic pain models. Notably, increased punishment sensitivity is a central prediction of the the fear avoidance model (Meulders, 2019; Vlaeyen et al., 2016). Previous investigations have focused on punishment generalisation (Meulders et al., 2015), which is a distinct component of avoidance in which aversive values spread to perceptually similar pain predictors. Importantly, both increased punishment sensitivity and aversive generalisation drive the same tendency towards enhanced avoidance. This is a critical component of mechanistic models of chronic pain, which posit that the brain forms a hierarchical internal representations of the injury to drive protective and recuperative behaviours (Seymour and Mancini, 2020). Although greater avoidance reduces the chance of worsening damage through new actions, it inherently affects the opportunity to learn if an injury has resolved (information restriction), and will tend to perpetuate internal representations of injury states (Seymour et al., 2023). Future longitudinal studies would need to determine whether punishment sensitivity is a manifestation of the pain state or an outcome predictor.

At a cortical level, neural activity associated with punishment prediction error (PPE) was enhanced in the right posterior insular cortex, putamen, pallidum, and dorsolateral prefrontal cortex (dlPFC, figure 3), in the inflammatory arthritis group, according to individual differences in punishment sensitivity. The PPE reflects the change of punishment value at each timepoint that is used for learning, which is then scaled by the punishment sensitivity constant. In previous instrumental learning studies on healthy participants, PPE activity was reported in the anterior insula (Gueguen et al., 2021), putamen (Seymour et al., 2007), dlPFC (Gueguen et al., 2021). The role of the insula is in keeping with the hypothesis that it plays a fundamental role in pain (Horing and Büchel, 2022; Isnard et al., 2011; Moayedi, 2014; Segerdahl et al., 2015) and fatigue states (Stefanov et al., 2020; Wylie et al., 2020). This can also be related to a proposed role in active inference of interoceptive (Barrett and Simmons, 2015; Fermin et al., 2022), somatosensory (Allen et al., 2016) and nociceptive signals (Fazeli and Büchel, 2018; Geuter et al., 2017; Horing and Büchel, 2022).

The insula is well placed to interface afferent nociceptive information with efferent control through connectivity to mesolimbic and mesocortical pathways, other cortical sites involved in learning and motivation (such as the anterior cingulate and medial prefrontal cortex) and descending pain control systems. In our connectivity analysis, we computed the network centrality to capture the importance of different insula subregions as a network hub, and this was evidenced by the the correlatation between posterior insula area 2 activity and subjective reports of daily fatigue and pain scores. We also note that this network metric was much more apparent during task performance than at rest. Although this finding needs to be cross-validated in new datasets, it might suggest that looking at brain connectivity in the context of motivated behaviour may better reveal distinguishing functional connections than pure resting state brain, when used for biomarker generation. We should also note that we restricted our analysis to the insula, based on our a priori hypothesis, but it may well be that other regions (such as PFC and ACC) show symptom-correlated network activity.

Clinically, pain and fatigue are often highly correlated in inflammatory chronic pain conditions. However, the methods used in this study could be applied to painless, fatigue-dominant disorders, such as chronic fatigue syndrome and long COVID, to dissect the effect of fatigue from that of pain on instrumental learning. In inflammatory arthritis, the two are nearly impossible to disentangle. The approach here provides a behaviourally and computationally informed approach to disentangling these in future studies across other pain and fatigue cohorts.

4 METHODS

4.1 Participants

We initially engaged approx. 15 local patients with chronic pain to help us design the study and iterated preliminary versions of the protocol using a systems engineering approach. The study was approved by the central NHS Health Research Authority ethics committee (REC: 17/SW/0113, IRAS: 216259). After the design was finalised, we screened 80 volunteers and recruited 30 patients with chronic rheumatoid

and psoriatic arthritis and 30 healthy age-matched controls. Patients were recruited at Rheumatology, Cambridge University Hospitals. All participants gave informed written consent according to the Helsinki Guidelines. Three participants did not complete the full study and were therefore excluded, leaving us with 57 participants in total (28 controls and 29 patients; 45 females). Patients were 53.83 ± 8.09 years old (range 36-66 years; 27 females), and controls were 51.50 ± 10.77 years old (range 33-69 years; 18 females).

Participants were screened with a non-standardized demographic questionnaire and the following standardized questionnaires: Beck Depression Inventory II, Brief Pain Inventory, McGill Pain Assessment, Functional Assessment of Chronic Illness Therapy Fatigue Scale, Short form PROMIS Sleep Scale. Trained staff carried out arthritis assessments with standardized joint assessment disease activity score (DAS), rheumatoid classification (ACR/EULAR criteria) and psoriatic classification. Patients with disease activity (DAS >2.6) or subjectively high fatigue scores (>4 on scale 0 = 'no fatigue' to 10 = 'worst fatigue imaginable', FACIT <35) were included in the study. Controls had no clinically significant disease activity (DAS <2.6) or subjective fatigue (<4 on scale 0 = no fatigue to 10 = worst fatigue imaginable, FACIT >30). Participants with other explanations for fatigue (anaemia, unstable thyroid, heart conditions, diabetes, recent cancer treatment, recent surgery) and contra-indications for MRI safety (WBIC safety questionnaire) were considered not eligible to the study. The demographic and clinical details of the participants who completed the study are reported in Supplementary Table 1.

4.2 Procedure

Participants took part in an initial screening and training session, and in a further MRI session. The two sessions were separated by 5-73 days, depending on participant, site and scanner availability. Throughout both sessions and in between tasks/runs, participants were asked to rate their fatigue and pain on a scale of 0 to 10 (0 = no fatigue/pain, 10 = worst fatigue/pain imaginable).

4.3 4-armed bandit task

Participants underwent a 4-armed bandit decision-making task during MRI, following some practice. In each trial, participants could choose between 4 abstract boxes presented on the screen and try to win green tokens and simultaneously, but independently, try to avoid red tokens. The green and red signalled winning and loosing £1, respectively. The sum of positive minus negative tokens was exchanged for money at the end of the experiment. The probability of win/loss in each box changes gradually over trials as a gaussian random walk. The participants completed 240 trials over 4 runs.

4.4 Inflammatory markers

Blood samples were obtained for all participants in the second visit and the following inflammatory markers were evaluated: ESR, IFNγ, IL-10, IL-12p70, IL-13, IL-1β, IL-2, IL-4, IL-6, IL-8, TNFα. General health was assessed in full blood count and liver function samples.

4.5 Actigraphy and daily ratings

Participants recorded daily ratings and wore an ActiGraph GT9X Link (Actigraph, LLC) to the wrist, 24 hours a day, over the course of 5-7 days. The wearable device contains a piezo-electric sensor that generates a voltage when the device undergoes a change in acceleration (3 axial). It was set to record 60s epochs. Participants removed the wristbands only for showering or swimming. During this week, participants rated their fatigue and pain ("how much fatigue/pain do you feel right now on a scale of 0 to 10, 0 = no fatigue/pain, 10 = worst fatigue/pain imaginable). The participants set their alarms in the morning (9:00/ 9:30) and evening (19:30/20:00) and noted their current rating down on a questionnaire that was previously provided by the experimenters. Participants also recorded their daily consumption of caffeine and alcohol on a questionnaire. They noted how many units were consumed and the details of the product (e.g. filter coffee, paracetamol, wine etc). Analysis and results of actigraphy is reported in the Supplementary Methods and Results document.

4.6 Modelling of choice behaviour

We simulated and fitted five Reinforcement Learning (RL) models based on non-probabilistic delta rules, whereby the learning rates are fixed and driven by discrepancies between the estimate of the expected value

and observed values. In all models, we tracked the Q value of the four cues, as the sum of the Q values for rewards (Q^r) and punishments (Q^p). We calculated the reward and punishment prediction errors as the difference between the expected and observed value (Eq. 1, 2):

$$\delta^p{}_{t(i)} = O^p_{t(i)} - Q^p_{t(i)} \tag{1}$$

$$\delta^r{}_{t(i)} = O^r_{t(i)} - Q^r_{t(i)} \tag{2}$$

In all models, the prediction error was scaled by a free, sensitivity parameter (reward sensitivity R or punishment sensitivity P) (Eq. 3, 4), as in previous studies (Aylward et al., 2019):

$$\delta^p{}_{t(i)} = P \times (O^p_{t(i)} - Q^p_{t(i)}) \tag{3}$$

$$\delta^r{}_{t(i)} = R \times (O^r_{t(i)} - Q^r_{t(i)}) \tag{4}$$

The models differed also according to the update rule, and the presence of lapses and forgetting in choice behaviour. In all models the learning rate was constant across trials. In models 1 and 2, the learning rate for rewards was calculated as the sum of two free parameters, $\alpha + \beta$, and the learning rate for punishments was calculated as the difference of these free parameters, $\alpha - \beta$:

$$Q^r_{t(i)} = Q^r_{t(i)} + (\alpha + \beta) \times \delta^r_{t(i)} \tag{5}$$

$$Q^p_{t(i)} = Q^p_{t(i)} + (\alpha - \beta) \times \delta^p_{t(i)} \tag{6}$$

In models 3-5, the learning rates for rewards and punishments were estimated as two independent free parameters, i.e. α^r and α^p:

$$Q^r_{t(i)} = Q^r_{t(i)} + \alpha^r \times \delta^r_{t(i)} \tag{7}$$

$$Q^p_{t(i)} = Q^p_{t(i)} + \alpha^p \times \delta^p_{t(i)} \tag{8}$$

Models 1 and 3 additionally had a decay free parameter d to describe the rate of forgetting in the tracking of Q^r and Q^p (Eq. 9, 10), as in (Aylward et al., 2019):

$$Q^p_{t(i)} = (1-d) \times Q^p_{t-1(i)} \tag{9}$$

$$Q^r_{t(i)} = (1-d) \times Q^r_{t-1(i)} \tag{10}$$

All models used a softmax policy to determine choice probability $\mathbb{P}$ by combining the Q values of rewards and punishments:

$$\mathbb{P}_{t(i)} = \frac{exp(Q^r_{t(i)} + Q^p_{t(i)})}{\sum_j exp(Q^r_{t(j)} + Q^p_{t(j)})} \tag{11}$$

Model ID	Sensitivity	Reward LR	Punishment LR	Decay	Lapses
1	R, P	$\alpha + \beta$	$\alpha - \beta$	d	ξ
2	R, P	$\alpha + \beta$	$\alpha - \beta$	n/a	ξ
3	R, P	α^r	α^p	d	ξ
4	R, P	α^r	α^p	n/a	ξ
5	R, P	α^r	α^p	n/a	n/a

Table 5. Free parameters of the models that were fitted to the behavioural data.

Models 1-4 softmax policy included a free parameter ξ that characterised lapses in choice behaviour, similarly to (Aylward et al., 2019):

$$\mathbb{P}_{t(i)} = \frac{exp(Q^r_{t(i)} + Q^p_{t(i)})}{\sum_j exp(Q^r_{t(j)} + Q^p_{t(j)})} \times (1 - \xi) + \frac{\xi}{4} \tag{12}$$

²⁶¹ Table 5 summarises the free parameters of the different models we simulated and fitted to the data.

²⁶² We used hierarchical Bayesian methods (Hamiltonian Monte Carlo) to estimate model parameters, with
²⁶³ the HBayesDM and Stan package for R (Stan Development Team, 2019; Ahn et al., 2017) in R. For each
²⁶⁴ model, we fit 4 chains with 1000 burn in samples and 2000 samples to the following data: choice (1:4), win
²⁶⁵ (0,1) and loss (0,-1) for each trial.

²⁶⁶ Parameters for all models were fit separately for each group, as well as for all participants pooled in a
²⁶⁷ common group. The winning model was defined as the model with the lowest Leave-One-Out Information
²⁶⁸ Criterion (LOOIC). We also estimated the difference in models' expected predictive accuracy through the
²⁶⁹ difference in expected log point-wise predictive density (ELPD) (Vehtari et al., 2017). By taking the ratio
²⁷⁰ between the ELPD difference and the standard error (SE) of the difference, we obatin the sigma effect,
²⁷¹ which is a heuristic for significance of such model differences.

²⁷² We compared parameter estimates from the winning model across the two groups using 95 % high density
²⁷³ intervals (HDI). In particular, for each comparison, we calculated the difference in the hyper parameters
²⁷⁴ between groups and reported the 95 % HDI of the difference. In the Bayesian scenario, a significant group
²⁷⁵ difference is indicated by the interval not containing the value 0.

4.7 MRI data acquisition

²⁷⁷ First, we collected a T1-weighted MPRAGE structural scan (voxel size 1 mm isotropic) on a 3T Siemens
²⁷⁸ Magnetom Prisma scanner (Siemens Healthcare), equipped with a 64-channel head coil, at the Wolfson
²⁷⁹ Brain Imaging Centre, Cambridge. Then we collected a resting state scan of 240 volumes and 4 task fMRI
²⁸⁰ sessions of 264 volumes using the same multiband echo planar imaging (EPI) sequence (TR = 2000 ms,
²⁸¹ TE = 30 ms, flip angle = 82 deg, slices per volume = 64, multiband acceleration factor 2, interleaved
²⁸² slice mode, Grappa 2, voxel size 2 mm isotropic, A>P phase-encoding; this included four dummy volumes,
²⁸³ in addition to those pre-discarded by the scanner). In order to correct for inhomogeneities in the static
²⁸⁴ magnetic field, we imaged 4 volumes using an EPI sequence identical to that used in task fMRI, inverted in
²⁸⁵ the posterior-to-anterior phase encoding direction. Full sequence metadata are available XX.

4.8 MRI data analyses

4.8.1 Preprocessing

²⁸⁸ The anatomical images were reoriented to standard orientation, automatically cropped and bias-field
²⁸⁹ corrected using 'fsl_anat' (FMRIB's Software Library, www.fmrib.ox.ac.uk/fsl). The brain was extracted
²⁹⁰ with ANTs tool 'antsBrainExtraction' (https://github.com/ANTsX/ANTs). Functional images had the first
²⁹¹ four volumes removed, were motion corrected using FSL tool 'mcflirt', 3D time shifted using AFNI program
²⁹² '3dTshift' and a Fourier method (https://afni.nimh.nih.gov/). Fieldmap correction was applied using HySCO,
²⁹³ Hyperelastic Susceptibility Artefact Correction (Ruthotto et al., 2013). Non-brain data were removed using
²⁹⁴ FSL tool BET. Spatial smoothing was applied using a Gaussian kernel of FWHM 8.0mm. The entire 4D
²⁹⁵ dataset was grand-mean intensity normalised by a single multiplicative factor and highpass temporal filtering
²⁹⁶ was applied (Gaussian-weighted least-squares straight line fitting, with sigma=50.0s) using FSL.

4.8.2 Generalised Linear Modelling

First and second level GLM analyses were conducted using FEAT (FMRI Expert Analysis Tool) Version 6.00, part of FSL. All univariate imaging results were obtained from a single GLM model. We investigated neural correlates using model 1. All model predictors were generated with the group mean fitted parameters in order to minimise noise. First level regressors included onset times and values for the reward outcomes, punishment outcomes, reward prediction error, punishment prediction error (modelled at the time of the outcome) and regressors of no interest, such as the cue onset and keypress time. Sessions within subject were not concatenated, but modelled separately and then averaged using a mixed effect model (second-level analysis). Third level analyses were Two-Group Difference models, in which we compared the positive and negative means of reward or punishment prediction errors between the arthritis and control groups; as covariates, we included the sensitivity parameter and age, demeaned across all subjects. Finally, we applied a cluster threshold z>2.6, p<0.05 and we report only clusters that survived this multiple-comparison correction.

4.8.3 Connectivity analyses

Filtered functional images, preprocessed as per section 4.8.1 were registered to standard format (MNI template) and then parcellated into 180 regions of interest using the Glasser functional atlas (Glasser et al., 2016). Each region of interest combines the symmetric region in the brain, i.e. right and left regions jointly form one parcel. The timeseries for each parcel were extracted as the mean of the timeseries' corresponding to all voxels in that parcel. This process was performed for all four runs of task-fMRI and one run of rest-fMRI, and we performed elementwise summation of the four task-fMRI timeseries runs, making it comparable with the rest-fMRI.

Fully connected networks were constructed with edge strength between two nodes being the Pearson correlation between the timeseries corresponding to the nodes. We then thresholded the Pearson's full correlation matrices to a produce binary adjacency matrix (consisting of 1's and 0's) for each participant, as done in our previous studies(Mano et al., 2018). Each of the correlation matrices was thresholded in an adaptive manner to produce an adjacency matrix with a 10% link density. This value was chosen based on previous network topology studies that have found such a link density to provide optimal discriminative ability (Mano et al., 2018; Achard et al., 2012; Itahashi et al., 2014; Mansour et al., 2016; Termenon et al., 2016). We then computed the following nodal centrality measures for parcels corresponding to insular sub-regions: degree centrality, eigen-vector centrality, betweenness centrality, closeness centrality and load centrality.

The Glasser's atlas, being a functional atlas, has several parcels, some of which may overlap with the anatomical insular region. We found the Glasser parcels which correspond to the anatomical insula region by cross-checking the regions which intersect/overlap with the left and right insula regions in the AAL atlas, in FSLeyes. This resulted in 10 parcels of the Glasser atlas, reported in Figure 5C with their corresponding area descriptions from the Glasser atlas. Then, we performed Bayesian Pearson correlation of insular nodal centrality metrics with clinical scores and model parameters, only for the patient group. We did not include any healthy subjects because they have normal clinical scores, and no pain, and correlations would have risked being confounded by group differences.

We correlated the nodal centrality metrics of these insular subregions with several scores and model variables: Daily average pain, Daily average fatigue, Daily average sleep quality, DAS score, BDI score, GAD score, BPI severity, BPI interferance, McGill questionnaire score, Facit score, Promis (raw score), Immunological variables (ESR, IFNγ, IL-10, IL-12p70, IL-13, IL-1β, IL-2, IL-4, IL-6, IL-8, TNFα) and model parameters.

5 DATA AND CODE AVAILABILITY

All code and data will be available open source, released upon acceptance of the paper.

6 ACKOWLEDGEMENTS

The study was funded by a Versus Arthritis (21537) grant to B.S. and M.L., a MRC Career Development Award to FM (MR/T010614/1) and a UKRI Advanced Pain Discovery Platform grant to both F.M. and B.S. (MR/W027593/1). B.S. was also funded by Wellcome (214251/Z/18/Z) and IITP (MSIT 2019-0-01371). This work has been performed using resources provided by the Cambridge Tier-2 system operated by the University of Cambridge Research Computing Service (www.hpc.cam.ac.uk) funded by EPSRC Tier-2 capital grant (EP/T022159/1). HPC access was additionally funded by an EPSRC research infrastructure grant to F.M.. We are grateful to all the study participants and the Wolfson Brain Imaging Centre for their support. For the purpose of open access, the author has applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising from this submission.

7 AUTHOR CONTRIBUTIONS

BS, FM, ML and NS designed the study. AG and IS collected the data. AG, JO, FM, PM analysed the data. BS, FM and PM wrote the article.

8 DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

Achard, S., Delon-Martin, C., Vértes, P. E., Renard, F., Schenck, M., Schneider, F., Heinrich, C., Kremer, S., and Bullmore, E. T. (2012). Hubs of brain functional networks are radically reorganized in comatose patients. *Proceedings of the National Academy of Sciences*, 109(50):20608–20613.

Ahn, W.-Y., Haines, N., and Zhang, L. (2017). Revealing neurocomputational mechanisms of reinforcement learning and decision-making with the hBayesDM package. *Computational Psychiatry*, 1:24–57.

Allen, M., Fardo, F., Dietz, M. J., Hillebrandt, H., Friston, K. J., Rees, G., and Roepstorff, A. (2016). Anterior insula coordinates hierarchical processing of tactile mismatch responses. *Neuroimage*, 127:34–43.

Aylward, J., Valton, V., Ahn, W.-Y., Bond, R. L., Dayan, P., Roiser, J. P., and Robinson, O. J. (2019). Altered learning under uncertainty in unmedicated mood and anxiety disorders. *Nature human behaviour*, 3(10):1116–1123.

Barrett, L. F. and Simmons, W. K. (2015). Interoceptive predictions in the brain. *Nature reviews neuroscience*, 16(7):419–429.

Cole, R. J., Kripke, D. F., Gruen, W., Mullaney, D. J., and Gillin, J. C. (1992). Automatic sleep/wake identification from wrist activity. *Sleep*, 15(5):461–469.

Daw, N. D., O'doherty, J. P., Dayan, P., Seymour, B., and Dolan, R. J. (2006). Cortical substrates for exploratory decisions in humans. *Nature*, 441(7095):876–879.

Dunsmoor, J. E. and Paz, R. (2015). Fear generalization and anxiety: behavioral and neural mechanisms. *Biological psychiatry*, 78(5):336–343.

Fazeli, S. and Büchel, C. (2018). Pain-related expectation and prediction error signals in the anterior insula are not related to aversiveness. *Journal of Neuroscience*, 38(29):6461–6474.

Fermin, A. S., Friston, K., and Yamawaki, S. (2022). An insula hierarchical network architecture for active interoceptive inference. *Royal Society Open Science*, 9(6):220226.

Freedson, P. S., Melanson, E., and Sirard, J. (1998). Calibration of the computer science and applications, inc. accelerometer. *Medicine and science in sports and exercise*, 30(5):777–781.

Gatchel, R. J., Peng, Y. B., Peters, M. L., Fuchs, P. N., and Turk, D. C. (2007). The biopsychosocial approach to chronic pain: scientific advances and future directions. *Psychological bulletin*, 133(4):581.

Geuter, S., Boll, S., Eippert, F., and Büchel, C. (2017). Functional dissociation of stimulus intensity encoding and predictive coding of pain in the insula. *Elife*, 6:e24770.

Glasser, M. F., Coalson, T. S., Robinson, E. C., Hacker, C. D., Harwell, J., Yacoub, E., Ugurbil, K., Andersson, J., Beckmann, C. F., Jenkinson, M., et al. (2016). A multi-modal parcellation of human cerebral cortex. *Nature*, 536(7615):171–178.

Gueguen, M. C., Lopez-Persem, A., Billeke, P., Lachaux, J.-P., Rheims, S., Kahane, P., Minotti, L., David, O., Pessiglione, M., and Bastin, J. (2021). Anatomical dissociation of intracerebral signals for reward and punishment prediction errors in humans. *Nature communications*, 12(1):3344.

Horing, B. and Büchel, C. (2022). The human insula processes both modality-independent and pain-selective learning signals. *PLoS Biology*, 20(5):e3001540.

Huys, Q. J., Pizzagalli, D. A., Bogdan, R., and Dayan, P. (2013). Mapping anhedonia onto reinforcement learning: a behavioural meta-analysis. *Biology of mood & anxiety disorders*, 3(1):1–16.

Isnard, J., Magnin, M., Jung, J., Mauguière, F., and Garcia-Larrea, L. (2011). Does the insula tell our brain that we are in pain? *PAIN®*, 152(4):946–951.

Itahashi, T., Yamada, T., Watanabe, H., Nakamura, M., Jimbo, D., Shioda, S., Toriizuka, K., Kato, N., and Hashimoto, R. (2014). Altered network topologies and hub organization in adults with autism: a resting-state fmri study. *PloS one*, 9(4):e94115.

Lister, K. C., Bouchard, S. M., Markova, T., Aternali, A., Denecli, P., Pimentel, S. D., Majeed, M., Austin, J.-S., Williams, A. C. d. C., and Mogil, J. S. (2020). Chronic pain produces hypervigilance to predator odor in mice. *Current Biology*, 30(15):R866–R867.

Mano, H., Kotecha, G., Leibnitz, K., Matsubara, T., Sprenger, C., Nakae, A., Shenker, N., Shibata, M., Voon, V., Yoshida, W., et al. (2018). Classification and characterisation of brain network changes in chronic back pain: A multicenter study. *Wellcome open research*, 3.

Mansour, A., Baria, A. T., Tetreault, P., Vachon-Presseau, E., Chang, P.-C., Huang, L., Apkarian, A. V., and Baliki, M. N. (2016). Global disruption of degree rank order: a hallmark of chronic pain. *Scientific reports*, 6(1):1–17.

Meulders, A. (2019). From fear of movement-related pain and avoidance to chronic pain disability: a state-of-the-art review. *Current Opinion in Behavioral Sciences*, 26:130–136.

Meulders, A., Jans, A., and Vlaeyen, J. W. (2015). Differences in pain-related fear acquisition and generalization: an experimental study comparing patients with fibromyalgia and healthy controls. *Pain*, 156(1):108–122.

Meulders, A., Meulders, M., Stouten, I., De Bie, J., and Vlaeyen, J. W. (2017). Extinction of fear generalization: a comparison between fibromyalgia patients and healthy control participants. *The Journal of Pain*, 18(1):79–95.

Moayedi, M. (2014). All roads lead to the insula. *Pain*, 155(10):1920–1921.

Ruthotto, L., Mohammadi, S., Heck, C., Modersitzki, J., and Weiskopf, N. (2013). Hyperelastic susceptibility artifact correction of dti in spm. In *Bildverarbeitung für die Medizin 2013: Algorithmen-Systeme-Anwendungen. Proceedings des Workshops vom 3. bis 5. März 2013 in Heidelberg*, pages 344–349. Springer.

Segerdahl, A. R., Mezue, M., Okell, T. W., Farrar, J. T., and Tracey, I. (2015). The dorsal posterior insula subserves a fundamental role in human pain. *Nature neuroscience*, 18(4):499–500.

Seymour, B., Barbe, M., Dayan, P., Shiner, T., Dolan, R., and Fink, G. R. (2016). Deep brain stimulation of the subthalamic nucleus modulates sensitivity to decision outcome value in parkinson's disease. *Scientific Reports*, 6(1):1–12.

Seymour, B., Crook, R., and Chen, Z. (2023). Post-injury pain and behaviour: a control theory perspective. *Nature Reviews Neuroscience*.

Seymour, B., Daw, N., Dayan, P., Singer, T., and Dolan, R. (2007). Differential encoding of losses and gains in the human striatum. *Journal of Neuroscience*, 27(18):4826–4831.

Seymour, B., Daw, N. D., Roiser, J. P., Dayan, P., and Dolan, R. (2012). Serotonin selectively modulates reward value in human decision-making. *Journal of Neuroscience*, 32(17):5833–5842.

Seymour, B. and Mancini, F. (2020). Hierarchical models of pain: Inference, information-seeking, and adaptive control. *NeuroImage*, 222:117212.

Stan Development Team (2019). RStan: the R interface to Stan. R package version 2.21.0.

Stefanov, K., McLean, J., McColl, A., Basu, N., Cavanagh, J., and Krishnadas, R. (2020). Mild inflammation in healthy males induces fatigue mediated by changes in effective connectivity within the insula. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, 5(9):865–874.

Termenon, M., Achard, S., Jaillard, A., and Delon-Martin, C. (2016). The "hub disruption index," a reliable index sensitive to the brain networks reorganization. a study of the contralesional hemisphere in stroke.

Frontiers in computational neuroscience, page 84.

Uddin, L. Q., Nomi, J. S., Hébert-Seropian, B., Ghaziri, J., and Boucher, O. (2017). Structure and function of the human insula. *Journal of clinical neurophysiology: official publication of the American Electroencephalographic Society*, 34(4):300.

Vehtari, A., Gelman, A., and Gabry, J. (2017). Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Statistics and Computing*, 27(5):1413–1432. arXiv:1507.04544 [stat].

Vlaeyen, J. W., Crombez, G., and Linton, S. J. (2016). The fear-avoidance model of pain. *Pain*, 157(8):1588–1589.

Walters, E. T. (1994). Injury-related behavior and neuronal plasticity: an evolutionary perspective on sensitization, hyperalgesia, and analgesia. *International review of neurobiology*, 36:325–427.

Walters, E. T. and Williams, A. C. d. C. (2019). Evolution of mechanisms and behaviour important for pain.

White, C. N., Servant, M., and Logan, G. D. (2018). Testing the validity of conflict drift-diffusion models for use in estimating cognitive processes: A parameter-recovery study. *Psychonomic Bulletin & Review*, 25(1):286–301.

Wilson, R. C. and Collins, A. G. (2019). Ten simple rules for the computational modeling of behavioral data. *eLife*, 8:e49547.

Woolf, A. D. and Pfleger, B. (2003). Burden of major musculoskeletal conditions. *Bulletin of the world health organization*, 81(9):646–656.

Wylie, G., Yao, B., Genova, H., Chen, M., and DeLuca, J. (2020). Using functional connectivity changes associated with cognitive fatigue to delineate a fatigue network. *Scientific reports*, 10(1):21927.

Zhao, W., Makowski, C., Hagler, D. J., Garavan, H. P, Thompson, W. K., Greene, D. J., Jernigan, T. L., and Dale, A. M. (2023). Task fmri paradigms may capture more behaviorally relevant information than resting-state functional connectivity. *NeuroImage*, 270:119946.

SUPPLEMENTARY METHODS AND RESULTS

CLINICAL AND DEMOGRAPHIC DATA

Clinical and demograhic data of theb 57 partcipants who completed the full study are reported in Table S1. As expected, patients scored higher than controls on the Brief Pain Inventory (severity score: $t_{55}=11.188$, $p<0.001$, Cohen's d=1.860, BF10=$4.495\times10^{(+12)}$; interference score: $t_{55}=8.074$, $p<0.001$, BF10=$9.935\times10^{(+7)}$), McGill Pain Questionnaire ($t_{55}=7.807$, $p<0.001$, Cohen's d=2.068, BF10=$3.873\times10^{(+7)}$). Patients reported more fatigue (FACIT scale: $t_{55}=-9.055$, $p<0.001$, Cohen's d=-2.399, BF10=3.132e+9) and poorer sleep quality than controls (PROMIS sleep scale: $t_{55}=4.784$, $p<0.001$, Cohen's d=1.268, BF10=1302.373). This is paralleled by higher mean ratings of pain ($t_{55}=12.299$, $p<0.001$, Cohen's d=3.259, BF10=1.650e+14) and fatigue ($t_{55}=8.672$, $p<0.001$, Cohen's d=2.298, BF10=8.177e+8), and lower ratings of sleep quality ($t_{55}=2.948$, $p=0.005$, Cohen's d=0.788, BF10=8.667), collected using diaries over a continuous period of 5 days, relative to controls. Average levels of pain and fatigue were highly correlated across participants (Pearson's r=0.84, BF10=$2.210\times10^{1}3$). Interestingly, patients adopted higher severity ratings to describe their fatigue levels relative to their pain levels (mean fatigue diary rating ± SD, 5.461 ± 1.765; mean pain diary rating ± SD, 4.804 ± 1.698; $t_{28}=2.780$, $p=0.010$, Cohen's d=0.516, BF10=4.710). Unsurprisingly, patients consumed more caffeine per day ($t_{55}=3.444$, $p=0.001$, Cohen's d=0.913, BF10=28.578), but comparable units of alcohol than controls ($t_{55}=0.221$, $p=0.826$, Cohen's d=0.059, BF10=0.274). Finally, patients reported greater depression symptoms at the Beck Depression Inventory (mean score ± SD, 8.429±5.295; group difference, $t_{55}=6.961$, $p<0.001$, Cohen's d=1.860, BF10=1.798e+6) and generalised anxiety symptoms at the GAD7 test (mean score ± SD, 7.857±4.897; group difference, $t_{55}=6.560$, $p<0.001$, Cohen's d=1.753, BF10=448796.707). Overall, we found strong evidence for an associations between pain and fatigue questionnaires (BPI, McGIll and FACIT) and self-reported measures of depression and generalised anxiety (all correlations p < 0.001).

Table S1. Demographic and clinical data of the participants who completed the study.

ID	Group	Gender	Age	Diagnosis	DAS/DAPSA score	BPI severity	BPI interference	McGill	Facit	Diary avg pain	Diary avg fatigue	BDI score	GAD score	Promis (raw score)	Diary avg sleep	CRP	ESR	RheumatoidFactor	IFNγ	IL-10	IL-12p70	IL-13	IL-1β	IL-2	IL-4	IL-6	IL-8	TNFα
P08	Control	Male	60	n/a	0.55	0.25	0.00	0	49	0.43	2.05	0	0	14	3.73	<4	2	<10	3.33	0.21	0.33	0.91	0.13	0.46	<0.1	0.81	4.87	1.65
P10	Control	Female	66	n/a	2.26	0	0.00	0	52	0.45	1.91	3	1	10	0.67	<4	16	<10	1.88	0.37	0.87	1.94	0.10	0.80	<0.1	0.64	3.04	0.66
P12	Control	Female	43	n/a	1.46	2.75	0.29	8	28	1.33	3.93	3	1	25	7.29	<4	6	10	4.23	0.22	<0.2	<0.8	0.10	0.52	<0.1	0.32	9.12	2.46
P13	Control	Male	55	n/a	1.97	0.25	0.43	0	46	0.64	0.64	1	4	11	1.29	<4	11	<10	18.08	0.20	0.41	<0.8	<0.1	0.65	<0.1	0.65	3.76	1.86
P14	Control	Male	65	n/a	2.02	0	0.00	0	43	0.00	0.67	1	4	15	1.43	<4	18	<10	2.96	0.08	0.56	<0.8	<0.1	1.70	<0.1	0.45	6.19	2.24
P15	Control	Female	62	n/a	1.27	0	0.00	0	47	0.00	3.54	0	4	23	6.71	<4	6	<10	13.94	0.57	<0.2	1.15	0.38	0.53	<0.1	0.51	9.23	4.37
P20	Control	Female	50	n/a	0.49	0.75	0.29	4	47	2.26	2.36	0	1	28	5.14	<4	4	10	7.70	0.09	0.47	1.03	<0.1	<0.2	<0.1	0.79	6.27	1.38
P23	Control	Male	51	n/a	1.53	0	0.29	0	43	0.71	1.14	2	2	21	4.86	<4	2	11	6.29	0.32	0.53	1.03	0.16	0.43	<0.1	0.55	4.36	31.99
P32	Control	Male	50	n/a	1.36	0	0.00	4	52	0.00	2.07	0	0	12	1.43	<4	8	<10	3.04	0.25	<0.2	<0.8	<0.1	2.57	<0.1	0.96	3.61	1.75
P37	Control	Female	50	n/a	1.46	1	0.00	0	52	0.00	1.23	5	2	16	3.29	<4	7	<10	4.15	0.20	0.47	0.85	<0.1	0.41	<0.1	1.11	5.77	2.84
P43	Control	Female	33	n/a	1.25	1	0.29	0	49	1.20	1.73	1	0	23	3.13	<4	4	11	7.18	0.52	0.20	1.28	0.50	0.63	<0.1	0.63	8.48	2.87
P46	Control	Female	42	n/a	1.39	1	0.00	0	50	0.00	2.13	0	3	25	3.50	<4	6	11	2.56	0.25	0.20	<0.8	0.11	0.24	<0.1	0.63	3.45	1.23
P47	Control	Female	67	n/a	1.2	1	0.43	4	46	0.64	1.64	4	3	14	3.33	n/a	5	11	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
P49	Control	Female	38	n/a	1.54	1	0.00	0	52	0.17	1.92	0	0	14	7.29	<4	9	<10	14.92	0.55	0.48	1.68	0.47	0.32	<0.1	2.43	8.16	23.14
P50	Control	Male	69	n/a	n/a	1	0.00	2	45	0.00	3.21	0	5	14	1.00	n/a	n/a	n/a	2.38	0.29	0.40	0.98	0.15	0.44	<0.1	0.68	4.95	2.03
P55	Control	Female	68	n/a	2.33	1	0.00	0	48	1.53	2.80	5	6	16	3.29	n/a	n/a	n/a	4.04	0.23	0.61	0.81	<0.1	1.22	<0.1	1.22	4.52	2.50
P58	Control	Female	59	n/a	n/a	0	0.00	0	52	0.60	0.73	4	1	14	1.00	<4	4	<10	2.65	0.33	0.61	1.31	<0.1	0.34	<0.1	0.86	2.49	1.90
P59	Control	Male	65	n/a	1	0	0.00	0	50	0.18	1.45	1	1	16	2.67	<4	5	<10	10.30	0.18	0.33	<0.8	<0.1	0.33	<0.1	1.28	3.31	2.65
P60	Control	Female	64	n/a	1.13	0	0.00	0	49	0.07	0.29	1	1	16	4.29	<4	4	<10	12.56	0.37	0.46	1.97	0.16	0.65	<0.1	1.65	5.06	2.05
P68	Control	Female	58	n/a	0.96	0	0.00	0	42	0.00	1.25	0	0	16	6.50	<4	8	10	2.07	0.41	0.28	1.37	<0.1	0.34	<0.1	0.55	3.22	1.96
P71	Control	Male	42	n/a	1.46	0	0.00	4	50	0.00	0.80	2	0	17	2.00	<4	2	10	9.96	0.12	0.47	<0.8	<0.1	1.56	<0.1	0.48	6.82	2.05
P72	Control	Male	54	n/a	0.56	0	0.00	0	51	0.00	4.29	0	0	20	5.43	<4	5	<10	5.68	0.74	0.79	<0.8	0.18	0.45	<0.1	0.55	6.67	1.91
P74	Control	Male	44	n/a	1.27	1.5	0.00	0	48	0.00	1.57	0	0	12	2.71	<4	5	<10	2.77	0.24	0.32	2.30	<0.1	0.34	<0.1	1.36	4.00	1.79
P75	Control	Female	61	n/a	2.02	0	0.57	9	48	2.17	0.83	2	0	14	1.14	<4	12	<10	8.20	0.32	<0.2	<0.8	0.10	0.40	<0.1	1.32	4.93	1.71
P76	Control	Female	46	n/a	1.13	0	0.00	0	45	0.00	5.47	2	2	28	5.43	<4	5	<10	7.61	0.32	0.46	0.96	0.18	0.42	<0.1	2.77	4.11	2.02
P78	Control	Male	39	n/a	0.97	0	0.00	0	50	2.10	2.70	1	2	21	4.00	<4	4	<10	2.95	0.71	0.58	1.14	<0.1	0.55	<0.1	0.45	4.47	39.77
P79	Control	Female	48	n/a	1.36	0	0.00	0	47	0.00	1.00	0	4	23	4.86	<4	7	<10	16.39	0.28	0.22	1.06	0.16	0.46	<0.1	0.67	5.26	3.81
P80	Control	Female	57	n/a	n/a	0	0.00	0	52	0.00	0.00	0	0	22	2.43	n/a	n/a	n/a	4.78	0.33	2.23	<0.8	<0.1	<0.2	0.30	0.73	12.59	18.17
P01	Patient	Female	52	n/a	4.95	6.5	4.29	23	27	7.31	7.31	0	4	22	5.71	<4	28	36	31.74	0.33	3.62	6.36	1.67	0.85	0.30	8.65	3.92	4.51
P02	Patient	Female	56	RA	4.96	5.25	6.57	13	36	5.08	6.27	2	16	28	5.88	<4	11	36	3.17	0.27	0.74	<0.8	<0.1	0.56	<0.1	1.76	5.68	1.97
P03	Patient	Female	50	RA	3.5	3.25	2.29	9	38	4.00	3.73	7	8	33	2.00	<4	11	<10	2.25	0.29	0.36	0.95	<0.1	0.39	<0.1	1.10	7.03	1.69
P05	Patient	Male	60	RA	2.62	2.25	2.43	4	29	1.99	4.55	6	5	20	3.70	34	16	56	10.36	1.36	0.57	1.31	0.30	0.39	<0.1	12.08	5.17	3.52
P07	Patient	Female	36	RA	6.94	7	7.00	20	12	7.10	7.00	19	15	33	5.50	<4	19	101	3.20	0.28	0.53	<0.8	<0.1	0.46	<0.1	0.54	5.31	2.62
P16	Patient	Female	55	RA	2.66	1.25	1.14	2	33	2.00	2.93	2	6	19	2.43	<4	9	<10	3.20	0.12	0.29	<0.8	<0.1	0.37	<0.1	6.50	5.54	2.46
P17	Patient	Female	37	RA	1.63	1	0.00	2	49	2.33	2.93	5	9	22	3.14	<4	9	12	3.36	0.66	0.38	<0.8	<0.1	0.31	<0.1	0.37	5.51	1.90
P18	Patient	Female	59	RA	3.75	4.75	2.43	22	28	4.87	2.07	1	2	17	n/a	n/a	n/a	25	2.62	0.14	0.25	<0.8	<0.1	<0.2	<0.1	0.67	5.51	2.05
P19	Patient	Female	54	RA	3.28	3	2.00	13	40	3.00	2.53	10	5	22	5.43	30	12	80	6.66	0.27	<0.2	1.04	0.12	0.43	<0.1	3.07	3.18	2.01
P24	Patient	Female	56	RA	3.27	2.5	1.57	6	41	3.53	3.33	12	18	26	6.43	<4	11	44	9.46	0.21	0.96	1.48	0.26	0.27	<0.1	0.52	4.23	2.65
P27	Patient	Female	65	RA	4.74	1.5	0.43	4	29.25	2.40	4.33	5	6	25	4.63	<4	8	10	5.20	0.28	0.77	0.81	<0.1	0.60	<0.1	0.34	3.20	1.90
P28	Patient	Female	37	RA	4.19	4.75	5.57	11	36	4.73	6.80	12	15	25	6.29	<4	13	67	5.71	0.31	0.58	3.89	<0.1	0.99	<0.1	12.19	13.82	3.02
P29	Patient	Female	47	RA	4.4	2.5	1.29	3	36	3.42	4.68	6	14	14	3.86	8	11	17	2.73	0.26	0.58	<0.8	<0.1	0.39	<0.1	0.34	5.30	2.18
P33	Patient	Female	57	RA	4.26	3.5	1.57	17	7	3.60	5.80	13	5	28	5.67	8	23	252	2.73	0.13	0.21	0.81	<0.1	<0.2	<0.1	2.80	3.51	1.74
P34	Patient	Female	54	RA	6.75	6	4.29	21	16	7.80	7.92	22	14	17	1.60	<4	18	19	10.89	0.29	0.26	1.09	<0.1	0.38	<0.1	0.25	9.06	2.64
P35	Patient	Female	60	RA	6	4.25	6.57	17	22	6.09	4.07	9	9	22	5.20	16	18	19	6.11	0.29	0.39	<0.8	0.14	0.21	<0.1	1.30	3.28	1.99
P38	Patient	Female	57	RA	5.91	5	6.57	4	34	5.69	5.67	5	6	38	7.17	<4	8	143	4.61	0.29	0.77	<0.8	<0.1	0.37	<0.1	0.57	4.39	2.77
P40	Patient	Female	53	RA	7.18	5.75	4.57	10	33	3.85	7.00	14	6	23	2.33	17	41	73	10.22	0.73	<0.2	<0.8	<0.1	<0.2	<0.1	1.61	6.44	2.16
P42	Patient	Female	35	PsA	5.63	3.75	7.00	17	31	3.36	4.23	11	7	23	4.57	21	12	11	4.54	0.22	0.31	<0.8	0.11	0.27	<0.1	0.50	4.14	1.63
P44	Patient	Female	52	RA	3.51	6	7.00	20	21	4.77	4.77	10	16	38	7.00	21	18	<10	4.08	0.64	0.77	1.04	<0.1	0.29	<0.1	0.86	3.46	2.35
P54	Patient	Female	45	RA	5.46	6.75	8.43	26	21	6.47	7.77	13	13	32	5.00	22	35	368	2.94	0.33	2.02	<0.8	<0.1	0.99	<0.1	0.48	3.62	1.88
P56	Patient	Female	45	RA	5.66	6.75	5.00	15	27	4.94	6.38	4	0	32	4.50	19	21	<10	3.67	0.10	0.22	<0.8	<0.1	<0.2	<0.1	0.95	3.95	2.00
P57	Patient	Female	60	RA	4.76	3.25	4.43	21	32	5.60	8.30	8	10	39	9.20	<4	18	<10	2.28	0.97	0.26	<0.8	<0.1	0.86	<0.1	2.23	6.00	2.65
P61	Patient	Male	57	PsA	3.2	5.25	2.29	4	34	5.46	6.31	4	3	24	6.29	n/a	n/a	n/a	<1.7	0.70	0.39	1.09	<0.1	0.22	<0.1	0.97	2.68	1.35
P64	Patient	Female	62	RA	3.28	4.25	6.43	1	25	7.64	6.21	6	5	31	6.14	29	47	<10	4.40	0.24	<0.2	<0.8	0.14	0.37	<0.1	1.57	4.39	2.61
P65	Patient	Female	62	PsA	5	6.25	6.43	24	3	5.00	6.60	16	5	23	3.50	n/a	9	114	2.25	0.27	0.31	<0.8	0.11	0.29	<0.1	0.50	4.03	1.65
P66	Patient	Female	62	RA	4.24	6.25	6.43	19	32	7.50	8.00	16	5	36	6.67	<4	32	<10	4.97	0.22	0.62	0.93	0.23	0.93	<0.1	0.86	3.95	2.00
P69	Patient	Female	39	RA	6.93	3.75	8.43	24	25	7.50	8.00	16	6	36	6.67	<4	32	<10	4.97	0.22	0.77	1.52	0.11	0.56	<0.1	0.45	4.15	2.16
P73	Patient	Female	45	PsA	4.12	3	0.71	2	34	4.33	5.13	8	6	26	4.29	18	25	10	6.72	0.38	0.35	<0.8	0.14	0.59	<0.1	1.15	4.76	27.27

ADDITIONAL TASK RESULTS

Patients won a comparable amount of money (net £4.62±2.50) than controls (net £5.18±2.53) in the 4-armed bandit task (t55=0.837, p=0.406, Cohen's d=0.222, BF10=0.359). Reaction times were comparable between groups (t55=0.794, p=0.430, Cohen's d=0.210, BF10=0.349).

Patients and controls were similarly likely to switch choice, irrespective of the outcome (t55=0.528, p=0.6, Cohen's d=0.140, BF10=0.301). However, patients were overall slower at making a choice than controls (p=0.013) and we found a nominally-significant interaction between group and choice type (2 levels: repeated, switched; p=0.027); the choice to switch was slower in patients than controls (t55=2.478, p=0.016, Cohen's d=0.657, BF10=3.243), whereas the choice to stick to the previous decision had similar latency in the two groups (t55=1.484, p=0.144, Cohen's d=0.393, BF10=0.667).

Figure S1. Task performance. A. Net gains (GBP); B. Probability of switching choice irrespective of the outcome; C. Latency of the switched choice; D. Latency of the repeated choice in controls (green) and patients (red). For each variable, the probability density functions, scatterplot of individual data and boxplot are shown.

We evaluated whether the frequency of choice switches was affected by the outcome of the preceding trial (4 levels: reward only, punishment only, simultaneous reward and punishment, no outcome) and by the diagnosis (2 levels: inflammatory arthritis, healthy controls). A mixed-effects ANOVA provided evidence for an effect of condition (F1.45,79.75=110.824, p<0.001, eta2=0.637), but no evidence for main effect of group (F1,55=0.907, p=0.345, eta2=0.016) nor interaction (F1.45,79.75=0.517, p=0.540, eta2=0.003). Irrespective of their group, participants were more likely to change their choice following a trial that yielded either a loss or no outcome than in trials that yielded either a win or a simultaneous win and loss (all ps < 0.001, Cohen's d > 1.5). Moreover, the frequency of choice switches was similar after a loss and no outcome (p=0.062, Cohen's d=0.253); participants switched choices less frequently after affording a win rather than an ambiguous outcome (p=0.017, Cohen's d=0.360). Finally, the total frequency of choice switches were comparable across groups (t55=0.952, p=0.345, Cohen's d=0.252, BF10=0.391).

SLEEP ACTIGRAPHY

We used ActiLife v6.13.3 to extract sleep measures. Automatic detection of sleep periods was implemented with Tudor-Locke automatic sleep period detection. If there was a discrepancy of >2 hours between the automatic detected timing and the self-documented bedtime or out-of-bed time, the self-documented times would be entered manually as sleep periods. The Cole-Kripke algorithm was applied to the data to determine sleep parameters (Cole et al., 1992). Data from 5 participants (3 patients) with <5 nights of recording were excluded from further analyses, leaving 52 participants for analysis (25 patients).

Controls and patients showed comparable results in the sleep parameters, total sleep time (t(50)=1.597, p=.117, Cohen's d=0.443, BF10=0.788), movement index (t(50)=-1.684, p=0.098, Cohen's d=0.467, BF10=0.884), sleep fragmentation (t(50)=-1.648, p=0.106, Cohen's d=0.457, BF10=0.842), sleep latency (t(50)=-0.296, p=0.768, Cohen's d=0.082, BF10=0.289), sleep efficiency (t(50)=0.794, p=0.431, Cohen's d=0.220, BF10=0.361) and wake-after-sleep-onset (t(50)=-0.171, p = 0.865, Cohen's d=0.047, BF10=0.282).

DAILY ACTIVITY

We used ActiLife v6.13.3 to extract daily activity measures, implementing the default wear time validation and (Freedson et al., 1998) algorithm to detect activity bouts. Days with < 1080 epochs (i.e., 18 hours) of data were excluded from the analyses (154 days removed in total). Similar to the sleep analysis, any participants with < 5 days of recording were removed from further analysis (N=5, 2 patients), leaving 52 participants for analysis (26 patients).

Patients and controls showed comparable results in the activity parameters, with mean daily steps (mean ±SD, patients: 12114.340 ±4323.304, controls: 14191.425 ± 3714.191; t(50)= 1.858, p=.069, Cohen's d=0.515, BF10=1.131), proportion spent in sedentary activity (mean ± SD, patients: 0.536 ± 0.108, controls: 0.498 ±0.098; t(50)=-1.347, p=.184, Cohen's d=0.374, BF10=0.585), proportion spent in light activity (mean ± SD, patients: 0.355 ± 0.083, controls: 0.371 ± 0.072; t(50)=0.768, p=.446, Cohen's d=0.213, BF10=0.355), proportion spent in moderate activity (mean ± SD, patients: 0.109 ± 0.062, controls: 0.131 ± 0.043; t(50)=1.487, p=.143, Cohen's d=0.413, BF10=0.688), neither group spending any time in vigorous activity, number of bouts (mean ± SD, patients: 8.371 ± 5.703, controls: 9.667 ± 4.585; t(50)=0.903, p=.371, Cohen's d=0.251, BF10=0.389), total time of bouts (mean ± SD, patients: 137.213 ± 100.688, controls: 162.494 ±77.400; t(50)=1.015, p=.315, Cohen's d=0.282, BF10=0.425), and total activity bout count (mean ± SD, patients: 462028.583 ± 350630.645, controls: 596035.778 ± 253689.269; t(50=1.579, p=.121, Cohen's d=0.438, BF10=0.770).

SUPPLEMENTARY MODELLING ANALYSES

8.1 Parameter recovery

To asses the reliability of our modelling analysis (Wilson and Collins, 2019), for each model we performed parameter recovery analysis, where we simulated participants' responses using newly drawn individual-level parameters from the group-level distributions, for each group and for a pooled dataset.

We then fit the same model to the simulated data and calculated Pearson correlation coefficients r between the generated and estimated individual-level parameters. We recovered each individual parameter (for each member of the population dataset) 50 times and calculated the mean and SD of the correlation between the true and recovered parameter values. See results in S2-S4. The higher the coefficient r, the more reliable the estimates are, which can be categorised as: poor (if r <0.5); fair (if 0.5< r <0.75); good (0.75< r <0.9); excellent (if r >0.9) (White et al., 2018).

Moreover, to assess the number of simulations needed, we calculated the average error of the parameter recovery correlation (and its error) for each model, parameter, and dataset as a function of increasing number of simulation, as plotted in Figure S2. The average was obtained from the 1000 randomly chosen permutations of different simulations at each n (out of 50).

Table S2. Pearson correlation coefficients (SD) for parameter recovery analysis for each parameter from five RL models for a pooled dataset.

Model ID	LR1	LR2	R	P	ξ	d
1	0.91 (0.03)	0.75 (0.09)	0.87 (0.04)	0.78 (0.06)	0.67 (0.08)	0.92 (0.02)
2	0.90 (0.11)	0.75 (0.16)	0.83 (0.09)	0.79 (0.13)	0.61 (0.17)	
3	0.92 (0.02)	0.26 (0.11)	0.85 (0.04)	0.91 (0.03)	0.65 (0.10)	0.92 (0.02)
4	0.95 (0.02)	0.88 (0.04)	0.88 (0.03)	0.90 (0.03)	0.70 (0.11)	
5	0.95 (0.02)	0.88 (0.03)	0.94 (0.02)	0.92 (0.03)		

Table S3. Pearson correlation coefficients (SD) for parameter recovery analysis for each parameter from five RL models for patients.

Model ID	LR1	LR2	R	P	ξ	d
1	0.93 (0.03)	0.78 (0.11)	0.84 (0.05)	0.61 (0.11)	0.76 (0.09)	0.93 (0.03)
2	0.90 (0.11)	0.73 (0.20)	0.76 (0.16)	0.73 (0.17)	0.70 (0.19)	
3	0.90 (0.06)	0.20 (0.16)	0.83 (0.06)	0.90 (0.04)	0.69 (0.10)	0.91 (0.05)
4	0.96 (0.02)	0.91 (0.05)	0.80 (0.07)	0.81 (0.07)	0.73 (0.11)	
5	0.95 (0.03)	0.91 (0.04)	0.94 (0.03)	0.86 (0.05)		

Table S4. Pearson correlation coefficients (SD) for parameter recovery analysis for each parameter from five RL models for healthy controls.

Model ID	LR1	LR2	R	P	ξ	d
1	0.90 (0.05)	0.14 (0.22)	0.92 (0.04)	0.92 (0.03)	0.29 (0.16)	0.90 (0.04)
2	0.88 (0.11)	0.72 (0.17)	0.88 (0.11)	0.88 (0.08)	0.27 (0.25)	
3	0.91 (0.04)	0.22 (0.18)	0.91 (0.03)	0.93 (0.03)	0.23 (0.18)	0.89 (0.05)
4	0.92 (0.03)	0.83 (0.07)	0.94 (0.04)	0.94 (0.02)	0.26 (0.17)	
5	0.92 (0.03)	0.80 (0.10)	0.95 (0.02)	0.94 (0.03)		

Figure S2. Parameter recovery mean error of the correlation for each model and each parameter. The average error is plotted as a function of simulation number averaged across 1000 permutations of 50 simulations.

Figure S3. Neural correlates of punishment prediction error shared across the arthritis and control groups. Colorscale represents z scores ranging from 0 to 5, thresholded at z>2.6, p<0.05.

8.2 Model recovery

We also performed model recovery analysis (Wilson and Collins, 2019), where we first simulated responses using each model and then fit each model to the model-specific dataset (50 simulations for each pair). We then counted the number of times a model fit the simulated data best (according to the LOOIC rule), effectively creating an $M \times M$ confusion matrix, where M is the number of models. In the case where we have a diagonal matrix of ones, the models are perfectly recoverable and hence as reliable as possible. See the model recovery results for the pooled sample in Table S5.

Table S5. Confusion matrix from the model recovery analysis of five RL models for a pooled dataset.

		fit with				
		M1	M2	M3	M4	M5
generated	M1	0.52	0.08	0.28	0.10	0.02
	M2	0.22	0.08	0.22	0.34	0.14
	M3	0.44	0.02	0.36	0.10	0.08
	M4	0.22	0.14	0.22	0.22	0.20
	M5	0.12	0.16	0.14	0.28	0.30

SUPPLEMENTARY IMAGING RESULTS

Cluster ID	Voxels	P	Z peak	X	Y	Z
2a	16180	5.48e-30	5.62	42	36	34
2b			5.34	-32	0	56
2c			4.97	2	-64	54
2d			4.91	-44	20	38
2e			4.83	4	-72	50
2f			4.83	-4	-66	56
1a	8416	3.53e-19	4.81	32	-70	-44
1b			4.76	-34	-48	-30
1c			4.59	42	-66	-38
1d			4.58	-40	-56	-44
1e			4.57	-26	-66	-40
1f			4.57	20	-78	-20

Table S6. Activation clusters associated with punishment prediction error, common activity across patient and control groups. Coordinates in MNI space.

Figure S4. Neural correlates of reward prediction error shared across the arthritis and control groups. Colorscale represents z scores ranging from 0 to 5, thresholded at z>2.6, p<0.05.

Cluster ID	Voxels	P	Z peak	X	Y	Z
5a	34881	0	7.08	12	-86	-14
5b			6.59	-26	-76	-16
5c			6.52	26	-82	-16
5d			6.38	22	-84	-16
5e			6.35	-26	-84	-18
5f			6.22	26	-92	2
4a	2285	4.92E-17	4.47	2	-12	12
4b			3.83	8	-12	12
4c			3.79	8	-22	10
4d			3.79	28	4	24
4e			3.78	-10	-12	12
4f			3.72	28	10	18
3a	1381	9.49E-12	4.38	-30	2	54
3b			3.9	-38	0	24
3c			3.71	-40	4	24
3d			3.7	-42	4	34
3e			3.7	-38	28	24
3f			3.53	-38	30	38
2a	712	5.36E-07	3.81	-32	56	-6
2b			3.71	-20	52	-12
2c			3.54	-32	50	-8
2d			3.47	-28	42	-8
2e			3.46	-28	40	-12
2f			3.46	-26	60	6
1a	202	0.0346	3.98	10	56	-12
1b			3.94	12	58	-8
1c			3.6	8	60	-6
1d			3.35	26	60	-6
1e			3.09	26	56	-8
1f			2.92	20	52	-10

Table S7. Activation clusters associated with reward prediction error, positive mean activity across patient and control groups. Coordinates in MNI space.

Figure S5. Network centrality correlation results for other network centralities. Please refer to the supplementary file with network analyses results for precise values of Pearson's R and LogBF10.