Dietary protein source alters gut microbiota composition and function

Running Title: Protein source alters the gut microbiota

Authors: J. Alfredo Blakeley-Ruiz†*[1], Alexandria Bartlett*[1,2], Arthur S. McMillan[3], Ayesha Awan[1,3], Molly Vanhoy Walsh[1], Alissa K. Meyerhoffer[1], Simina Vintila[1], Jessie L. Maier[1], Tanner Richie[1], Casey M. Theriot[3], Manuel Kleiner†[1]

1. Department of Plant and Microbial Biology, College of Agricultural Sciences, North Carolina State University, Raleigh, NC, USA
2. Department of Molecular Genetics and Microbiology, Duke University, Durham, NC, USA
3. Department of Population Health and Pathobiology, College of Veterinary Medicine, North Carolina State University, NC, USA

* These authors contributed equally to this manuscript
† Please refer correspondence to J. Alfredo Blakeley-Ruiz (jablakel@ncsu.edu) or Manuel Kleiner (manuel_kleiner@ncsu.edu)

Orcid:
JABR: https://orcid.org/0000-0001-7638-5849
AB: https://orcid.org/0000-0003-4350-0542
ASM: https://orcid.org/0000-0002-0115-0225
AA: https://orcid.org/0009-0002-3421-4798
MVW: https://orcid.org/0000-0003-2723-7297
AKM: https://orcid.org/0009-0001-7635-1275
SV: https://orcid.org/0000-0003-0018-0016
JLM: https://orcid.org/0009-0001-8575-5386
TR: https://orcid.org/0000-0002-2554-6005
CMT: https://orcid.org/0000-0002-1895-8941
MK: https://orcid.org/0000-0001-6904-0287

ABSTRACT

The source of protein in a person's diet affects their total life expectancy. However, the mechanisms by which dietary protein sources differentially impact human health and life expectancy are poorly understood. Dietary choices have major impacts on the composition and function of the intestinal microbiota that ultimately modulate host health. This raises the possibility that health outcomes based on dietary protein sources might be driven by interactions between dietary protein and the gut microbiota. In this study, we determined the effects of seven different sources of dietary protein on the gut microbiota of mice using an integrated metagenomics-metaproteomics approach. The protein abundances measured by metaproteomics can provide microbial species abundances, and evidence for the molecular phenotype of microbiota members because measured proteins indicate the metabolic and physiological processes used by a microbial community. We showed that dietary protein source significantly altered the species composition and overall function of the gut microbiota. Different dietary protein sources led to changes in the abundance of microbial proteins involved in the degradation of amino acids and the degradation of glycosylations conjugated to dietary protein. In particular, brown rice and egg white protein increased the abundance of amino acid degrading enzymes. Egg white protein increased the abundance of bacteria and proteins usually associated with the degradation of the intestinal mucus barrier. These results show that dietary protein sources can change the gut microbiota's metabolism, which could have major implications in the context of gut microbiota mediated diseases.

Keywords: Gut Microbiome, Metaproteomics, Metagenomics, Dietary Intervention, Mus musculus, Mice

INTRODUCTION

Source of dietary protein has a major impact on human health. People who consume high amounts of animal protein have higher mortality rates than those who consume mostly plant-based protein [1, 2]. Egg protein and red meat protein in particular have been shown to lead to increased mortality rates among humans [3] and a diet high in red meat protein has been shown to increase inflammation in a mouse model of colitis [4]. Replacing animal protein sources with plant protein sources in humans reduces mortality rates [3]. Currently, we have a limited understanding of the underlying causes, but the gut microbiota has been implicated as potentially having a major role in the differential health impacts of different dietary protein sources [5, 6]. Diet has been shown to change the gut microbiota's composition and function in

ways that can be detrimental or beneficial to health in both mice and humans[7–10]. For

example, protein fermentation by the gut microbiota generates a number of toxins including

ammonia, putrescine, and hydrogen sulfide [6, 11], whereas fermentation of fiber and certain

amino acids produces anti-inflammatory short-chain fatty acids [12]. Previous studies in mice

demonstrate that the amount of protein can have a greater impact on the gut microbiota's

composition than other macronutrients [13], and that source of dietary protein alters gut

microbiota composition[14]. There is, however, limited data showing the mechanisms by which

individual sources of dietary protein affect the gut microbiota's composition and function, which

could mediate the consumption and production of compounds beneficial or detrimental to the

host.

Metaproteomics represents a powerful tool for characterizing the mechanisms

underlying dietary effects on gut microbiota [7, 9]. Metaproteomics is defined as the large-scale

characterization of the proteins present in a microbiome [15]. Protein abundances measured by

metaproteomics simultaneously provide microbial species abundances [16], and evidence for

the metabolic and physiological phenotype of microbiota members [17–19]. Metaproteomes are

usually measured using a shotgun proteomics approach where proteins extracted from a

sample are digested into peptides, separated by liquid chromatography, and measured on a

mass spectrometer [20]. Proteins are then identified and quantified using a database search

algorithm, which matches the measured peptides to a database of protein sequences [21]. Due

to the heterogeneous nature of complex microbial communities it is usually best to construct the

protein database using gene predictions from metagenomes measured from the same samples

[21]. When metaproteomics is coupled to a genome-resolved metagenomic database it is

possible to evaluate strain and species level function even if the microbes have not been

previously characterized [22, 23]. We call this approach integrated metagenomics-

metaproteomics.

We used an integrated metagenomic-metaproteomic approach to investigate the effects

of dietary protein source on gut microbiota composition and function. We hypothesized that

dietary protein source affects the abundance of amino acid metabolizing enzymes from the gut

microbiota, altering the abundance of pathways involved in the production of toxins detrimental

to host health. We found that the source of dietary protein not only alters the abundance of

amino acid degrading enzymes, but has an even greater impact on the abundance of glycan

degrading proteins among other functions, indicating that dietary protein sources can have wide

ranging effects on the gut microbiota.

105 MATERIALS AND METHODS

Figure 1: Source of dietary protein alters the gut microbiota's composition. (a) Diagram showing the experimental design,
with number of cages, and order of the diets fed. Colors depicting the diets are used throughout the manuscript. Each row of arrows
represents one cage. We collected 10-12 samples for each experimental diet and 5-6 samples for each control diet. (b) A diagram
illustrating the integrated metagenomic-metaproteomics method used to analyze the samples along with raw metrics: quantifiable
species and number of proteins. (c) Bray-Curtis dissimilarity between the initial 20% soy diet (teal) or 20% casein diet (red) and all
other diets. Error bars reflect 95% confidence intervals for all line graphs as calculated by the Rmisc package in R, but the
PERMANOVA values from Table 1 and Extended Data Table 5 are the true tests of significance. (d) Ratio of spectra assigned to
microbes versus the host; boxes represent 95% confidence intervals calculated on a linear mixed effect model (Extended Data
Table 2). (e) Shannon diversity index of the gut microbiota across all diets; boxes represent 95% confidence intervals calculated on
a linear mixed effect model; boxes that do not overlap are significantly different (Extended Data Table 2). (f) Abundances of the two
most abundant bacterial classes *Bacteroidia* and *Clostridia* based on summed protein abundance. Colors represent diets and the
legend is (a). Error bars are 95% confidence intervals calculated using a linear-mixed effects model and error bars that do not
overlap represent significant differences (Extended Data Table 2). (g) A hierarchically clustered (ward.D2 algorithm on euclidean
distances) heatmap depicting the clustering by species group abundance of the 36 most abundant species in the study. Species
were considered abundant if they had at least 5% of the microbial biomass in at least one sample.

Experimental overview, description of diets, animal housing and preliminary results

We used twelve C57BL/6J mice in two groups (six males, six females, Jackson Labs,

Bar Harbor) aged three months at the beginning of the study. The males and females originated

from different mouse rooms at the Jackson Labs and thus were expected to have different

background microbiomes. Mice from both groups were housed in two separate cages (3

mice/cage) with a 12 h light/dark cycle. We autoclaved bedding, performed all cage changes in

a laminar flow hood and maintained an average temperature of 70°F and 35% humidity. We

conducted our animal experiments in the Laboratory Animal Facilities at the NCSU CVM

campus (Association for the Assessment and Accreditation of Laboratory Animal Care

accredited), which are managed by the NCSU Laboratory Animal Resources. Animals assessed

as moribund were humanely euthanized via CO_2 asphyxiation. NC State's Institutional Animal

Care and Use Committee approved all experimental activity (Protocol # 18-034-B).

We fed the mice a series of nine fully defined diets (Fig. 1a, Supplementary table 1)

formulated and purchased from Envigo Teklad (inotiv, Madison, WI, USA) based on the diets

used a previous study . We worked with Envigo to formulate 20% or 40% protein (by weight)

diets, which required slight variations in the amount of protein source added for each protein

source. The torula yeast protein source has large amounts of heavily glycosylated mannoprotein

which resulted in a higher amount of the yeast protein source being added to achieve a 20%

protein content (Supplementary table 1). We chose 20% and 40% as the protein amounts for

this study partially due to the results of a previous study which showed that most of the effects

of amount of protein on the gut microbiota happened between 6% and 20% and that there was

not much of an effect from 20% to 40% [24]. Because the goal of the study was to investigate

the effects of source of dietary protein, we selected 20% protein as our main dietary protein

amount to avoid any unwanted microbiota effects due to small variations in amount of protein.

We included two additional diets with 40% as the amount of both to serve as an internal control

by providing the dietary protein source again and also to show that the increase from 20% to

40% would not have too much of an effect on the microbiota. In order of feeding, the diets were

20% soy protein, 20% casein protein, 20% brown rice protein, 40% soy protein, 20% yeast

protein, 40% casein protein, 20% pea protein, 20% egg white protein, and 20% chicken bone

protein. To control for the succession effects of the serial dietary intervention and the age of the

mice, we fed the mice the 20% soy diet or the 20% casein diet as an additional control at the

end of the diet series. In order of feeding, the diets were 20% soy protein, 20% casein protein,

20% brown rice protein, 40% soy protein, 20% yeast protein, 40% casein protein, 20% pea

protein, 20% egg white protein, and 20% chicken bone protein. To control for the succession
effects of the serial dietary intervention and the age of the mice, we fed the mice the 20% soy
diet or the 20% casein diet as an additional control at the end of the diet series. The chicken
bone diet caused the mice to lose weight, so we discontinued the diet after 3 days and the mice
consumed a standard chow diet for the rest of that week. No fecal samples were collected for
the chicken bone diet. All defined diets were sterilized by ɣ-irradiation and mice were provided
sterile water (Gibco). After feeding each diet for 7 days, we collected fecal samples, prior to
replacing food with the next diet. We collected samples in NAP preservation solution at a 1:10
sample weight to solution ratio, and roughly homogenized the sample with a disposable pestle
prior to freezing at -80°C [25]. We had to sacrifice one mouse during the second diet (20%
casein) so no additional samples were collected. We also were unable to collect a sample from
one of the mice during the brown rice and egg white diets so only 10 samples were collected for
those diets. We analyzed samples from all mice and diets using an integrated metagenomic-
metaproteomic approach [7, 23] (Fig. 1b).
172
Metagenomic DNA sequencing
To create a database for metaproteomic analysis, we pooled fecal samples from each
cage to create four cage specific metagenomes. The metagenomes were only used to generate
a protein sequence database for metaproteomics, and all results presented in the manuscript
are derived from protein quantities acquired by metaproteomic measurements. We gathered
one fecal sample from each cage for four different diets (20% rice, 40% soy, 20% yeast, 40%
casein) for a total of 16 samples.To remove the preservation solution from the samples, we
added 5 mL of 1X Phosphate Buffered Saline solution (VWR) to the samples and centrifuged
them (17,000 x g, 5 min) to pellet solids and bacterial cells in suspension. We removed the
preservation solution and resuspended the fecal pellets in 1 mL of InhibitEX Buffer in Matrix E
(MP Biomedicals) bead beating tubes. We beat the samples at 3.1 m/s for 3 cycles with 1
minute of ice cooling between each cycle using a Bead Ruptor Elite 24 (Omni International). We
isolated DNA from the resulting lysate using the Qiagen QIAamp Fast DNA stool mini kit (cat.
No. 51604) [26]. Samples were extracted individually and pooled by cage with each sample
contributing a total of 200 ng of DNA.
We submitted genomic DNA to the North Carolina State Genomic Sciences Laboratory
(Raleigh, NC, USA) for Illumina NGS library construction (TruSeq Nano Library kit) and
sequencing (NovaSeq 6000 sequencer). We obtained between 51,152,549 and 74,618,259
paired-end reads for each of the 4 samples.

Metagenomic assembly and protein database construction

We assembled raw reads using a genome resolved metagenomics approach. We
removed PhiX174 (NCBI GenBank accession CP004084.1) and mouse genome (mm10)
contaminating sequences using BBSplit and removed adapters using BBDuk (BBMap, Version
38.06), parameters: mink = 6, minlength =20 [27]. We assembled decontaminated reads
individually using MetaSPAdes (v3.12.0) -k 33,55,99 [28] and co-assembled them using
MEGAHIT (v1.2.4) --kmin 31 --k-step 10 [29]. We mapped reads from all four samples to all five
assemblies using bbmap, and binned the contigs using MetaBAT (v2.12.1) [30]. We assessed
the quality of the bins using CheckM (v1.1.3) [31] and automatically accepted medium quality
bins with a completion score greater than 50% and a contamination score of less than 10% [32].
Because the purpose of metagenomics in our study was to generate a comprehensive protein
sequence database and to assign proteins to species, we further accepted bins that were
greater than >30% complete and <5% contaminated. We do not think that genomes of lower
than quality than medium, should be contributed to major public databases like those hosted by
NCBI and ENA so we have made these MAGs available in Dryad instead (see data availability
section). We clustered the bins into species groups by 95% ANI using dRep (v2.6.2) [33, 34]
and assigned taxonomy using GTDB-Tk (v1.3.0, ref r95) [35].

We assembled the protein database by combining gene annotations from the
metagenome with mouse and dietary protein databases [21]. For the metagenome, we
annotated the assemblies prior to binning and then for each bin individually using PROKKA
(Version 1.14.6)[36]. If the contig was binned, we compiled the annotations from the bins. We
then used CD-HIT-2D (Version 4.7), with a 90% identity cutoff, to compare the genes from the
unbinned PROKKA output to the binned gene annotation [37]. If a gene was not present in a bin
we added it to the database as an unbinned sequence. Once we compiled the microbial protein
database we assigned each protein sequence a species code if it was species specific or an
ambiguous, low-quality, or unbinned code if it was assigned to more than one species group,
belonged to a low-quality bin, or was not present in a bin. In addition to the microbial
sequences, we added a *Mus musculus* proteome (UP000000589, Downloaded 19Feb20), and
the relevant dietary protein database for each sample: *Glycine max* (UP000008827,
Downloaded 19Feb20), *Bos taurus* (UP000009136, Downloaded 19Feb20), *Cyberlindnera*
jadinii (UP000094389, Downloaded 25May20), *Oryza sativa* (UP000059680, Downloaded
25May20) and *Gallus gallus* (UP000000539, Downloaded 25May20). Due to the lack of a
reference proteome for the yellow pea diet, we created a custom pea reference with all available

UniProtKB [38] protein sequences for *Pisum sativum* (Taxon ID: 388 Downloaded 25Apr20) and

the reference proteome of *Cajanus cajan* (UP000075243, Downloaded 25May20). For T0

samples taken when mice were fed a standard chow diet, we added proteomes from the protein

sources likely to be in the diet based on the ingredient list (corn UP000007305, fish

UP000000437, soy UP000008827, wheat UP000019116, Downloaded 19Feb20). We clustered

the mouse and diet reference proteomes individually at a 95% identity threshold. We only

searched samples against their respective dietary database. The databases ranged in size from

597,215 to 809,468 protein sequences depending on the dietary components, each database

had the same number of microbial and host proteins.

In order to identify all sequences from the species *Bacteroides thetaiotaomicron* and

Lactococcus lactis we downloaded all the sequences matching these species from UniProt [38].

We then used diamond BLASTp to identify all sequences in the protein database that matched

with 95% identity or greater. The species code for these sequences was changed to BT or LAC

if they were found to be *B. thetaiotaomicron* or *L. lactis* respectively.

Metaproteomic sample processing

We extracted protein using a modified FASP protocol[39]. We pelleted fecal samples by

centrifugation (21,000 x g, 5 min) and removed the preservation solution. We suspended dietary

and fecal pellets in SDT lysis buffer [4% (w/v) SDS, 100 mM Tris-HCl pH 7.6, 0.1 M DTT] in

Lysing Matrix E tubes (MP Biomedicals) and bead beat the samples (5 cycles of 45 s at 6.45

m/s, 1 min between cycles). After bead beating we heated the lysates to 95°C for 10 minutes.

We mixed 60 µL of the resulting lysates with 400 µL of UA solution (8 M urea in 0.1 M Tris/HCl

pH 8.5), loaded the sample onto a 10 kDa 500 µL filter unit (VWR International) and centrifuged

at 14,000 x g for 30 minutes. We repeated this step up to three times to reach filter capacity.

After loading, we added another 200 µL of UA buffer and centrifuged at 14,000 x g for another

40 minutes. We added 100 µL of IAA solution (0.05 M iodoacetamide in UA solution) to the filter

and incubated at 22°C for 20 minutes. We removed IAA by centrifuging the filter at 14,000 x g

for 20 minutes. We then washed the filter 3 times by adding 100 uL of UA buffer and

centrifuging at 14,000 x g for 20 minutes. We then washed 3 more times by adding 100 uL of

ABC buffer (50 mM Ammonium Bicarbonate) and centrifuging at 14,000 x g for 20 minutes. To

digest the isolated protein, we added 0.95 µg of MS grade trypsin (Thermo Scientific Pierce,

Rockford, IL, USA) mixed in 40 µL of ABC to each filter and incubated at 37°C for 16 hours. We

eluted the peptides by centrifugation at 14,000 x g for 20 minutes. We eluted again with 50 uL of

0.5 M NaCL and centrifuged at 14,000 x g for another 20 minutes. We quantified the abundance

of the peptides using the Pierce Micro BCA assay (Thermo Scientific Pierce, Rockford, IL, USA)

following the manufacturer's instructions.

We analyzed the samples by 1D-LC-MS/MS. Samples were run in randomized block

design. For each run, we loaded 600 ng of peptides onto a 5 mm, 300 µm ID C18 Acclaim®

PepMap100 pre-column (Thermo Fisher Scientific) using an UltiMate 3000 RSLCnano Liquid

Chromatograph (Thermo Fisher Scientific) and desalted on the pre-column. After desalting, the

pre-column was switched in line with a 75 cm x 75 µm analytical EASY-Spray column packed

with PepMap RSLC C18, 2 µm material (Thermo Fisher Scientific), which was heated to 60 °C.

The analytical column was connected via an Easy-Spray source to a Q Exactive HF Hybrid

Quadrupole-Orbitrap mass spectrometer. Peptides were separated using a 140 minute reverse

phase gradient [24]. We acquired spectra using the following parameters: m/z 445.12003 lock

mass, normalized collision energy equal to 24, 25 s dynamic exclusion, and exclusion of ions of

+1 charge state. Full MS scans were acquired for 380 to 1600 m/z at a resolution of 60,000 and

a max IT time of 200 ms. Data-dependent MS^2 spectra for the 15 most abundant ions were

acquired at a resolution of 15,000 and max IT time of 100 ms.

Metaproteomic data processing

We searched raw MS spectra against the diet specific protein databases using the run

calibration, SEQUEST HT and percolator nodes in Proteome Discoverer 2.3 (Thermo Fisher

Scientific). We used the following setting for search: trypsin (full), 2 missed cleavages, 10 ppm

precursor mass tolerance, 0.1 Da fragment mass tolerance. We included the following dynamic

modifications: oxidation on M (+15.995 Da), deamidation on N,Q,R (0.984 Da) and acetyl on the

protein N terminus (+42.011 Da). We also included the static modification carbamidomethyl on

C (+57.021 Da). We filtered identified peptides and proteins at a false discovery rate (FDR) of

5%. Additionally, we only included proteins that had at least one protein unique peptide

identified. Proteins were quantified by peptide spectral match (PSM) count (spectral counting).

Statistical analysis and visualization

Whenever possible we tested significance of changes in abundance by applying an

ANOVA on a linear mixed effects model with the interacting fixed effects being mouse group

and diet, and the random effect being the individual mouse (lme4 version 4.3.1) [40]. For

multiple comparisons we calculated 95% confidence intervals for each diet using the emmeans

R package (version 1.8.8) [41]. The exceptions were PERMANOVA analysis for testing

significance of microbiota compositional changes (Table 1; Extended Data Table 5) and Welch's

t-tests to compare differences between yeast and egg white protein diets (Extended Data Table

17, 18 and 20). For each analysis, we controlled for multiple-hypothesis testing by converting *P*

values to FDR-level based q-values, unless all *P* values in the analysis were below 0.05 using

Benjamini-Hochberg procedure [42, 43]. By definition, if all the *P* values are less than 0.05 than

the FDR is less than 0.05. Visualizations were produced using ggplot2 (version 3.4.3) [44],

pheatmap (version 1.0.12) [45], RawGraphs [46], Microsoft Excel and Adobe Illustrator. All

boxes and error bars represent 95% confidence intervals. Boxes or error bars that do not

overlap denote significance. If no error bars are present then significance is denoted by letters

or asterisk. In the case of beta-diversity analysis the error bars are 95% confidence intervals

(Fig. 1c), but significance was tested separately by PERMANOVA.

Compositional profiling of gut microbiota

We calculated the abundances of gut microbes using proteinaceous biomass [16].

Briefly, we filtered for proteins with at least 2 protein unique peptides and summed their spectra

into their assigned taxonomy: microbial species, mouse, diet, ambiguous, low quality bins,

unbinned bacteria. We calculated the microbe to host ratio by summing the spectral count

assigned to microbial species, multiple microbial species (ambiguous), low quality bins and

unbinned bacteria proteins and dividing them by the number of spectral counts assigned to

mouse proteins. We considered a microbial species quantifiable if we could identify at least one

protein with 2 protein unique peptides unambiguously assigned to the species. If a protein

matched to more than one species, it was not included in the quantification. We showed

previously that these filtering and inclusion/exclusion criteria allow for accurate biomass

estimates [16].We calculated per sample species richness by simply counting the number of

quantifiable species per sample. We calculated alpha (Shannon Diversity Index) and beta

diversity (Bray-Curtis) metrics using the vegan (version 2.6-4) package in R (version 4.3.1) [47,

48] on a table of the quantifiable microbial species (statistics as described above). We also

evaluated gut microbiota composition using principal component analysis and hierarchical

clustering. For principal component analysis we normalized the quantified species using

centered-log ratio transformation and calculated principal components using the prcomp

function in base R on all the mice and separately on each mouse group. Principal components

were rendered using the ggplot2 (version 3.4.3) package in R [44]. For hierarchical clustering,

we focused on the species that were most abundant, representing at least 5% of the microbial

species biomass in at least one sample. We calculated the percent biomass for all the species

and then extracted the species that fit the abundant species criteria. We calculated the

individual significance of each abundant species using linear mixed effects models as described

above. We hierarchically clustered log transformed values of these species using the R package

pheatmap (version 1.0.12), using the ward.D2 algorithm and euclidean distances [45]. To

compare broad taxonomic changes at the class level, for all quantifiable species, we summed

the abundance of the assigned class by GTDB-Tk.

Functional profiling of gut microbiota

For analyses of functional categories at the level of the whole microbiota we calculated

the normalized spectral abundance factor (NSAF%) for each protein, which provides the relative

abundance for each protein as a percentage of the summed abundance of microbiota proteins

[49]. We annotated functions for all microbial proteins in our database using EggNOG-mapper

[50], MANTIS [51], and Microbe Annotator [52]. We assigned glycoside hydrolase protein family

identifiers from the CAZy database using dbCAN2 [53, 54]. We manually curated these

annotations by searching a subset of these proteins against the Swiss-Prot [38] and InterPro

[55] databases between February 2023 and June 2023 . If the Swiss-Prot or InterPro

annotations matched the automated tool annotations we extrapolated the assigned protein

name to all proteins with the same automated annotation. Alternatively, if the annotations from

the automated tools were in agreement, we consolidated the annotation into a consensus

annotation. We then assigned broad functional categories, detailed functional categories, and

specific names to each validated protein set. To evaluate functional changes due to diet, we

summed all microbiota proteins assigned to a broad or detailed functional category, or enzyme

name and applied a linear mixed-effects model to each function.

In vivo proteomic analysis of B. thetaiotaomicron

To analyze the *B. thetaiotaomicron* proteome, we calculated the orgNSAF by extracting

all proteins assigned to the species *B. thetaiotaomicron* detected in the metaproteomes, and

then calculating NSAF% [17]. We then compared abundances of *B. thetaiotaomicron* proteins

detected in the yeast and egg protein diets using the Welch's t-test in the Perseus software

(version 1.6.14.0) [56]. To visualize polysaccharide utilization loci (PULs) we mapped the reads

from one of our metagenomic samples to all the contigs that were assigned *B. thetaiotaomicron*

proteins using BBSplit (BBMap, Version 38.06). We then assembled all the mapped reads using

metaSPAdes. The genes in this newly assembled genome overlapped exactly with the previous

set of identified *B. thetaiotaomicron* genes, and this *B. thetaiotaomicron* genome was uploaded

to the RAST server for further analysis [57]. PULs were detected in the metaproteome by

identifying proteins labeled SusC, SusD, or TonB. The rest of the PUL was identified by

visualizing the gene neighborhood in RAST. The identified genes were then cross referenced

against PULDB to assign literature described PUL numbers [58].

In vitro growth and proteomics of B. thetaiotaomicron

We cultured *B. thetaiotaomicron* VPI-5482 in two biological replicates and at least 4

technical replicates using a defined *Bacteroides* medium [59]. *B. thetaiotaomicron* cultures were

grown statically at 37°C in a Coy anaerobic chamber (2.5 % H_2 /10 % CO_2 /88.5 % N_2) in

minimal medium (100 mM KH_2PO_4, 8.5 mM $[NH_2]_4SO_4$, 15 mM NaCl, 5.8 µM vitamin K_3, 1.44

µM $FeSO_4 \cdot 7H_2O$, 1 mM $MgCl_2$, 1.9 µM hematin, 0.2 mM L-histidine, 3.69 nM vitamin B_{12}, 208

µM L-cysteine, and 7.2 µM $CaCl_2 \cdot 2H_2O$). The four dietary protein sources: soy protein

(CA.160480), yeast protein (CA.40115), casein protein (CA.160030), and egg white protein

(CA.160230), were purchased from Envigoand were the same as the protein sources used in

the corresponding diets. Porcine muc2 mucin (Sigma CA. M2378) was also tested alongside

controls of glucose and no carbon source. To aid in suspension in aqueous media, we pre-

prepared the proteins in 200 mM NaOH water at 37°C for four days; the glucose control was

also dissolved in 200 mM NaOH water. We then added the protein or glucose solution to the

pre-prepared media at 0.5% (wt/v). Cultures were grown overnight in minimal media

supplemented with 0.5% (wt/v) glucose before being washed and inoculated into experimental

conditions at 0.01 OD and incubated at 37°C anaerobically with shaking every hour. Colony

forming units (CFUs) per mL of culture were enumerated by drip plating at 0 and 24 hr post

inoculation. Solid media for *B. thetaiotaomicron* was Brain-Heart Infusion agar (Difco CA.

241830) supplemented with 10% Horse Blood (LAMPIRE CA. 7233401) (BHI-HB).

To obtain samples for proteomics, we repeated the experiment for the glucose, yeast,

egg white, mucin and soy media. After 8 hours, CFUs were enumerated to confirm growth. We

pelleted cells by centrifuging at 4,000 g for 10 minutes. We then extracted the supernatant and

froze the pellets at -80°C. Protein was extracted by the same FASP protocol described above

but with two differences. We lysed pellets by adding 120 uL of SDT buffer and then heating at

95°C. We used PES 10kDa filters (MilliporeSigma). We also used a similar Mass Spectrometry

procedure, except the samples were run on an Exploris 480 mass spectrometer (Thermo Fisher

Scientific) and 1 ug of peptide were analyzed for each sample. We searched raw MS spectra

using the same Proteome Discoverer 2.3 workflow using the *B. thetaiotaomicron* proteome

downloaded from UniProt (UP000001414 downloaded January 9, 2024) as the protein

sequence database. We cross referenced PULs between the metaproteome and the *in vitro*

proteome to compare them.

RESULTS

Source of dietary protein alters gut microbiota composition

Shotgun sequencing of the fecal samples using a genome-resolved metagenomics

pipeline [33, 60] resulted in 454 metagenome-assembled genomes (MAGs) organized into 180

species groups. We used high-resolution mass spectrometry based metaproteomics to identify

and quantify proteins in each sample using a protein sequence database derived from the

metagenome and augmented with mouse and diet protein sequences [20, 21]. In total, we

identified 35,588 proteins, each distinguished as microbial, host, or dietary proteins (Extended

Data Table 1). We quantified the proteinaceous biomass for each species using

metaproteomics data and obtained measurements for 161 distinct species (Extended Data

Table 3) [16]. PERMANOVA analysis on the 161 quantified species revealed that protein source

and mouse group best explained the variance in gut microbiota composition with R^2 values of

0.39 and 0.29 respectively, whereas amount of protein in the diet and age of the mouse had

little effect (R^2 values 0.01 or less (Table 1). Pairwise comparisons using Bray-Curtis showed

that the source of dietary protein significantly changed the composition of the gut microbiota in

43 out of 49 comparisons (FDR controlled PERMANOVA q < 0.01; Extended Data Table 5), and

that the yeast and egg white diets had the most dissimilar gut microbial communities(Fig. 1c). In

all cases our internal controls 40% soy and casein diets in the middle of the study, and the 20%

soy and casein control feedings at the end of the study had the least dissimilar Bray-Curtis

values when compared to the initial 20% soy or 20% casein diets respectively, which helped to

validate the initial rationale described in the methods. We also assessed the ratio of microbial to

host proteins as proxy for bacteria load and the alpha diversity (within sample diversity) using

the Shannon Diversity index. We found that bacterial load significantly increased in the yeast

diet (Fig. 1d) and that within sample diversity significantly decreased in the yeast and egg white

diets (Fig. 1e).

The large differences in microbial composition in the egg white and yeast protein diets

were driven by a decrease in the abundances of species from the class *Clostridia* in favor of

species from the class *Bacteroidia* (Fig. 1f). Because we observed fewer species in the class

Bacteroidia overall, it makes sense that a drop in *Clostridia* in favor of *Bacteroidia* would result

in a lower alpha diversity in the yeast and egg white diets (Fig. 1g; Extended Data Table 3).

Table 1: PERMANOVA analysis of the factors that explain the variance proteinaceous biomass gut microbiota species. Table of proteinacieous biomass of species was analysed using the adonis2 function in the vegan package in R. Similarity matrix was calculated using Bray-Curtis and the factors were protein source, amount of protein, mouse group, and age in weeks. The strata function was used to account for repeated measures. R^2 describes how much of the variance is explained by the factor, whereas F determines if the difference is significant.

Factor	df	SumOfSqs	R^2	F	P
Protein source	7	12.087	0.39	14.95	0.001
Amount of protein	2	0.565	0.01	2.2	0.003
Mouse group	1	7.118	0.21	56.9188	0.001
Mouse age	1	0.317	0.0095	2.533	0.017

We analyzed the most abundant microbial species (>5% of the microbial protein biomass in at least one sample) and hierarchically clustered them by abundance across the different dietary protein sources/groups (Fig. 1g). This revealed three major clusters separating most samples by mouse group with the exception of the yeast and egg white diets which together formed a separate cluster that internally showed separation by mouse group. The T0 samples fell into the major mouse group clusters which indicates that the two mouse groups had distinct gut microbial communities at the start of the experiment. Within the mouse group clusters the abundance of gut microbes clustered by source of dietary protein, which was also observed in principal component analysis (Supplementary Fig. 1, Supplementary Results Section A). In the yeast diet, *B. thetaiotaomicron* dominates gut microbiota composition regardless of mouse group. *B. thetaiotaomicron* abundance also increased in response to the egg white diet. There were additional species specific to each mouse group that also increased in abundance in response to the egg white diet (Fig. 1g; Supplementary Fig. 2). In group 1, these species were *Akkermansia muciniphila* and *Atopobiaceae* bacterium AB25-9, but in group 2 these species were *Paramuribaculum* sp. and *Dubosiella newyorkensis.* Both *A. muciniphila* [61] and *Paramuribaculum* sp. [62] have been reported to forage on intestinal mucin and *B. thetaiotaomicron* has been shown to switch towards mucin foraging when fed a low-fiber diet [63]. These results show that the source of dietary protein changes the gut microbiota's composition and suggests that an egg white diet could promote mucin-foraging bacteria.

Source of dietary protein alters gut microbiota function

We used the normalized abundances of gut microbiota proteins as a measure of the

investment into metabolic and physiological functions [17, 18, 64]. We first used automated

annotation tools to assign functions to proteins. Because the annotations from these tools were

not always accurate, we manually curated the annotations of 3,959 proteins and then

extrapolated the functions to 14,547 similarly annotated proteins, which in total represented

between 74 and 86 percent of the total microbial protein abundance in each sample (Extended

Data Table 6). Based on the annotations, we assigned broad functional categories, such as

amino acid metabolism, gene expression, glycan degradation, and monosaccharide metabolism

and more detailed functional categories, such as ribosomal proteins and glycolysis to each of

these proteins (Extended Data Table 6; Supplementary Fig. 3; Supplementary Results Section

B). We then used the relative protein abundances to determine the investment of gut microbes

into each of these functions. All of the broad functional categories, except for secondary

metabolism, had significant changes in abundance due to dietary protein (ANOVA, P value <

0.05; Extended Data Table 7; Fig. 2; Supplementary Fig. 4), which indicates that the source of

dietary protein changes the gut microbiota's metabolism and physiology. Hierarchical clustering

of all samples by abundances of broad functional categories revealed that the yeast and egg

white diets clustered separately from all the other diets (Supplementary Fig. 5), similar to the

results from the taxonomic clustering (Fig. 1g); however, a similar analysis at the detailed

functional level revealed separate yeast, rice, and egg white clusters, with some outliers

(Supplementary Fig. 6).

The two abundant broad functional categories (>1% of the total protein biomass) that

had the greatest effect size due to diet were amino acid metabolism and glycan degradation,

with F values of 29 and 93 respectively (Fig. 2; Extended Data Table 7). Amino acid metabolism

increased in the brown rice and egg white diets relative to all other diets except the 40% casein

diet and glycan degradation significantly increased in yeast and egg white diets relative to all

other diets (Fig 2). Significant changes in the abundance of amino acid metabolism supported

our initial expectation that the response of gut microbiota to different dietary protein sources

would likely relate to amino acid metabolism; however, we also found that the abundance of

glycan degrading enzymes responded more strongly to the source of dietary protein than did

enzymes for amino acid metabolism. This suggests that glycan degradation instead of amino

acid metabolism may be the major driver of taxonomic and functional changes in the gut

microbiota in response to dietary protein source. We discuss these two functions in detail in

subsequent sections. In addition, we observed specific changes in the abundance of enzymes

associated with the gene expression, monosaccharide metabolism, fermentation, and stress

and cell protection functional categories (Supplementary Results Section B; Supplementary Fig
3).

Figure 2: Broad functional categories of microbial proteins change significantly in abundance due to the source of dietary protein. Abundance of broad functional categories that represent at least 1% of the microbial protein abundance in at least one diet. The abundance is a modeled mean calculated from mixed effects models and the error bars represent 95% confidence intervals calculated from these models. All the categories shown here had a *P* value for the diet factor below 0.05; *P* value < 2.2x10^{-16} is the lower limit of the method. For underlying data see Extended Data Tables 7 and 8. For higher resolution of functional categories, e.g. fermentation, see Supplementary Figs. 3-4.

Source of dietary protein alters the abundance of amino acid degrading enzymes

We manually classified 911 proteins (Extended Data Table 9) representing 68 enzyme

functions (Extended Data Table 10) according to their involvement in the degradation (Fig. 3a),

synthesis (Fig. 3b), interconversion (Fig. 3c), or reversible (Fig. 3d) reactions of specific amino

acid pathways (Supplementary Figs. 7-18). In all diets except the yeast and standard chow

diets, we observed that gut microbiota trended towards amino acid degradation instead of

synthesis. We found that amino acid degrading enzymes were on average 2- to 6-fold more

abundant than amino acid biosynthesis enzymes (Fig. 3a and 3b). Amino acid degrading

enzymes were significantly more abundant in the rice and egg diets as compared to all the other

diets (Fig. 3a), which is consistent with the observation that dietary proteins were significantly

more abundant in the fecal samples of the brown rice and egg diets as compared to all other

diets (Fig. 3e), suggesting that there may be a connection between the digestibility of dietary

protein and amino acid degradation by the gut microbiota. Though amino acid synthesis

enzymes were generally less abundant, we did observe a trend towards an increase in amino

acid synthesis enzymes in the yeast protein diet relative to the other diets. This trend was not

significant (ANOVA, P value = 0.09; Extended Data Table 11;Fig. 3b), but we observed several

individual synthesis enzymes to be significantly increased in the yeast protein diet relative to

other diets. These enzymes were involved in the synthesis of branched-chain amino acids

(Suppl. Fig. 8), cysteine (Suppl. Fig. 9), lysine (Suppl. Fig. 15), proline (Suppl. Fig. 17), or

tyrosine (Suppl. Fig 18).

Not all amino acid degrading enzymes increased in both the brown rice and egg white

diets; sometimes they increased in one or the other (Supplementary Figs. 7-18). For example,

enzymes associated with the degradation of threonine were more abundant in the egg white diet

(Supplementary Fig. 12), whereas enzymes associated with tryptophan degradation were

increased in the brown rice diet (Supplementary Fig. 18). Brown rice and egg were not the only

diets in which the abundances of specific amino acid degrading enzymes increased. Alanine

dehydrogenase increased in the 40% soy diet relative to the pea, yeast, 20% soy, and 20%

casein diets (Supplementary Fig. 9) and cysteine desulfurase increased in the 40% casein and

casein control diets relative to most other diets (Supplementary Fig. 9).

Changes in amino acid degradation by the gut microbiota have potential implications for

host health by directly affecting local tissues or through interactions along the gut-brain axis

depending on the metabolites produced by amino acid degradation pathways or the enzyme

pathways themselves [6, 65]. For example, ammonia produced by amino acid deamination and

hydrogen sulfide produced from methionine and cysteine degradation are toxic [66–68], indoles

produced by tryptophanase and γ-aminobutyric acid (GABA) produced by glutamate

decarboxylase are neurotransmitters [69, 70], branched-chain fatty acids produced by

branched-chain amino acid degradation have been suggested to be anti-inflammatory [71, 72],

and proline metabolism has been linked to the depression [73] and enteric infections [74]. We

found ammonia producing enzymes to be significantly more abundant in the brown rice diet as

compared to all other diets, and also more abundant in the egg white and 40% casein diets as

compared to the standard chow, 20% soy, yeast, pea, and control diets (Fig. 3f and 3g).

Tryptophanase significantly increased in the brown rice diet relative to all other diets, whereas

glutamate decarboxylase increased in the egg white diet relative to all other diets except brown

rice, pea, and the control diets (Fig. 3h and Supplementary Fig. 7). We observed that branched-

chain amino acid degrading enzymes were significantly increased in the egg white protein diet

relative to all other diets (Fig. 3i), and proline degrading enzymes were increased in the brown

rice diet relative to other diets, except the 40% soy and 40% casein diets where we also

observed proline degradation to be significantly increased relative to the standard chow, yeast,

and pea diets (Supplementary Fig. 17). These results show that the source of dietary protein

can alter overall amino acid metabolism in the gut microbiome, as well as the abundance of

different pathways. These changes have the potential to affect host physiology and health.

Figure 3: Amino acid degradation increases in the rice and egg diets. Box plots depict the percent abundance (out of the total microbial protein abundance) of different categories of microbial amino acid metabolism proteins. The exception is the dietary proteins, which are based on the percent protein abundance of the total metaproteome. The boxes represent the 95% confidence interval of the mean (line) for each diet from a complete mixed-effects model, and the q-values represent the FDR controlled *P* values for the diet factor from an ANOVA on these models (q<0.05 indicates significance)(Extended Data Tables 11-12). Boxes that do not overlap indicate statistical significance. Circle dots represent actual values per sample and are colored by mouse group. Abundance of proteins classified as (a) degrading an amino acid, (b) synthesizing an amino acid, (c) converting between two amino acids, and (d) reversible. (e) Abundance of all dietary proteins detected in each condition. (f) Abundance of enzymes that are likely to produce ammonia. Enzymes classified as degrading or reversible were included as long as ammonia was one of the potential products. Summed abundance of all proteins classified as (g) urease, (h) tryptophanase, and (i) involved in branched-chain amino acid (BCAA) degradation to branched-chain fatty acid (BCFA) (includes branched-chain amino acid aminotransferase or ketoisovalerate oxidoreductase).

Gut microbes express distinct glycoside hydrolases to grow on different sources of dietary protein

Glycan degrading enzymes (glycoside hydrolases) showed the largest overall changes

in response to dietary protein source (Fig. 2, Extended Data Table 7). Specifically, these

enzymes increased significantly in abundance in the yeast and egg white diets compared to the

other diets. To further investigate the interaction of these glycan degrading enzymes with dietary

protein we manually curated the functional assignments and potential substrate specificity of the

1,059 microbial glycoside hydrolases detected in our metaproteomes (Extended Data Table 13).

We grouped the validated glycoside hydrolases into 91 families based on the CAZy

database (Extended Data Table 14) [53]. Of these families, 54 significantly changed in

abundance between the different dietary protein sources (ANOVA, q<0.05) (Extended Data

Table 15). Different glycoside hydrolase families increased in abundance in the soy, casein,

brown rice, yeast, and egg white diets suggesting that distinct glycans drive their abundance

changes across the different diets (Fig. 4a, Extended Data Table 14-16, Supplementary Results

Section C). The most abundant glycoside hydrolase families, GH18 in the case of egg white and

GH92 in the case of yeast, have previously been associated with the degradation of glycans

conjugated onto proteins (glycosylations) as part of polysaccharide utilization loci (PULs). PULs

are operons that contain all the proteins necessary to import and degrade a specific glycan

structure [75]. These GH18s are endo-β-N-acetylglucosaminidases that break the bond

between two acetylglucosamine residues attached to asparagine in N-linked glycoproteins. This

reaction releases the glycan from the glycoprotein [76]. Meanwhile, GH92s, which are alpha-

mannosidases, have been previously associated with the release of mannose residues from the

glycosylations on yeast mannoproteins [76].

We found that the total abundance of glycoside hydrolases increased from <1% in the

majority of diets to >2.5% in the yeast and egg diets (Fig. 4b). Additionally, we observed a

general trend towards an increased abundance of glycoside hydrolases in all defined diets

compared to the T0 (standard chow) diet; however, the increase was only significant for the soy,

yeast and egg diets (Fig. 4b). The majority of the glycoside hydrolases in the yeast and egg

diets came from *B. thetaiotaomicron* (Fig. 4b). Because *B.thetaiotaomicron* is one of the primary

drivers of the changes in microbiota composition in these diets (Fig. 1g), this suggests that

glycoside hydrolases are closely associated with the observed changes in microbiota

composition.

To examine the specific role of *B. thetaiotaomicron*in glycan degradation in the yeast

and egg white diets, we compared the abundances of all *B. thetaiotaomicron* proteins in the

metaproteome between the two diets. Out of 1,420 detected *B. thetaiotaomicron* proteins, the

abundances of 592 proteins significantly differed between the two diets (q < 0.05, Welch's t-test)

(Fig. 4c; Extended Data Table 17). Many of the significant proteins that were the most abundant

and had the greatest fold-change between the two diets came from PULs (Fig. 4c;

Supplementary Fig. 19). Between 10% and 25% of the total protein abundance of *B.*

*thetaiotaomicron*in in the yeast and egg white diets came from these PULs (Extended Data

Table 18). The proteins belonging to each PUL tended to be expressed together either being

significantly increased in egg white or the yeast diet (Figure 4c).

Figure 4: Glycosylations on dietary proteins drive shifts in microbial composition. (a) Mean summed protein abundance per
diet of glycoside hydrolases with significantly different abundances between diets (q<0.05 in mixed-effects ANOVA models). (b)
Mean combined protein abundance of proteins predicted to be glycoside hydrolases. The proportion of these proteins that belong to
B. thetaiotaomicron is highlighted. Diets that do not have overlapping letters also have non-overlapping 95% confidence intervals for
each diet calculated from a complete mixed-effects model. (c) Volcano plot of -log$_{10}$ *P* values (Welch's t-test; FDR controlled at
q<0.05) versus the log$_2$ fold-change of *B. thetaiotaomicron* proteins under the yeast and egg white protein diets *in vivo* after
recalculating the protein abundance based on proteins only assigned to *B. thetaiotaomicron*. Filled circle symbols, indicating
individual proteins, were colored based on the polysaccharide utilizing locus (PUL) operon to which the protein belongs. We only
colored the proteins from PULs that had an absolute difference of 0.5% or greater between the yeast and egg diets. (d) Colony
forming units per mL (CFU/mL) of *B. thetaiotaomicron* grown in defined media with dietary proteins as the sole carbon source. The
dotted line indicates T0 CFU/mL. Media that do not share letters are significantly different based on ANOVA and Tukey HSD
multiple comparisons after log transformation (*P* value < 0.05). (e) Hierarchical clustering (ward.D2 on Euclidean distances) of the *in*
vitro B. thetaiotaomicron proteome under different media. (f) *In vivo* and *in vitro* comparison of the summed protein abundance of
PULs. The bottom axis depicts the log$_2$ fold-change between egg white and yeast protein or mucin and yeast protein. The top axis
depicts the mean protein abundance of the PULs *in vivo* in the yeast diet on the left and in the egg diet on the right. A Welch's t-test
(with FDR control) was performed between each comparison to detect significant changes in PUL protein abundances (*** = q <
0.01, ** = q < 0.05, * = q < 0.1) (Extended Data Tables 18 and 20).

Several of the PULs that increased when we fed mice the yeast diet have previously
been shown to specifically degrade the glycosylations on yeast cell wall proteins. PULs 68/92
(BT3773-3792) and 69 (BT3854-3862) (Fig. 4c; Supplementary Fig. 19) degrade ɑ-mannans
attached to yeast mannoproteins in *Saccharomyces cerevisiae* [76], whereas PUL 56 (BT3310-
3314), degrades yeast β-glucans also attached to yeast cell wall mannoproteins [77].
Conversely, the majority of the PULs increased when we fed mice the egg white diet had been
previously linked to growth on mucin glycan conjugates: PUL14 (BT1032-1051), PUL6 (BT3017-
0318), PUL16 (1280-1285), PUL 80 (BT4295-BT4299), and PUL12 (BT0865-0867)
(Supplementary Fig. 19) [78]. An additional abundant PUL, PUL72 (BT3983-BT3994), has been
previously implicated in the degradation of mannoproteins of mammalian origin [76] and our
result suggests that PUL72 is also involved in the degradation of mannoproteins from non-
mammalian vertebrates.
To test if *B. thetaiotaomicron* could grow on yeast and egg white protein as predicted
from the *in vivo* data, and if the expression of PULs was driven by direct responses to the
dietary protein sources, we characterized *B. thetaiotaomicron* growth and its proteome on
dietary protein sources *in vitro*. We used a defined culture media and supplemented purified
dietary protein sources as the sole carbon source to determine if this supported *B.*
thetaiotaomicron growth. We found that *B. thetaiotaomicron* grew in the presence of glucose
(control), yeast protein, egg white protein, soy protein, and intestinal mucin (Fig. 4d, Tukey HSD
adj $P < 0.05$). We analyzed the proteomes of *B. thetaiotaomicron* in these five different
conditions *in vitro* to determine if PULs played a role in growth (Extended Data Table 19). An
overall comparison of the proteome between the media supplemented with 4 different protein
sources revealed that egg white protein and mucin had the most similar proteome, and the
proteome from the glucose control clustered separately from the protein-sources (Fig. 4e;
Supplementary Fig. 20). We observed that 15 out of 24 PULs that were significantly different in
abundance between the egg white and yeast diets *in vivo* were also significantly different in the
same direction *in vitro* (Fig. 4f). In addition, 12 of these 15 PULs showed the same expression
pattern in both the mucin and egg white protein media as compared to the yeast protein medium
(Fig. 4f, Extended Data Tables 18 and 20). The relationship between mucin and egg white
metabolism in microbiota species *in vivo* is further supported by the fact that five of the six
species with greater than 5% abundance in an egg white sample (*B. thetaiotaomicron*, *A.*
muciniphila, *Atopobiaceae* bacterium AB25_9 *Paramuribaculum sp.*, *D. newyorkensis*) had
abundant enzymes associated with the metabolism of sugars usually thought to be derived from
mucin. These enzymes, of which several were among the top 100 most abundant proteins of

these organisms, catalyze the metabolism of sialic acid (N-acetylneuraminate lyase, N-
acylglucosamine 2-epimerase), N-acetylglucosamine (N-acetylglucosamine-6-phosphate
deacetylase, glucosamine-6-phosphate deaminase, PTS system N-acetylglucosamine-specific,
or fucose (fucosidase, fucose isomerase) (Extended Data Table 6). In summary, these results
indicate that the glycosylations on yeast and egg white proteins drive the increase in abundance
of *B. thetaiotaomicron* in the yeast and egg white diets, and that egg white proteins and
intestinal mucin share similar glycosylations leading to the expression of similar PULs for their
degradation.
669
DISCUSSION
In this study, we sought to characterize how dietary protein source affects the gut
microbiota's composition and function by measuring species-resolved proteins using integrated
metagenomics-metaproteomics. We showed that source of dietary protein significantly alters the
gut microbiota's composition, more so than amount of protein, and that yeast and egg white
protein had the greatest effect on the composition driven by an increase in the relative
abundance of *B. thetaiotaomicron* and a decrease of bacteria from the class *Clostridia*. We also
showed that the source of dietary protein altered the overall functional profile of the gut
microbiota as reflected by changes in the abundance of microbial proteins assigned to broad
functional categories. In particular, proteins involved in amino acid metabolism increased in
abundance in the brown rice and egg white diets, whereas enzymes assigned to glycan
degradation increased in the yeast and egg white diets.
The increase in amino acid metabolizing enzymes in the brown rice and egg white diets
was driven by amino acid degrading enzymes. Previous studies across multiple species have
shown that increasing the amount of protein fed to animals leads to an increase in the ammonia
concentration in stool [79–81], which suggests that increased protein availability leads to
increased amino acid deamination or urease activity in the gut. Here we show that, regardless
of the amount of protein, the source of protein itself can lead to increases in amino acid
deaminating enzymes and ureases from the intestinal microbiota. Gut microbiota urease activity
and amino acid deamination have been linked to serious diseases like hepatic encephalopathy
when liver function is disrupted [82]. Replacement of bacteria that produce these deaminating
enzymes and ureases with bacteria that do not has been suggested as a potential treatment
[83], our results suggest that adjustments in dietary protein source could be considered as well.
Because amino acids are the backbone of protein, we expected to observe changes in
the abundance of amino acid degrading enzymes between the different sources of dietary

protein; however, the effect of dietary protein source on the abundance of glycan degrading
proteins was even greater than the effect on amino acid degrading enzymes. Our results
suggest that the increase in glycan degrading proteins in the yeast and egg white diets is due to
the glycosylations conjugated to these proteins. Yeast and egg white proteins have distinct
glycan conjugate structures [84–86]. In the presence of yeast dietary protein we were able to
show, *in vivo* and *in vitro,* increased expression of PULs associated with the degradation of
yeast mannoprotein glycan conjugates. In the presence of egg white protein, we observed an
increase in PULs previously linked to the degradation of the glycan conjugates of mucin. This
combined with increases in mucin foraging bacteria *Akkermansia muciniphila* [61] and
Paramuribaculum sp. [62] suggests that egg white protein promotes the abundance of mucin
foraging bacteria and their proteins. The link between the foraging of mucin and egg white
protein in retrospect makes sense, as egg white protein contains mucins called ovomucin and
other proteins: ovalbumin, ovotransferrin, and ovomucoid, which have been previously shown to
be N-glycosylated with acetylglucosamine and mannose containing glycans [84, 87]. Previous
studies in mice have shown that diets, which promote bacteria and their enzymes that degrade
mucins, can make the host more susceptible to enteric inflammation and infection [72, 88].
Because egg white protein also promotes these functions, these results suggest that diets high
in egg protein may be detrimental to gastrointestinal health, which could explain the prior results
from population level studies that eggs lead to increased mortality rates among humans [3].
Our study has at least two limitations preventing direct translation of microbiota
responses to dietary protein sources into a human health context. First, we used purified dietary
proteins, which differ from commonly consumed dietary proteins in that regular dietary protein
sources also provide some amount of additional major dietary components such as fats,
carbohydrates, and fiber. For example, plant proteins usually come with a relevant amount of
fiber, whereas animal proteins are often low in fiber and have higher content fats [8]. Second,
we used fully defined diets which allowed us to track effects to specific protein sources, but we
anticipate that the dietary context of protein sources such as co-consumption of multiple protein,
fiber, fat and carbohydrate sources will strongly influence the interactions of dietary protein
sources with gut microbiota. For example, these diets only had cellulose as the fiber source,
which cannot be metabolized by many microbes that are commonly considered fiber degraders.
Thus, the low diversity in fiber sources of these diets likely amplified the effects of yeast and egg
white protein in this study. The power of our study lies in our ability to confirm that the source of
dietary protein does impact gut microbiota function and should be considered when thinking
about how diet impacts gut microbiota and its implications for host health. Future studies that

determine how the effect of dietary protein source on the gut microbiota impacts gastrointestinal
diseases are needed.
731
732   **Declarations**
733
Ethics approval and consent to participate
We conducted our animal experiments in the Laboratory Animal Facilities at the NCSU CVM
campus (Association for the Assessment and Accreditation of Laboratory Animal Care
accredited), which are managed by the NCSU Laboratory Animal Resources. Animals assessed
as moribund were humanely euthanized via CO_2 asphyxiation, as approved by NC State's
Institutional Animal Care and Use Committee (Protocol # 18-034-B).
740
Data availability
The mass spectrometry proteomics data have been deposited to the ProteomeXchange
Consortium via the PRIDE [89] partner repository with the dataset identifier PXD041586
(metaproteomic data) and PXD050296 (*B. thetaiotaomicron* in vitro proteomics data).
Metagenomic raw reads were submitted to NCBI SRA under the bioproject identifier
PRJNA1026909. All metagenome assembled genomes (MAGs) with accompanying metadata
were submitted to DRYAD DOI: 10.5061/dryad.x0k6djhq5. [Reviewer link:
https://datadryad.org/stash/share/QagcDe_b_b0GbbyQ7mPOxBapFL3QbaXt3-fhiZRvDCM]
749

750   **Competing interests**
751   The authors declare that there are no competing interests.
752
753   **Funding**
754   This work was supported by the National Institutes of Health through awards R35GM138362
755   (MK), T32DK007737 (JABR) and P30 DK034987, and by the USDA National Institute of Food
756   and Agriculture, Hatch project 7002782. The content is solely the responsibility of the authors
757   and does not necessarily represent the official views of the National Institutes of Health.
758

Acknowledgments
We thank Brandon C. Iker for inspiring this research and initial concepts, Tjorven Hinzke for
advice on statistics, Heather Maughan for editing of the manuscript, Erin Baker for advice on
methods, and Lawrence David and Balfour Sartor for advice and consultation on the project.

All LC-MS/MS measurements were made in the Molecular Education, Technology, and

Research Innovation Center (METRIC) at North Carolina State University.

Author Contributions

JABR: Experimental design, data collection, data processing, data analysis, author of original

manuscript, editing

AB: Conceptualization of the study, experimental design, data collection, data processing,

editing

ASM: Experimental design, data collection, editing

AA: Data processing, analysis, editing

MVW: Data processing, analysis, editing

AKM: Data processing

JM: Coding, graphic design

SV: Data collection

TR: Data collection

CMT: Experimental design, data collection, editing

MK: Funding, conceptualization of the study, experimental design, data collection, data

processing, data analysis, writing, editing

References

1. Budhathoki S, Sawada N, Iwasaki M, Yamaji T, Goto A, Kotemori A, et al. Association of

Animal and Plant Protein Intake With All-Cause and Cause-Specific Mortality in a

Japanese Cohort. *JAMA Intern Med* 2019; **179**: 1509–1518.

2. Song M, Fung TT, Hu FB, Willett WC, Longo VD, Chan AT, et al. Association of Animal

and Plant Protein Intake With All-Cause and Cause-Specific Mortality. *JAMA Intern Med*

2016; **176**: 1453–1463.

3. Huang J, Liao LM, Weinstein SJ, Sinha R, Graubard BI, Albanes D. Association Between

Plant and Animal Protein Intake and Overall and Cause-Specific Mortality. *JAMA Intern*

Med 2020; **180**: 1173–1184.

4. Ahn E, Jeong H, Kim E. Differential effects of various dietary proteins on dextran sulfate

sodium-induced colitis in mice. *Nutr Res Pr* 2022; **16**: 700–715.

5.   Bartlett A, Kleiner M. Dietary protein and the intestinal microbiota: An understudied

relationship. *iScience* 2022; **25**: 105313.

6.   Oliphant K, Allen-Vercoe E. Macronutrient metabolism by the human gut microbiome:

major fermentation by-products and their impact on host health. *Microbiome* 2019; **7**: 91.

7.   Blakeley-Ruiz JA, McClintock CS, Shrestha HK, Poudel S, Yang ZK, Giannone RJ, et al.

Morphine and high-fat diet differentially alter the gut microbiota composition and metabolic

function in lean versus obese mice. *ISME Commun* 2022; **2**: 66.

8.   David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, et al. Diet

rapidly and reproducibly alters the human gut microbiome. *Nature* 2014; **505**: 559–563.

9.   Patnode ML, Beller ZW, Han ND, Cheng J, Peters SL, Terrapon N, et al. Interspecies

Competition Impacts Targeted Manipulation of Human Gut Bacteria by Fiber-Derived

Glycans. *Cell* 2019; **179**: 59-73.e13.

10.   Perler BK, Friedman ES, Wu GD. The Role of the Gut Microbiota in the Relationship

Between Diet and Human Health. *Annu Rev Physiol* 2023; **85**: 449–468.

11.   Yao CK, Muir JG, Gibson PR. Review article: insights into colonic protein fermentation, its

modulation and potential health implications. *Aliment Pharmacol Ther* 2016; **43**: 181–196.

12.   Sun M, Wu W, Chen L, Yang W, Huang X, Ma C, et al. Microbiota-derived short-chain fatty

acids promote Th1 cell IL-10 production to maintain intestinal homeostasis. *Nat Commun*

2018; **9**: 3555.

13.   Faith JJ, McNulty NP, Rey FE, Gordon JI. Predicting a Human Gut Microbiota's Response

to Diet in Gnotobiotic Mice. *Science* 2011; **333**: 101–104.

14.   Zhu Y, Shi X, Lin X, Ye K, Xu X, Li C, et al. Beef, Chicken, and Soy Proteins in Diets

Induce Different Gut Microbiota and Metabolites in Rats. *Front Microbiol* 2017; **8**: 1395.

15.   Wilmes P, Bond PL. The application of two-dimensional polyacrylamide gel electrophoresis

and downstream analyses to a mixed community of prokaryotic microorganisms. *Environ*

Microbiol 2004; **6**: 911–920.

16. Kleiner M, Thorson E, Sharp CE, Dong X, Liu D, Li C, et al. Assessing species biomass

contributions in microbial communities via metaproteomics. *Nat Commun* 2017; **8**: 1558.

17. Mueller RS, Denef VJ, Kalnejais LH, Suttle KB, Thomas BC, Wilmes P, et al. Ecological

distribution and population physiology defined by proteomics in a natural microbial

community. *Mol Syst Biol* 2010; **6**: 374.

18. Kleiner M, Wentrup C, Lott C, Teeling H, Wetzel S, Young J, et al. Metaproteomics of a

gutless marine worm and its symbiotic microbial community reveal unusual pathways for

carbon and energy use. *Proc Natl Acad Sci* 2012; **109**: E1173.

19. Blakeley-Ruiz JA, Erickson AR, Cantarel BL, Xiong W, Adams R, Jansson JK, et al.

Metaproteomics reveals persistent and phylum-redundant metabolic functional stability in

adult human gut microbiomes of Crohn's remission patients despite temporal variations in

microbial taxa, genomes, and proteomes. *Microbiome* 2019; **7**: 18.

20. Salvato F, Hettich RL, Kleiner M. Five key aspects of metaproteomics as a tool to

understand functional interactions in host-associated microbiomes. *PLOS Pathog* 2021;

17: e1009245.

21. Blakeley-Ruiz JA, Kleiner M. Considerations for constructing a protein sequence database

for metaproteomics. *Comput Struct Biotechnol J* 2022; **20**: 937–952.

22. Brooks B, Mueller R, Young J, Morowitz M, Hettich R, Banfield J. Strain-resolved microbial

community proteomics reveals simultaneous aerobic and anaerobic function during

gastrointestinal tract colonization of a preterm infant. *Front Microbiol* 2015; **6**: 654.

23. Xiong W, Brown CT, Morowitz MJ, Banfield JF, Hettich RL. Genome-resolved

metaproteomic characterization of preterm infant gut microbiota development reveals

species-specific metabolic shifts and variabilities during early life. *Microbiome* 2017; **5**: 72.

24. Reese AT, Pereira FC, Schintlmeister A, Berry D, Wagner M, Hale LP, et al. Microbial

nitrogen limitation in the mammalian large intestine. *Nat Microbiol* 2018; **3**: 1441–1450.

25. Mordant A, Kleiner M. Evaluation of Sample Preservation and Storage Methods for

Metaproteomics Analysis of Intestinal Microbiomes. *Microbiol Spectr* 2021; **9**: e0187721.

26. Knudsen BE, Bergmark L, Munk P, Lukjancenko O, Priemé A, Aarestrup FM, et al. Impact

of Sample Type and DNA Isolation Procedure on Genomic Inference of Microbiome

Composition. *mSystems* 2016; **1**: 10.1128/msystems.00095-16.

27. Bushnell B. BBMap: a fast, accurate, splice-aware aligner. 2014. Lawrence Berkeley

National Lab.(LBNL), Berkeley, CA (United States).

(https://sourceforge.net/projects/bbmap/).

28. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. metaSPAdes: a new versatile

metagenomic assembler. *Genome Res* 2017; **27**: 824–834.

29. Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. MEGAHIT: an ultra-fast single-node solution

for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics*

2015; **31**: 1674–1676.

30. Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, et al. MetaBAT 2: an adaptive binning

algorithm for robust and efficient genome reconstruction from metagenome assemblies.

PeerJ 2019; **7**: e7359–e7359.

31. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the

quality of microbial genomes recovered from isolates, single cells, and metagenomes.

Genome Res 2015; **25**: 1043–1055.

32. Bowers RM, Kyrpides NC, Stepanauskas R, Harmon-Smith M, Doud D, Reddy TBK, et al.

Minimum information about a single amplified genome (MISAG) and a metagenome-

assembled genome (MIMAG) of bacteria and archaea. *Nat Biotechnol* 2017; **35**: 725–731.

33. Olm MR, Brown CT, Brooks B, Banfield JF. dRep: a tool for fast and accurate genomic

comparisons that enables improved genome recovery from metagenomes through de-

replication. *ISME J* 2017; **11**: 2864–2868.

34. Olm MR, Crits-Christoph A, Diamond S, Lavy A, Matheus Carnevali PB, Banfield JF.

Consistent Metagenome-Derived Metrics Verify and Delineate Bacterial Species

Boundaries. *mSystems* 2020; **5**.

35. Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify

genomes with the Genome Taxonomy Database. *Bioinformatics* 2020; **36**: 1925–1927.

36. Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinforma Oxf Engl* 2014; **30**:

2068–2069.

37. Li W, Godzik A. Cd-hit: a fast program for clustering and comparing large sets of protein or

nucleotide sequences. *Bioinformatics* 2006; **22**: 1658–1659.

38. The UniProt Consortium. UniProt: the Universal Protein Knowledgebase in 2023. *Nucleic*

Acids Res 2023; **51**: D523–D531.

39. Wiśniewski JR, Zougman A, Nagaraj N, Mann M. Universal sample preparation method for

proteome analysis. *Nat Methods* 2009; **6**: 359–362.

40. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using lme4. *J*

Stat Softw 2015; **67**: 1–48.

41. Lenth RV. emmeans: Estimated Marginal Means, aka Least-Squares Means. 2023.

42. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful

Approach to Multiple Testing. *J R Stat Soc Ser B Methodol* 1995; **57**: 289–300.

43. Storey JD, Taylor JE, Siegmund D. Strong Control, Conservative Point Estimation and

Simultaneous Conservative Consistency of False Discovery Rates: A Unified Approach. *J*

R Stat Soc Ser B Stat Methodol 2004; **66**: 187–205.

44. Wickham H. ggplot2: Elegant Graphics for Data Analysis. 2016. Springer-Verlag New York.

45. Kolde R. pheatmap: Pretty Heatmaps. 2019. (https://CRAN.R-

project.org/package=pheatmap).

46.   Mauri M, Elli T, Caviglia G, Uboldi G, Azzi M. RAWGraphs: A Visualisation Platform to

Create Open Outputs. *Proc. 12th Biannu. Conf. Ital. SIGCHI Chapter.* 2017. Association

for Computing Machinery, New York, NY, USA.

47.   R Core Team. R: A Language and Environment for Statistical Computing. 2023. R

Foundation for Statistical Computing, Vienna, Austria. (https://www.R-project.org).

48.   Oksanen J, Simpson GL, Blanchet FG, Kindt R, Legendre P, Minchin PR, et al. vegan:

Community Ecology Package. 2022. (https://CRAN.R-project.org/package=vegan).

49.   Florens L, Carozza MJ, Swanson SK, Fournier M, Coleman MK, Workman JL, et al.

Analyzing chromatin remodeling complexes using shotgun proteomics and normalized

spectral abundance factors. *Methods San Diego Calif* 2006; **40**: 303–311.

50.   Huerta-Cepas J, Forslund K, Coelho LP, Szklarczyk D, Jensen LJ, von Mering C, et al.

Fast Genome-Wide Functional Annotation through Orthology Assignment by eggNOG-

Mapper. *Mol Biol Evol* 2017; **34**: 2115–2122.

51.   Queirós P, Delogu F, Hickl O, May P, Wilmes P. Mantis: flexible and consensus-driven

genome annotation. *GigaScience* 2021; **10**: giab042.

52.   Ruiz-Perez CA, Conrad RE, Konstantinidis KT. MicrobeAnnotator: a user-friendly,

comprehensive functional annotation pipeline for microbial genomes. *BMC Bioinformatics*

2021; **22**: 11.

53.   Lombard V, Golaconda Ramulu H, Drula E, Coutinho PM, Henrissat B. The carbohydrate-

active enzymes database (CAZy) in 2013. *Nucleic Acids Res* 2014; **42**: D490–D495.

54.   Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, et al. dbCAN2: a meta server for

automated carbohydrate-active enzyme annotation. *Nucleic Acids Res* 2018; **46**: W95–

W101.

55.   Blum M, Chang H-Y, Chuguransky S, Grego T, Kandasaamy S, Mitchell A, et al. The

InterPro protein families and domains database: 20 years on. *Nucleic Acids Res* 2021; **49**:

D344–D354.

56.   Tyanova S, Temu T, Sinitcyn P, Carlson A, Hein MY, Geiger T, et al. The Perseus

computational platform for comprehensive analysis of (prote)omics data. *Nat Methods*

2016; **13**: 731–740.

57.   Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, et al. The RAST Server:

Rapid Annotations using Subsystems Technology. *BMC Genomics* 2008; **9**: 75.

58.   Terrapon N, Lombard V, Drula É, Lapébie P, Al-Masaudi S, Gilbert HJ, et al. PULDB: the

expanded database of Polysaccharide Utilization Loci. *Nucleic Acids Res* 2018; **46**: D677–

D683.

59.   Rogowski A, Briggs JA, Mortimer JC, Tryfona T, Terrapon N, Lowe EC, et al. Glycan

complexity dictates microbial resource allocation in the large intestine. *Nat Commun* 2015;

6: 7481.

60.   Parks DH, Rinke C, Chuvochina M, Chaumeil P-A, Woodcroft BJ, Evans PN, et al.

Recovery of nearly 8,000 metagenome-assembled genomes substantially expands the

tree of life. *Nat Microbiol* 2017; **2**: 1533–1542.

61.   Derrien M, Vaughan EE, Plugge CM, de Vos WM. Akkermansia muciniphila gen. nov., sp.

nov., a human intestinal mucin-degrading bacterium. *Int J Syst Evol Microbiol* . 2004.

Microbiology Society. , **54**: 1469–1476

62.   Lagkouvardos I, Lesker TR, Hitch TCA, Gálvez EJC, Smit N, Neuhaus K, et al. Sequence

and cultivation study of Muribaculaceae reveals novel species, host preference, and

functional potential of this yet undescribed family. *Microbiome* 2019; **7**: 28.

63.   Sonnenburg JL, Xu J, Leip DD, Chen C-H, Westover BP, Weatherford J, et al. Glycan

Foraging in Vivo by an Intestine-Adapted Bacterial Symbiont. *Science* 2005; **307**: 1955–

1959.

64.   Kleiner M. Metaproteomics: Much More than Measuring Gene Expression in Microbial

Communities. *mSystems* 2019; **4**: e00115-19.

65.   O'Keefe SJD. Diet, microorganisms and their metabolites, and colon cancer. *Nat Rev*

946   *Gastroenterol Hepatol* 2016; **13**: 691–706.

947   66.   Cagnon L, Braissant O. Hyperammonemia-induced toxicity for the developing central

948   nervous system. *Brain Res Rev* 2007; **56**: 183–197.

949   67.   Stummer N, Feichtinger RG, Weghuber D, Kofler B, Schneider AM. Role of Hydrogen

950   Sulfide in Inflammatory Bowel Disease. *Antioxidants* 2023; **12**.

951   68.   Rangroo Thrane V, Thrane AS, Wang F, Cotrina ML, Smith NA, Chen M, et al. Ammonia

952   triggers neuronal disinhibition and seizures by impairing astrocyte potassium buffering. *Nat*

953   *Med* 2013; **19**: 1643–1648.

954   69.   Jaglin M, Rhimi M, Philippe C, Pons N, Bruneau A, Goustard B, et al. Indole, a Signaling

955   Molecule Produced by the Gut Microbiota, Negatively Impacts Emotional Behaviors in

956   Rats. *Front Neurosci* 2018; **12**.

957   70.   Luscher B, Shen Q, Sahir N. The GABAergic deficit hypothesis of major depressive

958   disorder. *Mol Psychiatry* 2011; **16**: 383–406.

959   71.   Czumaj A, Śledziński T, Mika A. Branched-Chain Fatty Acids Alter the Expression of

960   Genes Responsible for Lipid Synthesis and Inflammation in Human Adipose Cells.

961   *Nutrients* 2022; **14**.

962   72.   Pereira GV, Boudaud M, Wolter M, Alexander C, De Sciscio A, Grant ET, et al. Opposing

963   diet, microbiome, and metabolite mechanisms regulate inflammatory bowel disease in a

964   genetically susceptible host. *Cell Host Microbe* 2024.

965   73.   Mayneris-Perxachs J, Castells-Nobau A, Arnoriaga-Rodríguez M, Martin M, de la Vega-

966   Correa L, Zapata C, et al. Microbiota alterations in proline metabolism impact depression.

967   *Cell Metab* 2022; **34**: 681-701.e10.

968   74.   Fletcher JR, Pike CM, Parsons RJ, Rivera AJ, Foley MH, McLaren MR, et al. Clostridioides

969   difficile exploits toxin-mediated inflammation to alter the host nutritional landscape and

970   exclude competitors from the gut microbiota. *Nat Commun* 2021; **12**: 462.

75. Grondin JM, Tamura K, Déjean G, Wade DW, Brumer H. Polysaccharide Utilization Loci:

Fueling Microbial Communities. *J Bacteriol* 2017; **199**: 10.1128/jb.00860-16.

76. Cuskin F, Lowe EC, Temple MJ, Zhu Y, Cameron E, Pudlo NA, et al. Human gut

Bacteroidetes can utilize yeast mannan through a selfish mechanism. *Nature* 2015; **517**:

165–169.

77. Temple MJ, Cuskin F, Baslé A, Hickey N, Speciale G, Williams SJ, et al. A Bacteroidetes

locus dedicated to fungal 1,6-β-glucan degradation: Unique substrate conformation drives

specificity of the key endo-1,6-β-glucanase. *J Biol Chem* 2017; **292**: 10639–10650.

78. Martens EC, Chiang HC, Gordon JI. Mucosal glycan foraging enhances fitness and

transmission of a saccharolytic human gut bacterial symbiont. *Cell Host Microbe* 2008; **4**:

447–457.

79. Badri DV, Jackson MI, Jewell DE. Dietary Protein and Carbohydrate Levels Affect the Gut

Microbiota and Clinical Assessment in Healthy Adult Cats. *J Nutr* 2021; **151**: 3637–3650.

80. Cummings J, Hill M, Bone E, Branch W, Jenkins DJA. The effect of meat protein and

dietary fiber on colonic function and metabolism II. Bacterial metabolites in feces and

urine1. *Am J Clin Nutr* 1979; **32**: 2094–2101.

81. Pinna C, Vecchiato CG, Bolduan C, Grandi M, Stefanelli C, Windisch W, et al. Influence of

dietary protein and fructooligosaccharides on fecal fermentative end-products, fecal

bacterial populations and apparent total tract digestibility in dogs. *BMC Vet Res* 2018; **14**:

106.

82. Rose CF. Ammonia-Lowering Strategies for the Treatment of Hepatic Encephalopathy.

Clin Pharmacol Ther 2012; **92**: 321–331.

83. Shen T-CD, Albenberg L, Bittinger K, Chehoud C, Chen Y-Y, Judge CA, et al. Engineering

the gut microbiota to treat hyperammonemia. *J Clin Invest* 2015; **125**: 2841–2850.

84. Hwang HS, Kim BS, Park H, Park H-Y, Choi H-D, Kim HH. Type and branched pattern of

N-glycans and their structural effect on the chicken egg allergen ovotransferrin: a

comparison with ovomucoid. *Glycoconj J* 2014; **31**: 41–50.

85. Kollár R, Reinhold BB, Petráková E, Yeh HJC, Ashwell G, Drgonová J, et al. Architecture

of the Yeast Cell Wall: β(1→6)-GLUCAN INTERCONNECTS MANNOPROTEIN, β(1→3)-

GLUCAN, AND CHITIN*. *J Biol Chem* 1997; **272**: 17762–17775.

86. Orlean P. Architecture and Biosynthesis of the Saccharomyces cerevisiae Cell Wall.

Genetics 2012; **192**: 775–818.

87. Thaysen-Andersen M, Mysling S, Højrup P. Site-Specific Glycoprofiling of N-Linked

Glycopeptides Using MALDI-TOF MS: Strong Correlation between Signal Strength and

Glycoform Quantities. *Anal Chem* 2009; **81**: 3933–3943.

88. Desai MS, Seekatz AM, Koropatkin NM, Kamada N, Hickey CA, Wolter M, et al. A Dietary

Fiber-Deprived Gut Microbiota Degrades the Colonic Mucus Barrier and Enhances

Pathogen Susceptibility. *Cell* 2016; **167**: 1339-1353.e21.

89. Perez-Riverol Y, Bai J, Bandla C, García-Seisdedos D, Hewapathirana S, Kamatchinathan

S, et al. The PRIDE database resources in 2022: a hub for mass spectrometry-based

proteomics evidences. *Nucleic Acids Res* 2022; **50**: D543–D552.

1012