**Multiplexed transcriptomic analyses of the plant embryonic hourglass**

Hao Wu[1,*,#], Ruqiang Zhang[1,#], Karl J. Niklas[1], Michael J. Scanlon[1,*]

[1]School of Integrative Plant Science, Plant Biology Section, Cornell University, Ithaca, NY,

USA

[#]These authors contributed equally

[*]Co-corresponding authors

**Zoologists have adduced morphological convergence among embryonic stages of  closely**

**related taxa, which has been called the phylotypic stage of embryogenesis. Transcriptomic**

**analyses reveal a phylotypic hourglass pattern of gene expression during plant as well as**

**animal embryogenesis, characterized by the accumulation of evolutionarily older and**

**conserved transcripts during mid-embryogenesis, whereas younger less-conserved**

**transcripts predominate at earlier and later embryonic stages.  However, widespread**

**comparisons of embryonic gene expression across animal phyla describe an inverse**

**hourglass pattern, where gene expression is correlated during early and late stages but not**

**during mid-development.  Here, multiplexed spatial-transcriptomic analyses is used to**

**investigate embryo ontogeny and homology in maize, which has novel, grass-specific**

**morphology. An embryonic-organ genetic network is identified, replete for**

**ancient/conserved genes manifesting a phylotypic hourglass during mid-embryogenesis.**

**Transcriptomic comparisons of grass embryo ontogeny with that of a moss *Physcomitrium***

***patens* identify a conserved, inverse hourglass pattern across plant phyla, as in animals .**

**The data indicate that the plant phylotypic stage and that of animals is characterized by**

**ancient gene network expression during early histo- and morphogenesis and polarized**

**embryonic development.  The data reveal an ancient, convergent mechanism for the**

**evolution of morphological novelty.**

The embryo is defined as the developmental stage spanning the period between fertilization and

the appearance of primary tissues and organs. Post-fertilization, flowering plant embryogenesis

consists of three conserved processes: (1) polarized and asymmetric cell divisions to create the

embryo proper, (2) patterning of discrete histological tissue-layers, and (3) organogenesis of

lateral organs and the formation of apical meristems. Persistent meristematic reservoirs

comprising the shoot apical meristem (SAM) and root apical meristem (RAM) enable vascular

plants to extend organogenesis beyond embryogenesis. The gene regulatory networks regulating

embryogenesis, however, are largely undescribed.

The embryos of the eudicot *Arabidopsis thaliana* and the monocot *Zea mays* ssp. *mays* involve

asymmetric cell divisions that establish the embryo proper and the basal, haustorial suspensor

that supplies nutrients to the embryo[1-3] (Fig. 1a). Subsequent histological layering creates an

outer epidermal layer surrounding internal tissues, and *Arabidopsis* embryos develop two

embryonic leaves (cotyledons) and shoot and root apical meristems, whereupon organogenesis is

suspended until seed germination[1, 4]. The grass embryo differs from that of *Arabidopsis*[2, 3, 5-9] in

several important respects. The first lateral organ formed in the maize embryo is the scutellum

(Fig. 1b), which at maturity forms a shield-shaped, haustorial organ that absorbs nutrients from

the endosperm[7, 10]. The coleoptile then emerges (Fig. 1c) to form a tubular-like sheath that

protects the shoot during germination. Unlike *Arabidopsis*, grasses extend embryo organogenesis

to develop several foliar leaves contained within the coleoptile before seed quiescence.

In contrast, the embryos of the moss *Physcomitrium patens* have no apical meristems *sensu*

*stricto* or lateral organs. After fertilization, a transverse zygotic cell division ultimately gives rise

to a basal "foot" and a cell mass that develops into the seta and the spore capsule [11, 12.] The latter

consists of a histologically layered sporangium, wherein meiosis results in thousands of haploid

spores. Fundamental differences in embryonic growth polarity are also noted between non-

vascular and vascular plants, e.g., non-vascular plant embryos develop exoscopically outward

from the subtending gametophyte, whereas vascular plant embryos develop endoscopically into

the gametophyte[13]. Indeed, embryo morphometric  diversity is widespread across the land plants,

although comparative transcriptomic analyses of embryogenesis are few[10, 14, 15].

The diversity of land plant embryo development stands in contrast with that of animals, which

appears to converge on to a period of maximum embryo morphological similarity among related

vertebrate taxa[16-18] during "mid-embryogenesis" in contrast to "earlier" and "later" stages that

are dissimilar morphologically. The result is an hourglass pattern. The convergent mid-embryo

stage has been called the phylotypic stage[19].  Recent transcriptomic analyses of embryogenesis,

first performed in fruit flies (*Drosophila*), the zebrafish (*Danio*), and later *Arabidopsis*, have

identified a phylotypic stage of gene expression featuring an enhanced accumulation of more

ancient and conserved transcripts during mid-embryogenesis as compared to earlier and later

ontogenetic stages[20-22].  Plots of transcriptomic gene age and divergence conform to the

"hourglass model" first described in animals[16-18].

However, more widespread comparisons of ontogenetic gene expression across ten diverse

animal phyla identify an inverse hourglass pattern, wherein the most prominent transcripts

during early and later stages of development are more conserved among phyla as compared to

genes expressed during the presumptive "phylotypic stage"[23].

An important issue in all embryological comparisons across diverse taxa involves

temporal homology — what is meant by early-, mid-, and late-embryogenesis? This issue has

added relevance when comparing metazoans and land plants, each of which is adduced to be

monophyletic yet morphologically extremely diverse. This concern applies even to broad

comparisons among land plant embryos, which manifest what appear to be three, distinct

patterns of embryogenesis: the nonvascular bryophytes, the vascular seedless pteridophytes, and

the seed plants[13]. Yet, another concern in this field of enquiry is the temporal delay between gene

expression and embryological-morphogenetic response, which likely differs among genes within

an individual organisms and among phylogenetically diverse taxa. Neither of these concerns has

been adequately address, and, as yet a transcriptomic phylotypic analysis is reported for only a

single land plant genus (*Arabidopsis*) [22]. No such study has been reported across the divergent

land plant lineages.

Here we present multiplexed analyses of gene expression during maize embryogenesis, a grass

that evolved more than 150 million years after the appearance of flowering plants[24].  Single cell

sequencing (scRNA-seq), spatial RNA-seq, laser-microdissection RNA-seq (LM RNAseq), and

multiplexed RNA-targeting *in situ* hybridizations were utilized to characterize gene expression

networks in developing maize embryos at high resolution.  An embryonic-organ-initiation

genetic network is identified for all lateral organs of the Stage 1 maize embryo, attesting to their

collective homology. This embryonic gene network is conserved in *Arabidopsis* and comprises

ancient and shared genes expressed in a phylotypic hourglass pattern at mid-embryo stages.

Across-phylum comparisons of embryonic gene expression in the flowering plants maize and

*Arabidopsis*, to that of a nonvascular moss, reveals inverse hourglass patterns equivalent to those

described in animals. The data indicate that the plant and animal phylotypic stage reveals the

ancestral morphometric processes of tissue histogenesis and meristematic growth of embryonic

organs. An across plant-animal kingdom model for the evolution of morphological novelty via

the innovative expression of newer-evolved, sequence-conserved gene networks is presented.

**Results**

**Multiplexed and single-cell profiling of the transcriptomic networks in the Stage 1 maize**

**embryo.**

A combinatorial sc-RNAseq approach was utilized to identify the gene networks during maize

embryogenesis and investigate phylotypic trends and homology of the grass embryo[25, 26]. Stage

1 maize embryos are approximately 1 mm long and contain a prominent, ovately-flattened

scutellum, an emerged coleoptile that is beginning to surround the SAM, and the newly-initiated

primordium of the first foliar leaf[5] (Fig. 1d). Stage 1 embryos contain all the lateral organ types

of the maize embryonic shoot and thus were an appropriate stage to analyze the organ-specific

genetic networks among grass embryonic organs.

10X Genomics Chromium™ was used to profile the single-cell transcriptomes of 11,467

protoplasts digested from over 400 Stage 1 embryos in two replicates (Fig. 1e). Twenty-three cell

clusters were identified; median values of 10,594 UMI per cell and 4,030 genes per cell were

obtained.

The first challenge encountered in analyses of the sc-RNAseq data was the identification of the

tissue/organ-specific origins of each cell-cluster. Prior RNA-seq analyses in maize have focused

on seedlings or inflorescences[27], such that cell-clusters derived from the SAM, embryonic stem,

and vasculature[28] were readily identified, as was the residual starchy endosperm (Fig. 1e and

Clusters 6, 15 and 20 respectively). In contrast, relatively few transcriptomic analyses have

centered on the grass scutellum or coleoptile, such that the origins of several Stage 1 embryonic

cell-clusters were not immediately identifiable. To annotate all the UMAP cell-clusters, a

multiplexed transcriptomic approach was used combining the sc-RNAseq data with spatial-

transcriptomics and laser-microdissection RNA-sequencing (LM-RNAseq) to determine spatial

information for all Stage 1 embryo cell-clusters, as described below (Fig. 2 and  Supplementary

Fig. 1-2, Supplementary Table 1, 2).

To validate the identification of Stage 1 embryo UMAP single cell clusters, eight individual

Stage 1 embryo sections were processed using the 10X Genomics Visium[TM] pipeline for spatial

transcriptomics (Fig. 2a-d and Supplementary Table 3). Nine spatially resolved cell-clusters were

identified and categorized into four supergroups: (1) shoot-root axis, (2) scutellum epidermis, (3)

internal scutellum, and (4) endosperm/embryo boundary (Fig. 2e). Hypergeometric testing was

used to analyze the overlap of cluster marker genes between the single-cell and spatial-

transcriptomic data, revealing that 14 of the 23 single-cell clusters (0-5, 7-9, 11, 13, 14, 18 and

19) are derived from the scutellum (Fig. 2h). Comparisons of transcript localizations of known

marker genes within the UMAP and the spatial-transcriptomic plots indicated that the resolution

of Visium[TM] technology comprises 5-10 embryonic cells. Thus, it was not possible to spatially-

distinguish transcripts specifically expressed in the diminutive coleoptile, SAM, or leaf 1. To

increase transcriptomic resolution, LM-RNAseq used to microdissect gene expression in the

SAM, adaxial scutellum, upper/lower coleoptile, and leaf primordium of Stage 1 embryos (Fig.

2f-g). Principal component analyses (PCA) showed that the transcriptomic profiles of these five,

maize embryonic organ/tissues are distinct (Supplementary Fig. 2a). In particular, 9,913 genes

were identified as differentially-expressed in at least one embryonic structure, and that 89, 1058,

246, 84, and 53 genes were preferentially expressed in these same five embryonic organ/tissues

(Supplementary Fig. 2b-f and Supplementary Table 4-6). Thus, the multiplexed transcriptomic

approach enabled the identification of the tissue/organ origin of each cell-cluster in the Stage 1

maize embryonic UMAP (Fig. 1e, 2i).

**Spatiotemporal analyses of embryo ontogeny and homology via multiplexed RNA-targeting**

Homologous organs are thought to employ shared gene networks regulating developmental

patterning[25, 29]. With an emphasis on regulatory genes differentially expressed in our

transcriptomic analyses of Stage 1 maize embryos, a total100 transcripts were targeted for

spatiotemporal analyses of multiplexed, *in-situ* hybridization of transcript localization during

embryo ontogeny, using the 10X Xenium[TM] protocols (Supplementary Table 7). Samples

included longitudinal sections of (i) transition-staged embryos just forming a SAM and scutellum

(Fig. 1b), (ii) coleoptilar-staged embryos (Fig. 1c), and (iii) Stage 1 embryos initiating the first

foliar leaf (Fig. 1d). Correlations of the sc-RNAseq data, spatial-transcriptomic plots, and

Xenium[TM] *in situ* RNA-targeting data were examined via comparisons of 16 genes preferentially

expressed in one-or-two single-cell UMAP clusters (Supplementary Fig. 1). In all cases, the *in*

*situ* hybridization patterns were consistent with the spatial expression data.

In light of the comparative consistencies of the multiplexed *in situ* hybridization data, the

dynamic co-localization patterns of some key regulators of maize shoot development during

serial stages in grass embryo ontogeny were examined. Consistent with prior studies[30, 31], the

spatial-temporal expression of the indeterminacy marker *KNOTTED1* (*KN1*; Fig. 3a-d) was first

detected during SAM initiation at the maize Transition stage (Fig. 1b). At later stages, *KN1* was

observed to accumulate in the SAM and subtending embryonic stem, but was downregulated in

lateral organs[30, 31]. *NARROW SHEATH1* (*NS1*) and *WUSCHEL-RELATED HOMEOBOX 3A*

(*WOX3A*), which encode homeodomain transcription factors required for mediolateral expansion

of maize lateral organs[29], were co-expressed at the edges of the developing coleoptile and

initiating leaf primordia as described previously[7, 25, 32, 33] (Fig. 3b-c, f-g). Undetected in prior

studies of grass embryo ontogeny, Fig. 3a reveals that *NS1* transcripts accumulate in the

initiating scutellum at the Transition stage.  Moreover, transcripts of *CUP-SHAPED*

*COTYLEDON2* (*CUC2*), which mark developmental boundaries throughout maize ontogeny[34],

were detected between the expression domains of *KN1* and *NS1* during scutellum initiation (Fig.

3a), and in the boundaries between the SAM and later-staged embryonic lateral organs (Fig. 3b-

c,e).  *DROOPING LEAF1* (*DRL1*), which regulates leaf architecture and is expressed in incipient

and developing foliar leaf primordia[35], was not detected in the coleoptile or scutellum at any

embryo stage examined (Fig. 3c,h). These combinatorial *in situ* RNA-targeting analyses enabled

the identification of previously undescribed combinatorial patterns of dynamic, gene expression

during maize embryo development.

Other candidate genes examined using Xenium[TM] displayed patterns of transcript accumulation

that offer insights into the developmental homologies of maize embryonic lateral organs. For

example, *BLADE-ON-PETIOLEa* (*BOPa*) accumulated throughout the developing coleoptile and

in the proximal region of the initiating leaf at Stage 1 (Fig. 3i-k), consistent with the proposed

function of *BOP* homologs as leaf proximal-identity genes[36]. Notably, *BOPa* was undetected

throughout scutellum ontogeny, in a manner similar to the accumulation patterns noted for the

*BOPa* paralog *TASSELS REPLACE UPPER EARS1* (*TRU1*) (Supplementary Fig. 3). Transcripts

of the *DRL1*-paralog *YABBY14* (*YAB14*) densely accumulated in the apical tip of the scutellum at

the transition stage and in the elongating scutellum and newly emerged coleoptile at the

coleoptilar stage. During Stage 1, a new *YAB14* expression maximum was organized at the

primordial leaf tip (Fig. 3l-n). Moreover, the pleiotropic gene *INDETERMINATE*

*GAMETOPHYTE1* (*IG1*), reported to function during leaf polarity and patterning of maize

leaves[37, 38], accumulated on the dorsal (i.e. adaxial) surfaces of initiating leaf primordia during

Stage 1, and also in the adaxial layers of the emerging coleoptile and in the upper-adaxial region

of the transition-staged scutellum (Fig. 3o-t).  No significant *IG1* expression was observed

during later stages of embryonic scutellum development (Fig. 3p-q, s-t). Another leaf polarity

regulator, *AUXIN RESPONSE FACTOR 3* (*ARF3*)[39], was expressed on the abaxial region of the

transition-staged embryo during scutellum initiation, and in the abaxial domains of the newly-

emerged coleoptile and leaf during later embryo stages (Fig. 3u-z). Interestingly, the expression

of *ARF3* in the scutellum moved from the ventral (i.e., abaxial) region in the transition stage to

adaxial region at Stage 1 (Fig. 3z).

**The initiation of maize embryonic lateral organs utilizes a shared genetic network**

By Stage 1, the three grass embryonic lateral organs initiate morphological divergence and

continue to do so before seed quiescence[5]. Therefore, the determination of organ homology and

of a transcriptomic phylotypic stage in maize requires the examination of genetic networks

throughout embryo ontogeny[26]. Toward this end, the LM-RNAseq data from Stage 1 embryonic

organs were combined with the LM-RNAseq data from serial stages of maize embryonic

development[28] to generate a more comprehensive sampling throughout embryogenesis.

Specifically, these prior analyses generated transcriptomic data precisely from the newly initiated

embryonic organ and the adjacent SAM, which were serially sampled at the transition stage

(scutellum +SAM), the coleoptilar stage (coleoptile +SAM) and Stage 1 (leaf 1 +SAM). Fig. 2i

shows that the expression levels of several leaf developmental genes were highest in Stage 1

leaves (L) and steadily decreased in the lower coleoptile (LC), upper coleoptile (UC), and

scutellum (S). Notably, the expression of these genes during the transition stage and the

coleoptilar stage was similar to that of leaf 1 during Stage 1 (Supplementary Table 4)[28]. These

data provide evidence that the gene networks involved in foliar leaf initiation at Stage 1 are also

associated with the initiation of the scutellum and coleoptile at successively earlier stages in

embryo ontogeny.

To further investigate this hypothesis, a weighted gene co-expression network analysis

(WGCNA)[40, 41] was performed on the Stage 1 embryonic LM-RNAseq data to identify co-

expression modules correlated with the expression pattern $L > LC > UC > S$. A total of 5,797

gene transcripts with high expression variability were selected. With the exception of one non-

clustered transcript module (MEgrey module), these were grouped into twelve co-expression

modules (ME1 through ME12). Two co-expression modules (ME2 and ME3) comprising 963

genes emerged as correlated with the L > LC > UC > S  pattern. (Supplementary Fig. 4a and

Supplementary Table 8). Intersection of these 963 genes with transcripts identified by LM-

RNAseq in the initiating scutellum, coleoptile and leaf 1[28] (excluding putative SAM-specific

genes) identified 130 genes comprising the maize embryonic-organ-initiation gene network (Fig.

4a and Supplementary Fig. 4b and Supplementary Table 9). Expression of this gene network was

significantly higher during initiation of these embryonic organs as compared to later

morphometric stages (Fig. 4b, Supplementary Fig. 4b-e).

In light of the controversy regarding the homology of the grass cotyledon (i.e., whether the

coleoptile is a foliar leaf[8], the entire cotyledon[9], or the proximal component of a bipartite, single,

grass cotyledon[3, 33]), a second WGNCA analysis was performed on the LM-RNAseq data

obtained from the Stage 1 embryonic shoot apex. In addition, published LM-RNAseq data[42] of

the Plastochron 0 (P0) and P1 leaves, from 14-day after planting maize seedlings, were

integrated as a reference for later leaf ontogeny. Batch effects were removed (Supplementary

Fig. 5a,b), and the merged dataset was filtered to select genes with variability in transcript

accumulation levels across embryonic organs. In the analysis, 5,128 out of 5,137 genes were

selected and clustered into 15 co-expression modules (LM1, LM2, …, LM15). The remaining 9

transcripts were resolved into an unclustered module (Supplementary Table 10).  Transcriptomic

relationships among the co-expression modules are shown in a tree graph (Fig. 4c), in which the

y-axis (height) scales the length of organ-specific branches, indicating transcriptomic distances

between these modules (Supplementary Fig. 5a,b). Pearson correlation tests between module

eigengenes and organ-specific transcriptomes identified associations between specific co-

expression modules and the corresponding embryonic organs (Fig. 4d and Supplementary Fig.

5c). Two major branches comprise (1) the Stage 1 leaf, SAM, P0 and P1, and (2) the coleoptile

and the scutellum.

To further investigate the transcriptomic relationships among all Stage 1 maize embryonic lateral

organs at single-cell resolution, high-dimensional WGCNA (hdWGCNA[43]) was performed on

the scRNAseq data. A total of 5,026 genes transcripts were clustered into 11 co-expression

networks (M1, M2, …, M11; Supplementary Table 11). A hypergeometric test was used to

analyze the overlap between the putative single-cell cluster marker-gene transcripts and those

highly correlated with corresponding co-expression modules. The data indicate that the SAM and

the leaf cluster marker-genes significantly overlap with the closely related modules M9 and M10,

respectively. In contrast, marker genes from the scutellum and the coleoptile cluster overlap with

multiple co-expression modules (M5, M7 and M8; Fig. 4d). The single-cell cluster tree shows

that the coleoptile together with the scutellum and the leaf is next to the SAM, consistent with

the LM-RNAseq WGCNA data. (Fig. 4c).

The WGCNA clustering results linking the scutellum and coleoptile were supported by genetic

analysis. *LONELY GUY7* (*LOG7*), which encodes a cytokinin-activating required for maize SAM

maintenance[42], was expressed at the SAM tip in Stage 1 embryos (Supplementary Fig. 6a-c).

Null mutations in *log7* resulted in SAM termination[42]. Histological examination of *log7* mutant

Stage 1 embryos revealed the absence of a detectable SAM or foliar leaf, although the scutellum

and coleoptile were morphologically normal (Supplementary Fig. 6d-h).  These data expose

morphogenetic differences between leaves, and the scutellum and coleoptile.

**The embryonic-organ-initiation genetic network stage is conserved in Arabidopsis embryos**

To investigate the transcriptomic similarities of embryonic lateral organs of maize and those of

the eudicot model Arabidopsis, comparative genomic toolkits [GRAMENE

(https://www.gramene.org/), TAIR (https://www.arabidopsis.org/), BLAST tools

(https://blast.ncbi.nlm.nih.gov/Blast.cgi)] were used to select *Arabidopsis* orthologs of the 130

transcripts in the maize embryonic-organ-initiation network, and to investigate the gene

expression atlas of *Arabidopsis* embryo development[44]. Embryonic nomenclature differs between

eudicots and grasses[1].  In *Arabidopsis*, histological layering and SAM initiation occur during the

globular stage, cotyledon initiation and emergence defines the heart stages, whereas the

cotyledons elongate during the torpedo and bent embryo stages.

Out of 130 genes in the maize embryonic-organ-initiation network, forty-nine had corresponding

*Arabidopsis* orthologs expressed during embryo development, commencing during cotyledon

initiation at the early heart stage and persisting until the late cotyledon stage (Supplementary

Table 12). Notably, the stage-specific accumulation patterns of these homologous maize

transcripts in the emerging scutellum (Transition stage), coleoptile (Coleoptilar stage), and foliar

leaf (Stage 1) were significantly correlated with the reported expression of the corresponding

*Arabidopsis* orthologs during specific stages in cotyledon initiation and development in this

eudicot embryo[44] (Fig. 4e, Supplementary Fig. 7). In addition, transcripts accumulating in the

expanding Stage 1 scutellum correlated with expression of *Arabidopsis* orthologs during the later

stages of cotyledon growth (i.e. late-torpedo stage[44]). Likewise, maize transcripts accumulating

in the elongating Stage 1 coleoptile were significantly correlated with expression of orthologous

*Arabidopsis* genes during earlier stages of cotyledon expansion (i.e., the early heart stage[44]; Fig.

4e). These data were interpreted to indicate that despite the morphological dissimilarities of these

grass and eudicot "mid-staged" embryos, the development of the maize and *Arabidopsis*

embryonic lateral organs share a homologous regulatory network.

**The embryonic organ initiation network comprises a phylotypic stage**

The data reported in this study support the hypothesis that the maize embryonic-organ-initiation

network is conserved in *Arabidopsis thaliana* and correlates with the transcriptomic signatures of

the eudicot heart and torpedo stage embryos. Prior studies report that during mid-embryogenesis,

old and conserved genes are more highly expressed than young and divergent genes, marking the

transcriptomic phylotypic stage of *Arabidopsis* embryogenesis[22]. To investigate if the maize

embryonic-organ-initiation network also comprises a transcriptomic phylotypic period in grass

embryogenesis, the age and divergence of the genes in the maize embryonic-organ-initiation

network were studied.

To evaluate gene age, a phylostratigraphic analyses obtained 13 phylostrata (PS), PS1-PS13 (Fig.

5a and Supplementary Table 13) using statistical genetic methods to identify homologous

founder genes across lineages (including 350 species) and date their phylogenetic origin

(Supplementary Table 14)[45]. Significantly, 102 of the 130 transcripts in this phylotypic maize

embryonic-organ-initiation-network appear in one of three phylostrata, i.e.,  phylostratum PS1

(29; unicellular eukaryotes and prokaryotes), PS2 (36; all eukaryotes), or PS4 (37; Bryophyta).

Less than 10 percent of this maize embryonic network comprises newer transcripts (from PS5-

PS13) that appear to have arisen later in plant evolution (Supplementary Figs. 8a-c). Notably, the

majority of genes in this network appear within the Bryophyte phylostratum PS4, which marks

the first appearance of the plant embryo[24]. In contrast, just 12 embryonic transcripts (9.23%) are

identified in the algal phylostratum (PS3), which consists of plants lacking embryos. Gene

expression numbers during the non-phylotypic Proembryo stage are dominated by later evolved

genes from PS9-PS11 (i.e., the Poaceae or grasses, the panicoid grasses, and the genus *Zea*; Fig.

5a), whereas transcripts from Stage 2 embryos and seedling leaves are enriched for newly

evolved genes from PS12 and PS13 and the genus *Zea* (Supplementary Table 15).

The transcriptome age index (TAI) measures gene age adjusted by gene expression level for each

embryogenic stage (Supplementary Table 16). As shown in Fig. 5b and Supplementary Fig. 8c,

the TAI displays a "classic" hourglass pattern featuring a significant decrease beginning at the

proembryo stage that reaches a minimum value during Stage 1, followed by an increased TAI at

Stage 2 that continues in the newly initiated leaves of later, germinated maize seedlings. The

transcriptome divergence index (TDI) measures the average sequence divergence of genes,

weighted by transcript accumulation level, at different developmental stages (Supplementary

Table 17). Consistent with the TAI results, the TDI also showed an hourglass pattern that

commences at the Proembryo stage and extends to Stage 1, followed by a notable increase at

Stage 2 and in germinated seedling leaf primordia (Fig. 5c and Supplementary Fig. 8d-e). Thus,

measures of both transcriptional gene age and gene sequence divergence reveal a mid-

embryogenic, phylotypic "hourglass" pattern during maize embryogenesis. The data also reveal

that the phylotypic-staged gene networks found in maize and *Arabidopsis* are homologous (Fig.

4e), as is predicted for two seed plants from the same morphological phylotype[13, 23].

**Transcriptomic phylotypic comparisons across plant phyla reveal an inverse hourglass pattern.**

A comprehensive transcriptomic survey of embryonic/sporophytic development in the non-vascular plant *Physcomitrium patens* is available for use in across-phylum analyses of phylotypic gene expression in plants[46]. Ortiz-Ramirez et al., (2016)[46] sampled gene expression in the zygote and young embryo housed within the gametophytic archegonium, and from four subsequent ontogenetic increments in moss embryo development (labeled S1-3, and SM). As described in Janzen (1929)[11] and modeled in Fig. 6a, key morphogenic events in *P. patens* sporophyte development include the establishment of a multicellular embryo during S1 [5-6 days after fertilization (AF)] and subsequent proliferative growth of the seta and spore capsule during both the S2 (9-11 days AF) and S3 stages (18-20 days AF) (Fig. 6a). Formation of the sporangial epidermal layers occurs during S3, followed by meiosis in the mature sporangium during the SM stage (28-30 days AF). It is important to note that the archegonium samples consist of primarily gametophytic, not embryonic/sporophytic, tissues[46].

Five phylostrata were used to determine the ages of genes enriched in *P. patens* embryos, whereas their sequence divergence ($K_a/K_s$) ratios were calculated by comparing gene homologs in *Physcomitrellopsis africana* (Fig. 6b and Supplementary Table 18-20). TAI and TDI were rendered for transcripts enriched during each of the five developmental increments: archegonia, and sporophyte stages S1, S2, S3, and SM.  The data indicate an increasing TAI and TDI pattern from the archegonium stage to S1, whereas older embryonic transcripts increase markedly during S2, reaching a phylotypic peak at S3, followed by increasing TAI and TDI at SM reflecting the

use of relatively younger genes in the mature sporophyte (Fig. 6c). Thus, embryos of this moss

also manifest a phylotypic hourglass pattern of gene expression at mid-embryogenesis.

To test if embryos of *P. patens* utilize a similar phylotypic gene network as the seed plants

*Arabidopsis* and maize, correlations of embryonic gene expression were made across phyla. The

data reveal an inverse hourglass pattern (Fig. 6d), with little correlation between mid-embryonic

phylotypic-staged transcripts of the moss (S2-S3) and genes enriched during the phylotypic

periods of maize or *Arabidopsis*. In contrast, early (S1) and later-staged (SM) embryonic

networks in the moss show increased correlations with genes expressed in the seed plant

embryos. Thus, as described in animal embryos, across-phylum comparisons of plant embryonic

gene expression yield an inverse hourglass pattern characterized by the expression of distinct,

phylotypic gene networks.

**Discussion**

A multiplexed transcriptomic approach was applied to investigate embryogenesis in a monocot

grass plant, which has evolved novel embryonic structures with previously undefined gene

expression networks. Spatial transcriptomic technologies complement the ambiguities

encountered in cell-cluster identification that can otherwise complicate scRNA-seq analyses,

whereas the precision of LM-RNAseq can delineate the tissue/organ-specific transcriptomes of

embryonic organs that are unresolved using spatial transcriptomics alone (Fig. 2). Moreover,

high-throughput, multiplexed *in situ* RNA-targeting technologies enable localizations of multiple

transcripts simultaneously and combinatorically, throughout embryo ontogeny. The RNA-

targeting analyses presented in this study support the expression of a gene network that functions

during the initiation of the grass scutellum, coleoptile, and foliar leaf (Fig. 3). Thus, the

combinatorial application of multivariate transcriptomic approaches enables a high-throughput

and spatially-informed transcriptomic profile of grass embryo development at cellular resolution

(Fig. 1).

Although questions about the evolution and homology of the cotyledon remain unresolved,

genetic and morphological models indicate that the cotyledon evolved via the modification of a

first embryonic leaf[1, 4, 47]. In support of these models, our transcriptomic comparisons of maize

embryonic lateral organs identify a conserved embryonic-organ-initiation network (Fig. 4a),

consistent with the proposition that the scutellum, coleoptile, and leaf 1 are indeed homologous

organs. Likewise, our co-expression analyses, RNA-targeting assays, and genetic analyses of

meristem mutants collectively support the hypothesis that the cotyledon is a bipartite organ

comprising an apical, haustorial scutellum fused to a basal, emergent and sheathing coleoptile[3,

33]. The scutellum and coleoptile cluster together in WGCNA analyses (Fig. 4c) and distant

from foliar leaves, reflecting the expression of genetic networks that are more closely-shared

within these first-elaborated embryonic organs as compared to leaves. Genetic analyses also

reveal that *Zmlog7* mutants fail to make a SAM or leaves, although the scutellum and coleoptile

are unaffected, likewise consistent with morphogenetic differences between leaves and both the

scutellum and coleoptile (Supplementary Fig. 6). Similar phenotypes are reported for null

mutations of the meristem maintenance gene *KNOTTED1,* wherein the SAM and leaf are

absent, but the coleoptile and scutellum are unperturbed by loss of leaf and meristem integrity[30].

Furthermore, expression of the proximal, sheath marker genes *BOPa* and *TRU1*[34,38, 48] is seen in

the Stage 1 coleoptile but not in the scutellum (Fig. 3i-k and Supplementary Fig. 3), in support

of models where the coleoptile is purported to be homologous to the proximal, leaf sheath and

the scutellum comprises the highly modified distal component of a fused, bipartite, grass

cotyledon[3, 33].

Further, our multiplexed transcriptomic analyses of maize embryogenesis identify a genetic

network expressed during initiation of embryonic organs that describes a phylotypic period (Fig

5b-c). As previously reported in the eudicot *Arabidopsis* and in the embryos of fruit flies

(*Drosophila melanogaster*) and zebrafish (*Danio rerio*)[20-22], the transcriptomic phylotypic period

is defined by the preferential expression of evolutionarily ancient and sequence-conserved genes.

We note that the transcriptomic phylotypic period in maize embryogenesis commences during

the late proembryo stage and forms an extended hourglass shape corresponding to successive

initiations of the scutellum (at transition stage), coleoptile (at coleoptilar stage), and leaf 1 (at

Stage 1). Significant decreases in both transcript age and conservation occur during later stages

of maize embryogenesis (Stage 2), and upon the initiation of seedling leaves after germination

(Fig. 5b-c).

As in the seed plants maize and *Arabidopsis*, embryos of the non-vascular moss *Physcomitrium*

*patens* display transcriptomic phylotypic hourglass patterns that peak mid-embryonically (Fig.

6c). In this context, it is worth noting that the phylotypic-stage gene network in *P. patens*

consists of transcripts that are markedly distinct from the gene networks expressed during

phylotypic mid-embryogenesis in *Arabidopsis* and maize. Indeed, across-phylum transcriptomic

comparisons display an inverse hourglass pattern, where the least homologous embryonic gene

expression networks occur at mid-embryonic stages, whereas early and later stages of moss

sporophyte/embryo development show increased co-expression with maize and *Arabidopsis*

embryonic networks (Fig. 6d). These findings are consistent with across phylum phylotypic

comparisons in animals[23], and likewise support predictions that non-vascular plants and seed

plants comprise distinct, embryonic phylotypes[13].

Historically, morphological and transcriptomic phylotypic embryonic periods are reported to

occur at mid-embryogenic in plants and animals[19-22], although such chronological staging is

especially problematic in plants where embryo-morphogenic outcomes can vary dramatically

among taxa[22]. Nonetheless, the question arises as to why older, sequence-conserved genes are

preferentially utilized during mid-embryogenesis in both plants and animals?  What

morphogenetic characteristics embody this mid-embryogenic stage favoring the utilization of

ancestral and conserved gene networks?

One concern when answering this question is the temporal delay between gene expression curves

and the developmental phenomenology that expression evokes. The maize phylotypic period

begins its steep decline in transcriptomic age (TAI) in the proembryo where the inner-and-outer

embryonic layers are formed, then culminates during mid-embryogenic Stage 1 when the

scutellum and coleoptile are expanding primordia, and the first foliar leaf is just initiating (Fig.

1d). Similarly in *Arabidopsis*, the transcriptomic phylotypic hourglass begins its decent during

histological patterning at the globular stage, and peaks during the torpedo stage when the two

cotyledon primordia exhibit expansive growth[22]. Moreover, the transcriptomic phylotypic peak

of the *P. patens* embryo begins during S2 and peaks during S3, stages comprising expansive

growth of the spore capsule from a central meristematic layer and histogenesis of the sporangium

(Fig 6). Thus, although no lateral organs are formed in the moss embryo, the transcriptomic

phylotypic hourglass in *P. patens* correlates with expansive embryonic growth and histogenesis.

Likewise in *Drosophila* embryogenesis[20,21], gene age (TAI) peaks during germband elongation[20],

at the onset of embryo segmentation and initiation of the organogenic imaginal discs[49]. Notably,

the trend toward the usage of older gene transcripts in the fruit fly begins earlier in

embryogenesis, during gastrulation and formation of tissue layers [20]. A similar pattern is

observed in zebrafish (*Danio rerio*), wherein the phylotypic hourglass begins its descent toward

older gene transcripts during gastrulation (histogenesis) and then peaks in the subsequent late

segmentation/early pharyngula stages, when organ primordia initiate and grow[20]. A later

reduction in relative transcriptomic gene age is noted during the zebrafish larval stage, which is

accompanied by additional morphogenesis. Taken in summary, the mid-embryogenic, phylotypic

period of plants and animals correlates with tissue layer organization and expansive growth of

embryonic organs.

Despite their evolutionary distances, we demonstrate that in maize and a moss, as in *Arabidopsis*,

fruit flies, and fish, the processes of embryonic tissue-layer histogenesis and primordial growth

utilize old and conserved genes, comprising anciently evolved, developmental genetic networks

that are proposed to initiate the fundamental body plan[20-22]. Importantly, these phylotypic gene

networks are not homologous across phyla and across kingdoms, but rather appear to comprise

disparate, old and conserved gene networks that are predicted to generate the variable,

phylotypic-staged embryo morphologies seen within animal and plant phylotypes[13, 23]. At the

end of this embryonic phylotypic period, plant and animal organs express newer and divergent

gene networks in species-specific ways, to generate the vast morphological diversity found in

nature.

Our phylotypic analyses of the relatively recently-evolved grass embryo are interpreted to

indicate that an evolutionary mechanism for the innovation of morphological novelty exists in

plants as well as animals. The maize embryo phylotypic period displays significantly higher TAI

values (expression of newer genes) at the Transition and Coleoptilar stages than during Stage 1

(Fig 5b). Notably, no such deviation in the hourglass, phylotypic shape is detected during

cotyledon initiation in *Arabidopsis* embryogenesis[22]. These data likely reflect the relatively

recent evolution of the bipartite grass cotyledon (the scutellum and coleoptile), which arose long

after the cotyledon evolved from an ancestral embryonic leaf 1[3, 4] and whose evolution involved

newer, non-ancestral gene networks. At the same time, the initiating scutellum uses highly-

conserved gene networks corresponding to the TDI maximum of maize embryogenesis (Fig. 5c).

These data indicate that after the scutellum first appeared, the expression of highly-conserved

gene networks during scutellum initiation ensured that this morphologically novel, embryonic

organ was retained in subsequent generations.  Relatedly speaking, the slight decrease in the use

of conserved genes (TDI value) during the Coleoptilar stage may be attributable to

morphogenetic differences in the initiating coleoptile ensuring that the bipartite components of

the single maize cotyledon (scutellum and coleoptile) are morphologically distinct. We interpret

these data to reveal a conserved, testable mechanism involving inverse trends in TAI and TDI

values during the innovation of evolutionary novelty in plants and animals.

**Methods**

**Plant Materials and growth conditions**

B73 inbred lines,  the *ns* 1:1 lines (segregate 1/2 wild type NS1+/*ns1 ns2/ns2* plants and 1/2

*ns1/ns1 ns2/ns2* double mutant[50], and *log7*/+ (obtained from the Maize Genetics COOP Stock

Center: UFMu-02863 mu1030680::Mu) were grown in the Gutermann greenhouse facility,

Cornell University (Ithaca, NY). Transition staged, coleoptilar staged, Leaf 1-staged, and Leaf 2-

staged embryos were collected at 8-9 DAP, 10-11 DAP, 12-13 DAP, 14-15 DAP, respectively.

*ns* 1:1 seeds were planted in soil consisting of a 1:1 mixture of Turface and LM111 in Percival

A100 growth chamber at Weill Hall, Cornell University (Ithaca, NY) with the set condition: day

29.4 °C/16 hours, night 23.9 °C/8 hours, humidity 50%. The seedlings were collected on the 7$^{th}$

day after planting for coleoptile comparisons between WT and *ns1;2* mutants.

**Single-cell transcriptomic analysis**

Two batches of B73 Leaf1 staged embryos were collected in summer 2021 and winter 2022. For

each batch, 200 embryos were dissected from developing seeds for protoplast isolation as

previously described[51]. Protoplast (1000-2000 cells per µL)  were loaded into the 10X Genomics

Chromium platform to generate a Gel Bead-In-Emulsion (GEM). The GEM-captured RNA was

reverse transcribed and amplified to make 3′ cDNA libraries, followed by the high throughput

RNA sequencing on the Illumina NextSeq 2000 sequencer at the Cornell University

Biotechnology Resource Center.

The resulting raw base call (BCL) files were demultiplexed to generate FASTQ files via

CellRanger mkfastq v6.0 (10x Genomics, Pleasanton, CA). FASTQ reads were trimmed, aligned

(to B73 reference genome V3[52]) and assigned to cell barcodes to generate expression count

matrices via CellRanger count with default settings. The expression count matrices were then

analyzed using Seurat v4[53-55]. In detail, the matrices were filtered with the setting (number of

genes > 1500, UMI > 6000) to remove low quality cells. The filtered matrices were normalized

and log transformed via SCTransform function. The normalized matrices were merged, and

batch effects were removed via R package Harmony v1.2.0[56]. The cell-cycle associated genes

were scored and regressed via CellCycleScoring and ScaleData function. Matrix dimension was

performed using RunPCA. The cell clusters and the Uniform Manifold Approximation and

Projection (UMAP) plot were generated with the setting (dim = 1:40, resolution = 1, n. neighbors

= 40, min. dist = 0.01, spread = 3). The differential expressed genes and cell cluster markers

were identified via FindAllMarkers function with the log2 fold change threshold = 0.2.

**Spatial transcriptomic analysis**

For use in spatial transcriptomics, B73 Leaf1 staged embryos were processed for either paraffin-

embedding and microtome sectioning, or cryoembedding and cryosectioning.  For paraffin -

embedding,  two embryos were fixed in Farmer's Fixative (Ethanol:Acetic acid = 3:1), followed

by dehydration in an ethanol concentration gradient (75%, 85%, 95%, 100%) and clearing with

Ethanol:Xylene gradient of 3:1, 1:1, 1:3, and 100% Xylene at room temperature. Cleared

embryos were embedded in Paraplast Plus® (Leica Biosystems, Deer Park, IL). The embedded

blocks were microtome-sectioned at 10 um at room temperature; sections were subsequently

mounted onto 10x Visium™ Spatial Gene Expression Slides (10x Genomics, Pleasanton, CA).

Slides were deparaffinized with two changes of 100% Xylene and 95% Ethanol for 10 min and 2

min, respectively. For cryoembedding, two kernels were dissected longitudinally to remove left

and right margin of the kernel; the central region containing the embryo was embedded in

Tissue-Tek® Optimal Cutting Temperature (O.C.T., Sakura Finetek, Torrance, CA), and snap-

frozen in the dry ice. Frozen tissue blocks were cryosectioned at10 µm, and the sections were

mounted on 10x Visium™ Spatial Gene Expression Slide (10x Genomics, Pleasanton, CA).

The 10x Visium™ Spatial Gene Expression Slides were processed for spatial library

construction following the manufacturer instructions. Briefly, the sections were baked at 37 $^{\circ}$C

for 1 min followed by re-fixation in 100% methanol at -20 $^{\circ}$C for 30 min. Subsequently, the

sections were stained with Toluidine Blue for histological imaging. Tissue permeabilization was

performed at 37 $^{\circ}$C for 6 min to release RNA from the fixed tissue, followed by first strand

cDNA synthesis. Subsequently, second strand cDNA was synthesized and amplified for library

construction and sequencing. High-throughput RNA sequencing was conducted on an Illumina

NextSeq 2000 sequencer at the Cornell University Biotechnology Resource Center.

Raw reads were demultiplexed, mapped to the Toluidine Blue stained images, and aligned to the

B73 reference genome V3[52] to generate spatial expression count matrices using SpaceRanger

v2.0 (10x Genomics, Pleasanton, CA). The matrices were analyzed using Seurat v4[53-55]. In detail,

the matrices were normalized and log transformed via SCTransform function. The normalized

matrices were merged followed by dimensionality reduction. Cell clusters and the UMAP were

generated with the following settings (dim = 1:40; resolution = 1.5; neighbors = 10; minimum

distance =0.01). The spatial data were visualized with DimPlot and SpatialPlot; differential

expressed genes and cell cluster markers were identified via FindAllMarkers function with the

log2 fold change threshold = 0.1.

**Laser-microdissection RNAseq analysis**

Sixty maize Leaf1 staged kernels in two replicates were dissected longitudinally to remove the

left and right margins of the kernels. This embryo-containing central region was fixed in

Farmer's Fixative (Ethanol:Acetic acid=3:1). After fixation, the samples were dehydrated,

cleared, embedded and sectioned as described previously[57, 58]. Tissue sections were mounted on

PEN membrane slides (Leica Microsystems, Deer Park, IL) pre-treated at 180 $^{o}$C for 4 hours

followed by UV irradiation at 254 nm for 30 minutes. The sections were deparaffinized with two

changes of 100% Xylene and 100% Ethanol for 10 min and 2 min, respectively. Laser

microdissection was performed using a PALM Laser Microbeam System (Zeiss Group,

Oberkochen, Germany). Adaxial scutellum, SAM, leaf, as well as upper and lower coleoptiles

were microdissected (total area for each captured organ is over 0.3 mm$^2$) and RNA extraction

was performed using the Arcturus$^{TM}$ PicoPure$^{TM}$ RNA Isolation Kit (Thermo Fisher Scientific,

Waltham, MA). The extracted RNA was amplified using Arcturus$^{TM}$ RiboAmp$^{TM}$ HS PLUS

RNA Amplification Kit (Thermo Fisher Scientific, Waltham, MA). RNA quality control, library

construction and sequencing were performed at the Cornell University Biotechnology Resource

Center. The raw reads quality control was conducted via FastQC

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Raw reads were mapped to the.

B73 reference genome V3[3] by HISAT2 (v2.1.0)[59, 60]. Read counts and the normalized number of

transcripts (transcript per million reads, TPM) were calculated as previously described[61]. DEseq2

(v1.26.0)[62] was used to identify differentially expressed genes (DEGs) with the cutoff criteria of

log2 fold change >1 or <1 with the false discovery rate (FDR) < 0.05.

**Multiplex *in situ* hybridization**

Transition staged, coleoptilar staged and Leaf1 staged kernels (two replicate for each stage) were

dissected longitudinally to remove the left and right kernel margins. The central region

containing the embryo was fixed in 4% paraformaldehyde (PFA); fixed samples were

dehydrated, cleared, embedded and sectioned as described previously[63, 64]. Sections were

mounted on 10x Genomics Xenium[TM] slides, deparaffinized and de-crosslinked following the

manufacturer's instructions. The Xenium custom gene expression panel containing probes of 99

maize genes were applied on each section to perform hybridization, ligation and rolling circle

amplification, followed by cycles of fluorescent probe hybridization, imaging and decoding. The

data was visualized by Xenium Explorer v1.3. The Xenium[TM] multiplex *in situ* hybridization

assays were conducted at Memorial Sloan Kettering Cancer Center (New York, NY).

**Histology and immunohistolocalizations**

Developing kernels from the proembryo stage, transition stage, coleoptilar stage, and the Stage 1

were fixed in FAA, dehydrated, cleared, embedded in paraplast, sectioned and stained as

described previously[64]. Immunohistolocalizations were performed using an antibody to maize

Class I KNOX proteins, KNOTTED1 and ROUGH SHEATH1, as described[65].

**WGCNA and hdWGCNA**

For WGCNA to pursue to initiation network, log2-transformed TPM data from LM-RNAseq

(SAM data was excluded) were filtered following these criteria: row mean > 0.2 and coefficient

of variation > 0.4. The general pipeline was performed as per R package instructions[40] with

minor modifications as follows. The network type was set to "signed". The soft power, the

minimum module size ,and the merging threshold were set to 13, 20 and 0.3, respectively. For

the analysis to pursue transcriptomic relationship between organs, the LM-RNAseq data in this

study was merged with the LM-RNAseq data of the 14-day seedling P0 and P1 data[42]. The data

was normalized and batched effect removed as described, and filtered following the same criteria

as the WGCNA for the initiation network. The network type was set to "signed". The soft power,

the minimum module size ,and the merging threshold were set to 16, 20 and 0.4, respectively.

For hdWGCNA, the single-cell RNAseq data was used to perform the analysis. The data was

SCTransformed and normalized. The general pipeline was conducted as the R package

instructions[43] with the following minor modifications. The network type was set to "signed", and

the soft power and minimum module size were set to 10 and 30, respectively. Hypergeometric

test was performed to analyze the correlation between the hdWGCNA modules and the single-

cell clusters following the previously described procedures[66].

**Statistical analysis**

Hypergeometric tests were performed to analyze the correlation between single-cell clusters and

spatial transcriptomic clusters, as well as between the single-cell clusters and hdWGCNA

modules. We followed previously algorithms[66] to statistically evaluate if the two groups of data

were significantly correlated by calculating the number of overlapping gene relative to the total

number of protein-coding genes in the maize B73 reference genome V3[52]. The p values were

transformed by –log10p, and visualized as heatmap built by the R package gplots

(https://github.com/cran/gplots).

LM-RNAseq read count data were normalized to fit the generalized linear model (GLM), and

Empirical Bater shrinkage estimation was used for calculating dispersions, expression fold

changes and adjust p values. In addition, Pearson correlation test was used to calculate the

correlation between the coexpression modules and the corresponding clusters (microdissected

organs or putative initiation network) in WGCNA, and between the maize and Arabidopsis

embryogenesis genes.

To statistically compare the expression of the initiation network between the transition,

coleoptilar and the stage1, the TPM data was log2 transformed [log2(TPM+1)] and compared

using Tukey-Kramer HSD test by JMP Pro 17 (SAS Institute, Cary, NC) with default settings.

For the correlation test, the transition, coleoptilar and the stage1 TPM data was double log2

transformed, i.e. log2[log2(TPM+1)+1], to fix the data skewness caused by the outliers. The

Arabidopsis TPM data was log2 transformed. The Pearson correlation was performed in R v4.3.2

with the significance cutoff value $p<0.05$.

**Calculating TAI and TDI for phylotypic period**

The procedures of constructing a phylostratigraphic map, calculating TAI and TDI, and

statistical tests, have been presented previously[20,22,45,67]. For each of the thirteen phylostrata for

maize (Fig. 5a), the amino acid sequences of all 350 species (Supplementary Dataset 15) with

completely sequenced genomes were extracted from NCBI, Ensembl genomes, Treegene

(https://treegenesdb.org), Fernbase (https://fernbase.org), or Maizegdb  (https://maizegdb.org).

An equivalent  strategy was applied for the five *P. patens* phylostrata f(Fig. 6b), encompassing

666 174 species (Supplementary Dataset 17). Each of the 63,031 amino acid sequences in *Zea mays*

667 ssp. *mays* (v3) and 24, 002 sequences in *P. patens* (v1.6) with minimum length of 30 amino acids

668 was analyzed by BLASTp (BLAST version 2.15.0) with an E-value cut-off of $10^{-5}$. Each such

669 gene of *Zea mays* ssp. *mays* was assigned to the phylogenetically most distant (oldest)

670 phylostratum containing at least one species with at least one blast hit.  In circumstances where

671 no blast hit was identified in older phylostrata, the *Zea mays* ssp. *mays* gene was assigned to

672 phylostratum 13 (PS13; Fig. 5a). In this way, each gene of *P. patens* was likewise assigned to

673 one *P. patens* phylostratum, from PS1 to PS5 (Fig 6b).

674 To evaluate gene sequence divergence, the $K_a/K_s$ ratio was calculated between maize and its

675 sister species *Zea diploperennis* or *Zea Mexicana*, between *Physcomitrium patens* and the

676 closely-related species *Physcomitrellopsis africana*[68]. $K_a$ and $K_s$ refer to the number of non-

677 synonymous and synonymous substitutions, respectively, between each orthologous pair.

678 Orthologous gene pairs were determined with the method of best hits using BLASTp. Amino

679 acid sequence alignments of each pair generated with MAFFT[69] (L-INS-i option) were used for

680 codon alignments generated with PAL2NAL[70] to compute sequence divergence levels ($K_a/K_s$)

681 with KaKs Calculator (version 3.0)[71]. Gene pairs with $K_a<0.5$, $K_s<5$ and $K_a/K_s$ ratios $< 2$ were

682 retained.

683 Transcriptomic data for maize embryonic developmental stages were collected from four data

684 sources 1) LM-RNAseq data of proembryo stage[28], 2) organ initiation network (this study), 3)

685 RNAseq data of Stage 2 embryo[72] and 4) LM-RNAseq data of 14-day after germination seedling

686 P0 and P1 leaves[42]. The datasets were processed with batch effect removal and normalization as

687 described above. The TAI and the TDI are weighted means of evolutionary age and sequence

divergence, respectively, and defined previously[20, 22]. The transcriptome age index $TAI_s$ and the

transcriptome divergence index $TDI_s$ of developmental stage s (s = proembryo, transition,

coleoptilar, stage 1 leaf, stage 1 scutellum, stage 1 coleoptile, seedling leaves P0 and seedling

leaves P1) are the weighted mean of the evolutionary age (phylostratum) weighted by the

expression level of each gene at developmental stage s, and the weighted mean of the Ka/Ks

ratio of each gene weighted by the expression level of each gene at developmental stage *s*,

respectively. Transcriptomic data for embryo/sporophyte stages in *P. patens* were from Ortiz-

Ramirez et al., (2016)[46]. Low/high PS values correspond to evolutionarily old/young genes, so

low/high TAI values correspond to evolutionarily old/young transcriptomes. Low/high $K_a/K_s$

ratios correspond to conserved/divergent genes, so low/ high TDI values correspond to

conserved/divergent transcriptomes. For gene models with multiple predicted peptides, TAI was

calculated with the most conserved phylostratum assigned to that locus and TDI was calculated

with the conserved codon divergence strata assigned to that locus.

To determine the statistical significance of the TAI and TDI profiles, the flat line test[22], reductive

hourglass test and reductive early conservation test were performed via myTAI package

(v1.0.1)[67]. The flat line test is a permutation test based on the variance of the TAI or TDI values

of a given TAI or TDI profile. For the reductive hourglass test in maize, the embryonic stages

were partitioned into three modules, early (proembryo stage), mid (transition stage, coleoptilar

stage, Stage 1 leaf, Stage 1 scutellum, and Stage 1 coleoptile), and late (Stage 2 and seedling

leaves), and a permutation test was performed for the minimum differences of the mean TAI or

TDI values between early and mid-modules, and between late and mid modules. For the

reductive early conservation test in maize, a permutation test was performed for the minimum

differences of the mean TAI or TDI values between mid and early modules, and between late

and early modules. For the reductive hourglass test and reductive early conservation tests on *P.*

*patens*, the three modules comprised the archegonium and S1 stage (early), S2 and S3 (middle),

and SM (late).  Detailed procedures are described[67].

**Data availability**

The single-cell RNAseq, Visium^TM spatial transcriptomics, and LM-RNAseq data were

deposited in the NCBI Sequence Read Archive (SRA) with the BioProject ID: PRJNA1045558.

The Xenium^TM multiplex *in situ* hybridization data was deposited in Science Data Bank with the

link: https://www.scidb.cn/s/F7ZbEz, which could be visualized by Xenium Explorer v1.3.

**Code availability**

The code for TAI and TDI calculations was uploaded to

https://gitlab.com/rz444/maizephylper.git.

**Figure Legends**

**Fig. 1 Multiplexed spatial transcriptomics of the maize Stage 1 embryo.**

Differential interference contrast (DIC) microscopy of maize embryos at the Proembryo stage

(**a**), Transition stage (**b**) Coleoptilar stage, (**c**) and Stage 1 (**d**). Immunohistological analyses of

the Class I KNOX proteins mark the shoot apical meristem (SAM). ep, embryo proper; en,

endosperm; su, suspensor; sc, scutellum; c, coleoptile; L, leaf; *, SAM. Scale bar: 50 μm (**a** and

**b**) and 100 μm (**c** and **d**). **e**. Uniform Manifold Approximation and Projection (UMAP) of Stage1

embryo single-cell transcriptomic profiling with cell/tissue types assigned for each cell cluster.

**Fig. 2 Spatial transcriptomics of maize Stage 1 embryo**

Longitudinal cryosections of Stage 1 embryos embedded in OCT (optimal cutting temperature

compound; **a-b**) or in paraffin (**c-d**) for 10x Visium[TM] spatial transcriptomics analysis. **a, c**

Toluidine Blue-stained sections for 10x Visium[TM] spatial transcriptomics assay are overlayed

with the10x Visium[TM] spatial map (**b, d**). The various color-coded circles correspond to the

capture spots where spatial cDNA libraries were generated. Scale bar = 100 μm. **e.** Longitudinal

sections of paraffin-embedded Stage 1 embryo for use in laser microdissection RNA-seq analysis

(LM-RNAseq). **f.** Enlarged view of a Stage 1 embryo showing regions microdissected for LM-

RNAseq). Figure labels: scu, scutellum; uc, upper coleoptile; lc, lower coleoptile. **e, f** Scale

bar=100 μm. **g**. UMAP of the Visium[TM] spatial transcriptomic analysis of Stage 1 embryos cell

clusters. The spatial, embryonic origins of each cluster are indicated by the color-coded embryo

cartoon; numbers correspond to the cluster designations on the UMAP. **h**. Heatmap illustrating

the correlation between the single-cell (sc) clusters (y-axis) and Visium[TM] spatial clusters

(x=axis). The significance level is marked by $-\log_{10}$p. **i**. Expression of candidate genes identified

in microdissected embryonic organs. Cell clusters exhibiting elevated candidate gene expression

of the specific markers are indicated; expression values are $\log_2$ transformed, i.e., $\log_2(TPM+1)$.

**Fig. 3 RNA-targeting analyses of Transition-staged, Coleoptilar-staged, and Stage 1**

**embryos**

**a-c.** Signal point map of *KN1, CUC2, NS1, WOX3A* and *DRL1* at the Transition stage (**a**),

Coleoptilar stage (**b**) and Stage 1 (**c**). **d-h.** Expression density map of *KN1* (**d**)*, CUC2* (**e**)*, NS1*

(**f**)*, WOX3A* (**g**) and *DRL1* (**h**) at Stage 1. **i-k, l-n, o-q** and **u-w**. Signal point map of *BOPa* (**i-k**),

*YAB14* (**l-n**), *IG1* (**o-q**), and *ARF3* (**u-w**) at the Transition stage, Coleoptilar stage and Stage 1. **r-**

**t** and **x-z**. Expression density map of *IG1* (**r-t**) and *ARF3* (**x-z**) at the corresponding stages.

Figure labels: T, Transition stage; C, Coleoptilar stage; L: Stage 1. The point colors are

arbitrarily determined. Yellow asterisks mark the adaxial expression of *IG1* and *ARF3*, and red

asterisks mark the abaxial expression of *ARF3*. The green asterisk in panel **k** marks the

expression maximum of *BOPa* at the emerging leaf sheath. The yellow asterisks in panel **r-t** and

**z** mark the expression maxima of *IG1* at the adaxial side of the scutellum (**r**), the coleoptile (**s**),

the coleoptile and the leaf (**t**), as well as the expression maxima of *ARF3* at the adaxial side of

the scutellum (**z**), respectively. The red asterisks in panel **x-z** mark the abaxial expression of

*ARF3* at the scutellum (**x, y**), the coleoptile (**y, z**) and the leaf (**z**). Scale bar for all panels = 50

μm.

**Fig. 4 Dynamics of the maize embryo organ initiation network**

**a**. A Venn diagram of embryonic transcriptomics identifies 130 genes in the embryo-organ-

initiation-network. **b**. Normalized expression of these 130 genes at the Proembryo stage (PE),

Transition stage (T), Coleoptilar stage (C), Stage 1 scutellum (Scu), Stage 1 upper coleoptile

(UC), Stage 1 lower coleoptile (LC), Stage 1 leaf (L), as well as P0 and P1 of the 14 day-after-

germination seedlings. The letters above the boxplot represent significance levels (non-

overlapping letters indicate significant differences at $p < 0.05$) using the Tukey-Kramer HSD test.

**c.** WGCNA tree graph of the transcriptomic relationships among organ-specific co-expression

modules. The height (y-axis) scales the length of organ-specific branches, indicating the

transcriptomic distances between modules. **d**. Heatmap illustrating correlations between single-

cell (sc) clusters (y-axis) and hdWGCNA modules (x=axis). The significance levels are marked

by $-\log_{10}p$. **e**. Heatmap illustrating the correlation of the initiation network expression between

Maize (y-axis) and Arabidopsis (x-axis) across the embryogenesis. Heatmap colors are

determined by correlation coefficient (r) via Pearson correlation test with significance level

$p < 0.05$ (*) or $p < 0.01$ (**). Scu, Scutellum; Col, coleoptile; hypoc, hypocotyl.

**Fig. 5 Phylotypic stage analysis of the maize embryogenesis**

**a**. Phylostratigraphic map of maize. All maize protein-coding transcripts were assigned to

thirteen phylostrata (PS1-PS13). Parentheses designate the number of genes per phylostratum;

taxonomic groupings are marked above the corresponding PS. **b-c**. The transcriptome age index

(TAI; **b**) and transcriptome divergence index profiles (TDI; **c**) during maize embryogenesis and

formation of seedling leaves after germination. Figure labels: PE, proembryo stage; T, transition

stage; C, coleoptilar stage; L, Stage 1 leaf; Scu, Stage 1 scutellum; Col, Stage 1 coleoptile; P0

and P1, Plastochron stages of the 15th and 14th leaves in the 14-day after planted seedlings. The

gray lines represent the standard deviation estimated by bootstrap analysis as described[22]. The

overall patterns of TAI and TDI profiles are significant, as measured by the flat line test

($P_{TAI}=3.56\times10^{-6}$, $P_{TDI}=4.80\times10^{-4}$), reductive hourglass test ($P_{TAI}=1.49\times10^{-4}$, $P_{TDI}=1.53\times10^{-3}$),

and reductive early conservation test ($P_{TAI}=1$, $P_{TDI}=0.998$).

**Fig. 6 Across-phylum phylotypic analyses of *Physcomitrium patens*.**

**a.** Development of embryo/sporophyte in the moss *P. patens*.  **b**. Phylostratigraphic map of *P.*

*patens*. All *P. patens* protein-encoding transcripts were assigned to one of five phylostrata (PS1-

PS5). Parentheses refer to the number of genes per phylostratum; taxonomic groupings are noted.

**c**. The transcriptome age index (TAI) and transcriptome divergence index (TDI) profiles of *P.*

*patens* archegonia; sporophyte stages S1, S2, S3 and SM. Colored bars illustrate the duration and

timing of different S stages along the 30 day time scale. **d**. Comparative expression levels of

stage-specific embryonic genes between *Arabidopsis* and *P. patens,* and maize and *P. patens*.

Figure labels: (in moss) AR, archegonium; S1, sporophyte 1 stage; S2, sporophyte 2 stage; S3,

sporophyte 3 stage; SM, mature sporophyte stage; (in maize) PE, proembryo stage; T, transition

stage; C, coleoptilar stage; L, Stage 1 leaf; Scu, Stage 1 scutellum; Col, Stage 1 coleoptile; S2,

Stage 2. The gray lines represent the standard deviation estimated by bootstrap analysis as

described[22]. The overall patterns of TAI and TDI profiles are significant, as measured by the flat

line test ($P_{TAI}$=0.0131, $P_{TDI}$=4.73×10$^{-3}$), reductive hourglass test ($P_{TAI}$=0.0332, $P_{TDI}$=0.0249),

and reductive early conservation test ($P_{TAI}$=0.998, $P_{TDI}$=0.973). Images in (**a**) and (**c**) are

reproduced from Ortiz-Ramirez et al., (2016)[46].

**Supplementary figure legends**

**Supplementary Fig. 1 Verification of single-cell cluster marker genes.**

Left panel: single-cell cluster marker expression projected as a UMAP. Purple dots mark cells

with corresponding gene-specific expression. Middle panel: verification of single-cell cluster

marker genes by Visium$^{TM}$. Red to green spots mark the locations of gene expression with high

to low level, respectively. Right panel: verification of single-cell cluster marker genes by

multiplexed RNA-targeting *in situ* hybridization [data not available for *GRMZM2G166141* (**k**),

*GRMZM2G094353* (**m**) and *GRMZM2G099454* (**p**)]. Scale bar = 100 μm.

**Supplementary Fig. 2 Differential expression between laser microdissected RNAseq**

**samples.**

**a**. Principal component analysis of laser-microdissected RNAseq samples. Each dot represents a

laser-microdissected embryonic shoot sample. **b-f**. Pairwise comparison of the gene expression

between laser-microdissected RNAseq samples. The number of overlapping, differential-

expressed genes (DEGs) is showed in the Venn diagram. The circled numbers represent the

genes preferentially expressed in the SAM (**b**), upper coleoptile (**c**), scutellum (**d**), leaf (**e**) and

lower coleoptile (**f**).

**Supplementary Fig. 3 Multiplex *in situ* hybridization of *TRU1*.**

Spatio-temporal expression of *TRU1* displayed as a signal point map (**a-c**) and as a transcript

abundance correlated signal density map (**d-f**) at the transition stage (**a and d**), coleoptilar stage

(**b and e**) and Stage 1 (**c and f**). The red arrow in (f) marks the expression of *TRU1* at the

emerging leaf sheath of leaf 1. Scale bar = 100 μm.

**Supplementary Fig. 4 Expression of the embryo organ initiation network.**

**a**. WGCNA identifies modules with overall highest expression at Stage 1 leaf (L), less in the

lower coleoptile (LC), and steadily decreased expression in the upper coleoptile (UC), scutellum

(S). Heatmap shows the correlation coefficients (r) and the corresponding p values (parentheses)

of modules ME1-12. **b**. Expression of the 130 genes comprising the embryonic-organ-initiation

network at the transition stage (T), coleoptilar stage (C) and the Stage 1 scutellum (Scu), upper

coleoptile (UC), lower coleoptile (LC) and leaf 1 (L). **c-e**. Correlations of the expression of these

130 genes during initiation of the scutellum, coleoptile, and leaf 1.

**Supplementary Fig. 5 WGCNA Module-organ correlation.**

**a-b**. Correlations of transcriptomic datasets from embryonic organs and leaves initiated post-

germination before and after batch effect removal. Correlations of the normalized overall gene

expression between the post- germination seedling leaves and embryonic stages before (**a**) and

after (**b**) batch effect removal. **c**. Heatmap highlights correlations between co-expression

modules (y-axis) and corresponding organs (x-axis). Colors are graded based on correlation

coefficients (r) and p values (in parentheses).

**Supplementary Fig. 6 Genetic analyses of *log7* mutant embryos.**

Expression and genetic analysis of *LOG7*. **a.** Conventional *in situ* hybridization of *LOG7*

transcript accumulation in a longitudinal section of the Stage 1 embryo. Multiplexed RNA-

targeting point map (**b**) and density map (**c**) of *LOG7* accumulation in longitudinal sections of

Stage 1 embryos. Scale bar = 100 µm. **d .**  Longitudinal median section of a WT Stage 2

embryo. The red dashed line marks the transverse section plane depicted in (**g**). **e-f.**

Longitudinal median sections of the *log7* mutant Stage 2 embryos. The red dashed line marks

the transverse section plane depicted in (**h**). Longitudinal off-median section of the *log7* mutant

Stage 2 embryo. All scale bars = 100 µm.  Asterisks (*) mark the SAM; sc, scutellum; c,

coleoptile; L, leaf.

**Supplementary Fig. 7 Correlation between the maize initiation network genes and the**

**Arabidopsis orthologs.**

The correlation of expression between the maize initiation network genes at transition (**a**) and

coleoptilar (**b**) stage, and the Arabidopsis orthologs at early heart stage.

**Supplementary Fig. 8 TAI and TDI for embryo stages under different BLASTp E-value**

**cutoffs.**

**a-c**. Distribution of all maize genes and genes in each embryo stage across phylostrata and TAI

profiles for different E-value cut-offs in BLASTp homology searches (ranging from 1E-2 to 1E-

20). This analysis investigates whether and how different E-values affect the hourglass pattern. **a**.

Variation of the E-value cut-off for the BLASTp analysis demonstrates minimal effects on the

distribution of maize genes among the phylostrata, indicating an overall pattern independent of

the E-values cut-offs. The y-axis is presented on a logarithmic scale. **b**. Similarly, the variation

of the E-value cut-off for the BLASTp analysis does not severely alter the distribution of maize

genes in each embryo stage among the phylostrata. **c**. The hourglass pattern of the TAI profile

remains consistent across various E-value cut-offs. **d-e**. Transcriptome divergence indices based

on two comparisons of genomes from three relatives of *Zea mays*: *Zea mays* ssp. Mays vs. *Zea*

*mays* ssp. mexicana (**d**); *Zea mays* ssp. mays vs. *Zea diploperennis* (**e**). The overall patterns of

TDI profiles are significant, as measured by the flat line test ($P_{\text{TDI-Mexicana}} = 1.07\times10^{-3}$, $P_{\text{TDI-}}$

$_{\text{diploperennis}} = 4.80\times10^{-4}$), reductive hourglass test ($P_{\text{TDI-Mexicana}} = 0.027$, $P_{\text{TDI-diploperennis}} = 1.53\times10^{-3}$),

and reductive early conservation test ($P_{\text{TDI-Mexicana}} = 0.68$, $P_{\text{TDI-diploperennis}} = 0.998$).

**Supplementary information.**

Supplementary Table 1. Single-cell RNAseq cluster marker genes

Supplementary Table 2. Identity of single-cell clusters

Supplementary Table 3. Spatial transcriptomic cluster markers

Supplementary Table 4. Normalized gene expression (TPM) of LM-RNAseq

Supplementary Table 5. Normalized gene expression (log2 (TPM+1)) of tissue specific classic

genes

Supplementary Table 6. Gene expression of selected leaf genes in the transition, coleoptilar and

Leaf 1 stage in the LM-RNAseq data

Supplementary Table 7. Xenium$^{TM}$ multiplex in situ hybridization probes and result

Supplementary Table 8. WGCNA: modules with gene expression pattern: Scu < UpCol < LoCol

< Leaf

Supplementary Table 9. Putative initiation network and the log2-transformed TPM in the lateral

organs of the transition, coleoptilar, Stage 1 (L1) and 14-day after germination seedlings

Supplementary Table 10. WGCNA: Cell-type specific modules

Supplementary Table 11. hdWGCNA module table

Supplementary Table 12. Maize initiation network vs Arabidopsis orthologs

Supplementary Table 13. Genomes used for phylogeny reconstruction and phylostratigraphic

map of maize

Supplementary Table 14. Phylostrata of maize genes

Supplementary Table 15. The gene numbers of each phylostratum in maize embryo stages

Supplementary Table 16. Gene expression for TAI and TDI calculation in maize

Supplementary Table 17. *Ka/Ks* ratio of maize genes expressed at the embryogenesis and 14-day

seedling

Supplementary Table 18. Genomes used for phylogeny reconstruction and phylostratigraphic

map of *Physcomitrium patens*

Supplementary Table 19. Phylostrata of *Physcomitrium patens* genes

Supplementary Table 20. Ka/Ks of *Physcomitrium patens* genes

**References**

1.  Kaplan, D. R., & Cooke, T. J. Fundamental concepts in the embryogenesis of dicotyledons: a morphological interpretation of embryo mutants. *Plant Cell* **9**, 1903–1919 (1997).

2.  Randolph, L. F. Developmental morphology of the caryopsis in maize. *J Agric Res* **53**, 881-916 (1936).

3.  Kaplan, D. R., & Specht, C. D. *Kaplan's Principles of Plant Morphology*. New York, NY: CRC Press (2022).

4.  Chandler, J. W. Cotyledon organogenesis. *J Exp Bot* **59**, 2917-2931 (2008).

5.  Abbe, E. C., Stein, O. L. The growth of the shoot apex in maize: embryogeny. *American Journal of Botany* **41**, 285-293 (1954).

6.  Bommert, P., & Werr, W. Gene expression patterns in the maize caryopsis: clues to decisions in embryo and endosperm development. *Gene* **271**, 131-142 (2001).

7.  Nardmann, J., & Werr, W. Patterning of the maize embryo and the perspective of evolutionary developmental biology. In: *Handbook of Maize: Its Biology* (eds Jeff L. Bennetzen, Hake SC). Springer (2009).

8.  Weatherwax, P. Position of Scutellum and Homology of Coleoptile in Maize. *Botanical Gazette* **69**, 179-182 (1920).

9.  Boyd, L. Evolution in the monocotyledonous seedling: a new interpretation of the morphology of the grass embryo. *Transactions of the Botanical Society of Edinburgh* **30**, 286-303 (1931).

10. Chen, M. *et al.* Comparative analysis of embryo proper and suspensor transcriptomes in plant embryos with different morphologies. *Proc Natl Acad Sci U S A* **118** (2021).

11. Janzen, P. Die Entwicklung und Biologie von Funaria hygrometrica. In: L. Loeske, ed., Die Laubmoose Europas. II. Funariaceae. pp, 72 -111. *Verlag Max Lande, Berlin-Schoneberg* (1929).

941 *12.* Reski, R. *Development, Genetics and Molecular Biology of Mosses. Botanica Acta* **111**, 1-15

942  (1998).

943 13. Niklas, K. J. et al. Haeckel's biogenetic law and the land plant phylotypic stage. *BioScience* **66**(6),

944  510-519 (2016).

945 14. Dresselhaus, T., & Jürgens, G. Comparative embryogenesis in angiosperms: activation and

946  patterning of embryonic cell lineages. *Annual Review of Plant Biology* **72**, 641-676 (2021).

947 15. Frank, M. H., & Scanlon, M. J. Transcriptomic evidence for the evolution of shoot meristem

948  function in sporophyte-dominant land plants through concerted selection of ancestral

949  gametophytic and sporophytic genetic programs. *Molecular Biology and Evolution* **32**(2), 355-

950  367 (2015).

951 16. Haeckel, E. *Generelle Morphologie der Organismen. Allgemeine Grundzüge der Organischen*

952  *Formen-Wissenschaft, Mechanisch begründet durch die von Charles Darwin reformierte*

953  *Descendenz-Theorie, vols I and II*. Verlag Georg Reimer, Berlin (1866).

954 17. Meckel, J.F. *Entwurf einer Darstellung der zwischen dem Embryozustande der höheren Tiere und*

955  *dem permanenten der niederen stattfindenden Parallele: Beyträge zur vergleichenden Anatomie*.

956  Carl Heinrich Reclam, Leipzig (1811).

957 18. Serrés, E. R. A. Théorie des formations organiques, ou rescherches d'anatomie transcendante sur

958  les lois de l'organogénie, appliquée à l'anatomie pathologique. *Annales de la Societe des Sciences*

959  *Naturelles et d'Archeologie de Toulon et du Var* **21**, 5-49 (1827).

960 19. Sander, K. *In Development and Evolution: the sixth Symposium of the British Society for*

961  *Developmental Biology*. Cambridge University Press. (1983).

962 20. Domazet-Loso, T., & Tautz, D. A phylogenetically based transcriptome age index mirrors

963  ontogenetic divergence patterns. *Nature* **468**, 815-818 (2010).

21.  Kalinka, A. T. *et al.* Gene expression divergence recapitulates the developmental hourglass model. *Nature* **468**, 811-814 (2010).

22.  Quint, M., et al. A transcriptomic hourglass in plant embryogenesis. *Nature* **490**, 98-101 (2012).

23.  Levin, M., et al. The mid-developmental transition and the evolution of animal body plans. *Nature* **531**, 637-641 (2016).

24.  Soltis, P.S. et al. The diversification of flowering plants. In: *Assembling the tree of life,* (eds Cracraft J, Michael J D). Oxford University Press (2004).

25.  Satterlee, J.W.*, et al.* A Wox3-patterning module organizes planar growth in grass leaves and ligules. *Nat Plants* **9**, 720-732 (2023).

26.  Kaplan, D. R. The concept of homology and its central role in the elucidation of plant systematic relationships. In: *Cladistics: Perspectives on the Reconstruction of Evolutionary History* (eds Duncan T, Stuessy TF). First Edition edn. Columbia University Press (1984).

27.  Xu, X. *et al.* Single-cell RNA sequencing of developing maize ears facilitates functional analysis and trait candidate gene discovery. *Dev Cell* **56**, 557-568 e556 (2021).

28.  Takacs, E. M. *et al.* Ontogeny of the maize shoot apical meristem. *The Plant Cell* **24**, 3219-3234 (2012).

29.  Poethig, R. S., & Szymkowiak, E. J. Clonal analysis of leaf development in maize. *Maydica* **40**, 67–76 (1995).

30.  Smith, L. G. et al. Expression of knotted1 marks shoot meristem formation during maize embryogenesis. *Developmental Genetics* **16**, 344-348 (1995).

31.  Vollbrecht, E. et al. Shoot meristem size is dependent on inbred background and presence of the maize homeobox gene, knotted1. *Development* **127**, 3161-3172 (2000).

32.    Nardmann, J. et al. The maize duplicate genes narrow sheath1 and narrow sheath2 encode a

conserved homeobox gene function in a lateral domain of shoot apical meristems. *Development*

**131**, 2827-2839 (2004).

33.    Wu, H. et al. Genetic analyses of embryo homology and ontogeny in the model grass *Zea mays* L.

subsp. mays. *New Phytol*, *in press*.

34.    Johnston, R. et al. Transcriptomic analyses indicate that maize ligule development recapitulates

gene expression patterns that occur during lateral organ initiation. *The Plant Cell* **26**, 4718-4732

(2014).

35.    Strable, J*, et al.* Maize YABBY genes drooping leaf1 and drooping leaf2 regulate plant

architecture. *The Plant Cell* **29**, 1622-1641 (2017).

36.    Toriba, T. *et al.* BLADE-ON-PETIOLE genes temporally and developmentally regulate the

sheath to blade ratio of rice leaves. *Nat Commun* **10**, 619 (2019).

37.    Evans, M. M. The indeterminate gametophyte1 gene of maize encodes a LOB domain protein

required for embryo Sac and leaf development. *The Plant Cell* **19**, 46-62 (2007).

38.    Leiboff, S. et al. Network analyses identify a transcriptomic proximodistal prepattern in the maize

leaf primordium. *New Phytol* **230**, 218-227 (2021).

39.    Fukushima, K. & Hasebe, M. Adaxial-abaxial polarity: the developmental basis of leaf shape

diversity. *Genesis* **52**, 1-18 (2014).

40.    Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network analysis.

*BMC Bioinformatics* **9**, 559 (2008).

41.    Langfelder, P. & Horvath, S. Eigengene networks for studying the relationships between co-

expression modules. *BMC Syst Biol* **1**, 54 (2007).

42.    Knauer, S. *et al.* A high-resolution gene expression atlas links dedicated meristem genes to key

architectural traits. *Genome Res* **29**, 1962-1973 (2019).

43.    Morabito, S. et al. hdWGCNA identifies co-expression networks in high-dimensional

transcriptomics data. *Cell Rep Methods* **3**, 100498 (2023).

44.    Hofmann, F. et al. The embryonic transcriptome of *Arabidopsis thaliana*. *Plant Reprod* **32**, 77-91

(2019).

45.    Domazet-Loso, T. et al. A phylostratigraphy approach to uncover the genomic history of major

adaptations in metazoan lineages. *Trends in Genetics* **23**(11), 533–539 (2007).

46.    Ortiz-Ramírez, C. et al. A transcriptome atlas of *physcomitrella patens* provides insights into the

evolution and development of land plants. *Mol Plant.* **9**(2), 205-220 (2016).

47.    Spurr, A. R. Histogenesis and organizationof the embryo in Pinusstrobus L. . *Am J Bot* **36**, 629-

641 (1949).

48.    Richardson, A. E. *et al.* Evolution of the grass leaf by primordium extension and petiole-lamina

remodeling. *Science* **374**, 1377-1381 (2021).

49.    Gilbert, S. F. *Developmental Biology*, 6th edn. Sunderland (MA): Sinauer Associates (2000).

50.    Scanlon, M. J. & Freeling, M. The narrow sheath leaf domain deletion : a genetic tool used to

reveal developmental homologies among modified maize organs. *Plant Journal* **13**, 547-561

(1998).

51.    Satterlee, J. W. & Scanlon, M. J. Protoplast isolation from undifferentiated maize seedling shoot

tissue. *Methods Mol Biol* **2464**, 123-130 (2022).

52.    Schnable, P. S. *et al.* The B73 maize genome: complexity, diversity, and dynamics. *Science* **326**,

1112-1115 (2009).

53.    Hao, Y. *et al.* Dictionary learning for integrative, multimodal and scalable single-cell analysis.

*Nat Biotechnol* **42**, 293-304 (2023).

54.    Stuart, T. *et al.* Comprehensive integration of single-cell data. *Cell* **177**, 1888-1902 e1821 (2019).

55.    Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587 e3529

(2021).

56.    Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with Harmony. *Nat*

*Methods* **16**, 1289-1296 (2019).

57.    Scanlon, M. J. et al. Laser microdissection-mediated isolation and in vitro transcriptional

amplification of plant RNA. *Curr Protoc Mol Biol* **Chapter 25**, Unit 25A 23 (2009).

58.    Zhang, S. et al. Laser-capture microdissection of maize kernel compartments for rna-seq-based

expression analysis. *Methods Mol Biol* **1676**, 153-163 (2018).

59.    Kim, D. et al. Graph-based genome alignment and genotyping with HISAT2 and HISAT-

genotype. *Nat Biotechnol* **37**, 907-915 (2019).

60.    Kim, D. et al. HISAT: A fast spliced aligner with low memory requirements. *Nature Methods* **12**,

357-360 (2015).

61.    Wu, H. *et al.* NAKED ENDOSPERM1, NAKED ENDOSPERM2, and OPAQUE2 interact to

regulate gene networks in maize endosperm development. *The Plant Cell* **36**, 19-39 (2023).

62.    Love, M. I. et al. Moderated estimation of fold change and dispersion for RNA-seq data with

DESeq2. *Genome Biology* **15**, 550 (2014).

63.    Laureyns, R. *et al.* An *in situ* sequencing approach maps PLASTOCHRON1 at the boundary

between indeterminate and determinate cells. *Plant Physiol* **188**, 782-794 (2022).

64.    Jackson, D. In situ hybridization in plants. In: *Molecular Plant Pathology: A Practical Approach,*

(eds D. J. Bowles, S. J. Gurr, McPherson M). Oxford University Press (1991).

65.    Scanlon, M. J. et al. The maize mutant narrow sheath fails to establish leaf margin identity in a

meristematic domain. *Development* **122**, 1683-1691 (1996).

66.    Zhan, J. *et al.* RNA sequencing of laser-capture microdissected compartments of the maize kernel

identifies regulatory modules associated with endosperm cell differentiation. *The Plant Cell* **27**,

513-531 (2015).

67.    Drost, H. G. et al. Evidence for active maintenance of phylotranscriptomic hourglass patterns in

animal and plant embryogenesis. *Mol Biol Evol* **32**, 1221-1231 (2015).

68.    Vuruputoor, V. S. et al. Crossroads of assembling a moss genome: navigating contaminants and

horizontal gene transfer in the moss *Physcomitrellopsis africana*. *G3 Genes|Genomes|Genetics*,

jkae104 (2024).

69.    Katoh, K. et al. MAFFT version 5: improvement in accuracy of multiple sequence alignment.

*Nucleic Acids Res* **33**, 511-518 (2005).

70.    Suyama, M. et al. PAL2NAL: robust conversion of protein sequence alignments into the

corresponding codon alignments. *Nucleic Acids Res* **34**, W609-612 (2006).

71.    Zhang, Z. KaKs_Calculator 3.0: Calculating selective pressure on coding and non-coding

sequences. *Genomics Proteomics Bioinformatics* **20**, 536-540 (2022).

72.    Chen, J. *et al.* Dynamic transcriptome landscape of maize embryo and endosperm development.

*Plant Physiol* **166**, 252-264 (2014).

1071

**Acknowledgements**

We thank L. Evans  and R. Ragas for helpful discussions of the data and comments on the

manuscript. Special thanks to Cornell Genomics Facility for performing single-cell and Visium™

sequencing, to the Cornell Transcriptional Regulation and Expression Facility for RNA

sequencing, and the Cornell Guterman Bioclimatic Labs for our plants. Thanks to the Integrated

Genomics Operation (IGO) of Memorial Sloan Kettering Cancer Center for performing Xenium

assays. H. Wu was supported by a grant from the National Science Foundation (IOS-2016021).

R. Zhang is supported by another grant from the National Science Foundation (IOS-2210259).

1080

**Author contributions**

1082     Author Contributions M.J.S., H.W., R.Z., and K.J.N. conceived the study. H.W., M.J.S., and

1083     R.Z. generated and analyzed the data. M.J.S, H.W., R.Z. and K.J.N. interpreted the results, and

1084     M.J.S, H.W., R.Z., and K.J.N. wrote and edited the manuscript.

1085

1086

**Ethics declaration**

1088     Competing interests
1089     The authors declare no competing interests.

1090

**Corresponding author**

1092     Michael J. Scanlon: mjs298@cornell.edu

1093     Hao Wu: haowu2021@gmail.com

1094

1095 **Figures and Supplementary Figures**

0,3,8,11,13 Scutellum internal tissue
1,2,4,5,7,14 Scutellum epidermis
6 SAM and hypocotyl
9,15 Vasculature
Coleoptile epidermis
Embryo-endosperm boundary
Boundary of lateral organs
Coleoptile internal tissue
Scutellum tip
Apical scutellum or suspensor
Endosperm (residue)
Leaf
Suspensor

1096

**Fig. 1 Multiplexed spatial transcriptomics of the maize Stage 1 embryo.**

Differential interference contrast (DIC) microscopy of maize embryos at the Proembryo stage

(**a**), Transition stage (**b**) Coleoptilar stage, (**c**) and Stage 1 (**d**). Immunohistological analyses of

the Class I KNOX proteins mark the shoot apical meristem (SAM). ep, embryo proper; en,

endosperm; su, suspensor; sc, scutellum; c, coleoptile; L, leaf; *, SAM. Scale bar: 50 μm (**a** and

**b**) and 100 μm (**c** and **d**). **e**. Uniform Manifold Approximation and Projection (UMAP) of Stage1

embryo single-cell transcriptomic profiling with cell/tissue types assigned for each cell cluster.

1104

1105

**Fig. 2 Spatial transcriptomics of maize Stage 1 embryo**

Longitudinal cryosections of Stage 1 embryos embedded in OCT (optimal cutting temperature

compound; **a-b**) or in paraffin (**c-d**) for 10x Visium$^{TM}$ spatial transcriptomics analysis. **a, c**

Toluidine Blue-stained sections for 10x Visium$^{TM}$ spatial transcriptomics assay are overlayed

with the10x Visium$^{TM}$ spatial map (**b, d**). The various color-coded circles correspond to the

capture spots where spatial cDNA libraries were generated. Scale bar = 100 μm. **e.** Longitudinal

sections of paraffin-embedded Stage 1 embryo for use in laser microdissection RNA-seq analysis

(LM-RNAseq). **f.** Enlarged view of a Stage 1 embryo showing regions microdissected for LM-

RNAseq). Figure labels: scu, scutellum; uc, upper coleoptile; lc, lower coleoptile. **e, f** Scale

bar=100 μm. **g**. UMAP of the Visium$^{TM}$ spatial transcriptomic analysis of Stage 1 embryos cell

clusters. The spatial, embryonic origins of each cluster are indicated by the color-coded embryo

cartoon;  numbers correspond to the cluster designations on the UMAP. **h**. Heatmap illustrating

the correlation between the single-cell (sc) clusters (y-axis) and Visium$^{TM}$ spatial clusters

(x=axis). The significance level is marked by $-\log_{10}$p. **i**. Expression of candidate genes identified

in microdissected embryonic organs. Cell clusters exhibiting elevated candidate gene expression

of the specific markers are indicated; expression values are $\log_2$ transformed, i.e. $\log_2$ (TPM+1).

**Fig. 3 RNA-targeting analyses of Transition-staged, Coleoptilar-staged, and Stage 1**

**embryos**

**a-c.** Signal point map of *KN1, CUC2, NS1, WOX3A* and *DRL1* at the Transition stage (**a**),

Coleoptilar stage (**b**) and Stage 1 (**c**). **d-h.** Expression density map of *KN1* (**d**)*, CUC2* (**e**)*, NS1*

(**f**)*, WOX3A* (**g**) and *DRL1* (**h**) at Stage 1. **i-k, l-n, o-q** and **u-w**. Signal point map of *BOPa* (**i-k**),

*YAB14* (**l-n**), *IG1* (**o-q**), and *ARF3* (**u-w**) at the Transition stage, Coleoptilar stage and Stage 1. **r-**

**t** and **x-z**. Expression density map of *IG1* (**r-t**) and *ARF3* (**x-z**) at the corresponding stages.

Figure labels: T, Transition stage; C, Coleoptilar stage; L: Stage 1. The point colors are

arbitrarily determined. Yellow asterisks mark the adaxial expression of *IG1* and *ARF3*, and red

asterisks mark the abaxial expression of *ARF3*. The green asterisk in panel **k** marks the

expression maximum of *BOPa* at the emerging leaf sheath. The yellow asterisks in panel **r-t** and

**z** mark the expression maxima of *IG1* at the adaxial side of the scutellum (**r**), the coleoptile (**s**),

the coleoptile and the leaf (**t**), as well as the expression maxima of *ARF3* at the adaxial side of

the scutellum (**z**), respectively. The red asterisks in panel **x-z** mark the abaxial expression of

*ARF3* at the scutellum (**x, y**), the coleoptile (**y, z**) and the leaf (**z**). Scale bar for all panels = 50

μm.

1142

**Fig. 4 Dynamics of the maize embryo organ initiation network**

**a**. A Venn diagram of embryonic transcriptomics identifies 130 genes in the embryo-organ-

initiation-network. **b**. Normalized expression of these 130 genes at the Proembryo stage (PE),

Transition stage (T), Coleoptilar stage (C), Stage 1 scutellum (Scu), Stage 1 upper coleoptile

(UC), Stage 1 lower coleoptile (LC), Stage 1 leaf (L), as well as P0 and P1 of the 14 day-after-

germination seedlings. The letters above the boxplots represent significance levels (non-

overlapping letters indicate significant differences at $p<0.05$) using the Tukey-Kramer HSD test.

**c**. WGCNA tree graph of transcriptomic relationships among organ-specific co-expression

modules. The height (y-axis) scales the length of organ-specific branches, indicating the

transcriptomic distances between modules. **d**. Heatmap illustrating correlations between single-

cell (sc) clusters (y-axis) and hdWGCNA modules (x=axis). Significance levels are marked by

$-\log_{10}p$. **e**. Heatmap illustrating dynamic correlations of the initiation network expression in

Maize (y-axis) and Arabidopsis (x-axis) during embryogenesis. Heatmap colors are determined

by correlation coefficient (r) via Pearson correlation test with significance level $p<0.05$ (*) or

$p<0.01$ (**). Scu, Scutellum; Col, coleoptile; hypoc, hypocotyl.

1160

**Fig. 5 Phylotypic analyses of maize embryogenesis**

**a**. Phylostratigraphic map of maize. All maize protein-coding transcripts were assigned to 13

phylostrata (PS1-PS13). Parentheses designate the number of genes per phylostratum, and

the taxonomic groupings are marked above the corresponding PS. **b-c**. The transcriptome age

index (TAI; **b**) and transcriptome divergence index profiles (TDI; **c**) of maize embryogenesis

and seedling leaves. Figure labels: PE, proembryo stage; T, transition stage; C, coleoptilar

stage; L, Stage 1 leaf; Scu, Stage 1 scutellum; Col, Stage 1 coleoptile; P0 and P1, Plastochron

stages of the 15th and 14th leaves in the 14-day after planted seedlings. The gray lines

represent the standard deviation estimated by bootstrap analysis as described[22]. The overall

patterns of TAI and TDI profiles are significant, as measured by flat line test ($P_{TAI} = 3.56\times10^{-}$

$^{6}$, $P_{TDI} = 4.80\times10^{-4}$), reductive hourglass test ($P_{TAI} = 1.49\times10^{-4}$, $P_{TDI} = 1.53\times10^{-3}$), and

reductive early conservation test ($P_{TAI} = 1$, $P_{TDI} = 0.998$).

1174

1175

1176

1177

1178

1179

1180

**Fig. 6 Across-phylum phylotypic analyses of *Physcomitrium patens*.**

**a.** Development of embryo/sporophyte in the moss *P. patens*.  **b**. Phylostratigraphic map of *P.*

*patens*. All *P. patens* protein-encoding transcripts were assigned to one of five phylostrata (PS1-

PS5). Parentheses refer to the number of genes per phylostratum; taxonomic groupings are noted.

**c**. The transcriptome age index (TAI) and transcriptome divergence index (TDI) profiles of *P.*

*patens* archegonia; sporophyte stages S1, S2, S3 and SM. Colored bars illustrate the duration and

timing of different S stages along the 30 day time scale. **d**. Comparative expression levels of

stage-specific embryonic genes between *Arabidopsis* and *P. patens,* and maize and *P. patens*.

Figure labels: (in moss) AR, archegonium; S1, sporophyte 1 stage; S2, sporophyte 2 stage; S3,

sporophyte 3 stage; SM, mature sporophyte stage; (in maize) PE, proembryo stage; T, transition

stage; C, coleoptilar stage; L, Stage 1 leaf; Scu, Stage 1 scutellum; Col, Stage 1 coleoptile; S2,

Stage 2. The gray lines represent the standard deviation estimated by bootstrap analysis as

described[22]. The overall patterns of TAI and TDI profiles are significant, as measured by the flat

line test ($P_{TAI} = 0.0131$, $P_{TDI} = 4.73 \times 10^{-3}$), reductive hourglass test ($P_{TAI} = 0.0332$, $P_{TDI} =$

0.0249), and reductive early conservation test ($P_{TAI} = 0.998$, $P_{TDI} = 0.973$). Images in (**a**) and (**c**)

are reproduced from Ortiz-Ramirez et al., 2016.

1197

1198

1199

1200

1201

1202

1203

1204

1205

1206

**Supplementary Fig. 1 Verification of single-cell cluster marker genes.**

Left panel: single-cell cluster marker expression projected as a UMAP. Purple dots mark cells

with corresponding gene-specific expression. Middle panel: verification of single-cell cluster

marker genes by Visium[TM]. Red to green spots mark the locations of gene expression with high

to low level, respectively. Right panel: verification of single-cell cluster marker genes by

multiplexed RNA-targeting *in situ* hybridization (data not available for *GRMZM2G166141* (**k**),

*GRMZM2G094353* (**m**) and *GRMZM2G099454* (**p**)). Scale bar = 100 μm.

1214

1215

1216

**Supplementary Fig. 2 Differential expression between laser microdissected RNAseq**

**samples.**

**a**. Principal component analysis of laser-microdissected RNAseq samples. Each dot represents a

laser-microdissected embryonic shoot sample. **b-f**. Pairwise comparison of the gene expression

between laser-microdissected RNAseq samples. The number of overlapping, differential-

expressed genes (DEGs) is showed in the Venn diagram. The circled numbers represent the

genes preferentially expressed in the SAM (**b**), upper coleoptile (**c**), scutellum (**d**), leaf (**e**) and

lower coleoptile (**f**).

1225

1226 **Supplementary Fig. 3 Multiplex *in situ* hybridization of *TRU1*.**

1227 Spatio-temporal expression of *TRU1* displayed as a signal point map (**a-c**) and as a transcript

1228 abundance correlated signal density map (**d-f**) at the transition stage (**a and d**), coleoptilar stage

1229 (**b and e**) and Stage 1 (**c and f**). The red arrow in (f) marks the expression of *TRU1* at the

1230 emerging leaf sheath of leaf 1. Scale bar = 100 μm.

1231

1232

1233

1234

1235

1236

1237

1238

1239

1240

1241

1242

1243

**Supplementary Fig. 4 Expression of the embryo organ initiation network.**

**a**. WGCNA identifies modules with overall highest expression at Stage 1 leaf (L), less in the

lower coleoptile (LC), and steadily decreased expression in the upper coleoptile (UC), scutellum

(S). Heatmap shows the correlation coefficients (r) and the corresponding p values (parentheses)

of modules ME1-12. **b**. Expression of the 130 genes comprising the embryonic-organ-initiation

network at the transition stage (T), coleoptilar stage (C) and the Stage 1 scutellum (Scu), upper

coleoptile (UC), lower coleoptile (LC) and leaf 1 (L). **c-e**. Correlations of the expression of these

130 genes during initiation of the scutellum, coleoptile, and leaf 1.

1252

1253

1254

1255

1256

1257

1258

1259

1260

1261

1262

1263

1264

1265

1266

1267

**Supplementary Fig. 5 WGCNA Module-organ correlation.**

**a-b**. Correlations of transcriptomic datasets from embryonic organs and leaves initiated post-germination before and after batch effect removal. Correlations of the normalized overall gene expression between the post- germination seedling leaves and embryonic stages before (**a**) and after (**b**) batch effect removal. **c**. Heatmap highlights correlations between co-expression modules (y-axis) and corresponding organs (x-axis). Colors are graded based on correlation coefficients (r) and p values (in parentheses).

**Supplementary Fig. 6 Genetic analyses of *log7* mutant embryos.**

Expression and genetic analysis of *LOG7*. **a.** Conventional *in situ* hybridization of *LOG7*

transcript accumulation in a longitudinal section of the Stage 1 embryo. Multiplexed RNA-

targeting point map (**b**) and density map (**c**) of *LOG7* accumulation in longitudinal sections of

Stage 1 embryos. Scale bar = 100 μm. **d .** Longitudinal median section of a WT Stage 2

embryo. The red dashed line marks the transverse section plane depicted in (**g**). **e-f.**

Longitudinal median sections of the *log7* mutant Stage 2 embryos. The red dashed line marks

the transverse section plane depicted in (**h**). Longitudinal off-median section of the *log7* mutant

Stage 2 embryo. All scale bars = 100 μm. Asterisks (*) mark the SAM; sc, scutellum; c,

coleoptile; L, leaf.

1291

1292

1293 **Supplementary Fig. 7 Correlation between the maize initiation network genes and the**

1294 **Arabidopsis orthologs.**

1295 The correlation of expression between the maize initiation network genes at transition (**a**) and

1296 coleoptilar (**b**) stage, and the Arabidopsis orthologs at early heart stage.

**Supplementary Fig. 8. TAI and TDI for embryo stages under different BLASTp E-value cutoffs**

**a-c**. Distribution of all maize genes and genes in each embryo stage across phylostrata and TAI profiles for different E-value cut-offs in BLASTp homology searches (ranging from 1E-2 to 1E-20). This analysis investigates whether and how different E-values affect the hourglass pattern. **a**. Variation of the E-value cut-off for the BLASTp analysis demonstrates minimal effects on the distribution of maize genes among the phylostrata, indicating an overall pattern independent of the E-values cut-offs. The y-axis is presented on a logarithmic scale. **b**. Similarly, the variation of the E-value cut-off for the BLASTp analysis does not severely alter the distribution of maize genes in each embryo stage among the phylostrata. **c**. The hourglass pattern of the TAI profile remains consistent across various E-value cut-offs. **d-e**. Transcriptome divergence indices based on two comparisons of genomes from three relatives of *Zea mays*: *Zea mays* ssp. Mays vs. *Zea mays* ssp. mexicana (**d**); *Zea mays* ssp. mays vs. *Zea diploperennis* (**e**). The overall patterns of TDI profiles are significant, as measured by the flat line test ($P_{\text{TDI-Mexicana}}=1.07\times10^{-3}$, $P_{\text{TDI-diploperennis}}=4.80\times10^{-4}$), reductive hourglass test ($P_{\text{TDI-Mexicana}}=0.027$, $P_{\text{TDI-diploperennis}}=1.53\times10^{-3}$), and reductive early conservation test ($P_{\text{TDI-Mexicana}}=0.68$, $P_{\text{TDI-diploperennis}}=0.998$).