A multiplexed transcriptomic analysis of a plant embryonic hourglass

Hao Wu[1*], Ruqiang Zhang[1], Michael J. Scanlon[1*]

[1]School of Integrative Plant Science, Cornell University, Ithaca, NY, USA

[*] Corresponding authors

Abstract

The embryo defines the primordial developmental stage of multicellular organisms, from

fertilization of the zygote to the formation of tissue layers and organs. Animal zoologists [1, 2, 3]

described a period of maximum morphological similarity between embryos of different species

within related taxa, later termed the phylotypic stage[4] of mid-embryogenesis. Modern

transcriptomic studies identified a phylotypic pattern of gene expression in Drosophila,

zebrafish, and the flowering plant Arabidopsis, featuring enhanced accumulation of

evolutionarily older and conserved transcripts during mid-embryogenesis[5, 6, 7]. Such plots of

transcriptomic age and divergence conform to the "hourglass models" of morphological

similarities described by 19[th] century embryologists. Chronological descriptions of the

phylotypic stage are problematic for plants, where embryogenesis extends throughout the life

cycle and the mid-point of seed maturation is morphologically and chronologically variable

among taxa[7]. Here we use multiplexed spatial-transcriptomic analyses to investigate the

homology of the grass embryo, which evolved ~ 150 million years since the divergence of

eudicots and grasses[8]. Our data support a model where the single maize embryonic leaf

(cotyledon) is a novel, bi-partite organ derived within the monocotyledonous grasses, after the

evolution of the ancestral angiosperm cotyledon from a modified embryonic leaf [9, 10]. We

describe a maize embryonic-organ-initiation genetic-network that uses ancient and conserved

genes, comprising a phylotypic pattern of "mid embryogenesis" during the ancestrally conserved

initiation of histogenesis and organogenic growth in plant and animal embryos.

Introduction

Following fertilization and formation of the diploid zygote, embryogenesis in flowering plants

comprises three widely conserved stages[11]: (1) polarized and asymmetric cell divisions to create

the embryo proper; (2) patterning of discrete histological tissue-layers; and (3) organogenesis of

lateral organs alongside the formation of pluripotent meristems. Persistent stem-cell reservoirs

comprising the shoot apical meristem (SAM) and root apical meristem (RAM) enable plants to

extend organogenesis long-beyond embryonic stages. Seed maturation and quiescence, the onset

of which is widely-variable and correlated with the mode of seed dispersion, is a commonly-

utilized but not universally-conserved strategy of plant embryogenesis[11, 12].

Embryos from the eudicot *Arabidopsis thaliana* and the monocot grass *Zea mays* ssp. *mays* both

use asymmetric cell divisions to establish the embryo proper that will generate the body plan and

the suspensor (Fig. 1a), a temporary supporting structure that is later consumed by programmed

cell death[13]. Histological layering of the globular embryo proper forms an outer protoderm

surrounding layers of internal tissue(s) in grass and eudicot embryos[14, 15, 16, 17]. Thereafter,

Arabidopsis embryos develop paired, embryonic leaves called cotyledons and organogenic

meristems in the shoot and root poles, whereafter organogenesis is suspended until seed

germination[10, 11]. In contrast, the morphology of the grass embryo is dissimilar to that of

Arabidopsis. The first lateral organ formed in the maize embryo is the scutellum (Fig. 1b), which

at maturity forms a shield-shaped, glandular, non-photosynthetic structure that absorbs nutrients

from the germinating seed[13, 17]. Subsequently, the coleoptile emerges as the second maize

embryonic structure (Fig. 1c), and later forms a cylindrical sheath that protects the underlying,

fragile shoot during seedling germination. Unlike Arabidopsis, grasses extend embryonic

organogenesis to form several foliar leaves before seed quiescence. Upon seed germination,

lateral organ production resumes at the SAM, and continues throughout the life cycle[14, 15]. The

homology of the single cotyledon in monocotyledonous grasses is an unresolved topic of

botanical debate[10, 18, 19, 20].

Result

Single-cell profiling of the transcriptomic landscapes in Stage 1 maize embryo

Historically, homology is inferred on the basis of developmental morphology and/or ontogenetic

expression of conserved genetic networks[21, 22]. Stage 1 maize embryos are approximately 1 mm

long and contain a prominent, ovately-flattened scutellum, an emerged coleoptile that is

beginning to surround the SAM, and the newly-initiated primordium of the first foliar leaf[15] (Fig.

1d). 10X Genomics Chromium™ was used to profile the single-cell transcriptomes of 11,467

protoplasts digested from over 400 Stage 1 embryos in two replicates (Fig. 1e). Twenty-three cell

clusters were identified; median values of 10,594 UMI per cell and 4,030 genes per cell were

obtained.

A first challenge encountered in analyses of sc-RNAseq data is the identification of the

tissue/organ-specific origin of each cell-cluster. Since prior RNA-seq analyses in maize have

focused on seedlings or inflorescences[23], cell-clusters derived from the SAM, embryonic stem,

and vasculature[24] are readily identified, as is the residual starchy endosperm (Fig. 1e; Clusters 6,

15 and 20 respectively). Relatively few studies have analyzed the transcriptomic patterns within

the grass scutellum or coleoptile, such that the origins of several cell-clusters were not

immediately identifiable. To annotate the UMAP cell-clusters, we utilized a multiplexed

transcriptomic approach combining these sc-RNAseq data with spatial-transcriptomics and laser-

microdissection Ranse (LM-RNAseq) to obtain spatial information for all the cell-clusters, as

described below (Fig. 2; Extended Data Fig. 1-2, Extended Data Table 1).

Eight individual Stage 1 embryo sections were processed using the 10X Genomics Visium[TM]

spatial transcriptomics pipeline (Fig. 2a-d). Nine spatially resolved cell-clusters were identified

and categorized into four supergroups: (1) shoot-root axis; (2) scutellum epidermis; (3) internal

scutellum; and (4) endosperm/embryo boundary (Fig. 2e). Hypergeometric testing analyzed the

overlap of cluster marker genes between the single-cell and spatial-transcriptomic data,

suggesting that 14 of the 23 single-cell clusters (0-5, 7-9, 11, 13, 14, 18 and 19) are derived from

the scutellum (Fig. 2h). Comparisons of transcript localizations of known marker genes within

the UMAP and the spatial-transcriptomic plots suggest that the resolution of Visium[TM]

technology may comprise 5-10 embryonic cells. Thus, it was not yet possible to spatially-

distinguish transcripts among the coleoptile, the SAM, and leaf1. To increase the resolution of

our embryo transcriptomics, LM-Ranse was utilized in micro-dissected transcriptomic analyses

of the SAM, adaxial scutellum, upper/lower coleoptile, and leaf primordium in Stage 1 embryos

(Fig. 2f-g). Principal component analysis (PCA) shows that the transcriptomic profiles of these

five embryonic organ/structures are well-separated (Extended Data Fig. 2a). In particular, 9,913

genes are identified as differential-expressed in at least one embryonic structure, and there are

89, 1058, 246, 84 and 53 genes preferentially expressed in these same five embryonic regions

(Extended Data Fig. 2b-f). Thus, this multiplexed transcriptomic approach enabled identification

of the tissue/organ origin of each cell-cluster in the maize Stage 1 embryonic UMAP (Fig. 1e;

Fig. 2i).

Temporal and spatial analysis of gene expression via multiplexed RNA-targeting

Homologous organs utilize homologous genes regulating developmental pattern[21, 25]. With an

emphasis on candidate regulatory genes differentially expressed in our transcriptomic analyses of

Stage 1 embryos, we targeted 100 maize transcripts for temporal and spatial analyses of

multiplexed transcript *in-situ* localization using the 10X Xenium[TM] protocol on longitudinal

sections of (i) Transition-staged embryos just forming a SAM and scutellum (Fig. 1b), (ii)

Coleoptilar-staged embryos (Fig. 1c), and (iii) Stage 1 embryos initiating a leaf (Fig. 1d). We

first examined the correlations of our sc-RNAseq data, spatial-transcriptomic plots, and

Xenium[TM] *in situ* RNA-targeting data via comparisons of 16 genes preferentially expressed in

one or two single-cell clusters (Extended Data Fig. 1). In all cases examined, the *in situ*

hybridization patterns are consistent with the spatial expression data. Satisfied with the

consistency of these multiplexed *in situ* data, we next analyzed the dynamic co-localization

patterns of some key regulators of maize shoot development during serial stages in grass embryo

ontogeny.

Consistent with prior studies[26, 27], the spatial-temporal expression of the indeterminacy marker

KNOTTED1 (*KN1*; Fig. 3a-d) is first-detected during SAM initiation at the Transition stage (Fig.

1b). At later stages, *KN1* accumulates in the SAM and subtending embryonic stem, but is

downregulated in lateral organs[26, 27]. *NARROW SHEATH1* (*NS1*) and *WUSCHEL-RELATED*

HOMEOBOX 3A (*WOX3A*) encode homeodomain transcription factors required for mediolateral

expansion of maize lateral organs[25], and are co-expressed at edges of the developing coleoptile

and initiating leaf primordia as described previously[17, 21, 28] (Fig. 3b-c, f-g). Undetected in prior

studies of embryo ontogeny, Fig. 3a reveals *NS1* transcript accumulation in the initiating

scutellum at the Transition stage. *CUP-SHAPED COTYLEDON2* (*CUC2*) marks developmental

boundaries throughout maize ontogeny[29]; *CUC2* transcripts are detected in-between the

expression domains of *KN1* and *NS1* during scutellum initiation (Fig. 3a), and in the boundaries

between the SAM and later-staged embryonic lateral organs (Fig. 3b-c,e). *DROOPING LEAF1*

(*DRL1*) regulates leaf architecture and is expressed in incipient and developing foliar leaf

primordia[30] but is not detected in the coleoptile or scutellum at any embryo stage examined (Fig.

3c,h). These combinatorial *in situ* RNA-targeting analyses enable identification of previously

undescribed combinatorial patterns of dynamic, marker gene expression during maize embryo

development.

Other candidate genes analyzed by Xenium[TM] displayed patterns of transcript accumulation

during embryo ontogeny that offer insights into the developmental homologies of maize

embryonic lateral organs. For example, *BLADE-ON-PETIOLEa* (*BOPa*) accumulates throughout

the developing coleoptile and in the proximal region of the initiating leaf at Stage 1 (Fig. 3i-k),

consistent with the proposed function of *BOP* homologs as leaf proximal-identity genes[31].

Notably, *BOPa* is predominantly undetected throughout scutellum ontogeny; similar

accumulation patterns are noted for the *BOPa* paralog *TASSELS REPLACE UPPER EARS1*

(*TRU1*) (Extended Data Fig. 3). Transcripts of the *DRL1*-paralog *YABBY14* (*YAB14*) are densely

accumulated in the apical tip of the scutellum at the transition stage and in the elongating

scutellum and newly emerged coleoptile at the coleoptilar stage. During Stage 1, a new *YAB14*

expression maximum is organized at the primordial leaf tip (Fig. 3l-n). Moreover, the pleiotropic

gene *INDETERMINATE GAMETOPHYTE1* (*IG1*) is implicated to function during leaf polarity

and patterning of maize leaves[32, 33]. *IG1* accumulates on the top (i.e. adaxial) surfaces of

initiating leaf primordia during Stage 1, but also in adaxial layers of the emerging coleoptile and

in the upper-adaxial region of the transition-staged scutellum (Fig. 3o-t) No significant *IG1*

expression is noted during later stages of embryonic scutellum development (Fig. 3p-q, s-t).

Another leaf polarity regulator, *AUXIN RESPONSE FACTOR 3* (*ARF3*)[34], is expressed on the

abaxial region of the transition-staged embryo during scutellum initiation, and in the abaxial

domains of the newly-emerged coleoptile and leaf during later embryo stages (Fig. 3u-z).

Interestingly, the expression of *ARF3* in the scutellum moves from abaxial domains in transition

stage to adaxial region at Stage 1 (Fig. 3z).

The initiation of maize embryonic lateral organs utilizes a shared genetic network

Homologous organs are inferred to utilize evolutionarily-conserved developmental genetic

networks[22]. By Stage 1, the three grass embryonic lateral organs have already begun to diverge

morphologically, and will each diverge further before seed quiescence[15]. Interrogations of lateral

organ homology require investigation throughout embryo ontogeny[22]. We therefore combined

the LM-RNAseq data from the Stage 1 embryonic organs generated in the current study with our

previous LM-RNAseq data from serial stages of maize embryonic development[24], to generate a

more complete sampling of grass embryogenesis. Specifically, these prior analyses generated

transcriptomic data precisely from the newly initiated embryonic organ and the adjacent SAM,

which were serially sampled at the transition stage (scutellum +SAM), the coleoptilar stage

(coleoptile +SAM) and Stage 1 (leaf 1 +SAM). Analyses of pairwise differential gene expression

among the five embryonic organs reveals a patterned expression of 53 leaf-preferential genes in

Stage 1 embryos (Extended Data Fig. 2e), wherein their overall expression is highest in leaf 1 (L)

and less in the lower coleoptile (LC), with steadily decreased expression in the upper coleoptile

(UC) and scutellum (S) (i.e. L > LC > UC > S).

Notably, the expression of those leaf-preferential genes during the transition stage and the

coleoptilar stage is similar to that discovered in leaf 1 during Stage 1 (Extended Data Table 2)[24].

These data suggest that genes involved in foliar leaf initiation at Stage 1 are also associated with

the initiation of the scutellum and coleoptile at successively earlier stages in embryo ontogeny.

To further interrogate this putative maize embryonic-organ-initiation network, a weighted gene

co-expression network analysis (WGCNA)[35, 36] was performed on our Stage 1 embryonic LM-

RNAseq data to identify co-expression modules correlated with the expression pattern L > LC >

UC > S. A total of 5,797 gene transcripts with high expression variability were selected; all

except one non-clustered transcript (MEgrey module) were clustered into twelve co-expression

modules (ME1 through ME12). Two co-expression modules (ME2 and ME3) comprising 963

genes emerged as correlated (Extended Data Fig. 4a; Fig. 4a). Intersection of these 963 genes

with transcripts previously found via LM-RNAseq of the transition-staged apex[24] (excluding

putative SAM-specific genes) identified 130 genes comprising the maize embryonic-organ-

initiation network (Fig. 4a, Extended Data Fig. 4b). Expression of these 130 network genes is

significantly and positively correlated with the initiation of the transition-staged scutellum, the

coleoptilar-staged coleoptile, and the Stage 1 leaf and lower-coleoptile (Extended Data Fig. 4c-

e). During Stage 1, the expression levels of this embryonic-organ-initiation network significantly

decrease in the scutellum and upper coleoptile, but not in the lower coleoptile or leaf 1 (Extended

Data Fig. 4b, Fig. 4b). Gene ontology (GO) enrichment analyses showed that these network

genes are replete for the GO term Organ Development, whereas the transcriptomic networks of

the Stage 1 scutellum and coleoptile are enriched for the GO terms Carbohydrate and Lipid

Metabolism, and Organ Development/Lipid Metabolism, respectively (Fig. 4c). The data

suggests that during initiation, all maize embryonic lateral organs utilize a similar genetic

network, the expression of which decreases in later embryonic stages as the scutellum,

coleoptile, and leaf differentiate to perform distinct functions in later plant development (Fig.

4d).

A central question in the controversial homology of grass embryonic organs is the identity of the

coleoptile as either a foliar leaf[18], the entire grass cotyledon[19], or the proximal component of a

bipartite, single grass cotyledon[20]. A second WGNCA analysis was performed on the LM-

RNAseq data obtained from the Stage 1 embryonic shoot apex. In addition, we integrated

published LM-RNAseq data[37] of the Plastochron 0 (P0) and P1 leaf from 14-day after planting

maize seedlings as a reference for later leaf ontogeny. Batch effects were removed (Extended

Data Fig. 5a,b), and the merged dataset was filtered to select genes with variability in transcript

accumulation levels across embryonic organs. In the analysis, 5,128 out of 5,137 selected genes

were clustered into 15 co-expression modules (LM1, LM2, …, LM15), whereas the remaining 9

transcripts resolved to an unclustered module (Extended Data Fig. 5c; Fig 4d).

Pearson correlation tests between module eigengenes and organ-specific transcriptomes

identified associations between specific co-expression modules and the corresponding embryonic

organs (Fig. 4d, Extended Data Fig. 5c). Transcriptomic relationships among the co-expression

modules are reflected in a tree graph (Fig. 4d), in which the y-axis (height) scales the length of

organ-specific branches, indicating transcriptomic distances between these modules (Fig. 5a).

Two major branches comprise (1) the Stage 1 leaf, SAM, P0 and P1, and (2) the coleoptile and

the scutellum.

To investigate the transcriptomic relationships among all Stage 1 maize embryonic lateral organs

at single-cell resolution, we then performed hdWGCNA[38] on the scRNAseq data. A total of

5,026 genes transcripts were clustered into 11 co-expression networks (M1, M2, …, M11). A

hypergeometric test analyzed the overlap between the putative single-cell cluster marker-gene

transcripts and those highly correlated with corresponding co-expression modules. The data

illustrate that the SAM and the leaf cluster marker-genes significantly overlap with the closely

related modules M9 and M10, respectively. In contrast, marker genes from the scutellum and the

coleoptile cluster overlap with multiple co-expression modules (M5, M7 and M8; Fig. 5b). In

summary, these hdWGCNA analyses of embryo scRNA-seq data suggest close transcriptomic

relationships between the initiating/expanding foliar leaves and the SAM, and between the

scutellum and the coleoptile.

The embryonic-organ-initiation genetic network stage is conserved in Arabidopsis embryos

To investigate the transcriptomic similarities of embryonic lateral organs of maize and those of

the eudicot model Arabidopsis, we used comparative genomic toolkits [GRAMENE

(https://www.gramene.org/), TAIR (https://www.arabidopsis.org/), BLAST tools

(https://blast.ncbi.nlm.nih.gov/Blast.cgi)] to select Arabidopsis orthologs of the 130 transcripts

identified to comprise the maize, embryonic-organ-initiation network, and interrogated the

temporal, gene expression atlas of Arabidopsis embryo development[39]. Nomenclature describing

embryo stages differs in eudicots and grasses[11]. In Arabidopsis, histological layering and SAM

initiation occur during the globular stage, cotyledon initiation and emergence defines the heart

stages, whereafter the cotyledons elongate during the torpedo and bent embryo stages.

Out of 130 genes in the maize embryonic-organ-initiation network, forty-nine have

corresponding Arabidopsis orthologs expressed during embryo development, commencing

during cotyledon initiation at the early heart stage and persisting until the bent cotyledon stage

(Supplementary Dataset; Extended Data Fig. 6). Notably, the stage-specific accumulation

patterns of these homologous maize transcripts in the emerging scutellum (Transition stage),

coleoptile (Coleoptilar stage), and foliar leaf (Stage 1) are significantly correlated with the

reported expression of the corresponding Arabidopsis orthologs during specific stages in

cotyledon initiation and development in this eudicot embryo[39] (Fig. 4f, Extended Data Fig. 6). In

addition, transcripts accumulating in the expanding Stage 1 scutellum correlate with expression

of Arabidopsis orthologs during later stages of cotyledon growth (i.e. late-torpedo stage[39]).

Likewise, maize transcripts accumulating in the elongating Stage 1 coleoptile are significantly

correlated with expression of orthologous Arabidopsis genes during earlier stages of cotyledon

expansion (i.e. early heart stage[39]) (Fig. 4f). These data suggest that despite the morphological

dissimilarities of these grass and eudicot "mid-staged" embryos, the development of the maize

and Arabidopsis embryonic lateral organs share a homologous regulatory network.

The embryonic organ initiation network comprises a phylotypic stage

We show that the maize embryonic-organ-initiation network is conserved in *Arabidopsis*

thaliana, where it correlates with the transcriptomic signatures of the eudicot heart and torpedo

stage embryos. Prior studies revealed that during mid-embryogenesis, old and conserved genes

are more highly expressed than young and divergent genes, marking the phylotypic stage of

Arabidopsis embryogenesis[7]. To investigate if the maize embryonic-organ-initiation network is

associated with a phylotypic stage, we analyzed the age (TAI) and divergence (TDI) of the genes

in the maize embryonic-organ-initiation network.

In detail, to evaluate gene age, we assigned a phylostratum to each protein-coding gene in the

maize genome (PS) by performing BLAST sequence alignment analyses on each protein

sequence in the embryonic-organ-initiation network to the protein sequences of 350 other

species, while searching for the most distant phylogenetic node containing at least one species

with a homologous protein domain.

We obtained 13 phylostrata, PS1-PS13 (Fig. 5a; Supplementary Dataset), from the oldest genes

with domains homologous to prokaryotic genes to the youngest genes with no homologs

identified in any other species. We utilized the transcriptome age index (TAI) to measure the

average gene age, adjusted by level of gene expression, at each developmental stage sampled. As

shown in Fig 5b and Extended Data Fig. 7a-c, the TAI undergoes a dramatic decrease beginning

at the proembryo stage and reaches a minimum value in the Stage 1 scutellum. This hourglass

pattern, characterized by low TAI values, continues through the Coleoptilar stage and Stage 1

followed by an increase at Stage 2 of embryogenesis and continued increased TAI values in the

newly initiated leaves of germinated maize seedlings (Fig. 5b).

To evaluate the gene sequence divergence, calculated the K_a/K_s ration between maize and its

sister species *Zea diploperennis*. K_a and K_s refer to the number of non-synonymous and

synonymous substitutions, respectively, between each orthologous pair. The transcriptome

divergence index (TDI) measures the average divergence of genes weighted, by transcript

accumulation level, at different developmental stages. Consistent with the TAI results, the TDI

also showed a reduction from the proembryo stage to the initiation of the first lateral organs at

the Transition and Coleoptilar stages, and until Stage 1, followed by a dramatic increase at Stage

2 and in germinated seedling leaf primordia (Fig. 5c; Extended Data Fig. 7d-e). Thus, measures

of both transcriptional gene age and gene sequence divergence show an "hourglass" pattern

during maize embryogenesis, indicating that the processes of tissue layer organization and lateral

organ initiation are associated with evolutionarily older and more conserved genes.

Discussion

A multiplexed transcriptomic approach was applied to investigate the developmental homologies

of grass embryonic organs. Spatial transcriptomics complements the ambiguities encountered in

cell-cluster identification that can complicate scRNA-seq analyses; the precision of LM-RNAseq

can differentiate the tissue/organ-specific transcriptomes of embryonic micro-structures that are

unresolved using spatial transcriptomics alone (Fig. 2). Moreover, high-throughput, multiplexed

in situ RNA-targeting technologies enable localizations of multiple transcripts simultaneously

and combinatorically, throughout embryo ontogeny. RNA-targeting analyses confirm the

expression of several key regulators comprising a transcriptomic network that functions at the

initiation of the grass scutellum, coleoptile, and foliar leaf (Fig. 3). Thus, the combinatorial usage

of these four transcriptomic approaches complements one another to build a high throughput,

highly resolved, and spatially informed transcriptomic profile of grass embryo development.

Although the evolution of the plant cotyledon from conifer and/or dicot ancestral leaves is still

unclear[10], genetic and morphological evidence support models wherein the cotyledon evolved via

modification of the first embryonic leaf[11, 40]. Comparisons of transcriptomic profiles from all

Stage 1 embryonic lateral organs identified a conserved embryonic-organ-initiation network,

suggesting that the scutellum, coleoptile, and leaf 1 are homologous organs (Fig. 4a; Extended

Data Fig. 4a-b). Accumulation of transcripts comprising this embryonic-organ-initiation network

is diminished during later developmental stages (Fig. 4b; Extended Data Fig. 4a-b), whereupon

each embryonic lateral organ is specifically modified to perform discrete functions during seed

germination[13, 17]. Organ-specific changes in gene expression during later embryonic stages are

observed in coleoptile and leaf primordia, wherein expression of the embryonic-organ-initiation

network decreases as these organs exercise differential gene-expression to form functionally

divergent organs.

Classical plant morphologists have proposed disparate models for the homology of maize

embryonic lateral organs[reviewed in 10]. One model suggests that the scutellum is homologous to the

cotyledon, and the coleoptile is a modified foliar leaf[18, 19]. Others proposed that the coleoptile

comprises the maize cotyledon, and the scutellum is an evolutionarily unique haustorium that

arose within the grasses[19]. A synthetic, bipartite hypothesis suggests that the scutellum and the

coleoptile together, comprise the single grass cotyledon, a derived character that arose within the

grasses. In this view the scutellum comprises the apical region, and the coleoptile forms the

sheathing base of the singular cotyledon[20].

Quality characteristics of Arabidopsis cotyledons include their unique status as the first-initiated

embryonic organs and their ultimate function in supporting the germinating seedling. The

scutellum is the first lateral organ initiated in grass embryos, whereas both the scutellum and

coleoptile satisfy the second criterion of seedling support at maturity, albeit in very different

ways[15, 41]. Transcriptomic analyses of Arabidopsis cotyledon development reveal significantly

correlated expression of orthologs of the maize embryonic-organ-initiation network, beginning

during embryo histogenesis in the globular-staged embryo and persisting until later-staged

cotyledon growth and expansion (Fig. 4f). Expression of the embryonic-organ-initiation network

during maize coleoptile and scutellum development at Stage 1 is significantly correlated with

that of mid-stages of Arabidopsis cotyledon development, suggesting that the Arabidopsis

cotyledon is homologous to the maize scutellum *and* coleoptile.

Our WGCNA analyses support a model where the maize cotyledon is a bi-partite embryonic

organ comprising the scutellum and the coleoptile (Fig. 4). The evolution of a single grass

cotyledon as a composite embryonic organ raises interesting questions. Grass phyllotaxy is

distichous; successive organs initiate ~180° apart and in two hierarchical ranks about the stem[41].

If the scutellum and coleoptile were separate embryonic organs initiated successively from the

embryonic axis, they are predicted to emerge from opposite sides of shoot. However, the

scutellum and coleoptile are both initiated on the same side of the grass embryo, although the

scutellum precedes the coleoptile[15] (Fig. 1a-d). Our composite data suggest that the scutellum

and coleoptile comprise different domains of the single grass cotyledon[42] (Fig. 6).

Foliar leaves of grasses contain a distal, photosynthetic blade and a proximal sheath that

surrounds and overlaps the stem[41]. A morphological model for grass cotyledon homology[19, 20]

proposes that the scutellum and coleoptile are homologous to the distal blade and the sheathing

base of the foliar leaf, respectively (Fig. 6). Consistent with this model, cells contributing to the

distal leaf blade emerge from the shoot apex prior to the emergence of leaf sheath progenitor

cells[25, 43, 44, 45]. This pattern mirrors the ontogenetic timing of scutellum and coleoptile initiation.

Furthermore, expression of the paralogous sheath marker genes *BOPa* and *TRU1* is restricted to

the maize leaf sheath[29, 46] and to the Stage 1 coleoptile (Fig. 3i-k, Extended Data Fig. 4), in

support of models for a bipartite grass cotyledon. Higher-order *Osbop* mutants delete the rice

leaf sheath[31], supporting hypotheses for BOP-mediated development of sheathing leaf bases.

Taken together, these data suggest that the coleoptile and the leaf sheath comprise the sheathing

bases of homologous lateral organs in grasses.

Our multiplexed transcriptomic analysis of maize embryogenesis identifies a network of genes

expressed during initiation of embryonic organs that comprises a phylotypic period. As

previously described in the eudicot plant Arabidopsis and in animal embryos of fruit flies

(*Drosophila melanogaster*) and zebrafish (*Danio rerio*)[5, 6, 7], the transcriptomic phylotypic period

is defined by the preferential expression of conserved and evolutionarily ancient genes (Fig 5b-

c). We note that this transcriptomic phylotypic period in maize embryogenesis commences when

tissue layers are established during the late proembryo stage and forms an extended hourglass

shape corresponding to successive initiations of the scutellum (T stage), coleoptile (C stage), and

leaf 1 (Stage 1). Striking decreases in both transcript age and conservation occur during later

stages of maize embryogenesis (Stage 2) and upon initiation of seedling leaves post-germination

(Fig. 5b-c). Considering the number of genes represented, most of the 130 transcripts in this

phylotypic maize embryonic-organ -initiation-network evolved in either phylostratum PS1

(unicellular eukaryotes and prokaryotes), PS2 (all eukaryotes), or PS4 (Bryophyta, i.e. the first

land plants). Less than 10 percent of this maize embryonic network comprises newer transcripts,

from PS5-PS13, that arose later in plant evolution (Extended Data Fig. 7b; Extended Data Fig. 8a;

Supplementary Dataset). Notably, the majority of genes in this network evolved within PS4,

which marks the first appearance of the plant embryo[8]. In contrast, just 12 embryonic transcripts

(9.23%) are identified from algae (PS3), comprising plants that do not contain embryos.

Relatively, gene expression numbers during the non-phylotypic Proembryo stage are more

dominated by later evolved genes from PS9-PS11(i.e. the Poaceae or grasses, the panicoid

grasses, and the genus *Zea*; Fig. 5a), whereas transcripts from Stage 2 embryos and seedling

leaves are enriched for newly evolved genes from PS12 and PS13 and the genus *Zea* (Extended

Data Fig. 8a). Thus, as in the eudicot Arabidopsis and in the animal models Drosophila and

Zebrafish, the transcriptomic hourglass pattern of the phylotypic period of maize is decidedly

embryonic, and is not identified during later, seemingly equivalent organogenic events that

continue throughout the life cycle of plants.

The maize phylotypic period displays higher TAI values (i.e. expression of newer genes) at the

Transition and Coleoptilar stages than during Stage 1 (Fig 5b; Extended Data Fig. 8b-i). These

data likely reflect the relatively recent innovation of the bipartite grass cotyledon (the scutellum

and coleoptile) that arose long after the evolution of the plant cotyledon from ancestral

embryonic leaves[10,20]. In contrast, the initiating scutellum at the Transition stage uses highly

conserved genes, corresponding to the phylotypic trough (Fig. 5c). These data suggest that once

the scutellum first appeared, the expression of highly conserved genes during scutellum initiation

ensured that this morphological novel, embryonic organ was retained in subsequent generations.

Relatedly, the slight decrease in TDI value during the Coleoptilar stage is attributed to changes in

gene expression level in the initiating coleoptile, which may ensure that the bipartite components

of the single maize cotyledon are morphologically distinct.

Historically, the morphological and transcriptomic phylotypic stages are described as occurring

during mid-embryogenesis in plants and animals[4, 5, 6, 7], although this chronological staging is

especially confounding in plants where embryo-morphogenic outcomes can vary dramatically

among taxa[7]. The phylotypic peak for transcriptomic age (TAI) in maize occurs during Stage 1, a

mid-embryogenic stage when the scutellum and coleoptile are young primordia and the first

foliar leaf is just initiating (Fig. 1d). Similarly in Arabidopsis, the transcriptomic phylotypic

hourglass began its decent during histological patterning at the globular stage, and peaked during

the torpedo stage when the two cotyledon primordia are expanding[7]. Likewise in Drosophila

embryogenesis[5, 6], gene age (TAI) peaked during germband elongation[5], at the onset of embryo

segmentation and initiation of the organogenic imaginal discs[47]. Notably, the trend toward usage

of older gene transcripts in Drosophila also began earlier in embryogenesis, during gastrulation

and the consummate formation of tissue layers[5]. In this way, the phylotypic period of

Drosophila melanogaster also resembles an extended hourglass. A similar pattern is observed in

zebrafish (*Danio rerio*), wherein the phylotypic hourglass begins its descent toward older gene

transcripts during gastrulation and then peaks in the subsequent late segmentation/early

pharyngula stages, when organ primordia initiate and grow[5]. A later reduction in relative

transcriptomic gene age is noted during the zebrafish larval stage, which is accompanied by

additional morphogenesis. Taken in summary, the phylotypic period in plants and animals

commences during tissue layer organization and extends to include organogenesis and initial

growth of embryonic organs.

Plants and animals evolved before the advent of multicellularity. Notably and perhaps

surprisingly, the earliest steps in animal and plant embryogenesis, after fertilization and during

the establishment of apical basal polarity, are not conserved within taxa and utilize new and

divergent genes to form an embryo proper (Fig. 5) [5, 6, 7]. We suggest that embryo histogenesis

and organogenesis correlates with asymmetric cell divisions that generate morphology during the

phylotypic period. In contrast, after apical-basal polarization, the initial cell divisions that

generate the globular embryo proper are primarily symmetrical and non-phylotypic. Despite their

evolutionary distances, we demonstrate that in maize, as in Arabidopsis, fruit flies, and fish, the

processes of embryonic histogenesis and initial organogenesis utilize old and conserved genes,

comprising anciently evolved, developmental genetic networks that help form the fundamental

body plan. At the end of this phylotypic period, plant and animal organs each express newer and

divergent genes in species-specific ways, to generate the vast morphological diversity found in

nature.

References

1.   Haeckel E. *Generelle Morphologie der Organismen. Allgemeine Grundzüge der Organischen*

438   *Formen-Wissenschaft, Mechanisch begründet durch die von Charles Darwin reformierte*

439   *Descendenz-Theorie, vols I and II*. Verlag Georg Reimer, Berlin (1866).

440   2.   Meckel JF. *Entwurf einer Darstellung der zwischen dem Embryozustande der höheren Tiere und*

441   *dem permanenten der niederen stattfindenden Parallele: Beyträge zur vergleichenden Anatomie*.

442   Carl Heinrich Reclam, Leipzig (1811).

443   3.   Serrés ERA. Théorie des formations organiques, ou rescherches d'anatomie transcendante sur les

444   lois de l'organogénie, appliquée à l'anatomie pathologique. *Annales de la Societe des Sciences*

445   *Naturelles et d'Archeologie de Toulon et du Var* **21**, 5-49 (1827).

446   4.   Sander K. *In Development and Evolution: the sixth Symposium of the British Society for*

447   *Developmental Biology.* . Cambridge University Press. (1983).

448   5.   Domazet-Loso T, Tautz D. A phylogenetically based transcriptome age index mirrors ontogenetic

449   divergence patterns. *Nature* **468**, 815-818 (2010).

450   6.   Kalinka AT*, et al.* Gene expression divergence recapitulates the developmental hourglass model.

451   *Nature* **468**, 811-814 (2010).

452   7.   Quint M, Drost HG, Gabel A, Ullrich KK, Bonn M, Grosse I. A transcriptomic hourglass in plant

453   embryogenesis. *Nature* **490**, 98-101 (2012).

454   8.   Soltis PS, Soltis DE, Chase MW, Endress PK, Crane PR. The diversification of flowering plants.

455   In: *Assembling the tree of life,* (eds Cracraft J, Michael J D). Oxford University Press (2004).

456   9.   Dean E. Upcoming changes in flowering plant family names: those pesky taxonomists are at it

457   again! *Fremontia* **30**, 3-12 (2002).

10.   Chandler JW. Cotyledon organogenesis. *J Exp Bot* **59**, 2917-2931 (2008).

11.　Kaplan DR, Cooke TJ. Fundamental concepts in the embryogenesis of dicotyledons: a

morphological interpretation of embryo mutants. *Plant Cell* **9**, 1903–1919 (1997).

12.　Kaplan DR. The monocotyledons: their evolution and comparative biology vii. the problem of

leaf morphology and evolution in the monocotyledons. *The Quarterly Review of Biology* **48**, 437-

457 (1973).

13.　Chen M*, et al.* Comparative analysis of embryo proper and suspensor transcriptomes in plant

embryos with different morphologies. *Proc Natl Acad Sci U S A* **118**, (2021).

14.　Randolph LF. Developmental morphology of the caryopsis in maize. *J Agric Res* **53**, 881-916

(1936).

15.　Abbe EC, Stein OL. The growth of the shoot apex in maize: embryogeny. *American Journal of*

Botany **41**, 285-293 (1954).

16.　Bommert P, Werr W. Gene expression patterns in the maize caryopsis: clues to decisions in

embryo and endosperm development. *Gene* **271**, 131-142 (2001).

17.　Nardmann J, Werr W. Patterning of the maize embryo and the perspective of evolutionary

developmental biology. In: *Handbook of Maize: Its Biology* (eds Jeff L. Bennetzen, Hake SC).

Springer (2009).

18.　Weatherwax P. Position of Scutellum and Homology of Coleoptile in Maize. *Botanical Gazette*

69, 179-182 (1920).

19.　Boyd L. Evolution in the monocotyledonous seedling: a new interpretation of the morphology of

the grass embryo. *Transactions of the Botanical Society of Edinburgh* **30**, 286-303 (1931).

20.　Kaplan DR. Early plant development: From seed to seedling to established plant. In: *Principles of*

Plant Morphology). Copy Central (1996).

21.   Satterlee JW, *et al.* A Wox3-patterning module organizes planar growth in grass leaves and

483   ligules. *Nat Plants* **9**, 720-732 (2023).

484   22.   Kaplan DR. The concept of homology and its central role in the elucidation of plant systematic

485   relationships. In: *Cladistics: Perspectives on the Reconstruction of Evolutionary History* (eds

486   Duncan T, Stuessy TF). First Edition edn. Columbia University Press (1984).

487   23.   Xu X, *et al.* Single-cell RNA sequencing of developing maize ears facilitates functional analysis

488   and trait candidate gene discovery. *Dev Cell* **56**, 557-568 e556 (2021).

489   24.   Takacs EM, *et al.* Ontogeny of the maize shoot apical meristem. *Plant Cell* **24**, 3219-3234 (2012).

490   25.   Poethig RS, Szymkowiak EJ. Clonal analysis of leaf development in maize. *Maydica* **40**, 67–76

491   (1995).

492   26.   Smith LG, Jackson D, Hake S. Expression of knotted1 marks shoot meristem formation during

493   maize embryogenesis. *Developmental Genetics* **16**, 344-348 (1995).

494   27.   Vollbrecht E, Reiser L, Hake S. Shoot meristem size is dependent on inbred background and

495   presence of the maize homeobox gene, knotted1. *Development* **127**, 3161-3172 (2000).

496   28.   Nardmann J, Ji J, Werr W, Scanlon MJ. The maize duplicate genes narrow sheath1 and narrow

497   sheath2 encode a conserved homeobox gene function in a lateral domain of shoot apical

498   meristems. *Development* **131**, 2827-2839 (2004).

499   29.   Johnston R, Wang M, Sun Q, Sylvester AW, Hake S, Scanlon MJ. Transcriptomic analyses

500   indicate that maize ligule development recapitulates gene expression patterns that occur during

501   lateral organ initiation. *Plant Cell* **26**, 4718-4732 (2014).

502   30.   Strable J, *et al.* Maize YABBY genes drooping leaf1 and drooping leaf2 regulate plant

503   architecture. *Plant Cell* **29**, 1622-1641 (2017).

504   31.   Toriba T*, et al.* BLADE-ON-PETIOLE genes temporally and developmentally regulate the sheath

505   to blade ratio of rice leaves. *Nat Commun* **10**, 619 (2019).

506   32.   Evans MM. The indeterminate gametophyte1 gene of maize encodes a LOB domain protein

507   required for embryo Sac and leaf development. *Plant Cell* **19**, 46-62 (2007).

508   33.   Leiboff S, Strable J, Johnston R, Federici S, Sylvester AW, Scanlon MJ. Network analyses

509   identify a transcriptomic proximodistal prepattern in the maize leaf primordium. *New Phytol* **230**,

510   218-227 (2021).

511   34.   Fukushima K, Hasebe M. Adaxial-abaxial polarity: the developmental basis of leaf shape

512   diversity. *Genesis* **52**, 1-18 (2014).

513   35.   Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network analysis.

514   *BMC Bioinformatics* **9**, 559 (2008).

515   36.   Langfelder P, Horvath S. Eigengene networks for studying the relationships between co-

516   expression modules. *BMC Syst Biol* **1**, 54 (2007).

517   37.   Knauer S*, et al.* A high-resolution gene expression atlas links dedicated meristem genes to key

518   architectural traits. *Genome Res* **29**, 1962-1973 (2019).

519   38.   Morabito S, Reese F, Rahimzadeh N, Miyoshi E, Swarup V. hdWGCNA identifies co-expression

520   networks in high-dimensional transcriptomics data. *Cell Rep Methods* **3**, 100498 (2023).

521   39.   Hofmann F, Schon MA, Nodine MD. The embryonic transcriptome of Arabidopsis thaliana.

522   *Plant Reprod* **32**, 77-91 (2019).

523   40.   Spurr AR. Histogenesisand organizationof the embryo in Pinusstrobus L. . *Am J Bot* **36**, 629-641

524   (1949).

41.   Kiesselbach TA. The structure and reproduction of corn. *Uni of Nebraska Agric Stat Res Bull* **161**,

1-96 (1949).

42.   Zimmermann R, Werr W. Transcription of the putative maize orthologue of the Arabidopsis

DORNROSCHEN gene marks early asymmetry in the proembryo and during leaf initiation in the

shoot apical meristem. *Gene Expr Patterns* **7**, 158-164 (2007).

43.   Poethig RS. Cellular parameters of leaf morphogenesis in maize and tobacco. In: *Contemporary*

problems in plant anatomy (eds White R, Dickison W). Academic Press (1984).

44.   Scanlon MJ. Developmental complexities of simple leaves. *Curr Opin Plant Biol* **3**, 31-36 (2000).

45.   Scanlon MJ. The polar auxin transport inhibitor N-1-naphthylphthalamic acid disrupts leaf

initiation, KNOX protein regulation, and formation of leaf margins in maize. *Plant Physiol* **133**,

597-605 (2003).

46.   Richardson AE*, et al.* Evolution of the grass leaf by primordium extension and petiole-lamina

remodeling. *Science* **374**, 1377-1381 (2021).

47.   Gilbert SF. *Developmental Biology*, 6th edn. Sunderland (MA): Sinauer Associates (2000).

48.   Ciceri P, Locatelli F, Genga A, Viotti A, Schmidt RJ. The activity of the maize Opaque2

transcriptional activator is regulated diurnally. *Plant Physiol* **121**, 1321-1328 (1999).

Figure Legends

Fig. 1 Multiplexed spatial transcriptomics of the maize Stage 1 embryo.

Differential interference contrast (DIC) microscopy of maize embryos at the Proembryo stage

(**a**), Transition stage (**b**) Coleoptilar stage, (**c**) and Stage 1 (**d**). Immunohistological analyses of

the Class I KNOX proteins mark the shoot apical meristem (SAM). ep, embryo proper; en,

endosperm; su, suspensor; sc, scutellum; c, coleoptile; L, leaf; *, SAM. Scale bar: 50 μm (**a** and

b) and 100 μm (**c** and **d**). **e**. Uniform Manifold Approximation and Projection (UMAP) of Stage1

embryo single-cell transcriptomic profiling with cell/tissue types assigned for each cell cluster.

Fig. 2 Spatial transcriptomics of maize Stage 1 embryo

Longitudinal cryosections of Stage 1 embryos embedded in OCT (optimal cutting temperature

compound; **a-b**) or in paraffin (**c-d**) for 10x Visium^{TM} spatial transcriptomics analysis. **a, c**

Toluidine Blue-stained sections for 10x Visium^{TM} spatial transcriptomics assay are overlayed

with the10x Visium^{TM} spatial map (**b, d**). The various color-coded circles correspond to the

capture spots where spatial cDNA libraries were generated. Scale bar = 100 μm. **e.** Longitudinal

sections of paraffin-embedded Stage 1 embryo for use in laser microdissection RNA-seq analysis

(LM-RNAseq). **f.** Enlarged view of a Stage 1 embryo showing regions microdissected for LM-

RNAseq). Figure labels: scu, scutellum; uc, upper coleoptile; lc, lower coleoptile. **e, f** Scale

bar=100 μm. **g**. UMAP of the Visium^{TM} spatial transcriptomic analysis of Stage 1 embryos cell

clusters. The spatial, embryonic origins of each cluster are indicated by the color-coded embryo

cartoon; numbers correspond to the cluster designations on the UMAP. **h**. Heatmap illustrating

the correlation between the single-cell (sc) clusters (y-axis) and Visium^{TM} spatial clusters

(x=axis). The significance level is marked by $-\log_{10}p$. **i**. Expression of candidate genes identified

in microdissected embryonic organs. Cell clusters exhibiting elevated candidate gene expression

of the specific markers are indicated; expression values are $\log_2$ transformed, i.e. $\log_2(TPM+1)$.

Fig. 3 RNA-targeting analyses of Transition-staged, Coleoptilar-staged, and Stage 1

embryos

a-c. Signal point map of *KN1, CUC2, NS1, WOX3A* and *DRL1* at the Transition stage (**a**),

Coleoptilar stage (**b**) and Stage 1 (**c**). **d-h.** Expression density map of *KN1* (**d**)*, CUC2* (**e**)*, NS1*

(**f**)*, WOX3A* (**g**) and *DRL1* (**h**) at Stage 1. **i-k, l-n, o-q** and **u-w**. Signal point map of *BOPa* (**i-k**),

YAB14 (**l-n**), *IG1* (**o-q**), and *ARF3* (**u-w**) at the Transition stage, Coleoptilar stage and Stage 1. **r-**

t and **x-z**. Expression density map of *IG1* (**r-t**) and *ARF3* (**x-z**) at the corresponding stages.

Figure labels: T, Transition stage; C, Coleoptilar stage; L: Stage 1. The point colors are

arbitrarily determined. Yellow asterisks mark the adaxial expression of *IG1* and *ARF3*, and red

asterisks mark the abaxial expression of *ARF3*. The green asterisk in panel **k** marks the

expression maximum of *BOPa* at the emerging leaf sheath. The yellow asterisks in panel **r-t** and

z mark the expression maxima of *IG1* at the adaxial side of the scutellum (**r**), the coleoptile (**s**),

the coleoptile and the leaf (**t**), as well as the expression maxima of *ARF3* at the adaxial side of

the scutellum (**z**), respectively. The red asterisks in panel **x-z** mark the abaxial expression of

ARF3 at the scutellum (**x, y**), the coleoptile (**y, z**) and the leaf (**z**). Scale bar for all panels = 50

μm.

Fig. 4 Dynamics of the maize embryo organ initiation network

a. Venn diagram identifies 130 genes in the embryo-organ-initiation-network. **b**. Normalized

expression of these 130 genes at the Proembryo stage (PE), Transition stage (T), Coleoptilar

stage (C), Stage 1 scutellum (Scu), Stage 1 upper coleoptile (UC), Stage 1 lower coleoptile (LC),

Stage 1 leaf (L), as well as P0 and P1 of the 14 day-after-germination seedling. The letters above

the boxplot represent significance levels (non-overlapping letters indicate significant differences

at p<0.05) by Tukey-Kramer HSD test. **c**. Heatmap of the GO terms of the putative initiation

network, as well as the single-cell (sc) cluster of the Stage 1 scutellum, coleoptile and leaf. The

heatmap color is determined by $-\log_{10}$FDR (false discovery rate). **d.** WGCNA tree graph of the

transcriptomic relationships among the co-expression modules specific to corresponding maize

organs. The height (y-axis) scales the length of organ-specific branches, indicating the

transcriptomic distance between the modules. **e**. Heatmap illustrating the correlation between the

single-cell (sc) clusters (y-axis) and hdWGCNA modules (x=axis). The significance level is

marked by $-\log_{10}$p. **f**. Heatmap illustrating the dynamic correlation of the initiation network

expression between Maize (y-axis) and Arabidopsis (x-axis) across the embryogenesis. The

heatmap color is determined by correlation coefficient (r) via Pearson correlation test with the

significance level p<0.05 (*) or p<0.01 (**). Scu, Scutellum; Col, coleoptile; hypoc, hypocotyl.

Fig. 5 Phylotypic stage analysis of the maize embryogenesis

a. Phylostratigraphic map of maize. All maize protein-coding transcripts were assigned to 13

phylostrata (PS1-PS13), Numbers in parenthesis refer to the number of genes per phylostratum,

and the taxonomic groupings are marked above the corresponding PS. **b-c**. The transcriptome

age index (TAI; **b**) and transcriptome divergence index profiles (TDI; **c**) of maize embryogenesis

and seedling leaves. Figure labels: PE, proembryo stage; T, transition stage; C, coleoptilar stage;

L, Stage 1 leaf; Scu, Stage 1 scutellum; Col, Stage 1 coleoptile; P0 and P1, Plastochron 0 and 1

leaf of the 14-day after planted seedlings. The gray lines represent the standard deviation

estimated by permutation analysis. The overall patterns of TAI and TDI profiles are significant,

as measured by flat line test ($P_{TAI}=3.56\times10^{-6}$, $P_{TDI}=4.80\times10^{-4}$), reductive hourglass test

($P_{TAI}=1.49\times10^{-4}$, $P_{TDI}=1.53\times10^{-3}$), and reductive early conservation test ($P_{TAI}=1$, $P_{TDI}=0.998$).

Fig. 6 Model for the homology of the grass cotyledon

The model describes that the single maize cotyledon as composed of two domains, the apical

scutellum (homologous to the foliar leaf blade; dashed arrow line) and the basal coleoptile

(homologous to the foliar leaf sheath; solid arrow line)

Methods

Plant Materials and growth conditions

B73 inbred lines, the *ns* 1:1 lines (segregate 1/2 wild type NS1+/*ns1 ns2/ns2* plants and 1/2

ns1/ns1 ns2/ns2 double mutant[1], and *log7*/+ (obtained from the Maize Genetics COOP Stock

Center: UFMu-02863 mu1030680::Mu) were grown in the Gutermann greenhouse facility,

Cornell University (Ithaca, NY). Transition staged, coleoptilar staged, Leaf1 staged and Leaf2

staged embryos were collected at 8-9 DAP, 10-11 DAP, 12-13 DAP, 14-15 DAP, respectively.

ns 1:1 seeds were planted in soil consisting of a 1:1 mixture of Turface and LM111 in Percival

A100 growth chamber at Weill Hall, Cornell University (Ithaca, NY) with the set condition: day

29.4 °C/16 hours, night 23.9 °C/8 hours, humidity 50%. The seedlings were collected on the 7[th]

day after planting for coleoptile comparisons between WT and *ns1;2* mutants.

Single-cell transcriptomic analysis

Two batches of B73 Leaf1 staged embryos were collected in summer 2021 and winter 2022. For

each batch, 200 embryos were dissected from developing seeds for protoplast isolation as

previously described[2]. Protoplast (1000-2000 cells per µL) were loaded into the 10X Genomics

Chromium platform to generate a Gel Bead-In-Emulsion (GEM). The GEM-captured RNA was

reverse transcribed and amplified to make 3′ cDNA libraries, followed by the high throughput

RNA sequencing on the Illumina NextSeq 2000 sequencer at the Cornell University

Biotechnology Resource Center.

The resulting raw base call (BCL) files were demultiplexed to generate FASTQ files via

CellRanger mkfastq v6.0 (10x Genomics, Pleasanton, CA). FASTQ reads were trimmed, aligned

(to B73 reference genome V3[3]) and assigned to cell barcodes to generate expression count

matrices via CellRanger count with default settings. The expression count matrices were then

analyzed using Seurat v4[4, 5, 6]. In detail, the matrices were filtered with the setting (number of

genes > 1500, UMI > 6000) to remove low quality cells. The filtered matrices were normalized

and log transformed via SCTransform function. The normalized matrices were merged and batch

effects were removed via R package Harmony v1.2.0[7]. The cell-cycle associated genes were

scored and regressed via CellCycleScoring and ScaleData function. Matrix dimension was

performed using RunPCA. The cell clusters and the Uniform Manifold Approximation and

Projection (UMAP) plot were generated with the setting (dim = 1:40, resolution = 1, n. neighbors

= 40, min. dist = 0.01, spread = 3). The differential expressed genes and cell cluster markers

were identified via FindAllMarkers function with the log2 fold change threshold = 0.2.

Spatial transcriptomic analysis

For use in spatial transcriptomics, B73 Leaf1 staged embryos were processed for either paraffin-

embedding and microtome sectioning, or cryoembedding and cryosectioning. For paraffin -

embedding, two embryos were fixed in Farmer's Fixative (Ethanol:Acetic acid = 3:1), followed

by dehydration in an ethanol concentration gradient (75%, 85%, 95%, 100%) and clearing with

Ethanol:Xylene gradient of 3:1, 1:1, 1:3, and 100% Xylene at room temperature. Cleared

embryos were embedded in Paraplast Plus® (Leica Biosystems, Deer Park, IL). The embedded

blocks were microtome-sectioned at 10 um at room temperature; sections were subsequently

mounted onto 10x Visium^{TM} Spatial Gene Expression Slides (10x Genomics, Pleasanton, CA).

Slides were deparaffinized with two changes of 100% Xylene and 95% Ethanol for 10 min and 2

min, respectively. For cryoembedding, two kernels were dissected longitudinally to remove left

and right margin of the kernel; the central region containing the embryo was embedded in

Tissue-Tek® Optimal Cutting Temperature (O.C.T., Sakura Finetek, Torrance, CA), and snap-

frozen in the dry ice. Frozen tissue blocks were cryosectioned at10 µm, and the sections were

mounted on 10x Visium^{TM} Spatial Gene Expression Slide (10x Genomics, Pleasanton, CA).

The 10x Visium^{TM} Spatial Gene Expression Slides were processed for spatial library

construction following the manufacturer instructions. Briefly, the sections were baked at 37 ^{o}C

for 1 min followed by re-fixation in 100% methanol at -20 ^{o}C for 30 min. Subsequently, the

sections were stained with Toluidine Blue for histological imaging. Tissue permeabilization was

performed at 37 ^{o}C for 6 min to release RNA from the fixed tissue, followed by first strand

cDNA synthesis. Subsequently, second strand cDNA was synthesized and amplified for library

construction and sequencing. High-throughput RNA sequencing was conducted on an Illumina

NextSeq 2000 sequencer at the Cornell University Biotechnology Resource Center.

Raw reads were demultiplexed, mapped to the Toluidine Blue stained images, and aligned to the

B73 reference genome V3[3] to generate spatial expression count matrices using SpaceRanger v2.0

(10x Genomics, Pleasanton, CA). The matrices were analyzed using Seurat v4[4, 5, 6]. In detail, the

matrices were normalized and log transformed via SCTransform function. The normalized

matrices were merged followed by dimensionality reduction. Cell clusters and the UMAP were

generated with the following settings (dim = 1:40; resolution = 1.5; neighbors = 10; minimum

distance =0.01). The spatial data were visualized with DimPlot and SpatialPlot; differential

expressed genes and cell cluster markers were identified via FindAllMarkers function with the

log2 fold change threshold = 0.1.

Laser-microdissection RNAseq analysis

Sixty maize Leaf1 staged kernels in two replicates were dissected longitudinally to remove the

left and right margins of the kernels. This embryo-containing central region was fixed in

Farmer's Fixative (Ethanol:Acetic acid=3:1). After fixation, the samples were dehydrated,

cleared, embedded and sectioned as described previously[8, 9] . Tissue sections were mounted on

PEN membrane slides (Leica Microsystems, Deer Park, IL) pre-treated at 180 $^{\circ}$C for 4 hours

followed by UV irradiation at 254 nm for 30 minutes. The sections were deparaffinized with two

changes of 100% Xylene and 100% Ethanol for 10 min and 2 min, respectively. Laser

microdissection was performed using a PALM Laser Microbeam System (Zeiss Group,

Oberkochen, Germany). Adaxial scutellum, SAM, leaf, as well as upper and lower coleoptiles

were microdissected (total area for each captured organ is over 0.3 mm^2) and RNA extraction

was performed using the Arcturus^{TM} PicoPure^{TM} RNA Isolation Kit (Thermo Fisher Scientific,

Waltham, MA). The extracted RNA was amplified using Arcturus^{TM} RiboAmp^{TM} HS PLUS

RNA Amplification Kit (Thermo Fisher Scientific, Waltham, MA). RNA quality control, library

construction and sequencing were performed at the Cornell University Biotechnology Resource

Center. The raw reads quality control was conducted via FastQC

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Raw reads were mapped to the.

B73 reference genome V3[3] by HISAT2 (v2.1.0)[10, 11]. Read counts and the normalized number of

transcripts (transcript per million reads, TPM) were calculated as previously described[12]. DEseq2

(v1.26.0)[13] was used to identify differentially expressed genes (DEGs) with the cutoff criteria of

log2 fold change >1 or <1 with the false discovery rate (FDR) < 0.05.

Multiplex *in situ* hybridization

Transition staged, coleoptilar staged and Leaf1 staged kernels (two replicate for each stage) were

dissected longitudinally to remove the left and right kernel margins. The central region

containing the embryo was fixed in 4% paraformaldehyde (PFA); fixed samples were

dehydrated, cleared, embedded and sectioned as described previously[14, 15]. Sections were

mounted on 10x Genomics Xenium[TM] slides, deparaffinized and decrosslinked following the

manufacturer's instructions. The Xenium custom gene expression panel containing probes of 99

maize genes were applied on each section to perform hybridization, ligation and rolling circle

amplification, followed by cycles of fluorescent probe hybridization, imaging and decoding. The

data was visualized by Xenium Explorer v1.3. The Xenium[TM] multiplex *in situ* hybridization

assays were conducted at Memorial Sloan Kettering Cancer Center (New York, NY).

Histology and immunohistolocalizations

Developing kernels from the proembryo stage, transition stage, coleoptilar stage, and the Stage 1

were fixed in FAA, dehydrated, cleared, embedded in paraplast, sectioned and stained as

described previously[15]. Immunohistolocalizations were performed as described[1].

WGCNA and hdWGCNA

For WGCNA to pursue to initiation network, log2-transformed TPM data from LM-RNAseq

(SAM data was excluded) were filtered following these criteria: row mean > 0.2 and coefficient

of variation > 0.4. The general pipeline was performed as per R package instructions[16] with

minor modifications as follows. The network type was set to "signed". The soft power, the

minimum module size ,and the merging threshold were set to 13, 20 and 0.3, respectively. For

the analysis to pursue transcriptomic relationship between organs, the LM-RNAseq data in this

study was merged with the LM-RNAseq data of the 14-day seedling P0 and P1 data[17]. The data

was normalized and batched effect removed as described, and filtered following the same criteria

as the WGCNA for the initiation network. The network type was set to "signed". The soft power,

the minimum module size ,and the merging threshold were set to 16, 20 and 0.4, respectively.

For hdWGCNA, the single-cell RNAseq data was used to perform the analysis. The data was

SCTransformed and normalized. The general pipeline was conducted as the R package

instructions[18] with the following minor modifications. The network type was set to "signed", and

the soft power and minimum module size were set to 10 and 30, respectively. Hypergeometric

test was performed to analyze the correlation between the hdWGCNA modules and the single-

cell clusters following the previously described procedures[19].

Gene Ontology (GO) enrichment analysis

GO enrichment analysis of the putative initiation network as well as the single-cell scutellum,

coleoptile and leaf clusters was performed by AgriGOv2

(http://systemsbiology.cau.edu.cn/agriGOv2/) with the locus ID v3.30 (Gramene Release 50)

datasets as GO annotation background. The significant enrichment cutoff FDR was set to 0.05.

Statistical analysis

Hypergeometric tests were performed to analyze the correlation between single-cell clusters and

spatial transcriptomic clusters, as well as between the single-cell clusters and hdWGCNA

modules. We followed previously algorithms[19] to statistically evaluate if the two groups of data

were significantly correlated by calculating the number of overlapping gene relative to the total

number of protein-coding genes in the maize B73 reference genome V3[3]. The p values were

transformed by –log10p, and visualized as heatmap built by the R package gplots

(https://github.com/cran/gplots).

LM-RNAseq read count data were normalized to fit the generalized linear model (GLM), and

Empirical Bater shrinkage estimation was used for calculating dispersions, expression fold

changes and adjust p values. In addition, Pearson correlation test was used to calculate the

correlation between the coexpression modules and the corresponding clusters (microdissected

organs or putative initiation network) in WGCNA, and between the maize and Arabidopsis

embryogenesis genes.

To statistically compare the expression of the initiation network between the transition,

coleoptilar and the stage1, the TPM data was log2 transformed [log2(TPM+1)] and compared

using Tukey-Kramer HSD test by JMP Pro 17 (SAS Institute, Cary, NC) with default settings.

For the correlation test, the transition, coleoptilar and the stage1 TPM data was double log2

transformed, i.e. log2[log2(TPM+1)+1], to fix the data skewness caused by the outliers. The

Arabidopsis TPM data was log2 transformed. The Pearson correlation was performed in R v4.3.2

with the significance cutoff value $p<0.05$.

Calculating TAI and TDI for phylotypic period

The procedures of constructing a phylostratigraphic map, calculating TAI and TDI, and

statistical tests, have been presented previously[20, 21, 22, 23]. For each of the 13 phylostrata shown in

Fig. 5a, the amino acid sequences of all 350 species with completely sequenced genomes were

extracted from NCBI, Ensembl genomes, Treegene (https://treegenesdb.org), Fernbase

(https://fernbase.org), or Maizegdb (https://maizegdb.org). A database of these sequences was

generated, and each of the 63,031 amino acid sequences of *Zea mays* ssp. *mays* (v3) with a

minimum length of 30 amino acids was blasted against this database using BLASTp (BLAST

version 2.15.0) with an E-value cut-off of 10^{-5}. The corresponding gene of *Zea mays* ssp. *mays*

was assigned to the phylogenetically most distant (oldest) phylostratum containing at least one

species with at least one blast hit, otherwise the corresponding gene of *Zea mays* ssp. *mays* was

assigned to phylostratum 13 (PS13) where no blast hit was identified.

Orthologous gene pairs of *Zea mays* ssp. *mays* and *Zea mays* ssp. *mexicana* were determined

with the method of best hits using BLASTp. Amino acid sequence alignments of each pair

generated with MAFFT[24] (L-INS-i option) were used for codon alignments generated with

PAL2NAL[25] to compute sequence divergence levels (K_a/K_S) with KaKs Calculator (version

3.0)[26]. Gene pairs with $K_a<0.5$, $K_s<5$ and K_a/K_s ratios < 2 were retained.

Expression levels of maize development stages were collected from 4 data sources 1) LM-

RNAseq data of proembryo stage[27], 2) organ initiation network (this study), 3) RNAseq data of

Stage 2 embryo[28] and 4) LM-RNAseq data of 14-day after germination seedling P0 and P1

leaves[17]. The datasets were processed with batch effect removal and normalization as described

above. The TAI and the TDI are weighted means of evolutionary age and sequence divergence,

respectively, and defined previously[20, 22]. The transcriptome age index TAI_s and the

transcriptome divergence index TDI_s of developmental stage s (s = proembryo, transition,

coleoptilar, stage 1 leaf, stage 1 scutellum, stage 1 coleoptile, seedling leaves P0 and seedling

leaves P1) are the weighted mean of the evolutionary age (phylostratum) weighted by the

expression level of each gene at developmental stage s, and the weighted mean of the Ka/Ks

ratio of each gene weighted by the expression level of each gene at developmental stage *s*,

respectively. Low/high PS values correspond to evolutionarily old/young genes, so low/high TAI

values correspond to evolutionarily old/young transcriptomes. Low/high K_a/K_S ratios correspond

to conserved/divergent genes, so low/ high TDI values correspond to conserved/divergent

transcriptomes. For gene models with multiple predicted peptides, TAI was calculated with the

most conserved phylostratum assigned to that locus and TDI was calculated with the conserved

codon divergence strata assigned to that locus.

To determine the statistical significance of the TAI and TDI profiles, the flat line test[22], reductive

hourglass test and reductive early conservation test were performed via myTAI package

(v1.0.1)[23]. The flat line test is a permutation test based on the variance of the TAI or TDI values

of a given TAI or TDI profile. To reductive hourglass test, the developmental stages were

partitioned into three modules, early (proembryo stage), mid (transition stage, coleoptilar stage,

Stage 1 leaf, Stage 1 scutellum, and Stage 1 coleoptile), and late (stage 2 and seedling leaves),

and permutation test was performed for the minimum differences of the mean TAI or TDI values

between early and mid-modules, and between late and mid modules. To reductive early

conservation test, permutation test was performed for the minimum differences of the mean TAI

or TDI values between mid and early modules, and between late and early modules. The detailed

procedures were described in Drost, et al., 2015[23].

Data accessions

The single-cell RNAseq, Visium™ spatial transcriptomics, and LM-RNAseq data were

deposited in the NCBI Sequence Read Archive (SRA) with the BioProject ID: PRJNA1045558.

The Xenium™ multiplex *in situ* hybridization data was deposited in Science Data Bank with the

link: https://www.scidb.cn/s/F7ZbEz, which could be visualized by Xenium Explorer v1.3. The

code for TAI and TDI calculations were uploaded to https://gitlab.com/rz444/maizephylper.git.

Methods references

1.   Scanlon MJ, Freeling M. The narrow sheath leaf domain deletion : a genetic tool used to

reveal developmental homologies among modified maize organs. *Plant Journal* **13**, 547-

561 (1998).

2.   Satterlee JW, Scanlon MJ. Protoplast isolation from undifferentiated maize seedling

shoot tissue. *Methods Mol Biol* **2464**, 123-130 (2022).

3.   Schnable PS*, et al.* The B73 maize genome: complexity, diversity, and dynamics. *Science*

326, 1112-1115 (2009).

4. Hao Y*, et al.* Dictionary learning for integrative, multimodal and scalable single-cell

analysis. *Nat Biotechnol*, (2023).

5. Stuart T*, et al.* Comprehensive integration of single-cell data. *Cell* **177**, 1888-1902 e1821

(2019).

6. Hao Y*, et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587

e3529 (2021).

7. Korsunsky I*, et al.* Fast, sensitive and accurate integration of single-cell data with

Harmony. *Nat Methods* **16**, 1289-1296 (2019).

8. Scanlon MJ, Ohtsu K, Timmermans MC, Schnable PS. Laser microdissection-mediated

isolation and in vitro transcriptional amplification of plant RNA. *Curr Protoc Mol Biol*

Chapter 25, Unit 25A 23 (2009).

9. Zhang S, Thakare D, Yadegari R. Laser-capture microdissection of maize kernel

compartments for rna-seq-based expression analysis. *Methods Mol Biol* **1676**, 153-163

(2018).

10. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome alignment and

genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**, 907-915 (2019).

11. Kim D, Langmead B, Salzberg SL. HISAT: A fast spliced aligner with low memory

requirements. *Nature Methods* **12**, 357-360 (2015).

12. Wu H*, et al.* NAKED ENDOSPERM1, NAKED ENDOSPERM2, and OPAQUE2

interact to regulate gene networks in maize endosperm development. *Plant Cell*, (2023).

13.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biology* **15**, 550 (2014).

14.   Laureyns R*, et al.* An in situ sequencing approach maps PLASTOCHRON1 at the

boundary between indeterminate and determinate cells. *Plant Physiol* **188**, 782-794

(2022).

15.   Jackson D. In situ hybridization in plants. In: *Molecular Plant Pathology: A Practical*

Approach, (eds D. J. Bowles, S. J. Gurr, McPherson M). Oxford University Press (1991).

16.   Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network

analysis. *BMC Bioinformatics* **9**, 559 (2008).

17.   Knauer S*, et al.* A high-resolution gene expression atlas links dedicated meristem genes

to key architectural traits. *Genome Res* **29**, 1962-1973 (2019).

18.   Morabito S, Reese F, Rahimzadeh N, Miyoshi E, Swarup V. hdWGCNA identifies co-

expression networks in high-dimensional transcriptomics data. *Cell Rep Methods* **3**,

100498 (2023).

19.   Zhan J*, et al.* RNA sequencing of laser-capture microdissected compartments of the

maize kernel identifies regulatory modules associated with endosperm cell differentiation.

The Plant Cell **27**, 513-531 (2015).

20.   Domazet-Loso T, Tautz D. A phylogenetically based transcriptome age index mirrors

ontogenetic divergence patterns. *Nature* **468**, 815-818 (2010).

21.   Domazet-Loso T, Brajkovic J, Tautz D. A phylostratigraphy approach to uncover the

genomic history of major adaptations in metazoan lineages. *Trends Genet* **23**, 533-539

(2007).

22.   Quint M, Drost HG, Gabel A, Ullrich KK, Bonn M, Grosse I. A transcriptomic hourglass

in plant embryogenesis. *Nature* **490**, 98-101 (2012).

23.   Drost HG, Gabel A, Grosse I, Quint M. Evidence for active maintenance of

phylotranscriptomic hourglass patterns in animal and plant embryogenesis. *Mol Biol Evol*

32, 1221-1231 (2015).

24.   Katoh K, Kuma K, Toh H, Miyata T. MAFFT version 5: improvement in accuracy of

multiple sequence alignment. *Nucleic Acids Res* **33**, 511-518 (2005).

25.   Suyama M, Torrents D, Bork P. PAL2NAL: robust conversion of protein sequence

alignments into the corresponding codon alignments. *Nucleic Acids Res* **34**, W609-612

(2006).

26.   Zhang Z. KaKs_Calculator 3.0: Calculating selective pressure on coding and non-coding

sequences. *Genomics Proteomics Bioinformatics* **20**, 536-540 (2022).

27.   Takacs EM, *et al.* Ontogeny of the maize shoot apical meristem. *Plant Cell* **24**, 3219-

3234 (2012).

28.   Chen J, *et al.* Dynamic transcriptome landscape of maize embryo and endosperm

development. *Plant Physiol* **166**, 252-264 (2014).

Acknowledgements

We thank L. Evans and R. Ragas for helpful discussions of the data and comments on the manuscript. Special thanks to Cornell Genomics Facility for performing single-cell and Visium™ sequencing, to the Cornell Transcriptional Regulation and Expression Facility for RNA sequencing, and the Cornell Guterman Bioclimatic Labs for our plants. Thanks to the Integrated Genomics Operation (IGO) of Memorial Sloan Kettering Cancer Center for performing Xenium assays. H. Wu was supported by a grant from the National Science Foundation (IOS-2016021). R. Zhang is supported by another grant from the National Science Foundation (IOS-2210259).

Author contributions

Author Contributions M.J.S, H.W. and R.Q. conceived the study. M.J.S. supervised the project. H.W and R.Q. generated and analyzed the data. M.J.S, H.W. and R.Q. interpreted the results, and M.J.S, H.W. and R.Q wrote the manuscript.

Ethics declaration

Competing interests
The authors declare no competing interests.

Supplementary information

Supplementary Dataset 1. Single-cell RNAseq cluster marker genes

Supplementary Dataset 2. Spatial transcriptomic cluster markers

Supplementary Dataset 3. Normalized gene expression (TPM) of LM-RNAseq

Supplementary Dataset 4. Normalized gene expression (log2(TPM+1)) of tissue specific classic genes

925 Supplementary Dataset 5. Xenium^{TM} multiplex in situ hybridization probes and result

Supplementary Dataset 5. Xenium^{TM} multiplex in situ hybridization probes and result

Supplementary Dataset 6. WGCNA: modules with gene expression pattern:

Scu<UpCol<LoCol<Leaf

Supplementary Dataset 7. Putative initiation network and the log2-transformed TPM in the

lateral organs of the transition, coleoptilar, Stage 1 (L1) and 14-day after germination seedlings

Supplementary Dataset 8. Gene ontology term category and corresponding FDR

Supplementary Dataset 9. WGCNA: Cell-type specific modules

Supplementary Dataset 10. hdWGCNA module table

Supplementary Dataset 11. Maize initiation network vs Arabidopsis orthologs

Supplementary Dataset 12. Gene expression for TAI and TDI calculation

Supplementary Dataset 13. Phylostrata of maize genes

Supplementary Dataset 14. *Ka/Ks* ratio of maize genes expressed at the embryogenesis and 14-

day seedling

Supplementary Dataset 15. Genomes used for phylogeny reconstruction and phylostratigraphic

map.

Supplementary Dataset 16. The gene numbers of each phylostratum in embryo stages.

Corresponding author

Michael J. Scanlon: mjs298@cornell.edu

Hao Wu: haowu2021@gmail.com

Extended data figure legends

Extended Data Fig. 1 Verification of single-cell cluster marker genes.

Left panel: the single-cell cluster marker expression on a UMAP. Purple dots mark cells with

corresponding gene expression data. Middle panel: the verification of the single-cell cluster

marker genes by Visium^{TM}. Red to green spots marks the location of the gene expression with

high to low level, respectively. Right panel: the verification of the single-cell cluster marker

genes by multiplex *in situ* hybridization. Note: that *GRMZM2G166141* (**k**), *GRMZM2G094353*

(**m**) and *GRMZM2G099454* (**p**) lack the multiplex *in situ* hybridization data. Scale bar = 100 μm.

Extended Data Fig. 2 Differential expression between laser microdissected RNAseq

samples.

a. Principal component analysis of laser-microdissected RNAseq samples. Each dot represents a

laser-microdissected embryonic shoot sample. **b-f**. Pairwise comparison of the gene expression

between laser-microdissected RNAseq samples. The number of overlapping, differential-

expressed genes (DEGs) is showed in the Venn diagram. The circled numbers represent the

genes preferentially expressed in the SAM (**b**), upper coleoptile (**c**), scutellum (**d**), leaf (**e**) and

lower coleoptile (**f**).

Extended Data Fig. 3 Multiplex *in situ* hybridization of *TRU1*.

The spatial expression of *TRU1* is showed in signal point map (**a-c**) and density map (**d-f**) from

the transition stage (**a and d**), the coleoptilar stage (**b and e**) and the Stage 1 (**c and f**). The red

arrow marks the expression of *TRU1* at the emerging leaf sheath. Scale bar = 100 μm.

Extended Data Fig. 4 Expression of the embryo organ initiation network.

a. WGCNA to identify modules with overall highest expression in the Stage 1 leaf (L) and less in

the lower coleoptile (LC), with steadily decreased expression in the upper coleoptile (UC) and

scutellum (S) (i.e. L > LC > UC > S). The heatmap shows the correlation coefficient (r) and the

corresponding p value (in the parentheses) of each module (ME1-12). **b**. The expression of the

130 genes in the initiation network at the transition stage (T), coleoptilar stage (C) and the Stage

1 scutellum (Scu), upper coleoptile (UC), lower coleoptile (LC) and leaf (L). **c-e**. The correlation

of the expression of the 130 genes between initiation stage of the scutellum, coleoptile and leaf.

Extended Data Fig. 5 WGCNA Module-organ correlation.

a-b. Correlation between datasets of the 14-day after pollination seedling and the embryo before

and after batch effect removal. The correlation of the normalized overall gene expression

between the 14 day after germination (DAG) seedling and embryo datasets before (**a**) and after

(**b**) batch effect removal. **c**. Heatmap illustrating the correlation between co-expression modules

(y-axis) and corresponding organs (x-axis. The colors are determined by the correlation

coefficient (r) with the p value (numbers in the parentheses)

Extended Data Fig. 6 Correlation between the maize initiation network genes and the

Arabidopsis orthologs.

The correlation of expression between the maize initiation network genes at transition (**a**) and

coleoptilar (**b**) stage, and the Arabidopsis orthologs at early heart stage.

Extended Data Fig. 7. TAI and TDI for embryo stages under different conditions

a-c. Distribution of all maize genes and genes in each embryo stage across phylostrata and TAI

profiles for different E-value cut-offs (ranging from 1E-2 to 1E-20). This analysis investigates

whether and how E-values affect the hourglass pattern. **a**. Variation of the E-value cut-off for the

BLASTp analysis demonstrates minimal import on the distribution of maize genes among the

phylostrata, indicating an overall pattern independent of the E-values. The y-axis is presented on

a logarithmic scale. **b**. Similarly, the variation of the E-value cut-off for the BLASTp analysis

does not much severely alter the distribution of maize genes in each embryo stage among the

phylostrata. **c**. the hourglass pattern of the TAI profile remains consistent across various E-value

cut-offs. **d-e**. Transcriptome divergence indices based on two different genome comparisons:

Zea mays ssp. mays vs. *Zea mays* ssp. mexicana (**d**); *Zea mays* ssp. mays vs. *Zea diploperennis*

(**e**). The overall patterns of TDI profiles are significant, as measured by flat line test ($P_{\text{TDI-}}$

$_{\text{Mexicana}}=1.07\times10^{-3}$, $P_{\text{TDI-diploperennis}}=4.80\times10^{-4}$), reductive hourglass test ($P_{\text{TDI Mexicana}}=0.027$, $P_{\text{TDI-}}$

$_{\text{diploperennis}}=1.53\times10^{-3}$), and reductive early conservation test ($P_{\text{TDI Mexicana}}=0.68$, $P_{\text{TDI-diploperennis}}$

$=0.998$).

1007

Extended Data Fig. 8. The gene expression in each embryo stage per phylostratum.

a. The stacked bars in percentage of gene numbers of each phylostratum in embryo stages. **b-i.**

The relative gene expression (normalized to the maximal level in each embryo stage) of maize

embryo genes, including in proembryo stage (**b, c**), in Initiation-Network stages (transition stage,

coleoptilar stage, and stage 1 leaf) (**d, e**), in stage 1 (stage 1 leaf, stage 1 scutellum, stage 1

coleoptile) (**f, g**), and after stage 2 and in seedling leaves (**h, i**). The data to after stage 2 is from

Chen et al., 2014, and were normalized in data source in (**h**).

1015

1016 **Extended Data Table 1 Identity of single-cell clusters.**

Single cell clusters	Identity	Evidence
0	Basal scutellum	Fig. 1h; Extended Data Fig. 1a
1	Abaxial or adaxial scutellum epidermis	Fig. 1h; Extended Data Fig. 1b
2	Abaxial scutellum epidermis	Fig. 1h; Extended Data Fig. 1c
3	Apical scutellum	Fig. 1h; Extended Data Fig. 1d
4	Adaxial scutellum epidermis	Fig. 1h; Extended Data Fig. 1e
5	Abaxial scutellum epidermis	Fig. 1h; Extended Data Fig. 1f
6	SAM and hypocotyl	Fig. 1h; Extended Data Fig. 1g; known marker genes[24, 29]
7	Abaxial scutellum epidermis	Fig. 1h; Extended Data Fig. 1f
8	Apical scutellum	Fig. 1h; Extended Data Fig. 1d
9	Possible vasculature	Fig. 1h; Extended Data Fig. 1h
10	Coleoptile epidermis	Fig. 1h,n; Extended Data Fig. 1i
11	Basal scutellum	Fig. 1h; Extended Data Fig. 1a
12	Possible embryo-endosperm boundary region	Fig. 1h; Extended Data Fig. 1j
13	Apical scutellum	Fig. 1h; Extended Data Fig. 1k
14	Abaxial scutellum epidermis	Fig. 1h; Extended Data Fig. 1l
15	Vasculature	Extended Data Fig. 1m; known marker genes[24]
16	Boundary of the lateral organs	Fig. 1h; Extended Data Fig. 1n

17	Coleoptile internal tissue	Fig. 1h,i,j,k,n; Extended Data Fig. 1o
18	Possible Tip of scutellum	Fig. 1h; Extended Data Fig. 1p
19	Apical scutellum or suspensor (residue)	Fig. 1h; Extended Data Fig. 1q
20	Endosperm (residue)	Fig. 1h; known marker genes[48]
21	Leaf	Fig. 1h,n; Extended Data Fig. 1r
22	Suspensor (residue)	Fig. 1h; Extended Data Fig. 1s

1017   Evidence based on previous published marker genes, Visium[TM] and multiplex *in situ*

1018   hybridization (Extended Data Fig. 1)

1019

1020   **Extended Data Table 2 Gene expression of selected leaf genes in the transition, coleoptilar**

1021   **and Leaf1 stage in the LM-RNAseq data.**

Genes	Transition	Coleoptilar	Leaf1
YAB14	60.85	59.62	82.24
FPF1L	83.16	178.63	103
IG1	33.85	21.49	26.94
NS1	8.37	19.55	10.28

1022

1023

1024

1025

1026

1027

1028

1029

1030

1031

1032

1033

1034   **Figures and Extended Data Figures**

Fig. 1 Multiplexed spatial transcriptomics of the maize Stage 1 embryo.

0,3,8,11,13 Scutellum internal tissue
1,2,4,5,7,14 Scutellum epidermis
6 SAM and hypocotyl
9,15 Vasculature
Coleoptile epidermis
Embryo-endosperm boundary
Boundary of lateral organs
Coleoptile internal tissue
Scutellum tip
Apical scutellum or suspensor
Endosperm (residue)
Leaf
Suspensor

Differential interference contrast (DIC) microscopy of maize embryos at the Proembryo stage

(**a**), Transition stage (**b**) Coleoptilar stage, (**c**) and Stage 1 (**d**). Immunohistological analyses of

the Class I KNOX proteins mark the shoot apical meristem (SAM). ep, embryo proper; en,

endosperm; su, suspensor; sc, scutellum; c, coleoptile; L, leaf; *, SAM. Scale bar: 50 μm (**a** and

b) and 100 μm (**c** and **d**). **e**. Uniform Manifold Approximation and Projection (UMAP) of Stage1

embryo single-cell transcriptomic profiling with cell/tissue types assigned for each cell cluster.

1043

1044

Fig. 2 Spatial transcriptomics of maize Stage 1 embryo

Longitudinal cryosections of Stage 1 embryos embedded in OCT (optimal cutting temperature

compound; **a-b**) or in paraffin (**c-d**) for 10x Visium^{TM} spatial transcriptomics analysis. **a, c**

Toluidine Blue-stained sections for 10x Visium^{TM} spatial transcriptomics assay are overlayed

with the10x Visium^{TM} spatial map (**b, d**). The various color-coded circles correspond to the

capture spots where spatial cDNA libraries were generated. Scale bar = 100 μm. **e.** Longitudinal

sections of paraffin-embedded Stage 1 embryo for use in laser microdissection RNA-seq analysis

(LM-RNAseq). **f.** Enlarged view of a Stage 1 embryo showing regions microdissected for LM-

RNAseq). Figure labels: scu, scutellum; uc, upper coleoptile; lc, lower coleoptile. **e, f** Scale

bar=100 μm. **g**. UMAP of the Visium^{TM} spatial transcriptomic analysis of Stage 1 embryos cell

clusters. The spatial, embryonic origins of each cluster are indicated by the color-coded embryo

cartoon; numbers correspond to the cluster designations on the UMAP. **h**. Heatmap illustrating

the correlation between the single-cell (sc) clusters (y-axis) and Visium^{TM} spatial clusters

(x=axis). The significance level is marked by $-\log_{10}p$. **i**. Expression of candidate genes identified

in microdissected embryonic organs. Cell clusters exhibiting elevated candidate gene expression

of the specific markers are indicated; expression values are $\log_2$ transformed, i.e. $\log_2(TPM+1)$.

1061

1062

1063

1064

Fig. 3 RNA-targeting analyses of Transition-staged, Coleoptilar-staged, and Stage 1

embryos

a-c. Signal point map of *KN1, CUC2, NS1, WOX3A* and *DRL1* at the Transition stage (**a**),

Coleoptilar stage (**b**) and Stage 1 (**c**). **d-h.** Expression density map of *KN1* (**d**)*, CUC2* (**e**)*, NS1*

(**f**)*, WOX3A* (**g**) and *DRL1* (**h**) at Stage 1. **i-k, l-n, o-q** and **u-w**. Signal point map of *BOPa* (**i-k**),

YAB14 (**l-n**), *IG1* (**o-q**), and *ARF3* (**u-w**) at the Transition stage, Coleoptilar stage and Stage 1. **r-**

t and **x-z**. Expression density map of *IG1* (**r-t**) and *ARF3* (**x-z**) at the corresponding stages.

Figure labels: T, Transition stage; C, Coleoptilar stage; L: Stage 1. The point colors are

arbitrarily determined. Yellow asterisks mark the adaxial expression of *IG1* and *ARF3*, and red

asterisks mark the abaxial expression of *ARF3*. The green asterisk in panel **k** marks the

expression maximum of *BOPa* at the emerging leaf sheath. The yellow asterisks in panel **r-t** and

z mark the expression maxima of *IG1* at the adaxial side of the scutellum (**r**), the coleoptile (**s**),

the coleoptile and the leaf (**t**), as well as the expression maxima of *ARF3* at the adaxial side of

the scutellum (**z**), respectively. The red asterisks in panel **x-z** mark the abaxial expression of

ARF3 at the scutellum (**x, y**), the coleoptile (**y, z**) and the leaf (**z**). Scale bar for all panels = 50

μm.

1081

1082

Fig. 4 Dynamics of the maize embryo organ initiation network

a. Venn diagram identifies 130 genes in the embryo-organ-initiation-network. **b**. Normalized

expression of these 130 genes at the Proembryo stage (PE), Transition stage (T), Coleoptilar

stage (C), Stage 1 scutellum (Scu), Stage 1 upper coleoptile (UC), Stage 1 lower coleoptile (LC),

Stage 1 leaf (L), as well as P0 and P1 of the 14 day-after-germination seedling. The letters above

the boxplot represent significance levels (non-overlapping letters indicate significant differences

at p<0.05) by Tukey-Kramer HSD test. **c**. Heatmap of the GO terms of the putative initiation

network, as well as the single-cell (sc) cluster of the Stage 1 scutellum, coleoptile and leaf. The

heatmap color is determined by $-\log_{10}$FDR (false discovery rate). **d.** WGCNA tree graph of the

transcriptomic relationships among the co-expression modules specific to corresponding maize

organs. The height (y-axis) scales the length of organ-specific branches, indicating the

transcriptomic distance between the modules. **e**. Heatmap illustrating the correlation between the

single-cell (sc) clusters (y-axis) and hdWGCNA modules (x=axis). The significance level is

marked by $-\log_{10}$p. **f**. Heatmap illustrating the dynamic correlation of the initiation network

expression between Maize (y-axis) and Arabidopsis (x-axis) across the embryogenesis. The

heatmap color is determined by correlation coefficient (r) via Pearson correlation test with the

significance level $p<0.05$ (*) or $p<0.01$ (**). Scu, Scutellum; Col, coleoptile; hypoc, hypocotyl.

1100

1104

Fig. 5 Phylotypic stage analysis of the maize embryogenesis

a. Phylostratigraphic map of maize. All maize protein-coding transcripts were assigned to 13

phylostrata (PS1-PS13), Numbers in parenthesis refer to the number of genes per

phylostratum, and the taxonomic groupings are marked above the corresponding PS. **b-c**. The

transcriptome age index (TAI; **b**) and transcriptome divergence index profiles (TDI; **c**) of

maize embryogenesis and seedling leaves. Figure labels: PE, proembryo stage; T, transition

stage; C, coleoptilar stage; L, Stage 1 leaf; Scu, Stage 1 scutellum; Col, Stage 1 coleoptile;

P0 and P1, Plastochron 0 and 1 leaf of the 14-day after planted seedlings. The gray lines

represent the standard deviation estimated by permutation analysis. The overall patterns of

TAI and TDI profiles are significant, as measured by flat line test ($P_{\text{TAI}}=3.56\times10^{-6}$,

$P_{\text{TDI}}=4.80\times10^{-4}$), reductive hourglass test ($P_{\text{TAI}}=1.49\times10^{-4}$, $P_{\text{TDI}}=1.53\times10^{-3}$), and reductive

early conservation test ($P_{\text{TAI}}=1$, $P_{\text{TDI}}=0.998$).

1117

1118

1119

1120

1121 **Fig. 6 Model for the homology of the grass cotyledon**

1122 The model describes that the single maize cotyledon as composed of two domains, the apical

1123 scutellum (homologous to the foliar leaf blade; dashed arrow line) and the basal coleoptile

1124 (homologous to the foliar leaf sheath; solid arrow line)

1125

1126

1127

1128

1129

1130

1131

1132

1133

1134

Extended Data Fig. 1 Verification of single-cell cluster marker genes.

Left panel: the single-cell cluster marker expression on a UMAP. Purple dots mark cells with

corresponding gene expression data. Middle panel: the verification of the single-cell cluster

marker genes by Visium™. Red to green spots marks the location of the gene expression with

high to low level, respectively. Right panel: the verification of the single-cell cluster marker

genes by multiplex *in situ* hybridization. Note: that *GRMZM2G166141* (**k**), *GRMZM2G094353*

(**m**) and *GRMZM2G099454* (**p**) lack the multiplex *in situ* hybridization data. Scale bar = 100 μm.

1142

1143

1144

Extended Data Fig. 2 Differential expression between laser microdissected RNAseq

samples.

a. Principal component analysis of laser-microdissected RNAseq samples. Each dot represents a

laser-microdissected embryonic shoot sample. **b-f**. Pairwise comparison of the gene expression

between laser-microdissected RNAseq samples. The number of overlapping, differential-

expressed genes (DEGs) is showed in the Venn diagram. The circled numbers represent the

genes preferentially expressed in the SAM (**b**), upper coleoptile (**c**), scutellum (**d**), leaf (**e**) and

lower coleoptile (**f**).

1153

1154 **Extended Data Fig. 3 Multiplex *in situ* hybridization of *TRU1*.**

1155 The spatial expression of *TRU1* is showed in signal point map (**a-c**) and density map (**d-f**) from

1156 the transition stage (**a and d**), the coleoptilar stage (**b and e**) and the Stage 1 (**c and f**). The red

1157 arrow marks the expression of *TRU1* at the emerging leaf sheath. Scale bar = 100 μm.

1158

1159

1160

1161

1162

1163

1164

1165

1166

1167

1168

1169

1170

 Extended Data Fig. 4 Expression of the embryo organ initiation network.

a. WGCNA to identify modules with overall highest expression in the Stage 1 leaf (L) and less in

the lower coleoptile (LC), with steadily decreased expression in the upper coleoptile (UC) and

scutellum (S) (i.e. L > LC > UC > S). The heatmap shows the correlation coefficient (r) and the

corresponding p value (in the parentheses) of each module (ME1-12). **b**. The expression of the

130 genes in the initiation network at the transition stage (T), coleoptilar stage (C) and the Stage

1 scutellum (Scu), upper coleoptile (UC), lower coleoptile (LC) and leaf (L). **c-e**. The correlation

of the expression of the 130 genes between initiation stage of the scutellum, coleoptile and leaf.

1179

1180

1181

1182

1183

1184

1185

1186

1187

1188

1189

1190

1191

1192

1193

1194

Extended Data Fig. 5 WGCNA Module-organ correlation.

a-b. Correlation between datasets of the 14-day after pollination seedling and the embryo before

and after batch effect removal. The correlation of the normalized overall gene expression

between the 14 day after germination (DAG) seedling and embryo datasets before (**a**) and after

(**b**) batch effect removal. **c**. Heatmap illustrating the correlation between co-expression modules

(y-axis) and corresponding organs (x-axis. The colors are determined by the correlation

coefficient (r) with the p value (numbers in the parentheses)

1209

1210

1211 **Extended Data Fig. 6 Correlation between the maize initiation network genes and the**

1212 **Arabidopsis orthologs.**

1213 The correlation of expression between the maize initiation network genes at transition (**a**) and

1214 coleoptilar (**b**) stage, and the Arabidopsis orthologs at early heart stage.

Extended Data Fig. 7. TAI and TDI for embryo stages under different conditions

a-c. Distribution of all maize genes and genes in each embryo stage across phylostrata and TAI profiles for different E-value cut-offs (ranging from 1E-2 to 1E-20). This analysis investigates whether and how E-values affect the hourglass pattern. **a**. Variation of the E-value cut-off for the BLASTp analysis demonstrates minimal import on the distribution of maize genes among the phylostrata, indicating an overall pattern independent of the E-values. The y-axis is presented on a logarithmic scale. **b**. Similarly, the variation of the E-value cut-off for the BLASTp analysis does not much severely alter the distribution of maize genes in each embryo stage among the phylostrata. **c**. the hourglass pattern of the TAI profile remains consistent across various E-value cut-offs. **d-e**. Transcriptome divergence indices based on two different genome comparisons: *Zea mays* ssp. Mays vs. *Zea mays* ssp. mexicana (**d**); *Zea mays* ssp. mays vs. *Zea diploperennis* (**e**). The overall patterns of TDI profiles are significant, as measured by flat line test ($P_{\text{TDI-Mexicana}}=1.07\times10^{-3}$, $P_{\text{TDI-diploperennis}}=4.80\times10^{-4}$), reductive hourglass test ($P_{\text{TDI Mexicana}}=0.027$, $P_{\text{TDI-diploperennis}}=1.53\times10^{-3}$), and reductive early conservation test ($P_{\text{TDI Mexicana}}=0.68$, $P_{\text{TDI-diploperennis}}=0.998$).

Extended Data Fig. 8. The gene expression in each embryo stage per phylostratum.

a. The stacked bars in percentage of gene numbers of each phylostratum in embryo stages. **b-i.**

The relative gene expression (normalized to the maximal level in each embryo stage) of maize

embryo genes, including in proembryo stage (**b, c**), in Initiation-Network stages (transition stage,

coleoptilar stage, and stage 1 leaf) (**d, e**), in stage 1 (stage 1 leaf, stage 1 scutellum, stage 1

coleoptile) (**f, g**), and after stage 2 and in seedling leaves (**h, i**). The data to after stage 2 is from

Chen et al., 2014, and were normalized in data source in (**h**).