ROS-Induced Stress Promotes Enrichment and

Emergence of Antibiotic Resistance in Conventional

Activated Sludge Processes

*Bharat Manna[1], Xueyang Zhou[1], and Naresh Singhal[1,2] ***

[1]Department of Civil and Environmental Engineering, University of Auckland, Auckland 1142,

New Zealand

[2]Water Research Centre, University of Auckland, Auckland 1142, New Zealand

*Corresponding author: n.singhal@auckland.ac.nz

Abstract

Since the Great Oxidation Event 2.4 billion years ago, microorganisms have evolved
sophisticated responses to oxidative stress. These ancient adaptations remain relevant in modern
engineered systems, particularly in conventional activated sludge (CAS) processes, which serve
as significant reservoirs of antibiotic resistance genes (ARGs). While ROS-induced stress
responses are known to promote ARG enrichment/emergence in pure cultures, their impact on
ARG dynamics in wastewater treatment processes remains unexplored. Shotgun-metagenomics
analysis of two hospital wastewater treatment plants showed that only 35-53% of hospital
effluent resistome was retained in final effluent. Despite this reduction, approximately 30% of
ARGs in CAS showed higher abundance than upstream stages, of which 22% emerged *de novo*.
Beta-lactamases and efflux pumps constituted nearly 50% of these enriched ARGs. These ARGs
exhibited significant correlations ($p < 0.05$) with ROS stress response genes (*oxyR*, *soxR*, *sodAB*,
katG and *ahpCF*). The CAS resistome determined 58-75% of the effluent ARG profiles,
indicating treatment processes outweigh influent composition in shaping final resistome. Proof-
of-concept batch reactor experiments confirmed increased ROS and ARG levels under high
dissolved oxygen (8 mg/L) compared to low oxygen (2 mg/L). Untargeted metaproteomics
revealed higher expression of resistant proteins (*e.g.*, OXA-184, OXA-576, PME-1, RpoB2,
Tet(W/32/O)) under elevated ROS levels. Our findings demonstrate that CAS processes actively
shape effluent resistome through ROS-mediated selection, indicating that treatment processes,
rather than initial wastewater composition, determine final ARG profiles. This study indicates
that the emergence of ARGs needs to be considered as an integral aspect of wastewater treatment
design and operation to prevent antibiotic resistance dissemination.

33

34 **Keywords**: wastewater treatment, great oxygenation event, reactive oxygen species,

35 metagenomics, SWATH-metaproteomics

Highlights

- Approximately 30% of ARGs enriched in CAS versus upstream stages, of which 22% emerged *de novo*

- Beta-lactamases and efflux pumps comprise ~50% of enriched ARGs in CAS

- CAS resistome shapes 58-75% of effluent ARGs, outweighing influent composition

- High DO levels increase ROS production and ARG abundance in CAS systems

- Metaproteomics confirms higher expression of resistant proteins under elevated ROS levels

1. Introduction

Antibiotic resistance (AR) poses an escalating threat to global public health, responsible for approximately 1.27 million fatalities worldwide and contributing to almost an estimated 4·95 million (3·62–6·57) deaths associated with bacterial antimicrobial resistance in 2019 (Murray, Christopher JL, 2022). In the United States alone, infections afflict over 2.8 million individuals annually, leading to the deaths of more than 35,000 people (Centers et al., 2019). The financial burden of managing infections caused by six prevalent antibiotic-resistant bacteria (ARB) (Murray, Christopher JL, 2022) especially within healthcare settings exceeds $4.6 billion each year (Nelson et al., 2021). Antibiotic resistance is a critical One Health issue (McEwen and Collignon, 2018), profoundly affecting humans, animals, and the environment, impacting even the most fundamental resource for life: water.

The presence of antibiotic-resistant genes (ARGs) and ARBs often can be attributed to effluents from hospitals, farms, and sewage systems, illustrating the interconnectedness of human activities and the broader environmental impact of AR (Agga et al., 2015; Rowe et al., 2017). As wastewater treatment plants (WWTPs) operate at the human-environment interface, they play a crucial role in both managing and understanding the global health effects of AR (Alexander et al., 2020; Manaia et al., 2018). WWTPs receive and process diverse influents, including hospital effluents laden with high concentrations of antibiotics, ARBs, and ARGs (Y. Chen et al., 2020; Uluseker et al., 2021; Wang et al., 2022). This influx creates a potent selection pressure for ARGs and facilitates horizontal gene transfer (HGT) among bacterial communities (Karkman et al., 2018; Uluseker et al., 2021; Wei et al., 2021). Notably, hospital wastewaters harbor ARG concentrations several orders of magnitude higher than municipal wastewater (Korzeniewska et al., 2013), highlighting their alarming potential to seed and propagate resistance within WWTPs

treating such influents (Hocquet et al., 2016; Mehanni et al., 2023; Paulus et al., 2019).

Moreover, recent studies reported that different WWTP treatment processes could exhibit

varying impacts on ARG and ARB dynamics (Honda et al., 2023; Luo et al., 2021; Majeed et al.,

2021). Interestingly, the authors identified significant alteration in the ARGs composition in the

sludge compared to that of the influent resistome and suggested seasonality in antimicrobial use

to be key influencing factor behind this transition (Honda et al., 2023). A multi-omics study

revealed significant enrichment of both absolute and relative abundance of specific resistance

genes in CAS resistome compared to influent, particularly those conferring resistance to ARGs

drug classes (multidrug, beta-lactam, aminoglycoside, tetracycline, vancomycin etc.), suggesting

bacterial community composition and biomass concentration as key factors (Ju et al., 2019).

Moreover, another study showed that while the overall ARGs abundance decreased during 11 out

of 12 WWTPs, there were instances of specific ARGs being enriched in the effluents (Garner et

al., 2024). Many other reports also identified CAS processes as potential hotspots for ARG

enrichment (Bengtsson-Palme et al., 2016; Guo et al., 2017; Liu et al., 2018, 2019; Majeed et al.,

2021; Petrovich et al., 2020). These enriched ARGs in CAS contribute significantly to the

effluent resistome which is disseminated into the receiving environment (Honda et al., 2023; Ju

et al., 2019; Majeed et al., 2021). However, the specific mechanisms driving the enrichment and

emergence of ARGs within CAS processes remain poorly understood. Consequently,

understanding the underlying mechanisms behind the emergence of ARGs in CAS processes has

become an urgent need.

While antibiotics are relatively recent human inventions, microbes have developed resistance

mechanisms over thousands of years through their continuous evolutionary struggle for survival

(Dcosta et al., 2011). To understand bacterial resistance mechanisms, the perspective of oxygen

on Earth provides a critical evolutionary context. Prior to the Great Oxidation Event (GOE)

around 2.4-2.1 billion years ago, Earth's atmosphere was essentially oxygen-free, and early life

thrived in a reducing condition (Lyons et al., 2014). The dramatic rise of oxygen through

photosynthesis represented the first major oxidative stress that organisms had to adapt to, driving

the evolution of cellular mechanisms to cope with oxygen toxicity and its reactive species

(ROS). Recent evidence suggests a unifying mechanism behind the action of many bactericidal

antibiotics, where ROS generation plays a key role behind lethality (Dwyer et al., 2009; Lam et

al., 2020; Van Acker and Coenye, 2017). ROS like hydrogen peroxide (H_2O_2), hydroxyl radical

(OH^-), singlet oxygen ($^1O_2$), and superoxide anion (O_2^-) are inherent byproducts of bacterial

aerobic respiration (Imlay, 2019, 2003). Furthermore, ROS-mediated killing can trigger a

secondary stress response beyond the initial antibiotic-induced damage, involving factors like

activation of response regulators (*e.g.*, *ArcA*), metabolic reprogramming of the TCA cycle, and

disruption of the nucleotide pool with increased ATP demand (Li et al., 2021) (Figure S1). To

counter this oxidative stress and survive, bacteria have evolved diverse antioxidant defenses

through metabolic remodeling (Li et al., 2021). However, bacterial efforts to develop ROS

tolerance often come at a cost. The increased oxidative stress can lead to direct DNA damage,

potentially triggering novel mutations, and ultimately contributing to the emergence of AR

(Ariza et al., 1994; Jin et al., 2018; Mandsberg et al., 2009; Qi et al., 2023). Since ROS are

common to both antibiotic action and bacterial stress responses, resistance to ROS could be a key

shared characteristic underlying ARG development. We recently showed that ROS stress has

been linked to ARGs proliferation in freshwater microbial communities (Manna et al., 2024).

These ancient adaptations to oxidative stress remain fundamental to how modern

microorganisms handle ROS in highly aerated wastewater treatment systems. The significance of

ROS in CAS processes can be attributed to the constant aeration inherent to these systems, which creates a unique environment characterized by high dissolved oxygen (DO) levels and, consequently, elevated ROS generation. While existing research hints at the connection between ROS and ARG emergence in the context of WWTP, it primarily focuses on lab experiments and specific antibiotics or stressors. One study observed a significant increase in antibiotic resistance in *Escherichia coli* exposed to aeration tanks compared to primary or final sedimentation tanks (Sulfikar et al., 2018). Another study linked triclosan-induced oxidative stress to a rise in ARGs, particularly tetracycline and multidrug resistance genes (Tan et al., 2021). Similarly, ROS generated by phenolic compounds have been proven to promote resistance through increased membrane permeability and HGT (Ma et al., 2021). Moreover, antidepressant fluoxetine induced resistance to multiple antibiotics (*e.g.*, fluoroquinolone, aminoglycoside, β-lactams, tetracycline and chloramphenicol) in *Escherichia coli* via ROS-mediated mutagenesis (Jin et al., 2018). Despite these findings, the critical question of whether aeration-induced ROS in real-world WWTPs shape the multifaceted resistome in CAS processes, remains unanswered.

Building on this evolutionary and mechanistic foundation, we present a comprehensive investigation of how aeration-induced oxidative stress shapes antibiotic resistance in full-scale CAS processes. By combining whole-genome shotgun metagenomics analysis from two WWTPs treating hospital effluents and untargeted metaproteomics from controlled batch reactor experiments, we unravel the relationship between DO levels, ROS generation, and ARG dynamics, providing unprecedented insights into how oxidative stress drives resistance development in CAS processes.

2. Materials and methods

2.1. Sample data collection

In this work, we considered paired-end shotgun metagenomics samples obtained using a HiSeq

2500 instrument from two full-scale WWTPs processing university hospital effluents in

Göttingen (WWTP_Gö), and Greifswald (WWTP_Gw), Germany (Schneider et al., 2020). The

samples were collected from five different stages (abbreviated as S) from both the WWTPs, *i.e.*,

S1: Hospital effluent (a storage tank next to the university hospital), S2: WWTP Influent, S3:

Primary Sludge, S4: conventional activated sludge (CAS), and S5: WWTP effluent as described

elsewhere (Schneider et al., 2021, 2020). The details of the sample names, and NCBI SRA

accession used in this study, can be found in Table S1 and Table S2. The raw metagenomes can

be accessed in the NCBI Sequence Read Archive under the BioProject accession number

PRJNA524094 (Schneider et al., 2021).

2.2. Metagenomics Analysis

We performed preprocessing steps to eliminate low-quality sequences using Trimmomatic v0.39

(Bolger et al., 2014) with specific quality thresholds (Manna et al., 2024). For metagenomic

analysis, we employed the SqueezeMeta v1.5.2 pipeline (Tamames and Puente-Sánchez, 2019).

The read statistics per sample are provided in Table S1 and Table S2. Co-assembly was

performed using Megahit v1.2.9 (D. Li et al., 2015). Within the SqueezeMeta pipeline, Barrnap

v0.9 (Seemann, 2014) tool was used for predicting RNAs, while Prodigal v.2.6.3 (Hyatt et al.,

2010) was utilized for predicting ORFs. Moreover, the Comprehensive Antibiotic Resistance

Database (Alcock et al., 2020) release 3.2.4 was used for annotating the ARGs using Diamond

v.2.0.14 (Buchfink et al., 2014) with sequence identity of 70% and e-value threshold of 1×10^{-3}

(Manna et al., 2024; Petrovich et al., 2020). ARG abundances were normalized using the 16S-

rRNA gene abundance as per previous report (B. Li et al., 2015). ARGs conferring resistance to

more than one antibiotic was classified as 'multidrug resistant' (MDR). Similarity searches

against Kyoto Encyclopedia of Genes and Genomes (Kanehisa and Goto, 2000), were performed

for functional assignments of ORFs. The metagenomics data was analyzed using the SQMtools

v1.6.3 R package (Puente-Sánchez et al., 2020).

2.3. Bioinformatics Analysis and Visualization

To assess the diversity of ARG composition across the different treatment stages, we calculated a

Bray-Curtis dissimilarity matrix with PERMANOVA to determine the significance of clustering,

using MicrobiomeAnalyst v2.0 (Dhariwal et al., 2017). A heatmap was plotted to visualize

changes in the abundance of each ARG, utilizing a z-score with TBtools v2.007 (C. Chen et al.,

2020). Furthermore, differentially abundant (enriched/emerged) ARGs in CAS (S4) were

identified using EdgeR v.4.4.0 (Robinson Mark et al., 2010) with an adjusted p-value cutoff of

0.05. The correlation between ROS stressor genes and differentially abundant ARGs was

performed using a Mantel test (p-value < 0.05) using OmicStudio tools v1.41.0 (Lyu et al.,

2023). Graphics were generated using the ggplot2 v3.5.1 (Wickham, 2011) package in R (R Core

Team, 2022) version 4.2.1.

2.4. Batch reactor experiments with different DO levels

To validate our metagenomics-based findings of higher ROS generation under more oxidative

stress leading to increased ARGs, we performed batch reactor experiments with CAS samples

under low (2 mg/L) and high (8 mg/L) DO levels. The CAS samples were obtained from the

Māngere WWTP in Auckland, New Zealand. The experiments were conducted in reactors (n = 3

biological replicates per condition) with a capacity of 1 L, at a temperature of 20 °C for 48 hours.

Composition of the synthetic wastewater is provided in Table S3 and Table S4. Detailed

methodology for ROS measurement, DNA extraction for metagenomics analysis and untargeted

metaproteomics analysis are provided in Supplementary Methods. The raw metagenome samples

from these experiments can be accessed in the European Nucleotide Archive under the

BioProject accession number PRJEB74089 as provided in Table S5.

2.5. SWATH-MS based quantitative Metaproteomics Analysis

Untargeted metaproteomic analysis was performed on activated sludge samples to characterize

microbial protein expression under the two DO conditions. Proteins were extracted using SDS-

based lysis buffer, followed by sonication and TCA-acetone precipitation. Tryptic-digested

peptides were analyzed using a NanoLC 400 UPLC (Eksigent, USA) system coupled to a

TripleTOF 6600 mass spectrometer (Sciex, USA). Protein identification using

MetaProteomeAnalyzer v3.4 (Muth et al., 2015) with X-tandem revealed 1,110 distinct proteins

for functional analysis. Detailed methodology is available in Supplementary Methods. The

metaproteomics data has been submitted to ProteomeXchange Consortium with the dataset

identifier PXD044490.

3. Results and Discussion

3.1. Comparison of ARG abundances across different treatment stages

In this study, we systematically analyzed and compared metagenomic samples from various

stages of wastewater treatment processes to assess changes in the resistome profile. These stages

included hospital effluents (S1), WWTP influent (S2), primary treatment (S3), CAS (S4), and

WWTP effluents (S5) (Figure 1a). Our findings showed a considerable decrease in total ARG

abundances at different stages of WWTP$_{Gö}$ compared to the hospital effluents: 54.17% in S2,

53.38% in S3, 78.69% in S4, and 83.30% in S5 (Figure 1b). Moreover, 69.50%, 68.42%,

46.59%, and 35.14% of the ARGs detected in S1 were retained in S2, S3, S4 and S5,

respectively (Figure S2a).

Figure 1. Alteration of ARG abundance across different WWTP stages. (a) Schematic representation of the WWTP stages studied, including S1: Hospital effluent, S2: WWTP influent, S3: Primary sludge, S4: Activated sludge, and S5: WWTP effluent. The variation in ROS stressor levels across these stages is depicted below. (b) and (c) show the total relative ARG abundance at each stage for WWTP$_{Gö}$ and WWTP$_{Gw}$, respectively. Heatmaps display the trend of change in each ARG relative abundance across the stages in (d) WWTP$_{Gö}$ and (e) WWTP$_{Gw}$ with a prominent set of enriched (or emerged) ARGs (red) in both S4 and S5. RAS: Recycled activated sludge.

A similar analysis conducted in the WWTP$_{Gw}$ also revealed a reduction in ARG abundances of

39.70% in S2, 40.65% in S3, 63.55% in S4, and 61.99% in S5, compared to S1 (Figure 1c). The

numbers of ARGs detected in S1 were retained by 73.14% in S2, 67.55% in S3, 31.52% in S4,

and 53.42% in S5 (Figure S2c). The highest abundance of ARGs was observed in the hospital

effluent (S1) for both WWTPs, which is expected due to the prevalent use of antibiotics in

hospital settings, leading to a higher presence of ARGs. Across the treatment stages, a consistent

trend of ARG reduction was noted, indicating the effectiveness of the WWTP processes in

decreasing the overall ARG abundance in the effluent (S5). This pattern of ARG reduction in

CAS and effluents are in agreement with previous metagenomics-based studies (Bengtsson-

Palme et al., 2016; Honda et al., 2023; Lira et al., 2020).

Furthermore, the diversity of resistome profiles across various stages of the WWTP$_{Gö}$ and

WWTP$_{Gw}$ was evaluated using principal coordinate analysis (PCoA) with a Bray-Curtis

dissimilarity matrix (Figure S2b, d). The PCoA demonstrated clear clustering of resistomes

corresponding to the five treatment stages. Significant separation observed (R^2: 0.79329,

PERMANOVA F-value: 14.392, *p*-value: 0.001) in WWTP$_{Gö}$, indicating distinct ARG profiles

at each treatment stage. Similarly, WWTP$_{Gw}$'s resistome profiles were markedly segregated

demonstrating substantial differentiation (R^2: 0.83563, PERMANOVA F-value: 19.064, *p*-value:

0.001). The PCoA results showed that the resistomes from S1, S2, and S3 stages clustered

closely, suggesting similar ARG profiles, possibly due to the direct contribution of hospital

effluent to the influent. In contrast, samples from the CAS (S4) and treated effluent (S5) stages

were distinctly separate from the preceding stages. This delineation reflects a significant

transition in the ARG profile, attributable to the biological processes and selection pressure

within the activated sludge system and secondary clarifiers. Notably, the effluent resistome

consisted of 58.42% and 75.58% of the CAS resistome in the WWTP$_{Gö}$ and WWTP$_{Gw}$,

respectively, indicating a substantial overlap between the systems. The distinct clusters for the

S4 and S5 stages also suggest that the treatment process, rather than initial wastewater

composition, determines final effluent ARG profiles, with potential environmental implications

upon its release. Similar variations in resistome in influent, activated sludge and effluent have

been reported in previous studies (Honda et al., 2023; Majeed et al., 2021).

In addition to the overall trend analysis, a detailed heatmap of all ARGs based on normalized

abundance (z-scores) was generated to further elucidate the specific changes in ARG profiles at

each treatment stage, with a focus on the CAS (S4) process (Figure 1d, e). The heatmap

illustrated that while some ARGs were consistently abundant across all stages, others were more

stage specific. This abundance pattern reinforces the notion of stage-specific selective pressures

and microbial community dynamics shaping the ARG profile. Interestingly, despite the general

trend of decreasing total ARG abundance (Figure 1b, c), certain ARGs exhibited notable

persistence and remarkable enrichment/emergence in the CAS stage (S4) in both the WWTPs

(Figure 1d, e). This relative enrichment of ARGs in the CAS was previously reported using

combined metagenomics and metatranscriptomics study (Ju et al., 2019).

3.2. Identifying the enriched and emerged ARGs in the CAS process

To enhance our understanding of the enriched or emerged ARGs within the CAS process, we

conducted a detailed analysis of ARGs that were differentially more abundant in CAS than in the

upstream stages. Specifically, in the WWTP$_{Gw}$ facility, we observed that 111, 115, and 116

ARGs exhibited significantly (FDR < 0.05) higher abundance in CAS when compared to S1, S2,

and S3, respectively, as illustrated in Figure 2a. Similarly, for the WWTP$_{Gö}$ facility, 152, 138,

and 114 ARGs were found to be more abundant in CAS relative to the same comparator stages

(Figure S3a). Additionally, certain ARGs were identified as being significantly more abundant in

S4 compared to all other stages (S1, S2, and S3), with some ARGs being uniquely abundant to

specific treatment stages (Figure S4 and Supplemental Data 1). To facilitate downstream

analyses, we compiled a list of ARGs that were significantly abundant in the CAS process,

selecting those genes that displayed a higher average abundance in CAS compared to all

preceding treatment stages. This analysis yielded a total of 116 (29.59%) and 94 (36.43%) highly

abundant ARGs for WWTP$_{Gö}$ and WWTP$_{Gw}$, respectively, as detailed in Supplemental Data 2a.

Notably, 22.22% (WWTP$_{Gö}$) and 21.88% (WWTP$_{Gw}$) of the abundant ARGs were not detected

at any of the upstream stages, suggesting they emerged under the selection pressure of the CAS

process (Supplemental Data 2b). These emerged ARGs exhibited resistance to aminoglycoside

(*AAC(3)-IIIc*, *AAC(6')-Iih*), beta-lactam (*OXA-34*, *OXA-660*, *OXA-4*, *GOB-3*, *GOB-4*, *IDC-2*,

IMP-1, *IMP-13*, *IMP-28*), rifamycin (*iri*, *arr-1*), tetracycline (*tap*, *tet(V)*), glycopeptide (*vanD*,

vanI), and other antibiotics. Additionally, among the enriched ARGs in both WWTPs were those

belonging to various beta-lactamases, such as *OXA* (including *OXA-45*, *OXA-101*, *OXA-129*,

OXA-198) beta-lactamases. These ARGs confer resistance to antibiotics like carbapenems,

cephalosporins, and penams, highlighting a significant selection pressure for beta-lactam

resistance. This is especially concerning given the extensive use of beta-lactam antibiotics and

the potential for these resistance mechanisms to spread into the environment. Furthermore, we

identified multiple ARGs conferring resistance to other antibiotic classes, including tetracycline

(*tet(35)*, *tet(49)*, *tet(G)*, *tet(T)* etc.), glycopeptide (*vanC*, *vanF*, *vanS_in_vanC_cl*, etc.), and

aminoglycoside (*AAC(3)-Id*, *AAC(6')-32*, *AAC(6')-Ic*, *AAC(6')-IIa*, *AAC(6')-IIb*, *AAC(6')-IIc*,

etc.), among others. Resistance to rifamycin (*rpoB2*, *Bado_rpoB_RIF*, *Nfar_rox*, *rphA*, *rphB*,),

macrolide (*EreB*, *mphA*), fluoroquinolone (*QepA4*, *QnrS6*, *QnrS8*), and sulfonamide (*sul2*, *sul4*)

antibiotics were also observed, alongside ARGs resistant to multiple antibiotics, such as *AxyY*,

efrB, *MexF*, *mexP*, *mexQ*, *mexY*, *opmE*, *smeE*, and more, as provided in Supplemental Data 2a.

Amongst the enriched/emerged ARGs, beta-lactamases (WWTPGö: 24.14%, WWTPGw:

25.53%) and efflux pumps (WWTPGö: 23.28%, WWTPGw: 27.66%) represented the

predominant resistance mechanisms in both WWTPs. Our recent study also demonstrated that

oxidative stress conditions could exacerbate ARG proliferation, particularly efflux pump genes

(Manna et al., 2024). Consistent with our observations, a significant enrichment of various

resistance genes, including those encoding for beta-lactamases, macrolides, and tetracyclines,

was identified in recirculated activated sludge in comparison to primary sludge (Bengtsson-

Palme et al., 2016). This enrichment was suggested due to the selective pressure within the

wastewater treatment facility. Similarly, previous research documented an escalation in the

relative abundance of a wide array of ARGs, spanning multiple drug classes such as beta-

lactamases, aminocoumarins, glycopeptides, rifamycins, and sulfonamides, from influent to

effluent in a conventional WWTP (Majeed et al., 2021). Furthermore, another study reported an

increase in both the transcript and gene abundance of various ARG classes, including Multidrug,

beta-lactamases, aminoglycosides, tetracyclines, trimethoprim, and vancomycin, in secondary

effluent compared to the influent of post-primary clarifiers (Ju et al., 2019). Additionally, it was

found that aerobic biomass exhibited a higher overall abundance of ARGs compared to

anaerobic and anoxic biomass, with a notable increase in effluent abundances of fluoroquinolone

and tetracycline ARGs (Petrovich et al., 2020). These observations underscore the diverse

spectrum of resistant ARGs emerging within the CAS process, indicating that while the CAS

process effectively reduces the overall ARG load, it also inadvertently facilitates the proliferation

(enrichment and emergence) of multiple concerning ARGs.

Figure 2. Differentially enriched (or emerged) ARGs and their correlation network with ROS stressors in the CAS process. a) Volcano plots illustrate the differentially abundant ARGs in the CAS stage compared to S1, S2, and S3 in the WWTP$_{Gw}$ (log$_2$FC $\geq$ 1, FDR < 0.05). b) A network analysis depicting the correlations between selected enriched/emerged ARGs and ROS stress response genes in CAS process in WWTP$_{Gw}$.

3.3. Assessing the correlation between the ROS stressors and enriched or emerged ARGs

To further assess previously established connections between oxidative stress and antibiotic resistance, we analyzed correlations between ROS-response genes and the enriched/emerged ARGs detected in the CAS systems of both WWTPs. We focused on five crucial ROS-related

genes (Seo et al., 2015): two transcription factors, *oxyR* and *soxR*, which play key roles in the

cellular response to oxidative stress, and three genes encoding enzymes that help detoxify ROS,

namely *sodAB* (Superoxide dismutase), *katG* (Catalase-peroxidase), and *ahpCF* (alkyl

hydroperoxide reductase). Notably, previous reports revealed *oxyR* (Lau et al., 2005; Wei et al.,

2012) and *soxRS* (P. Wang et al., 2020) contributed to virulence in *E. coli*, *Pseudomonas*

aeruginosa, and *Salmonella enterica*. Our analysis revealed a significant correlation between

these ROS stress response genes and numerous ARGs in both WWTPs (Figure 2b, Figure S3b,

Supplemental Data 3). Specifically, in the WWTP$_{Gö}$, the number of ARGs positively correlated

with each ROS stressor were *oxyR* (7), *soxR* (35), *sodAB* (22), *katG* (31), and *ahpCF* (17).

Meanwhile, in WWTP$_{Gw}$, the numbers of ARGs linked to the same stressors were *oxyR* (60),

soxR (49), *sodAB* (52), *katG* (79), and *ahpCF* (17). In the case of WWTP$_{Gö}$, we identified several

ARGs showing correlations with different combinations of ROS stressors (Figure S5a). Most

notably, 4 ARGs (*AAC(6')-IIb*, *lmrC*, *OXA-4*, and *QepA4*) correlated with all ROS stressors

examined. The largest individual correlations were observed with *soxR*, which correlated with 10

ARGs, including tetracycline resistance genes (*tet(T)*, *tet(W/32/O)*), and various others

(*Abau_AbaQ*, *dfrG*, *HelR*, *mexN*, *otr(A)S.liv*, *TriC*, *vgaALC*). Furthermore, a significant group of

6 ARGs, including metallo-beta-lactamases (*IMP-13*, *IMP-28*), beta-lactamases (*OXA-9*), and

others (*aadA8*, *arr-1*, *vanS_in_vanO_cl*), showed correlation specifically with both *sodAB* and

katG, indicating a potential link between these antioxidant enzymes and antibiotic resistance

mechanisms. Additional notable correlations were observed with *katG* alone, which correlated

with 9 ARGs, including various resistance mechanisms (*adeF*, *dfrB2*, *mphA*, *MYO-1*, *OXA-198*,

smeE, *vanD*, *vanS_in_vanP_cl*, *vgaD*), suggesting a broad spectrum of antibiotic resistance

mechanisms being influenced by oxidative stress responses. Besides, within the WWTP$_{Gw}$, 10

ARGs were correlated with all ROS stressors, including genes responsible for multidrug

resistance (*adeF*), beta-lactamase activity (*OXA-4*, *OXA-5*, *PAU-1*, *IMP-5*), tetracycline

resistance (*tet(T)*, *tetB(P)*), and rifamycin resistance (*rpoB2*), among others (*rsmA*, *arr-4*) (Figure

S5b). Additionally, a noteworthy observation was the correlation of a considerable number of

ARGs with multiple ROS stressors, specifically 24 ARGs correlating with *soxR*, *oxyR*, *sodAB*,

and *katG* (including *AxyY*, *MuxC*, *OXA-145*, *mexY*, and others), and 11 ARGs correlating with

soxR, *oxyR*, and *katG* (including *mtrA*, *vanD*, *mphA*, and others). Comparative analysis of both

WWTPs revealed many common ARGs showing correlations with various ROS stressors,

including genes encoding beta-lactamases (*OXA-4*, *PAU-1*), tetracycline resistance (*tet(T)*,

tetA(60)), rifamycin resistance (*rpoB2*), and various multidrug resistance mechanisms (*adeF*,

AxyY, *smeE*), suggesting shared patterns of ROS-mediated antibiotic resistance regulation across

different wastewater treatment environments. These observed patterns align with and extend

previous research findings in the field. For instance, studies with *Escherichia coli* K12

demonstrated that strains resistant to multiple antibiotics (chloramphenicol, amoxicillin and

tetracycline) produced significantly higher levels of ROS compared to the wild-type (Jin et al.,

2018). Additionally, the resistant strains also exhibited significant overexpression of ROS

scavengers including *sodA* and *ahpF*. A recent study showed that exposure to bactericidal

antibiotics (amoxicillin, enrofloxacin, and kanamycin) increased ROS production in the *oxyR*

knocked out *E. coli* compared to the wild-type (Qi et al., 2023). Moreover, heme induced

electron transport chain enhanced the ROS production, leading to *de novo* acquired resistance in

Lactobacillus lactis (Qi et al., 2023). These reports support our findings on the pattern of

correlations and underscore the potential link between ROS stressors and ARG emergence. A

notable aspect of our findings is the identification of common and unique ARGs across different

ROS stressors. Certain ARGs appear frequently across multiple ROS stressor correlations,

suggesting that these ARGs might be more universally responsive to ROS-induced stress. In

contrast, unique ARGs correlating with specific ROS stressors indicate targeted responses to

particular oxidative stress conditions. Thus far, through meticulous examination of ROS stressor

genes and their correlations with differentially enriched ARGs in the CAS aeration process, we

uncovered patterns suggesting ROS as an instrumental factor in driving ARG

enrichment/emergence.

3.4. Experimental support for increased ARG abundance and ROS under high DO

To investigate the mechanistic relationship between ROS and ARG enrichment/emergence

within the CAS process, we conducted batch reactor experiments using CAS samples from the

Māngere WWTP (Auckland, New Zealand). This experimental setup was designed to simulate

the oxidative conditions characteristic of the CAS process, with specific emphasis on DO

concentration variations. Our experimental design incorporated two distinct DO concentrations:

DO2 (2 mg/L) and DO8 (8 mg/L), enabling assessment of ROS generation under elevated DO

conditions. Quantitative analysis revealed mean H_2O_2 concentrations of 32.05 ± 2.08 μM and

38.38 ± 0.86 μM for DO2 and DO8 conditions, respectively (Figure 3a). The DO8 treatment

yielded significantly higher H_2O_2 production compared to DO2 ($p < 0.05$). Analysis of oxidative

stress response genes (*soxR*, *oxyR*, *sodAB*, *katG*, and *ahpCF*) demonstrated consistent

upregulation under DO8 conditions (Figure 3b), indicating systematic adaptation to elevated

ROS levels associated with higher DO concentrations. Resistome profiling revealed ARG

patterns consistent with those observed in both WWTP_Gö and WWTP_Gw CAS systems (Figure

S6a, b). The prevalent ARG classes encompassed resistance to aminocoumarin antibiotics,

aminoglycosides, beta-lactams (including carbapenems, cephalosporins, penams),

fluoroquinolones, macrolides, tetracyclines, peptide antibiotics, rifamycins, and sulfonamides

(Figure S6c). Differential abundance analysis identified 11 significantly enriched ARGs under

DO8 conditions (Figure 3c), including beta-lactamases (*OXA-34*), sulfonamide resistance (*sul4*),

tetracycline resistance (*tet(43)*, *tet(W/32/O)*), rifamycin resistance (*Bado_rpoB_RIF*, *rphA*), and

MDR (*mexQ*).

Figure 3. DO concentration influences ROS generation and ARG abundance. a) Violin plots showing H_2O_2
concentrations under DO2 (2 mg/L) and DO8 (8 mg/L) conditions (*$p < 0.05$, paired t-test). b) Relative abundance
of oxidative stress response genes under DO2 and DO8 conditions. c) Volcano plot of differentially abundant ARGs
comparing DO8 versus DO2 conditions ($\log_2 FC \geq 1.5$, FDR < 0.05).

To establish causality beyond correlation and verify protein expression, we performed untargeted

SWATH metaproteomics analysis. Results demonstrated enhanced expression of several ARGs

under DO8 conditions, including beta-lactamases (OXA-184, OXA-576, PME1), rifamycin

resistance (RpoB2), tetracycline resistance (Tet(W/32/O)), and aminoglycoside resistance

(AAC(6')-Ia) (Figure 4). While some proteins showed elevated expression under DO2 conditions

(Bpse_Omp38, K15582), others exhibited comparable expression levels across both conditions

(BUT-1, JOHN-1, OXA-153). These findings provide mechanistic insights into ARG dynamics

within CAS processes. The correlation between elevated ROS levels under high DO conditions

and upregulation of oxidative stress response genes support that increased ROS production,

resulting from intensive aeration in CAS systems, may contribute to ARG emergence and

persistence.

Figure 4. Heatmap with the protein expression of ARGs under DO2 and DO8 conditions measured by SWATH-MS

based metaproteomics analysis. The protein names refer to the ARGs from CARD and KEGG orthologs.

3.5. Mechanism of ROS-induced ARG enrichment/emergence in CAS

Sublethal concentrations of endogenous ROS can trigger a series of direct or indirect cellular

responses that may contribute to the enrichment/emergence of ARGs in the CAS process (Li et

al., 2021). These responses include the activation of transcriptional regulators, expression of

oxidative defense enzymes, metabolic adaptations, DNA damage, activation of efflux pumps,

cell damage, and increased permeability and transformation potential (Figure S1). Firstly,

enhanced expression of transcription factors like *oxyR* and *soxRS* facilitate the production of

ROS-scavenging enzymes such as superoxide dismutase, catalase, thioredoxins and others

(Imlay, 2008). In both the WWTP$_{Gö}$ and WWTP$_{Gw}$ CAS processes, we observed a high

abundance of these transcription factors and ROS scavengers (Figure 5 a,b). Secondly, ROS

stress can disrupt the TCA cycle, leading to hyperactivation of respiration, as seen in certain

antibiotic treatments (Kohanski et al., 2008). Previous reports also suggested the disruption of

the nucleotide pool, leading to the increase of ATP demand elevating TCA cycle activity and

cellular respiration under quinolones treatments (Yang et al., 2019). This disruption may trigger

metabolic adaptations, such as activating the glyoxylate shunt (GS) (Lemire et al., 2017) to

reduce oxidative burden or redirecting flux to the PPP for increased NADPH production,

supporting antioxidant defense systems (Christodoulou et al., 2018). Our comparative analysis

revealed an higher abundance of genes involved in these metabolic pathways, ATP and NADPH

production, and consumption in the CAS system compared to other treatment processes (Figure 5

a,b). Moreover, sublethal ROS levels can cause DNA damage and potential mutations,

facilitating the emergence of *de novo* antibiotic resistance (Qi et al., 2023). Notably, DNA repair-

related genes were abundant in the CAS processes at both WWTPs. We also identified several

ARGs in the CAS process that were not detected in the upstream processes of both WWTPs

(Supplemental Data 2b). This indicates that ROS-induced mutations could be responsible for

their emergence. As a secondary defense, bacteria activate multidrug efflux pumps to expel

antibiotics from the cell, promoting antibiotic tolerance. The abundance of multidrug efflux

pump genes was high in the CAS processes (Figure 5 a,b). Additionally, oxidative damage can

weaken the bacterial cell envelope, potentially leading to the release of ARGs into the

environment. This process, coupled with enhanced transformation capabilities under oxidative

stress, increases the uptake of foreign DNA, including ARGs, through HGT. Increased cell

permeability can promote the uptake of foreign ARGs and eDNA through transformation (Y.

Wang et al., 2020). Previous reports suggested that *Streptococcus* sp. released eDNA under H_2O_2

stress (Itzek et al., 2011). A recent report revealed that phenolic compounds can promote ROS-

induced increased cell permeability and HGT (Ma et al., 2021). These simultaneous events

occurring in the CAS process under oxidative stress conditions may play a significant role in

enriching ARGs or helping ARG emergence. Overall, the proposed mechanisms by which ROS

can contribute to ARG enrichment/emergence in the CAS process include activation of oxidative

stress responses and metabolic adaptations, DNA damage and *de novo* resistance development,

increased efflux pump activity, and enhanced cell permeability promoting horizontal gene

transfer facilitated by oxidative damage. This multifactorial process underscores the complex

interplay between oxidative stress and antibiotic resistance in the CAS process.

Figure 5. Heatmap illustrating the change in gene abundances associated with the cellular processes across various
stages of the (a) WWTP$_{Gö}$, and (b) WWTP$_{Gw}$.

4. Conclusions

Our findings reveal a crucial paradox in CAS processes: while they effectively reduce overall

ARG loads from hospital wastewater (retaining only 35-53% in effluent), they simultaneously

create conditions that promote enrichment and emergence of specific ARGs. The demonstration

that 30% of ARGs in CAS show higher abundance, with 22% of these enriched ARGs emerging

de novo, challenges the traditional view of treatment efficiency. Beta-lactamases and efflux

pumps, constituting nearly 50% of enriched ARGs, highlight the selective pressures within CAS

systems. Most significantly, our finding that CAS resistome determines 58-75% of effluent ARG

profiles indicates that treatment processes, rather than influent composition, shape the final

resistome. The mechanistic link between ROS generation and ARG proliferation, validated

through batch experiments, demonstrates how ancient oxidative stress responses continue to

influence modern antibiotic resistance dissemination through engineered systems. Our findings

highlight the need to consider AMR emergence as an integral component of the design and

operation of biological wastewater treatment processes.

Associated content

Supporting Information

All the supplementary methods, tables and figures are provided in Supporting Information.pdf.

Additionally, detailed information on differentially abundant ARGs, ARG family and drug

classes, and ARG correlation with ROS genes are provided in Supporting Data.xlsx.

Author information

Corresponding Author

Naresh Singhal* Phone: +64 9 923 4512; e-mail: n.singhal@auckland.ac.nz

ORCID

Bharat Manna: 0000-0001-7394-8204

Naresh Singhal: 0000-0002-9461-688X

Xueyang Zhou: 0000-0003-2591-0783

CRediT authorship contribution statement

BM: Conceptualization, Investigation, Methodology, Formal Analysis, Writing - original
draft. XZ: Investigation, Methodology, Writing - review & editing. NS: Conceptualization,
Funding acquisition, Supervision, Writing - review & editing.

Funding Sources

The study was funded by Marsden Award from the Royal Society of New Zealand
(Contract Number: MFP-UOA2018) made to NS.

Notes

The authors declare no competing financial interest.

Acknowledgment

The authors acknowledge the use of New Zealand eScience Infrastructure (NeSI) high
performance computing facilities, consulting support and/or training services as part of this
research. New Zealand's national facilities are provided by NeSI and funded jointly by NeSI's
collaborator institutions and through the Ministry of Business, Innovation & Employment's
Research Infrastructure programme. URL https://www.nesi.org.nz. The authors also
acknowledge the Centre for eResearch at the University of Auckland for their help in facilitating
this research. http://www.eresearch.auckland.ac.nz.

REFERENCES

Agga, G.E., Arthur, T.M., Durso, L.M., Harhay, D.M., Schmidt, J.W., 2015. Antimicrobial-
resistant bacterial populations and antimicrobial resistance genes obtained from
environments impacted by livestock and municipal waste. PLoS One.
https://doi.org/10.1371/journal.pone.0132586

Alcock, B.P., Raphenya, A.R., Lau, T.T.Y., Tsang, K.K., Bouchard, M., Edalatmand, A., Huynh,
W., others, 2020. CARD 2020: Antibiotic resistome surveillance with the comprehensive
antibiotic resistance database. Nucleic Acids Res. https://doi.org/10.1093/nar/gkz935

Alexander, J., Hembach, N., Schwartz, T., 2020. Evaluation of antibiotic resistance
dissemination by wastewater treatment plant effluents with different catchment areas in
Germany. Sci. Rep. https://doi.org/10.1038/s41598-020-65635-4

Ariza, R.R., Cohen, S.P., Bachhawat, N., Levy, S.B., Demple, B., 1994. Repressor mutations in
the marRAB operon that activate oxidative stress genes and multiple antibiotic resistance in
Escherichia coli. J. Bacteriol. https://doi.org/10.1128/jb.176.1.143-148.1994

Bengtsson-Palme, J., Hammarén, R., Pal, C., Östman, M., Björlenius, B., Flach, C.F., Fick, J.,
Kristiansson, E., Tysklind, M., Larsson, D.G.J., 2016. Elucidating selection processes for
antibiotic resistance in sewage treatment plants using metagenomics. Sci. Total Environ.
https://doi.org/10.1016/j.scitotenv.2016.06.228

Bolger, A.M., Lohse, M., Usadel, B., 2014. Trimmomatic: A flexible trimmer for Illumina
sequence data. Bioinformatics. https://doi.org/10.1093/bioinformatics/btu170

Buchfink, B., Xie, C., Huson, D.H., 2014. Fast and sensitive protein alignment using
DIAMOND. Nat. Methods. https://doi.org/10.1038/nmeth.3176

Centers, U.S., Control, D., Fattouh, M., El-Din, A.N., Davies, J., 2019. Centers for Disease
Control and Prevention. Antibiotic resistance threats in the United States, 2019. Int. J. Curr.
Microbiol. Appl. Sci.

Chen, C., Chen, H., Zhang, Y., Thomas, H.R., Frank, M.H., He, Y., Xia, R., 2020. TBtools: An
Integrative Toolkit Developed for Interactive Analyses of Big Biological Data. Mol. Plant.
https://doi.org/10.1016/j.molp.2020.06.009

Chen, Y., Shen, W., Wang, B., Zhao, X., Su, L., Kong, M., Li, H., Zhang, S., Li, J., 2020.
Occurrence and fate of antibiotics, antimicrobial resistance determinants and potential
human pathogens in a wastewater treatment plant and their effects on receiving waters in
Nanjing, China. Ecotoxicol. Environ. Saf. https://doi.org/10.1016/j.ecoenv.2020.111371

Christodoulou, D., Link, H., Fuhrer, T., Kochanowski, K., Gerosa, L., Sauer, U., 2018. Reserve
Flux Capacity in the Pentose Phosphate Pathway Enables Escherichia coli's Rapid
Response to Oxidative Stress. Cell Syst. https://doi.org/10.1016/j.cels.2018.04.009

Dcosta, V.M., King, C.E., Kalan, L., Morar, M., Sung, W.W.L., Schwarz, C., Froese, D., Zazula,
G., Calmels, F., Debruyne, R., Golding, G.B., Poinar, H.N., Wright, G.D., 2011. Antibiotic
resistance is ancient. Nature. https://doi.org/10.1038/nature10388

Dhariwal, A., Chong, J., Habib, S., King, I.L., Agellon, L.B., Xia, J., 2017. MicrobiomeAnalyst:
A web-based tool for comprehensive statistical, visual and meta-analysis of microbiome
data. Nucleic Acids Res. https://doi.org/10.1093/nar/gkx295

Dwyer, D.J., Kohanski, M.A., Collins, J.J., 2009. Role of reactive oxygen species in antibiotic
action and resistance. Curr. Opin. Microbiol. https://doi.org/10.1016/j.mib.2009.06.018

Garner, E., Maile-Moskowitz, A., Angeles, L.F., Flach, C.-F., Aga, D.S., Nambi, I., Larsson,

D.G.J., Bürgmann, H., Zhang, T., Vikesland, P.J., Pruden, A., 2024. Metagenomic Profiling
of Internationally Sourced Sewage Influents and Effluents Yields Insight into Selecting
Targets for Antibiotic Resistance Monitoring. Environ. Sci. \& Technol. 58, 16547–16559.

Guo, J., Li, J., Chen, H., Bond, P.L., Yuan, Z., 2017. Metagenomic analysis reveals wastewater
treatment plants as hotspots of antibiotic resistance genes and mobile genetic elements.
Water Res. https://doi.org/10.1016/j.watres.2017.07.002

Hocquet, D., Muller, A., Bertrand, X., 2016. What happens in hospitals does not stay in
hospitals: antibiotic-resistant bacteria in hospital wastewater systems. J. Hosp. Infect.
https://doi.org/10.1016/j.jhin.2016.01.010

Honda, R., Matsuura, N., Sorn, S., Asakura, S., Morinaga, Y., Van Huy, T., Sabar, M.A.,
Masakke, Y., Hara-Yamamura, H., Watanabe, T., 2023. Transition of antimicrobial
resistome in wastewater treatment plants: impact of process configuration, geographical
location and season. npj Clean Water. https://doi.org/10.1038/s41545-023-00261-x

Hyatt, D., Chen, G.L., LoCascio, P.F., Land, M.L., Larimer, F.W., Hauser, L.J., 2010. Prodigal:
Prokaryotic gene recognition and translation initiation site identification. BMC
Bioinformatics. https://doi.org/10.1186/1471-2105-11-119

Imlay, J.A., 2019. Where in the world do bacteria experience oxidative stress? Environ.
Microbiol. https://doi.org/10.1111/1462-2920.14445

Imlay, J.A., 2008. Cellular defenses against superoxide and hydrogen peroxide. Annu. Rev.
Biochem. https://doi.org/10.1146/annurev.biochem.77.061606.161055

Imlay, J.A., 2003. Pathways of Oxidative Damage. Annu. Rev. Microbiol.
https://doi.org/10.1146/annurev.micro.57.030502.090938

Itzek, A., Zheng, L., Chen, Z., Merritt, J., Kreth, J., 2011. Hydrogen peroxide-dependent DNA
release and transfer of antibiotic resistance genes in Streptococcus gordonii. J. Bacteriol.
https://doi.org/10.1128/JB.05791-11

Jin, M., Lu, J., Chen, Z., Nguyen, S.H., Mao, L., Li, J., Yuan, Z., Guo, J., 2018. Antidepressant
fluoxetine induces multiple antibiotics resistance in Escherichia coli via ROS-mediated
mutagenesis. Environ. Int. https://doi.org/10.1016/j.envint.2018.07.046

Ju, F., Beck, K., Yin, X., Maccagnan, A., McArdell, C.S., Singer, H.P., Johnson, D.R., Zhang,
T., Bürgmann, H., 2019. Wastewater treatment plant resistomes are shaped by bacterial
composition, genetic exchange, and upregulated expression in the effluent microbiomes.
ISME J. https://doi.org/10.1038/s41396-018-0277-8

Kanehisa, M., Goto, S., 2000. KEGG: Kyoto Encyclopedia of Genes and Genomes. Nucleic
Acids Res. https://doi.org/10.1093/nar/28.1.27

Karkman, A., Do, T.T., Walsh, F., Virta, M.P.J., 2018. Antibiotic-Resistance Genes in Waste
Water. Trends Microbiol. https://doi.org/10.1016/j.tim.2017.09.005

Kohanski, M.A., Dwyer, D.J., Wierzbowski, J., Cottarel, G., Collins, J.J., 2008. Mistranslation of
Membrane Proteins and Two-Component System Activation Trigger Antibiotic-Mediated
Cell Death. Cell. https://doi.org/10.1016/j.cell.2008.09.038

Korzeniewska, E., Korzeniewska, A., Harnisz, M., 2013. Antibiotic resistant Escherichia coli in
hospital and municipal sewage and their emission to the environment. Ecotoxicol. Environ.
Saf. https://doi.org/10.1016/j.ecoenv.2013.01.014

Lam, P.L., Wong, R.S.M., Lam, K.H., Hung, L.K., Wong, M.M., Yung, L.H., Ho, Y.W., Wong,
W.Y., Hau, D.K.P., Gambari, R., Chui, C.H., 2020. The role of reactive oxygen species in
the biological activity of antimicrobial agents: An updated mini review. Chem. Biol.
Interact. https://doi.org/10.1016/j.cbi.2020.109023

Lau, G.W., Britigan, B.E., Hassett, D.J., 2005. Pseudomonas aeruginosa OxyR is required for
full virulence in rodent and insect models of infection and for resistance to human
neutrophils. Infect. Immun. https://doi.org/10.1128/IAI.73.4.2550-2553.2005

Lemire, J., Alhasawi, A., Appanna, V.P., Tharmalingam, S., Appanna, V.D., 2017. Metabolic
defence against oxidative stress: the road less travelled so far. J. Appl. Microbiol.
https://doi.org/10.1111/jam.13509

Li, B., Yang, Y., Ma, L., Ju, F., Guo, F., Tiedje, J.M., Zhang, T., 2015. Metagenomic and
network analysis reveal wide distribution and co-occurrence of environmental antibiotic
resistance genes. ISME J. https://doi.org/10.1038/ismej.2015.59

Li, D., Liu, C.M., Luo, R., Sadakane, K., Lam, T.W., 2015. MEGAHIT: An ultra-fast single-
node solution for large and complex metagenomics assembly via succinct de Bruijn graph.
Bioinformatics. https://doi.org/10.1093/bioinformatics/btv033

Li, H., Zhou, X., Huang, Y., Liao, B., Cheng, L., Ren, B., 2021. Reactive Oxygen Species in
Pathogen Clearance: The Killing Mechanisms, the Adaption Response, and the Side Effects.
Front. Microbiol. https://doi.org/10.3389/fmicb.2020.622534

Lira, F., Vaz-Moreira, I., Tamames, J., Manaia, C.M., Martínez, J.L., 2020. Metagenomic
analysis of an urban resistome before and after wastewater treatment. Sci. Rep.
https://doi.org/10.1038/s41598-020-65031-y

Liu, S.S., Qu, H.M., Yang, D., Hu, H., Liu, W.L., Qiu, Z.G., Hou, A.M., Guo, J., Li, J.W., Shen,
Z.Q., Jin, M., 2018. Chlorine disinfection increases both intracellular and extracellular
antibiotic resistance genes in a full-scale wastewater treatment plant. Water Res.
https://doi.org/10.1016/j.watres.2018.02.036

Liu, Z., Klümper, U., Liu, Y., Yang, Y., Wei, Q., Lin, J.G., Gu, J.D., Li, M., 2019. Metagenomic
and metatranscriptomic analyses reveal activity and hosts of antibiotic resistance genes in
activated sludge. Environ. Int. https://doi.org/10.1016/j.envint.2019.05.036

Luo, L., Yao, J., Liu, W., Yang, L., Li, H., Liang, M., Ma, H., Liu, Z., Chen, Y., 2021.
Comparison of bacterial communities and antibiotic resistance genes in oxidation ditches

and membrane bioreactors. Sci. Rep. https://doi.org/10.1038/s41598-021-88335-z

Lyons, T.W., Reinhard, C.T., Planavsky, N.J., 2014. The rise of oxygen in Earth's early ocean
and atmosphere. Nature. https://doi.org/10.1038/nature13068

Lyu, F., Han, F., Ge, C., Mao, W., Chen, L., Hu, H., Chen, G., Lang, Q., Fang, C., 2023.
OmicStudio: A composable bioinformatics cloud platform with real-time feedback that can
generate high-quality graphs for publication. iMeta. https://doi.org/10.1002/imt2.85

Ma, X., Zhang, Xiuwen, Xia, J., Sun, H., Zhang, Xuxiang, Ye, L., 2021. Phenolic compounds
promote the horizontal transfer of antibiotic resistance genes in activated sludge. Sci. Total
Environ. https://doi.org/10.1016/j.scitotenv.2021.149549

Majeed, H.J., Riquelme, M. V., Davis, B.C., Gupta, S., Angeles, L., Aga, D.S., Garner, E.,
Pruden, A., Vikesland, P.J., 2021. Evaluation of Metagenomic-Enabled Antibiotic
Resistance Surveillance at a Conventional Wastewater Treatment Plant. Front. Microbiol.
https://doi.org/10.3389/fmicb.2021.657954

Manaia, C.M., Rocha, J., Scaccia, N., Marano, R., Radu, E., Biancullo, F., Cerqueira, F.,
Fortunato, G., Iakovides, I.C., Zammit, I., Kampouris, I., Vaz-Moreira, I., Nunes, O.C.,
2018. Antibiotic resistance in wastewater treatment plants: Tackling the black box. Environ.
Int. https://doi.org/10.1016/j.envint.2018.03.044

Mandsberg, L.F., Ciofu, O., Kirkby, N., Christiansen, L.E., Poulsen, H.E., Høiby, N., 2009.
Antibiotic resistance in Pseudomonas aeruginosa strains with increased mutation frequency
due to inactivation of the DNA oxidative repair system. Antimicrob. Agents Chemother.
https://doi.org/10.1128/AAC.00428-08

Manna, B., Jay, E., Zhang, W., Zhou, X., Lyu, B., Thomas, G.M., Singhal, N., 2024. Short-Term
Warming Induces Cyanobacterial Blooms and Antibiotic Resistance in Freshwater Lake, as
Revealed by Metagenomics Analysis. Water 16, 2655.

McEwen, S.A., Collignon, P.J., 2018. Antimicrobial Resistance: a One Health Perspective.
Microbiol. Spectr. https://doi.org/10.1128/microbiolspec.arba-0009-2017

Mehanni, M.M., Gadow, S.I., Alshammari, F.A., Modafer, Y., Ghanem, K.Z., El-Tahtawi, N.F.,
El-Homosy, R.F., Hesham, A.E.L., 2023. Antibiotic-resistant bacteria in hospital
wastewater treatment plant effluent and the possible consequences of its reuse in
agricultural irrigation. Front. Microbiol. https://doi.org/10.3389/fmicb.2023.1141383

Murray, Christopher JL, et al., 2022. Global burden of bacterial antimicrobial resistance in
2019: a systematic analysis. Lancet. https://doi.org/10.1016/S0140-6736(21)02724-0

Muth, T., Behne, A., Heyer, R., Kohrs, F., Benndorf, D., Hoffmann, M., Lehtevä, M., Reichl, U.,
Martens, L., Rapp, E., 2015. The MetaProteomeAnalyzer: A powerful open-source software
suite for metaproteomics data analysis and interpretation. J. Proteome Res.
https://doi.org/10.1021/pr501246w

Nelson, R.E., Hatfield, K.M., Wolford, H., Samore, M.H., Scott, R.D., Reddy, S.C., Olubajo, B.,
Paul, P., Jernigan, J.A., Baggs, J., 2021. National estimates of healthcare costs associated
with multidrug-resistant bacterial infections among hospitalized patients in the United
States. Clin. Infect. Dis. https://doi.org/10.1093/cid/ciaa1581

Paulus, G.K., Hornstra, L.M., Alygizakis, N., Slobodnik, J., Thomaidis, N., Medema, G., 2019.
The impact of on-site hospital wastewater treatment on the downstream communal
wastewater system in terms of antibiotics and antibiotic resistance genes. Int. J. Hyg.
Environ. Health. https://doi.org/10.1016/j.ijheh.2019.01.004

Petrovich, M.L., Zilberman, A., Kaplan, A., Eliraz, G.R., Wang, Y., Langenfeld, K., Duhaime,
M., Wigginton, K., Poretsky, R., Avisar, D., Wells, G.F., 2020. Microbial and Viral
Communities and Their Antibiotic Resistance Genes Throughout a Hospital Wastewater
Treatment System. Front. Microbiol. https://doi.org/10.3389/fmicb.2020.00153

Puente-Sánchez, F., Garciá-Garciá, N., Tamames, J., 2020. SQMtools: Automated processing
and visual analysis of 'omics data with R and anvi'o. BMC Bioinformatics.
https://doi.org/10.1186/s12859-020-03703-2

Qi, W., Jonker, M.J., Teichmann, L., Wortel, M., ter Kuile, B.H., 2023. The influence of oxygen
and oxidative stress on de novo acquisition of antibiotic resistance in E. coli and
Lactobacillus lactis. BMC Microbiol. 23, 1–14. https://doi.org/10.1186/s12866-023-03031-
4

R Core Team, 2022. R Core Team 2021 R: A language and environment for statistical
computing. R foundation for statistical computing. https://www.R-project.org/. R Found.
Stat. Comput.

Robinson Mark, D., McCarthy Davis, J., Smyth Gordon, K., 2010. edgeR: a Bioconductor
package for differential expression analysis of digital gene expression data. Bioinformatics.

Rowe, W.P.M., Baker-Austin, C., Verner-Jeffreys, D.W., Ryan, J.J., Micallef, C., Maskell, D.J.,
Pearce, G.P., 2017. Overexpression of antibiotic resistance genes in hospital effluents over
time. J. Antimicrob. Chemother. https://doi.org/10.1093/jac/dkx017

Schneider, D., Aßmann, N., Wicke, D., Poehlein, A., Daniel, R., 2020. Metagenomes of
Wastewater at Different Treatment Stages in Central Germany. Microbiol. Resour.
Announc. https://doi.org/10.1128/mra.00201-20

Schneider, D., Zühlke, D., Poehlein, A., Riedel, K., Daniel, R., 2021. Metagenome-Assembled
Genome Sequences from Different Wastewater Treatment Stages in Germany. Microbiol.
Resour. Announc. https://doi.org/10.1128/mra.00504-21

Seemann, T., 2014. Prokka: Rapid prokaryotic genome annotation. Bioinformatics.
https://doi.org/10.1093/bioinformatics/btu153

Seo, S.W., Kim, D., Szubin, R., Palsson, B.O., 2015. Genome-wide Reconstruction of OxyR and
SoxRS Transcriptional Regulatory Networks under Oxidative Stress in Escherichia coli K-

12 MG1655. Cell Rep. https://doi.org/10.1016/j.celrep.2015.07.043

Sulfikar, Honda, R., Noguchi, M., Yamamoto-Ikemoto, R., Watanabe, T., 2018. Effect of
sedimentation and aeration on antibiotic resistance induction in the activated sludge process.
J. Water Environ. Technol. https://doi.org/10.2965/jwet.17-046

Tamames, J., Puente-Sánchez, F., 2019. SqueezeMeta, a highly portable, fully automatic
metagenomic analysis pipeline. Front. Microbiol. https://doi.org/10.3389/fmicb.2018.03349

Tan, Q., Chen, J., Chu, Y., Liu, W., Yang, L., Ma, L., Zhang, Y., Qiu, D., Wu, Z., He, F., 2021.
Triclosan weakens the nitrification process of activated sludge and increases the risk of the
spread of antibiotic resistance genes. J. Hazard. Mater.
https://doi.org/10.1016/j.jhazmat.2021.126085

Uluseker, C., Kaster, K.M., Thorsen, K., Basiry, D., Shobana, S., Jain, M., Kumar, G.,
Kommedal, R., Pala-Ozkok, I., 2021. A Review on Occurrence and Spread of Antibiotic
Resistance in Wastewaters and in Wastewater Treatment Plants: Mechanisms and
Perspectives. Front. Microbiol. https://doi.org/10.3389/fmicb.2021.717809

Van Acker, H., Coenye, T., 2017. The Role of Reactive Oxygen Species in Antibiotic-Mediated
Killing of Bacteria. Trends Microbiol. https://doi.org/10.1016/j.tim.2016.12.008

Wang, P., Zhang, H., Liu, Y., Lv, R., Liu, X., Song, X., Wang, J., Jiang, L., 2020. SoxS is a
positive regulator of key pathogenesis genes and promotes intracellular replication and
virulence of Salmonella Typhimurium. Microb. Pathog.
https://doi.org/10.1016/j.micpath.2019.103925

Wang, Y., Han, Y., Li, L., Liu, J., Yan, X., 2022. Distribution, sources, and potential risks of
antibiotic resistance genes in wastewater treatment plant: A review. Environ. Pollut.
https://doi.org/10.1016/j.envpol.2022.119870

Wang, Y., Lu, J., Engelstädter, J., Zhang, S., Ding, P., Mao, L., Yuan, Z., Bond, P.L., Guo, J.,
2020. Non-antibiotic pharmaceuticals enhance the transmission of exogenous antibiotic
resistance genes through bacterial transformation. ISME J. https://doi.org/10.1038/s41396-
020-0679-2

Wei, Q., Le Minh, P.N., Dötsch, A., Hildebrand, F., Panmanee, W., Elfarash, A., Schulz, S.,
Plaisance, S., Charlier, D., Hassett, D., Häussler, S., Cornelis, P., 2012. Global regulation of
gene expression by OxyR in an important human opportunistic pathogen. Nucleic Acids
Res. https://doi.org/10.1093/nar/gks017

Wei, Z., Feng, K., Wang, Z., Zhang, Y., Yang, M., Zhu, Y.G., Virta, M.P.J., Deng, Y., 2021.
High-Throughput Single-Cell Technology Reveals the Contribution of Horizontal Gene
Transfer to Typical Antibiotic Resistance Gene Dissemination in Wastewater Treatment
Plants. Environ. Sci. Technol. https://doi.org/10.1021/acs.est.1c01250

Wickham, H., 2011. ggplot2. Wiley Interdiscip. Rev. Comput. Stat.
https://doi.org/10.1002/wics.147

Yang, J.H., Wright, S.N., Hamblin, M., McCloskey, D., Alcantar, M.A., Schrübbers, L.,
Lopatkin, A.J., Satish, S., Nili, A., Palsson, B.O., Walker, G.C., Collins, J.J., 2019. A
White-Box Machine Learning Approach for Revealing Antibiotic Mechanisms of Action.
Cell. https://doi.org/10.1016/j.cell.2019.04.016