Cameroonian blackflies (Diptera: Simuliidae) harbour a plethora of (RNA) viruses

Lander De Coninck[1], Amber Hadermann[2], Ludovica Ingletto[1], Robert Colebunders[2], Kongnyu Gamnsi Njamnshi[3], Alfred Kongnyu Njamnshi[3], John L Mokili[4], Joseph Nelson Siewe Fodjo[2#], Jelle Matthijnssens[1#]

1. KU Leuven, Department of Microbiology, Immunology and Transplantation, Rega Institute, Laboratory of Viral Metagenomics, Leuven, Belgium
2. Global Health Institute, University of Antwerp, Antwerp, Belgium.
3. Brain Research Africa Initiative (BRAIN), Yaoundé, Cameroon & Geneva, Switzerland; Neuroscience Lab, Faculty of Medicine and Biomedical Sciences, The University of Yaoundé I, Yaoundé, Cameroon.
4. Viral Information Institute, Department of Biology, San Diego State University San Diego, CA, USA.

[#]Co-last authors

Abstract

Strong epidemiological evidence suggests that onchocerciasis may be associated with epilepsy – hence the name onchocerciasis-associated epilepsy (OAE). However, the pathogenesis of OAE still needs to be elucidated, as recent studies failed to detect *Onchocerca volvulus* in the central nervous system of persons with OAE. Therefore, it was suggested that a potentially neurotropic virus transmitted by blackflies could play a role in triggering OAE. To investigate this hypothesis, adult blackflies were collected in an onchocerciasis-endemic area with a high OAE prevalence in the Ntui Health District, Cameroon. A whole-genome shotgun sequencing approach was used to detect viral sequences in fifty-five pools of ten blackflies. A very high abundance of viral reads was detected across multiple (novel) viral families, including viral families associated with human disease. Although, no genomes closely related to known neurotropic viruses were found in the blackfly virome, the plethora of novel viruses representing novel species, genera and even families, warrant further exploration for their potential to infect vertebrates. These results could serve as a first step for studying the viruses associated with the hematophagous blackfly, which also could be present in their nematode host *O. volvulus*.

31 Exploring the diversity of viruses in blackflies should be included in the active surveillance of

32 zoonotic diseases.

Introduction

Human onchocerciasis, commonly known as river blindness, is a parasitic disease caused by

the filarial worm *Onchocerca volvulus* (Nematoda: Secernentea; Spirurida; Filariidae;

Onchocerca). Transmission of the parasite occurs through repeated bites of female

blackflies belonging to the *Simulium* species (Brattig, Cheke and Garms 2021).

Onchocerciasis is classified as one of the neglected tropical diseases recognized by the

World Health Organization (WHO).

Onchocerciasis-endemic regions with high ongoing or past *O. volvulus* transmission are

known to have high epilepsy prevalence (Pion and Boussinesq 2012; Levick *et al.* 2017;

Colebunders *et al.* 2018b; Siewe Fodjo *et al.* 2018; Boullé *et al.* 2019; Mukendi *et al.* 2019;

Raimon *et al.* 2021). While onchocerciasis is commonly thought to affect only the skin and

eyes, emerging epidemiological evidence suggests a potential direct or indirect association

between *O. volvulus* infection and epilepsy, giving rise to the term "Onchocerciasis-

Associated Epilepsy" (OAE), also known as "River Epilepsy" (Chesnais *et al.* 2018, 2020;

Colebunders *et al.* 2019, 2021; Hadermann *et al.* 2023). OAE presents with a wide spectrum

of epileptic seizures, including generalized tonic-clonic seizures, myoclonic seizures, absence

seizures and head nodding seizures (Colebunders *et al.* 2018a). Nodding and Nakalanga

syndrome are considered to be phenotypic presentations of OAE (Colebunders et al. 2018a;

Van Cutsem et al. 2023). Nakalanga syndrome is characterized by severe growth

retardation, delayed sexual development, mental impairment, facial deformation,

kyphoscoliosis, and epileptic seizures (Föger *et al.* 2017; Siewe Fodjo *et al.* 2019).

So far, the pathogenesis of OAE is not yet established. Prior to the introduction of

ivermectin mass drug distribution, *O. volvulus* microfilariae (mf) had been detected in the

cerebrospinal fluid (CSF) of individuals with onchocerciasis (Duke, Vincelette and Moore

1976). However, in more recent studies, neither *O. volvulus* mf nor DNA could be detected

in CSF (Winkler et al. 2013; Hotterbeekx et al. 2020) nor in post-mortem brain tissue of

persons with OAE (Hotterbeekx *et al.* 2019). Therefore, it was suggested that a potentially

neurotropic virus transmitted by blackflies, or an endosymbiont of the *O. volvulus* parasite could play a role in the pathogenesis of OAE (Colebunders et al. 2014; Hadermann et al. 2023).

Very little is known about the virome of blackflies. Kraberger *et al.* described the ssDNA virome of blackflies in New Zealand and reported high numbers of *Genomoviridae*, *Circoviridae* and *Microviridae* (Kraberger *et al.* 2019). Moreover, invertebrate iridoviruses (family *Iridoviridae*), have been identified in blackfly larvae across the globe (Piégu *et al.* 2013, 2014a, 2014b). In addition, the most studied arbovirus transmitted by blackflies is vesicular stomatitis virus (family *Rhabdoviridae;* genus *Vesiculovirus*), that typically infects cattle, horses, and swine (Howerth, Mead and Stallknecht 2002). Nonetheless, there have been no reports of blackflies transmitting arboviruses to humans to date. However, considering the 1700 *Simulium* species described worldwide, it is plausible that blackflies may transmit unidentified viruses from human and zoonotic origins. The aim of this study was to identify potential neurotropic viruses transmitted by blackflies, which could play a role in the pathogenesis of OAE.

Materials and Methods

Sample collection

Adult blackflies investigated were collected in the village of Nachtigal (coordinates of the breeding site: N 4°21.146, E 11°37.953; Garmin GPSMAP78). The gathering took place from 7:00 AM to 6:00 PM over three consecutive days in July 2021. Nachtigal is a rural community of the Ntui Health District, an onchocerciasis-endemic area with a high prevalence of OAE (Siewe Fodjo *et al.* 2022), in the forest-savannah transition zone in the Centre Region of Cameroon. This community lives close to the Nachtigal rapids on the Sanaga River, which constitutes one of the main sources of blackfly vectors in Cameroon (Siewe Fodjo *et al.* 2022). The human landing catch method was used to collect the blackflies and mouth aspirators were used to aspirate blackflies into an empty container every hour (Hendy *et al.* 2021). At the end of each catching day, blackflies were stored dry at -20°C in the BRAIN laboratory, Yaoundé, Cameroon until shipping on dry ice a few weeks later. The frozen blackfly samples were then sent to the Laboratory of Viral Metagenomics in Leuven,

89 Belgium, where they were stored at −80°C until the nucleic acids were extracted for
90 sequencing. A total of fifty-five pools of ten blackflies were analysed (see <mark>Supplementary</mark>
91 <mark>Table S1</mark> for details).

Sample processing

The NetoVIR protocol (Conceição-Neto *et al.* 2015) was adapted for the homogenization of
the blackfly pools. Homogenization was carried out in 2mL tubes with 2.8mm zirconium
oxide beads at 5000 rpm for 15 seconds. Negative controls were added in the beginning and
taken through the whole process (including sequencing) to make sure no contamination
happened during the processing of the samples. Paired-end sequencing (2x150bp) was
performed on the Novaseq 6000 SP platform at the Nucleomics Core sequencing facility (VIB
Leuven).

Raw Reads Processing and Virome Analysis

To process the raw paired-end Illumina reads, an in-house bioinformatics pipeline (ViPER
v1.1) (De Coninck 2021) was used with the 'triple assembly' option enabled. Briefly,
adapters and low-quality reads were trimmed with Trimmomatic (Bolger, Lohse and Usadel
2014), before a *de novo* assembly with metaSPAdes (Nurk *et al.* 2017). Resulting fasta files
with scaffolds were subsequently concatenated. To remove redundancy in the data, these
scaffolds were clustered on 95% nucleotide identity over 85% coverage of the shortest
sequence with the clustering algorithm available from CheckV (Nayfach *et al.* 2020). A
length filter of 1000 nucleotides was applied on the sequences before viruses were
identified in the data using a combined approach. This approach consisted of: 1) genomad
(Camargo *et al.* 2023); 2) DIAMOND blastx (Buchfink, Xie and Huson 2014) with NCBI's nr
database (accessed 23-03-2023); and 3) a Hidden Markov Model (HMM) search of the
scaffolds' open reading frames (ORFs), predicted by prodigal (Hyatt *et al.* 2010), against the
NeoRdRP HMM dataset (v1.0) (Sakaguchi *et al.* 2022) with HH-suite3 (Steinegger *et al.*
2019). Taxonomy assignment for blastx-identified sequences was decided based on a lowest
common ancestor approach from the best twenty-five hits. When possible, the genomad
taxonomy output was prioritized for a sequence over the blastx taxonomic assignment.
Subsequently, for quality control purposes, CheckV (Nayfach *et al.* 2020) was employed to

assess the genome completeness of the identified viral sequences. Sequences not belonging

to the *Riboviria* and less than 50% complete as predicted by CheckV were removed from

further analysis. *Riboviria* sequences predicted to be less than 50% complete were kept

because CheckV does not handle (segmented) RNA viruses well, which often results in

wrong completeness estimations for RNA viruses. Finally, using the sequencing data of the

negative controls, sequences that were predicted by the prevalence method of decontam

(Davis et al. 2018) to be a contaminant were removed from our dataset.

Phylogenetic Analysis of the *Orthornavirae*

In ORFs identified to code for an RNA-dependent RNA polymerase (RdRP) (*vide supra*), we

searched for the palmcore sequence using Palmscan2

(https://github.com/rcedgar/palmscan). This palmcore sequence contains the palmprint,

three conserved catalytic domains across all known RdRP sequences (domain A, B and C)

with two variable regions in between, flanked by ~150 amino acids on both sides (Babaian

and Edgar 2022). Furthermore, we added 76,164 palmcore sequences from the RCR90 set of

Neri *et al.* (Neri *et al.* 2022). This set contained previously published *Orthornavirae*

sequences which were the basis of ICTV's *Orthornavirae* classification (Wolf *et al.* 2018,

2020), as well as new RNA viruses and ten reverse transcriptase (RT) sequences to be used

as outgroup for the rooting of the phylogenetic tree. Next, palmcore sequences were

aligned a first time with Muscle5 (Edgar 2022) with default settings. After alignment,

columns with >90% gaps were removed with trimal (Capella-Gutiérrez, Silla-Martínez and

Gabaldón 2009), and gaps were deleted from the sequences to prepare the data for a

second alignment. From these trimmed sequences a diversified ensemble was created with

Muscle5 (Edgar 2022). This diversified ensemble constitutes 100 multiple sequence

alignments (MSAs) of the trimmed sequences with perturbed alignment parameters and

permuted guide trees (see Muscle5 documentation for more information). From these 100

MSAs, the alignment with the highest column confidence, calculated with muscle maxcc,

was picked to generate the final phylogenetic tree. Resampling of the diversified ensemble

to generate support values for the phylogenetic tree was obtained with muscle resample -

conf 0 -gapfract 0.5. Maximum likelihood (ML) phylogenetic trees from the resampled MSAs

were inferred with Fasttree (Price, Dehal and Arkin 2010) (-lg -gamma -fastest -nosupport).

These ML trees served as support for the tree that was generated from the maxcc MSA (also
generated with Fasttree and unaltered settings) which was calculated with newick conftree.

Viral host inference

To infer if the *Flaviviridae* and *Rhabdoviridae* viruses in our data could have an invertebrate
or vertebrate host, we calculated the relative dinucleotide abundance (RDA) of our
sequences and sequences with a known host (see Supplementary Table S2 for accession
numbers) with DinuQ (Lytras and Hughes 2020). Next, we performed a principal component
analysis in R on all dinucleotide RDA values and plotted the first against the second principal
component for all viruses.

Ethical approval

Approval was obtained from the ethics committee of the Cameroon Baptist Convention
Health Services (reference number IRB2021-03). We also obtained a research permit from
the   Ministry   of   Scientific   Research   and   Innovation   (Ref:
000144/MINRESI/B00/C00/C10/C13), as well as an ABS-Nagoya Protocol Prior Informed
Consent   (Ref: Decision   No.   00016/D/MINEPDED/CNA)   and   ABS   Permit   No.
00013/MINEPDED/CAN/NP-ABS/ABS-FP) from the Ministry of Environment, Protection of
Nature and Sustainable Development.

Results

Identification of more than thousand viral contigs

For the fifty-five blackfly pools sequenced, between 6,113,870 and 24,002,232 reads were
obtained per pool. In total, around 720 million raw reads were generated from all fifty-five
pools. After trimming, each pool retained between 4,544,534 and 20,790,370 reads,
resulting in a 17.5% loss of the initial raw reads (see Supplementary Table S1). Subsequently,
de novo assemblies were generated, resulting in a total of 59,645 scaffolds larger than 1000
bp from the 55 pools. The clustering of the assembled scaffolds from all samples based on
95% nucleotide identity over 85% coverage resulted in 45,308 non-redundant scaffolds (see
Supplementary Table S1). After removing contamination predicted with decontam, a total of

1,678 scaffolds were identified as viral with a combination of genomad, DIAMOND blastx
and an HMM search against the NeoRdRP dataset (see <mark>Figure 1A</mark>). Of these, 56 were
considered to be 100% complete, 128 were of high quality (>90% complete) and 190 of
medium quality (>50% complete), the remaining viral sequences were either of low quality
or their completeness could not be determined by CheckV (see <mark>Figure 1B</mark>). However, the
low-quality and not-determined groups consisted only of sequences with *Riboviria* or
unclassified taxonomic realm assignments. This suggests that these sequences are
significantly divergent and might thus be of a higher quality than what is estimated by
CheckV (also see Materials). Overall, we concluded that, due to the high convergence of our
three identification methods and the completeness estimation of CheckV, we have a robust
set of identified viruses.

<mark>Figure 1</mark>: A robust set of viruses was identified based on three different identification
methods and a viral genome completeness estimation. A) Venn diagram of the identified
viral sequences by the various methods. B) Pie chart showing the completeness estimation
of CheckV on the identified viral scaffolds.

The blackfly virome is dominated by RNA viruses

When examining the taxonomic distribution of our viral genome sequences, the majority (n=1207) were classified within the *Riboviria* realm, 301 could not be assigned into a specific realm and a small number belonged to the *Monodnavaria* and the *Duplodnaviria* (see Figure 2A). At the read level, a distinct picture arises with approximately two thirds being identified as belonging to the *Duplodnaviria* realm. This is mostly because the average viral genome length of *Duplodnaviria* is much higher than the average genome length for the *Riboviria*, which in our data recruit about one third of the viral reads (see Figure 2A). The 301 unclassified sequences recruit only a small fraction (~1 million) of the reads. As the diversity seemed the greatest in the *Riboviria* realm, which also contains the vast majority of arboviruses, we further investigated the taxonomic distribution of their scaffolds as well as their reads. Around half of the viral genome sequences belonged to the *Duplornaviricota* phylum (see Figure 2B), while this phylum constituted only one third of the *Riboviria* reads (see Figure 2B). Surprisingly, 81 further unclassified *Riboviria* sequences were responsible for another third of the viral reads in our blackfly samples. These results show that the blackflies contained a high number of completely novel RNA viruses that were present in high abundance. Moreover, we constructed rarefaction curves to assess the depth of sampling required to encompass the full spectrum of viral diversity (Figure 2C). Notably, the curves (total and *Riboviria*) plateaued after approximately fifty pools, suggesting saturation and the absence of novel viral taxa beyond this point. These findings indicate that our sampling efforts comprehensively cover the viral diversity within the blackflies from the onchocerciasis-endemic Nachtigal village (Cameroon).

Figure 2: **Blackfly viromes are dominated by RNA viruses.** A) Barplots showing the relative abundance of all identified viruses at contig and read level. Labels indicate the absolute numbers. B) Barplots showing the relative abundance of all identified *Riboviria* phyla at contig and read level. Labels indicate the absolute numbers. C) Rarefaction curves for each virus kingdom and the total. Light shaded areas indicate the minimum and maximum observed species across all iterations, the dark shaded areas indicate the 95% confidence around the mean.

To visualize the virome similarity of the fifty-five pools, a heatmap with the read count for
each viral family on a log$_2$ scale was constructed (see Figure 3). In total, forty eukaryotic viral
families were identified across all samples. In addition, eleven different taxonomic instances
of prokaryotic viruses were identified on the family level. Most diversity in the blackflies on
the family level is present in the *Riboviria* realm with thirty-five families of eukaryotic RNA
viruses detected.

The viral families *Caliciviridae, Flaviviridae, Hepeviridae, Orthomyxoviridae, Parvoviridae,*
Phenuiviridae, Picornaviridae, Spinareoviridae, Rhabdoviridae, and *Togaviridae* include
human pathogenic viruses, some of which may also be arboviruses. Notably, known viruses
causing neurological complications belong to the families of *Rhabdoviridae, Flaviviridae,*
Picornaviridae as well as *Togaviridae*.

Plant and fungal viruses were the most prevalent viruses, with members of *Ghabrivirales*
(i.e., *Chrysoviridae, Totiviridae*) and *Partitiviridae* being the most abundant. A virus of the
family *Flaviviridae* was present in most blackfly pools and in a relatively high abundance.
Furthermore, other eukaryotic viral families (main hosts according to ICTV are given in
brackets) that were prevalent, were *Curvulaviridae* (fungi), *Dicistroviridae* (invertebrates),
Iflaviridae (insects), *Genomoviridae* (human, mammals, birds, fungi), *Mymonaviridae* (fungi),
Orthomyxoviridae (human, birds, pigs), *Parvoviridae* (vertebrates, insects), *Polycipiviridae*
(insects), *Rhabdoviridae* (vertebrates, invertebrates, plants), *Totiviridae* (fungi) and
Virgaviridae (plants). Other viral families of potential clinical interest *Caliciviridae,*
Hepeviridae, Phenuiviridae, Picornaviridae and *Spinareoviridae* were present in lower
abundance and prevalence.

Detailed phylogenetic analysis for all above mentioned viral families has been conducted
and can be found, together with extra information on closely related viruses, in the
Supplemental Information.

Figure 3: Blackflies harbour a large number of viral families, including families of potential clinical interest. Heatmap of all viral families per blackfly pool. The heatmap shows the read count on a log2 scale for eukaryotic virus on family level from all the 55 adult blackfly pools. The number of identified scaffolds are indicated after each viral family name in the heatmap. Hierarchical clustering of the columns is based on the Bray-Curtis distance calculated from the read counts.

Blackfly virome members are found across all *Orthornavirae* phyla, including a potential new phylum of dsRNA viruses

To further explore the *Orthornavirae* diversity in our samples, we generated a ML phylogenetic tree based on the RdRP palmcore of all RdRPs identified in the blackflies, supplemented with previously identified RNA virus sequences (Wolf *et al.* 2018, 2020; Neri *et al.* 2022). In total we could identify 76,097 palmcore sequences with a complete palmprint, of this set 466 were derived from the blackfly data. The cladogram in Figure 4 shows that blackflies host RNA viruses across all *Orthornavirae* phyla (indicated with red dots). Most are situated within the *Negarnaviricota*, the *Duplornaviricota* and the dsRNA *Pisuviricota*. These last two phyla mostly harbour viruses infecting fungi. Interestingly, in our phylogenetic tree the dsRNA *Pisuviricota* were placed in a separate clade from the ssRNA(+) *Pisuviricota*, suggesting that, in contrast to what was previously observed (Wolf *et al.* 2018, 2020; Neri *et al.* 2022), the dsRNA *Pisuviricota* might represent a distinct phylum.

In our search for potential neurotropic viruses, we identified two novel viruses belonging to the *Rhabdoviridae* which are part of the ssRNA(-) *Negarnaviricota*. The most elusive neurotropic member of the *Rhabdoviridae* is rabies virus. Furthermore, also the *Flaviviridae* within the *Kitrinoviricota* harbor known neurotropic viruses (eg. West Nile virus) (Johnson 2016). Within this family we found one novel virus that was consistently present across the blackfly pools (see Figure 3).

Figure 4: Cameroonian blackflies contain viruses across all phyla of the *Orthornavirae.*
Circularized cladogram of *Orthornavirae* viruses based on the phylogeny of the RdRP palmcore domain. The ML tree is rooted on RT sequences as outgroup. Blackfly sequences are indicated with red dots on the leaves of the tree. The ICTV phylum classification of viruses is shown on the outside ring, sequences established in putative new orders/families by Neri *et al.* (2022) are coloured in light grey, while the blackfly sequences are coloured black. Viruses with a permuted catalytic RdRP domain (motif CAB) are reported with an orange star.

Novel blackfly rhabdo- and flaviviruses are most likely insect-specific

To investigate whether the *Rhabdoviridae* and *Flaviviridae* member viruses we found in the blackflies could also infect vertebrates, we compared the dinucleotide composition within their genomes with the genomes of viruses with a known host. After coloring the viruses by host, this revealed a clear clustering in vertebrate-infecting vs. insect-specific viruses for the *Flaviviridae* (see Figure 5A), meaning that this family of viruses adapted itself to avoid the vertebrate immune system by using similar dinucleotide abundances in their genome (Lytras and Hughes 2020; Fros *et al.* 2021). For the *Rhabdoviridae* this clustering is less clear, but there seems to be nevertheless a small difference. The flavivirus we found in the blackflies shows a dinucleotide profile suggestive of being an insect-specific virus and also the rhabdoviruses we found in the blackflies seem to be infecting invertebrates (see Figure 5B). Additionally, we added two other virus species recently identified in other studies to the *Rhabdoviridae* dinucleotide analysis: *Onchocerca volvulus* RNA virus 1 (OVRV1) is a rhabdovirus that was found in *Onchocerca volvulus* and elicits an immune response in the host (human as well as cattle and hamsters) (Quek *et al.* 2023), while Mundri virus was detected in a patient with nodding syndrome (Edridge *et al.* 2022). Interestingly, both multiple OVRV1 variants as well as Mundri virus seem to cluster with vertebrate infecting rhabdoviruses, suggesting that these viruses may also be capable of infecting vertebrates.

Figure 5: **Identified blackfly flavi- and rhabdoviruses are most likely insect-specific.** A) PCA plot based on the RDA of selected *Flaviviridae* sequences. The red dot shows the newly identified flavivirus in blackflies. Labels show a selection of neurotropic flaviviruses (TBEV=tick-borne encephalitis virus, JEV=Japanese encephalitis virus, WNV=West Nile virus, ZIKV=Zika virus). B) PCA plot based on the RDA of selected *Rhabdoviridae* sequences. The blue dots show the newly identified blackfly rhabdoviruses, the orange dots show different variants of OVRV1, the brown dot shows Mundri virus (MUNV). Other labelled dots show a selection of neurotropic rhabdoviruses (CHPV=Chandipura virus, LBV=Lagos bat virus, RABV=rabies virus).

Discussion

To our knowledge, this study represents the first comprehensive investigation of DNA and RNA viruses in blackflies utilizing a whole-genome sequencing approach. A very high number of virus genomes, with a dominance of RNA viruses, were identified in blackfly samples. Fifteen good quality (>50% complete) genomes of eukaryotic DNA viruses (eg. *Parvoviridae* and *Genomoviridae*), and 641 RNA viruses (based on the RdRP, of which 466 with a complete palmprint) were detected. A significant proportion of the novel viruses belonged

to viral families associated with human disease, eg. *Caliciviridae, Flaviviridae, Hepeviridae,*
Orthomyxoviridae, Parvoviridae, Phenuiviridae, Picornaviridae, Spinareoviridae,
Rhabdoviridae, and *Togaviridae.* However, no viral genomes closely related to known
neurotropic viruses were found in the blackfly virome (Supplementary files). Obviously, this
does not mean that none of the newly identified viruses could be arboviruses with a
neurotropic potential. Additionally, the real host origin for many of the newly discovered
viruses remains unknown and further investigations are necessary to determine whether
these viruses have the potential to replicate in vertebrate and/or invertebrate hosts, or
even if they have a prokaryotic host as could be the case for the *Partitiviridae* and
Picobirnaviridae (Neri *et al.* 2022; Sadiq, Holmes and Mahar 2024).

The only *Flaviviridae* virus in our analysis was observed at a relatively high abundance in
most blackfly samples (see Figure 3). Previous phylogenetic analyses and experimental
studies (Kuno and Chang 2005; Cook *et al.* 2013) have shown that the phylogenetic tree of
the *Flaviviridae* separates into two large clusters. The first cluster contains ISVs which
appear to be host-restricted and only replicate in invertebrate cells and primarily infect
mosquitoes (Hoshino *et al.* 2009; Guzman *et al.* 2018; Miranda *et al.* 2019; Baidaliuk *et al.*
2020). The second cluster comprises human pathogenic arboviruses. These previous studies
also indicate that arthropod-borne viruses of vertebrates may have originated from ISVs
that evolved from being limited to insects to acquiring the ability to infect vertebrate hosts
(Bolling *et al.* 2015). The dinucleotide analysis of the *Flaviviridae* indicated that the flavivirus
in our blackfly samples is most likely an insect-specific virus. Our phylogenetic analysis
revealed that the new blackfly flavivirus forms a deep evolutionary branch within the ISV
subcluster, distantly related to Hangzhou flavivirus 3, detected in a fly (see Supplemental
Information). These two viruses may form a distinct cluster of ISVs with the fly as the natural
host. Given the high abundance of this flavivirus in blackfly pools and the presence of ISVs
among them, it is possible that they may become new emerging pathogens.

As blackflies can act as intermediate hosts of pathogens affecting humans and animal health
due to their hematophagic habit, several studies (Mead *et al.* 2004; Smith *et al.* 2009;
Mesquita *et al.* 2017; Young *et al.* 2021) have focused on the transmission of vesicular
stomatitis virus (VSV), an arbovirus belonging to the family *Rhabdoviridae,* genus

Vesiculovirus, which is known to be transmitted by blackflies. This virus primarily infects

livestock, representing an agriculturally relevant pathogen, but zoonotic events have also

been reported, highlighting its potential impact on human health (Lichty *et al.* 2004). VSV

infections are generally asymptomatic in humans, however, mild flu-like symptoms have

been reported in some individuals and a single case of encephalitis in a 3-year-old boy,

which was potentially associated with VSV infection, has been reported (Quiroz *et al.* 1988).

In 2022, Edridge and colleagues discovered a divergent rhabdovirus in the bloodstream of a

15-year-old girl with Nodding syndrome from Mundri West Country in South Sudan (Edridge

et al. 2022). They named this virus "Mundri virus" (MUNV) and classified it as a novel virus.

Subsequent phylogenetic analysis revealed that MUNV belongs to the monophyletic clade of

tiroviruses within the family *Rhabdoviridae*. The *Tibrovirus* genus is part of a larger

phylogenetic group of arthropod-borne rhabdoviruses. Several *tiroviruses* have been

isolated from biting midges (*Culicoides* spp.) and cattle, and some instances of human

detection have been reported (Cybinski *et al.* 1980; Cybinski and Gard 1986; Gibbs *et al.*

1989; Grard *et al.* 2012; Stremlau *et al.* 2015). Human tibrovirus infections may accidentally

occur via midge bites (Kuhn *et al.* 2020). Consistent with the findings of Edridge *et al.*, our

phylogenetic analysis of *Rhabdoviridae* (see Supplemental Information) showed that MUNV

is closely related to the Tibrovirus genus, supported by a high bootstrap value. However, the

two sequences detected in the blackfly virome were not closely related to the *Tibrovirus* nor

Vesiculovirus genera. In addition, another novel rhabdovirus, named OVRV1, was discovered

by Quek *et al.* in public transcriptomes and viromes of *Onchocerca volvulus* nematodes

(Quek *et al.* 2023). Because OVRV1 was shown to elicit an immune response in the

vertebrate host of the nematode, it could potentially play a role in the pathogenesis of OAE

(Quek *et al.* 2023). We were, however, not able to find this virus back in the virome of the

blackflies. Our dinucleotide analysis for the *Rhabdoviridae*, suggested that the viruses we

found in the blackflies are invertebrate-specific viruses, in contrast to the previously

identified rhabdoviruses MUNV (Edridge *et al.* 2022) and OVRV1 (Quek *et al.* 2023).

However, a recent association between MUNV infection and nodding syndrome was

investigated in a case-control study in South Sudan. MUNV RNA was only detected in the

plasma of one of the 72 persons with Nodding syndrome and in none of the controls. MUNV

anti-nucleocapsid IgG antibodies were observed in 28% of Nodding syndrome cases, 22% of

household controls, and 16% of community controls without significant differences between

cases and both control groups. Therefore, it was concluded that MUNV may not be causally

associated with Nodding syndrome (Edridge *et al.* 2022). Similarly, it remains to be shown

whether OVRV1 may be associated with OAE.

Our study is the first to extensively study the DNA and RNA virome of blackflies from an

onchocerciasis-endemic area in Africa. Our findings reveal a relative high abundance of the

family *Genomoviridae* in the majority of blackfly samples, aligning with the results of

Kraberger *et al.*, which demonstrated that the majority of novel ssDNA viral genomes

detected in blackflies in New Zealand belonged to the *Genomoviridae* family (Kraberger *et*

al. 2019). However, in contrast with the blackflies in New Zealand (Kraberger *et al.* 2019),

our study identified *Parvoviridae*, in the blackfly virome. Although this viral family exhibited

a high abundance, it was only observed in a limited number of blackfly samples.

Phylogenetic analysis revealed that one of the *Parvoviridae* genomes detected in this study

clustered within the recently defined *Hamaparvovirinae* subfamily, containing many viruses

identified in vertebrate hosts, making it an interesting candidate to further study for its

ability to infect vertebrate cells.

Most of the detected viruses exhibited low sequence similarities with their closest relatives.

Consequently, based on the demarcation criteria established for viral classification, it is

reasonable to propose that the majority of sequences identified in our study represent new

virus species, genera, and/or families, which warrant further studies. In our project, plant

and fungal infecting viruses (mainly members of the *Chrysoviridae* and *Partitiviridae*

families) were the most abundant, possibly from the nectar diet of the blackflies. Some

members of viral families of clinical interest were detected, albeit at lower abundances and

no viral genomes closely related to neurotropic viruses were identified.

Our study has some limitations. First of all, it reports the composition of the blackfly virome,

including both DNA and RNA viruses but without metadata that would have allowed further

associations to be made with *O. volvulus* and potential environmental factors. Further

research should compare the virome of blackflies with and without *O. volvulus* infection,

and the sex and species of the blackflies should be taken into consideration. It is important
to note that only a few blackfly species consume a blood meal (Kraberger *et al.* 2019).
Additionally, as less than 1% of blackflies in Cameroon are currently infected with *O.*
volvulus (Shintouo et al. 2020), it is recommended to screen blackflies for *O. volvulus*
infection before sequencing data analyses, as previously suggested (Ekangouo *et al.* 2022).

Conclusion

Overall, we provide a glimpse into the viral diversity associated with blackfly vectors in
Cameroon. Although no genomes associated with neurotropic viruses were found, a
plethora of new virus species, genera, and families were identified. The presence of
dominant and under-represented viruses warrants future research to determine the role of
these viruses in their hosts and ecosystems. Exploring the diversity of viruses in blackflies
should be included in the active surveillance of zoonotic diseases. Our findings constitute a
starting point for investigating the viruses associated with the hematophagous blackfly and
potentially in their nematode host *O. volvulus*.

Funding

421 LDC, RC and JNSF were supported by the Research Foundation – Flanders (FWO, grant
422 numbers 11L1323N, G0A0522N and 1296723N, respectively). In addition, RC received
423 funding from the European Research Council (grant 671055) and blackflies were captured in
424 Nachtigal as part of an EDCTP2 project supported by the European Union (grant number
425 TMA2020CDF-3152-SCONE).

Data availability

427 Sequencing data are available in the SRA database under BioProject accession number
428 PRJNA1088476. Viral genomes are submitted to GenBank under accession numbers XXX-
429 XXX. The code used for this study is available in a GitHub repository:
430 https://github.com/Matthijnssenslab/2024_XXX_Blackfly_Virome.

Acknowledgements

We thank Jill Swinnen (KU Leuven) and Jara Wagemans (KU Leuven) for their assistance during the processing of the blackfly pools for Illumina sequencing.

References

Babaian A, Edgar R. Ribovirus classification by a polymerase barcode sequence. *PeerJ* 2022;**10**:e14055.

Baidaliuk A, Lequime S, Moltini-Conclois I *et al.* Novel genome sequences of cell-fusing agent virus allow comparison of virus phylogeny with the genetic structure of Aedes aegypti populations. *Virus Evol* 2020;**6**:1.

Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 2014;**30**:2114–20.

Bolling BG, Weaver SC, Tesh RB *et al.* Insect-specific virus discovery: Significance for the arbovirus community. *In Viruses (Vol* 2015;**7**:9.

Boullé C, Njamnshi AK, Dema F *et al.* Impact of 19 years of mass drug administration with ivermectin on epilepsy burden in a hyperendemic onchocerciasis area in Cameroon. *Parasit Vectors* 2019;**12**, DOI: 10.1186/S13071-019-3345-7.

Brattig NW, Cheke RA, Garms R. Onchocerciasis (river blindness) – more than a century of research and control. *Acta Trop* 2021;**218**:7.

Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND. *Nature Methods 2014 12:1* 2014;**12**:59–60.

Camargo AP, Roux S, Schulz F *et al.* Identification of mobile genetic elements with geNomad. *Nature Biotechnology 2023* 2023:1–10.

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 2009;**25**:1972.

Chesnais CB, Bizet C, Campillo JT *et al.* A Second Population-Based Cohort Study in Cameroon Confirms the Temporal Relationship Between Onchocerciasis and Epilepsy. *Open Forum Infect Dis* 2020;**7**:6.

Chesnais CB, Nana-Djeunga HC, Njamnshi AK *et al.* The temporal relationship between onchocerciasis and epilepsy: a population-based cohort study. *Lancet Infect Dis* 2018;**18**:1278–86.

Colebunders R, Abd-Elfarag G, Carter JY *et al.* Clinical characteristics of onchocerciasis-associated epilepsy in villages in Maridi County, Republic of South Sudan. *Seizure* 2018a;**62**:108–15.

Colebunders R, Hendy A, Nanyunja M *et al.* Nodding syndrome-a new hypothesis and new direction for research. *International Journal of Infectious Diseases* 2014;**27**:74–7.

Colebunders R, Njamnshi AK, Menon S *et al.* Onchocerca volvulus and epilepsy: A
comprehensive review using the Bradford Hill criteria for causation. *PLoS Negl Trop Dis*
2021;**15**:1.

Colebunders R, Siewe Fodjo JN, Hopkins A *et al.* From river blindness to river epilepsy:
Implications for onchocerciasis elimination programmes. *PLoS Negl Trop Dis* 2019;**13**:7.

Colebunders R, Y. Carter J, Olore PC *et al.* High prevalence of onchocerciasis-associated
epilepsy in villages in Maridi County, Republic of South Sudan: A community-based
survey. *Seizure* 2018b;**63**:93.

Conceição-Neto N, Zeller M, Lefrère H *et al.* Modular approach to customise sample
preparation procedures for viral metagenomics: A reproducible protocol for virome
analysis. *Sci Rep* 2015;**5**, DOI: 10.1038/SREP16532.

De Coninck L. ViPER. 2021, DOI: 10.5281/zenodo.5502204.

Cook S, Chung BYW, Bass D *et al.* Novel virus discovery and genome reconstruction from
field rna samples reveals highly divergent viruses in dipteran hosts. *PLoS One*
2013;**8**:11.

Van Cutsem G, Siewe Fodjo JN, Dekker MCJ *et al.* Case definitions for onchocerciasis-
associated epilepsy and nodding syndrome: A focused review. *Seizure* 2023;**107**:132–5.

Cybinski DH, Gard GP. Isolation of a new rhabdovirus in Australia related to Tibrogargan
virus. *Aust J Biol Sci* 1986;**39**:225–32.

Cybinski DH, St. George TD, Standfast HA *et al.* Isolation of tibrogargan virus, a new
Australian rhabdovirus, from Culicoides brevitarsis. *Vet Microbiol* 1980;**5**:301–8.

Davis NM, Proctor DiM, Holmes SP *et al.* Simple statistical identification and removal of
contaminant sequences in marker-gene and metagenomics data. *Microbiome*
2018;**6**:1–14.

Duke BO, Vincelette J, Moore PJ. Microfilariae in the cerebrospinal fluid, and neurological
complications, during treatment of onchocerciasis with diethylcarbamazine.
Tropenmed Parasitol 1976;**27**:123–32.

Edgar RC. Muscle5: High-accuracy alignment ensembles enable unbiased assessments of
sequence homology and phylogeny. *Nature Communications 2022 13:1* 2022;**13**:1–9.

Edridge AWD, Abd-Elfarag G, Deijs M *et al.* Divergent Rhabdovirus Discovered in a Patient
with New-Onset Nodding Syndrome. *Viruses* 2022;**14**:210.

Ekangouo AE, Djeunga HCN, Sempere G *et al.* Bacteriome Diversity of Blackflies' Gut and
Association with Onchocerca volvulus, the Causative Agent of Onchocerciasis in Mbam
Valley (Center Region, Cameroon). *Pathogens* 2022;**11**, DOI:
10.3390/PATHOGENS11010044.

Föger K, Gora-Stahlberg G, Sejvar J *et al.* Nakalanga Syndrome: Clinical Characteristics,
Potential Causes, and Its Relationship with Recently Described Nodding Syndrome.
PLoS Negl Trop Dis 2017;**11**, DOI: 10.1371/JOURNAL.PNTD.0005201.

Fros JJ, Visser I, Tang B *et al.* The dinucleotide composition of the Zika virus genome is
shaped by conflicting evolutionary pressures in mammalian hosts and mosquito
vectors. *PLoS Biol* 2021;**19**, DOI: 10.1371/JOURNAL.PBIO.3001201.

Gibbs EPJ, Calisher CH, Tesh RB *et al.* Bivens arm virus: a new rhabdovirus isolated from
Culicoides insignis in Florida and related to Tibrogargan virus of Australia. *Vet Microbiol*
1989;**19**:141–50.

Grard G, Fair JN, Lee D *et al.* A novel rhabdovirus associated with acute hemorrhagic fever in
central Africa. *PLoS Pathog* 2012;**8**, DOI: 10.1371/JOURNAL.PPAT.1002924.

Guzman H, Contreras-Gutierrez MA, Travassos da Rosa APA *et al.* Characterization of Three
New Insect-Specific Flaviviruses: Their Relationship to the Mosquito-Borne Flavivirus
Pathogens. *Am J Trop Med Hyg* 2018;**98**:410.

Hadermann A, Amaral LJ, Van Cutsem G *et al.* Onchocerciasis-associated epilepsy: an update
and future perspectives. *Trends Parasitol* 2023;**39**:126–38.

Hendy A, Krit M, Pfarr K *et al.* Onchocerca volvulus transmission in the Mbam valley of
Cameroon following 16 years of annual community-directed treatment with
ivermectin, and the description of a new cytotype of Simulium squamosum. *Parasit*
Vectors 2021;**14**:563.

Hoshino K, Isawa H, Tsuda Y *et al.* Isolation and characterization of a new insect flavivirus
from Aedes albopictus and Aedes flavopictus mosquitoes in Japan. *Virology*
2009;**391**:1.

Hotterbeekx A, Lammens M, Idro R *et al.* Neuroinflammation and not tauopathy is a
predominant pathological signature of nodding syndrome. *J Neuropathol Exp Neurol*
2019;**78**:1049–58.

Hotterbeekx A, Raimon S, Abd-Elfarag G *et al.* Onchocerca volvulus is not detected in the
cerebrospinal fluid of persons with onchocerciasis-associated epilepsy. *International*
Journal of Infectious Diseases 2020;**91**:119–23.

Howerth EW, Mead DG, Stallknecht DE. Immunolocalization of vesicular stomatitis virus in
black flies (Simulium vittatum). *Ann N Y Acad Sci* 2002;**969**:340.

Hyatt D, Chen GL, LoCascio PF *et al.* Prodigal: Prokaryotic gene recognition and translation
initiation site identification. *BMC Bioinformatics* 2010;**11**:1–11.

Johnson BW. Neurotropic flaviviruses. *Neurotropic Viral Infections: Volume 1: Neurotropic
RNA Viruses* 2016:229–58.

Kraberger S, Schmidlin K, Fontenele RS *et al.* Unravelling the single-stranded DNA virome of
the New Zealand blackfly. *Viruses* 2019;**11**, DOI: 10.3390/V11060532.

Kuhn JH, Pan H, Chiu CY *et al.* Human Tibroviruses: Commensals or Lethal Pathogens?
Viruses 2020;**12**:3.

Kuno G, Chang GJJ. Biological transmission of arboviruses: Reexamination of and new
insights into components, mechanisms, and unique traits as well as their evolutionary
trends. *In Clinical Microbiology Reviews (Vol* 2005;**18**:4.

Levick B, Laudisoit A, Tepage F *et al.* High prevalence of epilepsy in onchocerciasis endemic
regions in the Democratic Republic of the Congo. *PLoS Negl Trop Dis*
2017;**11**:e0005732.

Lichty BD, Power AT, Stojdl DF *et al.* Vesicular stomatitis virus: Re-inventing the bullet.
Trends Mol Med 2004;**10**:5.

Lytras S, Hughes J. Synonymous Dinucleotide Usage: A Codon-Aware Metric for Quantifying
Dinucleotide Representation in Viruses. *Viruses 2020, Vol 12, Page 462* 2020;**12**:462.

Mead DG, Howerth EW, Murphy MD *et al.* Black fly involvement in the epidemic
transmission of Vesicular stomatitis New Jersey virus (Rhabdoviridae: Vesiculovirus).
Vector-Borne and Zoonotic Diseases 2004;**4**:351–9.

Mesquita LP, Diaz MH, Howerth EW *et al.* Pathogenesis of Vesicular Stomatitis New Jersey
Virus Infection in Deer Mice (Peromyscus maniculatus) Transmitted by Black Flies (
Simulium vittatum). *Vet Pathol* 2017;**54**:1.

Miranda J, Mattar S, Gonzalez M *et al.* First report of Culex flavivirus infection from Culex
coronator (Diptera: Culicidae), Colombia. *Virol J* 2019;**16**, DOI: 10.1186/S12985-018-
1108-2.

Mukendi D, Tepage F, Akonda I *et al.* High prevalence of epilepsy in an onchocerciasis
endemic health zone in the Democratic Republic of the Congo, despite 14 years of
community-directed treatment with ivermectin: A mixed-method assessment. *Int J
Infect Dis* 2019;**79**:187–94.

Nayfach S, Camargo AP, Schulz F *et al.* CheckV assesses the quality and completeness of
metagenome-assembled viral genomes. *Nature Biotechnology 2020 39:5* 2020;**39**:578–
85.

Neri U, Wolf YI, Roux S *et al.* Expansion of the global RNA virome reveals diverse clades of
bacteriophages. *Cell* 2022;**185**:4023-4037.e18.

Nurk S, Meleshko D, Korobeynikov A *et al.* MetaSPAdes: A new versatile metagenomic
assembler. *Genome Res* 2017;**27**:824–34.

Piégu B, Guizard S, Spears T *et al.* Complete genome sequence of invertebrate iridescent
virus 22 isolated from a blackfly larva. *Journal of General Virology* 2013;**94**:9.

Piégu B, Guizard S, Spears T *et al.* Complete genome sequence of invertebrate iridovirus IIV-
25 isolated from a blackfly larva. *Arch Virol* 2014a;**159**:5.

Piégu B, Guizard S, Yeping T *et al.* Complete genome sequence of invertebrate iridovirus
IIV22A, a variant of IIV22, isolated originally from a blackfly larva. *Stand Genomic Sci*
2014b;**9**:3.

Pion SDS, Boussinesq M. Significant Association between Epilepsy and Presence of
Onchocercal Nodules: Case-Control Study in Cameroon. *Am J Trop Med Hyg*
2012;**86**:557.

Price MN, Dehal PS, Arkin AP. FastTree 2 – Approximately Maximum-Likelihood Trees for
Large Alignments. *PLoS One* 2010;**5**, DOI: 10.1371/JOURNAL.PONE.0009490.

Quek S, Hadermann A, Wu Y *et al.* The RNA virome of human and animal parasitic
nematodes. *Res Sq* 2023, DOI: 10.21203/RS.3.RS-3265990/V1.

Quiroz E, Moreno N, Peralta PH *et al.* A human case of encephalitis associated with vesicular
stomatitis virus (Indiana serotype) infection. *Am J Trop Med Hyg* 1988;**39**:3.

Raimon S, Dusabimana A, Abd-Elfarag G *et al.* High Prevalence of Epilepsy in an
Onchocerciasis-Endemic Area in Mvolo County, South Sudan: A Door-To-Door Survey.
Pathogens 2021;**10**, DOI: 10.3390/PATHOGENS10050599.

Sadiq S, Holmes EC, Mahar JE. Genomic and phylogenetic features of the Picobirnaviridae
suggest microbial rather than animal hosts. *bioRxiv* 2024:2024.02.04.578841.

Sakaguchi S, Urayama SI, Takaki Y *et al.* NeoRdRp: A Comprehensive Dataset for Identifying
RNA-dependent RNA Polymerases of Various RNA Viruses from Metatranscriptomic
Data. *Microbes Environ* 2022;**37**, DOI: 10.1264/JSME2.ME22001.

Shintouo CM, Nguve JE, Asa FB *et al.* Entomological assessment of onchocerca species
transmission by black flies in selected communities in the west region of cameroon.
Pathogens 2020;**9**:9.

Siewe Fodjo JN, Mandro M, Mukendi D *et al.* Onchocerciasis-associated epilepsy in the
Democratic Republic of Congo: Clinical description and relationship with microfilarial
density. *PLoS Negl Trop Dis* 2019;**13**, DOI: 10.1371/JOURNAL.PNTD.0007300.

Siewe Fodjo JN, Ngarka L, Njamnshi WY *et al.* Onchocerciasis in the Ntui Health District of
Cameroon: epidemiological, entomological and parasitological findings in relation to
elimination prospects. *Parasit Vectors* 2022;**15**, DOI: 10.1186/S13071-022-05585-0.

Siewe Fodjo JN, Tatah G, Tabah EN *et al.* Epidemiology of onchocerciasis-associated epilepsy
in the Mbam and Sanaga river valleys of Cameroon: Impact of more than 13 years of
ivermectin. *Infect Dis Poverty* 2018;**7**:1.

Smith PF, Howerth EW, Carter D *et al.* Mechanical transmission of vesicular stomatitis New
Jersey virus by Simulium vittatum (Diptera: Simuliidae) to domestic swine (Sus scrofa). *J*
Med Entomol 2009;**46**:6.

Steinegger M, Meier M, Mirdita M *et al.* HH-suite3 for fast remote homology detection and
deep protein annotation. *BMC Bioinformatics* 2019;**20**:1–15.

Stremlau MH, Andersen KG, Folarin OA *et al.* Discovery of novel rhabdoviruses in the blood
of healthy individuals from West Africa. *PLoS Negl Trop Dis* 2015;**9**, DOI:
10.1371/JOURNAL.PNTD.0003631.

Vinkeles Melchers NVS, Mollenkopf S, Colebunders R *et al.* Burden of onchocerciasis-
associated epilepsy: First estimates and research priorities. *Infect Dis Poverty* 2018;**7**,
DOI: 10.1186/S40249-018-0481-9.

Winkler AS, Friedrich K, Velicheti S *et al.* MRI findings in people with epilepsy and nodding
syndrome in an area endemic for onchocerciasis: An observational study. *Afr Health Sci*
2013;**13**:529–40.

Wolf YI, Kazlauskas D, Iranzo J *et al.* Origins and evolution of the global RNA virome. *mBio*
2018;**9**, DOI: 10.1128/MBIO.02329-18/ASSET/42CAF1D8-B7AA-433C-9128-
816182956BD1/ASSETS/GRAPHIC/MBO0061842030010.JPEG.

Wolf YI, Silas S, Wang Y *et al.* Doubling of the known set of RNA viruses by metagenomic
analysis of an aquatic virome. *Nature Microbiology 2020 5:10* 2020;**5**:1262–70.

Young KI, Valdez F, Vaquera C *et al.* Surveillance along the Rio Grande during the 2020
Vesicular Stomatitis Outbreak Reveals Spatio-Temporal Dynamics of and Viral RNA
Detection in Black Flies. *Pathogens* 2021;**10**:10.

Phylum
- *Duplornaviricota*
- *Kitrinoviricota*
- *Lenarviricota*
- *Negarnaviricota*
- *Pisuviricota*
- *Pisuviricota* (dsRNA)
- Reverse transcriptases
- Neri *et al.* Cell (2022)
- Blackflies

Catalytic domain order
- ABC
- ★ CAB

Rhabdoviridae

Flaviviridae