Drosophila Hamlet mediates epithelial tissue assembly of the reproductive system

Huazhen Wang[1], Ludivine Bertonnier-Brouty[2,3], Isabella Artner[2,3], Jiayu Wen[4,*] and Qi Dai[1,*]

1 Department of Molecular Bioscience, the Wenner-Gren Institute, Stockholm University,
Sweden

2 Lund Stem Cell Center, Lund University, Lund, Sweden

3 Lund University Diabetes Center, Lund University, Malmö, Sweden

4 Division of Genome Sciences and Cancer, The John Curtin School of Medical Research, The
Australian National University, and Australian Research Council Centre of Excellence for the
Mathematical Analysis of Cellular Systems.

* Co-corresponding authors

Abstract

Epithelial tissue fusion requires coordinated molecular events at the ends of two epithelial structures. Regulatory mechanisms controlling these events remain largely elusive. In the *Drosophila* reproductive system (RS), this fusion unites the gonad and the genital disc-derived tissues, into a continuous tube. This study unveils the pivotal role of Hamlet (Ham), a *Drosophila* PR-domain containing transcription factor, in orchestrating epithelial tissue fusion in the RS. Loss of *ham* leads to sterility and disconnection between the testes (TE) and seminal vesicles (SV). Systematic analysis of Ham-downstream genes reveals cytoskeletal, metabolic regulators and signaling pathway components. Ham activates genes for epithelial differentiation and remodeling, while repressing genes required for tissue growth and patterning. Using multiplexed in situ hybridization, we demonstrate spatial-temporal gene expression dynamics in contacting epithelia. Key Ham downstream effectors include E-Cadherin (E-Cad), Toll (Tl) and Wnt2 signaling pathways, regulating tissue interaction and fusion. Our findings present a comprehensive gene network crucial for heterotypic epithelial tissue fusion. Mammalian Ham orthologs PRDM3 and PRDM16 are highly expressed in epithelial tissues, suggesting a conserved role across species.

Introduction

Epithelial tissue fusion is a crucial process for building epithelial tubular network and uniting separate ends [1, 2]. Studies from different model systems, like neural tube fusion and *Drosophila* tracheal development, have provided several mechanistic insights into epithelial fusion process. For instance, specialized cells at the leading edge migrate and interact using cytoskeleton protrusions [3-9] and small GTPases like Rac and Cdc42 [10]. Upon contact, the leading cells re-establish polarity and connections with the help of extracellular matrix molecules, apical proteins and E-Cad [11]. However, the exact regulators and mechanisms controlling epithelial fusion remain to be determined, especially in a context-specific manner.

The reproductive system in animals comprises tissues from different origins, and must unite to form a connected system for producing and delivering gametes. For example, the human testis is directly linked to a complex tubular system consisting of the efferent and epididymal ducts [12]. Anomalies in the fusion of these ducts are a significant cause of cryptorchidism, affecting at least three percent of newborn boys. However, there is little understanding of the molecular mechanisms guiding the patterning and fusion of these ducts.

Similarly, the *Drosophila* male RS (**Fig. 1A**) includes the gonad-derived testis and the genital disc (GD)-formed seminal vesicle, accessory gland (AG), ejaculatory duct (EJD) and additional supportive structures. During development, the TE and SV grow toward each other and merge into a continues tubular system [13]. This involves two epithelial ends, the terminal epithelial cells in the testis and the SV epithelium. Thus, the developing *Drosophila* RS provides a representative *in vivo* context to dissect mechanisms underlying epithelial tissue assembly.

Ham belongs to the PR domain containing protein (PRDM) family whose members are evolutionarily conserved cell fate regulators [14, 15]. The PRDM proteins are characterized by the presence of an N-terminal PR (PRDI-BF1 and RIZ1 homology) domain which is similar to the catalytic SET domain shared by histone methyltransferases. Dysfunction of the human orthologs of Ham, PRDM16 (Mel1) and PRDM3 (Evi1, MECOM), has been found in a variety of human abnormalities including congenital diseases, metabolic disorders and cancers [16-23]. Moreover, in at least two types of epithelial cancers (kidney and pancreas cancers), PRDM16 was shown to play a tumor suppressive role [20, 23], indicating its importance in

maintaining epithelial integrity. However, it remains undermined how these PR factors regulate

epithelial tissue development and homeostasis.

Ham was exclusively studied in the nervous system where it is required for neuronal dendrite

morphogenesis [24] and cell fate maintenance [25-27]. Ham was shown to repress gene

expression through modulating chromatin states [28] or interacting with the co-repressor CtBP

[25, 28]. However, the molecular functions of PRDM16 and PRDM3 are versatile, with their

PR domains possessing intrinsic histone methyltransferase activity [17, 29] and the ability to

interact with various co-activators and co-repressors [14, 30]. It remains unknown whether

Ham plays a role outside of the nervous system and whether Ham also has dual function in

gene expression.

In this study, we discovered that Ham is a master regulator in guiding epithelial tissue

patterning and fusion in the *Drosophila* RS. Even a slight reduction of Ham disrupts epithelial

differentiation and tissue assembly in the RS, ultimately affecting sperm and egg delivery. To

explore new mechanisms governing epithelial tissue fusion in the RS, we conducted systematic

analyses using RNA-sequencing (RNA-seq), Cleavage Under Targets and Tagmentation

(CUT&TAG), multiplexed in situ hybridization data and genetic screening. These analyses led

to identification of gene regulatory networks required for epithelial tissue fusion. Ham activates

a gene network to promote epithelial cell differentiation and remodeling while suppressing

another gene network involved in cell specification and early patterning. These target genes

exhibit spatial-temporal expression dynamics during the fusion of the TE and SV. As a key

Ham downstream effector, Wnt2 signaling regulates epithelial cell communication and

interaction. By elucidating the regulatory network controlled by Ham, our study provides novel

insights into the molecular mechanisms underlying epithelial tissue fusion.

Results

Drosophila RS development is vulnerable to reduced Ham activity

While PRDM16 and MECOM are critical cell fate regulators in various developmental contexts in mammals, the genetic roles of Ham were mainly found in the central and periphery nervous systems [24-27]. To systematically study Ham function, we generated new frameshift alleles using CRISPR/Cas9. The sgRNA targets the third exon shared by the three long isoforms B, D and G (**Supplementary Fig. 1A**), producing new alleles specifically disrupting the three PR-domain containing isoforms. We therefore named them $ham^{PR\Delta}$. The original ham^1 and the null allele ham^{SK1} shifted open reading frame (ORF) for all predicted isoforms [24, 26].

We validated nucleotide deletions in the new alleles by sanger sequencing (**Supplementary Fig. 1B**) and protein deletion by western-blot using an antibody against Ham. The Ham antibody detected several bands in the blot (**Supplementary Fig. 1C**). The two bands at and above the 150KD marker position in the two heterozygotic controls were not detectable in the ham^{SK1} homozygous mutant, suggesting that these two bands correspond to the longest and the second longest isoforms respectively. The $ham^{PR\Delta}$ homozygous samples lack the largest band, verifying the depletion of the longest isoforms. We used one of them (-5nt) for all the later experiments.

We compared viability and sensory organ development of these alleles. In contrast to the original report showing that *ham* mutant animals died at the larval stage [24], we found that ham^{SK1}, $ham^{PR\Delta}$ and ham^1 alleles all survived to hatch when combined in trans with the *ham* deficiency line (Df) which deletes the entire *ham* locus (**Supplementary Fig. 1D**). While ham^1/Df and ham^{SK1}/Df animals all died right after hatching, about two thirds of the $ham^{PR\Delta}$/Df mutants were viable without obvious gross defects. Furthermore, Ham is known to specify internal cell types (the neuron and sheath) in the mechano-sensory organ lineage, as ham^1 mutants exhibited ectopic external cell types (the socket and shaft) with the expense of internal cells [25]. $ham^{PR\Delta}$ mutants displayed normal sensory bristles, in contrast to defective sensory organs in the ham^1 and ham^{SK1} alleles (**Supplementary Fig. 1E**). Thus, the PR-containing isoforms of Ham are dispensable for mechano-sensory organ development and viability.

However, *ham^PRΔ* mutant animals showed abnormal RS morphology (**Fig. 1B**) and impaired

fertility in both male and female animals (**Fig. 1C, Supplementary 2A-B**). The TE, AG and

SV structures were disrupted in males, while mature eggs were stuck in the mutant female

ovaries, indicating failed egg delivery. To confirm that the defects are induced by the *ham*

mutation, we performed a rescue experiment by utilizing a pBAC transgene that carries the

genomic region of *ham* with the super-folding GFP (sfGFP*)* coding sequence in frame with the

ham ORF. The pBAC transgene rescued the viability of all *ham* mutant conditions, and at least

partly restored their male fertility (**Fig. 1C**). Together, these results indicate that the *ham^PRΔ*

allele represents a hypomorphic condition which preserves Ham shorter isoforms and partial

activity. Thus, although viability and sensory organ development do not require the PR-

containing isoforms, formation of the RS depends on them.

Ham-expressing epithelial cells connect the gonadal and supportive tissues

To assess how Ham controls RS development, we examined Ham expression in the developing

RS using a V5 antibody which detects all Ham isoforms in the *ham_V5* allele [31]. Ham_V5

was detected in the connecting tube of the adult SV and TE (**Fig. 1D**), suggesting a direct

involvement of Ham in regulating RS formation.

During RS development, the TE terminal epithelial cells are specified in the embryo as male-

specific somatic gonadal precursors (msSGPs) [32]. The SV epithelium, together with AG,

develops from a group of mesenchymal cells in the GD which undergo mesenchymal-to-

epithelial transition (MET) during the larval stage (**Fig. 1E**) [33, 34]. Notably, in the L3 larval

male RS, Ham_V5 was detected in the TE terminal epithelial cells as well as the btl-Gal4-

expressing mesenchymal cells in the GD (**Fig. 1F**).

In the pupal RS, multiple events, including epithelial tube formation and fusion as well as

myoblast and pigment cell migration, are tightly coordinated (**Fig. 1G**) [35]. At this stage, a

high level of Ham persists in the TE terminal epithelial cells, SV and AG (**Fig. 1H-J**). In

contrast, Ham is barely detectable in the SV myoblasts/myotubes labeled by the reporter, mef2-

Gal4-driving UAS-GFP (**Fig. 1J**). Similarly, high levels of Ham_V5 were found at the joint

sites of female oviducts and ovaries during both pupal and adult stages (**Supplementary Fig**

2C). Thus, Ham is an intrinsic factor in epithelial cells at the connection site between the

gonadal- and genital-disc derived tissues in both males and females.

Ham is required for epithelial tissue fusion in the developing male RS

The specific expression of Ham and mutant phenotypes in the RS prompted us to investigate

its function in RS epithelial tissues. Here we focused on the male, as the assembly of the TE

and SV represents a remarkable process involving a series of tissue reorganization events (**Fig.**

1G). When the two ends of TE and SV get close, the epithelial cells at the leading edges meet

and form a common epithelial tube. In the meanwhile, myoblasts from the SV and pigment

cells from the TE migrate toward the opposite direction. Myoblasts further differentiate into

myotubes, forming a thin layer of muscular sheet and sculpting the TE into a "spirals" shape

[36].

We found that at 40-50h APF the TE and SV tubes in *ham^{PRΔ}/Df and hamSK1/Df* mutants did

not properly elongate and appeared shorter and wider compared to that in the wild-type (**Fig.**

2A). The mutant TE appeared as a sphere shape, indicating failed myotube migration. The SV

and AG were only partly elongated in the *ham^{PRΔ}* mutant, and even unseparated in the *ham* null

allele *hamSK1/Df*. This suggests that the severity of RS abnormality correlates with the levels

of reduction in *ham* gene products.

To examine epithelial integrity, we performed immunofluorescent staining using the antibodies

against the epithelial cell markers, E-Cad (adherent junctions) and Coracle (Cor, septate

junctions equivalent to mammalian tight junction) protein. We also incorporated an SV-

specific transgenic reporter, btlGal4-driving UAS-EGFP or UAS-CD8GFP, to the control and

ham mutant genotypes. E-Cad is enriched at the apical side (the lumen side, **Fig. 2B**) of the TE

and SV epithelial tubes. Notably, the TE terminal epithelium accumulates a higher level of E-

Cad compared to the SV epithelium, while Cor is mainly expressed at the SV side but not much

in the TE, suggesting a heterotypic nature of the TE terminal and SV epithelia.

Control animals always showed connected TE and SV, whereas ham mutant TE and SV tissues

were either separated from each other, or appeared contacted but only with disconnected

epithelial tubes (**Fig. 2B, Supplementary Fig. 3A**). Importantly, this phenotype can be fully

rescued by the *pBAC_ham_sfGFP* transgene, confirming that it is caused by the *ham* mutation.

These observations suggest that the epithelial ends of the TE and SV possess distinct properties

despite both being epithelial cell types and that Ham is required for TE and SV fusion.

Depletion of Ham abolishes TE terminal epithelial cell formation

A close inspection on the pupal TE revealed that most of TE epithelial cells were missing in

ham mutant animals (**Fig. 2B, Supplementary Fig. 3A**). *ham* mutant lava also either lacked

TE epithelial cells or showed abnormal location of these cells, which can be rescued by the

pBAC_ham_sfGFP transgene (**Supplementary Fig. 4A**). TE terminal epithelial cells derive

from male specific gonadal precursors (msSGPs) specified in the early embryo

(**Supplementary Fig. 4B**) [33] and expand during larval and early pupal stages. This result

indicates that Ham regulates specification or survival of these cells. Indeed, we detected Ham

expression in msSGPs and other SGPs in the embryonic gonad. The number of msSGPs labeled

by Abd-B or Sox100B-GFP was dramatically reduced in S13 and S15 *ham* mutant embryo

(**Supplementary Fig. 4C-G**). Our data suggest Ham to be a master regulator for the

specification and/ or survival of msSGPs, and the absence of msSGPs accounts for loss of TE

terminal epithelial cells in larval and pupal stages.

Ham regulates patterning and cell differentiation of the RS epithelial tubes

The epithelial cells in the SV and AG originate from a group of mesenchymal cells recruited

by the FGF pathway involving Btl and Branchless (Bnl) [33, 34]. These cells undergo MET to

form an epithelial mass which then cavitates to generate inner hollow space and eventually

elongates into an epithelial tube. In comparison with the control, the signal intensity of both E-

Cad and Crumbs (Crb) were significantly reduced in *ham* mutant SV epithelial cells, which

coincides with the shortening of septate junctions labeled by Cor (**Fig. 2C-E**). This result

suggests that Ham regulates epithelial cell differentiation in the SV. To check whether Ham

regulates epithelial cell differentiation in the TE, we used the FRT/FLP system (see Methods)

to generate mosaic somatic mutant clones. In these animals, *ham* mutant cells were GFP

negative, while wild-type and *ham* heterozygous cells expressed GFP (**Fig. 2F,**

Supplementary Fig. 3B). The E-Cad levels at the border between two mutant cells were

significantly lower than those between wild-type cells, indicating that *ham* depletion impaired

differentiation of TE terminal epithelial cells.

In summary, these phenotypic analyses suggest that Ham controls formation of TE terminal epithelial cells in the early embryo and regulates epithelial tube elongation and cell differentiation at the pupal stage. Thus, failed epithelium fusion in *ham* mutant RS could result from: 1) absence of TE terminal epithelial cells, 2) defects in epithelial tube elongation and cell differentiation of the SV and 3) a combination of both.

Ham directly controls epithelial fusion between the TE and SV

Next, we sought to assess whether and how Ham directly regulates the fusion process. We first attempted to generate a mild mutant condition which can bypass defects of TE epithelial cell loss and SV-AG tube elongation. We chose RNAi knockdown because knockdown efficiency can be enhanced by overexpressing Dicer2 (Dcr2) and modulated by temperature when RNAi is controlled by the GAL4/UAS system.

We first identified a Gal4 driver expressed in the SV and AG epithelial cells, after screening the available Gal4 transgenes driven by a series of genomic fragments spanning the *ham* locus [37] (**Supplementary Fig. 5A**). This transgene (thereafter named as *hamRSGAL4*) induces high expression of *UAS-GFP* in the TE and SV epithelial cells and a weak expression in the myoblasts (**Supplementary Fig. 5B**). Myotubes appeared at the junction site of the TE and SV and could migrate to the surface of the testis in the *ham* mutant (**Supplementary Fig. 5C-D**), confirming that loss of *ham* does not abolishes myoblast fusion or myotube migration. *hamRSGAL4,* when driving a *ham* RNAi line, nearly abolished Ham expression in animals carrying the *UAS-dcr2* transgene (**Fig. 2G-H,** the middle panels). These animals also had a reduced number of TE epithelial cells and displayed a ball-like structure of the GD, similar to the *ham* null allele. Without *UAS-dcr2*, *ham* RNAi knockdown reduced Ham expression but preserved TE epithelial cells and normal-appearing SV. Remarkably, the TE and SV epithelia still failed to fuse (**Fig. 2H,** the right panel), suggesting that Ham also controls the late step of fusion. Furthermore, E-Cad staining intensity was significantly reduced in *ham* RNAi TE but not in *ham* RNAi SV (**Fig. 2I-J, Supplementary 3C-D**), suggesting that expression of E-Cad in the TE is sensitive to even mild reduction of Ham protein levels.

Ham controls distinct molecular pathways to promote tissue assembly.

To identify mechanisms that control epithelial fusion, we started to investigate Ham
downstream target gene network. We performed RNA-seq experiments from dissected TE and
SV tissues during the stage of 18-25h APF. We used two pairs of genotypes, including the
wild-type (*wt*), and *ham^PRΔ* homozygous and the *wt/Df* and *hamSK1/Df* trans-heterozygous
(**Fig. 3A**). The comparison between the first pair will determine dysregulated genes in the
ham^PRΔ hypomorphic condition, while the comparison between the second pair will identify
genes that change expression between the null and *ham* heterozygous conditions.
304
The principal component analyses validated consistency between the three biological replicates
(**Supplementary Fig. 6A**). To not exclude small gene expression changes (*e.g.* in the leading
cells which are the minority within the whole tissue), we used a loose cutoff (FDR < 10% and
FC>1). We found more than two thousand up-regulated and down-regulated genes in the GD.
In the mutant TE, there were similar numbers of dysregulated genes (**Fig. 3B**). The genes that
only changed expression in one condition could result from unspecific genetic background,
differential dependence on the PR-containing isoforms or different sensitivity to *ham*
heterozygous condition.
313
Expression of many dysregulated genes can be indirectly affected by Ham. We next intended
to identify genes that are transcriptionally regulated by Ham. To this end, we determined Ham
genomic binding sites using Cleavage Under Targets and Tagmentation assays (CUT&TAG)
[38] on finely-dissected TE and grossly-dissected abdominal tissues in which Ham is mainly
expressed in the TE and SV epithelium (**Fig. 3A**). We used a Ham antibody for wildtype
tissues, a GFP antibody for tissues from the *pBAC_ham_SFGFP* transgene and IgG as a
negative control in parallel for all the samples. Using the SEACR software, we identified peaks
and intersections of peaks across the Ham and GFP antibody conditions (see Methods). We
performed principal component analyses and found the GFP antibody gave more consistent
peak calling from replicates (**Supplementary Fig. 6B**). For subsequent analyses, only the
common 3433 and 691 peaks identified by the two antibodies (Ham and GFP) were included
to ensure signal specificity (**Fig. 3C-D**).
326
To determine Ham target genes, we intersected the RNA-seq data from the hamSK1/Df versus
wt/Df comparison with the Cut&Tag data. Among the 2,546 up-regulated genes in the GD, 441
(17.3%) contained Ham CUT&TAG peaks. Among the 2379 downregulated genes, 569

(23.9%) showed Ham binding (**Fig. 3E**). In the TE, a lower fraction of dysregulated genes

contains Ham CUT&TAG peaks, including 4.7% (108/2274) upregulated genes and 7%

(144/2057) downregulated genes (**Fig. 3F**), suggesting that dysregulated genes in the mutant

TE are mainly indirectly controlled by Ham.

Both up- and down-regulated genes contain Ham binding peaks, suggesting that Ham can

repress and activate gene expression. To systematically explore the function of Ham-regulated

genes, we performed gene network analysis using STRING [39]. We focused on the genes

bound by Ham and also dysregulated in the genital disc (GD) in the null condition *hamSK1*,

considering that most of TE genes are only indirectly influenced in the *ham^{SK1}* mutant. The

SRING MSL clustering analysis generated 114 functional clusters from down-regulated genes

and 62 clusters from up-regulated genes (**Fig. 4A, Supplementary Table 1**). Interestingly,

many of downregulated genes clustered into several large groups including cytoskeleton

regulators, metabolic factors, cell adhesion and extracellular matrix proteins, suggesting that

Ham activates genes to promote epithelial differentiation and remodeling. In contrast,

upregulated genes are enriched for patterning factors, cell cycle genes and transcriptional and

translational regulators, suggesting that Ham suppresses early developmental processes.

Notably, metabolic factors are enriched in both categories, but these factors belong to distinct

functional groups. Thus, epithelial fusion requires complex and balanced metabolic processes.

Together, this network analysis confirms that Ham positions on top of distinct molecular

pathways to facilitate TE and SV assembly.

Ham was known for its repressive activity via interacting with the co-repressor CtBP and

influencing histone modifications [28]. Our finding of its activation function was unexpected.

To validate the activation activity of Ham, we chose three downregulated target genes (**Fig.**

4B), *dwnt2*, *shot* and *shg*, for transcriptional reporter assays. *shg* encodes E-cad, while Shot is

a member of the spectraplakin family of large cytoskeletal linker molecules simultaneously

binding to actin and microtubules. The function of *wnt2* was reported in the developing RS

development [40, 41]. We cloned four genomic fragments from the *wnt2* locus, two each from

shg and *shot* into a luciferase construct, and co-transfected each reporter with individual

plasmids expressing several variants of Ham. In addition to the Ham long isoform and a PR

truncation version of Ham, we included the NZF and the CZF domains fused with the VP16

activation domain (NZF_VP16 and CZF_VP16) (**Fig. 4C**). The two ZF domains of Ham are

putative DNA binding domains, but it remained undetermined whether they help to recruit Ham

to DNA. This experimental setting allows to assess which of the ZF domains is able to target

Ham-regulated enhancers. The full-length, the PR truncation and the NZF_VP16 fusion all

activated expression of the reporters driven by 7 out of 8 Ham-bound elements (**Fig. 4D-G,**

Supplementary Fig. 6C-D). By contrast, the CZF_VP16 had little activity on the reporters.

This result confirms that Ham can directly activate target gene expression and that the NZF

domain directly or indirectly recruits Ham to enhancers.

To further identify co-factors with which Ham interacts to regulate gene expression, we

performed *de novo* motif discovery from Ham CUT&TAG peaks in mixed TE and GD tissues

close to up-regulated and down-regulated genes in the *hamSK1* mutant using the MEME-ChIP

software. This resulted in nine and five significantly enriched motifs from up- and down-

regulated gene loci respectively (**Supplementary Fig. 6E**). While five motifs were shared

between these two sets, the motifs for Dpp/BMP signaling factor MAD, the zinc-finger protein

ZNF263 and two unknown factors were only enriched in the upregulated genes. The result

implies that Ham may switch activation to repression activity when interacting with these co-

factors in the context of RS development.

Ham downstream effector genes are required for male RS fertility.

Built upon our transcriptomic data, we next intended to identify genes crucial for the fusion

process. We reasoned that as the majority of TE terminal epithelial cells are lost in both *ham*

mutant alleles, the commonly downregulated genes in *ham^{PRΔ}* and *ham^{SK1}* TE (**Fig. 3B**) may

be expressed in the *wt* TE epithelial cells and can be functionally important. We thus focused

on the top 241 common downregulated genes whose expression showed a fold change greater

than 1.5, and the three validated Ham target genes: *shot*, *wnt2* and *shg*. Using all available

RNAi lines from the Bloomington stock center that cover 113 out of 244 candidate genes, we

performed a UAS-RNAi screen with the *UAS-dcr2*; *hamRSGal4* driver for male fertility (see

Methods).

The RNAi screen led to identification of 26 genes whose knockdown abolished or reduced

male fertility (**Supplementary Table 2**). Knocking down four of these genes, *shg*, *AP2-alpha*,

pebble (pbl) and *CG11406*, resulted in complete sterility, indicating their importance for male

fertility. We defined this category as 100% penetrance of sterility. Knockdown of other twenty-

two genes reduced fertility to different extent (**Fig. 5A-B**). Given that the *hamRSGal4* driver
is highly expressed in the TE and SV epithelia, we expect highly effective knockdown occurs
only in these epithelial cells. However, *hamRSGal4* also drives a weak expression in the
myoblasts before they differentiated into myotubes (**Supplementary Fig. 5B)**, which may
result in a non-tissue autonomous effect when knocking down the candidate genes expressed
in myoblasts.

Consistent with the gene network analysis (**Fig. 4A**), these genes include cytoskeleton
regulators (*pbl*, *shg*, *shark*) [42-44] and metabolism factors (*CG11406, Ph4alphaMP, GstE11,*
CG17477, CG11069, AdSL, CG17283, CG15117, CG7484). Other genes are known or
predicted to be involved in intracellular membrane trafficking (*AP-2alpha*, *CG31935* and
CG7456) [45] and signaling (*wnt2*, *Toll*, *AkhR*, *Idgf4* and *Hr46*) [46-49].

We confirmed that knocking down *pbl*, *AP-2alpha, CG11406* and *shg* led to failed TE and SV
fusion and abnormal morphology at the pupal stage (**Fig. 5C, Supplementary Fig. 7-8)**. *pbl*
encodes a Rho guanine nucleotide exchange factor, implicated in multiple processes involving
cytoskeleton reorganization[42, 50, 51]. The gene product of CG11406 is a predicted
triglyceride lipase, cleaving triacylglycerol (TAG) to produce diacylglycerol (DAG). AP-
2alpha, a subunit in the Adaptor complex 2, is involved in intracellular trafficking and secretion
[52]. Remarkably, *shg* knockdown also led to aberrant mobilization of TE epithelial cells to
the middle of the TE, while no such effect was found in the SV and AG. Downregulation of E-
Cad is a key event during epithelial to mesenchymal transition (EMT), and E-Cad promotes
cohesive cell migration of mesenchymal cells, but suppressing cell motility of epithelial cells
[53, 54]. Our result suggests that the SV and the TE terminal epithelia have distinct dependence
on E-Cad and that TE epithelial cells are prone to becoming motile when E-Cad is reduced.
Together, this screen identified a collection of known and novel factors involved in epithelial
tissue fusion, confirming that the fusion process requires coordinated activity of several classes
of molecules.

Ham downstream genes exhibit spatial- temporal expression dynamics in the male RS

During epithelial fusion, cells at the leading edge become specialized and express specific
genes required for tissue interaction and remodeling [1, 2]. Moreover, as we observed, the TE

terminal and SV epithelia have distinct properties, suggesting that these two sets of epithelial

cells may express distinct genes. We thus intended to assess spatial distribution of Ham

downstream genes. We conducted a high-resolution multiplexed RNA *in situ* hybridization

assay known as SCRINSHOT (single cell RNA *in situ* hybridization on tissues) [55] to

visualize expression of candidate genes. These genes include *wnt2*, *shg* and *shot* as these genes

were either known for male RS development (*wnt2 [40]*) or epithelial tissue fusion (*shg* [56]

and *shot [57]*), and the positive genes from the RNAi screen, along with the receptor genes in

Wnt2 signaling (*fz*, *fz2*, *fz3* and *fz4*) and ligand genes in Tl-signaling (*spz*, *spz4*, *spz5* and *spz6*)

(**Supplementary Table 3**). Some of these Wnt2 and Tl signaling genes displayed dysregulation

in at least one of the *ham* mutant alleles according RNA-seq data (**Supplementary Fig 9A-B**).

Other receptor genes in Wnt signaling and ligand genes in Tl signaling were omitted due to

low read numbers in the RNA-seq data, indicating negligible or minimal expression in the TE

and GD.

Each SCRINSHOT reaction took place within a mini-chamber, accommodating pairs of control

and *ham* mutant tissues, or GFP KD control and ham RNAi KD samples (refer to Methods for

details of probe design and experimental procedures). This setup was designed to minimize

technical variability between the genotypes. mRNAs are visualized as fluorescent dots in

SCRINSHOT images, with each dot representing a single mRNA molecule. However, due to

the densely packed nature of cells in the TE and SV tissues, quantifying mRNA molecules at

a single-cell resolution is challenging. Therefore, we opted to quantify fluorescent signal

intensity within defined areas (see Methods). In total, 37 genes were tested, including *ham*

itself (**Supplementary Table 3**). Probes for 7 genes gave either no or low signal in the TE or

GD, likely attributed to poor probe quality, given their RNA-seq read numbers not being low.

Resembling Ham protein expression (**Fig. 1F**), *ham* mRNAs were detected in the TE terminal

epithelium (**Fig. 6A, Supplementary Fig. 10A**), confirming the specificity of the

SCRINSHOT method. Similarly, *wnt2*, *shg* and *shot* all exhibited high expression in the TE

terminal epithelium (**Fig. 6A, Supplementary Fig. 10A**).

In total, 25 genes exhibited detectable SCRINSHOT signal in the wt SV and AG, while 29

displayed signals in the wt TE (**Supplementary Table 3**). Within the TE, 17 out of 29 were

specifically localized in the TE terminal epithelial cells, with the remaining 12 detected only

in other TE cells (**Fig. 6A, Supplementary Fig. 10-13, Supplementary Table 3**).

Subsequently, our analysis focused on the genes expressed in the TE terminal and SV epithelia.

While 17 genes exhibited expression in both epithelial ends (**Fig. 6A-B, Supplementary Fig.**

10A-B), 8 were exclusively expressed in SV epithelial cells (**Fig. 6F**, **Supplementary Table**

3), underscoring the heterotypic nature of the TE and SV epitheliums. Notably, certain genes

(CG11406, *AP-2alpha*, *GstE11*, *Tl, spz5* and *spz6*) demonstrated temporal dynamic expression

in the TE epithelium: spz5 and CG11406 are present during the 20-25h APF timeframe, prior

to fusion, but gone during the 28-32h APF period when the TE and SV are undergoing fusion;

AP-2alpha, *GstE11*, *Tl* and *spz6* showed the opposite trend, being expressed only later (**Fig. 6,**

Supplementary Fig. 10-12).

In the GD, while the majority of genes exhibited ubiquitous expression, *wnt2* and *spz* were

predominantly expressed in the leading cells within the SV, contrasting with their expression

in the remaining GD (**Fig. 6B**). Additionally, the expression patterns of *fz2*, *fz3* and *Tl* displayed

spatial specificity: fz2 exhibited higher expression levels in the SV compared to the AG, *fz3*

demonstrated the opposite pattern with higher expression in the AG, and *Tl* was primarily

expressed in the myoblasts (**Fig. 6B**). These findings imply that Wnt2 and Tl signaling may

have cell-type-specific roles in the fusion process. Thus, SCRINSHOT mapped 30 candidate

genes in the developing GD and TE, with 29 of them display spatial or temporal dynamics.

Wnt2 and Tl signaling pathways are key mediators of Ham function in RS tissue assembly

Next, we aimed to identify genes exhibiting altered expression in *ham* mutant SV and TE. As

hamSK1 mutant TE lacks terminal epithelial cells, we excluded TE samples and only quantified

SCRINSHOT signal levels in the developing SV between *wt* and the *hamSK1* mutant. In

contrast, *ham* RNAi knockdown resulted in fusion defects but only mildly impacted SV

morphology and TE terminal epithelial cell number (**Fig. 2G-H**). These samples allow to assess

gene expression alterations in both the SV and TE epithelia.

ham fluorescent intensity appeared comparable between *wt* and mutant samples (**Fig. 6B, D-**

E, Supplementary Fig. 10-13), suggesting that *ham* mutant RNAs, containing the CRISPR-

generated indels, remain stable at this stage. In the *hamSK1* mutant SV, seven genes became

downregulated, while two exhibited upregulation. The downregulated genes were four Wnt2

pathway genes (*wnt2, fz, fz2, fz4*), one Tl pathway gene (*spz*), and *shg* (**Fig. 6B-**C,

Supplementary Fig. 10). Notably, *wnt2*, *fz*, *spz* and *shg* also displayed downregulation in *ham*

RNAi SV (**Fig. 6D-E, Supplementary Fig. 10**), suggesting that these genes and their

associated signaling pathways may be directly involved in the fusion process downstream of

Ham. The upregulated genes in *hamSK1* mutant SV include *GstE11* and *CG11069*

(**Supplementary Fig.11C, E**), which exhibited low expression in the leading cells of wt SV

epithelium, but increased expression under the *hamSK1* condition. In *ham* RNAi knockdown

animals, *CG11069* was again upregulated, while *GstE11* did not change expression. Instead,

another ligand gene in Tl signaling (*spz5*) displayed increased expression (**Supplementary**

Fig.11D, F). On the TE side, only four genes showed expression alteration in *ham* RNAi

animals, including *ham* itself, *shg*, *fz* and *fz2* **Supplementary Fig.11G**), implicating functional

requirement of these genes in both TE and SV epithelia.

The observation that knocking down *Tl* resulted in decreased male fertility (**Fig. 5B**,

Supplementary Table 2), coupled with the expression of multiple *spz* ligand genes in the

developing TE and SV, suggests that the Tl signaling pathway might be required for assembling

the TE and SV. To investigate this possibility, we examined male flies with *spz* knockdown

driven by *hamRSGal4*. These animals produced a comparable number of progenies to control

animals. The lack of phenotype in this condition could be attributed to inefficient knockdown

or genetic compensation by another *spz* gene, like *spz5* whose expression increased in *ham*

RNAi animals when *spz* expression was reduced. In addition to Tl, two other Tl related genes

(Toll-7 and Toll-9) exhibited moderate number of reads in the RNA-seq data. We then tested

male fertility in *Toll-7* and *Toll-9* knockdown animals. *Toll-9* knockdown led to a drastic

reduction in progeny numbers compared to the RNAi control (**Supplementary Fig. 14A**).

Consistent with the sterility, some of the *Toll-9* knockdown males displayed abnormal TE

shape (**Supplementary Fig. 14B**). Remarkably the majority of Toll-9 knockdown animals

showed empty SV without sperms (**Supplementary Fig. 14C**), suggesting failed connection

between the TE and SV. Together, these results pinpoint an essential role of Tl signaling in

male RS development.

To validate the role of Wnt2 signaling in RS development, we investigated four genetic

conditions, aimed at modulating its level and activity: *wnt2^O* (a *wnt2* null allele), *wnt2*

overexpression driven by *hamRSGal4*, *fz* and *fz2* RNAi knockdown driven by *hamRSGal4*. The

wnt2^O mutant TE and GD never connect, suggesting that Wnt2 is essential for the fusion

process. The *wnt2^O* mutant TE lacks the pigment cell and myocyte layers (**Fig. 7A**), consistent

with its reported function in these two cell types [35, 40]. The *wnt2^O* mutant GD failed to form

elongated SV and AG, suggesting that Wnt2 is required for GD patterning and differentiation.

Moreover, overexpression of *wnt2* also resulted in fusion defects and mis-localization of TE

epithelial cells (**Fig. 7B**), a phenotype reminiscent of *shg* knockdown animals. Indeed, *wnt2-*

overexpressing cells displayed reduced levels of E-Cad, suggesting that Wnt2 signaling might

balance E-Cad levels to maintain TE epithelial cell properties. Consistently, *fz2* knockdown

animals exhibited comparable phenotypes, including mobilized TE epithelial cells and failed

tissue fusion (**Fig. 7C**). *fz2* and *fz* knockdown also decreased male fertility (**Fig. 7D**).

Collectively, these results suggest the importance of optimal Wnt2 signaling levels in male RS

development and epithelial tissue fusion.

Discussion

In this study we demonstrate that the transcription factor Ham plays a crucial role in controlling epithelial fusion during *Drosophila* RS development. Early in embryonic development, Ham determines the cell fate of msSGPs, which are progenitor cells for TE terminal epithelial cells. Later on, Ham directly regulates the differentiation of epithelial cells in both the TE and SV by orchestrating a gene expression program in epithelial cells. While our primary focus is on male RS, we found that Ham is also expressed in cells at the fusion site of the oviduct and ovary. Mutant females lacking functional Ham exhibit a non-functional oviduct epithelial tube, suggesting that Ham likely employs a similar mechanism to control epithelium fusion in females.

By examining *ham* mutant phenotypes and its downstream target genes, we identified several classes of novel factors involved in fusion process of SV and TE. These genes, directly or indirectly controlled by Ham, encompass a diverse array of molecular functions. This discovery provides a comprehensive gene network for future investigation of epithelial fusion processes. The cell-type-specific and tissue-specific spatial distribution of these genes suggest that they may regulate various aspects of fusion, including cell migration, intercellular communication between myoblasts/myotubes and epithelial cells, specification of the leading cells, and other steps in the fusion process. For instance, Tl and its related receptors is known to mediate cell adhesions [58]. We can speculate that Tl signaling and the molecule itself promote interaction between myoblasts and myoblasts and epithelial cells (**Fig. 7E**), based on high expression of Tl in myoblasts and the presence of multiple Spz ligands in the TE and SV epithelial cells. Moreover, *Wnt2*, *fz* and *fz2* are highly expressed in the leading cells of TE and SV epithelia, and also required for TE and SV connection, suggesting that Wnt2 signaling mediates epithelial cell communication, remodeling and fusion.

Despite being discovered more than twenty years ago [24], research on the function of Ham function has primarily focused on its roles in the nervous system. Our findings highlight a crucial function of Ham in epithelial tissues. Apart from the RS, we observed high expression of the Ham protein in other epithelial tissues including the Malpighian tubes (akin to the kidney in insects) and the intestine (data not shown). Furthermore, orthologs of Ham, PRDM16 and PRDM3/MECOM, exhibit high expression in epithelial tubular structures, such as Lung, pancreas, kidney, urinary bladder and gastrointestinal tract and lung, according to observations

in the Human Protein Atlas (**Supplementary Fig. 15**.
https://www.proteinatlas.org/ENSG00000142611-PRDM16/single+cell+type;
https://www.proteinatlas.org/ENSG00000085276-MECOM/single+cell+type) [59]. Notably,
Seminal vesicle and ductus deferens, the equivalent structure to fly seminal vesicle, are the top
one and three tissues with the highest PRDM16 expression (**Supplementary Fig. 15A**). This
suggests a potential role for these proteins in controlling epithelial integrity and fusion in
mammalian organogenesis. Indeed, we observed that the pancreas duct volume in mouse
embryos lacking *Prdm16* is significantly reduced (data not shown). Although the exact cellular
changes and the underlying molecular mechanisms in this context await further investigation,
this phenotype underscores a crucial function of *Prdm16* in pancreatic epithelial duct
formation.

We show that compared to Ham function in the nervous system, the PR-containing isoforms
are indispensable for RS epithelial fusion and differentiation, and that Ham can directly activate
epithelial gene expression. It remains to be determined whether Ham has distinct molecular
functions or it regulates a subset of common molecular pathways in these two contexts.
Dendrite morphogenesis involves outgrowth and branching, somewhat analogous to epithelial
tube growth and branching. For instance, molecular mechanisms regulating polarized protein
trafficking are crucial for both dendrite and epithelial tube branching [60]. Additionally, several
transcription factors like Sequoia [61, 62], Tramtrack [63, 64] and Cut [65, 66] play roles in
both dendrite morphogenesis and tracheal tube fusion. These examples suggest that there might
be more unrecognized common mechanisms between dendrite and epithelial morphogenesis.
It is of interest to test whether the effectors we identified in the RS epithelium are involved in
dendrite morphogenesis in the nervous system.

We observed a unique aspect of epithelial fusion in the *Drosophila* RS. Unlike many epithelial
fusion processes where two homologous ends merge, such as those in mammalian neural tube
closure and *Drosophila* tracheal formation *etc* [1, 8, 10], the two epithelial ends in the
Drosophila RS are heterotypic. Specifically, the TE terminal epithelial cells express higher
levels of E-Cad and lower levels of Cor compared to SV epithelial cells. In response to the
reduction of E-Cad or ectopic expression of *wnt2*, TE terminal epithelial cells become motile,
resembling a process akin to epithelial-to-mesenchymal transition. This suggests that TE
terminal epithelial cells may be more invasive during epithelial fusion. On the other hand, SV
epithelial cells, despite originating from mesenchymal cells, exhibit typical epithelial cell

characteristics, including regular Cor and E-Cad expression. Notably, many genes are expressed exclusively in the SV epithelium, distinguishing it from the TE epithelium. Heterotypic epithelial fusion occurs in many developmental processes, such as digestive system formation, vertebrate optic fissure closure, and mammalian urogenital development. Failure of epithelial fusion can lead to a wide variety of developmental diseases such as oesophageal atresia, spina bifida, ocular coloboma, congenital abnormalities of the kidney, urinary tract and persistent cloaca. Studying the basic processes and key players of normal epithelial tissue fusion in an *in vivo* context will help to understand the pathogenesis of these diseases.

Acknowledgements

We thank the imaging facility (IFSU) of Stockholm University, the BEA sequencing facility at Karolinska Insitute, and the National Genomic Institute of Scilife Laboratories, Sweden, for providing service and support. The Bloomington Stock Center and the *Drosophila* community provided important reagents and fly stocks. We thank Christos Samakovlis's lab for helping to set up SCRINSHOT and Eric Lai's lab for the initial Ham reagents. Funding from Australian Research Council Centre of Excellence for the Mathematical Analysis of Cellular Systems (CE230100001) and ANU Future Scheme to J.W. The project was supported by project grant from Swedish Research Council (Vetenskapsrådet, 2020-03543) and Swedish Cancerfonden (CAN 22 2161 Pj) to Q.D.

Author contributions

Conceptualization and project design by QD, experiments conduction by HW, preliminary test in pancreas by IA and LB, computation analysis by JW. Data interpretation by HW, JW and QD. Manuscript writing by QD with input from HW and JW.

Materials and methods

Fly strain culturing and generation of transgenes

All fly stocks were maintained at 22°C. The $ham^{PR\Delta}$ mutants were created using CRISPR: transgenic flies carrying single guide RNA targeting the PR domain containing isoforms were crossed into the vas-Cas9 transgenic flies. Frame-shift mutations were identified by PCR and Sanger sequencing. Transgenic flies were created at BestGene, Inc.

The ham Gal4 lines and RNAi stocks were obtained from Bloomington Stock Center. Knocking down experiments were performed at 25 °C and with a regular 12-hr light–dark cycle.

To generate ham null mutant somatic clones in the TE terminal epithelium, *hamsk1 FRT40A/CTG* males were crossed with virgin females carrying the genotype *hsFLP(X); hsGFP.nls FRT40A*. Embryos aged 0-5 hours were collected and developed at 25°C until the 2nd instar larval stage. GFP-negative larvae, indicating the genotype *hsFLP(X); hsGFP.nls/hamsk1 FRT40A*, were then selected. Head shock treatments at 37°C for 1 hour were administered three times during the 2nd and 3rd instar larval stages to induce somatic clones. Prior to dissection, another 1-hour head shock at 37°C was performed to induce GFP expression, marking the wild-type cells.

Fertility test

For ham mutants, single male or female with indicated genotype was mated with two w^{1118} virgin females or two w^{1118} males on day 0 of the experiment. Crosses were flipped every 2 days for 3 consecutive times. Progenies from the second vial with eggs laid between day 2-4 were counted after 2 weeks.

For the RNAi screen, individual UAS-RNAi males were crossed with females carrying UAS-dcr2 and hamRSGal4. The resultant F1 male progenies (2-5 days old) with the genotype UAS-dcr2; hamRSGal4>UAS-RNAi were crossed to w^{1118} virgin females (1-2 days old) on day 0 of the experiment. In each test, four males were crossed with three females. Crosses were flipped on day2 and day3 for pre-laying. After flipping on day3, crosses were kept in the new vial for egg laying for a specific time period (GroupA: 48hours; GroupB: 16-18h; GroupC: 7-8h). If at

702 least three males and two females were alive, then the flies were transferred to a new vial, the
703 old vial was kept, and all progenies were counted.
704
For fertility test of the Wnt2 and Tl signaling pathways components, UAS-RNAi males were
crossed to virgin females of the genotype UAS-dcr2; hamRSGal4. Two F1 male progenies (1-
2 days old) with the genotype UAS-dcr2; hamRSGal4>UAS.RNAi were crossed to three w[1118]
virgin females (1-2 days old) on day 0 of the experiment. Crosses were flipped every day for 4
consecutive times. Progenies from the fourth vial with eggs laid between day 3-4 were counted
after 2 weeks.
711
712 *Immunostaining*
713 For embryo staining: embryos were collected, dechorionated with bleach and fixed with 4%
714 formaldehyde in PEM (0.1M PIPES, 2mM EGTA, 1 mM MgSO4, pH 7) for 20 min. For
715 immunostaining, embryos were rehydrated, washed with PBT (0.3% Triton X-100 in PBS) and
716 blocked with PBSBT (0.3% Triton X-100, 1% BSA in PBS).
717
718 For tissue staining: tissues were dissected from stage defined larvae or pupae in 1xPBS within
719 30 min and fixed in 4% paraformaldehyde in PBS for 20min. For immunostaining, dissected
720 tissues were washed with PBT (0.1% Triton X-100 in PBS) and blocked with PBSBT (0.1%
721 Triton X-100, 1% BSA in PBS).
722
723 For both embryo and tissue staining, the primary antibodies were diluted with PBSBT (0.1%
724 Triton X-100, 1% BSA in PBS) and incubated at 4°C over night. After extensive washing with
725 PBSBT (15min x 4 times), the secondary antibodies were added in 1:500 dilution and incubated
726 at room temperature for 2h. After washing with PBT (0.1% Triton X-100 in PBS), the samples
727 were counter-stained with DAPI and mounted in Vectashield (Vector Laboratories).
728
The following primary antibodies were used: Rabbit anti-Ham (1:300) (Gift from Eric Lai's
lab), Ms anti-V5 (1:500, R960-25, invitrogen), chk anti-GFP (1:2000, ab13970, Abcam), Rat
anti-vasa (1:50, anti-vasa-c, Developmental Studies Hybridoma Bank, DSHB), Ms anti-ABD-
B (1:50, 1A2E9-c), Ms anti-SXL (1:50, M18-c), rat anti-DE-Cad (1:50, DCAD2, DSHB),
mouse anti-Crb (1:10, Cq4 DSHB), mouse anti-Coracle (1:100, C615.16 DSHB). Secondary
antibodies conjugated to Cy3 or Cy5 or Alexa Fluor-488 (Jackson ImmunoResearch) were
used.

Quantification of signal intensity from fluorescent immunostaining

For E-Cad and Crb in Figure 2, "raw intensity density" from manually defined regions (using free-hand drawing) along the apical side of the SV and TE terminal epitheliums was obtained in ImageJ. For each sample, 2 regions in the SV and 3 regions in the TE terminal epithelium were averaged. The regions used to determine relative E-Cad fluorescence in *hamSK1* TE somatic clones were at the apical side between cell junction. The length of Cor-stained cell junction was measured using ImageJ. A straight line was drawn over the Cor staining with the 'Freehand Line tool'. For each sample, the average length from 10 cell-cell junctions were used. In all experiments, images for the genotypes in comparison were always processed using exact settings. At least three biological replicates were included in statistical analysis.

Western Blot

Embryos from *w1118, ham$^{PR\Delta}$/CTG, ham^{SK1}/CTG* were collected and aged to embryonic stage S13 to S16. In each blot, 50-100st GFP+ or GFP- embryos were used. The selected embryos were homogenized directly in 1xSDS loading buffer (62.5 mM Tris-HCl pH 6.8, 2.5 % SDS, 0.002 % Bromophenol Blue, 0.7135 M (5%) β-mercaptoethanol, 10 % glycerol) and denatured at 98 °C heat blot for 5 min. After centrifuged, supernatant was taken for Western-Blotting with 8% of SDS-PAGE. Rabbit anti-Ham (1:2000) primary antibody was used in blotting. To minimize unspecific signal from IgG heavy chains, light chain specific secondary antibody of mouse anti-Rabbit IgG (1:40000 Jackson ImmunoResearch) was used, and the signals were developed with ECL Plus reagent (GE Healthcare).

Cell culture and Luciferase assays

To generate different variants of Ham, the open reading frames of HamFL, Ham-NZF-vp16, Ham-CZF-vp16, and *ham$^{PR\Delta}$* were PCR amplified and cloned into the pAC_V5 (nitrogen) vector. The enhancer fragments were cloned in to the p2xTetO-Firefly luciferase vector. All transfections were performed using *Drosophila* S2-R+ cells grown in Schneider *Drosophila* medium containing 10% fetal calf serum. Cells were co-transfected with the Ham expression construct, 2xTetO-Firefly luciferase and pAc-Renilla plasmids in 96-well plate using the Effectene Transfection kit (Qiagene). Luciferase assays were performed and measured using the Dual Luciferase Assay System (Promega). Expression was calculated as the ratio between

the firefly and *Renilla* luciferase activities. Three rounds of transfections were performed on
different days and different cell populations and considered as three biological replicates.
Within each biological replicate, 4 wells of cells were transfected with the same DNA mixture
and measured and thus considered as 4 technique replicates. In total, for each DNA
combination, the values from the 12 wells of transfection were calculated, averaged and plotted
in **Fig. 4F-G** and **Supplementary Fig. 6D**. Cloning oligos are provided in **Supplementary**
table 4.

SCRINSHOT

The Drosophila reproductive system was dissected at defined ages in 1xPBS and fixed in 4%
paraformaldehyde in PBS for 20min. After washing with PBST 0.1% (Triton X-100) for 2 x
15 min, fine dissected samples were placed on Poly-lysine coverslip and freezed at -80°C after
drying on a heater.

The SCRINSHOT experiments were carried out according to the published method [55]. In
brief, 3 padlock probes and 3 corresponding detection probes were designed for each gene of
interest. The slides with fine dissected samples were pretreated at 45°C to reduce moisture and
fixed in 4% PFA in 1X PBS, followed by washing in PBS Tween-20 0.05% twice. Samples
were then dehydrated with 70%, 85%, 100%ETOH and air dry. The SecureSeal hybridization
chamber (GRACE BIO-LABS, 621501) was then mounted to cover samples on the slide.
Permeabilization of tissues were first done by treating with 10µg/mL Proteinase K (Invitrogen)
in PBSTween-20 0.05% for 7 minutes followed by washing with Glycine (2mg/mL in PBS
Tween-20 0.05%) once and with PBS Tween-20 0.05% for 3 x 5 minutes. After this, samples
were further Permeabilized in 0.1M HCl for 3min, followed by washing in PBS Tween-20
0.05% twice. Samples were then blocked in a probe-free hybridization reaction mixture of
1XAmplifase buffer (Lucigen, A1905B), 0.05M KCl, 20% Formamide deionized (Millipore
S4117), 0.1uM Oligo-dT, 0.1ug/ul BSA (New England Biolabs, B9000S), 1U/ul RiboLock
(Thermo, EO0384), and 0.2ug/ul tRNAs (Ambion, AM7119). Then hybridization of padlock
probes was done by incubating samples with padlock probes (with the concentration of each
one 0.01uM) mixed in blocking reagents used before (no oligo dT used in this step). The slide
was then put into a PCR machine to denature at 55˚C for 15 minutes and hybridize at 45˚C for
120 minutes. Padlock probes were then ligated by using SplintR ligase (NEB M0375) at 25˚C
for 16 hours followed by RCA (rolling cycle amplification) at 30˚C for 16 hours by using phi29
polymerase (Lucigen, 30221–2) and RCA primer1. Then a fixation step was applied to stabilize
RCA product in 4% PFA for 15 minutes followed by washing in PBS Tween-20 0.05%. Then

the hybridization of the first 3 genes was done by mixing all 3 3' fluorophore-conjugated

detection probes of each gene in reaction reagent (2XSSC, 20% Formamide deionized, 0.1ug/ul

BSA, and 0.5 ng/µl DAPI (Biolegend, 422801) followed by hybridization at 30˚C for 1 hour.

The slides were washed in 20% formamide in 2X SSC and then in 6X SSC, followed by

dehydration in 70%, 85%, 100% Ethanol until the chamber was removed. Then the samples

were preserved in SlowFade Gold Antifade mountnant (Thermo, S36936) and kept in dark

before imaging. After image acquisition, the first 3 detection probes were removed by using

Uracil-DNA Glycosylase (Thermo, EN0362), and the slides were ready for the next round of

detection probe hybridization. The procedure was repeated until all genes were hybridized and

imaged. Images were acquired with an airyscan-equipped confocal microscope system (Zeiss

LSM 800, Carl Zeiss), and ZEN 2.5 Blue edition software was used for image processing.

Quantification of SCRINSHOT signals

Similar to quantification of immunofluorescent signal, fluorescence intensity of SCRINSHOT

for each gene was measured in defined areas within images of the SV, EjD, and TE in 2D

projections. Squares were drawn to define areas at the leading side of the SV, the middle region

of the Ejd and the TE terminal. The total fluorescence intensity was first normalized to area

size. The resultant value for *ham* mutant or RNAi samples was then contrasted to the

normalized intensity of control samples.

CUT&TAG

CUT&TAG was performed according to the published method [38]. In brief, TE and grossly

tissue of the posterior abdomen were dissected from 18-25h APF *w1118* and

pBAC_ham_sfGFP animals in PBS within 30 min. For grossly tissues, the dissected samples

were homogenized in 2ml NE1 buffer followed by incubation on ice for 10 min. After filtering

out the grossly cuticles, nuclei were collected by spinning at 1300xg and resuspended in 1ml

PBS. For TE, nuclei extraction was done by simply pipetted samples in 200ul NE1 up and

down for 20-30 times and incubated on ice for 10 min. Light fixation was performed with 0.1%

formaldehyde at RT for 2 minutes and neutralized with 75mM Glycine. Then the nuclei were

washed by PBS and wash buffer 1 time each and kept in 200ul wash buffer on ice until enough

samples were collected (30-50st animals per/antibody). Resuspended nuclei in 1.2ml wash

buffer and mixed with 20ul Concanavalin A-coated magnetic beads (Polyscience 86057) with

soft vortexing. Aliquoted samples into two parts immediately and incubated for 10 min with

rotation to allow the binding of the nuclei with the beads. The nuclei/beads mix was then

blocked with 1mL cold antibody buffer for 10 min and resuspended in 150ul cold antibody

buffer. For w1118 samples, 1uL Rabbit anti-Ham or 1ul Rabbit IgG (SIGMA-ALDRICH,

I5006) was added for an overnight incubation at 4°C. For *pBAC_ham_sfGFP* samples, 1ul

Rabbit anti-GFP (G10362, invitrogen) or Rabbit IgG was used. The beads/nuclei/antibody mix

was then washed with Dig-wash buffer and incubated with secondary antibody (1:100 dilution)

for 1 hour. After further washes with the Dig-wash buffer, a pA-Tn5 adaptor complex in Dig-

300 buffer was added to the beads for 1 hour reaction. The beads were then washed with Dig-

300 buffer before the incubation with 300ul Tagmentation buffer at 37°C for 1 hour. Then 10ul

0.5M EDTA, 3uL 10% SDS and 2.5uL 20 mg/mL proteinase K (Invitrogen) were used to stop

the reaction. Then the resultant DNA was purified with DNA clean & Concentrator kit (ZYMO

Research D4013) and eluted in 25uL Elution buffer. To generate libraries, 21uL fragment DNA

was mixed with 2uL 10uM Universal i5 primer, 2uL 10uM uniquely barcoded i7 primer and

25uL PCR master mix (NEB Phusion® High-Fidelity PCR Master Mix with HF Buffer,

M0531). The PCR condition was as follows: 72°C 5min, 98°C 30s, repeat 12 times (98°C 10s,

63°C 10s), 72°C 1 min and hold 4°C. The libraries were cleaned with standard Ampure XP

beads as previously described. Libraries from four biological replicates were produced for each

condition.

We mapped the CUT&TAG samples of *w1118* and *pBAC_ham_sfGFP* in TE and TG tissues

to the dm6 genome assembly using Bowtie2 with the following parameters: bowtie2 --end-to-

end --very-sensitive --no-mixed --no-discordant --phred33 -I 10 -X 700. The coverage for these

samples was generated using bedtools genomeCoverageBed and normalized to library depth to

yield reads per million per base (RPM). For peak calling, we employed SEACR [67], which is

tailored for sparse chromatin profiling data like CUT&TAG. We first used a relaxed FDR

cutoff of < 0.5 for each replicate (SEACR_1.3.sh normalized_coverage 0.5 norm stringent

output_peaks), we then identified common peaks across replicates for each condition. To select

confident peaks for *w1118* and *pBAC_ham_sfGFP*, we overlapped their peaks within TE and

TG separately, choosing the top 1800 peaks from TE and the top 6000 from TG. Peak

annotation was performed by HOMER's annotatePeaks.pl. Peak overlap analysis was

performed using HOMER's mergePeaks function with the default -d given parameter, ensuring

that only directly overlapping peaks were merged. This method preserved the integrity of the

original peak boundaries and maintained the specified distances between peaks from the initial

peak calls. *De novo* motif discovery for peaks in up-regulated or down-regulated genes,

positioned 2000 bp upstream and 500 bp downstream of the TSS in TE and TG tissues, was

performed using the MEME-ChIP software. The parameters used were: -ccut 100, -meme-p 5,

-dna, -meme-mod anr, -minw 5, -maxw 15, and -filter-thresh 0.05.

RNA-seq

Total RNA was extracted from finely dissected genital discs or testis samples from 18-25h APF

animals using Trizol reagent (Invitrogen). RNA quality was tested by Agilent Bioanalyzer.

RNA-seq libraries were made using the Illumina Truseq Total RNA library Prep Kit LT.

Sequencing was performed on the Illumina Hiseq2500 platform. Following the trimming of

adapter sequences with Trimmomatic, RNA-seq reads from replicated wild type (x3) and

mutant samples (x3) were aligned to the *Drosophila melanogaster* (dm6) genome assembly

utilizing HISAT2. Gene annotations were sourced from UCSC. The DESeq2 Bioconductor

package was employed to identify differentially expressed mRNAs between *w1118* and the

$ham^{PR\Delta}/ham^{PR\Delta}$ mutant, as well as between +/Df and ham^{SK1}/Df mutant, with a subsequent

estimation of the false discovery rate (FDR). Genes with an FDR < 10% and absolute

Log2FC>0 were considered differentially expressed. Details of the calculated differential

expression are provided in **Supplementary table 5**.

STRING gene network analyses

Ham activated (genes downregulated in *hamSK1* and bound by Ham in TG) and repressed

(genes upregulated in *hamSK1* and bound by Ham in TG) gene networks are constructed using

String, MCL clustering (find natural clusters the stochastic flow, inflation parameter, 2.

Disconnected genes are hidden. The position of the clusters is manually adjusted to fit related

GO categories. The links to the gene networks in STRING:

https://string-db.org/cgi/network?taskId=bbF8IgMTqf1M&sessionId=bzDIAZvMgWsi

https://string-db.org/cgi/network?taskId=bhKCcMoLlbwh&sessionId=bzDIAZvMgWsi

Data availability: All sequencing data produced in this study has been submitted to Gene

Expression Omnibus (accession number: GSE290310).

References

1.   Bernascone, I., M. Hachimi, and F. Martin-Belmonte, *Signaling Networks in Epithelial*
Tube Formation. Cold Spring Harb Perspect Biol, 2017. **9**(12).
2.   Chung, S. and D.J. Andrew, *The formation of epithelial tubes.* J Cell Sci, 2008. **121**(Pt
21): p. 3501-4.
3.   Samakovlis, C., et al., *Development of the Drosophila tracheal system occurs by a*
series of morphologically distinct but genetically coupled branching events.
Development, 1996. **122**(5): p. 1395-407.
4.   Samakovlis, C., et al., *Genetic control of epithelial tube fusion during Drosophila*
tracheal development. Development, 1996. **122**(11): p. 3531-6.
5.   Sutherland, D., C. Samakovlis, and M.A. Krasnow, *branchless encodes a Drosophila*
FGF homolog that controls tracheal cell migration and the pattern of branching. Cell,
1996. **87**(6): p. 1091-101.
6.   Tanaka-Matakatsu, M., et al., *Cadherin-mediated cell adhesion and cell motility in*
Drosophila trachea regulated by the transcription factor Escargot. Development,
1996. **122**(12): p. 3697-705.
7.   Gervais, L., G. Lebreton, and J. Casanova, *The making of a fusion branch in the*
Drosophila trachea. Dev Biol, 2012. **362**(2): p. 187-93.
8.   Jacinto, A., A. Martinez-Arias, and P. Martin, *Mechanisms of epithelial fusion and*
repair. Nat Cell Biol, 2001. **3**(5): p. E117-23.
9.   Vasioukhin, V., et al., *Directed actin polymerization is the driving force for epithelial*
cell-cell adhesion. Cell, 2000. **100**(2): p. 209-19.
10.   Pai, Y.J., et al., *Epithelial fusion during neural tube morphogenesis.* Birth Defects Res
A Clin Mol Teratol, 2012. **94**(10): p. 817-23.
11.   Caviglia, S. and S. Luschnig, *Tube fusion: making connections in branched tubular*
networks. Semin Cell Dev Biol, 2014. **31**: p. 82-90.
12.   de Mello Santos, T. and B.T. Hinton, *We, the developing rete testis, efferent ducts,*
and Wolffian duct, all hereby agree that we need to connect. Andrology, 2019. **7**(5):
p. 581-587.
13.   Jemc, J.C., *Somatic gonadal cells: the supporting cast for the germline.* Genesis, 2011.
49(10): p. 753-75.
14.   Fog, C.K., G.G. Galli, and A.H. Lund, *PRDM proteins: important players in*
differentiation and disease. Bioessays, 2012. **34**(1): p. 50-60.
15.   Chi, J. and P. Cohen, *The Multifaceted Roles of PRDM16: Adipose Biology and*
Beyond. Trends Endocrinol Metab, 2016. **27**(1): p. 11-23.
16.   Bard-Chapeau, E.A., et al., *Ecotopic viral integration site 1 (EVI) regulates multiple*
cellular processes important for cancer and is a synergistic partner for FOS protein in
invasive tumors. Proc Natl Acad Sci U S A, 2012. **109**(6): p. 2168-73.
17.   Zhou, B., et al., *PRDM16 Suppresses MLL1r Leukemia via Intrinsic Histone*
Methyltransferase Activity. Mol Cell, 2016.
18.   Corrigan, D.J., et al., *PRDM16 isoforms differentially regulate normal and leukemic*
hematopoiesis and inflammatory gene signature. J Clin Invest, 2018. **128**(8): p. 3250-
3264.

948	19.	Cibi, D.M., et al., *Prdm16 Deficiency Leads to Age-Dependent Cardiac Hypertrophy, Adverse Remodeling, Mitochondrial Dysfunction, and Heart Failure.* Cell Rep, 2020. **33**(3): p. 108288.
951	20.	Kundu, A., et al., *PRDM16 suppresses HIF-targeted gene expression in kidney cancer.* J Exp Med, 2020. **217**(6).
953	21.	Horvath, C. and C. Scheele, *PRDM16 stability and metabolically healthy adipose tissue.* Nat Metab, 2022. **4**(9): p. 1095-1096.
955	22.	Wang, Q., et al., *Post-translational control of beige fat biogenesis by PRDM16 stabilization.* Nature, 2022. **609**(7925): p. 151-158.
957	23.	Hurwitz, E., et al., *Antagonism between Prdm16 and Smad4 specifies the trajectory and progression of pancreatic cancer.* J Cell Biol, 2023. **222**(4).
959	24.	Moore, A.W., L.Y. Jan, and Y.N. Jan, *hamlet, a binary genetic switch between single- and multiple- dendrite neuron morphology.* Science, 2002. **297**(5585): p. 1355-8.
961	25.	Moore, A.W., et al., *Conversion of neurons and glia to external-cell fates in the external sensory organs of Drosophila hamlet mutants by a cousin-cousin cell-type respecification.* Genes Dev, 2004. **18**(6): p. 623-8.
964	26.	Rives-Quinto, N., et al., *Sequential activation of transcriptional repressors promotes progenitor commitment by silencing stem cell identity genes.* Elife, 2020. **9**.
966	27.	Eroglu, E., et al., *SWI/SNF complex prevents lineage reversion and induces temporal patterning in neural stem cells.* Cell, 2014. **156**(6): p. 1259-73.
968	28.	Endo, K., et al., *Chromatin modification of Notch targets in olfactory receptor neuron diversification.* Nat Neurosci, 2012. **15**(2): p. 224-33.
970	29.	Pinheiro, I., et al., *Prdm3 and Prdm16 are H3K9me1 methyltransferases required for mammalian heterochromatin integrity.* Cell, 2012. **150**(5): p. 948-60.
972	30.	Hohenauer, T. and A.W. Moore, *The Prdm family: expanding roles in stem cells and development.* Development, 2012. **139**(13): p. 2267-82.
974	31.	Gokcezade, J., G. Sienski, and P. Duchek, *Efficient CRISPR/Cas9 plasmids for rapid and versatile genome editing in Drosophila.* G3 (Bethesda), 2014. **4**(11): p. 2279-82.
976	32.	DeFalco, T., S. Le Bras, and M. Van Doren, *Abdominal-B is essential for proper sexually dimorphic development of the Drosophila gonad.* Mech Dev, 2004. **121**(11): p. 1323-33.
979	33.	Whitworth, C., E. Jimenez, and M. Van Doren, *Development of sexual dimorphism in the Drosophila testis.* Spermatogenesis, 2012. **2**(3): p. 129-136.
981	34.	Ahmad, S.M. and B.S. Baker, *Sex-specific deployment of FGF signaling in Drosophila recruits mesodermal cells into the male genital imaginal disc.* Cell, 2002. **109**(5): p. 651-61.
984	35.	Rothenbusch-Fender, S., et al., *Myotube migration to cover and shape the testis of Drosophila depends on Heartless, Cadherin/Catenin, and myosin II.* Biol Open, 2017. **6**(12): p. 1876-1888.
987	36.	Bischoff, M.C., et al., *Filopodia-based contact stimulation of cell migration drives tissue morphogenesis.* Nature Communications, 2021. **12**(1): p. 791.
989	37.	Jenett, A., et al., *A GAL4-driver line resource for Drosophila neurobiology.* Cell Rep, 2012. **2**(4): p. 991-1001.
991	38.	Kaya-Okur, H.S., et al., *CUT&Tag for efficient epigenomic profiling of small samples and single cells.* Nat Commun, 2019. **10**(1): p. 1930.

39. Szklarczyk, D., et al., *The STRING database in 2023: protein-protein association networks and functional enrichment analyses for any sequenced genome of interest.* Nucleic Acids Res, 2023. **51**(D1): p. D638-D646.

40. Kozopas, K.M., C.H. Samos, and R. Nusse, *DWnt-2, a Drosophila Wnt gene required for the development of the male reproductive tract, specifies a sexually dimorphic cell fate.* Genes Dev, 1998. **12**(8): p. 1155-65.

41. Deshpande, G., A. Nouri, and P. Schedl, *Wnt Signaling in Sexual Dimorphism.* Genetics, 2016. **202**(2): p. 661-73.

42. van Impel, A., et al., *Regulation of the Rac GTPase pathway by the multifunctional Rho GEF Pebble is essential for mesoderm migration in the Drosophila gastrula.* Development, 2009. **136**(5): p. 813-22.

43. Oda, H., et al., *A Drosophila homolog of cadherin associated with armadillo and essential for embryonic cell-cell adhesion.* Dev Biol, 1994. **165**(2): p. 716-26.

44. Ferrante, A.W., Jr., R. Reinke, and E.R. Stanley, *Shark, a Src homology 2, ankyrin repeat, tyrosine kinase, is expressed on the apical surfaces of ectodermal epithelia.* Proc Natl Acad Sci U S A, 1995. **92**(6): p. 1911-5.

45. Gonzalez-Gaitan, M. and H. Jackle, *Role of Drosophila alpha-adaptin in presynaptic vesicle recycling.* Cell, 1997. **88**(6): p. 767-76.

46. Weber, A.N., et al., *Binding of the Drosophila cytokine Spatzle to Toll is direct and establishes signaling.* Nat Immunol, 2003. **4**(8): p. 794-800.

47. Staubli, F., et al., *Molecular identification of the insect adipokinetic hormone receptors.* Proc Natl Acad Sci U S A, 2002. **99**(6): p. 3446-51.

48. Kawamura, K., et al., *A new family of growth factors produced by the fat body and active on Drosophila imaginal disc cells.* Development, 1999. **126**(2): p. 211-9.

49. Lam, G.T., C. Jiang, and C.S. Thummel, *Coordination of larval and prepupal gene expression by the DHR3 orphan receptor during Drosophila metamorphosis.* Development, 1997. **124**(9): p. 1757-69.

50. Prokopenko, S.N., et al., *A putative exchange factor for Rho1 GTPase is required for initiation of cytokinesis in Drosophila.* Genes Dev, 1999. **13**(17): p. 2301-14.

51. Nakamura, M., J.M. Verboon, and S.M. Parkhurst, *Prepatterning by RhoGEFs governs Rho GTPase spatiotemporal dynamics during wound repair.* J Cell Biol, 2017. **216**(12): p. 3959-3969.

52. Lin, Y.H., et al., *AP-2-complex-mediated endocytosis of Drosophila Crumbs regulates polarity by antagonizing Stardust.* J Cell Sci, 2015. **128**(24): p. 4538-49.

53. Campbell, K. and J. Casanova, *A role for E-cadherin in ensuring cohesive migration of a heterogeneous population of non-epithelial cells.* Nat Commun, 2015. **6**: p. 7998.

54. Chen, H., et al., *E-cadherin mediates adhesion and suppresses cell motility via distinct mechanisms.* J Cell Sci, 1997. **110 (Pt 3)**: p. 345-56.

55. Sountoulidis, A., et al., *SCRINSHOT enables spatial mapping of cell states in tissue sections with single-cell resolution.* PLoS Biol, 2020. **18**(11): p. e3000675.

56. Lee, M., et al., *Distinct sites in E-cadherin regulate different steps in Drosophila tracheal tube fusion.* Development, 2003. **130**(24): p. 5989-99.

57. Lee, S. and P.A. Kolodziej, *The plakin Short Stop and the RhoA GTPase are required for E-cadherin-dependent apical surface remodeling during tracheal tube fusion.* Development, 2002. **129**(6): p. 1509-20.

58. Umetsu, D., *Cell mechanics and cell-cell recognition controls by Toll-like receptors in tissue morphogenesis and homeostasis.* Fly (Austin), 2022. **16**(1): p. 233-247.

59. Karlsson, M., et al., *A single-cell type transcriptomics map of human tissues.* Sci Adv,
2021. **7**(31).

60. Corty, M.M., B.J. Matthews, and W.B. Grueber, *Molecules and mechanisms of*
dendrite development in Drosophila. Development, 2009. **136**(7): p. 1049-61.

61. Araujo, S.J. and J. Casanova, *Sequoia establishes tip-cell number in Drosophila*
trachea by regulating FGF levels. J Cell Sci, 2011. **124**(Pt 14): p. 2335-40.

62. Brenman, J.E., et al., *Sequoia, a tramtrack-related zinc finger protein, functions as a*
pan-neural regulator for dendrite and axon morphogenesis in Drosophila. Dev Cell,
2001. **1**(5): p. 667-77.

63. Araujo, S.J., C. Cela, and M. Llimargas, *Tramtrack regulates different morphogenetic*
events during Drosophila tracheal development. Development, 2007. **134**(20): p.
3665-76.

64. Li, S., et al., *Photoreceptor cell differentiation requires regulated proteolysis of the*
transcriptional repressor Tramtrack. Cell, 1997. **90**(3): p. 469-78.

65. Pitsouli, C. and N. Perrimon, *The homeobox transcription factor cut coordinates*
patterning and growth during Drosophila airway remodeling. Sci Signal, 2013. **6**(263):
p. ra12.

66. Grueber, W.B., L.Y. Jan, and Y.N. Jan, *Different levels of the homeodomain protein cut*
regulate distinct dendrite branching patterns of Drosophila multidendritic neurons.
Cell, 2003. **112**(6): p. 805-18.

67. Meers, M.P., D. Tenenbaum, and S. Henikoff, *Peak calling by Sparse Enrichment*
Analysis for CUT&RUN chromatin profiling. Epigenetics Chromatin, 2019. **12**(1): p. 42.
1062

1063 **Declaration of the use of AI**. ChatGPT was used to improve readability of the text, and the

1064 content was thoroughly checked and edited by the authors to ensure accuracy.

 Figure legends

 Figure 1. *ham* is expressed and required for *Drosophila* male RS development

 (**A**). The adult male RS. TE, testis; SV, seminal vesicle; AG, accessory gland; EJD, ejaculatory

 duct. (**B**). White-field images of adult male RS, showing defective morphology in the *ham*

 mutants. (**C**). Quantification of male fertility. Ham_sfGFP with the *ham* genomic region was

 used to rescue the sterility. Statistical significance was calculated using unpaired t-test (***,

 p<0.001; **, p< 0.01; *, p<0.05; n.s., non-significant). (**D**). Images of the adult RS stained with

 Ham_V5 in red. DNA is stained with DAPI in blue. The blue and white dashed lines highlight

 the TE and SV respectively. (**E**). Illustration of the developing male RS at the larval stage. The

 testis (TE) terminal epithelial cells localize at the opposite side of the germline stem cell (GSC)

 niche. The genital disc (GD) contains mesenchymal cells (MC) that will develop into the SV

 and accessory gland (ag). (**F**). Images of the larval TE (top) and GD (bottom) stained with

 Ham_V5 in red or magenta, the reporter in lime-green (btlGal4 driving UAS-GFP) and DAPI

 in blue. AEL, after egg laying. (**G**). Illustration of the TE and SV fusion process. APF, after

 pupal formation. (**H**). Images of the pupal GD stained with Ham_V5 in magenta, the reporter

 in lime-green (btlGal4 driving UAS-GFP) and DAPI in blue. (**I**) Images of the pupal RS with

 Ham_V5 in red and DAPI in blue. The blue and white dashed lines highlight the TE and SV

 respectively. (**J**). Higher magnification image of the joint site between the SV and TE, with

 myoblasts and myotubes labeled by mef2Gal4>GFP in lime-green.

 Figure 2. Ham promotes differentiation of the TE terminal and SV epithelial cells

 (**A**). Images of the male RS at the pupal stage, showing morphological defects of the TE and

 SV in the ham mutants. Epithelial cells are labeled with Coracle (Cor). The phenotype

 penetrance (%) and number of animals (n) counted are indicated as % (n). (**B**). High

 magnification images of the connection site of the TE and SV, showing that the ham mutant

 RS failed to form a continuous tube. Ham-sfGFP restored the normal morphology. Epithelial

 cells are marked by E-Cad in red, Cor in yellow, and the epithelium on the SV side also marked

 by btlGal4 driving UAS-mCD8GFP. DNA is labeled by DAPI in blue. (**C**). Images and

 quantification of the E-Cad signal in the *wt* and *hamPRΔ* mutant SV. (**D**). Images and

 quantification of the Crumbs signal in the *wt* and *hamPRΔ /Df* mutant SV. (**E**). Images and

 quantification of the length of Cor signal in the wt and *hamPRΔ* mutant SV. (**F**) Images of *ham*

 mutant mosaic clones in the TE terminal epithelial cells. Note, the mutant cells are GFP

negative and within the area surrounded by the white dashed line. The white arrows indicate

junctions between mutant cells, while the yellow arrows highlight the junctions between wild-

type cells. (**G**). Images of pupal male GD, showing that *hamRNAi*-mediated knockdown

reproduces *ham* mutant phenotypes. (**H**). Images of pupal male RS, stained with E-Cad in red,

Ham in yellow, GFP in green and DAPI in blue. UAS-dcr2 enhanced knockdown efficiency

and led to more severe morphological defects. (**I**). Images and quantification of E-Cad signal

in red and Cor length in blue in control and ham RNAi SV. (**J**) Images and quantification of E-

Cad signal in control and ham RNAi TE terminal epithelium. Statistical significance was

calculated using unpaired t-test except for TE mosaic clones which used paired t-test. (***,

$p<0.001$; **, $p< 0.01$; *, $p<0.05$; ns, non-significant).

1124

1125   **Figure 3. Identification of Ham-regulated genes in the developing TE and GD**

1126

(**A**). Illustration of the sample types and antibodies used in the RNA-seq and CUT&TAG

experiments. (**B**) Venn diagram of dysregulated genes between the two mutant conditions in

the GD (left) and TE (right) samples. (**C**) Heatmaps of Ham CUT&TAG peak density and IgG

control, ranked from high to low and centered at peak summits. (**D**) Venn diagrams of

CUT&TAG peaks from the two types of samples. (**E-F**). Volcano plots of dysregulated genes

in GD (E) and TE (F) in the *hamSK1/Df* mutant, and the ones with at least one Ham CUT&TAG

peak are highlighted.

1134

1135   **Figure 4. Ham activates expression of epithelial differentiation genes**

1136

(**A**) Ham gene network analysis from STRING. The position of MCL Clusters were manually

adjusted to place genes with similar function in close proximity. The functional categories were

determined by gene function tools in STRING. (**B**). A bar plot of the Log2 fold-change value

from RNA-seq for *wnt2*, *shot* and *shg*, showing downregulation of these genes in the

hamSK1/Df mutant. (**C**). Schemes of the Ham_FL and other truncation proteins, and illustration

of the luciferase reporter assay. (**D-E**). A genome browser snapshot of the *shg* and *wnt2* loci,

showing the CUT&TAG tracks from the TE and GD (TG) mixed samples and the enhancer

fragments in the luciferase assays. (**F-G**). Normalized luciferase activity to the empty reporter

and then to the control vector that expresses GFP and RFP proteins. Statistical significance was

calculated using unpaired t-test (****, $p<0.0001$; ***, $p<0.001$; **, $p< 0.01$; *, $p<0.05$; ns,

non-significant).

1148

Figure 5. A candidate RNAi screen identified Ham-downstream effectors in RS epithelial tissue fusion

(**A**). Scheme for the RNAi screen. (**B**) Summary of positive hits. The extent of sterility is determined by the number of progenies from the F1 male crossing to the wild-type female (see Methods). 100% means no progeny. >70%, the number of progenies being fewer than 30% of the number of progenies produced by the control cross (the same Gal4 driver x UAS-GFPRNAi). 40-70%, the number of progenies being between 30-60% of the number of progenies produced by the control. (**C**). Images of the control and gene-specific RNAi knockdown RS. The left panels show the fusion site stained with E-Cad in magenta, GFP in lime-green, Ham in grey and DAPI in blue, the middle panels are GD, and the right ones are TE. The orange arrows indicate mobilized epithelial cells in the *shg* knockdown TE.

Figure 6. Spatial-temporal expression dynamics of Ham-downstream genes

(**A**). Images from multiplexed in situ hybridization (the SCRINSHOT method), showing mRNA signal of the indicated genes in *wt* TE samples. DNA is marked by DAPI in grey. (**B**). Images from SCRINSHOT, showing mRNA signals of indicated genes in *wt/Df* and *hamSK1/Df* mutant GD. The last panel is an enlarged view from the sample with *Tl*, *ham* and *spz* probes. The yellow arrowheads indicate myoblast cells with Tl signal. (**C**). Quantification of signal intensity for genes shown in B in two areas, the SV and EjD. Statistical significance was calculated using unpaired t-test (***, p<0.001; **, p< 0.01; *, p<0.05; ns, non-significant). (**D**) Images from SCRINSHOT, showing mRNA signals of indicated genes in *GFP* and *ham RNAi* GD samples. (**E**). Quantification of signal intensity for genes shown in D in two areas, the SV and EjD. (**F**). A summary of genes expressed in the TE terminal and SV epithelium. Highlighted genes show temporal or spatial dynamic expression in the two tissues. Statistical significance was calculated using unpaired t-test (****, p<0.0001; ***, p<0.001; **, p< 0.01; *, p<0.05; ns, non-significant).

Figure 7. Wnt2 and its ligand genes are required for male fertility and RS development

(**A**) Images of *wnt2o* mutant TE and GD, showing disrupted morphology in these tissues. (**B**). Images of the TE and SV fusion site from the control and *wnt2* overexpression animals. The epithelial cells of the TE and SV are labeled with hamRSGal4 driving UAS-mCD8GFP in

lime-green and E-Cad in magenta and Ham in grey. The white arrows indicate mobilized GFP

positive cells in the middle of the TE, and these cells have a lower level of E-Cad. The blue

and white dashed lines highlight the TE and SV respectively. (**C**). Images of the TE and SV

fusion site from the control and *fz2* RNAi animals. The white arrows indicate mobilized GFP

positive cells in the middle of the TE, and these cells have a lower level of E-Cad. (**D**) Graphs

of progeny number from male fertility tests for *fz* and *fz2* RNAi lines in comparison to GFP

RNAi. Number of tested males in each genotype, 3. (**E**) Illustration of cell-cell interactions in

the developing RS, and a speculated model for Tl and Wnt2 signaling functions. Wnt2

signaling may be involved in epithelial cell interaction between the two ends. Tl signaling and

Tl may be involved in the interaction between epithelial cells and myoblasts/myotubes. In three

conditions (*shgRNAi*, *wnt2OE* and *fz2RNAi*), some TE terminal epithelial cells (illustrated in

orange) move up to the distal end of the testis.

1194

1195

1196

1197

A

Repressed — ⊣ Ham → Activated

○ Signaling and patterning factors
○ Metabolism
○ Translational regulation
○ Cell division and mitosis
○ Chitin binding and synthesis
○ Transcriptional regulation

○ Cytoskeleton regulators
○ Metabolism and signaling
○ Epithelium and tube morphogenesis
○ Membrane and ECM components
○ Protein modification and processing
○ RNA-binding and processing
○ Axon guidance

B

hamSK1/Df vs wt/Df TE

Log2(FC)

wnt2 shot shg

C

Ham-PB 1 ■———————————————— 990
Ham-PB_PRΔ 171 ———————————————— 990
Ham-NZF_VP16 171 ———— 465
Ham-CZF_VP16 763 ———— 990

■ PR ■ ZF ■ PTZ00121 ∖CtBP interacting motif ■ VP16 AD

actin enhancer [Ham] → luc Luciferase reporter
● Ham or the variant proteins
■ Candidate enhancers

D

shg_enh2-1 5 kb
shg *cpa* *CG15653*
HamGFP_TG1_GFP 63.7
HamGFP_TG2_GFP 57.7
wt_TG2_Ham 32.3

E

wnt2_enh1-4 5 kb
wnt2
HamGFP_TG1_GFP 25.5
HamGFP_TG2_GFP 39.9
wt_TG2_Ham 35.2

F

Normalized luciferase activity

■ GFP-RFP
■ Ham_FL
■ Ham_PRdel
■ Ham_NZF_VP16
■ Ham_CZF_VP16

shg_enh1 enh2

G

Normalized luciferase activity

■ GFP-RFP
■ Ham_FL
■ Ham_PRdel
■ Ham_NZF_VP16
■ Ham_CZF_VP16

wnt2_enh1 enh2 enh3 enh4

A

hamRSGal4>UAS-dcr2; UAS-mCD8GFP

X

UAS-RNAi
(135 lines for 113 genes)

↓

Screen for F1 male sterility

↓

26 positive gene hits

B

Sterility	Genes
100%	*pbl, AP-2alpha, CG11406, shg*
>70%	*GstE11, CG7456, Gr64c, AkhR, Idgf4*
40%-70%	*CG17477, CG11069, CG17186, CG4872, AdsL, Tl, CG17283, decay, PH4alphaMP, Hr46, CG15117, gsb-n, shark, CG17018, cag, CG7484, CG31935*

C

A

wnt2O 48h APF TE DAPI Cor E-Cad

100% (8)

50um

wnt2O 24-30h APF GD DAPI Cor E-Cad

100% (4)

50um

B

32-48h APF

UASCD8GFP DAPI hamRSGal4>CD8GFP E-Cad Ham

SV

50um

UASWnt2 DAPI hamRSGal4>CD8GFP E-Cad Ham

100% (10)

SV

50um

C

40-50h APF TE DAPI hamRSGal4>CD8GFP E-Cad

GFP RNAi

20um

fz2 RNAi

60% (15)

D

Number of progenies

GFP.RNAi fz2.RNAi_1 fz2.RNAi_2 fz.RNAi

**

*

*

E

wt

TE

Wnt2/Fz/Fz2

SV

TI

TI/Spz/Spz5

TE

SV

wnt2OE fz2RNAi shgRNAi

TE

SV

↔ Epithelial remodeling and interaction
↔ Myoblast-myoblast interaction
→ Myoblast migration
↔ Epithelial-myoblast interaction

0) Mobilized epithelial cell
Myoblast and myocyte
Pigment cell

Supplementary figure legend

Suppl._Figure 1. Ham gene locus and mutant alleles.

(**A**) Illustration of the ham genomic locus and predicted protein isoforms, highlighting the mutation sites for the three mutant alleles and the protein domains of each isoform. All Ham protein isoforms maintain a functionally unknown motif PTZ00121 and the C-terminal zinc finger (ZF) clusters, while the long isoforms B/D and G also comprise the PR domain, the N-terminal ZF clusters. Isoform H has a truncated PR domain, and the shortest isoform E lacks the PR domain and the N-terminal ZF clusters. (**B**) Flanking sequences of the sgRNA target site from wt and four alleles with indels. Red letters indicate the sgRNA sequence. The corresponding amino acid sequence is shown above. "*" indicates the stop codon. (**C**). An image from western-blot, showing the two Ham protein bands in the control samples and absence of the Ham band(s) in the indicated mutants. "*" indicates an unspecific band, used as the loading control. (**D**) A table showing viability and sensory organ (SO) defects in four ham alleles. Note: the -6nt allele has a 6 nucleotide in-frame deletion. (**E**). Mechano-sensory organ phenotypes in the three indicated alleles. Red arrowheads indicate double sockets and double shafts phenotype.

Suppl._Figure 2. Ham expression and function in the female RS

(**A**). Bright-field images of the adult female RS from the indicated genotypes, showing accumulated eggs in the ham mutant ovaries. (**B**). Quantification of the number of progenies produced by wild-type and ham mutant females after crossed with wild-type males. Statistical significance was calculated using unpaired t-test (***, p<0.001; **, p< 0.01; *, p<0.05; ns, non-significant). (**C**). Images of pupal and adult ovaries and oviducts, showing high expression of the Ham_V5 protein in the connecting sites of the ovaries and oviducts.

Suppl._Figure 3. Split channel images for images in Fig. 2

Suppl._Figure 4. Ham controls the formation of TE epithelial cells

(**A**). Images of the testes from the early L3 larval stage, showing the absence of or mis-polarized TE terminal epithelial cell cluster in the *ham* mutant. E-Cad marks epithelial cells in green and DAPI marks the nuclei. The white arrows indicate mis-

polarized epithelial cells. (**B**). Illustration of the precursor cell clusters of the male RS in the embryonic stages. PGC, primordium germ cells; SGPs, somatic gonadal precursors; msSGP, male-specific SGPs. (**C-E**) Images of stage S13 and S15 *wt* embryos, showing Ham staining in all SGPs. Abd-B and Sox100B label msSGPs, VASA labels PGCs. (**F-G**). Images of the *wt* and *ham* mutant embryos at S13 (F) and S15 (G), showing the reduction or absence of msSGPs in the *hamSK1* mutant. Vasa is a germ cell marker, and Sox100B labels msSGPs in lime green.

Suppl._Figure 5. An epithelial-specific Gal4 line in the ham locus
(**A**). The *ham* gene locus, indicating the fragments in the Gal4 transgenes. The one (hamRSGal4) that drives reporter gene expression in the RS epithelial cells is highlighted in orange. (**B**). Images of pupal genital disc and TE in animals with hamRSGal4 and UAS-mCD8GFP. Ham_V5 is stained in magenta, and DAPI in blue. (**C**). Images showing the presence of myotubes on the surface of the testis. The bar graph on the right shows the relative abundance of myotubes on the testis in wt (100%) and ham mutant condition quantified from five animals each. (**D**). Images showing the junction between the TE and SV, and myotubes are present even in the *ham* mutant. The white arrows indicate myotubes with two or three nuclei at the junction.

Suppl._Figure 6. Ham regulatory activity and potential co-factors
(**A**). Principle component analysis of the RNA-seq datasets. (**B**). Principle component analysis of the Cut&Tag datasets. Note: some samples had poor sequencing depth, which were excluded from the PCA and later analyses. (**C**). The *shot* locus, showing the CUT&TAG tracks from the TE and GD (TG) mixed samples and the enhancer fragments in the luciferase assays. (**D**). Normalized luciferase activity to the empty reporter and then to the control vector that expresses GFP and RFP proteins. Statistical significance was calculated using unpaired t-test (****, $p<0.0001$; ***, $p<0.001$; **, $p< 0.01$; *, $p<0.05$; ns, non-significant). (**E**). Significantly enriched motifs from Ham binding peaks in up-regulated genes (left) and downregulated genes (right). The dashed rectangles highlight MAD and ZNF263 motifs.

Suppl._Figure 7. Split channel images for images in Fig. 5

Suppl._Figure 8. Split channel images for images in Fig. 5

Suppl._Figure 9. Expression of genes tested by SCRINSHOT in RNA-seq data

(**A**) Log2FC (hamSK1/Df versus wt/Df and *hamPRΔ* versus wt) in TE. (**B**) Log2FC(hamSK1/Df versus wt/Df and *hamPRΔ* versus wt) in GD.

Suppl._Figure 10. Split channel images for images in Fig. 6

Suppl._Figure 11. Additional genes tested in SCRINSHOT

(**A**). Images from SCRINSHOT, showing mRNA signal of the indicated genes in *wt* TE samples. DNA is marked by DAPI in grey. (**B**). Images from SCRINSHOT, showing mRNA signal of the indicated genes in *wt* TE samples at a later pupal stage (28-32 APF). DNA is marked by DAPI in grey. (**C**). Images from SCRINSHOT, showing mRNA signals of indicated genes in *wt/Df* and *hamSK1/Df* mutant GDs. (**D**). Images from SCRINSHOT, showing mRNA signals of indicated genes in *GFP* and *ham RNAi* GD samples. (**E-F**). Quantification of signal intensity for genes shown in C-D in two areas, the SV and EjD. (**G**). Quantification of signal intensity for genes that changed expression in ham RNAi TE. Statistical significance was calculated using unpaired t-test (***, p<0.001; **, p< 0.01; *, p<0.05; ns, non-significant).

Suppl._Figure 12-13. Split channel images for images in Suppl._Figure 11

Suppl._Figure 14. Male fertility test for Toll-9

(**A**). Graphs of progeny number from male fertility test for hamRSGal4 driving *Toll-9* RNAi in comparison to GFP RNAi. The number of tested males =3. (**B**). White-field image of Toll-9 RNAi adult male RS, showing morphological defects of the TE and the phenotype penetrance (11% of 37 animals). (**C**). Images of adult RS from control GFP RNAi andToll-9 RNAi males. hamRSGal4>UAS-mCD8GFP is used to label the testis terminal epithelium and the SV, and DAPI stains DNA. The panels below are higher magnification views on the selected SV area, highlighted with white dashed squares. White arrows indicate sperms in the control SV, which are absent from the Toll-RNAi sample. The bar graph shows the percentage of SV with sperms in GFP and Toll-9 RNAi animals (N=10).

Suppl._Figure 15. PRDM16 and MECOM/PRDM3 expression in human tissues

(A). RNA expression data across human tissues through Cap analysis of gene expression (Fantom5 dataset from the Human Protein Atlas). **(B).** Expression in Single cell RNA-seq clusters in the indicated human tissues (from the Human Protein Atlas).

A

hamRS-Gal4

Scale 10 kb

chr2L:18,760,000

---TATAGGCGAGG
---TAatt--GAGG

hamPRΔ

hamSK1

ham1

lncRNA:CR43804

lncRNA:CR44904

ham-B/NM_001259152
ham-D/NM_001273614
ham-E/NM_001273615
ham-H/NM_001273616
ham-G/NM_165262

Isoform B/D — 990
Isoform E — 633
Isoform H — 1068
Isoform G — 1108

215   573   913

- PR (red)
- ZF (yellow)
- PTZ00121 (blue)
- CtBP interacting motif

B

```
        I  R  S  V  Q  D  D  C  I  G  E  A  N  L  Q  S
wt     ATCCGAAGCGTTCAGGACGATTGTATAGGCGAGGCTAATTTACAGAGT

        I  R  S  V  Q  D  D  C  N  *
-5+3   ATCCGAAGCGTTCAGGACGATTGTAATT--GAGGCTAATTTACAGAGT

        I  R  S  V  Q  D  D  C  I  G           *
-5     ATCCGAAGCGTTCAGGACGATTGTATAGGC-----TAATTTACAGAGT

        I  R  S  V  Q  A              R  L  I  Y  R
-13    ATCCGAAGCGTTCAGG-------------CGAGGCTAATTTACAGAGT

        I  R  S  V  Q  D  D  T        E  A  N  L  Q  S
-6     ATCCGAAGCGTTCAGGACGATT------GCGAGGCTAATTTACAGAGT
```

C

Control | PRΔ

-5/+ | -6/+ | SK1 | -5+3 | -5 | -13 | -6

150 KD —

anti-Ham

*

D

ham	Trans-heterozygous ham/ham Df	
	Lethality	SO defects
-5nts	semi-lethal (60.6% survival)	0
-6nts	Viable (100% survival)	0
ham[1]	die upon hatch	100%
ham[sk1]	die upon hatch	100%

E

ham[ΔPR_iso] /ham Df   *ham[1]/ham Df*   *ham[sk1]/ham Df*

A

wt   *hamPRΔ/hamPRΔ*   *hamPRΔ/Df*

B   **C**

A Fig2B split channels

DAPI btlGal4>GFP E-Cad Cor

wt

TE

SV

10 um

hamPRΔ/Df

DAPI Ham E-Cad

hamPRΔ/Df;
pBAC(ham-sfGFP)

TE

SV

10 um

B Fig2F split channels

DAPI GFP E-Cad

FRThamSK1
/FRTGFP

TE

10um

C Fig2I split channels

Cor hamRSGal4>GFP E-Cad DAPI

GFP RNAi

SV

20um

ham RNAi

SV

D Fig2J split channels

DAPI hamRSGal4>GFP E-Cad

GFP RNAi

TE

ham RNAi

TE

DAPI hamRSGal4>CD8GFP E-Cad Ham

A **Fig6A split channels**

20-30h APF wt TE   ham shg DAPI

20-25h APF wt TE   DAPI fz2 fz3 wnt2

20-25h APF wt TE   DAPI spz5 spz spz4

B **Fig6B split channels**   fz2 fz3 wnt2   fz4 fz

wt /Df

hamSK1 /Df

DAPI ham spz Tl   DAPI ham spz Tl

wt /Df

hamSK1 /Df

C **Fig6D split channels**   DAPI shg wnt2   DAPI ham spz Tl

GFP RNAi

ham RNAi

Supplementary Fig. 11A split channels

Supplementary Fig. 11B split channels

Supplementary Fig. 11D split channels

A

B

11% (37)

C

hamRSGal4>UAS-mCD8GFP

GFP DAPI

TE
SV
200 um

TE
SV

SV

SV

50 um

GFP RNAi   *Toll-9 RNAi*

N=10

N=10