Drosophila Hamlet mediates epithelial tissue assembly of the reproductive system

Huazhen Wang[1], Ludivine Bertonnier-Brouty[2,3], Isabella Artner[2,3], Jiayu Wen[4,*]and Qi Dai[1,*]

1 Department of Molecular Bioscience, the Wenner-Gren Institute, Stockholm University,
Sweden

2 Lund Stem Cell Center, Lund University, Lund, Sweden

3 Lund University Diabetes Center, Lund University, Malmö, Sweden

4 Division of Genome Sciences and Cancer, The John Curtin School of Medical Research, The
Australian National University, and Australian Research Council Centre of Excellence for the
Mathematical Analysis of Cellular Systems.

[*] Co-corresponding authors

Abstract

Epithelial tissue fusion requires coordinated molecular events at the ends of two epithelial structures. Regulatory mechanisms controlling these events remain largely elusive. In the *Drosophila* reproductive system (RS), this fusion unites the gonad and the genital disc-derived tissues, into a continuous tube. This study unveils the pivotal role of Hamlet (Ham), a *Drosophila* PR-domain containing transcription factor, in orchestrating epithelial tissue fusion in the RS. Loss of *ham* leads to sterility and disconnection between the testes (TE) and seminal vesicles (SV). Systematic analysis of Ham-downstream genes reveals cytoskeletal, metabolic regulators and signaling pathway components. Ham activates genes for epithelial differentiation and remodeling, while repressing genes required for tissue growth and patterning. Using multiplexed in situ hybridization, we demonstrate spatial-temporal gene expression dynamics in contacting epithelia. Key Ham downstream effectors include E-Cadherin (E-Cad), Toll (Tl) and Wnt2 signaling pathways, regulating tissue interaction and fusion. Our findings present a comprehensive gene network crucial for heterotypic epithelial tissue fusion. Mammalian Ham orthologs PRDM3 and PRDM16 are highly expressed in epithelial tissues, suggesting a conserved role across species.

Introduction

Epithelial tissue fusion is a crucial process for building epithelial tubular network and uniting separate ends [1, 2]. Studies from different model systems, like neural tube fusion and *Drosophila* tracheal development, have provided several mechanistic insights into epithelial fusion process. For instance, specialized cells at the leading edge migrate and interact using cytoskeleton protrusions [3-9] and small GTPases like Rac and Cdc42 [10]. Upon contact, the leading cells re-establish polarity and connections with the help of extracellular matrix molecules, apical proteins and E-Cad [11]. However, the exact regulators and mechanisms controlling epithelial fusion remain to be determined, especially in a context-specific manner.

The reproductive system in animals comprises tissues from different origins, and must unite to form a connected system for producing and delivering gametes. For example, the human testis is directly linked to a complex tubular system consisting of the efferent and epididymal ducts [12]. Anomalies in the fusion of these ducts are a significant cause of cryptorchidism, affecting at least three percent of newborn boys. However, there is little understanding of the molecular mechanisms guiding the patterning and fusion of these ducts.

Similarly, the *Drosophila* male RS includes the gonad-derived testis and the genital disc (GD)-formed seminal vesicle, accessory gland (AG), ejaculatory duct (EjD) and additional supportive structures. During development, the TE and SV grow toward each other and merge into a continues tubular system [13]. This involves two epithelial ends, the terminal epithelial cells in the testis and the SV epithelium. While terminal epithelial cells in the testis are specified already in the embryo as male-specific somatic gonadal precursors (msSGPs) [14]. The SV epithelium, together with AG, develops from a group of mesenchymal cells in the GD which undergo mesenchymal-to-epithelial transition (MET) during the larval stage (**Fig. 1A**) [15, 16]. In the developing RS, multiple events, including epithelial tube formation and fusion as well as myoblast and pigment cell migration, are tightly coordinated (**Fig. 1B**) [17]. Thus, the developing *Drosophila* RS provides a representative *in vivo* context to dissect mechanisms underlying epithelial tissue assembly.

Ham belongs to the PR domain containing protein family whose members are evolutionarily conserved cell fate regulators [18, 19]. Dysfunction of the human orthologs of Ham, PRDM16 (Mel1) and PRDM3 (Evi1, MECOM), has been found in a variety of human abnormalities

98   including congenital diseases, metabolic disorders and cancers [20-27]. Moreover, in at least
99   two types of epithelial cancers (kidney and pancreas cancers), PRDM16 was shown to play a
100 tumor suppressive role [24, 27], indicating its importance in maintaining epithelial integrity.
101 However, it remains undermined how these PR factors regulate epithelial tissue development
102 and homeostasis.
103
Ham was exclusively studied in the nervous system where it is required for neuronal dendrite
morphogenesis [28] and cell fate maintenance [29-31]. Ham was shown to repress gene
expression through modulating chromatin states [32] or interacting with the co-repressor CtBP
[29, 32]. However, the molecular functions of PRDM16 and PRDM3 are versatile, with their
PR domains possessing intrinsic histone methyltransferase activity [21, 33] and the ability to
interact with various co-activators and co-repressors [18, 34]. It remains unknown whether
Ham plays a role outside of the nervous system and whether Ham also has dual function in
gene expression.
112
In this study, we discovered that Ham is a master regulator in guiding epithelial tissue
patterning and fusion in the *Drosophila* RS. Even a slight reduction of Ham disrupts epithelial
differentiation and tissue assembly in the RS, ultimately affecting sperm and egg delivery. To
explore new mechanisms governing epithelial tissue fusion in the RS, we conducted systematic
analyses using RNA-sequencing (RNA-seq), Cleavage Under Targets and Tagmentation
(CUT&TAG), multiplexed in situ hybridization data and genetic screening. These analyses led
to identification of gene regulatory networks required for epithelial tissue fusion. Ham activates
a gene network to promote epithelial cell differentiation and remodeling while suppressing
another gene network involved in cell specification and early patterning. These target genes
exhibit spatial-temporal expression dynamics during the fusion of the TE and SV. As a key
Ham downstream effector, Wnt2 signaling regulates epithelial cell communication and
interaction. By elucidating the regulatory network controlled by Ham, our study provides novel
insights into the molecular mechanisms underlying epithelial tissue fusion.
131 **Results**

Drosophila RS development is vulnerable to reduced Ham activity

While PRDM16 and MECOM are critical cell fate regulators in various developmental
contexts in mammals, the genetic roles of Ham were mainly found in the central and periphery
nervous systems [28-31]. To systematically study Ham function, we generated new frameshift
alleles using CRISPR/Cas9. The sgRNA targets the third exon shared by the three long
isoforms B, D and G (**Supplementary Fig. 1A**), producing new alleles specifically disrupting
the three PR-domain containing isoforms. We therefore named them *ham^PRΔ*. The original *ham^1*
and the null allele *ham^SK1* shifted open reading frame (ORF) for all predicted isoforms [28, 30].

We validated nucleotide deletions in the new alleles by sanger sequencing (**Supplementary
Fig. 1B**) and protein deletion by western-blot using an antibody against Ham. The Ham
antibody detected several bands in the blot (**Supplementary Fig. 1C**). The two bands at and
above the 150KD marker position in the two heterozygotic controls were not detectable in the
ham^SK1 homozygous mutant, suggesting that these two bands correspond to the longest and the
second longest isoforms respectively. The *ham^PRΔ* homozygous samples lack the largest band,
verifying the depletion of the longest isoforms. We used one of them (-5nt) for all the later
experiments.

We compared viability and sensory organ development of these alleles. In contrast to the
original report showing that *ham* mutant animals died at the larval stage [28], we found that
ham^SK1, *ham^PRΔ* and *ham^1* alleles all survived to hatch when combined in trans with the *ham*
deficiency line (Df) which deletes the entire *ham* locus (**Supplementary Fig. 1D**). While
ham^1/Df and *ham^SK1*/Df animals all died right after hatching, about two thirds of the *ham^PRΔ*/Df
mutants were viable without obvious gross defects. Furthermore, Ham is known to specify
internal cell types (the neuron and sheath) in the sensory organ lineage, as *ham^1* mutants
exhibited ectopic external cell types (the socket and shaft) with the expense of internal cells
[29]. *ham^PRΔ* mutants displayed normal sensory organs, in contrast to defective sensory organs
in the *ham^1* and *ham^SK1* alleles (**Supplementary Fig. 1E**). Thus, the PR-containing isoforms of
Ham are dispensable for sensory organ development and viability.

However, *ham^{PRΔ}* mutant animals showed impaired fertility and abnormal morphology in both

male and female RS (**Fig. 1D-E, Supplementary 2A-B**). The TE, AG and SV structures were

disrupted in males, while mature eggs were stuck in the mutant female ovaries, indicating failed

egg delivery. To confirm that the defects are induced by the *ham* mutation, we performed a

rescue experiment by utilizing a pBAC transgene that carries the genomic region of *ham* with

the super-folding GFP (sfGFP*)* coding sequence in frame with the *ham* ORF. The pBAC

transgene rescued the viability of all *ham* mutant conditions, and at least partly restored male

fertility of these mutants. Together, these results indicate that the *ham^{PRΔ}* allele represents a

hypomorphic condition which preserves Ham shorter isoforms and partial activity. Thus,

although viability and sensory organ development do not require the PR-containing isoforms,

formation of the RS depends on them.

Ham-expressing epithelial cells connect the gonadal and supportive tissues

To examine whether Ham directly controls RS development, we examined Ham expression in

the developing RS using a V5 antibody which detects all Ham isoforms in the *ham_V5* allele

[35]. Ham_V5 was detected in the connecting tube of the adult SV and TE (**Fig. 1F**).

Correspondingly, at the late L3 stage Ham_V5 was detected in the terminal epithelial cells of

the TE as well as the btl-Gal4-expressing mesenchymal cells in the genital disc which will give

rise to the SV and AG (**Fig. 1G-H**). At the pupal stage, a high level of Ham persists in the TE

terminal epithelial cells, SV and AG. In contrast, Ham is barely detectable in the SV myoblasts

labeled by the reporter, mef2-Gal4-driving UAS-GFP (**Fig. 1I-J**). Similarly, in the female, high

levels of Ham_V5 were found at the joint sites of oviducts and ovaries during both pupal and

adult stages (**Supplementary Fig 2C**). Thus, Ham is an intrinsic factor in epithelial cells at the

connection site between the gonadal- and genital-disc derived tissues in both males and

females.

Ham is required for epithelial tissue fusion in the developing male RS

The specific expression of Ham and mutant phenotypes in the RS prompted us to investigate

its function in RS epithelial tissues. Here we focused on the male, as the assembly of the TE

and SV represents a remarkable process involving a series of tissue reorganization events (**Fig.**

1A-C). When the two ends of TE and SV get close, the epithelial cells at the leading edges

meet and form a common epithelial tube. In the meanwhile, myoblasts from the SV and

pigment cells from the TE migrate toward opposite directions, eventually forming distinct cell

layers covering both structures.

We found that at 40-50h APF the TE and SV tubes in *ham^PRA/Df and hamSK1/Df* mutants did

not properly elongate and appeared shorter and wider compared to that in the wild-type (**Fig.**

2A). The SV and AG were even unseparated in the *ham* null allele *hamSK1/Df*, suggesting that

the severity of RS abnormality correlates with the levels of reduction in *ham* gene products.

To examine epithelial integrity, we performed immunofluorescent staining using the antibodies

against the epithelial cell markers, E-Cad (adherent junctions) and Coracle (Cor, septate

junctions equivalent to mammalian tight junction) protein. We also incorporated an SV-

specific transgenic reporter, btlGal4-driving UAS-EGFP or UAS-CD8GFP, to the control and

ham mutant genotypes. E-Cad is enriched at the apical side (the lumen side, **Fig. 2B**) of the TE

and SV epithelial tubes. Notably, the TE terminal epithelium accumulates a higher level of E-

Cad compared to the SV epithelium, while Cor is mainly expressed at the SV side but not much

in the TE, suggesting a heterotypic nature of the TE terminal and SV epithelia.

Control animals always showed connected TE and SV, whereas *ham* mutant TE and SV tissues

were often separated from each other. And even in some mutants where TE and SV tubes were

in close contact, they were disconnected (**Fig. 2B**). Importantly, this phenotype can be fully

rescued by the *pBAC_ham_sfGFP* transgene, confirming that it is caused by the *ham* mutation.

These observations suggest that the epithelial ends of the TE and SV possess distinct properties

despite both being epithelial cell types and that Ham is required for TE and SV fusion.

Depletion of Ham abolishes TE terminal epithelial cell formation

A close inspection on the pupal TE revealed that most of TE epithelial cells were missing in

ham mutant animals (**Fig. 2B**). TE terminal epithelial cells derive from male specific gonadal

precursors (msSGPs) specified in the early embryo (**Supplementary Fig. 3B**) [15] and expand

during larval and early pupal stages. *ham* mutant lava either lacked TE epithelial cells or

showed abnormal location of these cells, which can be rescued by the *pBAC_ham_sfGFP*

transgene (**Supplementary Fig. 3A**). This result indicates that Ham regulates specification or

survival of these cells. Indeed, we detected Ham expression in msSGPs and other SGPs in the

embryonic gonad. The number of msSGPs labeled by Abd-B or Sox100B-GFP was

dramatically reduced in S13 *ham* mutant embryo (**Supplementary Fig. 3C-G**). Thus, Ham is

a master regulator for the specification and/ or survival of msSGPs, and the absence of msSGPs

accounts for loss of TE terminal epithelial cells in larval and pupal stages.

Ham regulates patterning and cell differentiation of the RS epithelial tubes

The epithelial cells in the SV and AG originate from a group of mesenchymal cells recruited

by the FGF pathway involving Btl and Branchless (Bnl) (**Fig. 1A** and **1H**) [15, 16]. These cells

undergo MET to form an epithelial mass which then cavitates to generate inner hollow space

and eventually elongates into an epithelial tube. In comparison with the control, the signal

intensity of both E-Cad and Crumbs (Crb) were significantly reduced in *ham* mutant SV

epithelial cells, which coincides with the shortening of septate junctions labeled by Cor (**Fig.**

2C-E). This result suggests that Ham regulates epithelial cell differentiation in the SV. To

check whether Ham regulates epithelial cell differentiation in the TE, we used the FRT/FLP

system (see Methods) to generate mosaic somatic mutant clones. In these animals, *ham* mutant

cells were GFP negative, while wild-type and *ham* heterozygous cells expressed GFP (**Fig.**

2F). The E-Cad levels at the border between two mutant cells were significantly lower than

those between wild-type cells, indicating that *ham* depletion impaired differentiation of TE

terminal epithelial cells.

In summary, these phenotypic analyses suggest that Ham controls formation of TE

terminal epithelial cells in the early embryo and regulates epithelial tube elongation and cell

differentiation at the pupal stage. Thus, failed epithelium fusion in *ham* mutant RS could result

from: 1) absence of TE terminal epithelial cells, 2) defects in epithelial tube elongation and cell

differentiation of the SV and 3) a combination of both.

Ham directly controls epithelial fusion between the TE and SV

Next, we sought to assess whether and how Ham directly regulates the fusion process. We first

attempted to generate a mild mutant condition which can bypass defects of TE epithelial cell

loss and SV-AG tube elongation. We chose RNAi knockdown because knockdown efficiency

can be enhanced by overexpressing Dicer2 (Dcr2) and modulated by temperature when RNAi

is controlled by the GAL4/UAS system.

We first identified a Gal4 driver expressed in the SV and AG epithelial cells, after screening

the available Gal4 transgenes driven by a series of genomic fragments spanning the *ham* locus

[36] (**Supplementary Fig. 4A**). This transgene (thereafter named as *hamRSGAL4*) induces

high expression of *UAS-GFP* in the TE and SV epithelial cells and neglectable expression in

the myoblasts (**Supplementary Fig. 4B**). *hamRSGAL4,* when driving a *ham* RNAi line, nearly

abolished Ham expression in animals carrying the *UAS-dcr2* transgene (**Fig. 2G-H,** the middle

panels). These animals also had a reduced number of TE epithelial cells and displayed a ball-

like structure of the GD, similar to the *ham* null allele. Without *UAS-dcr2*, *ham* RNAi

knockdown reduced Ham expression but preserved TE epithelial cells and normal-appearing

SV. Remarkably, the TE and SV epithelia still failed to fuse (**Fig. 2H,** the right panel),

suggesting that Ham also controls the late step of fusion. Furthermore, E-Cad staining intensity

was significantly reduced in *ham* RNAi TE but not in *ham* RNAi SV (**Fig. 2I-J**), suggesting

that expression of E-Cad in the TE is sensitive to even mild reduction of Ham protein levels.

Ham controls distinct molecular pathways to promote tissue assembly.

To identify mechanisms that control epithelial fusion, we started to investigate Ham

downstream target gene network. We performed RNA-seq experiments from dissected TE and

SV tissues during the stage of 18-25h APF. We used two pairs of genotypes, including the

wild-type (*wt*), and *ham^{PRΔ}* homozygous and the *wt/Df* and *hamSK1/Df* trans-heterozygous

(**Fig. 3A**). The comparison between the first pair will determine dysregulated genes in the

ham^{PRΔ} hypomorphic condition, while the comparison between the second pair will identify

genes that change expression between the null and *ham* heterozygous conditions.

To not exclude small gene expression changes (*e.g.* in the leading cells which are the minority

within the whole tissue), we used a loose cutoff (FDR < 10% and FC>1). We found more than

two thousand up-regulated and down-regulated genes in the GD. In the mutant TE, there were

similar numbers of dysregulated genes (**Fig. 3B**). The genes that only changed expression in

one condition could result from unspecific genetic background, differential dependence on the

PR-containing isoforms or different sensitivity to *ham* heterozygous condition.

Expression of many dysregulated genes can be indirectly affected by Ham. We next intended

to identify genes that are transcriptionally regulated by Ham. To this end, we determined Ham

genomic binding sites using Cleavage Under Targets and Tagmentation assays (CUT&TAG)

[37] on finely-dissected TE and grossly-dissected abdominal tissues in which Ham is mainly

expressed in the TE and SV epithelium (**Fig. 3A**). We used a Ham antibody for wildtype

tissues, a GFP antibody for tissues from the *pBAC_ham_SFGFP* transgene and IgG as a

negative control in parallel for all the samples. Using the SEACR software, we identified peaks

and intersections of peaks across the Ham and GFP antibody conditions (see Methods). For

subsequent analyses, only the common 3433 and 691 peaks identified by the two antibodies

(Ham and GFP) were included to ensure signal specificity (**Fig. 3C-D**).

To determine Ham target genes, we intersected the RNA-seq data from the hamSK1/Df versus

wt/Df comparison with the Cut&Tag data. Among the 2,546 up-regulated genes in the GD, 441

(17.3%) contained Ham CUT&TAG peaks. Among the 2379 downregulated genes, 569

(23.9%) showed Ham binding (**Fig. 3E**). In the TE, a lower fraction of dysregulated genes

contains Ham CUT&TAG peaks, including 4.7% (108/2274) upregulated genes and 7%

(144/2057) downregulated genes (**Fig. 3F**), suggesting that dysregulated genes in the mutant

TE are mainly indirectly controlled by Ham.

Both up- and down-regulated genes contain Ham binding peaks, suggesting that Ham can

repress and activate gene expression. To systematically explore the function of Ham-regulated

genes, we performed gene network analysis using STRING [38]. We focused on the genes

bound by Ham and also dysregulated in the genital disc (GD) in the null condition *hamSK1*,

considering that most of TE genes are only indirectly influenced in the *ham^{SK1}* mutant. The

SRING MSL clustering analysis generated 114 functional clusters from down-regulated genes

and 62 clusters from up-regulated genes (**Fig. 4A, Supplementary Table 1**). Interestingly,

many of downregulated genes clustered into several large groups including cytoskeleton

regulators, metabolic factors, cell adhesion and extracellular matrix proteins, suggesting that

Ham activates genes to promote epithelial differentiation and remodeling. In contrast,

upregulated genes are enriched for patterning factors, cell cycle genes and transcriptional and

translational regulators, suggesting that Ham suppresses early developmental processes.

Notably, metabolic factors are enriched in both categories, but these factors belong to distinct

functional groups. Thus, epithelial fusion requires complex and balanced metabolic processes.

Together, this network analysis confirms that Ham positions on top of distinct molecular pathways to facilitate TE and SV assembly.

Ham was known for its repressive activity via interacting with the co-repressor CtBP and influencing histone modifications [32]. Our finding of its activation function was unexpected. To validate the activation activity of Ham, we chose three downregulated target genes (**Fig. 4B**), *dwnt2*, *shot* and *shg*, for transcriptional reporter assays. *shg* encodes E-cad, while Shot is a member of the spectraplakin family of large cytoskeletal linker molecules simultaneously binding to actin and microtubules. The function of *wnt2* was reported in the developing RS development [39, 40]. We cloned four genomic fragments from the *wnt2* locus, two each from *shg* and *shot* into a luciferase construct, and co-transfected each reporter with individual plasmids expressing several variants of Ham. In addition to the Ham long isoform and a PR truncation version of Ham, we included the NZF and the CZF domains fused with the VP16 activation domain (NZF_VP16 and CZF_VP16) (**Fig. 4C**). The two ZF domains of Ham are putative DNA binding domains, but it remained undetermined whether they help to recruit Ham to DNA. This experimental setting allows to assess which of the ZF domains is able to target Ham-regulated enhancers. The full-length, the PR truncation and the NZF_VP16 fusion all activated expression of the reporters driven by 7 out of 8 Ham-bound elements (**Fig. 4D-G, Supplementary Fig. 5A-B**). By contrast, the CZF_VP16 had little activity on the reporters. This result confirms that Ham can directly activate target gene expression and that the NZF domain directly or indirectly recruits Ham to enhancers.

To further identify co-factors with which Ham interacts to regulate gene expression, we performed *de novo* motif discovery from Ham CUT&TAG peaks in mixed TE and GD tissues close to up-regulated and down-regulated genes in the *hamSK1* mutant using the MEME-ChIP software. This resulted in nine and five significantly enriched motifs from up- and down-regulated gene loci respectively (**Supplementary Fig. 5D**). While five motifs were shared between these two sets, the motifs for Dpp/BMP signaling factor MAD, the zinc-finger protein ZNF263 and two unknown factors were only enriched in the upregulated genes. The result implies that Ham may switch activation to repression activity when interacting with these co-factors in the context of RS development.

Ham downstream effector genes are required for male RS fertility.

Built upon our transcriptomic data, we next intended to identify genes crucial for the fusion
process. We reasoned that as the majority of TE terminal epithelial cells are lost in both *ham*
mutant alleles, the commonly downregulated genes in *ham^PRΔ* and *ham^SK1* TE (**Fig. 3B**) may
be expressed in the *wt* TE epithelial cells and can be and functionally important. We thus
focused on the top 241 common downregulated genes whose expression showed a fold change
greater than 1.5, and the three validated Ham target genes: *shot*, *wnt2* and *shg*. Using all
available RNAi lines from the Bloomington stock center that cover 113 out of 244 candidate
genes, we performed a UAS-RNAi screen with the *UAS-dcr2*; *hamRSGal4* driver for male
fertility (see Methods).

The RNAi screen led to identification of 26 genes whose knockdown abolished or reduced
male fertility (**Supplementary Table 2**). Knocking down four of these genes, *shg*, *AP2-alpha*,
pebble (pbl) and *CG11406*, resulted in complete sterility, indicating their importance for male
fertility. We defined this category as 100% penetrance of sterility. Knockdown of other twenty-
two genes reduced fertility to different extent (**Fig. 5A-B**).

Consistent with the gene network analysis (**Fig. 4A**), these genes include cytoskeleton
regulators (*pbl*, *shg*, *shark*) [41-43] and metabolism factors (*CG11406, Ph4alphaMP, GstE11,*
CG17477, CG11069, AdSL, CG17283, CG15117, CG7484). Other genes are known or
predicted to be involved in intracellular membrane trafficking (*AP-2alpha*, *CG31935* and
CG7456) [44] and signaling (*wnt2*, *Toll*, *AkhR*, *Idgf4* and *Hr46*) [45-48].

We confirmed that knocking down *pbl*, *AP-2alpha, CG11406* and *shg* led to failed TE and SV
fusion and abnormal morphology at the pupal stage (**Fig. 5C**). *pbl* encodes a Rho guanine
nucleotide exchange factor, implicated in multiple processes involving cytoskeleton
reorganization[41, 49, 50]. The gene product of CG11406 is a predicted triglyceride lipase,
cleaving triacylglycerol (TAG) to produce diacylglycerol (DAG). AP-2alpha, a subunit in the
Adaptor complex 2, is involved in intracellular trafficking and secretion [51]. Remarkably, *shg*
knockdown also led to aberrant mobilization of TE epithelial cells to the middle of the TE,
while no such effect was found in the SV and AG. Downregulation of E-Cad is a key event
during epithelial to mesenchymal transition (EMT), and E-Cad promotes cohesive cell
migration of mesenchymal cells, but suppressing cell motility of epithelial cells [52, 53]. Our
result suggests that the SV and the TE terminal epithelia have distinct dependence on E-Cad

and that TE epithelial cells are prone to becoming motile when E-Cad is reduced. Together,
this screen identified a collection of known and novel factors involved in epithelial tissue
fusion, confirming that the fusion process requires coordinated activity of several classes of
molecules.
400
Ham downstream genes exhibit spatial- temporal expression dynamics in the male RS
402
During epithelial fusion, cells at the leading edge become specialized and express specific
genes required for tissue interaction and remodeling [1, 2]. Moreover, as we observed, the TE
terminal and SV epithelia have distinct properties, suggesting that these two sets of epithelial
cells may express distinct genes. We thus intended to assess spatial distribution of Ham
downstream genes. We conducted a high-resolution multiplexed RNA *in situ* hybridization
assay known as SCRINSHOT (single cell RNA *in situ* hybridization on tissues) [54] to
visualize expression of candidate genes including *wnt2*, *shg*, *shot* and other positive genes from
the RNAi screen, along with the receptor genes in Wnt2 signaling (*fz*, *fz2*, *fz3* and *fz4*) and
ligand genes in Tl-signaling (*spz*, *spz4*, *spz5* and *spz6*) (**Supplementary Table 3**). Some of
these Wnt2 and Tl signaling genes displayed dysregulation in at least one of the *ham* mutant
alleles according RNA-seq data (**Supplementary Fig 6A-B**). Other receptor genes in Wnt
signaling and ligand genes in Tl signaling were omitted due to low read numbers in the RNA-
seq data, indicating negligible or minimal expression in the TE and GD.
416
Each SCRINSHOT reaction took place within a mini-chamber, accommodating pairs of control
and *ham* mutant tissues, or GFP KD control and ham RNAi KD samples (refer to Methods for
details of probe design and experimental procedures). This setup was designed to minimize
technical variability between the genotypes. mRNAs are visualized as fluorescent dots in
SCRINSHOT images, with each dot representing a single mRNA molecule. However, due to
the densely packed nature of cells in the TE and SV tissues, quantifying mRNA molecules at
a single-cell resolution is challenging. Therefore, we opted to quantify fluorescent signal
intensity within defined areas (see Methods). In total, 37 genes were tested, including *ham*
itself (**Supplementary Table 3**). Probes for 7 genes gave either no or low signal in the TE or
GD, likely attributed to poor probe quality, given their RNA-seq read numbers not being low.
Resembling Ham protein expression (**Fig. 1G**), *ham* mRNAs were detected in the TE terminal
epithelium (**Fig. 6A**), confirming the specificity of the SCRINSHOT method. Similarly, *wnt2*,
shg and *shot* all exhibited high expression in the TE terminal epithelium (**Fig. 6A**).

In total, 25 genes exhibited detectable SCRINSHOT signal in the wt SV and AG, while 29 displayed signals in the wt TE (**Supplementary Table 3**). Within the TE, 17 out of 29 were specifically localized in the TE terminal epithelial cells, with the remaining 12 detected only in other TE cells (**Fig. 6, Supplementary Fig. 7, Supplementary Table 3**).

Subsequently, our analysis focused on the genes expressed in the TE terminal and SV epithelia. While 17 genes exhibited expression in both epithelial ends (**Fig. 6A-B**), 8 were exclusively expressed in SV epithelial cells (**Fig. 6F**, **Supplementary Table 3**), underscoring the heterotypic nature of the TE and SV epitheliums. Notably, certain genes (CG11406, *AP-2alpha*, *GstE11*, *Tl, spz5* and *spz6*) demonstrated temporal dynamic expression in the TE epithelium: spz5 and CG11406 are present during the 20-25h APF timeframe, prior to fusion, but gone during the 28-32h APF period when the TE and SV are undergoing fusion; *AP-2alpha*, *GstE11*, *Tl* and *spz6* showed the opposite trend, being expressed only later (**Supplementary Fig. 7B**).

In the GD, while the majority of genes exhibited ubiquitous expression, *wnt2* and *spz* were predominantly expressed in the leading cells within the SV, contrasting with their expression in the remaining GD (**Fig. 6B**). Additionally, the expression patterns of *fz2*, *fz3* and *Tl* displayed spatial specificity: fz2 exhibited higher expression levels in the SV compared to the AG, *fz3* demonstrated the opposite pattern with higher expression in the AG, and *Tl* was primarily expressed in the myoblasts (**Fig. 6B**). These findings imply that Wnt2 and Tl signaling may have cell-type-specific roles in the fusion process. Thus, SCRINSHOT mapped 30 candidate genes in the developing GD and TE, with 29 of them display spatial or temporal dynamics.

Wnt2 and Tl signaling pathways are key mediators of Ham function in RS tissue assembly

Next, we aimed to identify genes exhibiting altered expression in *ham* mutant SV and TE. As *hamSK1* mutant TE lacks terminal epithelial cells, we excluded TE samples and only quantified SCRINSHOT signal levels in the developing SV between *wt* and the *hamSK1* mutant. In contrast, *ham* RNAi knockdown resulted in fusion defects but only mildly impacted SV morphology and TE terminal epithelial cell number (**Fig. 2G-H**). These samples allow to assess gene expression alterations in both the SV and TE epithelia.

ham fluorescent intensity appeared comparable between *wt* and mutant samples (**Fig. 6E,**
Supplementary Fig. 7E), suggesting that *ham* mutant RNAs, containing the CRISPR-
generated indels, remain stable at this stage. In the *hamSK1* mutant SV, seven genes became
downregulated, while two exhibited upregulation. The downregulated genes were four Wnt2
pathway genes (*wnt2, fz, fz2, fz4*), one Tl pathway gene (*spz*), and *shg* (**Fig. 6B**-C). Notably,
wnt2, *fz*, *spz* and *shg* also displayed downregulation in *ham* RNAi SV (**Fig. 6D-E**), suggesting
that these genes and their associated signaling pathways may be directly involved in the fusion
process downstream of Ham. The upregulated genes in *hamSK1* mutant SV include *GstE11*
and *CG11069* (**Supplementary Fig.7C, E**), which exhibited low expression in the leading cells
of wt SV epithelium, but increased expression under the *hamSK1* condition. In *ham* RNAi
knockdown animals, *CG11069* was again upregulated, while *GstE11* did not change
expression. Instead, another ligand gene in Tl signaling (*spz5*) displayed increased expression
(**Supplementary Fig.7D, F**). On the TE side, only four genes showed expression alteration in
ham RNAi animals, including *ham* itself, *shg*, *fz* and *fz2* **Supplementary Fig.7G**), implicating
functional requirement of these genes in both TE and SV epithelia.
479
The observation that knocking down Tl resulted in decreased male fertility (**Fig. 5B**,
Supplementary Table 2), coupled with the expression of multiple *spz* ligand genes in the
developing TE and SV, suggests that the Tl signaling pathway might be required for assembling
the TE and SV. To investigate this possibility, we examined male flies with *spz* knockdown
driven by *hamRSGal4*. These animals produced a comparable number of progenies to control
animals. The lack of phenotype in this condition could be attributed to inefficient knockdown
or genetic compensation by another *spz* gene, like *spz5* whose expression increased in *ham*
RNAi animals when *spz* expression was reduced. In addition to Tl, two other Tl related genes
(Toll-7 and Toll-9) exhibited moderate number of reads in the RNA-seq data. We then tested
male fertility in *Toll-7* and *Toll-9* knockdown animals. *Toll-9* knockdown led to a drastic
reduction in progeny numbers compared to the RNAi control (**Supplementary Fig. 8A**).
Consistent with the sterility, the adult SV in the *Toll-9* knockdown male lacked sperms
(**Supplementary Fig. 8A**), suggesting failed connection between the TE and SV. Together,
these results pinpoint an essential role of Tl signaling in male RS development.
494
495   To validate the role of Wnt2 signaling in RS development, we investigated four genetic
496   conditions, aimed at modulating its level and activity: *wnt2^O^* (a *wnt2* null allele), *wnt2*
497   overexpression driven by *hamRSGal4*, *fz* and *fz2* RNAi knockdown driven by *hamRSGal4*. The

wnt2^O mutant TE and GD never connect, suggesting that Wnt2 is essential for the fusion process. The *wnt2^O* mutant TE lacks the pigment cell and myocyte layers (**Fig. 7A**), consistent with its reported function in these two cell types [17, 39]. The *wnt2^O* mutant GD failed to form elongated SV and AG, suggesting that Wnt2 is required for GD patterning and differentiation. Moreover, overexpression of *wnt2* also resulted in fusion defects and mis-localization of TE epithelial cells (**Fig. 7B**), a phenotype reminiscent of *shg* knockdown animals. Indeed, *wnt2*-overexpressing cells displayed reduced levels of E-Cad, suggesting that Wnt2 signaling might balance E-Cad levels to maintain TE epithelial cell properties. Consistently, *fz2* knockdown animals exhibited comparable phenotypes, including mobilized TE epithelial cells and failed tissue fusion (**Fig. 7C**). *fz2* and *fz* knockdown also decreased male fertility (**Fig. 7D**). Collectively, these results suggest the importance of optimal Wnt2 signaling levels in male RS development and epithelial tissue fusion.

Discussion

In this study we demonstrate that the transcription factor Ham plays a crucial role in controlling epithelial fusion during *Drosophila* RS development. Early in embryonic development, Ham determines the cell fate of msSGPs, which are progenitor cells for TE terminal epithelial cells. Later on, Ham directly regulates the differentiation of epithelial cells in both the TE and SV by orchestrating a gene expression program in epithelial cells. While our primary focus is on male RS, we found that Ham is also expressed in cells at the fusion site of the oviduct and ovary. Mutant females lacking functional Ham exhibit a non-functional oviduct epithelial tube, suggesting that Ham likely employs a similar mechanism to control epithelium fusion in females.

By examining *ham* mutant phenotypes and its downstream target genes, we identified several classes of novel factors involved in fusion process of SV and TE. These genes, directly or indirectly controlled by Ham, encompass a diverse array of molecular functions. This discovery provides a comprehensive gene network for future investigation of epithelial fusion processes. The cell-type-specific and tissue-specific spatial distribution of these genes suggest that they may regulate various aspects of fusion, including cell migration, intercellular communication between myoblasts and epithelial cells, specification of the leading cells, and other steps in the fusion process. For instance, Tl and its related receptors is known to mediate cell adhesions [55]. We can speculate that Tl signaling and the molecule itself promote interaction between myoblasts and myoblast and epithelial cells (**Fig. 7E**), based on high expression of Tl in myoblasts and presence of multiple Spz ligands in the TE and SV epithelial cells. Moreover, *Wnt2*, *fz* and *fz2* are highly expressed in the leading cells of TE and SV epithelia, and also required for TE and SV connection, suggesting that Wnt2 signaling mediates epithelial cell communication, remodeling and fusion.

Despite being discovered more than twenty years ago [28], research on the function of Ham function has primarily focused on its roles in the nervous system. Our findings highlight a crucial function of Ham in epithelial tissues. Apart from the RS, we observed high expression of the Ham protein in other epithelial tissues including the Malpighian tubes (akin to the kidney in insects) and the intestine (data not shown). Furthermore, orthologs of Ham, PRDM16 and PRDM3/MECOM, exhibit high expression in epithelial tubular structures, such as Lung, pancreas, kidney, urinary bladder and gastrointestinal tract and lung, according to observations

in the Human Protein Atlas (**Supplementary Fig. 9**.
https://www.proteinatlas.org/ENSG00000142611-PRDM16/single+cell+type;
https://www.proteinatlas.org/ENSG00000085276-MECOM/single+cell+type) [56]. Notably,
Seminal vesicle and ductus deferens, the equivalent structure to fly seminal vesicle, are the top
one and three tissues with the highest PRDM16 expression (**Supplementary Fig. 9A**). This
suggests a potential role for these proteins in controlling epithelial integrity and fusion in
mammalian organogenesis. Indeed, we observed that the pancreas duct volume in mouse
embryos lacking *Prdm16* is significantly reduced (data not shown). Although the exact cellular
changes and the underlying molecular mechanisms in this context await further investigation,
this phenotype underscores a crucial function of *Prdm16* in pancreatic epithelial duct
formation.

We show that compared to Ham function in the nervous system, the PR-containing isoforms
are indispensable for RS epithelial fusion and differentiation, and that Ham can directly activate
epithelial gene expression. It remains to be determined whether Ham has distinct molecular
functions or it regulates a subset of common molecular pathways in these two contexts.
Dendrite morphogenesis involves outgrowth and branching, somewhat analogous to epithelial
tube growth and branching. For instance, molecular mechanisms regulating polarized protein
trafficking are crucial for both dendrite and epithelial tube branching [57]. Additionally, several
transcription factors like Sequoia [58, 59], Tramtrack [60, 61] and Cut [62, 63] play roles in
both dendrite morphogenesis and tracheal tube fusion. These examples suggest that there might
be more unrecognized common mechanisms between dendrite and epithelial morphogenesis.
It is of interest to test whether the effectors we identified in the RS epithelium are involved in
dendrite morphogenesis in the nervous system.

We observed a unique aspect of epithelial fusion in the *Drosophila* RS. Unlike many epithelial
fusion processes where two homologous ends merge, such as those in mammalian neural tube
closure and *Drosophila* tracheal formation *etc* [1, 8, 10], the two epithelial ends in the
Drosophila RS are heterotypic. Specifically, the TE terminal epithelial cells express higher
levels of E-Cad and lower levels of Cor compared to SV epithelial cells. In response to the
reduction of E-Cad or ectopic expression of *wnt2*, TE terminal epithelial cells become motile,
resembling a process akin to epithelial-to-mesenchymal transition. This suggests that TE
terminal epithelial cells may be more invasive during epithelial fusion. On the other hand, SV
epithelial cells, despite originating from mesenchymal cells, exhibit typical epithelial cell

characteristics, including regular Cor and E-Cad expression. Notably, many genes are expressed exclusively in the SV epithelium, distinguishing it from the TE epithelium. Heterotypic epithelial fusion occurs in many developmental processes, such as digestive system formation, vertebrate optic fissure closure, and mammalian urogenital development. Failure of epithelial fusion can lead to a wide variety of developmental diseases such as oesophageal atresia, spina bifida, ocular coloboma, congenital abnormalities of the kidney, urinary tract and persistent cloaca. Studying the basic processes and key players of normal epithelial tissue fusion in an *in vivo* context will help to understand the pathogenesis of these diseases.

Acknowledgements

We thank the imaging facility (IFSU) of Stockholm University, the BEA sequencing facility at Karolinska Insitute, and the National Genomic Institute of Scilife Laboratories, Sweden, for providing service and support. The Bloomington Stock Center and the *Drosophila* community provided important reagents and fly stocks. We thank Christos Samakovlis's lab for helping to set up SCRINSHOT and Eric Lai's lab for the initial Ham reagents. Funding from Australian Research Council Centre of Excellence for the Mathematical Analysis of Cellular Systems (CE230100001) and ANU Future Scheme to J.W. The project was supported by project grant from Swedish Research Council (Vetenskapsrådet, 2020-03543) and Swedish Cancerfonden (CAN 22 2161 Pj) to Q.D.

Author contributions

Conceptualization and project design by QD, experiments conduction by HW, preliminary test in pancreas by IA and LB, computation analysis by JW. Data interpretation by HW, JW and QD. Manuscript writing by QD with input from HW and JW.

Materials and methods

Fly strain culturing and generation of transgenes

All fly stocks were maintained at 22°C. The *ham^{PRΔ}* mutants were created using CRISPR: transgenic flies carrying single guide RNA targeting the PR domain containing isoforms were crossed into the vas-Cas9 transgenic flies. Frame-shift mutations were identified by PCR and Sanger sequencing. Transgenic flies were created at BestGene, Inc.

The ham Gal4 lines and RNAi stocks were obtained from Bloomington Stock Center. Knocking down experiments were performed at 25 °C and with a regular 12-hr light–dark cycle.

To generate ham null mutant somatic clones in the TE terminal epithelium, *hamsk1 FRT40A/CTG* males were crossed with virgin females carrying the genotype *hsFLP(X); hsGFP.nls FRT40A*. Embryos aged 0-5 hours were collected and developed at 25°C until the 2nd instar larval stage. GFP-negative larvae, indicating the genotype *hsFLP(X); hsGFP.nls/hamsk1 FRT40A*, were then selected. Head shock treatments at 37°C for 1 hour were administered three times during the 2nd and 3rd instar larval stages to induce somatic clones. Prior to dissection, another 1-hour head shock at 37°C was performed to induce GFP expression, marking the wild-type cells.

Fertility test

For ham mutants, single male or female with indicated genotype was mated with two w^{1118} virgin females or two w^{1118} males on day 0 of the experiment. Crosses were flipped every 2 days for 3 consecutive times. Progenies from the second vial with eggs laid between day 2-4 were counted after 2 weeks.

For the RNAi screen, individual UAS-RNAi males were crossed with females carrying UAS-dcr2 and hamRSGal4. The resultant F1 male progenies (2-5 days old) with the genotype UAS-dcr2; hamRSGal4>UAS-RNAi were crossed to w^{1118} virgin females (1-2 days old) on day 0 of the experiment. In each test, four males were crossed with three females. Crosses were flipped on day2 and day3 for pre-laying. After flipping on day3, crosses were kept in the new vial for egg laying for a specific time period (GroupA: 48hours; GroupB: 16-18h; GroupC: 7-8h). If at

least three males and two females were alive, then the flies were transferred to a new vial, the

old vial was kept, and all progenies were counted.

For fertility test of the Wnt2 and Tl signaling pathways components, UAS-RNAi males were

crossed to virgin females of the genotype UAS-dcr2; hamRSGal4. Two F1 male progenies (1-

2 days old) with the genotype UAS-dcr2; hamRSGal4>UAS.RNAi were crossed to three w[1118]

virgin females (1-2 days old) on day 0 of the experiment. Crosses were flipped every day for 4

consecutive times. Progenies from the forth vial with eggs laid between day 3-4 were counted

after 2 weeks. Crosses were flipped every day for 4 consecutive times. Progenies from the forth

vial with eggs laid between day 3-4 were counted after 2 weeks.

Immunostaining

For embryo staining: embryos were collected, dechorionated with bleach and fixed with 4%

formaldehyde in PEM (0.1M PIPES, 2mM EGTA, 1 mM MgSO4, pH 7) for 20 min. For

immunostaining, embryos were rehydrated, washed with PBT (0.3% Triton X-100 in PBS) and

blocked with PBSBT (0.3% Triton X-100, 1% BSA in PBS).

For tissue staining: tissues were dissected from stage defined larvae or pupae in 1xPBS within

30 min and fixed in 4% paraformaldehyde in PBS for 20min. For immunostaining, dissected

tissues were washed with PBT (0.1% Triton X-100 in PBS) and blocked with PBSBT (0.1%

Triton X-100, 1% BSA in PBS).

For both embryo and tissue staining, the primary antibodies were diluted with PBSBT (0.1%

Triton X-100, 1% BSA in PBS) and incubated at 4°C over night. After extensive washing with

PBSBT (15min x 4 times), the secondary antibodies were added in 1:500 dilution and incubated

at room temperature for 2h. After washing with PBT (0.1% Triton X-100 in PBS), the samples

were counter-stained with DAPI and mounted in Vectashield (Vector Laboratories).

The following primary antibodies were used: Rabbit anti-Ham (1:300) (Gift from Eric Lai's

lab), Ms anti-V5 (1:500, R960-25, invitrogen), chk anti-GFP (1:2000, ab13970, Abcam), Rat

anti-vasa (1:50, anti-vasa-c, Developmental Studies Hybridoma Bank, DSHB), Ms anti-ABD-

B (1:50, 1A2E9-c), Ms anti-SXL (1:50, M18-c), rat anti-DE-Cad (1:50, DCAD2, DSHB),

mouse anti-Crb (1:10, Cq4 DSHB), mouse anti-Coracle (1:100, C615.16 DSHB). Secondary

antibodies conjugated to Cy3 or Cy5 or Alexa Fluor-488 (Jackson ImmunoResearch) were
used.
702
703 *Quantification of signal intensity from fluorescent immunostaining*
704
For E-Cad and Crb in Figure 2, "raw intensity density" from manually defined regions (using
free-hand drawing) along the apical side of the SV and TE terminal epitheliums was obtained
in ImageJ. For each sample, 2 regions in the SV and 3 regions in the TE terminal epithelium
were averaged. The regions used to determine relative E-Cad fluorescence in *hamSK1* TE
somatic clones were at the apical side between cell junction. The length of Cor-stained cell
junction was measured using ImageJ. A straight line was drawn over the Cor staining with the
'Freehand Line tool'. For each sample, the average length from 10 cell-cell junctions were
used. In all experiments, images for the genotypes in comparison were always processed using
exact settings. At least three biological replicates were included in statistical analysis.
714
Western Blot
Embryos from *w1118, ham$^{PR\Delta}$/CTG, ham^{SK1}/CTG* were collected and aged to embryonic stage
S13 to S16. In each blot, 50-100st GFP+ or GFP- embryos were used. The selected embryos
were homogenized directly in 1xSDS loading buffer (62.5 mM Tris-HCl pH 6.8, 2.5 % SDS,
0.002 % Bromophenol Blue, 0.7135 M (5%) β-mercaptoethanol, 10 % glycerol) and denatured
at 98 °C heat blot for 5 min. After centrifuged, supernatant was taken for Western-Blotting
with 8% of SDS-PAGE. Rabbit anti-Ham (1:2000) primary antibody was used in blotting. To
minimize unspecific signal from IgG heavy chains, light chain specific secondary antibody of
mouse anti-Rabbit IgG (1:40000 Jackson ImmunoResearch) was used, and the signals were
developed with ECL Plus reagent (GE Healthcare).
725
Cell culture and Luciferase assays
To generate different variants of Ham, the open reading frames of HamFL, Ham-NZF-vp16,
Ham-CZF-vp16, and *ham$^{PR\Delta}$* were PCR amplified and cloned into the pAC_V5 (nitrogen)
vector. The enhancer fragments were cloned in to the p2xTetO-Firefly luciferase vector. All
transfections were performed using *Drosophila* S2-R+ cells grown in Schneider *Drosophila*
medium containing 10% fetal calf serum. Cells were co-transfected with the Ham expression
construct, 2xTetO-Firefly luciferase and pAc-Renilla plasmids in 96-well plate using the

Effectene Transfection kit (Qiagene). Luciferase assays were performed and measured using

the Dual Luciferase Assay System (Promega). Expression was calculated as the ratio between

the firefly and *Renilla* luciferase activities. Three rounds of transfections were performed on

different days and different cell populations and considered as three biological replicates.

Within each biological replicate, 4 wells of cells were transfected with the same DNA mixture

and measured and thus considered as 4 technique replicates. In total, for each DNA

combination, the values from the 12 wells of transfection were calculated, averaged and plotted

in Fig. 5D-F. Cloning oligos are provided in **Supplementary table 4**.

SCRINSHOT

The Drosophila reproductive system was dissected at defined ages in 1xPBS and fixed in 4%

paraformaldehyde in PBS for 20min. After washing with PBST 0.1% (Triton X-100) for 2 x

15 min, fine dissected samples were placed on Poly-lysine coverslip and freezed at -80°C after

drying on a heater.

The SCRINSHOT experiments were carried out according to the published method [54]. In

brief, 3 padlock probes and 3 corresponding detection probes were designed for each gene of

interest. The slides with fine dissected samples were pretreated at 45°°C to reduce moisture and

fixed in 4% PFA in 1X PBS, followed by washing in PBS Tween-20 0.05% twice. Samples

were then dehydrated with 70%, 85%, 100%ETOH and air dry. The SecureSeal hybridization

chamber (GRACE BIO-LABS, 621501) was then mounted to cover samples on the slide.

Permeabilization of tissues were first done by treating with 10μg/mL Proteinase K (Invitrogen)

in PBSTween-20 0.05% for 7 minutes followed by washing with Glycine (2mg/mL in PBS

Tween-20 0.05%) once and with PBS Tween-20 0.05% for 3 x 5 minutes. After this, samples

were further Permeabilized in 0.1M HCl for 3min, followed by washing in PBS Tween-20

0.05% twice. Samples were then blocked in a probe-free hybridization reaction mixture of

1XAmplifase buffer (Lucigen, A1905B), 0.05M KCl, 20% Formamide deionized (Millipore

S4117), 0.1uM Oligo-dT, 0.1ug/ul BSA (New England Biolabs, B9000S), 1U/ul RiboLock

(Thermo, EO0384), and 0.2ug/ul tRNAs (Ambion, AM7119). Then hybridization of padlock

probes was done by incubating samples with padlock probes (with the concentration of each

one 0.01uM) mixed in blocking reagents used before (no oligo dT used in this step). The slide

was then put into a PCR machine to denature at 55°C for 15 minutes and hybridize at 45°C for

120 minutes. Padlock probes were then ligated by using SplintR ligase (NEB M0375) at 25°C

for 16 hours followed by RCA (rolling cycle amplification) at 30°C for 16 hours by using phi29

polymerase (Lucigen, 30221–2) and RCA primer1. Then a fixation step was applied to stabilize

RCA product in 4% PFA for 15 minutes followed by washing in PBS Tween-20 0.05%. Then

the hybridization of the first 3 genes was done by mixing all 3 3' fluorophore-conjugated

detection probes of each gene in reaction reagent (2XSSC, 20% Formamide deionized, 0.1ug/ul

BSA, and 0.5 ng/μl DAPI (Biolegend, 422801) followed by hybridization at 30˚C for 1 hour.

The slides were washed in 20% formamide in 2X SSC and then in 6X SSC, followed by

dehydration in 70%, 85%, 100% Ethanol until the chamber was removed. Then the samples

were preserved in SlowFade Gold Antifade mountnant (Thermo, S36936) and kept in dark

before imaging. After image acquisition, the first 3 detection probes were removed by using

Uracil-DNA Glycosylase (Thermo, EN0362), and the slides were ready for the next round of

detection probe hybridization. The procedure was repeated until all genes were hybridized and

imaged. Images were acquired with an airyscan-equipped confocal microscope system (Zeiss

LSM 800, Carl Zeiss), and ZEN 2.5 Blue edition software was used for image processing.

Quantification of SCRINSHOT signals

Similar to quantification of immunofluorescent signal, fluorescence intensity of SCRINSHOT

for each gene was measured in defined areas within images of the SV, EjD, and TE in 2D

projections. Squares were drawn to define areas at the leading side of the SV, the middle region

of the Ejd and the TE terminal. The total fluorescence intensity was first normalized to area

size. The resultant value for *ham* mutant or RNAi samples was then contrasted to the

normalized intensity of control samples.

CUT&TAG

CUT&TAG was performed according to the published method [37]. In brief, TE and grossly

tissue of the posterior abdomen were dissected from 18-25h APF *w1118* and

pBAC_ham_sfGFP animals in PBS within 30 min. For grossly tissues, the dissected samples

were homogenized in 2ml NE1 buffer followed by incubation on ice for 10 min. After filtering

out the grossly cuticles, nuclei were collected by spinning at 1300xg and resuspended in 1ml

PBS. For TE, nuclei extraction was done by simply pipetted samples in 200ul NE1 up and

down for 20-30 times and incubated on ice for 10 min. Light fixation was performed with 0.1%

formaldehyde at RT for 2 minutes and neutralized with 75mM Glycine. Then the nuclei were

washed by PBS and wash buffer 1 time each and kept in 200ul wash buffer on ice until enough

samples were collected (30-50st animals per/antibody). Resuspended nuclei in 1.2ml wash

buffer and mixed with 20ul Concanavalin A-coated magnetic beads (Polyscience 86057) with

soft vortexing. Aliquoted samples into two parts immediately and incubated for 10 min with

rotation to allow the binding of the nuclei with the beads. The nuclei/beads mix was then blocked with 1mL cold antibody buffer for 10 min and resuspended in 150ul cold antibody buffer. For w1118 samples, 1uL Rabbit anti-Ham or 1ul Rabbit IgG (SIGMA-ALDRICH, I5006) was added for an overnight incubation at 4°C. For *pBAC_ham_sfGFP* samples, 1ul Rabbit anti-GFP (G10362, invitrogen) or Rabbit IgG was used. The beads/nuclei/antibody mix was then washed with Dig-wash buffer and incubated with secondary antibody (1:100 dilution) for 1 hour. After further washes with the Dig-wash buffer, a pA-Tn5 adaptor complex in Dig-300 buffer was added to the beads for 1 hour reaction. The beads were then washed with Dig-300 buffer before the incubation with 300ul Tagmentation buffer at 37°C for 1 hour. Then 10ul 0.5M EDTA, 3uL 10% SDS and 2.5uL 20 mg/mL proteinase K (Invitrogen) were used to stop the reaction. Then the resultant DNA was purified with DNA clean & Concentrator kit (ZYMO Research D4013) and eluted in 25uL Elution buffer. To generate libraries, 21uL fragment DNA was mixed with 2uL 10uM Universal i5 primer, 2uL 10uM uniquely barcoded i7 primer and 25uL PCR master mix (NEB Phusion® High-Fidelity PCR Master Mix with HF Buffer, M0531). The PCR condition was as follows: 72°C 5min, 98°C 30s, repeat 12 times (98°C 10s, 63°C 10s), 72°C 1 min and hold 4°C. The libraries were cleaned with standard Ampure XP beads as previously described. Libraries from four biological replicates were produced for each condition.

We mapped the CUT&TAG samples of *w1118* and *pBAC_ham_sfGFP* in TE and TG tissues to the dm6 genome assembly using Bowtie2 with the following parameters: bowtie2 --end-to-end --very-sensitive --no-mixed --no-discordant --phred33 -I 10 -X 700. The coverage for these samples was generated using bedtools genomeCoverageBed and normalized to library depth to yield reads per million per base (RPM). For peak calling, we employed SEACR [64], which is tailored for sparse chromatin profiling data like CUT&TAG. We first used a relaxed FDR cutoff of < 0.5 for each replicate (SEACR_1.3.sh normalized_coverage 0.5 norm stringent output_peaks), we then identified common peaks across replicates for each condition. To select confident peaks for *w1118* and *pBAC_ham_sfGFP*, we overlapped their peaks within TE and TG separately, choosing the top 1800 peaks from TE and the top 6000 from TG. Peak annotation was performed by HOMER's annotatePeaks.pl. Peak overlap analysis was performed using HOMER's mergePeaks function with the default -d given parameter, ensuring that only directly overlapping peaks were merged. This method preserved the integrity of the original peak boundaries and maintained the specified distances between peaks from the initial peak calls. *De novo* motif discovery for peaks in up-regulated or down-regulated genes,

positioned 2000 bp upstream and 500 bp downstream of the TSS in TE and TG tissues, was

performed using the MEME-ChIP software. The parameters used were: -ccut 100, -meme-p 5,

-dna, -meme-mod anr, -minw 5, -maxw 15, and -filter-thresh 0.05.

RNA-seq

Total RNA was extracted from finely dissected genital discs or testis samples from 18-25h APF

animals using Trizol reagent (Invitrogen). RNA quality was tested by Agilent Bioanalyzer.

RNA-seq libraries were made using the Illumina Truseq Total RNA library Prep Kit LT.

Sequencing was performed on the Illumina Hiseq2500 platform. Following the trimming of

adapter sequences with Trimmomatic, RNA-seq reads from replicated wild type (x3) and

mutant samples (x3) were aligned to the *Drosophila melanogaster* (dm6) genome assembly

utilizing HISAT2. Gene annotations were sourced from UCSC. The DESeq2 Bioconductor

package was employed to identify differentially expressed mRNAs between *w1118* and the

ham^{PRΔ}/*ham^{PRΔ}* mutant, as well as between +/Df and *ham^{SK1}*/Df mutant, with a subsequent

estimation of the false discovery rate (FDR). Genes with an FDR < 10% and absolute

Log2FC>0 were considered differentially expressed. Details of the calculated differential

expression are provided in **Supplementary table 5**.

STRING gene network analyses

Ham activated (genes downregulated in *hamSK1* and bound by Ham in TG) and repressed

(genes upregulated in *hamSK1* and bound by Ham in TG) gene networks are constructed using

String, MCL clustering (find natural clusters the stochastic flow, inflation parameter, 2.

Disconnected genes are hidden. The position of the clusters is manually adjusted to fit related

GO categories. The links to the gene networks in STRING:

https://string-db.org/cgi/network?taskId=bbF8IgMTqf1M&sessionId=bzDIAZvMgWsi

https://string-db.org/cgi/network?taskId=bhKCcMoLlbwh&sessionId=bzDIAZvMgWsi

References

1. Bernascone, I., M. Hachimi, and F. Martin-Belmonte, *Signaling Networks in Epithelial*
Tube Formation. Cold Spring Harb Perspect Biol, 2017. **9**(12).
2. Chung, S. and D.J. Andrew, *The formation of epithelial tubes.* J Cell Sci, 2008. **121**(Pt
21): p. 3501-4.
3. Samakovlis, C., et al., *Development of the Drosophila tracheal system occurs by a*
series of morphologically distinct but genetically coupled branching events.
Development, 1996. **122**(5): p. 1395-407.
4. Samakovlis, C., et al., *Genetic control of epithelial tube fusion during Drosophila*
tracheal development. Development, 1996. **122**(11): p. 3531-6.
5. Sutherland, D., C. Samakovlis, and M.A. Krasnow, *branchless encodes a Drosophila*
FGF homolog that controls tracheal cell migration and the pattern of branching. Cell,
1996. **87**(6): p. 1091-101.
6. Tanaka-Matakatsu, M., et al., *Cadherin-mediated cell adhesion and cell motility in*
Drosophila trachea regulated by the transcription factor Escargot. Development,
1996. **122**(12): p. 3697-705.
7. Gervais, L., G. Lebreton, and J. Casanova, *The making of a fusion branch in the*
Drosophila trachea. Dev Biol, 2012. **362**(2): p. 187-93.
8. Jacinto, A., A. Martinez-Arias, and P. Martin, *Mechanisms of epithelial fusion and*
repair. Nat Cell Biol, 2001. **3**(5): p. E117-23.
9. Vasioukhin, V., et al., *Directed actin polymerization is the driving force for epithelial*
cell-cell adhesion. Cell, 2000. **100**(2): p. 209-19.
10. Pai, Y.J., et al., *Epithelial fusion during neural tube morphogenesis.* Birth Defects Res
A Clin Mol Teratol, 2012. **94**(10): p. 817-23.
11. Caviglia, S. and S. Luschnig, *Tube fusion: making connections in branched tubular*
networks. Semin Cell Dev Biol, 2014. **31**: p. 82-90.
12. de Mello Santos, T. and B.T. Hinton, *We, the developing rete testis, efferent ducts,*
and Wolffian duct, all hereby agree that we need to connect. Andrology, 2019. **7**(5):
p. 581-587.
13. Jemc, J.C., *Somatic gonadal cells: the supporting cast for the germline.* Genesis, 2011.
49(10): p. 753-75.
14. DeFalco, T., S. Le Bras, and M. Van Doren, *Abdominal-B is essential for proper*
sexually dimorphic development of the Drosophila gonad. Mech Dev, 2004. **121**(11):
p. 1323-33.
15. Whitworth, C., E. Jimenez, and M. Van Doren, *Development of sexual dimorphism in*
the Drosophila testis. Spermatogenesis, 2012. **2**(3): p. 129-136.
16. Ahmad, S.M. and B.S. Baker, *Sex-specific deployment of FGF signaling in Drosophila*
recruits mesodermal cells into the male genital imaginal disc. Cell, 2002. **109**(5): p.
651-61.
17. Rothenbusch-Fender, S., et al., *Myotube migration to cover and shape the testis of*
Drosophila depends on Heartless, Cadherin/Catenin, and myosin II. Biol Open, 2017.
6(12): p. 1876-1888.
18. Fog, C.K., G.G. Galli, and A.H. Lund, *PRDM proteins: important players in*
differentiation and disease. Bioessays, 2012. **34**(1): p. 50-60.

19.   Chi, J. and P. Cohen, *The Multifaceted Roles of PRDM16: Adipose Biology and*
Beyond. Trends Endocrinol Metab, 2016. **27**(1): p. 11-23.

20.   Bard-Chapeau, E.A., et al., *Ecotopic viral integration site 1 (EVI1) regulates multiple*
cellular processes important for cancer and is a synergistic partner for FOS protein in
invasive tumors. Proc Natl Acad Sci U S A, 2012. **109**(6): p. 2168-73.

21.   Zhou, B., et al., *PRDM16 Suppresses MLL1r Leukemia via Intrinsic Histone*
Methyltransferase Activity. Mol Cell, 2016.

22.   Corrigan, D.J., et al., *PRDM16 isoforms differentially regulate normal and leukemic*
hematopoiesis and inflammatory gene signature. J Clin Invest, 2018. **128**(8): p. 3250-
3264.

23.   Cibi, D.M., et al., *Prdm16 Deficiency Leads to Age-Dependent Cardiac Hypertrophy,*
Adverse Remodeling, Mitochondrial Dysfunction, and Heart Failure. Cell Rep, 2020.
33(3): p. 108288.

24.   Kundu, A., et al., *PRDM16 suppresses HIF-targeted gene expression in kidney cancer.*
J Exp Med, 2020. **217**(6).

25.   Horvath, C. and C. Scheele, *PRDM16 stability and metabolically healthy adipose*
tissue. Nat Metab, 2022. **4**(9): p. 1095-1096.

26.   Wang, Q., et al., *Post-translational control of beige fat biogenesis by PRDM16*
stabilization. Nature, 2022. **609**(7925): p. 151-158.

27.   Hurwitz, E., et al., *Antagonism between Prdm16 and Smad4 specifies the trajectory*
and progression of pancreatic cancer. J Cell Biol, 2023. **222**(4).

28.   Moore, A.W., L.Y. Jan, and Y.N. Jan, *hamlet, a binary genetic switch between single-*
and multiple- dendrite neuron morphology. Science, 2002. **297**(5585): p. 1355-8.

29.   Moore, A.W., et al., *Conversion of neurons and glia to external-cell fates in the*
external sensory organs of Drosophila hamlet mutants by a cousin-cousin cell-type
respecification. Genes Dev, 2004. **18**(6): p. 623-8.

30.   Rives-Quinto, N., et al., *Sequential activation of transcriptional repressors promotes*
progenitor commitment by silencing stem cell identity genes. Elife, 2020. **9**.

31.   Eroglu, E., et al., *SWI/SNF complex prevents lineage reversion and induces temporal*
patterning in neural stem cells. Cell, 2014. **156**(6): p. 1259-73.

32.   Endo, K., et al., *Chromatin modification of Notch targets in olfactory receptor neuron*
diversification. Nat Neurosci, 2012. **15**(2): p. 224-33.

33.   Pinheiro, I., et al., *Prdm3 and Prdm16 are H3K9me1 methyltransferases required for*
mammalian heterochromatin integrity. Cell, 2012. **150**(5): p. 948-60.

34.   Hohenauer, T. and A.W. Moore, *The Prdm family: expanding roles in stem cells and*
development. Development, 2012. **139**(13): p. 2267-82.

35.   Gokcezade, J., G. Sienski, and P. Duchek, *Efficient CRISPR/Cas9 plasmids for rapid and*
versatile genome editing in Drosophila. G3 (Bethesda), 2014. **4**(11): p. 2279-82.

36.   Jenett, A., et al., *A GAL4-driver line resource for Drosophila neurobiology.* Cell Rep,
2012. **2**(4): p. 991-1001.

37.   Kaya-Okur, H.S., et al., *CUT&Tag for efficient epigenomic profiling of small samples*
and single cells. Nat Commun, 2019. **10**(1): p. 1930.

38.   Szklarczyk, D., et al., *The STRING database in 2023: protein-protein association*
networks and functional enrichment analyses for any sequenced genome of interest.
Nucleic Acids Res, 2023. **51**(D1): p. D638-D646.

39.   Kozopas, K.M., C.H. Samos, and R. Nusse, *DWnt-2, a Drosophila Wnt gene required*
for the development of the male reproductive tract, specifies a sexually dimorphic cell
fate. Genes Dev, 1998. **12**(8): p. 1155-65.

40.   Deshpande, G., A. Nouri, and P. Schedl, *Wnt Signaling in Sexual Dimorphism.*
Genetics, 2016. **202**(2): p. 661-73.

41.   van Impel, A., et al., *Regulation of the Rac GTPase pathway by the multifunctional*
Rho GEF Pebble is essential for mesoderm migration in the Drosophila gastrula.
Development, 2009. **136**(5): p. 813-22.

42.   Oda, H., et al., *A Drosophila homolog of cadherin associated with armadillo and*
essential for embryonic cell-cell adhesion. Dev Biol, 1994. **165**(2): p. 716-26.

43.   Ferrante, A.W., Jr., R. Reinke, and E.R. Stanley, *Shark, a Src homology 2, ankyrin*
repeat, tyrosine kinase, is expressed on the apical surfaces of ectodermal epithelia.
Proc Natl Acad Sci U S A, 1995. **92**(6): p. 1911-5.

44.   Gonzalez-Gaitan, M. and H. Jackle, *Role of Drosophila alpha-adaptin in presynaptic*
vesicle recycling. Cell, 1997. **88**(6): p. 767-76.

45.   Weber, A.N., et al., *Binding of the Drosophila cytokine Spatzle to Toll is direct and*
establishes signaling. Nat Immunol, 2003. **4**(8): p. 794-800.

46.   Staubli, F., et al., *Molecular identification of the insect adipokinetic hormone*
receptors. Proc Natl Acad Sci U S A, 2002. **99**(6): p. 3446-51.

47.   Kawamura, K., et al., *A new family of growth factors produced by the fat body and*
active on Drosophila imaginal disc cells. Development, 1999. **126**(2): p. 211-9.

48.   Lam, G.T., C. Jiang, and C.S. Thummel, *Coordination of larval and prepupal gene*
expression by the DHR3 orphan receptor during Drosophila metamorphosis.
Development, 1997. **124**(9): p. 1757-69.

49.   Prokopenko, S.N., et al., *A putative exchange factor for Rho1 GTPase is required for*
initiation of cytokinesis in Drosophila. Genes Dev, 1999. **13**(17): p. 2301-14.

50.   Nakamura, M., J.M. Verboon, and S.M. Parkhurst, *Prepatterning by RhoGEFs governs*
Rho GTPase spatiotemporal dynamics during wound repair. J Cell Biol, 2017. **216**(12):
p. 3959-3969.

51.   Lin, Y.H., et al., *AP-2-complex-mediated endocytosis of Drosophila Crumbs regulates*
polarity by antagonizing Stardust. J Cell Sci, 2015. **128**(24): p. 4538-49.

52.   Campbell, K. and J. Casanova, *A role for E-cadherin in ensuring cohesive migration of*
a heterogeneous population of non-epithelial cells. Nat Commun, 2015. **6**: p. 7998.

53.   Chen, H., et al., *E-cadherin mediates adhesion and suppresses cell motility via distinct*
mechanisms. J Cell Sci, 1997. **110 (Pt 3)**: p. 345-56.

54.   Sountoulidis, A., et al., *SCRINSHOT enables spatial mapping of cell states in tissue*
sections with single-cell resolution. PLoS Biol, 2020. **18**(11): p. e3000675.

55.   Umetsu, D., *Cell mechanics and cell-cell recognition controls by Toll-like receptors in*
tissue morphogenesis and homeostasis. Fly (Austin), 2022. **16**(1): p. 233-247.

56.   Karlsson, M., et al., *A single-cell type transcriptomics map of human tissues.* Sci Adv,
2021. **7**(31).

57.   Corty, M.M., B.J. Matthews, and W.B. Grueber, *Molecules and mechanisms of*
dendrite development in Drosophila. Development, 2009. **136**(7): p. 1049-61.

58.   Araujo, S.J. and J. Casanova, *Sequoia establishes tip-cell number in Drosophila*
trachea by regulating FGF levels. J Cell Sci, 2011. **124**(Pt 14): p. 2335-40.

59. Brenman, J.E., et al., *Sequoia, a tramtrack-related zinc finger protein, functions as a*
pan-neural regulator for dendrite and axon morphogenesis in Drosophila. Dev Cell,
2001. **1**(5): p. 667-77.
60. Araujo, S.J., C. Cela, and M. Llimargas, *Tramtrack regulates different morphogenetic*
events during Drosophila tracheal development. Development, 2007. **134**(20): p.
3665-76.
61. Li, S., et al., *Photoreceptor cell differentiation requires regulated proteolysis of the*
transcriptional repressor Tramtrack. Cell, 1997. **90**(3): p. 469-78.
62. Pitsouli, C. and N. Perrimon, *The homeobox transcription factor cut coordinates*
patterning and growth during Drosophila airway remodeling. Sci Signal, 2013. **6**(263):
p. ra12.
63. Grueber, W.B., L.Y. Jan, and Y.N. Jan, *Different levels of the homeodomain protein cut*
regulate distinct dendrite branching patterns of Drosophila multidendritic neurons.
Cell, 2003. **112**(6): p. 805-18.
64. Meers, M.P., D. Tenenbaum, and S. Henikoff, *Peak calling by Sparse Enrichment*
Analysis for CUT&RUN chromatin profiling. Epigenetics Chromatin, 2019. **12**(1): p. 42.
1019

1020 **Declaration of the use of AI**. ChatGPT was used to improve readability of the text, and the

1021 content was thoroughly checked and edited by the authors to ensure accuracy.

1022

1023

1024

1025

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

    Figure legends

    Figure 1. *ham* is expressed and required for *Drosophila* male RS development

    (**A**). Illustration of the developing male RS. The testis (TE) terminal epithelial cells localize at

    the opposite side of the germline stem cell niche. The genital disc (GD) contains mesenchymal

    cells (MC) that will develop into the SV and accessory gland (ag). (**B**). Illustration of the TE

    and SV fusion process. APF, after pupal formation. (**C**). The adult male RS. ejd, ejaculatory

    duct. (**D**). White-field images of adult male RS, showing defective morphology in the *ham*

    mutants. (**E**). Quantification of male fertility. Ham_sfGFP with the *ham* genomic region was

    used to rescue the sterility. Statistical significance was calculated using unpaired t-test (***,

    $p<0.001$; **, $p< 0.01$; *, $p<0.05$; n.s., non-significant). (**F**). Images of the adult RS stained with

    Ham_V5 in red. DNA is stained with DAPI in blue. (**G**). Images of the larval TE (top) and

    GD (bottom) stained with Ham_V5 in red, the reporter in green (btlGal4 driving UAS-GFP)

    and DAPI in blue. AEL, after egg laying. (**H**). Images of the pupal GD stained with Ham_V5

    in red, the reporter in green (btlGal4 driving UAS-GFP) and DAPI in blue. APF, after pupal

    formation. (**I**) Images of the pupal RS with Ham_V5 in red and DAPI in blue. (**J**). Higher

    magnification image of the joint site between the SV and TE, with myoblasts labeled by

    mef2Gal4>GFP.

    Figure 2. Ham promotes differentiation of the TE terminal and SV epithelial cells

    (**A**). Images of the male RS at the pupal stage, showing morphological defects of the TE and

    SV in the ham mutants. Epithelial cells are labeled with Coracle (Cor). (**B**). High magnification

    images of the connection site of the TE and SV, showing that the ham mutant RS failed to form

    a continuous tube. Ham-sfGFP restored the normal morphology. Epithelial cells are marked by

    E-Cad in red, Cor in yellow, and the epithelium on the SV side also marked by btlGal4 driving

    UAS-mCD8GFP. DNA is labeled by DAPI in blue. (**C**). Images and quantification of the E-

    Cad signal in the *wt* and *hamPRΔ* mutant SV. (**D**). Images and quantification of the Crumbs

    signal in the *wt* and *hamPRΔ /Df* mutant SV. (**E**). Images and quantification of the length of

    Cor signal in the wt and *hamPRΔ* mutant SV. (**F**) Images of *ham* mutant mosaic clones in the

    TE terminal epithelial cells. Note, the mutant cells are GFP negative and within the area

    surrounded by the white dashed line. The white arrows indicate junctions between mutant cells,

    while the yellow arrows highlight the junctions between wild-type cells. (**G**). Images of pupal

    male GD, showing that *hamRNAi*-mediated knockdown reproduces *ham* mutant phenotypes.

(**H**). Images of pupal male RS, stained with E-Cad in red, Ham in yellow, GFP in green and

DAPI in blue. UAS-dcr2 enhanced knockdown efficiency and led to more severe

morphological defects. (**I**). Images and quantification of E-Cad signal in red and Cor length in

blue in control and ham RNAi SV. (**J**) Images and quantification of E-Cad signal in control

and ham RNAi TE terminal epithelium. Statistical significance was calculated using unpaired

t-test except for TE mosaic clones which used paired t-test. (***, $p<0.001$; **, $p< 0.01$; *,

$p<0.05$; ns, non-significant).

1082

1083   **Figure 3. Identification of Ham-regulated genes in the developing TE and GD**

1084

(**A**). Illustration of the sample types and antibodies used in the RNA-seq and CUT&TAG

experiments. (**B**) Venn diagram of dysregulated genes between the two mutant conditions in

the GD (left) and TE (right) samples. (**C**) Heatmaps of Ham CUT&TAG peak density and IgG

control, ranked from high to low and centered at peak summits. (**D**) Venn diagrams of

CUT&TAG peaks from the two types of samples. (**E-F**). Volcano plots of dysregulated genes

in GD (E) and TE (F) in the *hamSK1/Df* mutant, and the ones with at least one Ham CUT&TAG

peak are highlighted.

1092

1093   **Figure 4. Ham activates expression of epithelial differentiation genes**

1094

(**A**) Ham gene network analysis from STRING. The position of MCL Clusters were manually

adjusted to place genes with similar function in close proximity. The functional categories were

determined by gene function tools in STRING. (**B**). A bar plot of the Log2 fold-change value

from RNA-seq for *wnt2*, *shot* and *shg*, showing downregulation of these genes in the

hamSK1/Df mutant. (**C**). Schemes of the Ham_FL and other truncation proteins, and illustration

of the luciferase reporter assay. (**D, F**). A genome browser snapshot of the *shg* and *wnt2* loci,

showing the CUT&TAG tracks from the TE and GD (TG) mixed samples and the enhancer

fragments in the luciferase assays. (**E, G**). Normalized luciferase activity to the empty reporter

and then to the control vector that expresses GFP and RFP proteins. Statistical significance was

calculated using unpaired t-test (****, $p<0.0001$; ***, $p<0.001$; **, $p< 0.01$; *, $p<0.05$; ns,

non-significant).

1106

1107   **Figure 5. A candidate RNAi screen identified Ham-downstream effectors in RS epithelial**

1108   **tissue fusion**

1109

(**A**). Scheme for the RNAi screen. (**B**) Summary of positive hits. The extent of sterility is

determined by the number of progenies from the F1 male crossing to the wild-type female (see

Methods). 100% means no progeny. >70%, the number of progenies being fewer than 30% of

the number of progenies produced by the control cross (the same Gal4 driver x UAS-

GFPRNAi). 40-70%, the number of progenies being between 30-60% of the number of

progenies produced by the control. (**C**). Images of the control and gene-specific RNAi

knockdown RS. The left panels show the fusion site stained with E-Cad in red, GFP in green,

Ham in yellow and DAPI in blue, the middle panels are GD, and the right ones are TE. The

orange arrows indicate mobilized epithelial cells in the *shg* knockdown TE.

1119

1120 **Figure 6. Spatial-temporal expression dynamics of Ham-downstream genes**

1121

(**A**). Images from multiplexed in situ hybridization (the SCRINSHOT method), showing

mRNA signal of the indicated genes in *wt* TE samples. DNA is marked by DAPI in grey. (**B**).

Images from SCRINSHOT, showing mRNA signals of indicated genes in *wt/Df* and

hamSK1/Df mutant GD. The last panel is an enlarged view from the sample with *Tl*, *ham* and

spz probes. The yellow arrowheads indicate myoblast cells with Tl signal. (**C**). Quantification

of signal intensity for genes shown in B in two areas, the SV and EjD. Statistical significance

was calculated using unpaired t-test (***, p<0.001; **, p< 0.01; *, p<0.05; ns, non-significant).

(**D**) Images from SCRINSHOT, showing mRNA signals of indicated genes in *GFP* and *ham*

RNAi GD samples. (**E**). Quantification of signal intensity for genes shown in D in two areas,

the SV and EjD. (**F**). A summary of genes expressed in the TE terminal and SV epithelium.

Highlighted genes show temporal or spatial dynamic expression in the two tissues. Statistical

significance was calculated using unpaired t-test (****, p<0.0001; ***, p<0.001; **, p< 0.01;

*, p<0.05; ns, non-significant).

1135

Figure 7. Wnt2 and its ligand genes are required for male fertility and RS development

(**A**) Images of *wnt2o* mutant TE and GD, showing disrupted morphology in these tissues. (**B**).

Images of the TE and SV fusion site from the control and *wnt2* overexpression animals. The

epithelial cells of the TE and SV are labeled with hamRSGal4 driving UAS-mCD8GFP in

green and E-Cad in red and Ham in yellow. The white arrows indicate mobilized GFP positive

cells in the middle of the TE, and these cells have a lower level of E-Cad. (**C**). Images of the

TE and SV fusion site from the control and *fz2* RNAi animals. The white arrows indicate

mobilized GFP positive cells in the middle of the TE, and these cells have a lower level of E-Cad. (**D**) Graphs of progeny number from male fertility tests for *fz* and *fz2* RNAi lines in comparison to GFP RNAi. Number of tested males in each genotype, 3. (**E**) Illustration of cell-cell interactions in the developing RS, and a speculated model for Tl and Wnt2 signaling functions. Wnt2 signaling may be involved in epithelial cell interaction between the two ends. Tl signaling and Tl may be involved in the interaction between epithelial cells and myoblasts. In three conditions (*shgRNAi*, *wnt2OE* and *fz2RNAi*), some TE terminal epithelial cells (illustrated in orange) move up to the distal end of the testis.

A Testis (TE)
Hub cells — GLCs
Pigment cells
Te epithelial cells
MC
GD

B Te epithelial cell
TE
SV epithelial cell
SV
24h APF

TE
SV
28h APF

TE
Pigment cell
Myoblast
SV
36h APF

C TE
sv
ag — ejd

D
TE
SV
AG
500uM
w1118 control   *hamPRΔ /hamPRΔ*   *hamSK1/Df*

E male fertility
Number of progenies
-50
***
***
n.s
***
**
***
***
W1118
hamPRΔ
hamPRΔ/Df
hamPRΔ; ham.sfGFP
hamPRΔ/Df; ham.sfGFP
hamSK1/Df; ham.sfGFP

F *RS Adult*
TE
sv
ejd
ag
50um
DAPI Ham_V5

G *Testis 109-120h AEL*
TE
50um
MC
GD
50um
DAPI btlGal4>GFP Ham_V5

H *GD 28h APF*
50um
DAPI btlGal4>GFP Ham_V5

I *RS 50h APF*
sv
TE
ag
50um
Ham_V5 DAPI

J *RS 50h APF*
sv
TE
50um
mef2Gal4>mCD8GFP
Ham_V5 DAPI

A

hamRSGal4>UAS-dcr2; UAS-mCD8GFP

X

UAS-RNAi
(135 lines for 113 genes)

↓

Screen for F1 male sterility

↓

26 positive gene hits

B

Sterility	Genes
100%	*pbl, AP-2alpha, CG11406, shg*
>70%	*GstE11, CG7456, Gr64c, AkhR, Idgf4*
40%-70%	*CG17477, CG11069, CG17186, CG4872, AdSL, Tl, CG17283, decay,PH4alphaMP, Hr46, CG15117, gsb-n, shark, CG17018, cag, CG7484, CG31935*

C

A

wnt2O

E-Cad Cor DAPI

48h APF TE

20um

24-30h APF GD

B

32-48h APF   E-Cad Ham mCD8GFP DAPI

hamRSGal4>
mCD8GFP

50um

hamRSGal4>
mCD8GFP
UASWnt2

50um

C

40-50h APF TE   E-Cad GFP DAPI

GFP RNAi

20um

fz2 RNAi

hamRSGal4>UASmCD8GFP

D

Number of progenies

GFP.RNAi   fz2.RNAi_1   fz2.RNAi_2   fz.RNAi

E

wt

TE

Wnt2/Fz/Fz2

SV

Tl

Tl/Spz/Spz5

TE

SV

wnt2OE fz2RNAi shgRNAi

TE

SV

↔ Epithelial remodeling and interaction
↔ Myoblast-myoblast interaction
→ Myoblast migration
↔ Epithelial-myoblast interaction

◖ ◗ Mobilized epithelial cell
◉ Myoblast and myocyte
◞ Pigment cell